

Interactive comment on “Response of *Nodularia spumigena* to pCO₂ – Part I: Growth, production and nitrogen cycling” by N. Wannicke et al.

J. Czerny

jczerney@ifm-geomar.de

Received and published: 17 March 2012

The approach of the experiment presented by Wannicke et al. investigating the effect of future elevated CO₂ concentrations on growth of *Nodularia* under phosphorus limited conditions can be seen as a reasonable continuation of the experiment presented by Czerny et al. (2009). In Czerny et al. (2009), cellular division and nitrogen fixation rates were found to decrease in response to increasing CO₂. With carbon and phosphorus cellular uptake rates being unaffected, lowered division rates resulted in enhanced intracellular accumulation of these elements. In the Czerny et al. study, low density semi-continuous batch cultures were grown in artificial YBC II media providing inorganic phosphorus, vitamins and trace metals in surplus. This approach offers a high level of reproducibility and measured effects can be attributed to CO₂ / pH treat-

C361

ments with a high certainty. However, investigating only exponential growth, results can at best be applied to natural pre-bloom conditions when growth is not limited by nutrients and cannot be applied to high density bloom situations under multiple growth limitation. Experiments simulating such conditions, like the present study, are urgently needed, but are technically more difficult to control. A more sophisticated experimental setup (chemostat) instead of simple batch cultures is generally recommended when CO₂ effects are studied under nutrient limiting conditions. In batch cultures, the physiological state of the stock culture can have substantial influence on the lag phase after inoculation. An acclimation of batch cultures to treatment conditions by a minimum of 5 generations at exponential growth should therefore be standard procedure in culture experiments. General guidelines for culture experiments are given in LaRoche et al. (2011) and work cited herein. It appears that the pre-culture media used in Wannicke et al. were not equilibrated with gas mixtures prior to inoculation, but were only bubbled during the three days of acclimation. It is important to know when and if treatment levels were established in the pre-cultures so that found results can be classified as short term or long term response. Information about phosphate concentration, pCO₂, pH and growth rate of the pre-cultures is missing. Experimental batch cultures were started with low amounts of inorganic phosphorus. After three days, initially supplied phosphate (0.35 μmol kg⁻¹) was obviously depleted and therefore another 0.35 μmol kg⁻¹ of phosphate was added. Overall, growth rates in this experiment were low (0.1–0.2 d⁻¹) relative to *Nodularia* cultures growing under nutrient replete conditions ($\mu=0.319$ /d-1 in Wannicke et al., 2009); $\mu\sim 0.3$ –0.5 d-1 in Czerny et al., 2009). It is not reported whether or when cultures entered the stationary growth phase. Moreover, it should be mentioned which cultures were already in a decaying state (indicated by decreasing cell numbers P 2493 L 9) during the last phase and whether they are included in the analyses. Another issue of the present study is the range, precision and documentation of the pCO₂ and pH treatment levels. The “Guide to best practices for ocean acidification research and data reporting” is referenced by Wannicke et al. for the choice of gas mixtures used to manipulate carbonate chemistry. Supplied CO₂ mixing ratios

C362

180, 380, and 750 ppm are used as treatment levels by Wannicke et al., but measured values (Table 1) range between ~ 480 and $\sim 590 \mu\text{atm CO}_2$ at the start of the experiment. Obviously, equilibration was not achieved after the three days of bubbling. Pure CO_2 , gas mixtures or acid can be used to adjust CO_2 treatments (Schulz et al., 2009), but it is mandatory to refer to achieved treatment levels when describing measured responses. It has to be stressed that treatment range is overall small and not directly comparable to present summer conditions in the Baltic (Thomas and Schneider, 1999) or to CO_2 scenarios by the IPCC (Meehl et al., 2007). As common summer surface pCO_2 concentrations in the Baltic Sea are usually below the applied concentrations, it is also misleading to refer to the treatments as “past”, “present” and “future”. Determining carbonate chemistry equilibrium using potentiometric pH measurement on the NBS scale results in high measurement uncertainty (Dickson, 2011). Therefore, it should be stated in the manuscript whether standard deviations presented in Table 1 are measurement uncertainties or differences in equilibration of the treatment replicates. If replicate bottles considerably differ in pCO_2 / treatments overlap, it would probably be better to use regression analyses evaluating each experimental unit at its individual treatment level instead of using ANCOVA, requiring replication of treatments. Presented data can hardly be compared to results of previous studies without appropriate reference to the physiological state of the cultures and the actual treatment CO_2 levels.

Literature:

Czerny, J., Barcelos e Ramos, J., and Riebesell, U.: Influence of elevated CO_2 concentrations on cell division and nitrogen fixation rates in the bloom-forming cyanobacterium *Nodularia spumigena*, *Biogeosciences*, 6, 10.5194/bg-6-1865-2009, 2009.

Dickson, A. G.: The carbon dioxide system in seawater: equilibrium chemistry and measurements, in: Guide to best practices for ocean acidification research and data reporting edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J. P., Publication Office of the European Union, Luxembourg, 260, 2011.

C363

LaRoche, J., Rost, B., and Engel, A.: Bioassays, batch culture and chemostat experimentation, in: Guide to best practices for ocean acidification research and data reporting edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J. P., Publication Office of the European Union, Luxembourg, 260, 2011.

Meehl, G. A. S., T. F., Collins, W. D., Friedlingstein, A. T., Gaye, A. T., Gregory, J. M., Kitch, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z. a.: Global climate projections., in: *Climate Change 2007, The physical science basis*, edited by: Solomon, S., D. Qin, M. Manning, Technical support unit, IPCC working group 1, Cambridge University Press, 748-845, 2007.

Schulz, K. G., Barcelos e Ramos, J., Zeebe, R. E., and Riebesell, U.: CO_2 perturbation experiments: similarities and differences between dissolved inorganic carbon and total alkalinity manipulations *Biogeosciences*, 6, 2145-2153, 2009.

Thomas, H., and Schneider, B.: The seasonal cycle of carbon dioxide in Baltic Sea surface water, *Journal of Marine Systems*, 22, 53-67, 1999.

Wannicke, N., Koch, B. P., and Voss, M.: Release of fixed N_2 and C as dissolved compounds by *Trichodesmium erythreum* and *Nodularia spumigena* under the influence of high light and high nutrient (P), *Anglais*, 57, 175-189, 2009.

Interactive comment on *Biogeosciences Discuss.*, 9, 2481, 2012.

C364