

Interactive
Comment

Interactive comment on “Feedbacks of CO₂ dependent dissolved organic carbon production on atmospheric CO₂ in an ocean biogeochemical model” by L. A. Bordelon-Katrynski and B. Schneider

L. A. Bordelon-Katrynski and B. Schneider

lb@gpi.uni-kiel.de

Received and published: 25 October 2012

REFeree: 1 General comments

The manuscript “Feedback of CO₂ -dependent dissolved organic carbon production on atmospheric CO₂ in an ocean biogeochemistry model” by L. Bordelon-Katrynski and B. Schneider is a contribution to the important question how changes in ocean biogeochemical cycling induced by rising atmospheric pCO₂ feed back on the ocean-atmosphere carbon cycle. The specific feedback that they study here is an increased

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



excretion rate of dissolved organic carbon (DOC) that has been hypothesized to explain findings from a mesocosm experiment (Riebesell et al., 2007): Here it was shown that increasing $p\text{CO}_2$ led to an increased rate of carbon drawdown relative to that of nitrogen, while the particulate C:N ratio remained unchanged. The hypothesized explanation was that the additional carbon taken up was routed into DOC. Besides thus changing the overall stoichiometry of organic matter this pathway has the potential to change particle sinking rates in the ocean, as part of the DOC pool, acidic polysaccharides, tend to aggregate and to also enhance aggregation of other particles into larger faster-sinking particles.

The authors study this feedback with a global ocean biogeochemical model with which they perform experiments where the excretion rate of DOC from phytoplankton changes over time following a simple relationship to the expected atmospheric $p\text{CO}_2$ increase. They compare this to a model experiment where the excretion rate stays constant over time; the difference between the two runs can then be discussed as following from the DOC feedback. Both runs are also performed with and without a parallel increase in $p\text{CO}_2$, so interactive effects can also be investigated.

The consequences of CO_2 induced changes in C:N stoichiometry have already been investigated before with a similar model (Tagliabue et al., 2011). However, the detailed assumptions how the additional carbon taken up is routed into dissolved or particulate biomass differ between the two studies, as are some key findings (increased vs. decreased particulate export under elevated $p\text{CO}_2$). The authors cite this as evidence that the sign of the feedback depends on the actual pathway the extra carbon is taking.

The general question of the manuscript concerning the magnitude of the DOC feedback as well as the discussion on how it depends on the way that they are implemented in biogeochemical model are worthwhile additions to the field. However, I have a problem with the assumptions that the authors made in parameterizing the $p\text{CO}_2$ -DOC feedback, a problem that was also shared by the other reviewer, and that I think also determines the outcome of the study in such a way that it cannot be interpreted

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

meaningfully. Much to my regret I therefore cannot recommend to publish the study in Biogeosciences, although, apart from the way of parameterization the whole study is methodologically well done and the manuscript is well written.

My problem is the following. The whole point in the mesocosm experiments by Riebesell et al. (2007) was that increasing the availability of CO₂ to phytoplankton led to an increased photosynthetic production of organic carbon by phytoplankton without an parallel increase in nitrogen uptake; and it seemed as if most of the excess carbon taken up was routed into dissolved organic carbon, thereby increasing the C:N ratio of dissolved organic matter (DOM). The model used in the present study, however, uses a constant C:N ratio both for particulate (phytoplankton/zooplankton/detritus) and dissolved organic matter compartments. At least this is what I took from the references that the authors give for their model, I have found no explicit statement on the C:N ratio in DOM in their manuscript. But there are several clear indicators that the C:N ratio in DOM is indeed fixed, e.g. the shallower depth of nitrate remineralization (p. 7991, l. 20-21). In a model with fixed C:N an increased relative excretion of DOC must be accompanied by an excretion of dissolved organic nitrogen, and thus by a reduction in the production of phytoplankton biomass. As the supply of inorganic nitrogen is what is often limiting the possible formation and sinking of biomass (at least in models) it is therefore no wonder that the authors observe that the increased formation of DOC is at the expense of the formation of POC rather than fostering particle aggregation (p. 7991, l. 11-15). So I would argue that

1. the main model result of decreased particle export is at least partly built into their model assumption, namely that the feedback operates through increased relative DOC excretion at fixed C:N ratio, and that

2. this assumption is inconsistent with the findings in the mesocosm experiments cited. Indeed I do not see a physiological reason why an increase in seawater pCO₂ should lead to an increase in organic carbon and nitrogen excretion at the expense of cell growth, while there are good physiological reasons why an increased CO₂ supply could

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

lead to an increased carbon fixation (e.g. the often low affinity of the carboxylating enzyme RuBisCO), and why excess carbon fixed without additional nitrogen uptake would lead to the formation of high C:N biomass, such as sugars.

My summary of the paper is thus that it is a valid sensitivity study for a physiological effect that does not exist. I think the authors could remedy this by allowing the stoichiometry of DOM to vary, while keeping that of the particulate biomass constant. That would, however, require to re-run all model runs, in effect it would be a new study.

I must say that, given the otherwise good methodological quality of the study, I would be happy if the authors could come up with convincing arguments why I am wrong and their study is meaningful, but at the moment I do not see any.

REPLY: We would like to thank the Referee for the very well pointed and fair critique of our manuscript. It probably needs to be explained more clearly in the method's section that the focus of our study is on the mechanism of particle aggregation (Arrigo 2007) not C:N decoupling (carbon overconsumption). In fact, we decided to use constant C:N ratios in our study, since several studies have done detailed analyses on the stoichiometry of the dissolved and particulate matter pools (Riebesell et al. 2007, Engel et al. 2008, Schulz et al. 2008, Kim et al. 2011) and none of them found significant changes in the stoichiometry of either pool in mesocosm studies. We are well aware that it is unfortunate that the fate of the increased carbon draw-down in the mesocosm experiments by Riebesell et al. (2007) could not be localized, which is also mentioned by Reviewer #2, who states: '... it seemed as if most of the excess carbon taken up was routed into dissolved organic carbon, thereby increasing the C:N ratio of dissolved organic matter (DOM).' However, in the absence of clear indications for stoichiometric shifts in DOM, both ours and the study of Tagliabue et al. 2011 serve as ideal complements with regard to the question of the fate of DOM and particle fluxes in the water column under ocean acidification.

We will be happy to more carefully address the differences between constant and de-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

coupled C:N ratios under DOM formation in our revised version. We are also eager to include the suggestions of reviewer #1 to perform a sensitivity analysis on critical concentrations of POM and DOM, explaining regionally enhancing particle export. Consequently, we do not see the need to perform new simulations with decoupled C:N ratios in DOM. Instead, this would be the repetition of a well-conducted study (Tagliabue et al. 2011), probably not providing many new insights.

Interactive comment on Biogeosciences Discuss., 9, 7983, 2012.

BGD

9, C5182–C5186, 2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C5186

