Biogeosciences Discuss., 9, C5204–C5211, 2012 www.biogeosciences-discuss.net/9/C5204/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD

9, C5204–C5211, 2012

Interactive Comment

Interactive comment on "Net sea-air CO₂ flux uncertainties in the Bay of Biscay based on the choice of wind speed products and gas transfer parameterizations" by P. Otero et al.

P. Otero et al.

pablo.otero@co.ieo.es

Received and published: 26 October 2012

First at all, we are grateful for your review and we really appreciate your careful reading of the manuscript. You state that this work is potentially interesting, however you suggest that "the present manuscript is not suitable for publication in Biogeosciences". From our point of view, this work deserves publication in this scientific journal, after a revision of some aspects that we will clarify below.

Your main disappointment concerns with the combination of different wind speed products (U) and different gas transfer coefficients (k). We agree with you when you refer to Section 6.1 of Takahashi et al. (2009) to highlight that, if a specific scale factor is



Full Screen / Esc

Printer-friendly Version



used to compute the gas transfer velocity as previous step to compute CO2 fluxes. Then the same wind product in the original study should only be used. We are aware that there is a dependence of the air-sea CO2 fluxes on the uncertainties of the wind speed products and the spatial and temporal resolution of the wind fields. Naegler et al. (2006) went deep into this issue re-assessing gas exchange scaling parameters for the Wanninkhof's parameterization (Table 1; Naegler et al., 2006) in order to correct the quadratic differences among wind fields (in situ measurements, satellite observations, climatology and reanalysis products...). As example, they obtained an overestimation of the NCEP wind speed. However this correction roughly appears inconsistent with the CO2 flux variability described in the Bay of Biscay. Using the Wanninkhof's parameterization as well, our results showed an overestimation of 16% of the fluxes estimated from QuikSCAT dataset. However, we suspect that if it would be possible to make the estimation, the normalization presented by Naegler et al. (2006) would lead to different results in coastal environments.

The high topographic variability of the continental margins are poorly represented by the terrain data models of the weather reanalysis models (e.g. NCEP-1 and NCEP-2) avoiding the correct estimation of land-sea interaction mechanisms. Continental margins also have particular impacts due to its spatial heterogeneity on the satellite wind observations (QuikSCAT and CCMP). In any case, coastline has an important impact on the wind speed measurements (Otero et al., 2008) in spite of being better represented by the new generation reanalysis models (ERA-Interim; Carvalho et al., 2012).

Precisely our main goal is evaluate the annual mean CO2 fluxes estimated in an arbitrary coastal region from five different wind speed data sets in order to going deep into the analyze of the uncertainty and variability of the coastal CO2 fluxes. In this sense, this effort is an useful case study of the discrepancy among coastal wind products, especially, when many of the articles computing CO2 fluxes in coastal regimes do not

BGD

9, C5204–C5211, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



take into account the recommendations described in Naegler et al. (2006).

Even the only scaling of air-sea CO2 fluxes in different continental margins (Cheng and Borges, 2009; Laurelle et al., 2010) do not take this into consideration in spite of showing large CO2 outgassing/uptake. Moreover, this analysis also gives an idea of the disagreement of using regional wind products (HIRLAM-AEMet...) as do biogeochemistry modellers in coastal regions and that cannot be evaluated following Naegler et al. (2006).

Therefore we will modify the manuscript to clarify the necessary correct selection of a wind speed product and the gas transfer parameterization. However we will keep the original analysis from the combination of gas transfer algorithms and different wind speed products.

1. The use of the Liss and Merlivat (1986) parameterization will be clarified in the revision. So, the previous paragraph:

" ... the main uncertainty remains in the estimation of the transfer velocity k which is, for its part, dependent on wind speed. there is not still consensus as to what is the adequate relatonship of k with de wind speed. .Various studies assume a linear {Liss, 1986 #248}, quadratic {Wanninkhof, 1992 #251}{Nightingale, 2000 #252}{Ho, 2006 #250}{Sweeney, 2007 #253} or cubic {Wanninkhof, 1999 #259}"

was replaced by this one:

"... the main uncertainty remains in the estimation of the transfer velocity k which is, for its part, dependent on wind speed. So, there is not still consensus as to what is the adequate relationship of k with de wind speed. Liss & Merlivat (1986) assumed three linear segments of gas transfer with different wind regimes from experiments in wind-wave tanks. Various studies suggested a quadratic dependence between gas transfer and wind speed, such as, Wanninkhof (1992) from the global bomb 14C inventory (Broecker et al. 1985) and Sweeney et al (2007) using more recent estimates

9, C5204–C5211, 2012

Interactive Comment



Printer-friendly Version

Interactive Discussion



of the carbon bomb radiocarbon inventory (Naegler and Levin, 2006). Nightingale et al (2000b) also computed a quadratic parameterization of the gas transfer velocity from field studies in coastal regions. Unlike these studies, McGillis et al (2001) described a gas transfer velocity controlled by breaking waves parameterized by cubic expression."

2. The equations of gas transfer velocity will be included.

3. The analysis of the temporal and spatial variations in pCO2 will be expanded in Section 3.3.

4. The value of 57% corresponding to the explained variability of the CO2 fluxes at long-term scale by the changes in the gas transfer velocity was estimated by Padin et al. (2009).

5. Comments of Fig.1 and Fig.4 have already been applied (see supplementary material to this response).

References:

Carvalho, D., Rocha, A., and Gómez-Gesteira, M. 2012. Ocean surface wind simulation forced by different reanalyses: Comparison with observed data along the Iberian Peninsula coast. Ocean Modelling, 56, 31-42.

Chen, C.T.A. and A.V. Borges. – 2009. Reconciling opposing views on carbon cycling in the coastal ocean: continental shelves as sinks and nearâĂŘshore ecosystems as sources of atmospheric CO2, Deep Sea Res., Part II, 56(8–10): 578-590, doi:10.1016/j.dsr2.2009.01.001.

Kanamitsu, M, W. Ebisuzaki, J. Woollen, S-K Yang, J .J. Hnilo, M. Fiorino, and G. L. Potter, 2002: NCEP-DOE AMIP-II Reanalysis (R-2),. Bull. Amer. Met. Soc., 83, 1631-1643.

Key, R. M., A. Kozyr, C. L. Sabine, K. Lee, R. Wanninkhof, J. L. Bullister, R. A. Feely, F. J. Millero, C. Mordy, and T.-H. Peng (2004), A global ocean carbon climatology:

9, C5204–C5211, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from Global Data Analysis Project (GLODAP), Global Biogeochem. Cycles, 18, GB4031, doi:10.1029/2004GB002247.

Laruelle, G.G., H.H. Dürr, C.P. Slomp and A.V. Borges. – 2010. Evaluation of sinks and sources of CO2 in the global coastal ocean using a spatiallyâĂŘexplicit typology of estuaries and continental shelves. Geophysical Research Letters, Vol. 37, L15607, doi:10.1029/2010GL043691.

Liss, P. S., and Merlivat, L.: Air–sea gas exchange rates: introduction and synthesis, in: The Role of Air-Sea Exchange in Geochemical Cycling, D. Reidel Publishing Company, 113-127, 1986.

Naegler, T., and I. Levin (2006), Closing the global radiocarbon budget 1945–2005, J. Geophys. Res., 111, D12311, doi:10.1029/2005JD006758.

Naegler, T., P. Ciais, K. B. Rodgers, and I. Levin (2006), Excess radiocarbon constraints on air-sea gas exchange and the uptake of CO2 by the oceans, Geophys. Res. Lett., 33, L11802, doi:10.1029/2005GL025408.

Otero, P. and Ruiz-Villarreal, M., 2008. Wind forcing of the coastal circulation off North and Northwest Iberia: comparison of atmospheric models, J. Geophys. Res., 113, C10019, doi:10.1029/2008JC004740.

Padin, X. A., Castro, C. G., Rios, A. F., and Perez, F. F.: fCO2sw variability in the Bay of Biscay during ECO cruises, Cont Shelf Res, 28, 904-914, 2008.

Padin, X.A., G. Navarro, M. Gilcoto, A.F. Rios and F.F. Pérez. – 2009 Estimation of air–sea CO2 fluxes in the Bay of Biscay based on empirical relationships and remotely sensed observations, J. Marine Sys., 75: 280-289.

Peacock, S. (2004), Debate over the ocean bomb radiocarbon sink: Closing the gap, Global Biogeochem. Cycles, 18, GB2022, doi:10.1029/2003GB002211.

Saha et al., 2010. The NCEP Climate Forecast System Reanalysis. American Meteo-

9, C5204–C5211, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rology Society, BAMS, 1015-1057. doi : 10.1175/2010BAMS 3001.1

Sweeney, C., Gloor, E. M., Jacobson, A. R., Key, R. M., Galen, M., Sarmiento, J. F., and Wanninkhof, R.: Constraining Global air-sea Gas Exchange for CO2 with recent Bomb 14C measurements, Global Biogeochem. Cycles, 21, GB2015, 10.1029/2006GB002784, 2007.

Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Kortzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO2, and net sea-air CO2 flux over the global oceans, Deep-Sea Res. Part II-Top. Stud. Oceanogr., 56, 554-577, 2009.

Wanninkhof, R.: Relationship Between Wind Speed and Gas Exchange Over the Ocean, J. Geophys. Res., 97, 7373-7382, 10.1029/92jc00188, 1992.

BGD

9, C5204–C5211, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactive comment on Biogeosciences Discuss., 9, 9993, 2012.

BGD

9, C5204-C5211, 2012

Interactive Comment



Printer-friendly Version

Interactive Discussion

Discussion Paper





Fig. 1. New Figure 1



Fig. 2. New Figure 4

BGD

9, C5204–C5211, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

