

Determination of the molecular signature of fossil conifers by experimental palaeochemotaxonomy. Part 1. The Araucariaceae family

Y. Lu¹, Y. Hautevelle¹, R. Michels¹

¹UMR 7359 CNRS GeoRessource, University of Lorraine, BP 239, 54506 Vandœuvre-lès-Nancy cedex, France.

Correspondence to: Y. HAUTEVELLE (yann.hautevelle@univ-lorraine.fr)

Abstract

Twelve species of the conifer family Araucariaceae family, including *Araucaria* (6 species), *Agathis* (3 species) and *Wollemia* (1 species) genera, were submitted to artificial maturation by confined pyrolysis. The aim of these experiments is to transform the biomolecules synthesized by these species into their homologous geomolecules in laboratory conditions. Determination of the diagenetic molecular signatures of Araucariaceae through experimentation on extant representatives allows to complete our knowledge in botanical palaeochemotaxonomy. Such knowledge is relevant to palaeoenvironmental environmental and archaeology purposes. All artificially diagenetic species of Araucariaceae are firstly characterized by a predominance of saturated tetracyclic diterpenoids including *ent*-beyerane, phyllocladanes and *ent*-kauranes. Moreover, the *Araucaria* genus shows a high relative abundance of bicyclic sesquiterpenoids, particularly the cadalane-type compounds accompanied by those of eudesmane-type, bisabolane-type as well as chamazulene, pentamethyl-dihydroindenes. Diterpenoids are of the labdane, isopimarane and abietane types (essentially derived from abietanoic acids) as well as isohexyl alkylaromatic hydrocarbons. Compared to the tetracyclic diterpenoids, these compounds show a relatively lower abundance, reaching trace levels in the case of saturated abietanes. Distributions of sesqui- and diterpenoids of *Agathis* show some similarities to that of *Araucaria*, with the exception of one species, in which the tetracyclic compounds are absent and the abietane-type (essentially derived from abietanoic acids) predominant. High similarities between the *Wollemia* and *Araucaria* genera are observed. Both are characterized by some high

relative abundance of tetracyclic compounds with no predominance of other specific diterpenoids.

1 Introduction

Numerous studies of the molecular composition of extant terrestrial plants have pointed out the chemotaxonomic values of many biological compounds bioterpenoids in particular (e.g. Aplin et al., 1963; Smith, 1976; Castro et al., 1996; Mongrand et al., 2001). This means that these biomolecules are synthesized by a restricted number of plant taxa and can be used as specific markers. While most bioterpenoids are degraded and their atomic constituents recycled in the surface processes of the Earth, a minor part is incorporated into sediments, thus joining the geological cycle. During this process, the bioterpenoids are transformed by diagenesis, leading to the formation of reaction products called geoterpenoids. Their initial chemotaxonomic value can be partially or totally retained (e.g. Simoneit, 1986; Otto and Simoneit, 2001). Geoterpenoids of ancient sediments may thus provide palaeochemotaxonomic information inherited from their biological precursors.

Botanical palaeochemotaxonomy has some specific attributes compared to palaeobotany and palynology in the reconstruction of palaeofloral and palaeoclimatic evolutions through geological times (e.g. Vliex et al., 1994; van Aarssen et al., 2000; Hautevelle et al., 2006a). Indeed, 1) plant biomarkers are more widespread in the stratigraphic record than well preserved plant macrofossils, 2) on the contrary to Palaeozoic and Mesozoic palynomorphs, they can be directly linked to specific taxa of plants and 3) biomarkers could be readily analyzed by usual organic geochemistry procedures on total rock samples.

In addition to be useful to palaeofloristic and palaeoclimatic studies, botanical palaeochemotaxonomy has also been recently applied in other instances. In environmental science, it proved to be helpful in the appreciation of past land-use and recent anthropogenic impacts to soil (e.g. Farella et al., 2001; Heim et al., 2010; Huang et al., 2011; Lavrieux et al., 2011) as well as to trace river pollution caused by paper mills (e.g. Leeming and Nichols, 1998; Wang et al., 2007). In archaeology,

palaeochemotaxonomy is used to trace dietary habits, to understand the use of natural products in craft and funeral rites (e.g. Colombini et al., 2005; Romanus et al., 2008).

Most of palaeochemotaxonomic investigations are derived from published chemotaxonomic data (Otto and Wilde, 2001) and to some parts from the studies of fossil plants found in sedimentary rocks (Otto and Simoneit, 2001; Otto et al., 2005; Dutta et al., 2011). Unfortunately, our knowledge on botanical palaeochemotaxonomy is still very scarce. As pointed by Hautevelle et al. (2006b), difficulties are related to 1) available chemotaxonomic data (generally focused on specific biomolecules or on particular substances, like resins or essential oils); 2) degradation and diagenetic reactions, which may significantly modify the initial molecular fingerprint, making it difficult to perform a direct chemotaxonomic relationship between an extant plant and its fossil counterpart; 3) the scarcity of reference collections of well preserved and identifiable fossil plants containing organic molecules.

In order to fill these gaps, within the 7 extant conifer families (Fig. 1), we investigated the palaeochemotaxonomy of several extant species of the Araucariaceae (Table 1) using an experimental method based on artificial maturation by confined pyrolysis (Hautevelle et al., 2006b). This procedure allows to simulate the conversion of biomolecules into their corresponding diagenetic geomolecules (Stankiewicz et al., 2000; Gupta et al., 2006, 2007). Aims of this study are 1) to determine the common palaeochemotaxonomic (diagenetic) signatures of all extant Araucariaceae species, 2) to evaluate the inter- and intra-generic differences within the family, 3) to highlight the molecular characteristics which should allow their distinction from other conifer families in ancient sediment samples. This contribution should serve as database to future palaeoenvironmental, archaeological and environmental research.

2 Some generalities concerning the Araucariaceae family

Araucariaceae is considered as a southern hemisphere conifer family and was among the first conifer families to be individualised from their "Veltziales" ancestors during Triassic (Stockey, 1994). Diversity and abundance of Araucariaceae are well preserved

up to nowadays despite some reduction of geographic distribution and the extinction of several species during evolution (Axsmith et al., 2004; Kunzmann, 2007a).

2.1 Taxonomy, geographical distribution and living environments

2.1.1 Extant Araucariaceae

Today, Araucariaceae species represent a dominant component of the South Hemisphere forests. The major native Araucariaceae species are restricted to South America, Southwest Asia and the Western Pacific region and cover a large rainfall region extending from subtropics to tropics (Enright and Hill, 1995). Nowadays, more than 40 species are described and consist of three well-defined genera which are (1) *Agathis* (Salisbury, 1807), represented by 21 species, (2) *Araucaria* (Jussieu, 1978) represented by 19 species and (3) *Wollemia* (Jones et al., 1995) represented by the solo species *Wollemia nobilis*. The latter, previously thought to be extinct, was rediscovered in 1994 in Australia, cloistered deep in the New South Wales (Jones et al., 1995).

These phylogenetic relationships of Araucariaceae species were also confirmed by Setoguchi et al. (1998) based on *rbcL* gene sequences:

- the three genera are all monophyletic (i.e. all species descend from a unique ancestor) ;
- *Araucaria* could be intragenerically classified into four sections: *Araucaria*, *Bunya*, *Eutacta*, and *Intermedia*. This sub-classification is in agreement with their morphological characteristics (Stockey, 1982) ;
- phylogenetic relationships within *Agathis* remain somewhat unclear ;
- *Wollemia* is the most basal extant taxon in the family, sharing morphologies with *Araucaria* and *Agathis*.

The geographic distribution of major Araucariaceae species is given in Table 1. Extant *Araucaria* species develop in the subtropical regions and extend into the marginal tropical regions where there is a lower climatic variability, while most of *Agathis* species occupy the tropical islands (Dettmann and Clifford, 2005). These two genera, extending within the equatorial region of New Guinea and Southeast Asia at lower latitude grow under a mesothermal climate limited to the lower montane zone (Kershaw

and Wagstaff, 2001). The monotypic genus *Wollemia* is by far the most endemic genus and its few wild species (about 40 adult plants) live deep in a wilderness rainforest of the Wollemi National Park in New South Wales (Australia).

2.1.2 Fossil Araucariaceae

Appearance and evolution of Araucariaceae are recorded by both macrofossils and pollen in sedimentary series all around the world. The first pollen record is from the Early Triassic of Australia (as ancestor) just after the Permo-Triassic extinction and the unequivocal Araucariaceae representatives were found in the Middle Jurassic North Yorkshire (Stockey, 1982; Kershaw and Wagstaff, 2001; Kunzmann, 2007b). Its maximum worldwide distribution was achieved in both hemispheres during the Cretaceous (Krassilov, 1978; Kunzmann, 2007b). Their distribution was then reduced during the Middle Palaeocene to the southern hemisphere, with an extension to the northern hemisphere (restricted to the Southeast Asia).

During the Jurassic, *Araucaria* (especially the *Eutacta* and the *Bunya* sections) was the first genus to appear and diversify. Its macrofossils and pollens broadened a wide range of habitats in the two hemispheres (Gondwana and Laurasia), where Jurassic climate condition was subtropical to warm-temperate (Kershaw and Wagstaff, 2001). During the Cretaceous, the first effective *Agathis* and *Wollemia* fossils appeared, but remained restricted to Australia and New Zealand (Hill and Bigwood, 1987; Stockey, 1994). This palaeogeographic limitation could have been due to the broken-up of palaeofloristic exchanges between Gondwana and Laurasia. Moreover, appearance of angiosperms during the Cretaceous also forced *Agathis* to be more competitive (Kershaw and Wagstaff, 2001), *Wollemia* probably evolved under similar circumstances. At the end of the Cretaceous, the *Araucaria* genus began to leave the northern hemisphere. As pointed out by Kunzmann (2007b), their retreat from northern hemisphere coincides with two major events in Earth history: 1) major changes in forest composition in Europe and North America; 2) rapid environmental changes across the K-Pg boundary.

Later, during the Cenozoic, *Wollemia* and especially *Agathis* became more prominent from the Early Eocene to the Early Oligocene. This development was synchronous with the diversification of angiosperms. Owing to their adaptation ability to many

unfavourable conditions, such as the ultramafic soils on peridotites and the unstable climate in islands, *Agathis* species began to colonize and diversify in New Caledonia and other Asia-Pacific islands (Jaffré, 1995). In parallel, there was a reduction of *Agathis* from south-eastern Australia due to the cooler conditions at high latitudes. Regarding *Araucaria* species, they regenerated from Early Miocene under subtropical to warm-temperate conditions (Kershaw and Wagstaff, 2001).

2.1.3 Usefulness of Araucariaceae from the past to present days

The worldwide patchy and sparse distribution of Araucariaceae in sedimentary deposits, as inferred by palaeobotanic and palynological data, can thus provide valuable palaeoclimatic information. Indeed, they preferentially grow under warm and wet conditions, mainly in equatorial, tropical or subtropical rainforests as well as peat swamps (Kershaw and Wagstaff, 2001).

Furthermore, many Araucariaceae produce resins having archaeological and economical values. *Agathis* is highly considered as a source of attractive, straight-grained, easily worked timber. The wood of *Agathis australis* has been intensively used in the past for the construction of canoes, houses and cult objects by native Maori people of New Zealand. Some of these objects were well preserved and are of archaeological interest nowadays. The resin of *Agathis* (kauri copal, kauri gum or kauri resin) was also widely used for the manufacture of paints, varnishes and linoleum during the XIX and XX centuries. Other *Agathis* living in Indonesia and Philippines, as *Agathis dammara*, produce resin having an important economic value (Manilla resin or Manilla copal). Copal has also been used for a long time in jewellery and related arts. Furthermore, *Araucaria* may not have an equivalent economical value as *Agathis* but provided wood for shipbuilding during several centuries.

2.2 Chemotaxonomy

Available data on the composition of monoterpenoids (C₁₀), sesquiterpenoids (C₁₅) and diterpenoids (C₂₀) of many Araucariaceae representatives are here summarised.

Monoterpenoids (C₁₀) are abundantly synthesized by Araucariaceae. Compounds, like pinene, thujene, limonene and cubebene are widespread in most of Araucariaceae

species (Brophy et al., 2000). They represent the major components of araucarian essential oils (Brophy et al., 2000; Staniek et al., 2009). Furthermore, owing to their lower molecular weight, these chemicals are generally highly volatile and not readily preserved in sediments.

Sesquiterpenoids are composed by monocyclic (e.g. elemene-, germacrene- and humulene-types), bicyclic (e.g., cadinene-, copaene-, muurolene- and caryophyllene-types) and tricyclic (e.g., aromadendrene-, cubebene- and gurjunene-types) compounds. These chemicals are mainly represented by unsaturated hydrocarbons, alcohols and acids, constituents of essential oil and resins (e.g., Pietsch and König, 2000; Brophy et al., 2000). However, they are common in conifers as well as in other plants like angiosperms and bryophytes (Otto and Wilde, 2001).

Regarding diterpenoids, they are represented by bicyclic, tricyclic and tetracyclic compounds. Bicyclic diterpenoids mainly composed by labdane-type compounds are common in all conifers. They are synthesized as unsaturated hydrocarbons (like sclarenes and biformenes), alcohols (like sclareols) as well as acids (like agathic and communic acids) and isomers may occur (Caputo and Mangoni, 1974; Caputo et al., 1974 a and b, 1976; Thomas, 1969, Pietsch and König, 2000; Otto and Wilde, 2001). Other bicyclic compounds, belonging to the clerodane-type, like clerodadienic acids seem to be restricted to two *Araucaria* species: *A. bidwillii* (Caputo and Mangoni, 1974) and *A. hunsteinii* (Otto and Wilde, 2001).

Tricyclic diterpenoids reported in Araucariaceae are composed essentially by the abietane- and pimarane-types. Tetracyclic diterpenoids belong to the beyerane-, kaurane-, phyllocladane-, trachylobane- and atisane-types (Brophy et al., 2000; Otto and Wilde, 2001). According to Thomas (1969), Brophy et al. (2000) and Pietsch and König (2000), both tri- and tetracyclic diterpenoids are synthesized as unsaturated hydrocarbons (like pimaradienes, abietadienes, kaurenes, etc.), acids (like abietic and isopimaric acids) and more rarely as alcohols (like phyllocladanol, phenolic abietanes which are only reported in few *Agathis* species).

Distribution of bioterpenoids varies from species to another in fresh plants (Thomas, 1969; Brophy et al., 2000). For instance, each species seems to be dominated by one or

eventually two compounds. Their molecular composition could thus be theoretically specific to each araucarian species and/or genus. These data originate from a compilation of several studies and some inconsistencies may occur. For instance, Thomas (1969) insists on the presence of compounds like agathic, communic, isopimaric and abietic acids in *Agathis* species while these compounds are not reported by Brophy et al. (2000). Such kind of contradictions is certainly due to the fact that the former studied the resins while the last studied essential oil. Nevertheless, chemotaxonomic data provide valuable knowledge to palaeochemotaxonomy.

3 Samples and experimental procedures

3.1 Samples

Leaves and twigs of about **12** Araucariaceae species were selected for this study including **3** *Agathis*, **8** *Araucaria* and **1** *Wollemia*. Choice of leaves and twigs as plant material in this study is done based on following reasons. 1) Most of sedimentary organic matter originating from conifers comes from leaves and twigs. 2) Plant biomarkers investigated in this paper derive from terpenoids that are of the major components of essential oils and resins. The terpenoids are transported with resins through vessels to all part of a plant, including leaves and twigs. 3) Analysis of the molecular composition of leaves and twigs provided currently available relevant chemotaxonomic and even some palaeochemotaxonomic data. More details of representatives, their phylogenetic relationships, native geographic distributions as well as their provenances are given in the Table 1. This sampling is believed to be well representative of the intrinsic variability of the Araucariaceae family.

3.2 Experimental and analytical procedures

This study focuses on the diagenetic molecular signature of Araucariaceae. Transformation of fresh material to its diagenetic counterpart was carried out by experimental artificial maturation by confined pyrolysis developed by Hautevelle et al. (2006b). Twigs and leaves of the selected plants were finely cut and dried under vacuum for 24 h in a desiccator at 45°C before being crushed. The plants to be pyrolyzed were introduced into gold tubes with and without LiAlH₄ (*Fluka* No 62420,

purity \geq 97%, powder form). The use of LiAlH_4 favors the generation of saturated over aromatic terpanes during pyrolysis. The filed gold tubes were then sealed under an argon atmosphere and introduced into autoclaves and heated at 280°C and 700 bars during 24 h. These parameters of pyrolysis were determined to obtain the broadest distribution of biomarkers (aliphatic, aromatic and polar). Biomolecules of the starting fresh plant samples of each representative were also studied.

The soluble organic compounds of fresh plants and their pyrolysates were extracted under 80°C and 100 bars using CH_2Cl_2 with an Accelerated Solvent Extractor (ASE 350, Dionex). The total extract was then fractionated into aliphatic, aromatic and polar fractions using liquid chromatography on alumina (separation of hydrocarbon and polar fraction) and silica (separation of aliphatic and aromatic fractions from the hydrocarbon fraction) columns. The fractions were successively eluted with CH_2Cl_2 and $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ (50/50 v/v) on the alumina column, then with pentane, followed by pentane/ CH_2Cl_2 (65/35 v/v) and $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ (50/50 v/v) on the silica column. The obtained fractions were diluted in hexane (4mg/ml for aliphatic fraction and 8mg/ml for the others) before analysis by GC-MS.

3.3 Gas Chromatography-Mass Spectrometry (GC-MS) and identification of compounds

Aliphatic and aromatic fractions were analysed using an HP 5890 Series II gas chromatograph coupled with an HP 5971 mass spectrometer (GC-MS). Polar fractions were silylated using *N,O*-bis-(trimethylsilyl)trifluoroacetamide (BSTFA) at 60°C during 15 min before analysis. The capillary column was a DB-5 J&W (60 m x 0.25 mm i.d., with 0.1 μm film thickness). The temperature programme was 70–315°C at 15°C/min to 130°C, and then 3°C/min followed by an isothermal stage at 315°C for 15 min. Helium was the carrier gas (1 ml/min flow rate). The MS operated in the electron impact mode (EI) at 70 eV ionization energy and mass spectra were scanned from 50 to 500 Da using a quadrupole detector. Data were acquired and processed using the Agilent ChemStation software. Compounds were identified by comparison of mass spectra with literature and library database (Wiley275) or by interpretation of mass

spectrometric fragmentation patterns. Relative abundance of compounds was measured by integration of peak area.

4 Results and discussion

Aliphatic, aromatic and polar fractions were studied for each *Araucariaceae* representative, including the extracts of fresh plants and of their pyrolysates. Since our study mainly focuses on palaeochemotaxonomy, the extracts of pyrolyzed plants were preferentially studied.

4.1 Molecular characteristics of artificially diagenetic *Araucaria* genus

4.1.1 Sesquiterpenoids

Distribution of aliphatic and aromatic sesquiterpenoids of *Araucaria* is respectively showed in Fig. 2 and Fig. 3. Proportion of detected sesquiterpenoids in each representative is given in Table 2 and Table 3, which respectively correspond to compositions of saturated and aromatic compounds obtained from pyrolysis with and without LiAlH_4 . Di- and tri- methyl-naphthalenes are ubiquitous in each representative. *Araucaria* species are also commonly characterized by :

- farnesane ($\text{C}_{15}\text{H}_{32}$) and bisabolane-type compounds. Acyclic farnesane, probably derived from isomers of farnesol (*trans*- and *cis*-) detected in the fresh plants, is identified in all *Araucaria* species. Bisabolane-type compounds, like isomers of saturated bisabolane ($\text{C}_{15}\text{H}_{30}$), *ar*-curcumene ($\text{C}_{15}\text{H}_{22}$) and dihydro-*ar*-curcumene ($\text{C}_{15}\text{H}_{24}$), are observed in the aliphatic and the aromatic fraction respectively. *Ar*-curcumene is often found in trace amounts for all species. Bisabolane-type compounds are generally derived from bisabolenes found in the fresh plants. They represent one of the major class of *Araucaria* sesquiterpenoids (especially in the aliphatic fractions), except for *A. bernieri* and *A. bidwillii* ;

- cadalane-type compounds. They are mainly derived from cadinol and cadinenes which are ubiquitous in vascular plants. In the aliphatic fractions, a variety of saturated cadalane-type compounds ($\text{C}_{15}\text{H}_{28}$) are identified like cadinanes, muurolanes and amorphanes. Their spectra are commonly characterized by a molecular ion at m/z 208, a

low abundance (even absent) of the M^+-15 ion at m/z 193, a relatively high abundance of ion at m/z 165 (corresponding to the loss of a isopropyl unit, M^+-43) and a base peak at m/z 109 (corresponding to a fragmentation of A/B-ring moiety after the loss of precedent units) (Fig. 4a). In the aromatic fraction, the cadalane-type compounds are represented by calamenene and calamene ($C_{15}H_{22}$, both partially aromatized), cadalene ($C_{15}H_{18}$, completely aromatized). One norcadalene [probably the 1-isopropyl-7-methylnaphthalene according to the spectrum from Singh et al., (1994)] is detected in all species (Fig. 4d). Its formation is the result of the demethylation of cadalane-type compounds, and no more precision to its structure was proposed. In addition, cadalane-type compounds show a high abundance in most of species samples and represent also one of major *Araucaria* sesquiterpenoids, except for *A. araucana* and *A. cunninghamii* ;

- eudesmane-type compounds. $4\alpha(H)$ -eudesmane and $4\beta(H)$ -eudesmane are identified according to Wiley275 database and the spectrum published by Alexander et al. (1983) (Fig. 4b and 4c). The relative abundances of these two compounds vary from one specie to another. It is higher in *A. bernieri* and at trace level in *A. heterophylla* as well as in *A. cunninghamii*. Moreover, eudesmanes-type compounds are widely distributed in fossil vascular plants (Alexander et al., 1983). However, they were not reported in the previous studies of Brophy et al., (2000) and Otto and Wilde, (2001), which focus on fresh material ;

- ionene ($C_{13}H_{18}$), with a molecular ion at m/z 174 and a base peak at m/z 159 according to Achari et al., (1973). It was detected for most of *Araucaria* species. It could be derived from β -carotene (like in sporopollenin (which derives from some pollen exines) (Achari et al., 1973; Simoneit, 1986). In our case, the ionene is probably directly derived from ionone that is abundant in the fresh plants ;

- chamazulene ($C_{14}H_{16}$) and other compounds identified as pentamethyl-2,3-dihydroindenes according to their spectra are also remarkable in the aromatic fraction. Chamazulene elutes after cadalene and its spectrum, according to the Wiley275 database, is quite similar to that of norcadalene but with a molecular ion (m/z 184) more intense than its base peak at m/z 169 (Fig. 4e). Due to its carbon skeleton composed of a ring composed of seven carbon atoms and a pentenic ring, it is probably derived from precursor of similar carbon skeleton, like aromadendrenes, alloaromadendrene and

spathulenol, which are widespread in the fresh species (Brophy et al., 2000; Olawore, 2005). Pentamethyl-2,3-dihydroindene, identified according to Wiley275 database, is characterized by a spectrum with a molecular ion at m/z 188 ($C_{14}H_{20}$) and a base peak at m/z 173 (M^+-15) (Fig. 4f). Pentamethyl-2,3-dihydroindene and chamazulene show a higher abundance than other aromatic bicyclic sesquiterpenoids with the exception of cadalene ;

- furthermore, 1,3,4-trimethyl-2-(4-methylpentyl)benzene ($C_{15}H_{24}$) with a molecular ion at m/z 204 and a base peak at m/z 133 was detected in all *Araucaria* species but with a low abundance. According to Ellis et al. (1996), it belongs to the family of isohexyl alkylaromatic hydrocarbons with opened A-ring (Fig. 5a). Its occurrence could be due to the alteration of some bicyclic compounds (like drimane) during the pyrolysis.

In addition, the general trend of *Araucaria* sesquiterpenoids, represented by the relative abundance of major class of sesquiterpenoids in each representative, is given in Table 4.

4.1.2 Diterpenoids

Diterpenoids in the extracts of pyrolysates of *Araucaria* species are characterized by a diversity of bicyclic, tricyclic and tetracyclic compounds. Distributions of saturated and aromatic diterpenoids are respectively shown in Fig. 6 and Fig. 7; relative proportion of each diterpenoid is given in Table 2 and Table 3.

- Bicyclic diterpenoids

Bicyclic diterpenoids are essentially composed by labdane-type compounds ($C_{20}H_{38}$). Several isomers of labdane are identified in *Araucaria* species, although they do not show a high abundance in each representative. As shown in the literature, they are generally derived from labdadiene, phenolic labdanes and communic acids present in the fresh plants (Caputo and Mangoni, 1974; Caputo et al., 1974ab, 1976; Otto and Wilde, 2001). Moreover, according to de Paiva Campello and Fonseca, (1975), the agathic acid could also be the origin of labdanes in the case of *A. angustifolia*.

A peak having a molecular ion at m/z 244 ($C_{18}H_{28}$) is detected in the aromatic fraction in all *Araucaria* species. It shows a significant abundance in most species, but in traces for *A. bidwillii*, *A. laubenfelsii* and *A. nemorosa*. According to its mass spectrum, its

base peak at m/z 159 matches very well to fragment composed of two rings (including an aromatic ring) and two methylene units. This fragment is probably formed via the loss of a branched-chain with 6 carbon atoms (Fig. 8a). This kind of fragmentation seems to be consistent with a monoaromatic labdane. No aromatic labdane has been described in the literature until now.

Another peak heaving a molecular ion at m/z 278 ($C_{20}H_{38}$) and a base peak at m/z 109 is detected with a low intensity in the aliphatic fraction of *A. bidwillii*. Its mass spectrum indicates that it could also be a bicyclic compound and has a similar carbon skeleton to labdane. Indeed, its fragment at m/z 193 matches to a configuration including two rings (A and B) and three methylene units, while its A/B-ring fragment (m/z 109) contains only two methylene units. The "third" methylene unit could thus only be bound at the carbon-9 position (Fig. 8b). This obtained configuration is consistent with a structure similar to that of clerodane. However, its report in extant conifers is limited to a few derivatives of *ent*-clerodadienes (Otto and Wilde, 2001) and is restricted to two *Araucaria* species including *A. bidwillii* (Caputo and Mangoni, 1974; Cox et al., 2007).

2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene are abundant for most species and in traces in *A. launbenfelsii*. These two compounds are identified from spectra published by Ellis et al. (1996) and given in Fig. 5b, 5c. They belong to the family of isohexyl alkylaromatic hydrocarbons just like the 1,3,4-trimethyl-2-(4-methylpentyl)benzene previously described. Indeed, their specific carbon skeleton with an opened A-ring could be due to some specific chemical reactions, like aromatisation, ring-opening processes and rearrangement, during the maturation (Ellis et al., 1996). Furthermore, the co-occurrence of 2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene suggests precursors like tricyclic diterpenoids (Ellis et al., 1996), for which the diagenetic products are structurally quite dissimilar to their precursors.

- Tricyclic diterpenoid

Few aliphatic but a high diversity of aromatic tricyclic diterpenoids are observed. Pimarane-type compound, like isopimarane ($C_{20}H_{36}$), is detected in the aliphatic fraction with a significant abundance for numerous species, like *A. bernieri*, *A. heterophylla*, *A.*

laubenfelsii and *A. nemorosa*. This kind of tricyclic aliphatic diterpenoids is probably derived from isopimara-7,15-dienes, 13-isopimardiene, isopimara-8(9),15-diene as well as 8 β -hydroxyisopimarene and 3 β -isopimadinol, that are detected in some of the fresh plants.

Other tricyclic diterpenoids, like abietane-type compounds, show a higher diversity, especially in the aromatic fraction. Related aromatic compounds are retene, dehydroabietane, 18- and 19-norabieta-8,11,13-trienes, tetrahydroretene, simonellite, bisnor-simonellite, 2- and 9-methylretenes. While the saturated abietane-type compounds like abietanes and fichetellite are detected at trace level.

These abietane-type compounds are the major diterpenoids of *Araucaria*. They are mostly derived from dehydroabietic acid that is observed in the polar fraction of pyrolysates and more generally from abietanoic acid in the fresh plants. As regards the phenolic abietanes including sugiol, hinokiol and ferruginol (Otto and Wilde, 2001; Otto and Simoneit, 2001), which are known as abietane-type compounds precursors for *Araucariaceae* as well, show a low relative abundance in the fresh plants and were mostly degraded during pyrolysis.

A peak having a similar spectrum to those of methylretenes is observed in the aromatic fractions of all *Araucaria* species (Fig. 8c). Compared to Alexander et al., (1995), the spectra of these compounds have the same molecular ion and mass fragments, but are different by the fragment intensities. They may rather be consistent with a pentamethyl phenanthrene.

- Tetracyclic diterpenoids

Tetracyclic diterpenoids, mainly *ent*-beyerane, 16 α (*H*)- and 16 β (*H*)-phyllocladanes and *ent*-16 α (*H*)- and *ent*-16 β (*H*)-kauranes, show relative high abundance in each representative. They are identified according to the spectra published by Noble et al. (1985). Their spectra are all characterized by a molecular ion at *m/z* 274 and a base peak at *m/z* 123 (C₂₀H₃₄). Distribution and abundance of these compounds for each representative are given in Fig. 6 and Tab. 2. They are probably derived mainly from beyerene, *ent*-kaurene, isokaurene and phylloclade-16-ene, as well as kauran-16-ol, phyllocladanol and kaur-16-en-19-ol, observed in the fresh plants. Furthermore,

413 beyerane-type compounds are detected in *A. araucana*, *A. bidwillii* and *A. cunninghamii*
414 in both their fresh plant (beyerene) and the pyrolysates (*ent*-beyerane). Co-occurrence of
415 phyllocladane-type and kaurane-type compounds is observed in most of *Araucaria*
416 species, except of *A. angustifolia* and *A. nemorosa*. Only phyllocladane-type or
417 kaurane-type compounds are observed in *A. angustifolia* and in *A. nemorosa*
418 respectively.

419 A peak with a spectrum characterized by a molecular ion at m/z 260 ($C_{19}H_{32}$) with a
420 base peak at m/z 123 is detected in *A. angustifolia*. This spectrum matches very well to
421 that of 17-nortetracyclic diterpane described by Noble et al., (1986) (Fig. 8d).
422 Formation of this C-19 diterpane is likely due to the demethylation of a C-20 tetracyclic
423 diterpane, while its precise molecular structure remains unclear. Furthermore, in the
424 case of *A. angustifolia*, as only phyllocladanes are observed, the C-19 tetracyclic
425 diterpane could rather be a 17-norphyllocladane.

426 Another peak, characterized by a molecular ion at m/z 288 ($C_{21}H_{32}$) and a base peak at
427 m/z 123 is detected in *A. angustifolia*, *A. bernieri* and *A. laubenfelsii* (Fig. 8e). It is
428 likely a methylated tetracyclic diterpane, because of the positive difference of molecular
429 weight of 14 Da to the well known tetracyclic diterpenoids corresponds to an additional
430 $-CH_2-$ group. This configuration is in correspondence to a C-21 tetracyclic diterpane.
431 Moreover, 1) according to the spectrum, the ion at m/z 231 is relatively more abundant
432 than the ions at m/z 259; 2) *A. angustifolia*, *A. bernieri* and *A. laubenfelsii* all show the
433 presence of phyllocladanes. This C-21 tetracyclic diterpane is probably a 16(*H*)-homo-
434 phyllocladane.

435 It is also interesting to point out that, according to Otto and Wilde (2001), tetracyclic
436 diterpenoids like trachylobane-type and atiserane-type compounds are those exclusive
437 for *Araucaria* genus among all conifers. Unfortunately, the diagenetic products of these
438 two compounds are still unknown to our knowledge. In our study, trachylobane and
439 atiserene/isoatiserene are identified in the fresh *A. araucana* and *A. nemorosa*. Yet, no
440 possible corresponding pyrolysates could be identified.

- Unidentified compounds

Numerous compounds remain unidentified in the extract of *Araucaria* pyrolysates and some of them, their mass spectra are given in the Fig. 5d, 5e and Fig. 9a – h. Tentative identification as proposed in Fig. 9a suggests saturated tricyclic diterpenoid. They have, to our knowledge, never been reported in previous studies on vascular plant biomarkers. Nevertheless, many of them are quite abundant in several species. For instance, it was observed that two peaks having spectra with a molecular ion at m/z 238 ($C_{18}H_{22}$; Fig. 5d) and 254 ($C_{19}H_{26}$; Fig. 5e), a base peak at m/z 168 and 184 respectively co-appear with 2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene. These two unidentified peaks show also respectively similar abundance to 2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene. The relationship of the compounds is not clear; while their co-occurrence and similar fragmentation tend each couple to a similar structure. However, these unidentified compounds may have some relevant palaeochemotaxonomic values and special attention should be paid to these biomarkers in future studies.

Relative abundance of major class of *Araucaria* diterpenoids (including saturated and aromatic compounds) in each representative is summarised in Table 4.

4.2 Molecular characteristics of artificially diagenetic *Agathis* genus

4.2.1 Sesquiterpenoids

Composition of sesquiterpenoids in *Agathis* (*Ag.*) species is shown in Fig. 10 (saturated) and Fig. 11 (aromatic). Proportion of each sesquiterpenoid is given in Table 2 (saturated) and Table 3 (aromatic). Some similarities to the molecular composition of *Araucaria* species are observed. Farnesane, cadalane-type compounds (saturated cadalanes, calamene, calamenene, cadalene and norcadalene), pentamethyl-dihydroindenes and chamazulene show some high relative abundances in each *Agathis* representative. 1,3,4-trimethyl-2-(4-methylpentyl)benzene is detected as well, but at trace level.

Furthermore, bisabolane-type compounds (bisabolanenes and dihydro-*ar*-curcumene), which are widespread in the *Araucaria* genus, are observed only in *Ag. australis* and

show a low abundance in the other *Agathis* species. Concerning the eudesmanes, they were only detected only in *Ag. australis*.

4.2.2 Diterpenoids

As sesquiterpenoids, compositions of diterpenoids of *Agathis* species (Fig. 11) show also several similarities to those of *Araucaria*. Distribution and abundance of major diterpenoids of the *Agathis* genus are given in Table 2 and 3.

Bicyclic diterpenoids are characterized by the presence of labdanes in the aliphatic fraction and monoaromatic labdane in the aromatic fractions, which essentially originate from agathic acid (Thomas, 1969). The tricyclic diterpenoids like isopimarane is observed in *Ag. australis* and *Ag. robusta*. The isohexyl alkylaromatic hydrocarbons [2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene] show high relative abundance in most of *Agathis* species and are found as traces for *Ag. robusta* (Fig. 5b and 5c). Abietane-type compounds show a high abundance in all species. Aromatic abietanes, like retene, dehydroabietane, 18- and 19-norabieta-8,11,13-trienes, tetrahydroretene, 2- and 9- methylretenes, are much more abundant in *Ag. moorei* and *Ag. robusta* than in *Ag. australis*. The methylretenes and norabieta-8,11,13-trienes are detected only at trace level in *Ag. australis*. Saturated abietanes are generally in trace in *Ag. australis* and *Ag. moorei*, but in higher abundance in *Ag. robusta*. As for *Araucaria*, the detected abietane-type compounds are in majority derived from dehydroabietic acid (detected in the pyrolysates) as well as from other abietanoic acids precursors and (cetono)phenolic abietanes (detected in the fresh plants). Only dehydroabietic acid is observed in the pyrolysats. Tetracyclic diterpenoids, the *ent*-beyerane is identified only in *Ag. australis*; 16 α (*H*)-phyllocladane, *ent*-16 α (*H*)- and *ent*-16 β (*H*)- kauranes, are detected in high abundance except for *Ag. robusta*. Despite of the presence and absence, these tetracyclic compounds represent as the major components within the *Agathis* species. Regarding to the unidentified compounds, like those of Fig. 8a to e, are also detected with a significant abundance in *Agathis* species.

Furthermore, some peaks show relatively high abundance and their spectra are respectively characterized by a molecular ion at *m/z* 264 and 274. A base peak at *m/z* 109 and 259 are observed in the aliphatic fraction of *Ag. australis* and *Ag. moorei*. Their chemical formulas are respectively C₂₀H₂₄ and C₂₀H₃₄, and their spectra are given in Fig.

9 (i and j). Since these spectra are neither referenced in literature nor in our spectral database, their molecular structures are not yet clear. However their frequent, significant but also unique presence (Fig. 9i) and predominance (Fig. 9j) in *Agathis* species could be interesting for future biomarker research.

In addition, the relative abundance of major class of sesqui- and diterpenoids in each *Agathis* representative is summarised in Table 4.

4.3 Molecular characteristics of artificially diagenetic of *Wollemia nobilis*

W. nobilis shows several similarities to other Araucariaceae genera (Fig. 2 and 3, Fig. 6 and 7, as well as Fig. 10 and 11). Relative proportion of sesqui- and di-terpenoids of this monotypic genus are given in Table 2 and 3.

For the sesquiterpenoids, *W. nobilis* is characterized by the presence of eudesmane-type [$4\alpha(H)$ - and $4\beta(H)$ - eudesmanes], bisabolane-type (bisabolane, *ar*-curcumene and dihydro-*ar*-curcumene), cadalane-type (saturated cadalanes, norcadalene, calamene, calamenene and cadalene) compounds. Pentamethyl-dihydroindene in *W. nobilis* is not as abundant as those observed in *Araucaria* and *Agathis* genera among the sesquiterpenoids. Chamazulene, which presents a relatively high abundance in both *Araucaria* and *Agathis* genera, is here absent. And do likewise 1,3,4-trimethyl-2-(4-methylpentyl)benzene.

Furthermore, *W. nobilis* shows a wide diversity of diterpenoids. On one hand, as the other Araucariaceae species, it is characterized by the presence of labdane-type compounds (like labdanes), isohexyl alkylaromatic hydrocarbons [like 2,6-dimethyl-1-(4-methylpentyl)naphthalene and 6-ethyl-2-dimethyl-1-(4-methylpentyl)-naphthalene], abietane-type compounds (like retene, dehydroabietane, 18- and 19- norabieta-8,11,13-trienes, tetrahydroretene, 2- and 9- methylretenes as well as dehydroabietic acid), tetracyclic diterpenoids (like *ent*-beyerane, *ent*-16 $\alpha(H)$ - and *ent*-16 $\beta(H)$ - kauranes). Phyllocladane-type compounds, being largely widespread in most of Araucariaceae species, are neither detected in fresh nor in pyrolysates of *W. nobilis*. Moreover, due to the high abundance of tetracyclic diterpenoids, the saturated abietanes are present at trace level in the aliphatic fraction. The unidentified compounds like those cited in paragraph 4.1.2. (Fig. 9) show a significant abundance. The diversity of diterpenoids of

W. nobilis is, on the other hand, demonstrated by a great variety of other unknown aromatic diterpenoids. The particularities of these compounds are that they are of similar abundance with no predominance of any specific compounds (Fig. 12).

Relative abundances of major classes of sesqui- and diterpenoids in *Wollemia nobilis* are summarised in Table 4.

5 Conclusion

Palaeochemotaxonomy of the Araucariaceae family is evaluated using artificial maturation by confined pyrolysis of extant species. The Araucariaceae family is characterized by the remarkable predominance of saturated tetracyclic diterpenoids, including beyerane, phyllocladanes as well as kauranes. This family is also characterised by a relatively high abundance of sesquiterpenoids (like cadalanes, bisabolane, eudesmane as well as chamazulene) and diterpenoids like abietanes. The palaeochemotaxonomy of the Araucariaceae could be summarized as follows :

- the *Araucaria* genus is mainly characterized by the high abundance of sesquiterpenoids, like eudesmanes, bisabolanes, chamazulene, pentamethyl-dihydroindenes and more particularly by cadalane-type compounds. For the diterpenoids, saturated tetracyclic compounds are largely predominant. Compared to the tetracyclic diterpenoids, other diterpenoids, like labdane, isopimarane, abietanes as well as isohexyl alkylaromatic hydrocarbons, show a relatively lower abundance. Moreover, isopimarane is only detected in the *Eutecta* section species.

- the *Agathis* genus is characterized by the high abundance of sesquiterpenoids, like cadalane-type compounds, chamazulene and pentamethyl-dihydroindenes. Compared to the *Araucaria* genus, the bisabolanes and eudesmanes are occasionally present in only some species. Concerning the diterpenoids, most of species are highly similar to those of *Araucaria* species, which are characterised by a high abundance of tetracyclic diterpanes and isohexyl alkylaromatic hydrocarbons, but a low abundance of saturated abietanes. This is the case for *Ag. australis* and *Ag. Moorei*, but not for *Ag. robusta*. The latter is characterised by the absence of tetracyclic diterpane and a relatively lower abundance of isohexyl alkylaromatic hydrocarbons. The saturated abietanes represent

thus a significant fraction of the aliphatics. Moreover, the aromatic abietanes show constantly a high relative abundance in all *Agathis* representatives.

- *Wollemia nobilis*, presents high similarities to the *Araucaria* genus. Presence of cadalane-types, bisabolanes, eudesmanes, penatamethyl-dihydroindenes and also the absence of chamazulene characterize the sesquiterpenoids signature of *Wollemia nobilis*. The tetracyclic compounds, like *ent*-beyerane and *ent*-kauranes, are predominant among the diterpenoids. Other recurrent diterpenoids in the *Araucaria* and *Agathis* genera, like labdane and isopimarane, are at trace amount or absent. The recurrent abietane-type compounds are identified. The aromatic abietanes do not seem as abundant as those observed in the two other genera. Moreover, there is a great variety of unidentified aromatic diterpenoids and are of same relative abundance as much as the other well known biomarkers, like the aromatic abietane-type compounds and the isohexyl alkylaromatic hydrocarbons.

Acknowledgements

We thank Alain Rouiller, Frédéric Lannuzel, Bastien Farre for technical assistance. We thank Frederic Thévenard (UMR 5276 LGLTPE, University Claude Bernard, Lyon I), Anne Allaria (Botanical Garden of the Villa Thuret, INRA, Antibes), Denis Larpin (Museum Nationale d'Histoire Naturelle, Paris), Romaric Pierrel and Marc Rémy (Botanical Garden of Montet, Vandoeuvre-lès-Nancy) for their help for providing samples. We also thank the Wollemi Pine France (<http://france.wollemipine.com>) and the Kientzler society for providing branches of the Wollemi Pine in 2006 during the propagation program of this species in several botanical gardens around the world. We also wish to thank the handling editor of Biogeoscience (Mr. X. Wang), Dr Sawada and a second anonymous reviewer for their appreciation, as well as for their numerous helpful suggestions, which improved the quality of our manuscript. This research benefited from the support of the Ministry of Higher Education and Research of France (MESR) through a PhD grant to Yueming LU.

References

- Achari, R. G., Shaw, G., and Holleyhead, R.: Identification of ionene and other carotenoid degradation products from the pyrolysis of sporopollenins derived from some pollen exines, a spore coal and the Green River shale, *Chemical Geology*, 12, 229-234, 1973.
- Alexander, R., Bastow, T. P., Fisher, S. J., and Kagi, R. I.: Geosynthesis of organic compounds: II. Methylation of phenanthrene and alkylphenanthrenes, *Geochimica et Cosmochimica Acta*, 59, 4259-4266, 1995.
- Alexander, R., Kagi, R., and Noble, R.: Identification of the bicyclic sesquiterpenes drimane and eudesmane in petroleum, *Journal of the Chemical Society, Chemical Communications*, 5, 226-228, 1983.
- Aplin, T.T., Cambie, R.C., and Rutledge, P.S.: The taxonomic distribution of some diterpene hydrocarbons. *Phytochemistry* 2, 205–214, 1963.
- Axsmith, B. J., Krings, M., and Waselkov, K.: Conifer pollen cones from the Cretaceous of Arkansas: implications for diversity and reproduction in the Cheirolepidiaceae, *Journal of Paleontology*, 78, 402-409, 2004.
- Brophy, J. J., Goldsack, R. J., Wu, M. Z., Fookes, C. J. R., and Forster, P. I.: The steam volatile oil of *Wollemia nobilis* and its comparison with other members of the Araucariaceae (*Agathis* and *Araucaria*), *Biochemical Systematics and Ecology*, 28, 563-578, 2000.
- Caputo, R., and Mangoni, L.: Diterpenes from *Araucaria bidwilli*, *Phytochemistry*, 13, 467-470, 1974.
- Caputo, R., Dovinola, V., and Mangoni, L.: New diterpenes from *Araucaria cunninghami*, *Phytochemistry*, 13, 475-478, 1974a.
- Caputo, R., Dovinola, V., and Mangoni, L.: New labdane diterpenes from *Araucaria cooki*. *Phytochemistry* 13, 471–474, 1974b.

- 27 Caputo, R., Mangoni, L., Monaco, P., Pelosi, L., and Previtera, L.: Neutral diterpenes
28 from *Araucaria bidwilli*, *Phytochemistry*, 15, 1401-1402, 1976.
- 29 Castro, M. A., Gordaliza, M., Miguel Del Corral, J. M., and Feliciano, A. S.: The
30 distribution of lignanoids in the order coniferae, *Phytochemistry*, 41, 995-1011, 1996.
- 31 Colombini, M. P., Giachi, G., Modugno, F., and Ribechini, E.: Characterisation of
32 organic residues in pottery vessels of the Roman age from Antinoe (Egypt), *Microchem*
33 *J.*, 79, 83-90, 2005.
- 34 Cox, R. E., Yamamoto, S., Otto, A., and Simoneit, B. R. T.: Oxygenated di- and
35 tricyclic diterpenoids of southern hemisphere conifers, *Biochemical Systematics and*
36 *Ecology*, 35, 342-362, 2007.
- 37 de Paiva Campello, J., and Fonseca, S. F.: Diterpenes from *Araucaria angustifolia*,
38 *Phytochemistry*, 14, 2299-2300, 1975.
- 39 Dettmann, M. E., and Clifford, H. T.: Biogeography of Araucariaceae, Australia and
40 New Zealand Forest Histories: N°2 Araucarian Forests. (Ed. John Dargavel), 2, 2005.
- 41 Dutta, S., Mallick, M., Kumar, K., Mann, U., and Greenwood, P. F.: Terpenoid
42 composition and botanical affinity of Cretaceous resins from India and Myanmar,
43 *International Journal of Coal Geology*, 85, 49-55, 2011.
- 44 Ellis, L., Singh, R. K., Alexander, R., and Kagi, R. I.: Formation of isohexyl
45 alkylaromatic hydrocarbons from aromatization-rearrangement of terpenoids in the
46 sedimentary environment: A new class of biomarker, *Geochimica et Cosmochimica*
47 *Acta*, 60, 4747-4763, 1996.
- 48 Enright, N. J., Ogden, J., and Rigg, L. S.: Dynamics of forests with Araucariaceae in the
49 western Pacific, *Journal of Vegetation Science*, 10, 793-804, 1999.
- 50 Farella, N., Lucotte, M., Louchouart, P., and Roulet, M.: Deforestation modifying
51 terrestrial organic transport in the Rio Tapajos, Brazilian Amazon, *Organic*
52 *Geochemistry*, 32, 1443-1458, 2001.

- 53 Gupta, N. S., Michels, R., Briggs, D. E. G., Evershed, R. P., and Pancost, R. D.: The
54 organic preservation of fossil arthropods: an experimental study, *Proceedings of the*
55 *Royal Society B: Biological Sciences*, 273, 2777-2783, 2006.
- 56 Gupta, N. S., Michels, R., Briggs, D. E. G., Collinson, M. E., Evershed, R. P., and
57 Pancost, R. D.: Experimental evidence for the formation of geomacromolecules from
58 plant leaf lipids, *Organic Geochemistry*, 38, 28-36, 2007.
- 59 Hautevelle, Y., Michels, R., Malartre, F., and Trouiller, A.: Vascular plant biomarkers
60 as proxies for palaeoflora and palaeoclimatic changes at the Dogger/Malm transition of
61 the Paris Basin (France), *Organic Geochemistry*, 37, 610-625, 2006a.
- 62 Hautevelle, Y., Michels, R., Lannuzel, F., Malartre, F., and Trouiller, A.: Confined
63 pyrolysis of extant land plants: A contribution to palaeochemotaxonomy, *Organic*
64 *Geochemistry*, 37, 1546-1561, 2006b.
- 65 Heim, A., Hofmann, A., and Schmidt, M. W. I.: Forest-derived lignin biomarkers in an
66 Australian oxisol decrease substantially after 90 years of pasture, *Organic*
67 *Geochemistry*, 41, 1219-1224, 2010.
- 68 Hill, R.S., and Bigwood, A.J.: Tertiary gymnosperms from Tasmania: Araucariaceae.
69 *Alcheringa* 11, 325–335, 1987.
- 70 Huang, Z. Q., Davis, M. R., Condon, L. M., and Clinton, P. W.: Soil carbon pools,
71 plant biomarkers and mean carbon residence time after afforestation of grassland with
72 three tree species, *Soil Biol. Biochem.*, 43, 1341-1349, 2011.
- 73 Jaffré, T.: Distribution and ecology of the conifers of New Caledonia, in: *Ecology of the*
74 *Southern Conifers*, N. J. Enright and R. S. Hill ed., Cambridge University Press,
75 Cambridge, 171–196, 1995.
- 76 Jones, W. G., Hill, K. D., and Allen, J. M.: *Wollemia nobilis*, a new living Australian
77 genus and species in the Araucariaceae, *Telopea* 6, 173–176, 1995.
- 78 Jussieu, A. L. d.: *Genera plantarum secundum ordines naturales disposita, iuxta*
79 *methodum in Horto regio Parisiensi exaratam anno 1774*, L'Herissant, Paris, 1778.

- 80 Kershaw, P.: The southern conifer family Araucariaceae: History, status, and value for
81 paleoenvironmental reconstruction, *Annual Review of Ecology and Systematics*, 32,
82 397-414, 2001.
- 83 Krassilov, V.A.: Araucariaceae as indicators of climate and paleolatitudes. Review of
84 Palaeobotany and Palynology, 26, 113-124, 1978.
- 85 Kunzmann, L.: Neue Untersuchungen zu *Araucaria* Jussieu aus der europäischen
86 Kreide, *Palaeontographica Abteilung B: Palaeophytologie* 276, 97-131, 2007a.
- 87 Kunzmann, L.: Araucariaceae (Pinopsida): Aspects in palaeobiogeography and
88 palaeobiodiversity in the Mesozoic, *Zoologischer Anzeiger - A Journal of Comparative*
89 *Zoology*, 246, 257-277, 2007b.
- 90 Lavrieux, M., Jacob, J., LeMilbeau, C., Zocatelli, R., Masuda, K., Bréheret, J.-G., and
91 Disnar, J.-R.: Occurrence of triterpenyl acetates in soil and their potential as
92 chemotaxonomical markers of Asteraceae, *Organic Geochemistry*, 42, 1315-1323,
93 2011.
- 94 Leeming, R., and Nichols, P. D.: Determination of the sources and distribution of
95 sewage and pulp-fibre-derived pollution in the Derwent Estuary, Tasmania, using sterol
96 biomarkers, *Mar. Freshw. Res.*, 49, 7-17, 1998.
- 97 Mongrand, S., Badoc, A., Patouille, B., Lacomblez, C., Chavent, M., Cassagne, C., and
98 Bessoule, J.-J.: Taxonomy of gymnospermae : multivariate analyses of leaf fatty acid
99 composition. *Phytochemistry*, 58, 101–115, 2001.
- 100 Noble, R. A., Alexander, R., Kagi, R. I., and Knox, J.: Tetracyclic diterpenoid
101 hydrocarbons in some Australian coals, sediments and crude oils, *Geochimica et*
102 *Cosmochimica Acta*, 49, 2141-2147, 1985.
- 103 Noble, R. A., Alexander, R., Kagi, R. I., and Nox, J. K.: Identification of some
104 diterpenoid hydrocarbons in petroleum, *Organic Geochemistry*, 10, 825-829, 1986.
- 105 Olawore, N. O.: Analysis of the leaf oil of *Araucaria cunninghamii* sweet. Grown in
106 Nigeria, *The Journal of essential oil research*, 17, 459-461, 2005.

- 107 Otto, A., and Simoneit, B. R. T.: Chemosystematics and diagenesis of terpenoids in
108 fossil conifer species and sediment from the Eocene Zeitz formation, Saxony, Germany,
109 *Geochimica et Cosmochimica Acta*, 65, 3505-3527, 2001.
- 110 Otto, A., and Wilde, V.: Sesqui-, di-, and triterpenoids as chemosystematic markers in
111 extant conifers—A review, *The Botanical Review*, 67, 141-238, 2001.
- 112 Otto, A., Simoneit, B. R. T., and Rember, W. C.: Conifer and angiosperm biomarkers in
113 clay sediments and fossil plants from the Miocene Clarkia Formation, Idaho, USA,
114 *Organic Geochemistry*, 36, 907-922, 2005.
- 115 Pietsch, M., and König, W. A.: Enantiomers of sesquiterpene and diterpene
116 hydrocarbons in *Araucaria* species, *Phytochemical Analysis*, 11, 99-105, 2000.
- 117 Romanus, K., Van Neer, W., Marinova, E., Verbeke, K., Luypaerts, A., Accardo, S.,
118 Hermans, I., Jacobs, P., De Vos, D., and Waelkens, M.: Brassicaceae seed oil identified
119 as illuminant in Nilotic shells from a first millennium AD Coptic church in Bawit,
120 Egypt, *Analytical and Bioanalytical Chemistry*, 390, 783-793, 2008.
- 121 Salisbury, R. A.: XIV. The Characters of several Genera in the Natural Order of
122 Coniferæ: with Remarks on their Stigmata, and Cotyledons, *Transactions of the Linnean*
123 *Society of London*, 8, 308-317, 1807.
- 124 Setoguchi, H., Asakawa Osawa, T., Pintaud, J.-C., Jaffre, T., and Veillon, J.-M.:
125 Phylogenetic relationships within Araucariaceae based on rbcL gene sequences, *Am. J.*
126 *Bot.*, 85, 1507-1516, 1998.
- 127 Simoneit, B. R. T., Grimalt, J. O., Wang, T. G., Cox, R. E., Hatcher, P. G., and
128 Nissenbaum, A.: Cyclic terpenoids of contemporary resinous plant detritus and of fossil
129 woods, ambers and coals, *Organic Geochemistry*, 10, 877-889, 1986.
- 130 Singh, R. K., Alexander, R., and Kagi, R. I.: Identification and occurrence of
131 norcadalenes and related compounds in crude oils and sediments, *Organic*
132 *Geochemistry*, 21, 249-256, 1994.
- 133 Smith, P.: *The chemotaxonomy of plants*. Arnold, London, 1976.

- 134 Staniek, A., Muntendam, R., Woerdenbag, H. J., and Kayser, O.: Essential oil
135 constituents derived from different organs of a relictual conifer *Wollemia nobilis*,
136 *Biochemical Systematics and Ecology*, 38, 131-135, 2010.
- 137 Stockey, R. A.: The Araucariaceae: An evolutionary perspective, Review of
138 *Palaeobotany and Palynology*, 37, 133-154, 1982.
- 139 Stockey, R.: Mesozoic Araucariaceae: Morphology and systematic relationships,
140 *Journal of Plant Research*, 107, 493-502, 10.1007/bf02344070, 1994.
- 141 Thomas, B. R.: Kauri resins-modern and fossil. In: Eglinton, G., Murphy, M.T.J. (Eds.),
142 *Organic Geochemistry – Methods and Results*, Springer, Berlin, 599–618, 1969.
- 143 van Aarssen, B. G. K., Alexander, R., and Kagi, R. I.: Higher plant biomarkers reflect
144 palaeovegetation changes during Jurassic times, *Geochimica et Cosmochimica Acta*, 64,
145 1417-1424, 2000.
- 146 Vliex, M., Hagemann, H. W., and Püttmann, W.: Aromatized arborane/fernane
147 hydrocarbons as molecular indicators of floral changes in Upper Carboniferous/Lower
148 Permian strata of the Saar-Nahe Basin, southwestern Germany, *Geochimica et*
149 *Cosmochimica Acta*, 58, 4689-4702, 1994.
- 150 Wang, L., Lee, F. S. C., Wang, X. R., Yin, Y. F., and Li, J. R.: Chemical characteristics
151 and source implications of petroleum hydrocarbon contaminants in the sediments near
152 major drainage outfalls along the coastal of Laizhou Bay, Bohai Sea, China, *Environ.*
153 *Monit. Assess.*, 125, 229-237, 10.1007/s10661-006-9514-0, 2007.

1 **Table 1:** Phylogentic relationships, geographical range and provenance of Araucariaceae species selected for this study.

Genera	Section	Species	Geographical range	Samples provenance
Agathis		australis (D. Don) Loudon 1829 Syn. : <i>Dammara australis</i> , <i>Podocarpus zamiifolius</i> Com. : kauri, New Zealand kauri	New Zealand (Northland & Coromandel peninsulas)	Muséum National d'Histoire Naturelle (Paris, France)
		moorei (Lindl.) Mast. 1892 Syn. : <i>Dammara moorei</i> , <i>Dammara lanceolata</i> Com. : Moore kauri	New Caledonia	Botanical Garden of Lyon (France)
		robusta (C. Moore ex F. Muell.) Bailey 1883 Com. : Queensland kauri, smooth-bark kauri, kauri pine	Australia (Queensland)	Botanical Garden of Lyon (France)
Araucaria	Araucaria	angustifolia (Bertol.) Kuntze 1898 Syn. : <i>A. brasiliana</i> , <i>Columbea angustifolia</i> Com. : Parana pine, candelabra tree	South America (Brazil, North Argentina & Paraguay)	Botanical Garden of Lyon (France)
		araucana (Molina) K. Koch 1873 Syn. : <i>A. imbricata</i> Com. : monkeypuzzle tree	South America (S. Chile & SW. Argentina)	Botanical Garden of Montet (Nancy, France)
	Bunya	bidwillii Hooker 1843 Com. : bunya, bunya pine	Australia (Queensland)	Botanical Garden of the Villa Thuret (France)
	Araucaria	bernieri Buchholz 1949 Com. : Bernier's araucaria	New Caledonia	Botanical Garden of Lyon (France)
		cunninghamii Aiton ex Lambert 1837 Com. : Hoop pine, Moreton Bay pine	Australia (Queensland & North of New South Wales)	Botanical Garden of Montet (Nancy, France)
		heterophylla (Salisb.) Franco 1952 Syn. : <i>A. excelsa</i> , <i>Dombeya excelsa</i> , <i>Eutassa heterophylla</i> Com. : Norfolk Island pine	Norfolk Island	found in the trade
	Eutecta	laubenfelsii Corbasson 1968 Com. : De Laubenfels araucaria	South of New Caledonia	Botanical Garden of Lyon (France)
		nemorosa de Laubenfels 1969 Com. : Boise araucaria	South of New Caledonia	Botanical Garden of Lyon (France)
	Indermedia \		New Guinea	\
	Wollemia	nobilis Johns, Hill & Allen 1994 Com. : Wollemi pine	Australia (Wollemi National Park)	Wollemi Pine France

2 **Table 2:** Relative abundances (%; GC-MS peak integration normalized to the highest peak) of saturated sesqui- and di- terpenoids in
3 Araucariaceae species obtained from the pyrolysis with LiAlH₄.

Compounds	MW	Araucaria(%)								Agathis (%)			Wollemia (%)
		Angustifolia	Araucana	bernieri	bidwillii	cunninghamii	heterophylla	laubenfelsii	nemorosa	australis	moorei	robusta	nobilis
Sesquiterpanes													
Farnesane	212	3,5	6,8	-	-	1,0	8,1	0,6	0,2	1,9	10,1	3,3	0,7
Eudesmane													
4β(H)-edesmane	208	1,5	-	-	-	-	-	-	-	-	-	-	1,0
4α(H)-eudesmane	208	2,0	1,0	1,2	0,6	-	0,6	0,1	0,2	4,0	1,8	-	1,8
Bisabolane													
bisabolane N°1	210	3,8	6,6	-	0,3	2,0	3,0	0,4	0,2	1,1	-	-	1,0
bisabolane N°2	210	1,3	3,6	-	-	-	0,8	0,3	0,1	1,1	-	-	0,6
Cadalanes													
109(208)N°1	208	1,3	4,0	0,1	-	-	0,9	-	-	1,6	-	4,0	0,9
109(208)N°2	208	1,8	0,8	0,2	-	-	-	0,1	-	3,5	39,7	16,2	1,3
109(208)N°3	208	2,8	-	0,9	1,5	-	1,1	0,2	0,1	2,4	100,0	5,8	1,9
109(208)N°4	208	4,8	1,4	0,4	0,3	6,0	0,9	0,2	0,3	6,5	-	15,8	3,1
109(208)N°5	208	3,2	-	0,5	0,5	2,0	1,4	0,1	-	10,1	-	-	2,7
109(208)N°6	208	3,0	-	0,2	0,9	-	1,4	0,7	-	-	-	-	1,9
Diterpanes													
Clerodane (Fig. 6 b)	278	-	-	1,7	-	-	-	-	-	-	-	-	-
Labdane-type													
Labdane 1	278	4,6	4,8	7,3	9,6	86,4	7,5	0,0	14,6	6,8	7,7	66,6	-
Labdane 2	278	4,8	8,6	5,0	11,5	51,7	9,4	0,0	11,2	5,8	5,3	47,6	-
Labdane 3	278	3,8	-	4,1	-	45,7	-	0,4	2,5	-	-	-	2,5
Labdane 4	278	-	-	8,1	-	-	-	0,3	2,2	-	-	-	2,5
C19H36 (Fig. 8. i)	264	-	-	-	-	-	-	-	-	41,8	53,3	-	-
Abietane-type													
Fichtellite	262	-	-	-	-	-	-	-	-	-	-	100	-
Abietane	276	-	-	-	-	-	-	-	-	-	-	32,2	-
Ralated abietane-type													
C20H36 (Fig. 8. a)	276	-	-	-	-	50,0	-	-	100	-	-	-	-
C20H34 (Fig. 8. j)	274	-	-	-	-	-	-	-	-	10,8	6,0	-	-
Pimarane-type													
Isopimarane	276	-	-	6,0	-	-	2,2	8,1	0,8	13,0	-	28,1	-
Ent-beyerane	274	-	100	2,6	100	100	-	-	-	54,9	-	-	8,8
16β(H)-phyllocladanes	274	17,6	-	22,9	1,5	59,0	19,2	39,6	-	57,4	13,5	-	32,2
16α(H)-	274	100	4,5	41,0	2,4	18,0	100	100	-	100	49,2	-	100
17-norphylocladane (Fig. 6 d)	260	1,6	-	-	-	-	-	-	-	-	-	-	-
16(H)-homophyllocladane (Fig. 6 e)	288	2,6	-	1,9	-	-	-	1,2	-	-	-	-	1,4
Ent-16β(H)-kauranes	274	-	1,7	100	-	-	40,7	23,7	-	12,1	5,4	-	1,6
Ent-16α(H)	274	-	5,6	8,9	-	56,0	8,2	19,2	2,3	-	-	-	0,6
Tetracyclic diterpane (Fig. 8. j)	274	1,2	-	-	5,0	-	-	7,2	3,3	43,9	9,0	-	3,8

4 **Table 3:** Relative abundances (% , GC-MS peak integration normalized to the highest peak) of aromatic sesqui- and di- terpenoids in
5 Araucariaceae species obtained from the pyrolysis without LiAlH₄.

Compounds	MW	Araucaria (%)								Agathis (%)			Wollemia(%)	
		Angustifolia	Araucana	bernieri	bidwillii	cunninghamii	heterophylla	laubenfelsii	nemorosa	australis	moorei	robusta	nobilis	
Sesquiterpenoids														
Ionene	156	7,8	13,3	4,7	-	-	43,5	9,3	-	4,1	-	-	2,7	
<i>n</i> -methyl naphthalene														
	DMN	156	39,3	22,3	33,7	-	-	40,7	16,4	1,5	20,9	13,0	10,4	27,4
	TMN	170	100	100	56,8	33,6	66,3	100	24,3	43,1	3,2	39,9	100	50
Tetramethyl-octahydro-naphthalene	174	-	26,3	11,3	6,2	5,8	51,3	100	69,2	62,4	8,2	-	-	
Bisabolane-type														
Dihydro- <i>ar</i> -curcumene	204	19,7	27,0	9,3	1,9	3,8	33,9	19,6	1,8	2,3	2,7	3,1	-	
<i>ar</i> -curcumene	202	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr	
Isohexyl alkylaromatic hydrocarbons														
1,3,4-trimetyl-2-(4-methylpentyl)benzene, (Fig.4 a)	204	11,0	9,2	2,0	4,3	2,1	16,2	11,7	6,9	10,4	3,8	2,3	1,7	
Cadalane-type														
Calamenene	202	17,0	8,3	9,0	5,7	4,1	13,9	15,3	4,4	2,2	15,3	4,0	14,1	
Norcadalene	184	9,4	9,4	5,4	5,6	4,4	8,6	7,7	4,0	2,9	4,0	2,9	3,4	
Calamene	202	9,3	4,4	13,7	5,1	1,3	6,2	8,2	5,0	27,1	16,5	3,3	18,5	
Cadalene	198	51,4	14,1	100	21,9	12,1	26,8	17,8	34,8	5,7	100	20,8	100	
Pentamethyl-didydroindene 1	188	11,5	55,6	9,3	4,2	100	30,2	7,5	24,0	31,8	22,3	2,8	2,6	
Pentamethyl-dihydroindene 2	188	24,2	15,7	5,4	48,1	10,0	25,9	18,2	14,6	100	8,3	12,9	7,4	
Chamazulene	184	40,4	19,6	20,7	7,8	31,2	30,3	10,7	24,6	19,1	22,7	40,1	14,6	
Diterpenoids														
Labdane-type														
Monoaromatic labdane (Fig. 6. a)	244	39,9	28,0	-	6,6	8,6	51,7	-	33,8	39,2	13,1	-	7,7	
Abietane-type														
Bis-simonellite	224	-	-	-	-	-	-	6,3	16,5	-	-	-	19,1	
19-Norabietatriene	256	28,0	-	21,1	10,7	7,3	27,5	6,6	-	-	18,7	-	21,5	
18-Norabietatriene	256	20,1	-	19,9	39,7	7,1	33,2	4,9	-	-	13,8	3,3	12,7	
Dehydroabietane	270	17,9	-	20,3	18,2	11,9	26,0	41,4	82,5	46,2	-	-	-	
1-methyl-10,18-bisnorabieta-8,11,13-triene	256	15,7	15,7	-	28,9	6,3	18,8	-	43,9	11,8	-	-	18,3	
1,2,3,4-tetrahydroretene	238	23,1	4,7	10,7	49,1	11,6	31,7	13,5	29,6	-	25,4	15,3	26,9	
Simonellite	252	-	-	14,3	9,1	-	-	4,3	83,8	15,9	-	-	-	
Retene	234	8,3	-	22,3	100	19,8	38,0	10,2	18,4	4,2	31,9	43,2	33,3	
9-metyl-retene	248	8,6	-	3,0	19,4	2,6	3,7	1,2	-	-	3,5	5,3	5,7	
2-metyl-retene	248	3,5	-	1,7	4,3	2,2	-	-	6,0	-	1,8	4,1	-	

(continued on next page)

Compounds	MW	Araucaria (%)								Agathis (%)			Wollemia(%)
		Angustifolia	Araucana	bernieri	bidwillii	cunninghamii	heterophylla	laubenfelsii	nemorosa	australis	moorei	robusta	nobilis
Diterpenoids													
Related abietane-type													
C18H26 (Philp, 1985, N°94)	242	30,0	24,4	-	18,4	9,0	49,5	5,7	15,6	25,5	-	-	11,4
C19H28	256	-	-	-	-	-	-	34,7	100	36,4	-	-	-
C19H30 (Fig. 8. b)	258	29,8	18,8	-	38,4	11,4	25,2	-	13,0	26,3	-	-	13,8
C19H26 (Fig. 8. h)	254	-	-	18,7	48,3	5,7	-	-	-	-	12,8	-	-
C18H22 (Fig. 8.c)	238	23,1	10,9	16,3	23,3	12,5	31,3	-	21,0	12,7	24,1	-	-
C18H22 (Fig. 8.d)	238	49,0	-	8,4	37,3	9,5	13,9	6,4	-	15,8	12,8	9,7	-
C19H24 (Fig. 8.e)	252	32,4	11,6	14,8	26,5	12,4	22,2	-	20,0	6,2	8,5	4,4	20,7
C19H24 (Fig. 8.f)	252	20,1	-	9,0	11,5	11,9	12,0	-	12,1	6,2	7,4	-	6,5
C19H24 (Fig. 8.g)	252	14,1	-	5,5	24,1	8,6	9,7	2,3	-	-	16,0	9,2	17,5
Pentamethyl phenanthrene (Fig. 6. c)	248	8,1	-	2,3	7,3	8,8	9,6	5,4	-	-	6,3	17,2	15,3
Isohexyl alkylaromatic hydrocarbons													
2,6-dimetyl-1-(4-methylpentyl)naphthalene (Fig.4 b)	240	57,3	38,3	16,7	21,9	15,8	66,6	-	50,2	43,8	25,5	3,2	16,9
6-ethyl-2-dimetyl-1-(4-methylpentyl)naphthalene (Fig.4 c)	254	12,1	17,6	26,3	33,7	22,8	41,0	-	53,1	8,1	26,9	5,0	19,5
C18H22 (Fig.4 d)	238	47,9	28,4	24,2	46,8	22,3	65,9	-	43,6	41,5	19,2	5,5	6,9
C19H26 (Fig.4 e)	254	2,3	5,5	11,3	18,5	10,8	12,1	-	12,5	7,4	-	2,5	7,8

6

7

8 **Table 4:** Major sesqui- and di- terpenoids classes detected in Araucariaceae species after confined pyrolysis with (saturated compounds)
9 and without (aromatic compounds) LiAlH₄ as reducing agent.

Class of structure		<i>Agathis</i>			<i>Araucaria</i>								<i>Wollemia</i>
		<i>australis</i>	<i>moorei</i>	<i>robusta</i>	<i>angustifolia</i>	<i>araucana</i>	<i>bidwillii</i>	<i>bernieri</i>	<i>cunninghamia</i>	<i>heterophylla</i>	<i>laubenfelsii</i>	<i>nemorosa</i>	<i>nobilis</i>
Sesquiterpenoids	Farnesane-type	+++	++	+	+++	+++	+	++	+	+++	+++	++	++
	Bisabolane-type	+	tr	tr	+++	++++	+	+	tr	++	++	+	+
	Eudesmane-type	+++	tr	tr	++	+	++	+	tr	tr	+	+	++
	Cadalane-type	++++	++++	++	++++	++	++++	++++	++	+++	+++	+++	++++
	Ionene	+	tr	tr	+	+	tr	+	tr	++	tr	tr	tr
	Chamazulene	++	++	++	+++	++	++	+++	++	++	++	+++	/
	Pentametyl-dihydroindene	+	+	+	++	+	+	+	tr	++	++	++	+
Diterpenoids	Labdane-type	+	+	++	++	++	+++	+++	++	++	+	+++	+
	Isohexyl alkylaromatic hydrocarbons	+++	+++	+	++++	++++	++	+++	+++	++++	/	++	++
	Abietane-type												
	normal abietane (dehydroabietic acid)	+	+	+	+	+	+	+	+	+	+	+	+
	phenolic abietane (ferruginol, sugiol and hinokiol)	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr	tr
	diagenetic abietanes	++	+++	+++	+++	tr	++++	++++	+++	+++	+++	+++	++++
	Pimarane-type	+	/	/	/	/	/	+	/	+	++	+	/
	Ent-beyerane	+++	/	/	/	+++	+++	+	++	/	/	/	+
	Phyllocladane-type	+	+	/	++++	++	tr	++++	++++	+++	+++	/	/
	Ent-kaurane	++++	+++	/	/	++	/	++++	++	++++	++++	+	++++
	?C ₁₉ H ₂₄ (252Da)	+	++	+	++	++	+	++	++	++	+	++	++
	?C ₁₈ H ₂₂ (252Da)	tr	+	+	+	+	++	++	++	+	+	++	++

1 **Figure 1.** Phylogenetic classification of conifers.

2 **Figure 2.** Partial chromatograms (**sesquiterpenoids** retention time window) of aliphatic
3 fraction of pyrolysates of *Araucaria* species. The mass spectra of **4a** and **4b** are given in **Fig. 4**.
4 **x**: absence of compound; **tr**: at trace level ; *: cadalane-type compounds.

5 **Figure 3.** Partial chromatograms (**sesquiterpenoids** retention time window) of aromatic
6 fraction of pyrolysates of *Araucaria* species. The mass spectra of **4c**, **4d**, **4e** and **4f** are given in
7 **Fig. 4**. **tr**: at trace level.

8 **Figure 4.** Mass spectra of some identified Araucariaceae sesquiterpenoids. **a**: saturated
9 cadalane-type compounds; **b**: 4 α (*H*)-eudesmane (identified using Wiley275 database); **c**:
10 4 β (*H*)-eudesmane (identified from spectrum in Alexander et al., 1983); **d**: norcadalene
11 (identified from spectrum in Singh et al., 1994); **e**: chamazulene (identified using Wiley275
12 database); **f**: pentamethyl-2,3-dihydroindene (identified using Wiley275 database).

13 **Figure 5.** Mass spectra of identified isohexyl alkylaromatic hydrocarbons in Araucariaceae
14 species and mass spectral cleavage patterns (identified after Ellis et al., 1996). **a**: 1,3,4-trimethyl-
15 2-(4-methylpentyl)benzene; **b**: 2,6-dimethyl-1-(4-methylpentyl)- naphthalene; **c**: 6-ethyl-2-
16 dimethyl-1-(4-methylpentyl)-naphthalenemethylretene.

17 Mass spectra of some unidentified Araucariaceae diterpenoids but similar to some of the
18 identified isohexyl alkylaromatic hydrocarbons: **d** to **b** and **e** to **c**.

19 **Figure 6.** Partial chromatograms (**diterpenoids** retention time window) of aliphatic fraction of
20 pyrolysates of *Araucaria* species. The mass spectra of **8d** and **8e** are given in **Fig. 8**. The mass
21 spectrum of **9a** is given in **Fig. 9**. **★**: labdanes; **x**: absence of compound.

22 **Figure 7.** Partial chromatograms (**diterpenoids** retention time window) of aromatic fractions of
23 pyrolysates of *Araucaria* species. The mass spectra of **5b**, **5c**, **5d** and **5e** are given in **Fig. 5**.
24 The mass spectra of **8a** and **8c** are given in **Fig. 8**. The mass spectra of **9b**, **9c**, **9d**, **9e**, **9f** and **9g**
25 are given in **Fig. 9**. **tr**: in trace; **x**: absence of compound.

26 **Figure 8.** Mass spectra of some tentatively identified Araucariaceae diterpenoids and suggested
27 mass spectral cleavage patterns for **a**, **b**, **d** and **e**. **a**: monoaromatic labdane; **b**: clerodane; **c**:
28 methylretene (tentatively identified according to Alexander et al., 1995); **d**: 17-
29 norphyllocladane (identified according to Noble et al., 1986); **e**: 16(*H*)-homo-phyllocladane.

30 **Figure 9.** Mass spectra of some unidentified but recurrent Araucariaceae diterpenoids. **a** (only
31 for *A. nemorosa* and *A. cunninghamii* in **Fig. 7**), **i** and **j** (only for *Agathis*, in **Fig. 11**): in the
32 aliphatic fraction; **b**, **c**, **d**, **e**, **f**, **g** in the aromatic fraction (**Fig. 5b**, **8b**).

33 **Figure 10.** Partial chromatograms (**sesquiterpenoids** retention time windows) of aliphatic
34 (**left**) and aromatic (**right**) fractions of pyrolysates of *Agathis* species. The mass spectra of **4a**,
35 **4b**, **4e**, **4f** and **4f** are given in **Fig. 4**. **x**: absence of compound ; *: cadalane-type compounds.

36 **Figure 11.** Partial chromatograms (**diterpenoids** retention time windows) of aliphatic (**left**) and
37 aromatic (**right**) fractions of pyrolysates of *Agathis* species. The mass spectra of **5b** to **5e** are
38 given in **Fig. 5**. The mass spectra of **8a** and **8c** are given in **Fig. 8**. The mass spectra of **9b** to **9j**
39 are given in **Fig. 9**. **x**: absence of compound.

40 **Figure 12.** Partial chromatograms [**sesquiterpenoids** (**a**, **b**) and **diterpenoids** (**c**, **d**) retention
41 time windows] of aliphatic (**a**, **c**) and aromatic (**b**, **d**) fractions of pyrolysates of *Wollemia*
42 *nobilis*. The mass spectra of **4a**, **4b**, **4c**, **4d** and **4f** are given in **Fig. 4**. The mass spectra of **5b** to

43 **5e** are given in **Fig. 5**. The mass spectra of **8c** are given in **Fig. 8**. The mass spectra of **9b** to **9f**
44 are given in **Fig. 9**. *: cadalane-type compounds.