

Interactive comment on “Primary production and respiration of hypersaline microbial mats as a response for high and low CO₂ availability” by L. Bento et al.

L. Bento et al.

luizjb@gmail.com

Received and published: 1 March 2013

Dear editor,

We greatly appreciate the valuable comments and suggestions received from anonymous referees #1 and #3, and Dr. Alexandra Rao. Our reactions to their recommendations and the actions that directly were taken to address them are described in detail below.

Actions taken in response to the comments by the anonymous referee #1: 1. Anonymous referee #1: “Light intensity. P12742L26. “Excess” is quite relative. What light level is actually excessive (even in the strict sense of ‘supersaturating’) depends on
C8612

the photoacclimation state of the sample. Without knowing the photosynthetic light response it is not possible to determine if a certain light level is excessive. As this determines the interpretation of the results regarding the exposure to different light intensities, it would be important to know the light-response of MPB photosynthesis, when it saturates, when is supersaturating etc.” Comment: We agree. It was not an aim of our study to contribute to the complex question of microphytobenthos photoacclimation, a subject properly addressed by other studies (i.e. Serodio et al., 2008). We think that the most part of the problem correctly noted by anonymous referee #1 was induced by the use of the word “excess”. Furthermore, the light level used was of the same magnitude of other experimental studies (Serodio et al., 2005; Blanchard & Gall, 1994; Lefebvre et al., 2011; Cruz & Serodio, 2008), and field conditions can be even 2-3 times higher (Perkins et al., 2001; Serodio et al., 2008; Chevalier et al., 2010). Action: Removed the expression “excess light” from the text.

2. Anonymous referee #1: Replication. All figures show measurements of a single profile under each experimental condition. In fact, it seems that the whole paper is based on observations of a single sample per treatment. Is this true, or are the figures showing only representative cases? Comment: We agree that clarification is necessary. Each round of experiments was conducted in one spot, where profiles were recorded repeatedly to show short-term effects while was blocked out. It was not our goal to describe the spacial heterogeneity of the microbial mat, which may require > 100 replicates for correct estimation (Spilmout et al., 2011). Action: We included in the text the time lapse to reach steady-states and explain why we chose the experimental design with successive microprofiles to accomplish our objective.

3. Anonymous referee #1: Biofilm community composition. This is a main aspect, that could condition many topics of the discussion, and which is not adequately addressed. The authors simply state that the samples are dominated by diatoms and cyanobacteria, but will would be important to know, at least 1) which of these groups is in fact dominant (and what is their proportional abundance) 2) main genera present, if all

motile. Comment: We agree that more information on biofilm community composition could have supplemented the discussion, but the main, general conclusions are independent from the biofilm composition. The objective of this study was to demonstrate the short-term effect of atmospheric CO₂ on the microbial mat and that was clearly accomplished. Action: In Discussion we now list several parameters, including community composition, which would be very interesting and important to include in deeper investigations of the role of CO₂ limitation.

4. Anonymous referee #1: Microalgae biomass. This is also a crucial piece of information needed to adequately characterize the studied biofilms. Carbon depletion will ultimately depend on absolute photosynthetic rates, which naturally depend on the amount of photosynthetic organisms present. Comment: We agree that microalgae biomass rather than CO₂ or light in some places will be the more limiting factor for production. For the present place we qualitatively state that the phototrophs formed a dense mat, and the results confirmed that CO₂ became the limiting factor within the top 0.3 mm while high rates of GOP were found down to about 1 mm depth.

5. Anonymous referee #1: Some results seem counter-intuitive. What is the explanation for: - the photosynthetic activity of the bottom layer of the sediment being stimulated by lower atmospheric CO₂? (P12741 L11) - why a decrease in light intensity decreases respiration? (P12741L23-25) Comment: Evidently we have not explained all this very well and too many aspects with marginal connection to the data were addressed. Action: These discussions have been revised with emphasis on fewer and more straight forward interpretations, conclusions and suggestions. See also comment 12, Anonymous referee #1.

6. Anonymous referee #1: CO₂ concentrations. Were the tested CO₂ concentrations ecologically-relevant? Also, would be interesting if some information could be given regarding how long do changes in CO₂ in the atmosphere take to be translated into changes within the biofilm. Comments and actions: All six different CO₂ concentrations used in this study were ecologically or scientifically relevant. Five can be experienced

C8614

by microalgae depending on their microhabitat. For instance in the lakes located in the studied area water pCO₂ can vary from 3 to 20000 μ atm (Marotta et al., 2010) and even higher values may occur in the sediment. In the revised text it is discussed how the results with the applied CO₂ concentrations allow extrapolations to atmospheric changes in past and coming centuries. The effect of zero CO₂ provided a dramatic and simple illustration of how CO₂ limitation worked and is actually a favorite figure when explaining and teaching this subject. As now specified in the text we observed a time lapse of 30-60 minutes until CO₂ changes were translated into new steady state oxygen profiles. It is also discussed how careful you must be in extrapolating these results to other situations, as water height and numerous other physical, chemical, and biological parameters are probably important and deserve further investigations.

7. Anonymous referee #1: P12738 L18-20. A scheme would be very useful. Comment: We agree. Action: A scheme is included as Figure 2.

8. Anonymous referee #1: P12739 L14. Halogen lamps are known to deliver a lot of infrared radiation. Did the halogen lamp heat the sample surface significantly? Comment and action: As described in P12740L13 our light source was a slide projector and the cooler from the slide projector kept the heat away from the sample. The distance to the mat (21.5 cm) and the recorded temperature of the water column (25-26 C) is specified in the text.

9. Anonymous referee #1: P12739 L27-. How accurate was this technique? Were the CO₂ concentrations measured independently? Comment and action: The whole technique and the way by which the CO₂ concentration was controlled and calculated is now explained in M&M (See also comment 7 from Alexandra Rao). The complete elimination of NPP at 0 ppm confirmed the successful elimination of CO₂ from the air line and avoidance of any CO₂ intrusion in tubes and chambers further downstream. In fact this improvised field-lab technology turned out to work better than most of the commercial gas-mixing systems we use in our main laboratories.

C8615

10. Anonymous referee #1: P12742 L4. What does it mean exactly 'absolute carbon limitation'? This is presented as one of the key findings of this study, but the meaning of this expression is not clearly explained. Comment and action: We realize that "absolute carbon limitation" is a confusing expression and it is now eliminated. We now emphasize that decreasing atmospheric concentration translates directly into a growing zone with no NPP. It is well known that microphytes can use both CO₂ and HCO₃⁻ as carbon source, but the actual carbonate speciation and transport in this very dynamic system is not resolved, and as our data have little to add there, we avoid a lengthy discussion.

11. Anonymous referee #1: P12742 L22. It is hard to accept that CO₂ is the most important limiting factor for primary productivity – what about light? Also, wouldn't carbon limitation be dependent on microalgae biomass? Comment and action: Agree. In our efforts to draw attention to one overlooked and ignored factor we have ourselves been a little too narrow-minded. In the revised discussion we emphasize how many more interacting factors including light and abundance that have to be considered when evaluating primary productivity control at a specific place and time.

12. Anonymous referee #1: P12742 L23. Strickly speaking, a decrease in light intensity always causes an increase in photosynthetic efficiency. This is contradictory with the statement that elevated light intensities cause higher photosynthetic efficiencies (12743L11-12). Comment and action: This is a complex matter when dealing with a mixed community with a steep light gradient. As we did not study light adaption directly, most of this discussion has now been discarded. We do state that the observed CO₂ limitation should lead to dual selection for the most light tolerant and the most light efficient forms. But without further experimental support we refrain from mentioning that this might in turn lead to the counter-intuitive conclusion that higher light intensities favors a more light efficient population in the lower part of the mat. This is yet another interesting aspect of CO₂ limitation for future studies.

13. Anonymous referee #1: P12742 L24. Is photorespiration expected to occur in this particular communities? Another strong argument for knowing the composition of the

C8616

biofilm. Comment: Photorespiration did occur at high rates in this community and the complete elimination of NPP in some zones indicated that intracellular storages of inorganic carbon were not important. Diatoms are indeed known to use photorespiration (e.g. Glud et al. (1992), but a role of the cyanobacteria cannot be excluded without much more thorough investigations of community composition, growth, stratification, etc. This, however, will not affect the major conclusions.

14. Anonymous referee #1: P12743 L26-27 Meaning "can become carbon limited" (in the context of elevated CO₂)?? Comment: They can be carbon limited even with CO₂ concentrations in atmosphere much higher than actual. These correlations should now be much clearer in the revised results and discussions.

15. Anonymous referee #1: P12739 L16,20: I think the IS units are $\mu\text{mol m}^{-2} \text{s}^{-1}$ P12742 L11 "Direct" instead of "Directly"? P12740 L8 Why not 50 micrometers? Comment: We agree and the changes were made.

Actions taken in response to the comments by Alexandra Rao: 1. Alexandra Rao: The authors often discuss "respiration", but they mean "aerobic respiration". It should be stated clearly somewhere that anaerobic respiration is not measured, so their estimates probably significantly underestimate total respiration in sediment (mat) samples. Comment and action: We just follow the conventional term used in oxygen microsensors literature since the first paper published by Niels Peter Revsbech (Revsbech et al., 1981) and other recent papers (Ulstrup et al., 2011; Wangpraseurt et al., 2012; Sommer et al., 2010; Pringault et al., 2009; Walker et al., 2011). In some places we now do discuss anaerobic respiration and aerobic, lithotrophic respiration which can be deduced from the data, and this is clearly distinguished from conventional, organotrophic aerobic respiration.

2. Alexandra Rao: and used frequently (abstract line 11; p. 12741 l. 2; p. 12741 l. 21; p. 12742 l. 7). This is consistent with poor use of the English language throughout the paper (see Technical Corrections below), but this is particularly bad because it sounds

C8617

like respiration is responsible for part of the O₂ produced by primary production, when of course, respiration and photosynthesis have opposite effects on O₂ flux Comment: The English is hopefully better now. The wording “respiration represents XX% of GPP” was indeed incorrect and other expressions are now used when respiration is compared with GPP. For further clarification we have also renamed GPP to GOP (gross oxygen production) as CO₂ limitation was found to decouple oxygen production from carbon fixation, which must be the real meaning of primary production.

3. Alexandra Rao: I understand the importance of high CO₂, but I fail to see the point of testing the effect of low (0 and 100 ppm) atmospheric CO₂ concentrations. Comment: See comment 6 to referee #1.

4. Alexandra Rao: Section 2.2. Measurements of pH and alkalinity are mentioned here, but not presented anywhere in the text. If the results are not presented, then the methods should not be mentioned. If the authors decide to present these results, then it's also important to mention what kind of standards were used for pH and alkalinity measurements. The absence of this information is conspicuous. Comment: In situ pH and alkalinity are presented at table 1 and the methods are described in M&M. Alkalinity and in particular pH surely varied significantly in the mat and the overlying water during the experiments, but microsensors to record this were not available. When the chambers were opened after each of the two experimental rounds we used indicator strips to check pH. But realizing that these sparse data of poor quality are of virtually no use in the evaluation of the experiments, we have left them out in the new version. To our best knowledge no one has yet resolved the complex speciation and transport of inorganic carbon species in such a dynamic system, and this challenge is now put forward in Discussion.

5. Alexandra Rao: Section 2.2, line 20. How were the diatoms and cyanobacteria identified? Comment and action: They were identified by direct observation with dissection microscope and this is added to the text.

C8618

6. Alexandra Rao: Section 2.3. How were NPP rates calculated? Yes, it's useful to know NPP was calculated using a particular kind of software, but it's also important to provide more detail about the calculations. Was it simply determined as the sum of upward and downward diffusive O₂ fluxes in the euphotic surface layer of the mats? Comment and action: The approach and the use of the numerical simulation model is now explained much more detailed.

7. Alexandra Rao: Section 2.4. The sample chamber, covered with thin plastic film with an “opening” for the microsensor and no apparent CO₂ measurements for verification, just doesn't seem very well controlled. Is there an outflow port for the air that's being pumped in to keep chamber pressure constant, or does pressure fluctuate? Given that flow rates change significantly with the aging of pump tubing, how well regulated is the peristaltic pump flow? Given all these uncertainties and no measurement verification, it's difficult to believe the CO₂ levels calculated are at all precise! Comment: See response to comment 9 from Anonymous referee #1. Action: The technique to control CO₂ is now more accurately described and a drawing of the experimental chamber is added.

8. Alexandra Rao: Section 2.4. The authors mention high and low CO₂ levels, but precise concentrations should be listed in the Methods section. Comment and action: We agree and the concentration ranges are now specified in M&M .

9. Alexandra Rao: Section 2.4. Very little detail is provided about the microprofiling. Were profiles really measured “in the same spot” (p. 12740 l. 9)? If so, disturbances caused by microsensor insertion may well be responsible for the changes observed in microprofiles (Kühl and Revsbech 2001). Apparently, there is water overlying the microbial mats (p. 12740 l. 12), but there is no other mention of the overlying water – how deep is it? Was it stirred during measurements, to maintain a diffusive boundary layer (DBL)? How thick was the DBL? How was the sediment-water-interface discerned for flux measurements? How was the O₂ sensor calibrated? Was there an offset in the calibration according to calculated concentrations in the mat overlying water? Was

C8619

there a cross-calibration with Winkler titrations? How were sensors positioned at the mat-water-interface? How were the profiles aligned after measurement? How much waiting time was there at each depth before recording readings? How much time were the mats allowed to equilibrate after changing light and CO₂ levels? Comment and action: We apologize for the insufficient description of the experimental set up. All the requested details are now described and explained in the revised M&M. We do not agree that profiling in the same spot could be responsible for changes observed in the microprofiles, and in M&M we now explain why repeated profiling in one spot is actually a way to avoid disturbances and irrelevant variations. See also response to comments 2, 6 and 9 from reviewer #1.

10. Alexandra Rao: Section 2.4, p. 12740, l. 12. The absence of changes in pH in the overlying water, despite dramatic changes in chamber CO₂ levels (0 to 5000 ppm) is surprising, and suggests to me that there may be a big problem in the regulation of chamber CO₂ levels. See comment 7 above. Comment: As a confirmation of the gas mixing system we observed similar oxygen profiles under normal atmospheric air as under a gas mix with the same prescribed CO₂ concentration of 380 ppm.

11. Alexandra Rao: The terminology, “NPP participation in GPP” on p. 12741 line 24 is just another example of poor use of the English language, but this one really stands out. NPP doesn’t “take part in” GPP in any way, although it could be said that NPP represents a fraction of GPP after accounting for respiration. Comment: We agree. As with comment 2 we have carefully inspected and corrected these phrases.

12. Alexandra Rao: What is meant by “absolute” carbon limitation, on p. 12742 l. 4? Comment: See answer to Anonymous referee #1 (comment 10).

13. Alexandra Rao: Section 4, 1st paragraph. The authors suggest that phototrophs are carbon limited when CO₂ was removed from the chamber atmosphere, because respiration was equal to GPP, so there is a closed carbon cycle. This completely ignores anaerobic respiration, since the authors only measure O₂ consumption and

C8620

therefore can only consider aerobic respiration. This is a hypersaline lagoon, so I imagine anaerobic respiration is also important. Comment: The anaerobic processes are actually not hidden. The effect of anaerobic, organotrophic respiration on the oxygen balances depends on the fate of the reduced products; in this system primarily sulfur and iron species. If the reduced products are re-oxidized immediately, the net effect will be just as if it was aerobic, organotrophic respiration. If the reduced products pile up, i.e. as iron sulfide or pyrite, the excess CO₂ generation relative to oxygen reduction will result in a net production of oxygen in the lower part of the oxic and photic zone. In the opposite case where reduced products are oxidized faster than they are generated, the lower zone will have a net consumption of oxygen. The latter is actually what was observed, in particular during the first round of experiments. This is now pointed out and discussed in the text. Notice that this has no impact on the evaluation of CO₂ limitation in the upper production zone.

14. Alexandra Rao: The suggestion of a 2-layer system, in which the surface and deeper layers respond differently to changes in chamber CO₂ levels (p. 12742 l. 12-17; p. 12741 l. 3-14; p.12743 l. 15-18) could have been affected by disturbances associated with repeated profiling in the same spot – see comment 9 above. For the sake of argument, assume repeated profiles are being measured in the same spot, each time under lower CO₂ levels, from 5000 ppm to 380 ppm as in this study. Each time the sensor is inserted, porewater at this spot is mixed. It’s entirely conceivable that this could explain the changes shown, e.g. in Fig. 2. Comment: We do not agree. As explained earlier (comment 9) artifacts from repeated profiling are assumed to be negligible on these samples (please check cited literature). Each graph at figure 2 represents the steady-state of each treatment. At least 6 profiles were performed for each treatment, in a time series. The steady-state represents at least 3 profiles that were considered stabilized since they had the same shape.

15. Alexandra Rao: Figure 1 – This 3-panel map is not very useful. In particular, the expansion from the 2nd panel to the 3rd panel provides no extra information about the

C8621

relative dimensions of Visgueiro lagoon (depth, area) with respect to the neighboring lagoons. Either zoom in much more in the 3rd panel and provide more detail on Visgueiro and maybe a couple of surrounding lagoons, or eliminate the 3rd panel. Comment: We agree. Action: Figure 1 were taken out and the coordinates were provided.

16. Alexandra Rao: Figure 2 – I'm surprised the authors provide only 1 replicate profile at each CO₂ level. I'm also surprised that it seems like all measurements were done on only 2 mat samples (1 for high CO₂ measurements, 1 for low CO₂ measurements, see Section 2.4). With no replication, it's difficult to have any confidence in the results. Comment: Each CO₂ level represents a time series of profiles. Since we are describing a phenomenon that occurs in short temporal scale and measurements of gross primary production in a profile with the microsensors method takes at least 30-40 minutes to be made (Glud, 2008) this approach was chosen by the authors. It was not our goal to describe the spacial heterogeneity of the microbial mat, nor to use these measurements to illustrate a larger scale. See also comments to Anonymous referee #1, question 2. The phenomenon is confirmed since we have steady-state profiles of each treatment.

17. Alexandra Rao: If the issues mentioned above can be addressed, the manuscript also requires extensive language editing. Numerous spelling and grammatical mistakes, poor sentence structure and organization are evident throughout the text, making it difficult to understand in some cases. Comment: We agree. Action: We revised closely the text to improve clarity

18. Alexandra Rao: The figures are not discussed in order in the text. For example, in section 3.1, the text jumps from a reference to Fig. 2 to the following reference to Fig. 6. The order of the figures should be changed to match the order in which they are mentioned in the text. Comment: In the revised Result section the order of the figures fits more naturally. We could not avoid a few jumps as NPP is discussed somewhat separated from GOP, but the GOP profiles have to be shown close to their corresponding NPP profiles. Figure 7 integrates data from the previous 4 figures and

C8622

is now introduced after them as requested.

Actions taken in response to the comments by anonymous referee #3: 1. Anonymous referee #3: The major problem I identify is replication. All figures show measurements of a single profile. Averages and standard deviations should be presented if, in fact, replicates were used. From reading Materials and Methods I get the feeling no replication was used. Comment: See response to comment 2 of anonymous referee #1 and comment 16 of Alexandra Rao.

2. Anonymous referee #3: The microphytobenthic community is very poorly described. The authors refer only diatoms and filamentous cyanobacteria. What was the relative contribution of each group? What were the main genera/species present? Comment: See response to similar comment 3 of anonymous referee #1 and comment 5 of Alexandra Rao.

3. Anonymous referee #3: The authors refer this manuscript as the "first account of carbon limitation in a microbial mat". Previous studies refer carbon limitation in microphytobenthic biofilms, for example Cook and Roy (2006) (this paper is in the reference list of the manuscript). Comment: We agree that our formulation was in inappropriate. It is now specified that the novelty of our study is the control of MPB production by CO₂ in the air.

4. Anonymous referee #3: The English requires improvement. Comment: The text was carefully checked to improve clarity and language.

References cited:

Blanchard, G. F., & Cariou-Le Gall, V. (1994). Photo synthetic characteristics of microphytobenthos in Marennes-Oléron Bay, France: Preliminary results. *Journal of experimental marine biology and ecology*, 182(1), 1-14. Chevalier, E. M., Gévaert, F., & Créach, A. (2010). In situ photosynthetic activity and xanthophylls cycle development of undisturbed microphytobenthos in an intertidal mudflat. *Journal of Experimental Ma-*

C8623

rine Biology and Ecology, 385(1), 44-49. Cruz, S., & Serôdio, J. (2008). Relationship of rapid light curves of variable fluorescence to photoacclimation and non-photochemical quenching in a benthic diatom. *Aquatic Botany*, 88(3), 256-264. Glud, R. N., Ramsing, N., & Revsbech, N. (1992). Photosynthesis and photosynthesis-coupled respiration in natural biofilms quantified with oxygen microsensors. *Journal of phycology*, 28(1), 51-60. Glud, R. N. (2008). Oxygen dynamics of marine sediments. *Marine Biology Research*, 4(4), 243-289. Lefebvre, S., Mouget, J. L., & Lavaud, J. (2011). Duration of rapid light curves for determining the photosynthetic activity of microphytobenthos biofilm in situ. *Aquatic Botany*, 95(1), 1-8. Marotta, H., Duarte, C. M., Pinho, L., and Enrich-Prast, A. (2010). Rainfall leads to increased pCO₂ in Brazilian coastal lakes, *Biogeosciences*, 7, 1607-1614, doi:10.5194/bg-7-1607-2010 Perkins, R. G., Underwood, G. J. C., Brotas, V., Snow, G. C., Jesus, B., & Ribeiro, L. (2001). Responses of microphytobenthos to light: primary production and carbohydrate allocation over an emersion period. *Marine Ecology Progress Series*, 223, 101-112. Pringault, O., Tesson, S., & Rochelle-Newall, E. (2009). Respiration in the light and bacterioplankton coupling in a coastal environment. *Microbial ecology*, 57(2), 321-334. Revsbech, N. P., Jørgensen, B. B., & Brix, O. (1981). Primary Production of Microalgae in Sediments Measured by Oxygen Microprofile, H¹⁴CO₃-Fixation, and Oxygen Exchange Methods. *Limnology and Oceanography*, 26(4), 717-730. Serôdio, J., Vieira, S., Cruz, S., & Barroso, F. (2005). Short-term variability in the photosynthetic activity of microphytobenthos as detected by measuring rapid light curves using variable fluorescence. *Marine Biology*, 146(5), 903-914. Serôdio, J., Vieira, S., & Cruz, S. (2008). Photosynthetic activity, photoprotection and photoinhibition in intertidal microphytobenthos as studied in situ using variable chlorophyll fluorescence. *Continental Shelf Research*, 28(10), 1363-1375. Sommer, S., Linke, P., Pfannkuche, O., Niemann, H., & Treude, T. (2010). Benthic respiration in a seep habitat dominated by dense beds of ampharetid polychaetes at the Hikurangi Margin (New Zealand). *Marine Geology*, 272(1), 223-232. Spilmont, N., Seuront, L., Meziane, T., & Welsh, D. T. (2011). There's more to the picture than meets the eye: sampling microphytobenthos in a heterogeneous en-

C8624

vironment. *Estuarine, Coastal and Shelf Science*, 95-4, 470-476. Ulstrup, K. E., Kühl, M., Van Oppen, M. J., Cooper, T. F., & Ralph, P. J. (2011). Variation in photosynthesis and respiration in-geographically distinct populations of two reef-building coral species. *Aquatic Biology*, 12(3), 241-248. Walker, R. A., Hallock, P., Torres, J. J., & Vargo, G. A. (2011). Photosynthesis and respiration in five species of benthic foraminifera that host algal endosymbionts. *The Journal of Foraminiferal Research*, 41(4), 314-325. Wang-praseurt, D., Weber, M., Røy, H., Polerecky, L., de Beer, D., & Nuges, M. M. (2012). In situ oxygen dynamics in coral-algal interactions. *PLoS one*, 7(2), e31192.

Interactive comment on *Biogeosciences Discuss.*, 9, 12735, 2012.

C8625