



1 **High net CO₂ and CH₄ release at a eutrophic shallow lake**
2 **on a formerly drained fen**

3
4 **D. Franz¹, F. Koebsch¹, E. Larmanou¹, J. Augustin² and T. Sachs¹**

5 [1] {Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences, Telegrafenberg,
6 14473 Potsdam, Germany}

7 [2] {Institute for Landscape Biogeochemistry, Leibniz Centre for Agricultural Landscape Research
8 (ZALF), Eberswalder Str. 84, 15374 Müncheberg, Germany}

9 *Correspondence to:* D. Franz (daniela.franz@gfz-potsdam.de)

10

11 **Abstract**

12 Drained peatlands often act as carbon dioxide (CO₂) hotspots. Raising the groundwater table is
13 expected to reduce their CO₂ contribution to the atmosphere and revitalize their function as carbon
14 (C) sink in the long term. Without strict water management rewetting often results in partial flooding
15 and the formation of spatially heterogeneous, nutrient-rich shallow lakes. Uncertainties remain as to
16 when the intended effect of rewetting is achieved, as this specific ecosystem type has hardly been
17 investigated in terms of greenhouse gas exchange (GHG) exchange. In most cases, methane (CH₄)
18 emissions increase under anoxic conditions due to a higher water table and in terms of global warming
19 potential (GWP) outperform the shift towards CO₂ uptake, at least in the short-term.

20 Based on eddy covariance measurements we studied the ecosystem–atmosphere exchange of CH₄ and
21 CO₂ (NEE) at a shallow lake situated on a former fen grassland in Northeast (NE) Germany. The lake
22 evolved shortly after flooding, 9 years previous to our investigation period. The ecosystem consists of
23 two main surface types: open water (inhabited by submerged and floating vegetation) and emergent
24 vegetation (particularly including the eulittoral zone of the lake, dominated by *Typha latifolia*). To
25 determine the individual contribution of the two main surface types to the net CO₂ and CH₄ exchange
26 of the whole lake ecosystem, we combined footprint analysis with CH₄ modelling and NEE partitioning.

27 The CH₄ and CO₂ dynamics were strikingly different between open water and emergent vegetation.
28 Net CH₄ emissions from the open water area were around 4-fold higher than from emergent vegetation
29 stands, accounting for 53 and 13 g CH₄ m⁻² a⁻¹, respectively. In addition, both surface types were net



30 CO₂ sources with 158 and 750 g CO₂ m⁻² a⁻¹, respectively. Unusual meteorological conditions in terms
31 of a warm and dry summer and a mild winter might have facilitated high respiration rates. In sum, even
32 after 9 years of rewetting the lake ecosystem exhibited a considerable C loss and global warming
33 impact, the latter mainly driven by high CH₄ emissions. We assume the eutrophic conditions in
34 combination with permanent high inundation as major reasons for the unfavourable GHG balance.

35

36 1 Introduction

37 Peatland ecosystems play an important role in global greenhouse gas (GHG) cycles, although they
38 cover only about 3 % of the earth's surface (Frolking et al. 2011). Peat growth depends on the
39 proportion of carbon (C) sequestration and release. Pristine peatlands act as long-term C sinks and are
40 near-neutral (slightly cooling) regarding their global warming potential (GWP; Frolking et al. 2011),
41 dependent on rates of C sequestration and methane (CH₄) emissions. However, many peatlands
42 worldwide are used e.g. for agriculture, as are more than 85% of the peatlands in Germany and the
43 Netherlands (Silvius et al. 2008). Drainage is associated with shrinkage and internal phosphor
44 fertilisation of the peat (Zak et al. 2008). Moreover, the hydrology of the area as well as physical and
45 chemical peat characteristics are changing (Holden et al. 2004, Zak et al. 2008). Above all, drained and
46 intensively managed peatlands are known as strong sources of carbon dioxide (CO₂; e.g. Joosten et al.
47 2010, Hatala et al. 2012, Beetz et al. 2013). On the other hand, lowering the water table is typically
48 accompanied with decreasing CH₄ emissions (Roulet et al. 1993). Emission factors of 1.6 g CH₄ m⁻² a⁻¹
49 and 2235 g CO₂ m⁻² a⁻¹ were assigned to temperate deep-drained nutrient-rich grassland in the 2013
50 wetland supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2014).

51 In the last decades rewetting of peatlands attracted attention in order to stop soil degradation, reduce
52 CO₂ emissions and to recover their functions as C and nutrient sink and ecological habitat (Zak et al.
53 2015). Large rewetting projects were initiated, e.g. the Mire Restoration Program of the federal state
54 of Mecklenburg-West Pomerania in Northeast (NE) Germany (Berg et al. 2000) starting in 2000 and
55 involving 20 000 ha of formerly drained peatlands, thereby especially fens (Zerbe et al. 2013) e.g. in
56 the Peene river catchment. However, uncertainties remain as to when the intended effects of
57 rewetting are achieved. Only few studies exist on the temporal development of GHG emissions of
58 rewetted fens, especially on longer time scales. Augustin and Joosten (2007) discuss three very
59 different states following peatland rewetting based on observations at Belarusian mires, though
60 without specifying the individual lengths of the phases. Broad agreement exists concerning the CH₄
61 hot spot characteristic of newly rewetted peatlands (e.g. Meyer et al. 2001, Hahn-Schöfl et al. 2011,



62 Knox et al. 2015). However, a rapid recovery of the net CO₂ sink function is not consistently reported
63 (e.g. Wilson et al. 2007).

64 Peatlands develop a pronounced microtopography after drainage and subsequent subsidence.
65 Rewetting e.g. in the Peene river catchment resulted in the formation of large-scale shallow lakes in
66 the lower parts of the fens, with water depths usually below 1 m (Zak et al. 2015, Steffenhagen et al.
67 2012). These new ecosystems are nutrient-rich and most often strikingly different from natural
68 peatlands. They experience a rapid secondary plant succession (Zak et al. 2015). Helophytes are
69 expected to progressively enter the open water body over the time leading to the terrestrialisation of
70 the shallow lake and in the best case peat formation. However, this new ecosystem type and its
71 progressive transformation have hardly been investigated in terms of GHG dynamics. The ecosystem-
72 inherent spatial heterogeneity suggests complex patterns of GHG emissions due to distinct GHG source
73 or sink characteristics of the involved surface types (generally open water and the littoral zone)
74 resulting in measurement challenges. Site-specific heterogeneity implicitly has to be considered for
75 the evaluation of ecosystem scale flux measurements (e.g. Barcza et al. 2009, Hendriks et al. 2010,
76 Hatala Matthes et al. 2014). The importance of small open water bodies in wetlands as considerable
77 GHG sources was highlighted in previous studies (e.g. by Schrier-Uijl et al. 2011, Zhu et al. 2012, IPCC
78 2014) and in case of CH₄ even for landscape-scale budgets e.g. by Repo et al. (2007). In addition, the
79 littoral zone of lakes is often found to be a CH₄ hot spot (Juutinen et al. 2003, Wang et al. 2006) with a
80 contribution of up to 90 % to the whole-lake CH₄ release (Smith and Lewis 1992), albeit depending on
81 the lake size (Bastviken et al. 2004) and plant community. Rööm et al. (2014) measured the largest CH₄
82 (and CO₂) emissions of a temperate eutrophic lake at the helophyte zone within the littoral.

83 The objectives of this study are 1) to investigate the ecosystem-atmosphere exchange of CH₄ and CO₂
84 (NEE) of a nutrient-rich lake ecosystem emerged at a former fen grassland and 2) particularly infer the
85 individual GHG dynamics of the main surface types within the ecosystem and quantify their
86 contribution to the annual exchange rates. Therefore, we applied the eddy covariance technique from
87 May 2013 to May 2014 and used an analytical footprint model to downscale the spatially integrated,
88 half-hourly fluxes to the main surface types “open water” and “emergent vegetation”. The resulting
89 source area (i.e. spatial origin of the flux) fractions were then included in a temperature response (CH₄)
90 and NEE partitioning model (CO₂) in order to quantify the source strength of the two surface types.

91



92 2 Material and methods

93 2.1 Study site

94 The study site “Polder Zarnekow” is a rewetted, rich fen (minerotrophic peatland) located in the Peene
95 river valley (Mecklenburg-West Pomerania, NE Germany, 53°52.5′ N 12°53.3′ E, see Fig. 1), with less
96 than 0.5 m a.s.l. elevation. It is part of the Terrestrial Environmental Observatories Network (TERENO).
97 The temperate climate is characterised by a long-term mean annual air temperature and mean annual
98 precipitation of 8.7 °C and 584 mm, respectively (German Weather Service, meteorological station
99 Teterow, 24 km SW of the study site; reference period 1981–2010). The geomorphological character
100 of the area is predominantly a result of the Weichselian glaciation as the last period of the Pleistocene
101 (Steffenhagen et al. 2012). The fen developed with continuous percolating groundwater flow (Succow
102 2001). Peat depth partially reaches 10 m (Hahn-Schöfl et al. 2011). Drainage was initialized in the 18th
103 century and strongly intensified between 1960 and 1990 within an extensive melioration program
104 (Höper et al. 2008). The decline of the water table to > 1 m below surface and subsequent
105 decomposition and mineralisation of the peat (especially in the upper 30 cm, Hahn-Schöfl et al. 2011)
106 caused phosphor fertilisation (Zak et al. 2008) and soil subsidence to levels below that of adjacent
107 freshwater bodies (Steffenhagen et al. 2012, Zerbe et al. 2013). The latter simplified the rewetting
108 process which was initiated in winter 2004/2005 by opening the dikes.

109 In consequence of flooding the drained fen was converted into a spatially heterogeneous site of
110 emergent vegetation (on temporarily inundated soil) and permanent open water areas. In this study
111 we focus on a eutrophic shallow lake (open water body about 7.5 ha) as part of the rewetted area,
112 with water depths ranging from 0.1 to 0.7 m. During the study period the open water body of the lake
113 was inhabited by submerged and floating macrophytes, particularly *Ceratophyllum demersum*, *Lemna*
114 *minor*, *Spirodela polyrhiza* (Steffenhagen et al. 2012) and *Polygonum amphibium*, which rather
115 corresponds to the sublittoral zone in a typical lake zonation. *Ceratophyllum* and *Lemna* sp. were
116 already reported to colonise the lake in the second year of rewetting (Hahn-Schöfl et al. 2011). *Phalaris*
117 *arundinacea*, that dominated the fen before rewetting, died off in the first year of inundation (Hahn-
118 Schöfl et al. 2011) and has been limited to the non-inundated periphery of the ecosystem. Helophytes
119 (e.g. *Glyceria*, *Typha*) started the colonisation of lake margins and other temporarily inundated areas
120 in the third year of rewetting. The eulittoral zone of the lake is now dominated by *Typha latifolia* stands
121 gradually colonising the open water in the last years. Emergent vegetation stands also include sedges
122 as *Carex gracilis* (Steffenhagen et al. 2012). At the bottom of the shallow lake an up to 30 cm thick
123 layer of organic sediment evolved, initially fed by fresh plant material of the former vegetation and



124 since then continuously replenished by recent aquatic plants and helophytes after die-back (Hahn-
125 Schöfl et al. 2011).

126 2.2 Eddy covariance and additional measurements

127 We conducted eddy covariance (EC) measurements of CO₂ and CH₄ exchange on a tower placed on a
128 stationary platform at the NE edge of the shallow lake (see Fig. 1). Thereby we ensured to frequently
129 catch the signal from both the open water body and the *Typha latifolia* dominated belt of the shallow
130 lake (eulittoral zone). We defined an area of interest (AOI) in order to focus on an ecosystem
131 dominated by a shallow lake and to avoid a possible impact of the farm and grassland to the north of
132 the shallow lake. The EC measurement setup included: an ultrasonic anemometer for the 3D wind
133 vector (u , v , w) and sonic temperature (HS-50, Gill, Lymington, Hampshire, UK), an enclosed-path
134 infrared gas analyser (IRGA) and an open-path IRGA for CO₂/H₂O and CH₄ concentrations, respectively
135 (LI-7200 and LI-7700, LI-COR Biogeosciences, Lincoln NE, USA). Flowrate was about 10-11 l min⁻¹.
136 Measurement height was on average 2.63 m above the water surface at the position of the tower,
137 depending on the water level. We recorded raw turbulence and concentration data with a LI-7550
138 digital data logger system (LI-COR Biogeosciences, Lincoln NE, USA) at 20 Hz in half-hourly files. The
139 dataset is shown in Coordinated Universal Time (UTC), which is 1 hour behind local time (LT).

140 We further equipped the tower with instrumentation for net radiation, air temperature/humidity, 2D
141 wind direction and speed, incoming and reflected photosynthetic photon flux density (PPFD/PPFD_r)
142 and water level. Additional measurements in close proximity to the tower included precipitation, soil
143 heat flux as well as soil and water temperature. Soil temperature was measured below the water
144 column in depths of 10 cm, 20 cm, 30 cm, 40 cm and 50 cm and water temperature at the sediment-
145 water-interface. All non-eddy covariance-related measurements were logged as 1 min averages/sums
146 (precipitation). Gaps were filled with measurements of the Leibniz Centre for Agricultural Landscape
147 Research (ZALF, Müncheberg, Germany) at the same platform and a nearby climate station (Climate
148 station Karlshof, GFZ German Research Centre for Geosciences, 14 km distance from study site, Itzerott
149 2015).

150 A water density gradient was calculated based on the temperature at the water surface and at the
151 sediment-water interface. The water surface temperature was calculated based on the Stefan-
152 Boltzmann law (see e.g. Foken et al. 2008):

$$153 T_w = \sqrt[4]{\frac{I}{\epsilon_w \sigma_{SB}}} \quad (1)$$

154



155 where T_w is the water surface temperature (K), I is the long-wave outgoing radiation (W m^{-2}), ε_w is
156 the infrared emissivity of water (0.960) and σ_{SB} is the Stefan–Boltzmann constant ($5.67 \cdot 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$). We calculated the density of the air-saturated water at the water surface and the sediment-water
157 interface according to Bignell (1983):
158

$$159 \quad \rho_{as} = \rho_{af} - 0.004612 + 0.000106 * T \quad (2)$$

160 where ρ_{as} is the density of the respective air-saturated water (k m^{-3}), ρ_{af} is the density of the
161 respective air-free water (k m^{-3} ; see Wagner and Pruß 2002) at atmospheric pressure (1013 hPa) and
162 T is the respective water temperature ($^{\circ}\text{C}$). The gradient of the two water densities (air-saturated)
163 $\Delta\rho/\Delta z$ was calculated as difference of the water density (air-saturated) at the sediment-water
164 interface and the surface water density (air-saturated), divided by the distance (m) between the two
165 basic temperature measurements. Changes of the distance due to the fluctuating water level were
166 considered. Positive and negative gradients indicate periods of stratification and thermally induced
167 convective mixing of the water column, respectively.

168 **2.3 Flux computation and further processing**

169 For this analysis we used data from 14 May 2013 to 14 May 2014. We calculated half-hourly fluxes of
170 CO_2 and CH_4 based on the covariances between the respective scalar concentration and the vertical
171 wind velocity using the processing package EddyPro 5.2.0 (LI-COR, Lincoln, Nebraska, USA). Sonic
172 temperature was corrected for humidity effects according to van Dijk et al. (2004). Artificial data spikes
173 were removed from the 20 Hz data following Vickers and Mahrt 1997. We used the planar fit method
174 (Finnigan et al. 2003, Wilczak et al. 2001) for axis rotation and defined the sector borders according to
175 Siebicke et al. (2012). Block averaging was used to detrend turbulent fluctuations. For time lag
176 compensation we applied covariance maximization (Fan et al. 1990). Spectral losses due to crosswind
177 and vertical instrument separation were corrected according to Horst and Lenschow (2009). The
178 methods of Moncrieff et al. (2004) and Fratini et al. (2012) were used for the correction of high-pass
179 filtering and low-pass filtering effects, respectively. For fluctuations of CH_4 density we corrected
180 changes in air density according to Webb et al. (1980), considering LI-7700-specific spectroscopic
181 effects (McDermitt et al. 2011). According to the micrometeorological sign convention, positive values
182 represent fluxes from the ecosystem into the atmosphere (emission) and negative values fluxes from
183 the atmosphere into the ecosystem (ecosystem uptake).



184 **2.4 Quality assurance**

185 We filtered the averaged fluxes according to their quality as follows (see Table 1):

- 186 - We rejected fluxes with quality flag 2 (QC 2, bad quality) based on the 0-1-2 system of Mauder
187 and Foken (2004).
- 188 - CH₄ fluxes were skipped if the signal strength (RSSI) was below the threshold of 14 %. This
189 threshold was estimated according to Dengel et al. (2011).
- 190 - Fluxes with friction velocity (u^*) < 0.12 m s⁻¹ and > 0.76 m s⁻¹ were not included due to
191 considerably high fluxes beyond these thresholds, which were estimated similar to the
192 procedure described in Aubinet et al. (2012) based on binned u^* classes. The storage term was
193 calculated as described in Béziat et al. (2009).
- 194 - Unreasonably high positive and negative fluxes (0.2%/99.8% percentile) were discarded from
195 the CO₂ and CH₄ flux dataset.

196 Quality control (apart from EddyPro internal steps) and the subsequent processing steps were
197 performed with the free software environment R (R Core Team 2012).

198 **2.5 Footprint modelling**

199 We applied footprint analysis to determine the source area including the fractions of the surface types
200 of each quality-controlled half-hourly flux using a footprint calculation procedure following Göckede
201 et al. (2004). The source area functions were calculated based on the analytical footprint model of
202 Kormann and Meixner (2001). Roughness length and vegetation height were estimated with an
203 iterative algorithm (see also Barcza et al. 2009). Based on an aerial image (GoogleEarth,
204 <http://earth.google.com/>) the surface of our study site was classified into two main types and
205 implemented in a land cover grid: “open water” including in particular the open waterbody of the
206 shallow lake with 0.1 to 0.7 m water depth and “emergent vegetation” with a height up to 2 m and
207 including the eulittoral zone of the shallow lake dominated by *Typha latifolia*. The cumulative annual
208 footprint climatology was calculated following Chen et al. (2011). Fluxes were excluded where
209 footprint information was not available or more than 20 % of the source area was outside the AOI (see
210 Fig. 1 and Table 1). The fractional coverage within the AOI (A_i) was 21.7 % for open water.

211 Quasi-continuous source area information for the two surface types were achieved by gapfilling the
212 results of the footprint model with the means of the source area fractions of the surface types (Ω_i) for
213 1°-wind direction-intervals, separately for stable and unstable conditions. In case the sum of the Ω_i was



214 less than 100 %, when the source area exceeded the set borders, we assigned the remaining
215 contribution percentages to emergent vegetation, as the area beyond the borders is dominated by
216 emergent vegetation rather than open water.

217 2.6 Gapfilling

218 An enhanced lookup table (LUT) approach proposed by Reichstein et al. (2005), available as web tool
219 based on the R package REddyProc (<http://www.bgc-jena.mpg.de/REddyProc/brew/REddyProc.rhtml>)
220 was applied for gapfilling and partitioning of NEE measurements ($LUT_{CO_2nofoot}$), with air temperature as
221 temperature variable. For the gapfilling of CH_4 measurements non-linear regression (NLR) was applied
222 ($NLR_{CH_4nofoot}$):

$$223 F_{CH_4} = \exp(a + b_1 \cdot X_1 + \dots + b_j \cdot X_j) \quad (3)$$

224 where a and $b_1 \dots b_j$ are fitting parameters and $X_1 \dots X_j$ are environmental parameters. Several
225 environmental parameters, which were reported to be correlated with CH_4 flux on different time
226 scales, were tested to find the best bi- or multivariate NLR model for the ecosystem CH_4 flux: pressure
227 change, u^* , PAR, air temperature, soil heat flux, soil/peat temperature in different heights and
228 waterlevel. Only fluxes of the best quality (QC 0) were used to fit the NLR model and the LUT.

229 2.7 Calculation of the annual CO_2 and CH_4 budget and the global warming 230 potential (GWP)

231 We used the continuous flux datasets derived from gapfilling for the calculation of annual CO_2 and CH_4
232 budgets. The ecosystem GHG balance was calculated by summation of the net ecosystem exchange of
233 CO_2 and CH_4 using the global warming potential (GWP) of each gas at the 100-year time horizon (IPCC,
234 2013). According to the IPCC AR5 (IPCC, 2013) CH_4 has a 28-fold global warming potential compared
235 to CO_2 (without inclusion of climate-carbon feedbacks).

236 The uncertainty of the annual estimates was calculated as the square root of the sum of the squared
237 random error (measurement uncertainty) and gapfilling error within the one-year observation period
238 (see e.g. Hommeltenberg et al. 2014, Shoemaker et al. 2015). An estimation of the random uncertainty
239 due to the stochastic nature of turbulent sampling according to Finkelstein and Sims (2001) is
240 implemented in EddyPro 5.2.0. In case of the LUT approach the gapfilling error (standard error) was
241 calculated from the standard deviation of the fluxes used for gapfilling, provided by the web tool. For
242 budgets based on the NLR approach we used the residual standard error of the NLR model as gapfilling
243 error (following Shoemaker et al. 2001).



2.8 Estimation of surface type fluxes

To estimate the specific surface type fluxes, we combined footprint analysis with NEE partitioning (using NLR) to assign gross primary production (GPP) and ecosystem respiration (R_{eco}) to the two main surface types ($NLR_{CO2foot}$). R_{eco} and GPP were modelled as sum of the two surface type fluxes weighted by Ω_i (analogous to Forbrich et al. 2011). Night-time R_{eco} (global radiation $< 10 \text{ W m}^{-2}$) was estimated by the exponential temperature response model of Lloyd and Taylor (1994) assuming that night-time NEE represents the night time R_{eco} :

$$R_{eco} = \sum_{i=1}^2 \Omega_i \cdot (R_{ref_i} \cdot \exp(E_0_i (\frac{1}{T_{ref}-T_0} - \frac{1}{T_{air}-T_0}))) \quad (4)$$

where R_{eco} is the half-hourly measured ecosystem respiration ($\mu\text{mol}^{-1}\text{m}^{-2}\text{s}^{-1}$), Ω_i is the source area fraction of the respective surface type, R_{ref} is the respiration rate at the reference temperature T_{ref} (283.15 K), E_0 defines the temperature sensitivity, T_0 is the starting temperature constant (227.13 K) and T_{air} the mean air temperature during the flux measurement. The model parameters achieved for night time R_{eco} were applied for the modelling of day-time R_{eco} . GPP was calculated by subtracting daytime R_{eco} from the measured NEE. GPP was further modelled using a rectangular, hyperbolic light response equation based on the Michaelis–Menten kinetic (see e.g. Falge et al. 2001):

$$GPP = \sum_{i=1}^2 \Omega_i \cdot \left(\frac{GP_{max_i} \cdot \alpha_i \cdot PAR}{\alpha_i \cdot PAR + GP_{max_i}} \right) \quad (5)$$

where GPP is the calculated gross primary production ($\mu\text{mol}^{-1}\text{m}^{-2}\text{s}^{-1}$), Ω_i is the source area fraction of the respective surface type, GP_{max} is the maximum C fixation rate at infinite photon flux density of the photosynthetic active radiation PAR ($\mu\text{mol}^{-1}\text{m}^{-2}\text{s}^{-1}$), α is the light use efficiency ($\text{mol CO}_2 \text{ mol}^{-1}$ photons). We calculated one parameter set for R_{eco} and GPP per day based on a moving window of 28 days (method NLR_{nofoot}). In order to avoid over-parameterization we introduced fixed values of 150 for E_0 and -0.03 and -0.01 for α of emergent vegetation and water bodies, respectively, to get reasonable parameter values for R_{ref} and GP_{max} . We excluded parameter sets for R_{eco} or GPP, if one of the two R_{ref} and GP_{max} parameter values was insignificant ($p\text{-value} \geq 0.05$), negative or zero. In addition, the 1%/99% percentiles of GP_{max} were excluded. These gaps within the parameter set were filled by linear interpolation. Gaps remain in R_{eco} and GPP time series due to gaps in the environmental variables. Gaps up to 3 hours length were filled by linear interpolation. Larger gaps were filled with the mean of the flux during the same time of the day before and after the gap. Due to the moving window approach, we could not estimate model parameters for the first and last 14 days of our study period. Instead, we applied the first and last estimated parameter set, respectively. Modelled GPP and R_{eco} were summed up to half-hourly NEE fluxes and used for alternative NEE gapfilling.



275 As for NEE we expect different CH₄ emission rates of the two surface types. Thus, we extended the NLR
276 model (NLR_{CH₄nofoot}) in a way that the CH₄ flux is the sum of the two surface type fluxes weighted by Ω_i
277 (NLR_{CH₄foot}):

$$278 F_{CH_4} = \sum_{i=1}^2 \Omega_i \cdot \exp(a_i + b_{1i} \cdot X_1 + \dots + b_{ji} \cdot X_j) \quad (6)$$

279 where Ω₁ is the source area fraction of the respective surface type. Considering the principle of
280 parsimony, we combined up to three parameters besides the contribution of the surface types.
281 Remaining gaps were filled by interpolation. Surface type CO₂ and CH₄ fluxes were derived based on
282 the fitted NLR parameters.

283 We calculated the annual budgets of CO₂ and CH₄ for the EC source area, the surface types (assuming
284 source area fraction of 100 % for the respective surface type) and the AOI, the latter following Forbrich
285 et al. (2011) by applying Eq. 4 and Eq. 5 for CO₂ as well as Eq. 6 for CH₄ with the fitted parameters, but
286 A_i instead of Ω_i as weighting surface type contribution. The gapfilling error for the NLR_{CO₂foot} model was
287 based on the residual standard error of both R_{eco} and GPP.

288

289 **3 Results**

290 **3.1 Environmental conditions and fluxes of CO₂ and CH₄**

291 Mean annual air temperature and annual precipitation for the study period were 10.1 °C and 416.5
292 mm, respectively, indicating an unusual dry and warm measurement period compared to the long-
293 term average. The summer 2013 was among the 10 warmest since the beginning of the measurements
294 in 1881 (German Weather Service DWD). From June to August monthly averaged air temperature was
295 0.2 up to 0.9 °C higher and precipitation was 9.1 up to 38.1 mm less than the long-term averages. The
296 open water area of the shallow lake was densely vegetated with submerged and floating macrophytes.
297 A summertime algae slick accumulated in the NE part of the shallow lake. Winter 2013/2014 was
298 characterised by exceptionally mild temperatures and very sparse precipitation. However, a short cold
299 period (see Fig. 2) resulted in ice cover on the shallow lake between 21 January and 16 February 2014.
300 The water level of the shallow lake fluctuated between 0.36 and 0.77 m (at the position of the sensor)
301 and had its minimum at the end of August/beginning of September and its maximum in January. We
302 observed the exposure of normally inundated soil surface at emergent vegetation stands during the
303 dry period in summer 2013.

304 Both CO₂ and CH₄ flux measurement time series showed a clear seasonal trend with median CO₂ flux
305 of 0.57 μmol m⁻² s⁻¹ and a median CH₄ flux of 0.02 μmol m⁻² s⁻¹. CH₄ emissions peaked in mid-August



306 2013 with $0.57 \mu\text{mol m}^{-2} \text{s}^{-1}$. The highest net CO_2 uptake ($-15.34 \mu\text{mol m}^{-2} \text{s}^{-1}$) and release ($21.04 \mu\text{mol}$
 307 $\text{m}^{-2} \text{s}^{-1}$) were both observed in June 2013. A diurnal cycle of CO_2 fluxes with peak uptake around midday
 308 and peak release around midnight was obvious until November 2013 and beginning in March 2014
 309 (see Fig. 3). To investigate the potential presence of a diurnal cycle of CH_4 fluxes we normalized the
 310 mean half-hourly CH_4 fluxes per month with the respective median of the half-hourly fluxes of the
 311 specific month (minimum five 30 min fluxes per day; method modified from Rinne et al. 2007). We
 312 found a clear diurnal cycle of CH_4 fluxes from June to September 2013 and starting again in March 2014
 313 (April and May not shown as the sensor was dismantled) with daily peaks during night time (around
 314 midnight until early morning). The water density gradient indicates thermally induced convective
 315 mixing of the whole water column during the night (around midnight until early morning) from May
 316 until October 2013 and from February to May 2014. In May 2014 the diurnal pattern of the water
 317 density gradient was less pronounced than in May 2013.

318 3.2 Gapfilling performance and annual budgeting of CO_2 , CH_4 , C and GWP

319 The $\text{LUT}_{\text{CO}_2\text{nofoot}}$ approach explained 74 % of the variance in NEE (see Table 2). Median NEE accounted
 320 for $1.9 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$. The annual budget of gapfilled NEE ($\text{LUT}_{\text{CO}_2\text{nofoot}}$) between 14 May 2013 and 14
 321 May 2014 was $524.5 \pm 5.6 \text{ g CO}_2 \text{ m}^{-2}$ (see Table 3), characterising the site as strong CO_2 source with
 322 moderate rates of R_{eco} and GPP. We found a surprising CO_2 release strength during summer 2013,
 323 where already at the end of June daily R_{eco} often exceeded GPP. The highest CO_2 emission and uptake
 324 rates of $24.8 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and $-27.9 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ were both observed in the beginning of July 2013 (see
 325 Fig. 2). July 2013 accounted for 23.2 % and 25.8 % of the annual R_{eco} and GPP, respectively. In addition,
 326 net CO_2 release outside the growing season (definition of the growing season following Lund et al.
 327 2010; until 19 November 2013 and starting 26 February 2014) was $203.7 \text{ g CO}_2 \text{ m}^{-2}$ with a median of
 328 $2.2 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$.

329 The environmental variable giving the best NLR model for CH_4 was soil temperature in 10 cm depth
 330 (T_{s10}):

$$331 F_{\text{CH}_4} = \exp(-7.224 + 0.313 \cdot T_{s10}) \quad (7)$$

332 The model described 79 % of the variance in CH_4 flux (see Table 2). Including additional environmental
 333 variables to the regression function did not increase the model performance significantly. Cumulative
 334 CH_4 emissions were $40.5 \pm 0.2 \text{ g CH}_4 \text{ m}^{-2} \text{ a}^{-1}$ (see Table 3). Median CH_4 emissions were $41.9 \text{ mg m}^{-2} \text{ d}^{-1}$,
 335 peaked at the end of July 2013 with $0.6415 \text{ g CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ and were at the minimum in January 2014
 336 (see Fig. 2). The month with the highest proportion of annual CH_4 emissions was August 2013 (27.3 %).



337 Non-growing season CH₄ fluxes only accounted for a small proportion within the annual budget, about
 338 0.8 g CH₄ m⁻².

339 The site was an effective C and GHG source, accounting for 173.4 ± 1.7 g C m⁻² a⁻¹ and 1658.5 ± 11.2 g
 340 CO₂-Eq. m⁻² a⁻¹ for the EC source area (see Fig. 4). The proportion of CO₂ in the C and GWP budget was
 341 82.5 % and 31.6 %, respectively. Components of the annual net C balance other than CO₂ and CH₄
 342 fluxes, e.g. dissolved C, are not considered in this study. Our uncertainty estimates are within the range
 343 of similar studies (e.g. Shoemaker et al. 2015).

344 **3.3 Source area composition and spatial heterogeneity of CO₂ and CH₄** 345 **exchange**

346 Footprint analysis revealed the peak contribution in an average distance of 18 m from the tower and
 347 mainly from the open water area of the shallow lake (see Fig. 5). Open water covered on average 62.5
 348 % of the EC source area. The two surface types showed different emission rates in terms of higher CH₄
 349 fluxes and lower NEE rates with increasing Ω_{water} (see Fig. 6). Within the NLR_{CO₂foot} approach both
 350 surface types were denoted as sources of CO₂, but with about 4-fold stronger rates of GPP, R_{eco} and
 351 NEE for emergent vegetation compared to open water (see Fig. 7 and Table 3). The approach yielded
 352 a similar cumulative annual NEE for the whole EC source area including both surface types as the
 353 LUT_{CO₂nofoot} approach, but lower component fluxes (GPP and R_{eco}). As for CO₂, we implemented Ω_i as
 354 weighting factors within the NLR model for CH₄ (NLR_{CH₄foot}) to get the surface type specific fluxes of CH₄
 355 and fitted the parameters as follows:

$$356 F_{CH_4} = \Omega_{veg} \cdot \exp(-10.076 + 0.415 \cdot T_{s10}) + \Omega_{water} \cdot \exp(-6.449 + 0.286 \cdot T_{s10}) \quad (8)$$

357 Open water accounted for more than 4-fold higher emissions than the vegetated areas (see Fig. 7 and
 358 Table 3). The NLR_{CH₄foot} approach revealed a similar annual CH₄ budget as the NLR_{CH₄nofoot} approach.

359 Annual budgets of CO₂ (844 g CO₂ m⁻² a⁻¹) and CH₄ (22 g CH₄ m⁻² a⁻¹) for the AOI differed strongly from
 360 the budgets for the EC source area due to the contrasting emission rates of open water and emergent
 361 vegetation (see Table 3) and different fractional coverages of the surface types within the AOI and the
 362 EC source area. This resulted in a higher C loss (246.5 g C m⁻² a⁻¹) and a lower GWP (1452.9 g CO₂-Eq.
 363 m⁻² a⁻¹) than for the EC source area. In the following we will primarily discuss the budgets of the EC
 364 source area and the surface types.

365



366 4 Discussion

367 4.1 Diurnal variability of CH₄ emissions

368 In terms of its daily cycle, CH₄ exchange between wetland ecosystems and the atmosphere is not
369 generalisable, but rather dependent on the spatial characteristics of the wetland and thus, the impact
370 of the individual CH₄ emission pathways (diffusion, ebullition, plant-mediated transport). Our
371 measurements showed a diurnal cycle of CH₄ exchange from June to September 2013 and in March
372 2014, with the strongest emissions during night, as reported for shallow lakes (e.g. Podgrajsek et al.
373 2014) and wetland sites with a considerable fraction of open water (e.g. Godwin et al. 2013, Koebsch
374 et al. 2015). In comparison, wetland CH₄ emissions were also reported to show daily maxima at day-
375 time (e.g. Morrissey et al. 1993, Hendriks et al. 2010, Hatala Matthes et al. 2014), especially at sites with
376 high abundance of vascular plants. No diurnal pattern (e.g. Rinne et al. 2007, Forbrich et al. 2011,
377 Herbst et al. 2011) occurred especially at sites without large open water areas (Godwin et al. 2013).

378 We assume the process of convective mixing of the water column (e.g. Godwin et al. 2013, Poindexter
379 and Variano 2013, Podgrajsek et al. 2014, Sahlée et al. 2014, Koebsch et al. 2015) to be crucial for the
380 diurnal pattern of CH₄ emissions at our study site. This is indicated by the concurrent timing of
381 convective mixing and daily peak CH₄ emissions and a generally high fractional source area coverage
382 of the open water, which shows higher rates of CH₄ release than emergent vegetation. Furthermore,
383 closed chamber measurements likewise show night-time peak emissions on the shallow lake in
384 summer 2013 (Hoffmann et al. 2015). During the day, CH₄ is trapped in the lower (anoxic) layers of the
385 thermally stratified water column. Due to the heat release of the surface water to the atmosphere in
386 the night the surface water cools down, initiating convective mixing of the water column down to the
387 bottom. Diffusion is enhanced due to the buoyancy-induced turbulence, the associated increased gas
388 transfer velocity at the air-water interface (Eugster et al. 2003, MacIntyre et al. 2010, Podgrajsek et al.
389 2014) as well as the transport of CH₄ enriched bottom water to the surface (Godwin et al. 2013,
390 Podgrajsek et al. 2014). In addition, ebullition can be triggered by turbulence due to convective mixing
391 (Podgrajsek et al. 2014, Read et al. 2012). The daily pattern of the open water CH₄ release might
392 superimpose the reverse diurnal cycle of plant-mediated transport with peak emissions during day-
393 time, as the release of methane is dependent on the stomatal conductance of the plants (e.g. Morrissey
394 et al. 1993). This pathway is limited to plants with aerenchymatic tissue like *Typha latifolia*, which
395 dominates the eulittoral zone at our study site. CH₄ is transported from the soil to the atmosphere,
396 bypassing potential oxidation zones above the rhizosphere (chimney effect). Unusually for wetland



397 plants (Torn and Chapin 1993), complete stomatal closure during night was observed for *Typha latifolia*
398 (Chanton et al. 1993).

399 4.2 Annual CH₄ emissions

400 The CH₄ emissions of our studied ecosystem were within the range of other temperate fen sites
401 rewetted for several years (up to 63 g CH₄ m⁻² a⁻¹; e.g. Hendriks et al. 2007, Wilson et al. 2008, Günther
402 et al. 2013, Schrier-Uijl et al. 2014). This rate corresponds to twice the emission factor of 21.6 g CH₄ m⁻²
403 a⁻¹, that was assigned to rewetted temperate rich organic soils, which is in turn more than twice the
404 rate of the nutrient-poor complement (IPCC 2014). In contrast, newly rewetted fens emit its multiple.
405 In the first year after flooding, Hahn et al. (2015) observed an average net release of 260 g CH₄ m⁻² a⁻¹,
406 which is 186 times higher than before flooding, at a fen site in NE Germany. Two years later the CH₄
407 emissions were significantly lower (40 g CH₄ m⁻² per growing season; Koebisch et al. 2015). However,
408 natural fens release most often less CH₄ than the majority of rewetted fens (e.g. Bubier et al. 1993,
409 Nilsson et al. 2001), with some exceptions (e.g. Huttunen et al. 2003).

410 The two main surface types open water and emergent vegetation differed substantially in their CH₄
411 exchange rates. Open water contributed overproportionally to the measured ecosystem fluxes and
412 showed higher CH₄ release rates (52.6 g CH₄ m⁻² a⁻¹) than the emergent vegetation stands (13.2 g CH₄
413 m⁻² a⁻¹). However, closed-chamber measurements at the shallow lake show an even higher long-term
414 average annual CH₄ release rate (206 g CH₄ m⁻² a⁻¹) since rewetting with large interannual variability
415 and occasionally extreme high release rates (up to 400 g CH₄ m⁻² a⁻¹; Casares et al., in prep.).

416 We assume the permanent high inundation and high productivity due to eutrophic conditions, feeding
417 the organic mud deposited at the bottom of the open water body (which is typically for shallow lakes
418 in rewetted fens), to be of particular importance for high CH₄ emissions as substrate for
419 decomposition. The mud initially evolved as a mixture of sand and easily decomposable labile plant
420 litter from reed canary grass, which died-off after flooding and produced a large C pool for CH₄
421 production (Hahn-Schöfl et al 2011). During an incubation experiment with substrate from our study
422 site Hahn-Schöfl et al. (2011) observed that the new sediment layer has very high specific rates of
423 anaerobic CH₄ (and CO₂) production. In addition, Zak et al. (2015) emphasised the impact of litter
424 quality and reported a very high CH₄ production potential for litter of *Ceratophyllum demersum*, which
425 dominates the biomass in the open water at our study site. Due to the eutrophic character of the lake
426 and associated high productivity within the open water body and in the eulittoral zone, high amounts
427 of fresh labile organic matter continuously replenish the mud layer and thus the C pool. As the C
428 balance (CO₂ and CH₄) seems to be extremely unbalanced, we further assume lateral input of



429 allochthonous organic matter into the NE “bay” of the shallow lake, which is the area with the peak
430 contribution of our EC derived fluxes, especially during strong winds. The importance of fresh labile
431 organic matter provided by the die-back of the former vegetation as driving force for high CH₄
432 emissions was also discussed in Hahn et al. (2015). They measured the highest CH₄ emissions in sedge
433 stands suffering from strongest die-back.

434 For comparison annual budgets of CO₂ and CH₄ for other nutrient-rich lentic freshwater ecosystems in
435 terms of pristine, anthropogenically influenced and transient ecosystems are listed in Table 4. Studies
436 on nutrient-rich lakes generally revealed lower CH₄ release for open water. In contrast, beaver ponds
437 were partially reported to emit similar rates of CH₄. Similarly to our study site beaver ponds are at least
438 in the beginning disbalanced ecosystems due to a rapidly increased water level with associated
439 suffering and finally the die-back of former vegetation, which is not adapted to higher water levels. A
440 large C pool for CH₄ production develops. However, even for a beaver pond existing more than 30 years
441 CH₄ emissions still account for 40 g CH₄ m⁻² a⁻¹ (Yavitt et al. 1992).

442 Annual CH₄ emissions of the surface type emergent vegetation were about 4-fold lower than for open
443 water. This might be the result of increased CH₄ oxidation in the soil, as plants with aerenchymatic
444 tissue release oxygen into the rhizosphere, in reverse to the emission of CH₄ into the atmosphere
445 (Bhullar et al. 2013). Minke et al. (2015) highlight the difference in net CH₄ release for typical helophyte
446 stands with moderate emissions for *Typha* dominated sites. Besides the effect of the gas transport
447 within plants, lower water and sediment temperatures due to shading by the emergent vegetation
448 might yield lower CH₄ production than for open water. Furthermore, the soil of emergent vegetation
449 stands is generally only temporarily and partly inundated and the water table decreased additionally
450 during the unusual warm and dry summer 2013, probably resulting in a lower rate of anaerobic
451 decomposition to CH₄. This in turn might be a reason, that in comparison to other sites dominated by
452 *Typha* (rewetted wetlands, lake shores and freshwater marshes; see Table 4) the emergent vegetation
453 at our site is at the lower limit of reported CH₄ release rates and best comparable to closed chamber
454 measurements of *Typha latifolia* microsites at another rewetted fen site in NE Germany (Günther et
455 al. 2015).

456 **4.3 Annual net CO₂ release**

457 We observed high annual net release of CO₂ during the observation period, which is rather uncommon
458 for fens several years after rewetting (e.g. Hendriks et al. 2007, Schrier-Uijl et al. 2014, Knox et al.
459 2015). Surprisingly, net CO₂ budgets were similar to those of drained and degraded peatlands (e.g.
460 Hatala et al. 2012, Schrier-Uijl et al. 2014). Both surface types acted as net sources, with emergent



461 vegetation ($750 \text{ g CO}_2 \text{ m}^{-2} \text{ a}^{-1}$) showing a distinctively higher net budget as well as GPP and R_{eco} rates
462 than open water ($158 \text{ g CO}_2 \text{ m}^{-2} \text{ a}^{-1}$). Only few NEE rates are published for the open water body of
463 eutrophic shallow lakes. Ducharme-Riel et al. (2015) report $224 \text{ g CO}_2 \text{ m}^{-2} \text{ a}^{-1}$ as annual NEE of a
464 eutrophic lake in Canada (see Table 4). According to Kortelainen et al. (2006) Finnish lakes, which are
465 mainly small and shallow, continuously emit CO_2 during the ice-free period, positively correlated with
466 their trophic state.

467 Our study revealed a high annual net CO_2 release for emergent vegetation, which is in the wide range
468 of NEE rates for *Typha* sites reported in other studies, including both net CO_2 sources and sinks (see
469 Table 5). GPP and R_{eco} are generally high (especially at rewetted fen sites; both component fluxes most
470 often $> 3000 \text{ g CO}_2 \text{ m}^{-2} \text{ a}^{-1}$), characterising *Typha* stands as high turnover sites, usually resulting in net
471 CO_2 uptake. In contrast, R_{eco} and GPP rates at our study site are in the lower part of the reported range.
472 We assume the continuously high R_{eco} rates during winter 2013/2014, contributing to the high annual
473 net CO_2 emissions, to be the result of mild and dry meteorological conditions. In summer 2013, R_{eco}
474 exceeded GPP already in late June, indicating a significant contribution of heterotrophic respiration to
475 the CO_2 production. We cannot completely exclude a misestimation of the CO_2 exchange during
476 midsummer due to longer data gaps. However, unusual warm and dry conditions and associated water
477 table lowering during summer 2013 might have triggered a shift from anaerobic to aerobic
478 decomposition. This includes the exposed organic mud at former shallowly inundated soil of emergent
479 vegetation stands, e.g. at the edge of the lake. Besides CH_4 , Hahn-Schöfl et al. (2011) showed that the
480 new sediment layer at the bottom of inundated areas exhibits very high rates of anaerobic CO_2
481 production. The effect of water table lowering at *Typha* sites due to dry conditions is also shown by
482 Günther et al. (2015) and Chu et al. (2015): relative increase of R_{eco} rates, resulting in net CO_2 release.
483 This might be of special interest in terms of climate change, as a temperature increase and significantly
484 less precipitation in summer are expected for NE Germany. In addition, a considerable increase of
485 microbial activity and thus, generally increased decomposition due to high temperatures might be of
486 importance. Allochthonous organic matter import into the NE bay due to lateral transport, as discussed
487 for CH_4 , might have further enhanced decomposition (e.g. Chu et al. 2015).

488 **4.4 Global warming potential and the impact of spatial heterogeneity**

489 The lake ecosystem is characterised by a high GWP 9 years after rewetting, mainly driven by high CH_4
490 emissions. Based on our results the site can hardly be classified into any phase following peatland
491 rewetting discussed by Augustin and Joosten (2007). The slow development and shift of the ecosystem
492 to a C sink with reduced climate impact might be the result of the exceptional characteristics



493 represented by eutrophic conditions and lateral transport of organic matter within the open water
494 body. The trophic status of water and sediment is an important factor regulating GHG emissions, as
495 shown by Schrier-Uijl et al. (2011) for lakes and drainage ditches in wetlands. However, the unusual
496 meteorological conditions during our study period might have caused a comparable low GWP
497 compared to previous years due to lower CH₄ emissions at the expense of high net CO₂ release. In
498 comparison, e.g. Schrier-Uijl et al. (2014) report C uptake and a GHG sink function of a fen 10 years
499 after rewetting with water levels below or at the soil surface. In a study by Knox et al. (2015) a wetland
500 with mean water level above the soil surface was characterised by a near-neutral climate impact after
501 15 years of rewetting, where continued high CH₄ emissions were compensated by strong net CO₂
502 uptake. In the course of rewetting the water table is recommended to be held at or just below the soil
503 surface to prevent inundation and thus, the formation of organic mud (Couwenberg et al. 2011,
504 Joosten et al. 2012, Zak et al. 2015).

505 We calculated the “true” fluxes of CO₂ and CH₄ for the AOI by weighting the non-linear regression
506 functions for the two surface types with their fractional coverage inside the AOI. The inferred C budget
507 and global warming potential differs considerably from that of the EC source area, highlighting the
508 strikingly different emission rates of open water versus emergent vegetation. Thus, footprint analysis
509 providing the fractional coverage of the main surface types is imperative for the interpretation of
510 ecosystem flux measurements as provided by the EC technique at such a spatially heterogeneous site.
511 In addition, for an interannual comparison of EC derived budgets for such sites it is necessary to define
512 a fixed AOI, as the cumulative footprint climatology (representing the EC source area) changes
513 interannually. Inter-site comparisons (e.g. with other shallow lakes evolved during fen rewetting) are
514 challenging with regard to the site-specific spatial heterogeneity.

515

516 **5 Conclusions**

517 This study contributes to the understanding of eutrophic shallow lakes as a challenging ecosystem
518 often evolving during fen rewetting in NE Germany. Within the study period the ecosystem was a
519 strong source of CH₄ and CO₂. Both open water and emergent vegetation, particularly including the
520 eulittoral zone, were net emitters of CH₄ and CO₂, but with strikingly different release rates. This
521 illustrates the importance of footprint analysis for the interpretation of the EC measurements on a
522 rewetted site with distinct spatial heterogeneity. Our results show that the intended effects of
523 rewetting in terms of CO₂ emission reduction and C sink recovery are not yet achieved at this site. The
524 negative climate impact of the lake is dominated by considerable CH₄ release, particularly from the
525 open water section. In combination with the high net CO₂ release the C budget seems to be extremely



526 unbalanced. Measurements of lateral transport of organic substrate within the open water body and
527 a full C budget could give indication on a potential allochthonous input into the NE bay. Furthermore,
528 the effect of unusual meteorological conditions need further investigation. A comparison with existing
529 chamber measurements at the open water body for the same time period will be helpful for the
530 evaluation of our measurements and estimation for the surface type fluxes. The site is continuously
531 changing, with *Typha latifolia* progressively entering the open water body in the course of
532 terrestrialisation, probably resulting in peat formation and C uptake once the shallow lake is
533 replenished by organic sediments. Therefore, long-term measurements are necessary to evaluate the
534 impact of future ecosystem development on GHG emissions. Moreover, statements for the climate
535 impact of rewetted fens can only be provided by inclusion of additional sites varying in groundwater
536 table and vegetation type. We assume that shallow lakes represent a special case with regard to the
537 GHG dynamics and climate impact, with exceptionally high CH₄ release and occasionally high net CO₂
538 emissions. Inundation involves the risk of unpredictable and long-term high CH₄ emissions, especially
539 in case of nutrient-rich conditions, that counteract the actually intended lowering of the climate impact
540 of drained and degraded fens. We strongly recommend to consider this risk in future rewetting
541 projects and support the call of Lamers et al. (2015) for the need of well-conceived restoration
542 management instead of the trial-and-error approach, whereon restoration of wetland ecosystem
543 services was based on for a long time.

544

545 **Acknowledgements**

546 This work was supported by the Helmholtz Association of German Research Centres through a
547 Helmholtz Young Investigators Group to T.S. (grant VH-NG-821) and is a contribution to the Helmholtz
548 Climate Initiative REKLIM (Regional Climate Change). Infrastructure funding through the Terrestrial
549 Environmental Observatories Network (TERENO) is also acknowledged. We thank M. Hoffmann (ZALF,
550 Müncheberg, Germany) and C. Hohmann (GFZ Potsdam, Germany) for providing meteorological data.
551



552 **References**

- 553 Aubinet, M., Feigenwinter, C., Heinesch, B., Laffineur, Q., Papale, D., Reichstein, M., Rinne, J. and Van
554 Gorsel, E.: Nighttime flux correction, in: Eddy Covariance: A practical guide to measurement and
555 data analysis, Springer Atmospheric Sciences, edited by Aubinet, M., Vesala, T. and Papale, D.,
556 Springer Netherlands, Dordrecht, doi:10.1007/978-94-007-2351-1, 133-158, 2012.
- 557 Augustin, J. and Joosten, H.: Peatland rewetting and the greenhouse effect, International Mire
558 Conservation Group Newsletter, 3, 29-30, 2007.
- 559 Barcza, Z., Kern, A., Haszpra, L. and Kljun, N.: Spatial representativeness of tall tower eddy covariance
560 measurements using remote sensing and footprint analysis, Agr. Forest Meteorol., 149, 795-807,
561 doi: 10.1016/j.agrformet.2008.10.021, 2009.
- 562 Bastviken, D., Cole, J., Pace, M. and Tranvik, L.: Methane emissions from lakes: Dependence of lake
563 characteristics, two regional assessments, and a global estimate, Global Biogeochem. Cy., 18,
564 GB4009, doi: 10.1029/2004GB002238, 2004.
- 565 Beetz, S., Liebersbach, H., Glatzel, S., Jurasinski, G., Buczko, U. and Höper, H.: Effects of land use
566 intensity on the full greenhouse gas balance in an Atlantic peat bog, Biogeosciences, 10, 1067-
567 1082, doi:10.5194/bg-10-1067-2013, 2013.
- 568 Berg, E., Jeschke, L. and Lenschow, U.: Das Moorschutzkonzept Mecklenburg-Vorpommern, Telma, 30,
569 173-220, 2000.
- 570 Béziat, P., Ceschia, E. and Dedieu, G.: Carbon balance of a three crop succession over two cropland
571 sites in South West France, Agr. Forest Meteorol., 149, 1628-1645,
572 doi:10.1016/j.agrformet.2009.05.004, 2009.
- 573 Bhullar, G. S., Edwards, P. J. and Olde Venterink, H.: Variation in the plant-mediated methane transport
574 and its importance for methane emission from intact wetland peat mesocosms, J. Plant Ecol., 6,
575 298-304, doi: 10.1093/jpe/rts045, 2013.
- 576 Bignell, N.: The effect of dissolved air on the density of water, Metrologia, 19, 57-59, 1983.
- 577 Bonneville, M.-C., Strachan, I. B., Humphreys, E. R. and Roulet, N. T.: Net ecosystem CO₂ exchange in a
578 temperate cattail marsh in relation to biophysical properties, Agr. Forest Meteorol., 148, 69-81,
579 doi:10.1016/j.agrformet.2007.09.004, 2008.
- 580 Bubier, J. L., Moore, T. R. and Roulet, N. T.: Methane emissions from wetlands in the Midboreal Region
581 of Northern Ontario, Canada, Ecology, 74, 2240-2254, 1993.



- 582 Casper, P., Maberly, S. C., Hall, G. H. and Finlay, B. J.: Fluxes of methane and carbon dioxide from a
583 small productive lake to the atmosphere, *Biogeochemistry*, 49, 1-19, doi:
584 10.1023/A:1006269900174, 2000.
- 585 Chanton, J. P., Whiting, G. J., Happell, J. D. and Gerard, G.: Contrasting rates and diurnal patterns of
586 methane emission from different types of vegetation, *Aquat. Bot.*, 46, 111-128, 1993.
- 587 Chen, B., Coops, N. C., Fu, D., Margolis, H. A., Amiro, B. D., Barr, A. G., Black, T. A., Arain, M. A., Bourque,
588 C. P.-A., Flanagan, L. B., Lafleur, P. M., McCaughey, J. H. and Wofsy, S. C.: Assessing eddy-
589 covariance flux tower location bias across the Fluxnet-Canada Research Network based on remote
590 sensing and footprint modelling, *Agr. Forest Meteorol.*, 151, 87–100,
591 doi:10.1016/j.agrformet.2010.09.005, 2011.
- 592 Chu, H., Gottgens, J. F., Chen, J., Sun, G., Desai, A. R., Ouyang, Z., Shao, C. and Czajkowski, K.: Climatic
593 variability, hydrologic anomaly, and methane emission can turn productive freshwater marshes
594 into net carbon sources, *Glob. Change Biol.*, 21, 1165-1181, doi: 10.1111/gcb.12760, 2015.
- 595 Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Bärtsch, S., Dubovik, D., Liashchynskaya, N.,
596 Michaelis, D., Minke, M., Skuratovich, A. and Joosten, H.: Assessing greenhouse gas emissions
597 from peatlands using vegetation as a proxy, *Hydrobiologia*, 674, 67-89, doi:10.1007/s10750-011-
598 0729-x, 2011.
- 599 Dengel, S., Levy, P. E., Grace, J., Jones, S. K. and Skiba, U. M.: Methane emissions from sheep pasture,
600 measured with an open-path eddy covariance system, *Glob. Change Biol.*, 17, 3524-3533, doi:
601 10.1111/j.1365-2486.2011.02466.x, 2011.
- 602 Ducharme-Riel, V., Vachon, D., del Giorgio, P. A. and Prairie, Y. T.: The relative contribution of winter
603 under-ice and summer hypolimnetic CO₂ accumulation to the annual CO₂ emissions from
604 northern lakes, *Ecosystems*, 18, 547-559, doi:10.1007/s10021-015-9846-0, 2015.
- 605 Eugster, W., Kling, G., Jonas, T., McFadden, J. P., Wüest, A., MacIntyre, S. and Chapin III, F. S.: CO₂
606 exchange between air and water in an Arctic Alaskan and midlatitude Swiss lake: importance of
607 convective mixing, *J. Geophys. Res.* 108, 4362, doi: 10.1029/2002JD002653, 2003.
- 608 Falge, E., Baldocchi, D., Olson, R. J., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G., Ceulemans, R.,
609 Clement, R., Dolman, H., Granier, A., Gross, P., Grünwald, T., Hollinger, D., Jensen, N.-O., Katul, G.,
610 Keronen, P., Kowalski, A., Ta Lai, C., Law, B. E., Meyers, T., Moncrieff, J., Moors, E., Munger, J. W.,
611 Pilegaard, K., Rannik, Ü., Rebmann, C., Suyker, A., Tenhunen, J., Tu, K., Verma, S., Vesala, T.,



- 612 Wilson, K. and Wofsy, S. C.: Gap filling strategies for defensible annual sums of net ecosystem
613 exchange, *Agr. Forest Meteorol.* 107, 43–69, doi:10.1016/S0168-1923(00)00225-2, 2001.
- 614 Fan, S.-M., Wofsy, S. C., Bakwin, P. S., Jacob, D. J. and Fitzjarrald, D. R.: Atmosphere-biosphere
615 exchange of CO₂ and O₃ in the Central Amazon Forest, *J. Geophys. Res.*, 95, 16851-16864, 1990.
- 616 Finkelstein, P. L. and Sims, P. F.: Sampling error in eddy correlation flux measurements, *J. Geophys.*
617 *Res.-Atmos.*, 106, 3503-3509, doi:10.1029/2000JD900731, 2001.
- 618 Finnigan, J. J., Clement, R., Mahli, Y., Leuning, R. and Cleugh, H. A.: A re-evaluation of long-term flux
619 measurement techniques. Part I: Averaging and coordinate rotation, *Bound.-Lay. Meteorol.*, 107,
620 1-48, doi:10.1023/A:1021554900225, 2003.
- 621 Foken, T.: *Micrometeorology*. Springer, Berlin, 308 pp, doi:10.1007/978-3-540-74666-9, 2008.
- 622 Forbrich, I., Kutzbach, L., Wille, C., Becker, T., Wu, J. and Wilmking, M.: Cross-evaluation of
623 measurements of peatland methane emissions on microform and ecosystem scales using high-
624 resolution landcover classification and source weight modelling, *Agr. Forest Meteorol.*, 151, 864-
625 874, doi:10.1016/j.agrformet.2011.02.006, 2011.
- 626 Fratini, F., Ibrom, A., Arriga, N., Burba, G. and Papale, D.: Relative humidity effects of water vapour
627 fluxes measured with closed-path eddy-covariance systems with short sampling lines, *Agr. Forest*
628 *Meteorol.*, 165, 53-63, 2012.
- 629 Frohling, S., Talbot, J., Jones, M. C., Treat, C. C., Kauffman, J. B., Tuittila, E.-S. and Roulet, N. T.: Peatlands
630 in the earth's 21st century climate system, *Environ. Rev.*, 19, 371-396, doi:10.1139/a11-014, 2011.
- 631 Göckede, M., Rebmann, C. and Foken, T.: A combination of quality assessment tools for eddy
632 covariance measurements with footprint modelling for the characterisation of complex sites, *Agr.*
633 *Forest Meteorol.*, 127, 175-188, doi:10.1016/j.agrformet.2004.07.012, 2004.
- 634 Godwin, C. M., McNamara, P. J. and Markfort, C. D.: Evening methane emission pulses from a boreal
635 wetland correspond to convective mixing in hollows, *J. Geophys. Res.-Biogeo.*, 118, 994-1005, doi:
636 10.1002/jgrg.20082, 2013.
- 637 Günther, A. B., Huth, V., Jurasinski, G. and Glatzel, S.: Scale-dependent temporal variation in
638 determining the methane balance of a temperate fen, *Greenhouse Gas Measurement and*
639 *Management*, 4, 41-48, doi: 10.1080/20430779.2013.850395, 2013.



- 640 Günther, A. B., Huth, V., Jurasinski, G. and Glatzel, S.: The effect of biomass harvesting on greenhouse
641 gas emissions from a rewetted temperate fen, *GCB Bioenergy*, 7, 1092-1106, doi:
642 10.1111/gcbb.12214, 2015.
- 643 Hahn, J., Köhler, S., Glatzel, S. and Jurasinski, G.: Methane exchange in a coastal fen in the first year
644 after flooding – A systems shift, *PLoS ONE*, 10, e0140657, doi:10.1371/journal.pone.0140657,
645 2015.
- 646 Hahn-Schöfl, M., Zak, D., Minke, M., Gelbrecht, J., Augustin, J. and Freibauer, A.: Organic sediment
647 formed during inundation of a degraded fen grassland emits large fluxes of CH₄ and CO₂,
648 *Biogeosciences*, 8, 1539-1550, doi:10.5194/bg-8-1539-2011, 2011.
- 649 Hatala, J. A., Detto, M., Sonnentag, O., Deverel, S. J., Verfaillie, J. and Baldocchi, D. D.: Greenhouse gas
650 (CO₂, CH₄, H₂O) fluxes from drained and flooded agricultural peatlands in the Sacramento-San
651 Joaquin Delta, *Agr. Ecosyst. Environ.*, 150, 1-18, doi:10.1016/j.agee.2012.01.009, 2012.
- 652 Hatala Matthes, J., Sturtevant, C., Verfaillie, J., Knox, S. and Baldocchi, D.: Parsing the variability in CH₄
653 flux at a spatially heterogeneous wetland: Integrating multiple eddy covariance towers with high-
654 resolution flux footprint analysis, *J. Geophys. Res.-Biogeo.*, 119, 1322-1339,
655 doi:10.1002/2014JG002642 2014.
- 656 Hendriks, D. M. D., van Huissteden, J., Dolman, A. J. and van der Molen, M. K.: The full greenhouse gas
657 balance of an abandoned peat meadow, *Biogeosciences*, 4, 411-424, doi:10.5194/bg-4-411-2007,
658 2007.
- 659 Hendriks, D. M. D., van Huissteden, J. and Dolman, A. J.: Multi-technique assessment of spatial and
660 temporal variability of methane fluxes in a peat meadow, *Agr. Forest Meteorol.*, 150, 757-774,
661 doi:10.1016/j.agrformet.2009.06.017, 2010.
- 662 Herbst, M., Friborg, T., Ringgaard, R. and Soegaard, H.: Interpreting the variations in atmospheric
663 methane fluxes observed above a restored wetland, *Agr. Forest Meteorol.*, 151, 841-853,
664 doi:10.1016/j.agrformet.2011.02.002, 2011.
- 665 Hoffmann, M., Schulz-Hanke, M., Garcia Alba, J., Jurisch, N., Hagemann, U., Sachs, T., Sommer, M. and
666 Augustin, J.: Technical Note: A simple calculation algorithm to separate high-resolution CH₄ flux
667 measurements into ebullition and diffusion-derived components, *Biogeosciences Discussions*, 12,
668 12923-12945, doi:10.5194/bgd-12-12923-2015, 2015.



- 669 Holden, J., Chapman, P. J. and Labadz, J. C.: Artificial drainage of peatlands: hydrological and
670 hydrochemical process and wetland restoration, *Prog. Phys. Geog.*, 28, 95-123,
671 doi:10.1191/0309133304pp403ra, 2004.
- 672 Hommeltenberg, J., Mauder, M., Drösler, M., Heidbach, K., Werle, P. and Schmid, H. P.: Ecosystem
673 scale methane fluxes in a natural temperate bog-pineforest in southern Germany, *Agr. Forest
674 Meteorol.*, 198-199, 273-284, doi:10.1016/j.agrformet.2014.08.017, 2014.
- 675 Höper, H., Augustin, J., Cagampan, J. P., Drösler, M., Lundin, L., Moors, E., Vasander, H., Waddington,
676 J. M., and Wilson, D.: Restoration of peatlands and greenhouse gas balances, in: *Peatlands and
677 climate change*, edited by Strack, M., International Peat Society, Jyväskylä, 182-210, 2008.
- 678 Horst, T. W. and Lenschow, D. H.: Attenuation of scalar fluxes measured with spatially-displaced
679 sensors, *Bound.-Lay. Meteorol.*, 130, 275-300, doi:10.1007/s10546-008-9348-0, 2009.
- 680 Huttunen, J. T., Nykänen, H., Turunen, J. and Martikainen, P. J.: Methane emissions from natural
681 peatlands in the northern boreal zone in Finland, *Fennoscandia, Atmos. Environ.*, 37, 147-151,
682 doi:10.1016/S1352-2310(02)00771-9, 2003.
- 683 IPCC: *Climate Change 2013: The physical science basis. Contribution of Working Group I to the Fifth
684 Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F.,
685 Qin, D., Plattner, G.-K., Tignor, M. M. B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and
686 Midgley, P. M., Cambridge University Press, Cambridge, New York, 1535 pp,
687 doi:10.1017/CBO9781107415324, 2013.
- 688 IPCC: *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories:
689 Wetlands*, edited by Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M.
690 and Troxler, T.G., IPCC, Switzerland, 354 pp, 2014.
- 691 Itzerott, S.: TERENO (Northeast), Climate station Karlshof, Germany. Deutsches
692 GeoForschungsZentrum GFZ. <http://dx.doi.org/10.5880/TERENO.265>, 2015.
- 693 Joosten, H.: The global peatland CO₂ picture – Peatland status and drainage related emissions in all
694 countries of the world, *Wetlands International*, Ede, 36 pp, 2010.
- 695 Joosten, H., Tapio-Biström, M.-L. and Tol, S. (Eds.): *Peatlands – guidance for climate change mitigation
696 through conservation, rehabilitation and sustainable use*, 2nd edition, Food and Agriculture
697 Organization of the United Nations and Wetlands International, 100 pp, 2012.



- 698 Juutinen, S., Alm, J., Larmola, T., Huttunen, J. T., Morero, M., Martikainen, P. J. and Silvola, J.: Major
699 implication of the littoral zone for methane release from boreal lakes, *Global Biogeochem. Cy.*,
700 17, 1117, doi:10.1029/2003GB002105, 2003.
- 701 Kankaala, P., Ojala, A. and Käki, T.: Temporal and spatial variation in methane emissions from a flooded
702 transgression shore of a boreal lake, *Biogeochemistry*, 68, 297-311, doi:
703 10.1023/B:BI0G.0000031030.77498.1f, 2004.
- 704 Knox, S. H., Sturtevant, C., Hatala Matthes, J., Koteen, L., Verfaillie, J. and Baldocchi, D.: Agricultural
705 peatland restoration: effects of land-use change on greenhouse gas (CO₂ and CH₄) fluxes in the
706 Sacramento-San Joaquin Delta, *Glob. Change Biol.*, 21, 750-765, doi:10.1111/gcb.12745, 2015.
- 707 Koesch, F., Jurasinski, G., Koch, M., Hofmann, J. and Glatzel, S.: Controls for multi-scale temporal
708 variation in ecosystem methane exchange during the growing season of a permanently inundated
709 fen, *Agr. Forest Meteorol.*, 204, 94-105, doi:10.1016/j.agrformet.2015.02.002, 2015.
- 710 Kormann, R. and Meixner, F. X.: An analytical footprint model for non-neutral stratification, *Bound.-*
711 *Lay. Meteorol.*, 99, 207-224, doi:10.1023/A:1018991015119, 2001.
- 712 Kortelainen, P., Rantakari, M., Huttunen, J. T., Mattsson, T., Alm, J., Juutinen, S., Larmola, T., Silvola, J.
713 and Martikainen, P. J.: Sediment respiration and lake trophic state are important predictors of
714 large CO₂ evasion from small boreal lakes, *Glob. Change Biol.*, 12, 1554-1567, doi: 10.1111/j.1365-
715 2486.2006.01167.x. 2006.
- 716 Lamers, L. P. M., Vile, M. A., Grootjans, A. P., Acreman, M. C., van Diggelen, R., Evans, M. G., Richardson,
717 C. J., Rochefort, L., Koojiman, A. M., Roelofs, J. G. M. and Smolders, A. J. P. (2015): Ecological
718 restoration of rich fens in Europe and North America: from trial and error to an evidence-based
719 approach, *Biol. Rev.*, 90, 182-203, doi: 10.1111/brv.12102, 2015.
- 720 Lloyd, J. and Taylor, J. A.: On the temperature dependence of soil respiration, *Funct. Ecol.*, 8, 315-323,
721 doi:10.2307/2389824, 1994.
- 722 Lund, M., Lafleur, P. M., Roulet, N. T., Lindroth, A., Christensen, T. R., Aurela, M., Chojnicki, B. H.,
723 Flanagan, L. B., Humphreys, E. R., Laurila, T., Oechel, W., Olejnik, J., Rinne, J., Schubert, P. and
724 Nilsson, M. B.: Variability in exchange of CO₂ across 12 northern peatland and tundra sites, *Glob.*
725 *Change Biol.*, 16, 2436-2448, doi:10.1111/j.1365-2486.2009.02104.x, 2010.
- 726 MacIntyre, S., Jonsson, A., Jansson, M., Aberg, J., Turney, D. E. and Miller, S. D.: Buoyancy flux,
727 turbulence, and the gas transfer coefficient in a stratified lake, *Geophys. Res. Lett.*, 37, L24604,
728 doi:10.1029/2010GL044164, 2010.



- 729 Mauder, M. and Foken, T.: Documentation and instruction manual of the Eddy covariance software
730 package TK2. Arbeitsergebnisse Nr. 26, University of Bayreuth, Department of Micrometeorology,
731 42 pp, 2004.
- 732 McDermitt, D., Burba, G., Xu, L., Anderson, T., Komissarov, A., Riensche, B., Schedlbauer, J., Starr, G.,
733 Zona, D., Oechel, W., Oberbauer, S. and Hastings, S.: A new low-power, open-path instrument for
734 measuring methane flux by eddy covariance, *Appl. Phys. B*, 102, 391-405, doi:10.1007/s00340-
735 010-4307-0, 2011.
- 736 Meyer, K., Höper, H. and Blankenburg, J.: Spurengashaushalt und Klimabilanz von Niedermooren unter
737 dem Einfluß des Vernässungsmanagements, in: *Ökosystemmanagement für Niedermoore.*
738 *Strategien und Verfahren zur Renaturierung*, edited by Kratz, R. and Pfadenhauer, J., Ulmer,
739 Stuttgart, 104-111, 2001.
- 740 Minke, M., Augustin, J., Burlo, A., Yarmashuk, T., Chuvashova, H., Thiele, A., Freibauer, A., Tikhonov, V.
741 and Hoffmann, M.: Water level, vegetation composition and plant productivity explain
742 greenhouse gas fluxes in temperate cutover fens after inundation, *Biogeosciences Discussions*,
743 12, 17393-17452, doi:10.5194/bgd-12-17393-2015, 2015.
- 744 Moffat, A. M., Papale, D., Reichstein, M., Hollinger, D. Y., Richardson, A. D., Barr, A. G., Beckstein, C.,
745 Braswell, B. H., Churkina, G., Desai, A. R., Falge, E., Gove, J. H., Heimann, M., Hui, D., Jarvis, A. J.,
746 Kattge, J., Noormets, A. and Stauch, V. J.: Comprehensive comparison of gap-filling techniques for
747 eddy covariance net carbon fluxes, *Agr. Forest Meteorol.*, 147, 209-232,
748 doi:10.1016/j.agrformet.2007.08.011, 2007.
- 749 Moncrieff, J. B., Clement, R., Finnigan, J. and Meyers, T.: Averaging, detrending and filtering of eddy
750 covariance time series, in: *Handbook of micrometeorology: a guide for surface flux*
751 *measurements*, edited by Lee, X., Massman, W. J. and Law, B. E., Kluwer Academic, Dordrecht, 7-
752 31, doi:10.1007/1-4020-2265-4, 2004.
- 753 Morrissey, L. A., Zobel, D. B. and Livingston, G. P.: Significance of stomatal control on methane release
754 from *Carex*-dominated wetlands, *Chemosphere*, 26, 339-355, doi:10.1016/0045-6535(93)90430-
755 D, 1993.
- 756 Naiman, R. J., Manning, T. and Johnston, C. A.: Beaver population fluctuations and tropospheric
757 methane emissions in boreal wetlands, *Biogeochemistry*, 12, 1-15, doi: 10.1007/BF00002623,
758 1991.



- 759 Nilsson, M., Mikkilä, C., Sundh, I., Granberg, G., Svensson, B. H. and Ranneby, B.: Methane emission
 760 from Swedish mires: National and regional budgets and dependence on mire vegetation, *J.*
 761 *Geophys. Res.*, 106, 20847-20860, doi:10.1029/2001JD900119, 2001.
- 762 Petrescu, A. M. R., Lohila, A., Tuovinen, J.-P., Baldocchi, D. D., Desai, A. R., Roulet, N. T., Vesala, T.,
 763 Dolman, A. J., Oechel, W. C., Marcolla, B., Friborg, T., Rinne, J., Hatala Matthes, J., Merbold, L.,
 764 Meijide, A., Kiely, G., Sottocornola, M., Sachs, T., Zona, D., Varlagin, A., Lai, D. Y. F., Veenendaal,
 765 E., Parmentier, F.-J. W., Skiba, U., Lund, M., Hensen, A., van Huissteden, J., Flanagan, L. B.,
 766 Shurpali, N. J., Grünwald, T., Humphreys, E. R., Jackowicz-Korczyński, M., Aurela, M. A., Laurila, T.,
 767 Grüning, C., Corradi, C. A. R., Schrier-Uijl, A. P., Christensen, T. R., Tamstorf, M. P., Mastepanov,
 768 M., Martikainen, P. J., Verma, S. B., Bernhofer, C. and Cescatti, A.: The uncertain climate footprint
 769 of wetlands under human pressure, *PNAS*, 112, 4594-4599, doi:10.1073/pnas.1416267112, 2015.
- 770 Podgrajsek, E., Sahlée, E. and Rutgersson, A.: Diurnal cycle of lake methane flux, *J. Geophys. Res.-*
 771 *Biogeo.*, 119, 236-248, doi:10.1002/2013JG002327, 2014.
- 772 Poindexter, C. M. and Variano, E. A.: Gas exchange in wetlands with emergent vegetation: the effects
 773 of wind and thermal convection at the air–water interface, *J. Geophys. Res.-Biogeo.*, 118, 1297–
 774 1306, doi:10.1002/jgrg.20099, 2013.
- 775 R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical
 776 Computing, Vienna, Austria, 2012.
- 777 Read, J. S., Hamilton, D. P., Desai, A. R., Rose, K. C., MacIntyre, S., Lenters, J. D., Smyth, R. L., Hanson,
 778 P. C., Cole, J. J., Staehr, P. A., Rusak, J. A., Pierson, D. C., Brookes, J. D., Laas, A. and Wu, C. H.: Lake-
 779 size dependency of wind shear and convection as controls on gas exchange, *Geophys. Res. Lett.*
 780 39, L09405, doi:10.1029/2012GL051886, 2012.
- 781 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann,
 782 N., Gilmanov, T., Granier, A., Grünwald, T., Havránková, K., Ilvesniemi, H., Janous, D., Knohl, A.,
 783 Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.-M.,
 784 Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T.,
 785 Yakir, D. and Valentini, R.: On the separation of net ecosystem exchange into assimilation and
 786 ecosystem respiration: review and improved algorithm, *Glob. Change Biol.* 11, 1424-1439,
 787 doi:10.1111/j.1365-2486.2005.001002.x, 2005.
- 788 Repo, M. E., Huttunen, J. T., Naumov, A. V., Chichulin, A. V., Lapshina, E. D., Bleuten, W. and
 789 Martikainen, P. J.: Release of CO₂ and CH₄ from small wetland lakes in western Siberia, *Tellus* 59B,
 790 788-796, doi:10.1111/j.1600-0889.2007.00301.x, 2007.



- 791 Rinne, J., Riutta, T., Pihlatie, M., Aurela, M., Haapanala, S., Tuovinen, J.-P., Tuittila, E.-S. and Vesala, T.:
792 Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique,
793 Tellus 59B, 449-457, doi:10.1111/j.1600-0889.2007.00261.x, 2007.
- 794 Rocha, A. V. and Goulden, M. L.: Large interannual CO₂ and energy exchange variability in a freshwater
795 marsh under consistent environmental conditions, J. Geophys. Res., 113, G04019,
796 doi:10.1029/2008JG000712, 2008.
- 797 Rõõm, E.-I., Nõges, P., Feldmann, T., Tuvikene, L., Kisand, A., Teearu, H. and Nõges, T.: Years are not
798 brothers: Two-year comparison of greenhouse gas fluxes in large shallow Lake Võrtsjärv, Estonia,
799 J. Hydrol., 519, 1594-1606, doi:10.1016/j.jhydrol.2014.09.011, 2014.
- 800 Roulet, N. T., Ash, R. and Moore, T. R.: Low boreal wetlands as a source of atmospheric methane, J.
801 Geophys. Res., 97, 3739-3749, 1992.
- 802 Roulet, N.T., Ash, R., Quinton, W. and Moore, T. R.: Methane flux from drained northern peatlands:
803 Effect of a persistent water table lowering on flux, Global Biogeochem. Cy., 7, 749-769,
804 doi:10.1029/93GB01931, 1993.
- 805 Sahlée, E., Rutgersson, A., Podgrajsek, E. and Bergström, H.: Influence from surrounding land on the
806 turbulence measurements above a lake, Bound.-Lay. Meteorol., 150, 235-258,
807 doi:10.1007/s10546-013-9868-0, 2014.
- 808 Schrier-Uijl, A. P., Veraart, A. J., Leffelaar, P. A., Berendse, F. and Veenendaal, E. M.: Release of CO₂ and
809 CH₄ from lakes and drainage ditches in temperate wetlands, Biogeochemistry, 102, 265-279,
810 doi:10.1007/s10533-010-9440-7, 2011.
- 811 Schrier-Uijl, A. P., Kroon, P. S., Hendriks, D. M. D., Hensen, A., van Huissteden, J., Berendse, F. and
812 Veenendaal, E. M.: Agricultural peatlands: towards a greenhouse gas sink – a synthesis of a
813 Dutch landscape study, Biogeosciences, 11, 4559-4576, doi:10.5194/bg-11-4559-2014, 2014.
- 814 Shoemaker, W. B., Anderson, F., Barr, J. G., Graham, S. L. and Botkin, D. B.: Carbon exchange between
815 the atmosphere and subtropical forested cypress and pine wetlands, Biogeosciences, 12, 2285-
816 2300, doi:10.5194/bg-12-2285-2015, 2015.
- 817 Siebicke, L., Hunner, M. and Foken, T.: Aspects of CO₂ advection measurements, Theor. Appl. Climatol.,
818 109, 109-131, doi:10.1007/s00704-011-0552-3, 2012.
- 819 Silvius, M., Joosten, H. and Opdam, S.: Peatlands and people, in: Assessment on peatlands, biodiversity
820 and climate change: Main Report, edited by Parish, F., Sirin, A., Charman, D., Joosten, H.,



- 821 Minayeva, T., Silvius, M. and Stringer, L., Global Environment Centre, Kuala Lumpur, and Wetlands
822 International, Wageningen, 20-38, 2008.
- 823 Smith, L. K. and Lewis Jr., W. M.: Seasonality of methane emissions from five lakes and associated
824 wetlands of the Colorado Rockies, *Global Biogeochem. Cy.*, 6, 323-338, 1992.
- 825 Steffenhagen, P., Zak, D., Schulz, K., Timmermann, T. and Zerbe, S.: Biomass and nutrient stock of
826 submersed and floating macrophytes in shallow lakes formed by rewetting of degraded fens,
827 *Hydrobiologia*, 692, 99-109, doi:10.1007/s10750-011-0833-y, 2012.
- 828 Strachan, I. B., Nugent, K. A., Crombie, S. and Bonneville, M.-C.: Carbon dioxide and methane exchange
829 at a cool-temperate freshwater marsh, *Environ. Res. Lett.*, 10, 065006, doi:10.1088/1748-
830 9326/10/6/065006, 2015.
- 831 Succow, M.: Durchströmungsmoore, in: *Landschaftsökologische Moorkunde*, edited by Succow, M.
832 and Joosten, H., Schweizerbart, Stuttgart, 463-470, 2001.
- 833 Torn, M. S. and Chapin III, F. S.: Environmental and biotic controls over methane flux from arctic tundra,
834 *Chemosphere*, 26, 357-368, doi:10.1016/0045-6535(93)90431-4, 1993.
- 835 van Dijk, A., Moene, A. F. and de Bruin, H. A. R.: The principles of surface flux physics: Theory, practice
836 and description of the ECPack library, Meteorology and Air Quality Group, Wageningen University,
837 Wageningen, 98 pp, 2004.
- 838 Vickers, D. and Mahrt, L.: Quality control and flux sampling problems for tower and aircraft data, *J.*
839 *Atmos. Ocean. Tech.*, 14, 512-526, 1997.
- 840 Waddington, J. M. and Day, S. M.: Methane emissions from a peatland following restoration, *J.*
841 *Geophys. Res.*, 112, G03018, doi:10.1029/2007JG000400, 2007.
- 842 Wagner, W. and Pruß, A.: The IAPWS formulation 1995 for the thermodynamic properties of ordinary
843 water substance for general and scientific use, *J. Phys. Chem. Ref. Data*, 31, 387-535, 2002.
- 844 Wang, H., Lu, J., Wang, W., Yang, L. and Yin, C.: Methane fluxes from the littoral zone of hyper eutrophic
845 Taihu Lake China, *J. Geophys. Res.*, 111, D17109, doi:10.1029/2005JD006864, 2006.
- 846 Webb, E. K., Pearman, G. I. and Leuning, R.: Correction of flux measurements for density effects due
847 to heat and water vapor transfer, *Q. J. Roy. Meteor. Soc.*, 106, 85-100,
848 doi:10.1002/qj.49710644707, 1980.
- 849 Whiting, G. J. and Chanton, J. P.: Greenhouse carbon balance of wetlands: methane emission versus
850 carbon sequestration, *Tellus*, 53B, 521-528, doi: 10.1034/j.1600-0889.2001.530501.x, 2001.



- 851 Wilczak, J. M., Oncley, S. P. and Stage, S. A.: Sonic anemometer tilt correction algorithms, Bound.-Lay.
852 Meteorol., 99, 127-150, doi:10.1023/A:1018966204465, 2001.
- 853 Wilson, D., Tuittila, E.-S., Alm, J., Laine, J., Farrell, E. P. and Byrne, K. A.: Carbon dioxide dynamics of a
854 restored maritime peatland, Ecoscience 14, 71-80, doi: 10.2980/1195-
855 6860(2007)14[71:CDDOAR]2.0.CO;2, 2007.
- 856 Wilson, D., Alm, J., Laine, J., Byrne, K. A., Farrell, E. P. and Tuittila, E.-S.: Rewetting of cutaway
857 peatlands: are we re-creating hot plots of methane emissions?, Restor. Ecol., 17, 796-806,
858 doi:10.1111/j.1526-100X.2008.00416.x, 2008.
- 859 Yavitt, J. B., Angell, L. L., Fahey, T. J., Cirimo, C. P. and Driscoll, C. T.: Methane fluxes, concentrations,
860 and production in two Adirondack beaver impoundments, Limnol. Oceanogr., 37, 1057-1066,
861 1992.
- 862 Zak, D., Gelbrecht, J., Wagner, C. and Steinberg, C. E. W.: Evaluation of phosphorus mobilisation
863 potential in rewetted fens by an improved sequential chemical extraction procedure, Eur. J. Soil.
864 Sci., 59, 1191-1201, doi:10.1111/j.1365-2389.2008.01081.x, 2008.
- 865 Zak, D., Reuter, H., Augustin, J., Shatwell, T., Barth, M., Gelbrecht, J. and McInnes, R. J.: Changes of the
866 CO₂ and CH₄ production potential of rewetted fens in the perspective of temporal vegetation
867 shifts, Biogeosciences, 12, 2455-2468, doi:10.5194/bg-12-2455-2015, 2015.
- 868 Zerbe, S., Steffenhagen, P., Parakenings, K., Timmermann, T., Frick, A., Gelbrecht, J. and Zak, D.:
869 Ecosystem service restoration after 10 years of rewetting peatlands in NE Germany, Environ.
870 Manage., 51, 1194-1209, doi:10.1007/s00267-013-0048-2, 2013.
- 871 Zhu, D., Chen, H., Zhu, Q., Wu, Y. and Wu, N.: High Carbon Dioxide Evasion from an Alpine Peatland
872 Lake: The Central Role of Terrestrial Dissolved Organic Carbon Input, Water Air Soil Poll., 223,
873 2563-2569, doi:10.1007/s11270-011-1048-6, 2012.
- 874



875 Table 1: Data loss and final data coverage during the observation period. Percentage of CO₂ and CH₄
 876 flux data lost by power and instrument failure and maintenance as well as quality control and footprint
 877 analysis.

Filter criteria	Percentage of data [%]	
	CO ₂	CH ₄
Power and instrument failure, maintenance	15.0	46.4
Absence of sensor	-	11.2
QC 2	7.5	2.0
RSSI	-	2.1
u*	18.6	8.8
Unreasonably high fluxes	0.2	0.1
No footprint information/footprint > 20 % outside the AOI	13.2	6.5
Final data coverage	45.5	22.9

878



879 Table 2: Gapfilling model performance was estimated according to Moffat et al. (2007) with several
 880 measures ($n_{\text{CO}_2} = 6193$, $n_{\text{CH}_4} = 3386$, fluxes of best quality QC 0): the adjusted coefficient of
 881 determination R^2_{adj} for phase correlation (significant in all cases, $p\text{-value} < 2.2e^{-16}$), the absolute root
 882 mean square index (RMSE_{abs}) and the mean absolute error (MAE) for the magnitude and distribution
 883 of individual errors, as well as the bias error (BE) for the bias of the annual sums.

Method	R^2_{adj}	RMSE_{abs} ($\text{mg m}^{-2} 30\text{min}^{-1}$)	MAE ($\text{mg m}^{-2} 30\text{min}^{-1}$)	BE ($\text{g m}^{-2} \text{a}^{-1}$)
LUT _{CO2nofoot}	0.74	104.35	24.05	13.14
NLR _{CO2foot}	0.66	119.10	27.51	-2.12
NLR _{CH4nofoot}	0.79	1.36	0.83	-3.34
NLR _{CH4foot}	0.81	1.28	0.78	-2.54

884



885 Table 3: Annual balances of CO₂ and CH₄ derived by different methods for the whole EC source area,
 886 the area of interest (AOI) and the two surface types: LUT approach without footprint consideration
 887 (LUT_{CO2nofoot}), NLR approach without (NLR_{CH4nofoot}) and with (NLR_{CH4foot}, NLR_{CO2foot}) footprint
 888 consideration. Uncertainty was calculated as square root of the sum of squared random uncertainty
 889 (measurement uncertainty) and gapfilling uncertainty.

Source area	Flux (g m ⁻² a ⁻¹)	Method			
		CO ₂		CH ₄	
		LUT _{CO2nofoot}	NLR _{CO2foot}	NLR _{CH4nofoot}	NLR _{CH4foot}
Whole EC source area	NEE	524.5 ± 5.6	531.4 ± 13.0		
	GPP	-2380.5 ± 5.6	-2122.1 ± 16.7		
	R _{eco}	2863.6 ± 5.6	2603.6 ± 8.4		
	CH ₄			40.5 ± 0.2	39.8 ± 0.2
AOI	NEE		843.5 ± 13.0		
	GPP		-3192.2 ± 16.7		
	R _{eco}		4035.7 ± 8.4		
	CH ₄				21.8 ± 0.2
Emergent vegetation	NEE		750.3 ± 13.0		
	GPP		-4076.8 ± 16.7		
	R _{eco}		4827.2 ± 8.4		
	CH ₄				13.2 ± 0.2
Open water	NEE		158.2 ± 13.0		
	GPP		-1021.5 ± 16.7		
	R _{eco}		1179.7 ± 8.4		
	CH ₄				52.6 ± 0.2

890



891 Table 4: NEE and net CH₄ exchange at open water sites. The letters in parentheses indicate seasonal
 892 (S; May to October) and annual (A) budgets. Positive water level indicates inundated conditions. GHG
 893 flux measurement methods are denoted as: CH = chambers, CO = concentration profiles, TR = gas traps.

Reference	Location, ecosystem type	Dominant plant species	Study year	Average water depth (m)	NEE (g CO ₂ m ⁻² a ⁻¹)	CH ₄ (g CH ₄ m ⁻² a ⁻¹)
Huttunen et al. (2003), CH	Lake Postilampi, Finland: hypertrophic lake		1997	3.2		16 (A)
Casper et al. (2000), TR/CO	Priest Pot, UK: hypertrophic lake		1997	2.3		13 (A)
Ducharme-Riel et al. (2015), CO	Bran-de-Scie, Quebec: eutrophic lake		2007-2008	3.2	224 (A)	
Wang et al. (2006), CH	Taihu Lake, China, hypertrophic lake: - bare infralittoral zone - pelagic zone		2003-2004	0.5 to 1.8 1.8		3 (A) 4 (A)
Hendriks et al. (2007), CH	Horstermeer, The Netherlands: eutrophic ditches		2004-2006	> 0		5 (A)
Waddington and Day (2007), CH	Bois-des-Bel peatland, Quebec: - ponds - ditches		2000-2002	> 0 > 0		0.3 (S) 2.9 (S) ¹
Naimann et al. (1991), CH	Kabetogama Peninsula, Minnesota, beaver pond: - submergent aquatic plants - deep water	<i>Utricularia spp.</i> , <i>Potamogeton spp.</i>	1988	0.45 1.25		14 (A) 12 (A)
Roulet et al. (1992), CH	Low forest region, Ontario: beaver ponds		1990	0.2 to 0.4		7.6 (A)
Bubier et al. (1993), CH	Clay Belt, Ontario: beaver pond		1991	0.5 to 1.5		44 (A)
Yavitt et al. (1992), CH	New York, beaver ponds: - 3 years old - > 30 years old		1990	≤ 2 ≤ 2		34 (A) 40 (A)




894 Table 5: Annual (A)/seasonal (S) NEE, GPP, R_{eco} and net CH_4 exchange at *Typha* sites. Positive water
 895 level indicates inundated soil. GHG flux measurement methods are denoted as: CH = chambers, EC =
 896 eddy covariance.

Reference	Location, ecosystem type	Dominant plant species	Study year	Mean water level (m)	NEE (g CO_2 m ⁻² a ⁻¹)	GPP (g CO_2 m ⁻² a ⁻¹)	R_{eco}	CH_4 (g CH_4 m ⁻² a ⁻¹)
Kankaala et al. (2004), CH	Lake Vesijärvi, Finland: - inner cattail-reed zone	<i>Phragmites australis</i> , <i>Typha latifolia</i>	1997	< 0.1 to > 0.2				51 (S) ¹
			1998	< 0.1 to > 0.2				43 (S) ¹ , 6 (S) ²
Chu et al. (2015), EC	Lake Erie, Freshwater marsh	<i>Phragmites australis</i> , <i>Typha latifolia</i>	1997	< 0.1 to > 0.2				30 (S) ¹
			1998	< 0.1 to > 0.2				23 (S) ¹ , 7 (S) ²
Bonneville et al. (2008), EC Strachan et al. (2015), NEE: EC, CH: CH	Mer Bleue, Canada, freshwater marsh	<i>Typha angustifolia</i> , <i>Nymphaea odorata</i>	2011	0.3 to 0.6	-289 (A)	-3338 (A)	3049 (A)	58 (A)
			2012	0.3 to 0.6	109 (A)	-3490 (A)	3599 (A)	76 (A)
			2013	0.3 to 0.6	340 (A)	-2666 (A)	3006 (A)	70 (A)
			2005-2006 2005-2009	winter > summer ≈ 0	-967 (A) -462 to -1041 (A)	-3045 (A)	2078 (A)	170 (A)
Whiting and Chanton (2001), CH	Virginia, freshwater marsh	<i>Typha latifolia</i>	1992-1993	0.05 to 0.2	-3288 (A)			109 (A)
			1992 1993	0.05 to 0.2 0.05 to 0.2	-3587 (A) -4177 (A)			69 (A) 96 (A)
Rocha and Goulden (2008), EC	San Joaquin Freshwater Marsh Reserve, California: - freshwater marsh	<i>Typha latifolia</i>	1999 2000 2001 2012	winter +, midsummer - winter +, midsummer - winter +, midsummer - 1.07		-3994 (A) -6006 (A) 1887 (A) -1349 (A)	4811 (A) 5980 (A) 6721 (A)	71 (A)
Knox et al. (2015), EC	- wetland (rewetted 2010) - wetland (rewetted 1997)	<i>Schaenoplectus acutus</i> , <i>Typha</i> spp. <i>Schaenoplectus acutus</i> , <i>Typha</i> spp.	2012	0.26	-1455 (A)	-5519 (A)	4064 (A)	52 (A)
			2010	0.51	388 (A)			21 (A)
Petrescu et al. (2015), EC	- wetland (rewetted 2010)	?	2010-2011 2011-2012	1 0.7	553 (A) -414 (A)	-2825 (A) -3980 (A)	3375 (A) 3566 (A)	80 (A) 91 (A)
			Belarus, fen (rewetted 1985)					
Minke et al. (2015), CH	Giel'čykaŭ Kašy, Belarus, fen (rewetted 1985)	<i>Typha latifolia</i> , <i>Hydrocharis morsus-ranae</i>	2011 2012	0.02 -0.09	-156 (A) 345 (A)			13 (A) 4 (A)
			Trebbelal, Germany, fen (rewetted 1997)					
Günther et al. (2015), CH	Turreaun, Ireland, cutover bog (rewetted 1991)	<i>Typha latifolia</i>	2002 2003	0.07 0.03	975 (A) 1653 (A)	-3272 (A) -4357 (A)	4064 (A) 6010 (A)	39 (A) 29 (A)


¹ open water period

² winter



897
898 Figure 1: a) Polder Zarnekow is situated in NE Germany within the Peene River valley; map source and
899 copyright: <https://commons.wikimedia.org/wiki/File:Germanymap2.png> (modified). b) Footprint
900 climatology calculated according to Chen et al. (2011) on a Landsat image (6 Jun 2013, source: Google
901 Earth). White lines represent the isopleths of the cumulative annual footprint climatology, where the
902 area within the 95 isopleth indicates 95 % contribution to the annual flux. The white dot denotes the
903 tower position. The yellow box indicates the area of interest (AOI) as a filter criterion to focus on fluxes
904 of the shallow lake and to avoid the possible impact of a farm and grassland to the north of the shallow
905 lake. If the half-hourly flux source area exceeded the AOI by more than 20 % the flux was discarded.
906 The site is characterised by two main surface types: open water and emergent vegetation.

907



908


909 Figure 2: Temporal variability of environmental variables and ecosystem CO₂ and CH₄ exchange.

910 Seasonal course a) of water level (Wlevel), cumulative precipitation (Cum. Precip) and air temperature

911 (T_{air}), b) the daily CH₄ flux (gapfilled, NLR_{CH₄nofoot}) and c) the daily NEE (gapfilled LUT_{CO₂nofoot}) and

912 component fluxes (modelled R_{eco} and GPP, LUT_{CO₂nofoot}).

913



914

915 Figure 3: Average diurnal cycle of a) CO₂ flux, b) CH₄ flux and c) the water density gradient per month.

916 The numbers at the x-axis denote midnight (0) and midday (12). Midnight is also illustrated with a

917 dashed line. Black and grey lines represent the mean and the range, respectively. The CH₄ fluxes are

918 normalized with the monthly median of the half-hourly fluxes. Positive CO₂ fluxes represent the


919 dominance of R_{eco} against GPP, negative fluxes the dominance of GPP against R_{eco}. The period of ice-

920 cover was excluded from the calculation of the temperature gradient. A density gradient equal to or

921 below zero indicates thermally induced convective mixing down to the bottom of the open water body

922 of the shallow lake, positive gradients instead thermal stratification.

923




924

925 Figure 4: Cumulative GWP budgets of CO₂ (based on LUT_{CO2nofoot}), CH₄ (based on NLR_{CH4nofoot}) for the EC

926 source area and the sum of both during the observation period.

927




928

929 Figure 5: Source area fraction Ω_i of the two main surface types in dependence on the wind direction

930 (2° -bins).


931



932

933 Figure 6: Impact of the fractional coverage of open water (Ω_{water}) within the EC source area on the
934 measured fluxes of CO_2 and CH_4 . The variability of CO_2 flux rates decreased with increasing Ω_{water} ,
935 whereas the variability of the CH_4 flux increased.

936



937

938 Figure 7: Daily CH₄, NEE and component fluxes (R_{eco} and GPP) for the surface types: a) daily CH₄ flux of
 939 open water and emergent vegetation, b) daily NEE and component fluxes for open water, c) daily NEE
 940 and component fluxes for emergent vegetation, derived by NLR with the source area fractions of the
 941 surface types (Ω_i) as weighting factors (NLR_{CH_4foot} , NLR_{CO_2foot}).