



Ideas and perspectives: Heat stress: more than hot air

Hans J. De Boeck¹*, Helena Van De Velde^{1,2}*, Toon De Groote¹, Ivan Nijs¹

¹ Centre of Excellence PLECO (Plant and Vegetation Ecology), Department of Biology, Universiteit Antwerpen (Campus Drie Eiken), Universiteitsplein 1, B-2610 Wilrijk, Belgium

² Terrestrial Ecology Unit, Department of Biology, Universiteit Gent, K. L. Ledeganckstraat 35, B-9000 Ghent, Belgium

*equal contribution

Correspondence to: Hans J. De Boeck (hans.deboeck@uantwerp.be)

Abstract. Climate change models project an important increase in the frequency and intensity of heat waves. In gauging the impact on plant responses, much of the focus has been on air temperatures while a critical analysis of leaf temperatures during heat extremes has not been made. Nevertheless, direct physiological consequences from heat depend primarily on leaf rather than on air temperatures. We discuss how the interplay between various environmental variables and the plants' stomatal response affects leaf temperatures and the potential for heat stress by making use of both an energy balance model and field data. The results demonstrate that this interplay between plants and environment can cause leaf temperatures fluctuations in excess of 10 °C (for narrow leaves) to even 20 °C (for big broad leaves) at the same air temperature. In general, leaves tended to heat up when radiation was high and when stomates were closed, as expected. But perhaps counterintuitively, also high air humidity raised leaf temperatures, while humid conditions are typically regarded as benign with respect to plant survival since they limit water loss. High wind speeds brought the leaf temperature closer to the air temperature, which can imply either cooling or warming (i.e. abating or reinforcing heat stress) depending on other prevailing conditions. The results thus indicate that heat waves characterized by similar extreme air temperatures may pose little danger under some atmospheric conditions, but could be lethal in other cases. The trends illustrated here should give ecologists and agronomists a more informed indication about which circumstances are most conductive for heat stress to occur.

25 1 Introduction

Current climate change has made heat waves more likely as both the temperature mean and variability are increasing (Schär et al., 2004). Several well-documented heat waves have occurred during the past years such as those of 2003 (Europe), 2010 (Russia) and 2012 (North America), and the likelihood of such major events is expected to increase 5 to 10-fold within the next 40 years (Barriopedro et al., 2011). Heat stress in plants is usually observed when tissue temperatures exceed 40 °C, a threshold that is fairly stable across biomes (Larcher, 2003). Such excessive temperatures affect plant metabolism in multiple



ways, ultimately reducing growth and economic yield (Bastos et al., 2014; Chung et al., 2014). This seems at odds with the reported lack of significant single-factor effects in several ecological studies on heat waves (Poirier et al., 2012; Hoover et al., 2014; De Boeck et al., 2016). We examine here how these seemingly contrasting notions can be reconciled. The fundamental issue is that air temperature (T_a) is often considered as an important indicator of heat stress, while metabolic rates and physiological processes are affected much more directly by leaf (tissue) temperatures (T_l). The latter are influenced by a number of environmental conditions (apart from T_a , primarily through radiation, wind speed and air humidity) and the stomatal response of the plants. The extent to which these variables can decouple leaf from air temperatures and therefore increase or decrease the potential for heat stress during a heat wave of similar magnitude (in terms of air temperature, as it is usually considered) is discussed here by making use of both an energy balance model and field data.

10

2 Materials and Methods

The model used to calculate leaf temperature is based on the energy balance equation (Eq. 1):

$$R_{s,in} + R_{l,in} - R_{l,out} - H - \lambda E = 0 \quad (1)$$

15

The equation states that an equilibrium is reached under a certain set of environmental conditions (the flux of sensible heat H can be either incoming or outgoing), so that the sum of incoming energy (via shortwave radiation $R_{s,in}$ and longwave radiation $R_{l,in}$) and outgoing energy (outgoing longwave radiation $R_{l,out}$, and latent heat λE) is zero. The different terms are derived from other equations, which feature both environmental variables such as wind speed (u) and relative humidity (RH) 20 of the air, leaf-scale parameters such as stomatal conductance (g_s) and characteristic leaf dimension (d), and constants such as the Stefan Boltzman constant (σ , $5.67e^{-8} \text{ W m}^{-2} \text{ K}^{-4}$). For more details, we refer to De Boeck et al. (2012).

25

The leaf temperature is calculated in an iterative manner: as a starting situation it is assumed that leaf and air temperature are equal, in which case the energy budget equals zero. In any other situation, the model will assume T_l to be lower/higher than T_a if the energy budget is negative/positive. The iteration proceeds in a stepwise manner, until a precision of $0.01 \text{ }^\circ\text{C}$ is achieved. The model was validated earlier (De Boeck et al., 2012), demonstrating a deviation between measured and modelled leaf temperatures of less than $1.5 \text{ }^\circ\text{C}$ for over 90% of the cases.

30

In this study, we set T_a at $40 \text{ }^\circ\text{C}$ to approximate the general threshold for heat stress. Atmospheric pressure (which has limited influence) was kept constant at 100 kPa. Emissivity, reflectivity and absorptivity parameters for leaves and soil were used like in De Boeck et al. (2012). Major inputs were varied in a dichotomous manner (high or low): incident shortwave energy (R_s), stomatal conductance, wind speed, and relative humidity of the air. We focus on vegetation represented by species that have narrow leaves (like those found in many grasses) with a characteristic dimension of 0.5 cm, but we also consider the opposite end of the spectrum, namely very broad leaves with a d of 20 cm.



The modelled results are supported by data recorded on five sunny days during a heat wave in Belgium in 2015 (1-5 July). These data were collected at an experimental site in Wilrijk, Belgium on two homogeneous 10 cm tall young grass stands (sown five weeks earlier on homogenised soils). The grass was irrigated daily, with the exception of one day to test the impact of surface drying on the difference between T_a and T_l (via stomatal responses). Radiation sensors (SR03-05, 5 Hukseflux Thermal Sensors, Delft, The Netherlands) had been installed approximately 30 cm above the vegetation, with one sensor directed upwards, and one sensor directed downwards to measure absorbed radiation (the difference between the two readings). At the same height, canopy temperature was recorded with a non-contact thermometer (custom made). Air temperature and relative humidity were measured at 15 cm height in each plot using custom made sensors shielded from the sun by a thin wooden panel. To ensure that mostly data from times when direct sunshine reached the plots was used 10 (generally between 9 am and 7 pm CET), we omitted data points with absorbed radiation below 100 W m^{-2} . This was done to prevent artefacts from dew or times when stomates were still closed.

3 Results and discussion

Our results show that high radiation loads are an important prerequisite for heat stress, unless air temperatures exceed the 15 tissue heat stress threshold significantly. Without the extra energy provided by sunshine, plant tissues will almost always be cooler than the surrounding air (Fig. 1). In reality, heat waves usually feature clear sky conditions (De Boeck et al., 2010), implying that radiation loads during hot weather are highly probable. This also means that experiments in which high air temperatures are imposed in low-radiation environments, like under laboratory conditions or during overcast days, may underestimate impacts.

20 As highlighted in earlier studies, water availability or lack thereof is greatly relevant in gauging whether a heat wave will give rise to heat stress (Salvucci and Crafts-Brandner, 2004). If drought prompts a plant to conserve water by lowering stomatal conductance (g_s), it warms up as energy dissipation shifts from latent fluxes (providing cooling) to sensible fluxes (increasing temperatures). Because heat and drought often co-occur naturally (De Boeck et al., 2010), this effect is highly relevant in assessing heat wave impacts (Idso, 1982; De Boeck et al., 2016). The potentially misleading nature of T_a in 25 predicting heat stress under varying stomatal conductance is clearly highlighted in our results (Fig. 1).

Whenever other conditions alleviate some amount of heat stress (e.g. less radiation, higher g_s), more wind would counteract such beneficial effects (Fig. 1) through closer coupling between the plant and the air. This may seem counterintuitive as windiness is generally associated with heat dissipation, but the same process also works in the opposite case: when other environmental conditions would exacerbate heat stress, more wind reduces the increase of leaf temperatures. In other words, 30 windy conditions lead to avoidance of the most extreme cases of overheating. Obviously, higher wind speeds promote evapotranspiration, resulting in faster depletion of soil water reserves. This could subsequently lead to lower g_s and thus indirectly promote overheating. As wind speeds in laboratory conditions and/or enclosures are often far below those



observed outside (De Boeck et al., 2012), canopy warming may be significantly different from outside as calm conditions tend to exacerbate other effects (Fig. 1).

Also for relative air humidity (RH), the results are counterintuitive, with higher humidity more likely to give rise to heat stress. This is caused by slower heat dissipation via transpiration as the water vapour gradient between leaf and air is smaller
5 than in the case of drier air. In fact, the combination of low stomatal conductance and high air humidity causes the greatest warming of leaves above the air temperature (Fig. 1). A five-day period featuring air temperatures at vegetation height above 30 °C every day provided us with an opportunity to test whether increasing air humidity diminishes the cooling capacity of leaves. We indeed found a significant relationship between RH and $T_l - T_a$ (Fig. 2), with ± 0.84 °C change per 0.1 increase
10 in RH. In the same figure, the influence of decreasing stomatal conductance is also illustrated, with transpirational cooling clearly reduced on the only day during which irrigation was withheld.

The aforementioned trends were observed both for simulations using narrow (Fig. 1) and also for simulations using bigger leaves (Fig. S1). Any variable increasing the heat load (high radiation) or decreasing heat dissipation (high RH, low wind and g_s) led to higher temperature increases in big compared to in small leaves, however. This is no surprise as larger surfaces result in increased decoupling from air temperatures, which can lead to extreme temperature deviations. In cushion plants,
15 which physically act as a giant leaf, increases of tissue temperatures of 20 °C and more above the air temperature have been observed (Gauslaa, 1984), illustrating the importance of physical dimensions in energy balances.

In conclusion, we clearly demonstrated that exceedance of critical temperatures in plants depends on more variables than air temperature alone. Radiation, wind speed and relative humidity all affect tissue temperatures, depending on plant water status. This implies that heat waves characterized by the same extreme air temperatures may cause little plant damage under
20 some conditions, but could be detrimental to plant growth and survival in other cases. Although heat stress also depends on other factors, like hardening (Neuner and Buchner, 2012), the results from this study can help predict when the probability of heat stress occurring is most likely, and can stimulate ecologists and agronomists to shift the focus beyond merely air temperatures when considering heat waves.

25 Author contribution

H. De Boeck and I. Nijs conceptualised the study. T. De Groote developed the model code and H. Van De Velde performed the simulations. H. De Boeck carried out the field measurements. All authors worked together to prepare the manuscript.

Acknowledgements

30 H. Van De Velde was supported by FWO Vlaanderen. We thank F. Kockelbergh for technical assistance.



References

5 Barriopedro, D., Fischer, E. M., Luterbacher, J., Trigo, R., and Garcia-Herrera, R.: The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe, *Science*, 332, 220-224, 2011.

Bastos, A., Gouveia, C. M., Trigo, R. M., and Running, S. W.: Analysing the spatio-temporal impacts of the 2003 and 2010 extreme heatwaves on plant productivity in Europe, *Biogeosciences*, 11, 3421–3435, 2014.

10 Chung, U., Gbegbelegbea, S., Shiferawe, B., Robertson, R., Yun, J. I., Tesfaye, K., Hoogenboom, G., and Sonder, K.: Modeling the effect of a heat wave on maize production in the USA and its implications on food security in the developing world, *Weather Clim. Extremes*, 5–6, 67–77, 2014.

15 De Boeck, H. J., Dreesen, F. E., Janssens, I. A., and Nijs, I.: Climatic characteristics of heat waves and their simulation in plant experiments, *Glob. Change Biol.*, 16, 1992-2000, 2010.

De Boeck, H. J., De Groote, T., and Nijs, I.: Leaf temperatures in glasshouses and open-top chambers, *New Phytol.*, 194, 1155-1164, 2012.

20 De Boeck, H. J., Bassin, S., Verlinden, M., Zeiter, M., and Hiltbrunner, E.: Simulated heat waves affected alpine grassland only in combination with drought, *New Phytol.*, 209, 531-541, 2016.

Gauslaa, Y.: Heat resistance and energy budget in different Scandinavian plants, *Holarctic Ecol.*, 7, 1–78, 1984.

25 Hoover, D. L., Knapp, A. K., and Smith, M. D. Resistance and resilience of a grassland ecosystem to climate extremes, *Ecology*, 95, 2646–2656, 2014.

Idso, S. B. Non-water-stressed baselines: a key to measuring and interpreting plant water stress, *Agr. Meteorol.*, 27, 59-70, 30 1982.

Larcher, W.: *Physiological plant ecology*, 4th edition. Springer-Verlag, Berlin, 2003.



Neuner G. and Buchner, O.: Dynamics of tissue heat tolerance and thermotolerance of PS II in alpine plants, in: Plants in alpine regions: cell Physiology of adaptation and survival strategies, edited by: Lütz, C. Springer-Verlag, Vienna, 61-74, 2012.

5 Poirier, M., Durand, J. L., and Volaire, F.: Persistence and production of perennial grasses under water deficits and extreme temperatures: importance of intraspecific vs. interspecific variability, *Glob. Change Biol.*, 18, 3632–3646, 2012.

Salvucci, M. E. and Crafts-Brandner, S.J.: Inhibition of photosynthesis by heat stress: the activation state of Rubisco as a limiting factor in photosynthesis, *Physiol. Plantarum*, 120, 179-186, 2004.

10

Schär, C., Vidale, P. L., Lüthi, D., Frei, C., Häberli, C., Liniger, M. A., and Appenzeller, C.: The role of increasing temperature variability in European summer heatwaves, *Nature*, 427, 332–336, 2004.

15

20

25




Figure 1: Modelled leaf-to-air temperature difference depending on type of heat wave and stomatal conductance (g_s). Type of heat wave: high (A) or low (B) incident shortwave radiation (800 or 100 W m^{-2}), high or low relative humidity of the air ($\text{RH} = 0.90$ or 5 0.45), and calm or windy weather (wind speed 0.1 or 6 m s^{-1}). Air temperature was set to 40°C in all simulations, and leaf width to 0.005 m .




Figure 2: Differences between air (T_a) and leaf (T_l) temperature in function of relative air humidity (RH) measured on a homogeneous grass stand during 5 heat wave days (1-5 July 2015, Belgium). The grass was irrigated daily (white circles), with the exception of one day (black circles). The linear regression was significant at $p < 0.001$ ($R^2 = 0.13$).