

Responses to the comments of reviewers

Referee #1

The authors tested the growth and photophysiological responses of *Emiliana huxleyi* to PAR and UV in the presence of coccoliths, after removal of coccoliths and in a strain that lacks coccoliths. The data show that presence of coccoliths renders the cells less susceptible to inhibition by UV, and increases their capacity for non-photochemical quenching. The manuscript presents a tidy study on an important question, and is appropriate for BioGeoScience. I offer a few minor wording and reference comments for the author's consideration.

best regards, Doug Campbell

Abstract: Fine.

"...since decades..." is not incorrect, but is idiomatically odd. I suggest "...for decades...".

[Response: corrected as suggested.](#)

Introduction: "This notion is supported by the exceptionally high light tolerance of the surface layer dwelling species *Emiliana huxleyi* (Nanninga and Tyrell 1996; Gao et al., 2009)" Geider's group had a paper: Ragni M, Airs RL, Leonardos N, Geider RJ. 2008. PHOTOINHIBITION OF PSII IN EMILIANIA HUXLEYI (HAPTOPHYTA) UNDER HIGH LIGHT STRESS: THE ROLES OF PHOTOACCLIMATION, PHOTOPROTECTION, AND PHOTOREPAIR. *Journal of Phycology* 44: 670–683. and we had a paper: Loebel M, Cockshutt AM, Campbell DA, Finkel ZV. 2010. Physiological basis for high resistance to photoinhibition under nitrogen depletion in *Emiliana huxleyi*. *Limnology and Oceanography* 55: 2150–2160. both showing that the high PAR tolerance of *E. hux* related to very strong repair capacities, rather than intrinsic resistance to photoinactivation, per se. It would be worth noting that UV is a strong inhibitor of PSII repair, as well as acting through direct inhibition of PSII. So it could be that the coccoliths protect PSII repair from UV inhibition. I now read you briefly make this point in the discussion, citing Gao 2007.

[Response: We thank the reviewer for this constructive comments. We added a sentence in](#)

the last paragraph of introduction and cited extra references. The mentioned references above were cited in the introduction and discussion at lines 75 and 355.

Materials & Methods: Fine

Results: "Photochemical performance was measured for dark-adapted (15 min) cells in calcified, de-calcified or non-calcifying naked cells"

The table and figure abbreviation Cal-R does not obviously suggest 'de-calcified'. Why not 'D-Cal' or 'Cal-D'? More generally, why erect abbreviations? Why not just write out 'Calcified', 'De-calcified', and 'Naked'? In the text the naked strain is sometimes called naked, or sometimes 'non-calcifying'. Unify the terminology; pick a single name for each cell condition and use it throughout.

Response: As you suggested, we corrected them as 'Calcified', 'De-calcified', and 'Naked'

Anonymous Referee #2

Received and published: 2 June 2016

Coccolithophores are an ecologically important group of marine phytoplankton that characteristically produce calcium carbonate plates (liths) internally and then secrete them to the cell surface. Exactly why coccolithophores produce liths has been the subject of considerable debate, with a range of possibilities raised. The manuscript by Xu et al has set out to test the hypothesis that the calcite from which the liths are constructed will absorb enough UV radiation to protect the cells from damage.

The approach the authors have used is to compare the UV sensitivity (measured as growth, quantum yield of PSII and relative electron transport characteristics including non-photochemical quenching) in a calcified strain, a non-calcifying strain and the calcifying strain with the liths removed (which could be more clearly termed 'decalcified' in the text and figures/tables).

Response: Corrected as "de-calcified"

The experimental approach is sound and the results are presented clearly and discussed thoroughly. I really only have some minor points the authors might like to consider.

P 2 line 26: 'for' not 'since'

Response: Corrected.

P 3 line 52: delete 'by'

Response: deleted.

P 4 lines 72/72: shading effects could include scattering of light which is certainly a feature of coccolithophore blooms

Response: We added this in the text and cited two papers to support it at lines 84-87.

P 4 lines 73/74: a bit pedantic I know, but strictly speaking it is hard to see how liths could stimulate NPQ – presumably liths are affecting the light climate in some way that leads to up-regulation of NPQ

Response: We think it is possible that NPQ is indirectly affected, which coincides with calcification. This sentence was deleted in the text.

P 9 line 180 et seq.: I assume some of this loss of transmittance could be due to scattering by liths rather than absorbance? In Fig 1, the non calcifying strain has a lower transmittance in the UV than does the calcifying strain without liths - could this be because the non-calcifying strain employs another strategy (such as inducing UV screening compounds) to ameliorate UVB?

Response: We used a double beam UV-VIS-NIR spectrophotometer (PerkinElmer, Lambda950, USA) to obtain the absolute absorbance of liths, so that all the scattered light was recaptured. This was added in the Methods (lines 166-167). It might be one of strategies against UV damage for non calcifying strain to synthesize UV screening compounds, and we added this in the discussion at lines 320-323.

P 10 lines 201-203: A significant decrease in $rETR_{max}$ would, if α is unaffected, be expected to show a decrease in I_k (given the relationship $I_k = rETR_{max}/\alpha$) so the lack of an effect on I_k seems odd. In fact there does seem to be a difference but it is statistically non-significant.

Response: Considering the relationship $I_k = rETR_{max}/\alpha$, it would be true. In our results, there are differences in I_k between Calcified and De-calcified cells, which is only statistically different at $P > 0.05$. Since α is lower in De-calcified cells than that in calcified cells,

which is not statistically different at $P < 0.05$, $I_k = rETR_{max}/\alpha$ could not statistically be different.

P 11 discussion lines 233- 234: While I don't disagree with the authors conclusions, you have to be a bit careful in making claims that the liths are causing the differences in growth rate between the naked and calcifying strain based on only one strain of each phenotype (certainly when grown indoors). Would all naked strains grow more slowly than all calcified strains? Remember also that calcification leads to internal generation of CO₂ - although it has been shown elegantly by Bach et al 2013 that there is no obligatory coupling between calcification and photosynthesis. The differential impact of UVB on growth is though compelling!

Response: We have revised the conclusion, now it reads "It has to be noted that our experimental data is based on only two strains of a naked and calcified *E. huxleyi*. However, similar trends in photophysiology between naked and decalcified in comparison to calcified cells suggest that coccoliths of *E. huxleyi* play an important role in protecting this species against harmful solar radiation especially UV-A and UV-B. Furthermore, the reported absence of photoinhibition in this alga at high light levels also appears to be connected to the coccosphere of *E. huxleyi* or its calcification process. In view of ongoing ocean change, the projected shoaling of the upper mixed layer (UML) caused by global warming and progressive ocean acidification that reduces the thickness or the number of coccoliths per cell (Gao et al., 2009; De Bodt et al., 2010) could reduce *E. huxleyi* growth rates within the UML due to increased UVR exposure. "

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

**The role of coccoliths in protecting *Emiliana huxleyi* against stressful light and
UV radiation**

Running Title: Photoprotective role of coccoliths in *Emiliana huxleyi*

Juntian Xu^{1,2}, Lennart T Bach³, Kai G Schulz³, Wenyan Zhao¹, Kunshan Gao^{1*}, Ulf
Riebesell³

¹State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen,
Fujian, 361102 China;

²Key Laboratory of Marine Biotechnology of Jiangsu Province, Huaihai Institute of
Technology, Lianyungang, Jiangsu, 222005 China;

³GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, Kiel,
24105 Germany

*Author for Correspondence: ksgao@xmu.edu.cn (Kunshan Gao)

26 **Abstract**

27 Coccolithophores are a group of phytoplankton species which cover themselves
28 with small scales (coccoliths) made of calcium carbonate (CaCO₃). The reason why
29 coccolithophores form these calcite platelets has been a matter of debate for decades
30 but has remained elusive so far. One hypothesis is that they serve a role in light/UV
31 protection, especially in surface dwelling species like *Emiliana huxleyi* which can
32 tolerate exceptionally high levels of solar radiation. In this study, we tested this
33 hypothesis by culturing a calcified and a naked strain under different light conditions
34 with and without UV radiation. The coccoliths of *E. huxleyi* reduced the transmission
35 of visible radiation (400-700 nm) by 7.5%, UV-A (315-400 nm) by 14.1% and UVB
36 (280-315 nm) by 18.4%. Growth rates of the calcified strain (PML B92/11) were
37 about 2 times higher than those of the naked strain (CCMP 2090) under indoor
38 constant light levels in the absence of UV radiation. When exposed to outdoor
39 conditions (fluctuating sunlight with UV radiation), growth rates of calcified cells
40 were almost 3.5 times higher compared to naked cells. Furthermore, relative electron
41 transport rate was 114% higher and non-photochemical quenching (NPQ) 281%
42 higher in the calcified compared to the naked strain, implying higher energy transfer
43 associated with higher NPQ in the presence of calcification. When exposed to natural
44 solar radiation including UV radiation, maximal quantum yield of photosystem II was
45 only slightly reduced in the calcified but strongly reduced in the naked strain. Our
46 results reveal an important role of coccoliths in mitigating light and UV stress in *E.*
47 *huxleyi*.

48

49 **Key words:** coccoliths, *Emiliana huxleyi*, light protection, growth, photosynthetic
50 performance, UV radiation

删除的内容: since

删除的内容: calcifying

删除的内容: non-calcifying

删除的内容: calcifying

删除的内容: on-calcifying

删除的内容: calcifying

删除的内容: non-calcifying

删除的内容: calcifying

删除的内容: non-calcifying

60

61 **1 Introduction**

62 Coccolithophores are a group of marine phytoplankton species which are able to
63 precipitate CaCO₃ in the form of small calcitic scales (coccoliths) surrounding the
64 organic part of the cell. They contribute about 1-10% to marine primary production
65 (Poulton et al., 2007) and approximately 50% to pelagic deep ocean CaCO₃ sediments
66 (Broecker and Clark, 2009). Blooms of coccolithophores can cover up to 8 million
67 km² of the Earth's surface (Moore et al., 2012), and are considered to be important
68 drivers of biogeochemical cycling (Rost and Riebesell, 2004).

69 Despite intense research on coccolithophore calcification and its biogeochemical
70 relevance during the last decade, it is still an unresolved question why
71 coccolithophores calcify (Young, 1994; Raven and Crawford, 2012). One hypothesis
72 is that the layer of coccoliths surrounding the cell (coccosphere) protects the organism
73 from excess light and UV radiation. This notion is supported by the exceptionally
74 high light tolerance of the surface layer dwelling species *Emiliana huxleyi* (Nanninga
75 and Tyrrell, 1996; [Ragni et al., 2008](#); [Gao et al., 2009](#); [Loebl et al., 2010](#)).

76 Physiological studies investigating the light tolerance of *E. huxleyi* showed that the
77 radiation wavelength matters in this context. The coccosphere does not seem to
78 constitute a protection against very high intensities of photosynthetically active
79 radiation (PAR) since non-calcifying *E. huxleyi* cells are equally resistant to
80 photoinhibition as their calcifying counterparts (Nanninga and Tyrrell, 1996). This is
81 in clear contrast to the influence of stressful ultraviolet radiation (UVR) on the cells
82 where results from different physiological experiments support a protective role of the
83 coccoliths ([Gao et al., 2009](#); [Guan and Gao, 2010](#); [Gao et al., 2012](#)). Protection from
84 UVR or high light exposures by coccoliths may either work by [physically shading](#)

删除的内容: by

86 | intracellular organelles or by strongly scattering light which is certainly a feature of
87 | coccolithophore blooms (Balch et al., 1996; Voss et al., 1998). The underlying
88 | mechanisms, however, are not well understood and warrant further investigations.

89 | UVR strongly contributes to photoinhibition of photosystem II (e.g. Hakala-Yatkin
90 | et al., 2010) and effectively inhibits repair processes (Ragni et al., 2008). Therefore,
91 | it is likely that the coccoliths protect PSII repair from UV inhibition. In this study we
92 | explore in more detail how different PAR and UV radiation (280-400 nm) treatments
93 | affect calcified and naked *E. huxleyi* cells. Specifically we address the question
94 | whether the coccosphere of *E. huxleyi* helps the cells to withstand stressful levels of
95 | PAR and/or UV radiation and whether calcification influences photochemical
96 | performance.

98 | 2. Materials and Methods

99 | 2.1 Materials and pre-culture conditions

100 | Calcified *E. huxleyi* (PML B92/11 isolated in the Raunefjord area, Bergen, Norway)
101 | and naked cells (CCMP 2090 isolated in the South Pacific) were used in the
102 | experiments. Both strains were grown in triplicate cultures (300 ml square glass
103 | bottles) at 15°C in 0.2 µm filtered natural seawater (gathered from the Gulf of Biscay)
104 | at a photon flux density of 500 µmol photons m⁻² s⁻¹ on a 16/8 light/dark cycle. The
105 | natural seawater medium was enriched with 64 µmol L⁻¹ nitrate, 4 µmol L⁻¹ phosphate,
106 | f/8 concentrations of a trace metal and vitamin mixture (Guillard & Ryther 1962), and
107 | 10 nmol kg⁻¹ selenium. Pre-cultures and experimental incubations in semi-
108 | continuously diluted batch cultures (>8 generations) ensured exponential growth
109 | throughout the experiment.

110 | 2.2 Experimental setup

删除的内容: physically shading intracellular organelles

删除的内容: or by facilitating thermal dissipation through increased non-photochemical quenching (Xu and Gao, 2011).

删除的内容: It is known that

删除的内容: , PSII

删除的内容: can

带格式的: 字体: Times New Roman, 小四

删除的内容: strongly inhibit PSII repair as well as

带格式的: 字体: Times New Roman, 小四

删除的内容: damage PSII directly

带格式的: 字体: Times New Roman, 小四

删除的内容: (ref)

带格式的: 字体: Times New Roman, 小四

删除的内容:

删除的内容: calcifying

删除的内容: on-calcifying

删除的内容: k

删除的内容: Calcifying

删除的内容: on-calcifying

130 2.2.1 Indoor growth experiments

131 After pre-culture for at least 8 generations, the cells of calcified and naked strains
132 were inoculated in the same glass bottles of 300 ml and cultured under the same
133 condition as pre-cultures, maintaining the cell concentrations at exponential growth
134 within a range of $3-10 \times 10^4$ cells/ml.

删除的内容: calcifying

删除的内容: o-calcifying

135 2.2.2 Outdoor growth experiments

136 Following the indoor growth experiment, the cells were transferred into quartz
137 tubes (100 ml) for the outdoor growth experiment and were exposed to natural solar
138 radiation at the institution's pier. The cultures were maintained outside in a flow-
139 through water tank, where the seawater temperature was maintained within a range of
140 14-16°C. After the cells had acclimated for 7 days under the solar radiation, aliquots
141 of the cell cultures were transferred to new quartz tubes filled with fresh medium
142 before measurements were taken. For the outdoor cultures, the cells received 60% full
143 spectrum solar radiation (the quartz tubes wrapped with neutral density screens). The
144 daytime average intensities (from 7:00 am to 5:00 pm) of PAR, UV-A and UV-B
145 which the cells received during the outdoor experiment were about 260 $\mu\text{mol photons}$
146 $\text{m}^{-2} \text{s}^{-1}$ (about 53 W m^{-2}), 12.4 and 0.34 W m^{-2} , respectively.

147 2.2.3 Short-term incubation experiments

148 Short-term incubation experiments were carried out to test UV effects around noon
149 time on a cloudy day and sunny day, respectively. Three different radiation treatments
150 were implemented as follows: 1) Cells in uncovered quartz tubes, receiving the full
151 spectrum of solar radiation (above 280 nm, PAB treatment); 2) cells in quartz tubes
152 covered with Folex 320 (Montagefolie, Nr. 10155099, Folex, Dreieich, Germany),
153 exposed to UV-A and PAR (above 320 nm, PA treatment); and 3) cells receiving only
154 PAR (P treatment) in quartz tubes covered with Ultraphan film 395 (UV Opak,

157 Digefra, Munich, Germany). The transmission spectra of the quartz tubes and the cut-
158 off foils are given by Zheng and Gao (2009). A time-course experiment was also
159 conducted around noon under full solar spectrum conditions.

160 2.3 Absorptivity of coccoliths

161 We examined absorption spectra of the cells with or without coccoliths to get an
162 indication on how much light and/or UV are blocked by the coccosphere. Therefore,

163 calcified cells, ~~de-calcified cells~~ and cells of the naked strain, were filtered onto
164 Whatman GF/F glass fiber filters (25 mm), ~~and then~~ were subsequently placed at the
165 window near the detector of a double beam UV-VIS-NIR spectrophotometer
166 (PerkinElmer, Lambda950, USA) which can obtain the absolute absorbance of
167 coccoliths based on the recaptured scattered light. The absorption of the GF/F filter
168 was corrected with a control filter which was soaked with particle free culture
169 medium (Kishino et al., 1985).

170 2.4 Growth measurement

171 Cell densities were measured during a period of 7 days with a particle counter
172 (Coulter Z1, Beckman). The specific growth rate was calculated as: $\mu (d^{-1}) = (\ln N_t -$
173 $\ln N_0)/t$, where N_0 and N_t represent the cell concentrations at the beginning and the end
174 of the incubations and t is the incubation time in days.

175 2.5 Chlorophyll fluorescence measurement

176 Parameters of in vivo induced chlorophyll a fluorescence of photosystem II were
177 estimated by a phyto-~~pulse~~ amplitude modulated fluorometer (Phyto-PAM, Walz).
178 The maximum quantum yield of PSII (F_v/F_m) was calculated as: $F_v/F_m = (F_m -$
179 $F_o)/F_m$; where F_o is the basal fluorescence under measuring light of $0.2 \mu\text{mol}$
180 $\text{photons m}^{-2} \text{s}^{-1}$ and F_m the maximal fluorescence measured with a saturating light
181 pulse of $5000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (0.8 s) in dark-adapted (15 min) cells.

删除的内容: (Cal-C)

删除的内容: (Cal-R, see above)

删除的内容: (N-Cal)

删除的内容: which

删除的内容: -

187 In order to compare the transmission of the same strain with or without coccoliths
188 and to relate this to that of the naked strain, the calcified strain was de-calcified with
189 HCl (1 mol/L, the final concentration is 0.01 mol/L) for 10 s and subsequent recovery
190 of the pH with equimolar amounts of NaOH. Photochemical performance was
191 measured for dark-adapted (15 min) cells in calcified, de-calcified or naked cells. De-
192 calcified cells revealed Fv/Fm values similar to those obtained prior to de-
193 calcification. The actinic light levels were set at 533, 1077 and 2130 $\mu\text{mol photons m}^{-2}$
194 s^{-1} , respectively (growth light, saturated light and over-saturated light). Non-
195 photochemical quenching (NPQ) was calculated as: $\text{NPQ} = (F_m - F_m')/F_m'$, where F_m
196 was the maximum fluorescence yield after dark adaptation and F_m' the maximum
197 fluorescence yield under the actinic light levels.

删除的内容: non-calcifying

删除的内容: non-calcifying

198 To determine rapid light curves (RLCs, electron transport rate vs light), the cells
199 were exposed to 10 different PAR levels in sequence (87, 140, 263, 382, 449, 611,
200 778, 993, 1195 and 1391 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$), each of which lasted for 20 s. The
201 relative electron transport rate (rETR) was assessed as: $\text{rETR} = \text{Yield} \times 0.5 \times \text{PFD}$,
202 where the yield represents the effective quantum yield of PSII (F_v'/F_m'); the
203 coefficient 0.5 takes into account that roughly 50% of all absorbed quanta reach PSII;
204 and PFD is the photon flux density of the actinic light ($\mu\text{mol m}^{-2} \text{s}^{-1}$) (Genty et al.,
205 1989).

206 To examine immediate photochemical responses of the cells to UV radiation, the
207 cells were exposed to the three different solar radiations (see above) for 60 min during
208 noontime under natural solar radiation. The effective quantum yield was calculated as:
209 $F_v'/F_m' = (F_m' - F_t) / F_m'$, where F_m' and F_t are the maximal fluorescence and steady
210 state fluorescence in the light adapted cells, respectively.

211 2.6 Measurement of solar irradiances

214 Solar PAR was measured using a Quantum Scalar Laboratory Irradiance Sensor
215 (QSL-2100/ 2101, Biospherical Instruments, San Diego, USA). The measured values
216 were recorded every 10 s and saved on a computer. Solar UV-A and UV-B radiation
217 were measured with a radiometer (PMA 2100 Solar Light Co., Glenside, USA), the
218 mean irradiances of solar UV-A and UV-B during the experimental periods were
219 confirmed according to the ratios of UV-A /UV-B to PAR at the experimental
220 location.

221 2.7 Statistics

222 The data were expressed as the means \pm standard deviation (SD). Statistical
223 significance of the data was tested with software of Origin 9.0 (one way ANOVA,
224 Tukey's post-hoc test). A confidence level of 95% was used in all analyses.

225

226 3 Results

227 The coccolith layer of *E. huxleyi* absorbed both visible and UV radiation. It reduced
228 the transmission of visible radiation (400-700 nm) by 7.5%, UV-A (315-400 nm) by
229 14.1% and UVB by 18.4% (280-315 nm) relative to de-calcified cells and 6.5% for
230 PAR, 6.6% for UV-A and 5.1% for UV-B, relative to **naked** cells (Fig. 1). The
231 specific growth rate of calcifying *E. huxleyi* strain (PML B92/11) was about 2 times
232 higher than that of the **naked** strain (CCMP 2090) ($P < 0.05$) when grown at 500 μmol
233 photons $\text{m}^{-2} \text{s}^{-1}$ of PAR under indoor conditions (Fig. 2A). Growth rates of both
234 strains were significantly ($P < 0.05$) reduced when the cells were transferred outdoor
235 and exposed to natural solar radiation. However, under outdoor conditions, growth
236 rates of calcified cells were 3.5 times higher than those of the **naked** cells, indicating
237 that the latter was more harmed by the solar exposure than the former (Fig. 2A). The
238 cell diameter was not significantly different in the calcified cells between the indoor

删除的内容: non-calcifying

删除的内容: on-calcifying

删除的内容: on-calcifying

242 and outdoor conditions ($P > 0.05$), but an 18% increase was found in the naked cells 删除的内容: on-calcifying

243 after they had grown under the outdoor conditions for 7 days ($P < 0.05$) (Fig. 2B). The

244 maximal quantum yield (F_v/F_m) decreased when the cells were transferred from

245 indoor to the outdoor conditions, reflecting a harmful effect of solar radiation. The

246 decrease of F_v/F_m , however, was much more pronounced in the naked cells (27%) 删除的内容: on-calcifying

247 compared to calcified cells (11%) (Fig. 2C). 删除的内容: calcifying

248 Calcified cells had significantly higher apparent light use efficiency (α), maximal 删除的内容: ,

249 electron transport rate ($rETR_{max}$) and light saturation parameters (I_k) compared with 删除的内容: and higher

250 naked cells. The de-calcified cells of the calcified strain showed a remarkable 删除的内容: , but significantly lower

251 decrease of $rETR_{max}$ ($P < 0.05$), and also alpha and I_k decreased, however 删除的内容: calcifying

252 statistically not significantly (Fig. 3, Table 1). Increased actinic light levels 删除的内容: but did not show obvious changes in α and I_k

253 (acclimating light during the fluorescence measurement) led to higher NPQ in both

254 the calcified and naked strain (Fig. 4). Furthermore, calcified cells showed higher 删除的内容: calcifying

255 NPQ values compared to naked cells ($p < 0.05$). 删除的内容: on-calcifying

256 When exposed to full spectrum solar radiation, the quantum yield of calcified cells 删除的内容: non-calcifying

257 showed no significant change during the first 30 min ($P > 0.05$). After 30 minutes,

258 quantum yield quickly dropped from about 0.35 to 0.22 for ~20 min ($P < 0.05$)

259 followed by a slight recovery in the last 25 minutes. A similar trend was observed in

260 the de-calcified cells with the key difference that the sharp decrease already happened

261 during the first 10 min. Quantum yield of the naked cells decreased constantly for the 删除的内容: on-calcifying

262 first 50 minutes and remained at the low level thereafter (Fig. 5).

263 No effect of the radiation treatment (P, PA and PAB radiation) on the quantum

264 yield of calcified cells was observed after the cells grown under indoor condition were

265 transferred to outdoor solar radiation for 1h exposure (very cloudy day, average PAR,

266 UV-A and UV-B were $481\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, 22.1 and 0.7 W m^{-2} , respectively) ($P >$

280 | 0.05). Quantum yield was significantly higher in the naked cells, however, when they
281 | were exposed to UVA radiation (PA vs. P treatment, $P < 0.05$ Fig. 6A).

删除的内容: on-calcifying

282 | Similar responses were observed when the same test was done on a sunny day with
283 | average PAR, UV-A and UV-B of $1605 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, 69 and 2.4 W m^{-2} ,
284 | respectively. Here, the quantum yield of the calcified cells showed no significant
285 | difference between the different light treatments but it decreased significantly under
286 | PAB treatment compared to P treatments in the naked cells ($P < 0.05$) (Fig. 6B).

删除的内容: on-calcifying

287

288 | **4 Discussion**

289 | Various hypotheses were proposed for the possible functions of coccoliths, but
290 | none of them is supported by sufficient evidence (Young, 1994; Raven and Crawford,
291 | 2012). One important function of coccoliths for surface-dwelling species such as *E.*
292 | *huxleyi* could be the protection against high photon flux densities, especially UV
293 | radiation (Berge, 1962; Young, 1994; Gao et al., 2009).

294 | Some of our results support this hypothesis. The growth rate of the calcified cells of
295 | *E. huxleyi* grown under indoor conditions was about 2 times higher than that of naked
296 | cells. This difference came out even stronger, with growth rates 3.5 times higher in
297 | calcified versus naked cells, when the cells were exposed to full spectrum solar
298 | radiation (Fig. 2A). This could potentially be attributed to the screening of PAR, UV-
299 | A, and UV-B by coccoliths. Although the daytime PAR of solar radiation was
300 | reduced to about half of the light level of the indoor test, noon time PAR levels were
301 | higher than $500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, and the presence of UV could lead to more
302 | harms to the naked cells. Light protection by coccoliths is further supported by the
303 | Fv/Fm measurements. The maximum photochemical efficiency of PSII was only
304 | slightly reduced in calcified cells but significantly decreased in naked cells when they

删除的内容: on-calcifying

308 were exposed to natural solar PAR and UV radiation (Fig. 2C). Furthermore,
309 photochemical performance of de-calcified cells decreased significantly faster and
310 stronger with time compared calcified cells (Fig. 5).

311 The diameter of calcified cells did not significantly change when they were

312 exposed to the full spectrum of solar radiation. The diameter of the naked cells,

删除的内容: on-calcifying

313 however, increased significantly (Fig. 2B). Perhaps, the naked cells experienced more

删除的内容: on-calcifying

314 DNA damage and so did not enter the S phase regularly (Buma et al., 2000).

315 Alternatively, it may reflect a strategy to acclimatize to stressful solar UV radiation

316 since it is well known that smaller cells are usually more sensitive to UV than their

317 larger counterparts (Garcia-Pichel, 1994; Laurion and Vincent, 1998). Some field and

318 laboratory studies showed increased cell size with increased UV exposures (Buma et

319 al., 2000), which can be interpreted as adaptive or acclimation mechanism for

320 protecting the cells against UV radiation. Furthermore, the naked cells might also

删除的内容:

321 employ other strategies such as synthesizing UV screening compounds to ameliorate

删除的内容: Alternatively,

322 UV stress because the naked strain had a lower UV transmittance than the decalcified

删除的内容: another

323 strain.

删除的内容: y

删除的内容: (

删除的内容: s

删除的内容: in the UV than does the calcified strain without coccoliths.

删除的内容: .

324 Several studies found that coccoliths do not protect *E. huxleyi* from excess PAR

325 (Nanninga and Tyrrell, 1996; Houdan et al., 2005; Trimborn et al., 2007). However,

326 UV radiation was not considered in these experiments. Our results showed that the

327 naked cells were more sensitive to full spectrum solar radiation than calcified cells

删除的内容: on-calcifying

328 and even in the same strain, the photochemical performance of de-calcified cells

329 decreased significantly when comparing the calcified cells. This suggests that

330 coccoliths efficiently protect the cells from solar UV radiation.

331 On the other hand, *E. huxleyi* appears to be more sensitive to UV-B irradiances than

332 other phytoplankton species, and its growth rate and physiological performances were

345 highly inhibited by UV radiation (Peletier et al., 1996; Buma et al., 2000; Xu et al.,
346 2011). However, competition tests for community changes are rare, and longer-term
347 experiments with less extreme UVR would be more ecologically and evolutionarily
348 relevant (Raven and Crawford, 2012). In our work, UVR had no significant effect on
349 the quantum yield of calcified cells regardless of high or low light condition but it
350 showed inhibition in naked cells when they were exposed to high solar light (Fig. 6A,
351 B). This provides further evidence for protection by coccoliths against UV radiation.

352 On the cloudy day, no significant difference was observed among the treatments for
353 the calcified cells; on the sunny day, under the fluctuating light (data not shown)
354 calcified cells manage to refurbish damage to their photosynthetic apparatus by
355 balancing damage and repair (Gao et al., 2007; Ragni et al., 2008; Loebel et al., 2010).
356 For the naked cells, on the other hand, UV damage was not effectively repaired,
357 leading to the observed negative effect on photosynthetic performance.

358 It has to be noted that our experimental data is based on only two strains of a naked
359 and calcified *E. huxleyi*. However, similar trends in photophysiology between naked
360 and decalcified in comparison to calcified cells suggest that coccoliths of *E. huxleyi*
361 play an important role in protecting this species against harmful solar radiation
362 especially UV-A and UV-B. Furthermore, the reported absence of photoinhibition in
363 this alga at high light levels also appears to be connected to the coccosphere of *E.*
364 *huxleyi* or its calcification process. In view of ongoing ocean change, the projected
365 shoaling of the upper mixed layer (UML) caused by global warming and progressive
366 ocean acidification that reduces the thickness or the number of coccoliths per cell
367 (Gao et al., 2009; De Bodt et al., 2010) could reduce *E. huxleyi* growth rates within
368 the UML due to increased UVR exposure,

369 Acknowledgements

- 删除的内容: on-calcifying
- 删除的内容: calcifying
- 删除的内容: calcifying
- 删除的内容: on-calcifying
- 删除的内容:
- 删除的内容:
- 删除的内容: In conclusion, the
- 删除的内容: of
- 删除的内容: calcifying
- 删除的内容: calcified
- 删除的内容:
- 删除的内容: Since our data were only based on only one strain of each phenotype, cautions should be taken when drawing conclusion about the differential responses of calcifying and naked
- 删除的内容: strains. In
- 删除的内容: addition, since calcification leads to internal generation of CO₂, such an increased availability of CO₂ might aid to enhance growth, though it has been shown that there is no obligatory coupling between calcification and photosynthesis (Bach et al 2013)
- 删除的内容: .
- 带格式的: 字体: Times New Roman, 小四
- 删除的内容: Nevertheless,
- 删除的内容: T
- 删除的内容: is
- 删除的内容: most likely
- 删除的内容: the photoprotective role played by
- 删除的内容: the ecological implications
- 删除的内容: .
- 删除的内容: With
- 删除的内容: ,
- 删除的内容: d
- 删除的内容: threaten
- 删除的内容: ,
- 删除的内容: cells of
- 删除的内容: living
- 删除的内容: would be impacted
- 删除的内容: daily
- 删除的内容: s to solar radiation

417 This study was supported by National Natural Science Foundation (41430967;
418 41476097; 41120164007), State Oceanic Administration (National Programme on
419 Global Change and Air-Sea Interaction, GASI-03-01-02-04), Joint project of National
420 Natural Science Foundation of China and Shandong province (No. U1406403),
421 Strategic Priority Research Program of Chinese Academy of Sciences (No.
422 XDA1102030204). Visit of KG to Kiel was supported by DAAD. KGS is the
423 recipient of an Australian Research Council Future Fellowship (FT120100384).

424

425

426 **References**

427 Berge, G.: Discoloration of the sea due to *Coccolithus huxleyi* “bloom”, *Sarsia*, 6, 27-
428 40, 1962.

429 Balch W. M., Kilpatrick K. A., and Trees E. E.: The 1991 coccolithophore bloom in
430 the central North Atlantic. 1. Optical properties and factors affecting their
431 distribution. *Limnol. Oceanogr.*, 41: 1669-1683, 1996.

432 Broecker, W., and Clark, E.: Ratio of coccolith CaCO_3 to foraminifera CaCO_3 in late
433 Holocene deeper-sea sediments, *Paleoceanography*, 24, PA3205, 2009

434 Buma, A. G. J., van Oijen, T., van de Poll, W., Veldhuis, M. J. W., and Gieskes, W.
435 W. C.: The sensitivity of *Emiliana huxleyi* (Prymnesiophyceae) to ultraviolet-B
436 radiation, *J. Phycol.*, 36, 296-303, 2000.

437 De Bodt, C., Van Oostende, N., Harlay, J., Sabbe, K., and Chou, L.: Individual and
438 interacting effects of pCO_2 and temperature on *Emiliana huxleyi* calcification:
439 study of the calcite production, the coccolith morphology and the coccosphere
440 size, *Biogeosciences*, 7, 1401-1412, 2010.

441 Gao, K., Ruan, Z., Villafane, V. E., Gattuiso, J. P., and Helbling, E. W.: Ocean

442 acidification exacerbates the effect of UV radiation on the calcifying
443 phytoplankter *Emiliana huxleyi*, *Limnol. Oceanogr.*, 54, 1855-1862, 2009.

444 Gao, K., Helbling, E. W., Häder, D. P., and Hutchins, D. A.: Responses of marine
445 primary producers to interactions between ocean acidification, solar radiation,
446 and warming, *Mar. Ecol. Prog. Ser.*, 470, 167-189, 2012.

447 Gao, K., Wu, Y., Li, G., Wu, H, Villafañe, V. E., and Helbling, E. W.: Solar UV
448 radiation drives CO₂ fixation in marine phytoplankton: A double-edged sword.
449 *Plant Physiol.*, 144, 54-59, 2007.

450 Garcia-Pichel, F.: A model for internal self-shading in planktonic organisms and its
451 implications for the usefulness of ultraviolet sunscreens, *Limnol. Oceanogr.*, 39,
452 1704-1717, 1994.

453 Genty, B., Briantais, J. M., and Baker, N. R.: The relationship between the quantum
454 yield of photosynthetic electron-transport and quenching of chlorophyll
455 fluorescence, *Biochim. Biophys. Acta*, 990, 87-92, 1989.

456 Guan, W., and Gao, K.: Enhanced calcification ameliorates the negative effects of UV
457 radiation on photosynthesis in the calcifying phytoplankter *Emiliana*
458 *huxleyi*, *Chin. Sci. Bull.* 55, 588-593, 2010.

459 Guillard, R. R., and Ryther, J. H.: Studies of marine planktonic diatoms: I. *Cyclotella*
460 *nana* hustedt, and *Detonula confervacea* (cleve) gran, *Can. J. microbial.*, 8, 229-
461 239, 1962.

462 [Hakala-Yatkin, M., Mäntysaari, M., Mattila, H., and Tyystjärvi, E.: Contributions of](#)
463 [visible and ultraviolet parts of sunlight to photoinhibition. *Plant Cell physiol.*, 51:](#)
464 [1745-1753, 2010.](#)

465 Houdan, A., Probert, I., Van Lenning, K., and Lefebvre, S.: Comparison of
466 photosynthetic responses in diploid and haploid life-cycle phases of *Emiliana*

467 *huxleyi* (Prymnesiophyceae), Mar. Ecol. Prog. Ser., 292:139-146, 2005.

468 Kishino, M., Takahashi, M., Okami, N., and Ichimur S.: Estimation of the spectral
469 absorption coefficients of phytoplankton in the sea, Bull. Mar. Biol., 37: 634-642,
470 1985.

471 Laurion, I., and Vincent, W. F.: Cell size versus taxonomic composition as
472 determinants of UV-sensitivity in natural phytoplankton communities, Limnol.
473 Oceanogr., 43, 1774-1779, 1998.

474 [Loebl M., Cockshutt A. M., Campbell D. A., and Finkel Z. V.: Physiological](#)
475 [basis for high resistance to photoinhibition under nitrogen depletion in *Emiliana*](#)
476 [*huxleyi*, Limnol. Oceanogr., 55, 2150-2160, 2010.](#)

477 Moore, T. S., Dowell, M. D., and Franz, B. A.: Detection of coccolithophore blooms
478 in ocean color satellite imagery: a generalized approach for use with multiple
479 sensors, Remote Sens. Environ., 117, 249-263, 2012.

480 Nanninga, H. J., and Tyrrell, T.: Importance of light for the formation of algal blooms
481 by *Emiliana huxleyi*, Mar. Ecol. Progr. Ser., 136, 195-203, 1996.

482 Peletier, H., Gieskes, W. W. C., and Buma, A. G. J.: Ultraviolet-B radiation resistance
483 of benthic diatoms isolated from tidal flats in the Dutch Wadden Sea, Mar. Ecol.
484 Prog. Ser., 135, 163-168, 1996.

485 Poulton, A. J., Adey, T. R., Balch, W. M., and Holligan, P. M.: Relating
486 coccolithophore calcification rates to phytoplankton community dynamics:
487 regional differences and implications for carbon export, Deep-Sea Res. Part II,
488 54, 538-557, 2007.

489 [Ragni, M., Airs, R. L., Leonardos, N., and Geider, R. J.: Photoinhibition of PSII in](#)
490 [*Emiliana huxleyi* \(Haptophyta\) under high light stress: the roles of](#)
491 [photoacclimation, photoprotection, and photorepair, J. Phycol., 44, 670-683,](#)

492 | [2008.](#)
493 | Raven, J. A., and Crawford, K.: Environmental controls on coccolithophore
494 | calcification, Mar. Ecol. Progr. Ser., 470, 137-166, 2012.
495 | Rost, B., and Riebesell, U.: Coccolithophores and the biological pump: responses to
496 | environmental changes, In: Coccolithophores- from molecular processes to
497 | global impact, Thierstein, H. R., and Young, J. R. (eds), Springer, Berlin, 99-125,
498 | 2004.
499 | Trimborn, S., Langer, G., and Rost, B.: Effect of varying calcium concentrations and
500 | light intensities on calcification and photosynthesis in *Emiliana huxleyi*, Limnol.
501 | Oceanogr., 52, 2285-2293, 2007.
502 | [Voss K., Balch W. M., and Kilpatrick K. A.: Scattering and attenuation](#)
503 | [properties of *Emiliana huxleyi* cells and their detached coccoliths, Limnol.](#)
504 | [Oceanogr., 43: 870-876, 1998.](#)
505 | Xu, K., Gao, K., Villafane, V. E., Helbling, E. W.: Photosynthetic responses of
506 | *Emiliana huxleyi* to UV radiation and elevated temperature: roles of calcified
507 | coccoliths, Biogeosciences, 8, 1441-1452, 2011.
508 | Young, J. R.: Functions of coccoliths, In: Coccolithophores, Winter, A., and Siesser,
509 | W. G. (eds), Cambridge University Press, Cambridge, 63-82, 1994.
510 | Zheng, Y., and Gao, K.: Impacts of solar UV radiation on the photosynthesis, growth,
511 | and UV-absorbing compounds in *Gracilaria lemaneiformis* (Rhodophyta) grown
512 | at different nitrate concentrations, J. Phycol., 45, 314-323, 2009.
513 |
514 |
515 |
516 |
517 |

518

519 Figure captions

520 **Figure 1.** Transmission spectra of cells with ~~(calcified strain)~~ and without ~~(calcified~~
521 strain with coccoliths removed artificially, ~~de-calcified strain)~~ coccolith cover and
522 ~~naked~~ cells of *Emiliana huxleyi*.

删除的内容: Cal-C,

删除的内容: ying

删除的内容: Cal-R,

删除的内容: ying

删除的内容: non-calcifying

删除的内容: (N-Cal)

523

524 **Figure 2.** The specific growth rate (μ) (A), diameter (B) and maximum quantum yield

525 (C) of PSII (Fv/Fm) of the calcified and ~~naked~~ cells of *E. huxleyi* grown in indoor and
526 outdoor conditions. Different letters represent significant difference between the
527 indoor and outdoor experiments. Different horizontal lines represent significant
528 difference between the different strains.

删除的内容: (Cal-C)

删除的内容: on-calcifying (N-Cal)

529

530 **Figure 3.** The relative electron rate (rETR) of ~~calcified, de-calcified and naked~~ cells
531 of *E. huxleyi* grown under indoor conditions as function of PAR. The cells had been
532 grown for 12-22 generations under $500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ of PAR.

删除的内容: coccolith-covered

删除的内容: (Cal-C), coccolith-removed (Cal-R) and non-calcifying (N-Cal)

533

534 **Figure 4.** The non-photochemical quenching (NPQ) of ~~calcified and naked~~ cells of *E.*
535 *huxleyi* grown under indoor conditions. Different letters represent significant
536 difference among the light levels. Different horizontal lines represent significant
537 difference among the different type cells.

删除的内容: coccolith-covered (Cal-C) and non-calcifying (N-Cal)

538

539 **Figure 5.** The time course of quantum yield of ~~calcified, de-calcified and naked~~ cells
540 of *E. huxleyi* under full spectrum solar radiation (noontime, average PAR, UV-A and
541 UV-B were $1082 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, 48.1 and 1.6 W m^{-2} , respectively).

删除的内容: coccolith-covered (Cal-C), coccolith-removed (Cal-R) and non-calcifying (N-Cal)

542

删除的内容: calcified (Cal-C) and non-calcifying (N-Cal)

560 | **Figure 6.** The change of quantum yield of the calcified and naked cells of *E. huxleyi*
561 | when transferred from indoor to outdoor conditions, being exposed to PAR alone (P),
562 | PAR+UVA(PA) and PAR+UVA+B(PAB) for 60 min at around noon time. A,
563 | measured under a cloudy day (average PAR, UV-A and UV-B were 481 $\mu\text{mol photons}$
564 | $\text{m}^{-2} \text{s}^{-1}$, 22.1 and 0.7 W m^{-2} , respectively); B, measured under a sunny day (average
565 | PAR, UV-A and UV-B were 1605 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, 69 and 2.4 W m^{-2}). Different
566 | letters represent significant difference among the light treatments. Different horizontal
567 | lines represent significant difference between the different strains.

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606

Table 1. Photosynthetic parameters of relative electron transport rate (Figure 3) as a function of PAR, different letters represent significant difference ($P < 0.05$) among the calcified, de-calcified and naked cells.

	α	rETR _{max}	I _k
<u>Calcified</u>	0.23±0.02 ^a	90.6±9.0 ^a	1010.8±95.0 ^a
<u>De-calcified</u>	0.20±0.01 ^a	73.5±3.5 ^b	986.3±27.4 ^a
<u>naked</u>	0.17±0.02 ^b	42.3±8.5 ^c	621.8±111.1 ^b

删除的内容: he treatments

删除的内容: Cal-C

删除的内容: Cal-R


删除的内容: N-Cal

611

612

613

614



615


616

617

618

619

Fig. 1




620

621

622

Fig. 2


623
624
625
626



627
628
629
630
631
632
633
634
635
636
637
638


Fig. 3

639
640
641
642
643




644
645
646
647
648
649
650
651
652

Fig. 4



653

654



655

656

657

658

Fig. 5


659

660

661

662

663



664
 665
 666
 667
 668
 669
 670
 671

Fig. 6