

Blue carbon stocks in Baltic Sea eelgrass (*Zostera marina*) meadows

Maria Emilia Röhr^{1,2*}, Christoffer Boström¹, Paula Canal- Vergés³, Marianne Holmer²

¹Åbo Akademi University, Faculty of Science and Engineering, Environmental and Marine Biology, Artillerigatan 6, 20520 Åbo, Finland

²University of Southern Denmark, Department of Biology, Campusvej 55, 5230 Odense M, Denmark

³Danish Shellfish Centre, DTU Aqua, Technical University of Denmark, Ørøddevej 80, 7900 Nykøbing Mors, Denmark

* Corresponding author;

Emilia Röhr (mrohr@abo.fi)

Abstract

Although seagrasses cover only a minor fraction of the ocean seafloor, their carbon sink capacity accounts for nearly one-fifth of the total oceanic carbon burial and thus play a critical structural and functional role in many coastal ecosystems. We sampled 10 eelgrass (*Zostera marina*) meadows in Finland and 10 in Denmark to explore seagrass carbon stocks (Corg stock) and carbon accumulation rates (Corg accumulation) in the Baltic Sea area. The study sites represent a gradient from sheltered to exposed locations in both regions to reflect expected minimum and maximum stocks and accumulation. The Corg stock integrated over the top 25 cm of the sediment averaged 627g C m⁻² in Finland, while in Denmark the average Corg stock was over six times higher (4324 g C m⁻²). A conservative estimate of the total organic carbon pool in the regions ranged between 6.98-44.9 t C ha⁻¹. Our results suggest that the Finnish eelgrass meadows are minor carbon sinks compared to the Danish meadows, and that majority of the Corg produced in the Finnish meadows is exported. Our analysis further showed that > 40 % of

the variation in the Corg stocks was explained by sediment characteristics i.e. dry density, porosity and silt content. In addition, our analysis show, that the root: shoot- ratio of *Z. marina* explained > 12 % and contribution of *Z. marina* detritus to the sediment surface Corg pool explain > 10 % of the variation in the Corg stocks. The mean monetary value for the present carbon storage and carbon sink capacity of eelgrass meadows in Finland and Denmark, were 281 and 1809 € ha⁻¹, respectively. For a more comprehensive picture of seagrass carbon storage capacity, we conclude that future Blue Carbon studies should in a more integrative way, investigate the interactions between sediment biogeochemistry, seascape structure, plant species architecture and the hydrodynamic regime.

Keywords: Blue Carbon, eelgrass, seagrass, carbon stock, carbon accumulation, sequestration, carbon sink

40

41 Introduction

42

The atmospheric carbon dioxide (CO₂) enters the ocean via gas-exchange processes at the ocean-atmosphere interface. In the ocean dissolved inorganic carbon is fixed in photosynthesis by primary producers, and released again through respiration. A large percentage of this fixed carbon is stored and sequestered in the sediments of vegetated coastal ecosystems of which the three globally most significant are saltmarshes, mangrove forests and seagrass meadows (Herr et al. 2012). The carbon stored by these ecosystems is known as Blue Carbon (Duarte et al. 2005 Duarte et al. 2013a, Nellemann et al. 2009). Blue Carbon ecosystems function as carbon sinks, in which the rate of carbon sequestered by the ecosystem exceeds the rate of carbon lost through respiration and export.

Seagrass meadows play a critical structural and functional role in many coastal ecosystems (Orth et al. 2006). Although seagrass meadows only cover globally about 300 000 - 600 000 km² of the ocean sea floor, corresponding to 0.1 to 0.2 % of the total area, their carbon sink capacity

55 (capacity of seagrasses to absorb and store carbon in living and dead biomass and in the
56 sediments) may account for up to 18 % of the total oceanic carbon burial (Gattuso et al. 1998,
57 Duarte et al. 2005, Kennedy et al. 2010, Fourqurean et al. 2012). A large portion of the carbon
58 sequestered (captured and stored) by seagrasses is stored in sediments with a conservative
59 value of 10 Pg C in the top 1 meter of seagrass sediments (Fourqurean et al. 2012).
60 Consequently, recent global estimates imply that seagrass sediments store almost 25 200 to 84
61 000 t C km² (Fourqurean et al. 2012). More importantly, carbon in submerged sediments is
62 stored for timescales of millennia while terrestrial soils are usually less stable and only
63 sequester carbon up to decades (Hendriks et al. 2008).

64
65 The coasts of Scandinavia and the Baltic Sea are key distribution areas for eelgrass (*Zostera*
66 *marina* L.) meadows (Boström et al. 2002, Boström et al. 2014). The meadows extend from fully
67 saline (>30) along the Norwegian coast to brackish (5-6) archipelago areas of Finland. This
68 region is estimated to support >6 000 individual meadows covering at least 1 500 – 2 000 km²,
69 which is four times more than the combined eelgrass area of Western Europe (Spalding et al.
70 2003, Boström et al. 2014). Consequently, this region plays a key role in the coastal carbon
71 dynamics, but we presently lack estimates of the role of eelgrass for carbon storage in temperate
72 sediments. Seagrasses are lost at accelerating rates and it has been estimated that 29 % of global
73 seagrass area has disappeared since the initial recording of seagrasses in 1879 (Waycott et al.
74 2009). This decline could have severe consequences on the total capacity of marine ecosystems
75 to store and sequester carbon in addition to the other ecosystem services seagrass meadows
76 provide. Little is known about the magnitude of carbon emissions from degraded seagrasses
77 ecosystems, not to mention its economic implications. A recent study points out, that despite the
78 importance of these ecosystems in the global carbon budget, none of the three Blue Carbon
79 ecosystems have been included in global carbon market protocols (Pendleton et al. 2012).

80

81

82 Seagrasses exhibit marked differences in shoot architecture and grow under variable
83 environmental settings, making direct extrapolations between species and locations difficult.
84 Consequently, there is a pressing need to better understand which factors are causing variability
85 in carbon storage (Corg stocks, carbon stored in living and dead seagrass biomass and
86 sediments) and the capacity of seagrass meadows to sequester carbon (Corg accumulation) in
87 seagrass sediments. Indeed, recent studies show considerable influence of seagrass habitat
88 setting, sediment characteristics and species-specific traits on the variability in carbon storage
89 capacity in seagrass meadows (Duarte et al. 2013a, Lavery et al. 2013, Miyajima et al. 2015).
90 Such differences contribute to uncertainty in local and global estimates of the carbon storage
91 capacity and carbon dynamics in coastal seagrass areas.

92
93 In order to determine seagrass Corg stocks and Corg accumulation, knowledge on the sources of
94 the carbon stored in the sediments is also crucial. The different Corg sources vary in their
95 turnover compared to seagrasses (sources other than seagrasses being typically faster) and
96 volumes of standing stock (typically less) and thus affect the dynamics of the Corg stocks and
97 accumulation (Fry et al. 1977, Kennedy et al. 2004, Kennedy et al. 2010). Seagrasses are known
98 to be enriched in $\delta^{13}\text{C}$ compared to other potentially sources of Corg in the seagrass sediments,
99 such as plankton, macroalgae, allochthonous carbon material, seagrass epiphytes, and benthic
100 microalgae (Kennedy et al. 2004, Kennedy et al. 2010, Fry and Sherr 1984, Moncreiff and
101 Sullivan 2001, Bouillon et al. 2002, Bouillon and Boschker 2006, Macreadie et al. 2014). Thus,
102 the stable isotope signals of seagrasses and other potential Corg sources can be relatively easily
103 and reliably used as a proxy for identification of the origin of Corg in seagrass sediment carbon
104 pool (Kennedy et al. 2010). Unfortunately, the current knowledge base on how these factors
105 interact and influence carbon fluxes and storage is, at best, limited at both local and global
106 scales.

107

108 In this study, we contrast storage, accumulation rates and sources of the accumulated carbon in
109 eelgrass (*Zostera marina*) meadows in two regions differing in salinity, temperature and
110 seagrass productivity, namely Finland and Denmark. Specifically we asked;

- 111 (1) How large is the carbon storage capacity (Corg stocks) of Baltic Sea eelgrass meadows?
- 112 (2) Which are the environmental factors determining the variability of carbon storage (Corg
113 stocks) and accumulation (Corg accumulation) at local and regional scales?
- 114 (3) How do the sediment characteristics influence the carbon storage (Corg stocks) of
115 eelgrass meadows at local and regional scales?
- 116 (4) How much carbon (Corg stocks) is presently stored in Finnish and Danish eelgrass
117 meadows, respectively.
- 118 (5) What is the present and historically lost (only in Denmark) monetary carbon value?

119

120 Materials and methods

121 Study area

122

123 Plant and sediment samples were collected in June-September 2014 from 10 sites in Finland
124 (The Archipelago Sea) and 10 sites in Denmark (Funen and Limfjorden) (Fig. 1). The Baltic Sea
125 sediments are typically mineral sediments consisting of glaciofluvial deposits and only a small
126 fraction of the sediment carbon content consists of carbonates (Leipe et al. 2011). The inorganic
127 carbon content in our samples was low and contributed with 0.5-5 % of total carbon content (n=
128 10 sites per region) and therefore carbonates were not removed from the sediment samples
129 prior to the analysis to avoid analytical errors in low organic samples (Schlacher and Connolly
130 2014). However, when interpreting the data it should be kept in mind, that given the %IC
131 variation in the samples for the different sites (range 0.1-5.78 %; average 3.33%), the
132 inorganic carbon could cause bias in the stable isotope signals of the sediment surface
133 samples (range 0.16-1.17‰; average 0.76‰).

134 The study sites in each region spanned a gradient from sheltered to exposed areas. The
135 Archipelago Sea of southwestern Finland is a shallow (mean depth 23 m), brackish (salinity 5-6)
136 coastal area characterized by a complex mosaic of some 30.000 islands and skerries (Boström et
137 al. 2006, Downie et al. 2013). The region is heavily influenced by human pressures, especially
138 eutrophication, and exhibits naturally steep environmental gradients, as well as, strong
139 seasonality in temperature and productivity (Boström et al. 2014).

140
141 Limfjorden is a brackish water area in the Jutland peninsula connected to both North Sea and
142 Kattegat with salinity ranging from 17 to 35. The Fjord has a surface area of ~1500 km² and a
143 mean depth of 4.7 m (Olesen and Sandjensen 1994, Wiles et al. 2006, Petersen et al. 2013).
144 Funen is located between the Belt Seas in the transition zone where waters from Baltic Sea and
145 Kattegat meet. The salinity of the area ranges between 10 and 25 and the annual mean water
146 temperature ranges from 10-15° C (Rask et al. 1999). This study was conducted in shallow (< 10
147 m) fjords around Funen. Also the Danish areas are heavily influenced by human pressures,
148 especially eutrophication from intense agricultural farming. (DMU; Danmarks
149 Miljøundersøgelser, 2003).

150

151 Field sampling

152

153 All samples were collected from depths of 2.5-3 m by scuba diving. At all sites, three replicate
154 sediment cores (corer: length: 50 cm, diameter: 50 mm) were taken randomly at a minimum
155 distance of 15 m from each other. The corer was manually forced to a depth of 30-40 cm and the
156 sediment between 0-25 cm was used for the analysis. The cores were capped in both ends under
157 water, and kept in a vertical position during transport to the laboratory. Eelgrass production and
158 biomass were measured at all sites from four randomly chosen locations within the eelgrass
159 meadow. In the vicinity of each sediment core, shoot density was counted using a 0.25 m² frame,
160 and above- and belowground biomass samples were collected with a corer (diameter 19.7 cm)

161 and bagged underwater. Additionally, when present, samples of plants and algae (drift algae,
162 other angiosperms, phytoplankton and epiphytes) considered likely carbon sources in the
163 eelgrass meadows were collected from each site for identification and analysis of stable isotope
164 composition. Approximately 10 g wet material was collected for each species. Annual eelgrass
165 production was determined from estimates of previous growth by applying the horizontal
166 rhizome elongation technique (Short and Duarte 2001). From each site, five replicate rhizome
167 samples with the longest possible intact rhizome carefully removed, were collected and
168 transported to the laboratory for further analysis.

169

170 Seagrass variables

171

172 In the laboratory, the above- and belowground biomass was separated and eelgrass leaves and
173 rhizomes were cleaned from epiphytes, detritus and fauna with freshwater and gently scrubbed
174 with a scalpel. All plant material was dried to constant weight (48 h in 60° C). The belowground
175 biomass was separated into living and dead rhizomes and dried separately. Only the living
176 rhizomes were used for the belowground biomass measurements while samples of both living
177 and dead rhizomes were used for analysis of organic carbon (OC) and stable isotopic
178 composition of the organic carbon ($\delta^{13}\text{C}$). The root: shoot-ratio was calculated as the ratio
179 between below- and aboveground biomasses of *Z. marina* samples. A pooled sample of 2
180 youngest leaves from 10 randomly selected shoots were collected prior to drying from the
181 aboveground biomass samples and dried separately for analysis of OC and $\delta^{13}\text{C}$. All samples
182 were analyzed by Thermo Scientific, delta V advantage, isotope ratio mass spectrometer. The
183 measured isotope ratios were represented using the δ - notation with Vienna Peedee belemnite
184 as reference material.

185

186 Determination of annual eelgrass production was done by measuring length of each individual
187 internode of the rhizomes to the nearest millimeters. To obtain an estimate of the mean annual

production per site, internode length measurements of individual replicates (n= 5) were pooled. Due to lack of two annual production peaks in both regions the annual production was estimated based on the distance between shortest and longest measured internodes, assuming that they represent the time point when the water temperature was at its minimum and maximum average, respectively. The time points for the water temperatures were obtained from databases of the Finnish and Danish Meteorological Institutes, respectively.

Sediment variables

In the laboratory, sediment samples were sliced into sections of 2-5 cm, where the upper 10 cm layer was divided into 2 cm layers and the remaining part in 5 cm layers. From each subsample visible plant parts and fauna were removed before the sediment was homogenized. From the 0-2 cm section a subsample of 20 ml was taken for grain size analysis by a Malvern Mastersizer 3000 particle size analyzer. The sediment silt content was calculated as the fraction with particle size of 2-63 μm from the range of all particle sizes (Folk and Ward 1957). Sediment water content, dry bulk density and porosity were determined from a subsample of 5 ml that was taken using a cut-off 5 ml syringe and weighed before and after drying at 105°C for 6 h from all sediment layers. The dried sediment samples were homogenized in a mortar and divided into two subsamples from which one was used for analysis of sediment organic matter content (OM, as loss of ignition: 4 h in 520°C), and the other for analysis of sediment $\delta^{13}\text{C}$ and OC as described above for the plant materials. Inorganic carbon content was low in sediments from both regions (< 5 %) and considered insignificant compared to the organic fraction (1-2 order of magnitude higher).

Corg stock and Corg accumulation calculations

214 The depth integrated Corg stocks were calculated according to Lavery et al. (2013) by
215 multiplying the OC (OC mg gDW⁻¹) measured from different sections of the sediment core with
216 the corresponding sediment dry density (g cm⁻³). These numbers were then depth integrated
217 over 25 cm in order to estimate the depth integrated carbon density. To estimate sediment Corg
218 stock and Corg accumulation of Finnish and Danish eelgrass area we used averages from 10 sites
219 from each region in our calculations. The Corg (obtained by depth integration of the carbon
220 density (mg C cm⁻³) of the sampled region was multiplied with estimated seagrass area of the
221 region based on the most recent areal estimates (in km²) of seagrass distribution available in the
222 literature (Boström et al. 2014) and given as Corg in g C m⁻². In Finland the estimated areal
223 extent was 30 km², while in Denmark the extrapolations were based on the minimum and
224 maximum estimates of the areal extent, respectively (673 and 1345 km²; Boström et al. 2014).
225 Results for carbon accumulation (applied by multiplying the depth integrated Corg stock,
226 regional seagrass area and sediment accumulation rate estimate from literature) in each area
227 are given as Corg accumulation (t y⁻¹). Due to lack of long term monitoring of sediment
228 accumulation in eelgrass meadows, we used available minimum, average and maximum
229 sediment accumulation rates in seagrass meadows obtained from literature (Duarte et al. 2013b,
230 Serrano et al. 2014, Miyajima et al. 2015).

231
232 To calculate the total Corg pool in Danish and Finnish eelgrass sediments, we summed the
233 following three components: (1) the annual areal eelgrass carbon accumulation rate (Corg
234 accumulation in t C ha⁻¹y⁻¹, calculated by dividing the measured Corg stocks (C g DW m⁻²) in each
235 region with the time that it takes to accumulate this stock with a sedimentation rate of 2.02 mm
236 y⁻¹), (2) the total C in the average living aboveground and belowground *Z. marina* tissue (t C ha⁻¹
237 ¹), and (3) the mean Corg stocks (t C ha⁻¹) in eelgrass sediments in Denmark and Finland,
238 respectively. To calculate the present and lost economic value of eelgrass carbon stocks, we used
239 the social cost of carbon of 40.3 € t C⁻¹ (United States Government 2010) and multiplied this
240 value with the Corg stocks (t C km⁻²). To estimate the Danish eelgrass losses over the past 100

241 years in economic terms, we used the calculations above, but accounted for the annually lost
242 sequestration value by multiplying the rate by 100. We used the most recent loss estimates for
243 Denmark for the period 1900-2000, assuming that the present coverage constitutes 10 % or 20
244 % of the historical area, respectively (Boström et al. 2014).

245

246 Sediment carbon sources

247

248 The Isosource 1.3 isotope mixing model software (Phillips and Gregg 2003) was used to estimate
249 the contribution of different carbon sources to the sediment surface Corg stock. We ran the
250 Isosource model using the $\delta^{13}\text{C}$ obtained from stable isotope analysis of *Z. marina* leaves, living
251 and dead rhizomes and for samples of other abundant Corg sources within the meadow (n=1-5)
252 with increments of 1 % and tolerance of 0.1. Numbers are given as percentage contribution to
253 the sediment surface carbon pool.

254

255 Data analysis

256

257 All statistical analyses were performed using the PRIMER 6 PERMANOVA+ package (Anderson
258 et al. 2008). A 2-factor mixed model was used, where sampling sites and region (FIN, DEN) were
259 used as fixed factors for the biological response variable (sediment organic carbon stock, g C m⁻²).
260 In addition we ran reduced, countrywise DistLm models in order to better address the
261 possible differences in regional environmental drivers for Corg stock. Prior to analysis, the
262 environmental predictor variables (degree of sorting, sediment dry density, sediment water
263 content, sediment porosity, sediment silt content, sediment organic content, annual production,
264 root: shoot-ratio, shoot density and percentage of *Z. marina* detritus contribution to Corg) were
265 visually inspected for collinearity using Draftsman plots of residuals. Due to autocorrelation
266 between sediment variables (water content, porosity and dry density) sediment water content
267 was removed from the environmental variables. To achieve normality in the retained

environmental variables, data was log-transformed ($\log(X+1)$) and Euclidean distance was used to calculate the resemblance matrix. The biological response variable (Corg stock in g C m^{-2}) was square-root transformed and Bray-Curtis similarity was used to calculate the abundance matrix. The relative importance of different environmental variables was determined by use of DistLm, a distance-based linear model procedure (Legendre and Anderson 1999). DistLm model was constructed using a step-wise procedure that allows addition and removal of terms after each step of the model construction. AICc (Akaike Information Criterion with a correction for finite sample sizes) was chosen as information criterion as it enables to fit the best explanatory environmental variables from of relatively small biological dataset compared to number of environmental variables (Burnham and Anderson 2002). An alpha level of significance of 95 % ($p < 0.05$) was used for all the analysis. All means are reported as mean \pm SE (SEM).

Results

Seagrass meadow and sediment characteristics

In general, the Finnish meadows were found on exposed sandy bottoms while the environmental settings of the eelgrass meadows in Denmark were more variable (Fig. 2). Shoot density was nearly equal in both regions, averaging at 417 ± 75 (shoots m^{-2}) in Finland and 418 ± 32 (shoots m^{-2}) in Denmark. In Finland variation between sites (112-773 shoots m^{-2}) was greater than in Denmark (300-652 shoots m^{-2}). In Denmark the highest shoot density was found at the most exposed site (Nyborg), while in Finland the highest shoot density was found at Sackholm, a fairly sheltered site. The lowest shoot densities in Finland and Denmark were found in Tvärminne and Løgstør, respectively. The mean aboveground biomasses were 101 ± 3 and 145 ± 5 (g DW m^{-2}) and the mean belowground biomasses 79 ± 5 and 148 ± 13 (g DW m^{-2}) at Finnish and Danish sites, respectively. In Denmark, the mean fraction of OC in aboveground and belowground *Z. marina* tissue was 35 ± 0.32 % DW and 29 ± 1.10 % DW, respectively, while the corresponding numbers

for Finland were 38 ± 0.24 % DW and 36 ± 0.27 % DW, respectively. Given an average total *Z. marina* biomass (above- and belowground) of 293 ± 22.31 (Denmark) and 180 ± 9.60 g DW m⁻² (Finland), we estimate the Corg pool in bound in living seagrass biomass to 0.66 ± 0.005 and 0.94 ± 0.014 t C ha⁻¹ in Finland and Denmark, respectively. The root:shoot-ratio was slightly lower in Finland (0.87 ± 0.05) than in Denmark (1.14 ± 0.12), and varied between 0.29 to 3.29 and 0.15 to 6.45 in Finland and Denmark, respectively. The annual production of eelgrass for Finland (average 524 ± 62 g DW m⁻² y⁻¹) showed relatively low variation between sites (270-803 g DW m⁻² y⁻¹) being lowest at Jänisholm and highest at Ryssholmen. In Denmark, the mean annual eelgrass production was almost twice as high (928 ± 159 g DW m⁻² y⁻¹) with large variation (470-2172 g DW m⁻² y⁻¹). Production was lowest and highest at Dalby and Visby, respectively (Table 1).

The sediment characteristics varied significantly between Finland and Denmark. There was a significant difference ($F_{1,9} = 14.7$, $p < 0.003$) between regions in terms of silt content, which was generally lower at Finnish (6.3 ± 1 %) sites than at Danish sites (20.2 ± 3.9 %), although in Denmark the variation between sites ranged from 0.8 % at Nyborg to 31.6 % at Thurøbund (Table 1, Fig. 2). In Finland, the variation between sites was lower and ranged from 1.6% (Kolaviken) to 15.5 % (Sackholm). At the Finnish sites the mean sediment dry density was higher (1.35 ± 0.01 g cm⁻³) compared to the Danish sites (1.25 ± 0.02 g cm⁻³), and the Finnish sites exhibited lower within-region variability ranging from 1.1 g cm⁻³ at Lyddaren to 1.5 cm⁻³ at Långören, while the Danish sites varied from 0.3 g cm⁻³ at Thurøbund to 1.5 g cm⁻³ at Visby. The Finnish sites showed consistently lower pools of organic matter (LOI: 1.4 ± 0.3 %DW) compared to the average of Danish sites (LOI: 3.9 ± 1.5 % DW). Similarly, the mean OC content was lower in Finland (0.24 ± 0.033) than in Denmark (1.75 ± 0.563). Consequently, the mean water content was similarly lower in Finland (20.9 ± 0.4 %: range 16-29 %) than in Denmark (37.4 ± 1.8 %: range 17-76 %) (Table 1). Sediment porosity was similar in both regions, and ranged from 0.25 to 0.30 in Finland, and from 0.20 to 0.40 in Denmark. At the Finnish sites, the proxy (degree of sorting)

that was used to estimate exposure, varied from 0.8 to 1.5 (ϕ), with Kolaviken being the most exposed and Ängsö being the most sheltered site. In Denmark degree of sorting varied from 0.6 to 2.1 (ϕ) with Nyborg and Visby being the most exposed and sheltered sites, respectively (Fig. 2).

Organic carbon stocks

The profiles of carbon densities (g C cm^{-3}) in the upper 25 cm of the sediment showed marked differences both between and within the sampled regions. At the Finnish sites, where eelgrass typically grows at exposed locations, the sediment carbon density was low (mean $2.6 \pm 0.09 \text{ mg C cm}^{-3}$) and declined with depth at most of the 10 study sites (Fig. 3). At the Danish sites, however, the sediment carbon density was more variable (mean $17.45 \pm 9.42 \text{ mg C cm}^{-3}$) both within and between sites (Fig. 3). Depth integrated Corg stocks (0-25 cm, g C m^{-2} , Fig. 4) were particularly high at one sheltered site in Funen, namely Thurøbund. This site is characterized by soft sediments with high organic content, high annual eelgrass production and high belowground biomass (Table 1). The lowest eelgrass Corg stocks in Denmark were found at two relatively exposed and sandy sites, namely Nyborg and Dalby (Fig. 4). The estimate of average total Corg stock in Finland was $0.019 \pm 0.001 \text{ Mt C}$ taking the total area of eelgrass into account (30 km^2 ; Table 2). Using minimum and maximum estimates of the eelgrass area in Denmark the estimates for mean total sediment Corg stock in Denmark were $2.164 \pm 0.005 \text{ Mt C}$ or $5.868 \pm 0.014 \text{ Mt C}$, respectively (673 and 1345 km^2 ; Table 2).

Using an annual carbon accumulation value of 0.05 and 0.35 $\text{t C ha}^{-1} \text{ y}^{-1}$ for Finland and Denmark, respectively, and assuming sediment accumulation of 2.02 mm y^{-1} on average (Table 2), the total pool of Corg in the *Z. marina* meadows (Corg bound in living biomass, sediment Corg stock and Corg accumulation) corresponds to 6.98 t C ha^{-1} (698 t km^{-2}) and 44.9 t C ha^{-1} (4490 t km^{-2}) for Finland and Denmark, respectively. Using the social cost of carbon of 40.3 € t C^{-1}

(United States Government 2010), the present economic value of eelgrass carbon in Finnish and Danish eelgrass meadows is estimated at 281 and 1809 € ha⁻¹, respectively. Using an average of these values (1045 € ha⁻¹) and a conservative estimate of the eelgrass acreage in the Baltic Sea (2100 km²: Boström et al. 2014), we estimate a total monetary value of the present sequestration by eelgrass meadows to be 219.4 million euro. Given the total eelgrass loss in Denmark for the time period 1900-2000 is between 5381 km² (present area 20 % of historical distribution) and 6053 km² (present area 10 % of historical distribution), this equals to a Corg loss of 0.042 and 0.048 Gt C, respectively. Using the acreage value (1045 € ha⁻¹) these areal loss estimates corresponds to a lost economic value between 562 and 632 million euro, for the minimum and maximum areal loss estimates, respectively.

Corg accumulation

The estimates for annual Corg accumulation in the Finnish seagrass meadows (30 km²) were low (0.002, 0.016, 0.033 Mt C y⁻¹), when applying sediment accumulation rates of 0.32, 2.02 and 4.20 mm y⁻¹, respectively. The low Corg accumulation in Finnish meadows was a result of low mean Corg stocks and relatively small size of seagrass area compared to Denmark (Table 2). The estimates for annual Corg accumulation for the Danish sites differed between the two sub-regions Limfjorden (18 km²) and Funen (179 km²). At the sampling sites around Funen, the Corg accumulation was 0.139, 0.881 and 1.832 Mt C y⁻¹, while in Limfjorden the Corg accumulation was lower (0.006, 0.038 and 0.079 Mt C y⁻¹) and similar to Corg accumulation for Finland. Using upper and lower eelgrass areal estimates, total Corg accumulation based on 3 sediment accumulation rates in Denmark were more than four orders of magnitude higher (0.376, 2.373, 3.636 and 0.75, 4.741 and 9.859 Mt C y⁻¹) than the estimated total Corg accumulation in Finnish eelgrass meadows.

374 Carbon sources

375

376 The $\delta^{13}\text{C}$ values of the surface sediment within regions where quite homogenous ranging from -
377 22.8 to -18.9 ‰ and -17.6 to 13.5 ‰, in Finland and Denmark respectively. The analytical error
378 for the sediment $\delta^{13}\text{C}$ values was 2.8 ‰. The $\delta^{13}\text{C}$ in *Z. marina* tissues ranged from -11.4 to -8.5
379 ‰ and from -12.5 to -8.2‰, in Finland and Denmark, respectively. There was no significant
380 difference between living above- and belowground tissue and decomposed belowground tissue
381 and samples were pooled in the isotope mixing model. Although *Z. marina* was the dominant
382 seagrass species in Finland, the study sites included both monospecific and mixed seagrass
383 meadows. Mixed meadows typically contained pondweeds, e.g. *Potamogeton pectinatus* and
384 *Potamogeton perfoliatus*. In particular, *P. pectinatus* ($\delta^{13}\text{C}$ -11.3 to -7.6 ‰) and *P. perfoliatus*
385 ($\delta^{13}\text{C}$ -15.6 to -12.6 ‰) were both present at five of the Finnish study sites (Jänisholm,
386 Sackholm, Hummelskär, Tvärminne and Fårö) and *P. pectinatus* was present at Kolaviken,
387 Ryssholmen and Lyddaren. *Ruppia cirrhosa* (-11.5 to -8.8 ‰) was less abundant and found at
388 three of the Finnish sites (Sackholm, Ängsö, Kolaviken) and at one study site in Denmark
389 (Kertinge). The $\delta^{13}\text{C}$ for phytoplankton ranged from -24.6 to -22.6 ‰ and -18.6 to 16.4 ‰, in
390 Finland and Denmark, respectively. Drift algae was present at all Danish study sites, except
391 Thurøbund, and had $\delta^{13}\text{C}$ values from -17.9 to -13.5 ‰, but only at five Finnish sites (Ängsö,
392 Ryssholmen, Fårö, Långören and Hummelskär) with $\delta^{13}\text{C}$ values ranging from -20.0 to -16.3‰.

393

394 The isotope mixing model indicated that at all Finnish sites, phytoplanktonic material was the
395 major contributor (43-86 %) to the sediment surface Corg pool. In Denmark *Z. marina*
396 contributed with 13-81 % to the sediment surface Corg pool, contribution being lowest at the
397 most exposed site in Nyborg and highest in Visby. The corresponding numbers for Finland were
398 1.5-32 %, being lowest and highest in Tvärminne and Lyddaren, respectively (Fig. 5).

399

400 Environmental factors explaining carbon pools

401
402 The combined (FIN + DK) DistLm analysis showed, that three sediment variables (dry density,
403 silt content, porosity) and three plant variables (annual eelgrass production, the root:shoot-ratio
404 and *Z. marina* contribution to the sediment carbon pool) explained 67 % of the variation in the
405 sediment Corg stock (g C m^{-2}) (Table 3 and 4, Fig. 6). Specifically, sediment silt content alone
406 explained > 36 % of the variation in Corg stocks (Table 3). In both regions, exposed sites
407 characterized by sandy, low organic sediments and low silt content, had low Corg stocks. In
408 contrast, at sheltered sites like Thurøbund in Denmark, we measured the highest sediment Corg
409 stock along with highest silt and water content among all sites. Although sediment porosity and
410 sediment dry density also contributed to the model, they were of minor importance (~2 %
411 each).

412 The combined (FIN+ DK) DistLm analysis also showed that the *Z. marina* contribution to the
413 sediment surface carbon pool explained 10.9 % of the variation in the measured Corg stocks
414 (Fig. 6, Table 3 and Table 4). Drift algae was a significant contributor (72 %) to the sediment
415 surface Corg pool at the Danish sites, while it appeared to play only a minor role (0-21 %) in
416 Finland. The carbon sources were generally more mixed at the Danish study sites compared to
417 the Finnish sites where phytoplankton dominated (Fig. 5).

418
419 While the overall model including all sites explained almost 70 % of the variation in carbon
420 stocks (Table 3, 4), and indicated that the most relevant environmental variables were included
421 in this Baltic scale analysis, reduced, countrywise DistLM models revealed different results. In
422 particular, variability in Finnish carbon stocks were explained up to 50 % by geological variables
423 (porosity, sorting and sediment dry density), while the best sequential model for carbon stock
424 variability at Danish sites explained 75 % of the total variance. In contrast to Finland, the role of
425 eelgrass related variables (relative proportion of *Z. marina* in the sediment and the root:shoot

ratio) where most important and explained 40 % and 25 %, respectively of the carbon stock variability.

Discussion

Recent studies have shown considerable variation in the global estimates of carbon stocks (Corg stocks) and carbon accumulation rates (Corg accumulation) in seagrass meadows, indicating an incomplete understanding of factors influencing this variability (Fourqurean et al. 2012, Duarte et al. 2013a, Lavery et al. 2013, Miyayima et al. 2015). The Baltic Sea forms a key distribution area for eelgrass in Europe, but similarly to the global data sets, we have so far lacked estimates on seagrass carbon stocks and accumulation.

In our study, the Finnish eelgrass meadows showed consistently very low Corg stocks and Corg accumulation, and the meadows were minor carbon sinks compared to the Danish meadows. The Danish sites showed more variation in the sediment Corg stock and accumulation and Corg stock was particularly high at one site, Thurøbund ($26138 \pm 385 \text{ g C m}^{-2}$), which is a relatively sheltered site with high organic sediments. Expectedly, due to both larger overall eelgrass acreage and larger Corg stocks in the Danish meadows, the total Corg accumulation ($0.38\text{-}9.86 \text{ t C y}^{-1}$) was three to four orders of magnitude higher than in the Finnish meadows ($0.002\text{-}0.033 \text{ t C y}^{-1}$). As eelgrass in Finland generally grow in more exposed locations potentially due to increased interspecific competition with freshwater plants such as common reed (*Phragmites australis*) in sheltered locations (Boström et al. 2006), it is probable that most of the Corg produced in the Finnish meadows is exported, and thus incorporated in detrital food webs in deeper bottoms. This argument is supported when applying sediment accumulation rates from literature, as only 0.15 – 2.0 % of the annual production accumulated in Finnish meadows, while the corresponding numbers for Denmark were 0.6 -7.8 %. Duarte and Cebrian (1996) estimated

that on average 25 % of the global seagrass primary production is exported, and seagrass detritus may thus contribute significantly to Corg stocks in other locations, a fact that is often overlooked.

Extrinsic drivers of carbon sequestration in seagrass meadows

As proposed in previous work accumulation of fine grained size fractions in seagrass sediments, relative to those accumulated in bare sediments, appears to be one of the major factors influencing the carbon sink capacity of seagrass meadows (Kennedy et al. 2010, Miyajima et al. 2015), and may thus be a useful proxy for the sink capacity. In addition, it is well known, that seagrasses modify sediments by reducing water flow and consequently increasing particle trapping and sedimentation and reducing resuspension (Fonseca and Fisher 1986, Fonseca and Cahalan 1992, Gacia et al. 2002, Hendriks et al. 2008, Boström et al. 2010) and also increasing Corg (Kennedy et al. 2010).

In this study, the DistLm analysis showed, that contribution of *Z. marina* to the sediment surface carbon pool was an important driver (> 10.9 %) of the variation in the sediment Corg stock (Table 3 and 4, Fig. 6) when the model included both regions. Surprisingly, the reduced countrywise analysis revealed different results and showed that *Z. marina* contribution to the sediment surface carbon pool was an important driver for Corg stocks only in Denmark. We believe that the countrywise differences in explanatory variables might relate to the more pronounced influence of eelgrass for carbon stocks at Danish sites. Indeed, these sites exhibited on average 30 % higher aboveground biomasses, 45 % higher belowground biomasses, 24 % higher root:shoot- ratios and 44 % higher productivity compared to the Finnish sites. In Finland and the northern Baltic Sea, eelgrass meadows appears to be primarily physically controlled, and thus sediment variables play a relatively more pronounced role. The results from the model

were also supported by our data in which we found increasing Corg stocks at the Danish sites, where *Z. marina* was the major source of organic carbon, contributing with 13-81% to the surface sediment Corg, while in contrast, at the Finnish sites where only a minor fraction of carbon buried in sediments derive from eelgrass detritus (1.5-39.6 %) the Corg stocks were low. Correspondingly, the average $\delta^{13}\text{C}$ value (-16.2 ‰) in the Danish sediment samples was similar to the global median value (-16.3±0.2 ‰) reported by Kennedy et al. (2010) in which on average 51 % of the carbon was derived from seagrass detritus, whereas it was -20±0.6 ‰ in Finland indicative of higher contribution from other more negative carbon sources., such as phytoplankton. The importance of the *Z. marina* contribution to the Corg stocks may be explained by slow decomposition rates of seagrass tissue. Especially, the high proportion of refractory organic compounds in the seagrass belowground parts and high C:N:P-ratios of seagrass tissue in general make seagrasses less biodegradable than most marine plants and algae (Fourqurean and Schrlau 2003, Vichkovitten and Holmer 2004, Kennedy and Björk 2009, Holmer et al. 2011, Röhr et al in prep.). The slow decomposition rates are also a result of reduced sediment conditions commonly encountered in Danish seagrass meadows (Kristensen and Holmer 2001, Holmer et al. 2009, Pedersen et al. 2011). Despite the extensive distribution (2-29 ha), high biomasses (300-800 g DW m⁻²) and major impact of drifting algal mats on coastal ecosystem functioning (Norkko and Bonsdorff 1996, Salovius and Bonsdorff 2004, Rasmussen et al. 2013, Gustafsson and Boström 2014), the stable isotope composition of the sediments suggests that drift algae had a surprisingly minor influence on the sediment surface Corg pool in both regions. Thus, despite present on several sampling sites, drift algae is likely exported and mineralized in deeper sedimentation basins. Furthermore, we found that at all study sites in both regions, there were several other potential sources influencing the sediment surface Corg pool. Similarly, Bouillon et al. (2007) showed that in seagrass sediments adjacent to mangrove forests in Kenya, none of their sites had seagrass material as the sole source of Corg, and instead mangrove-derived detritus contributed significantly to the seagrass sediment Corg pool.

Similarly to the contribution of *Z. marina* to the sediment surface carbon pool, the root:shoot-ratio explained 12.7 % of the variation in the Corg stocks when both regions were included in the model, but in the reduced countrywise models it was important driver for Corg stocks in only Denmark. Accordingly, the highest Corg stocks, belowground biomass and root: shoot-ratio was found in Thurøbund (Denmark). In Finland, the highest root: shoot-ratio (2.07) was found at Kolaviken, with a relatively low Corg stock (397 g C m⁻²). Due to higher degree of exposure at the site (degree of sorting 0.7 ϕ) compared to Thurøbund (1.4 ϕ) it is likely that large portion of the eelgrass production was exported away from the meadow and not stored in the sediment. The mean shoot densities were almost identical between regions, and shoot density did not contribute to the model explaining Corg.

The annual eelgrass production explained only 2.3 % of the variation in the Corg stocks in the combined model. The annual production rates were almost twice as high at Danish sites compared to the Finnish sites. Regional differences in seagrass productivity may be caused by differences in e.g. the inorganic carbon concentration in water column and light availability between the regions (with higher values in Denmark), which both affect the photosynthetic capacity of the plant (Hellblom and Björk 1999, Holmer et al. 2009, Boström et al. 2014). Eelgrass production tend to be higher in physically exposed areas compared to more sheltered areas, which can be due to improved sediment oxygen conditions and hydrodynamical effects (Hemminga and Duarte 2000). This finding was not supported by our study, in which we found the highest annual eelgrass production rates at both the most sheltered and exposed sites, namely Visby and Nyborg (DK).

Geographical comparisons of carbon stocks and accumulation

Our estimated Corg stocks for the study sites were generally lower (627-4324 t C km⁻²) than estimates (25200-84000 t C km⁻²) found in the literature (Duarte et al. 2005, Nellemann et al.

2009, Mcleod et al. 2011, Fourqurean et al. 2012). In Duarte et al. (2005) the data set used for the calculations was gathered from various studies conducted at different temporal scales and habitat types, as well as different methods for determination of Corg accumulation. Additionally, several of the studies were conducted in the Mediterranean *P. oceanica* meadows - a habitat with superior carbon sequestration and storage capacity (Duarte et al. 2005, Lavery et al. 2013). The average sizes of Corg stocks in Finnish and Danish eelgrass meadows were also considerably lower than the mean values reported by Alongi et al. (2014) for tropical seagrass meadows (14270 t C km⁻²). In contrast, our estimate for the carbon stock in the top 25 cm for Danish and Finnish meadows (627-6005 g C m⁻²) are comparable to Australian (262-4833 g C m⁻²; Lavery et al. 2013) and Asian estimates (3800-12000 g C m⁻²; Miyajima et al. 2015).

Consequences of seagrass loss for carbon pools

Despite the importance of seagrasses, their global distribution has decreased by 29 % since 1879 primarily due to anthropogenic pressures (Waycott et al. 2009), thus weakening the carbon sink capacity of marine environments to sequester carbon (Duarte et al. 2005). Since the 1970s, the Baltic Sea has been subject to strong anthropogenic pressures (Conley et al. 2009) leading to eelgrass declines in several countries (Boström et al. 2014). In the 1930s, the Danish eelgrass meadows were significantly reduced by the wasting disease (Rasmussen 1977). These regime shifts in Denmark have resulted in a 80-90 % decline corresponding to 6726 km² in the beginning of 1900's to 673-1345 km² in 2005, using the minimum and maximum estimates for the current coverage area, respectively (Boström et al. 2014). Using the mean carbon density (17.45 mg C cm⁻³) measured at the Danish sites, the lost Corg stock is estimated to 23-27 Mt C and these large-scale seagrass declines, which are also found in Sweden and Germany, have eroded the Corg stocks in the Baltic Sea significantly (Table 2). In Finland there is a lack of long-term monitoring, but the meadows appear to be stable and cover at least 30 km² with no significant loss of Corg stocks.

559

560 Using a carbon monetary value of 40.3€ C⁻¹, we calculated the monetary value of the present
561 carbon storage and sequestration capacity of eelgrass meadows in Finland and Denmark to be
562 281 and 1809 € ha⁻¹, respectively. Pendleton et al. (2014) calculated a global estimated
563 economic cost of lost seagrass meadows to be 1.9-13.7 billion USD. This value was derived from
564 the cost of lost carbon sink capacity, ignoring other lost ecosystem services including e.g. coastal
565 protection, water quality management, food provision and the role of seagrasses as fisheries and
566 key habitats for marine species (Barbier et al. 2011, Atwood et al. 2015). Our estimate also only
567 considers lost carbon sink capacity and can be compared directly with Pendleton et al. (2014).
568 The present economic value of carbon storage and sequestration capacity of Baltic Sea eelgrass
569 meadows is thus between 1.7 and 12 % out of the global seagrass Blue Carbon value.

570

571 While useful, our and previous work still remain snap shots of complex processes causing local
572 and regional variability in estimates of seagrass Blue Carbon stocks and accumulation. Clearly, in
573 order to produce more reliable estimates of global seagrass carbon sequestration rates and
574 stocks, there is a need for more studies integrating and modeling the individual and joint role of
575 e.g. sediment biogeochemistry, seascape structure, plant species architecture and hydrodynamic
576 regime. Since seagrasses are lost at accelerating rates (Waycott et al. 2009), there is also an
577 urgent need for a better understanding of the fate of lost seagrass carbon (Macreadie et al. 2014)
578 and the development of the carbon sink capacity in restored seagrass ecosystems (Nellemann et
579 al. 2009, Greiner et al. 2013, Marba et al. 2015). Nelleman et al. (2009) proposed the use of
580 carbon trading programs using financial incentives for forest conservation, such as REDD+
581 (Reduced Emissions from Deforestation and Degradation) and NAMAs (Nationally Appropriate
582 Mitigation Actions), to include the blue carbon ecosystems as part of their environmental
583 protection protocol. Both of these carbon mitigation programs require ongoing monitoring of
584 organic carbon storage and emission in the different Blue Carbon ecosystems. In order to
585 manage seagrass meadows, mitigate climate change and produce information required for the

586 carbon trading programs, it is fundamental to understand factors influencing the capacity of
587 seagrass meadows to capture and store carbon. By solving these uncertainties, the conservation
588 and restoration of seagrass meadows can be implemented in the most beneficial manner by e.g.
589 giving priority to protection of the seagrass meadows and species with the highest carbon sink
590 capacity and foundation of restoration projects in areas most suitable for seagrass growth
591 (Duarte et al. 2013a).

592

593 Copyright statement

594

595 The article is distributed under the [Creative Commons Attribution 3.0](#) License. Unless otherwise
596 stated, associated published material is distributed under the same licence. Authors
597 grant Copernicus Publications a license to publish the article and identify itself as the original
598 publisher. Authors grant Copernicus Publications commercial rights to produce hardcopy
599 volumes of the journal for purchase by libraries and individuals. Authors grant any third party
600 the right to use the article freely under the stipulation that the original authors are given credit
601 and the appropriate citation details are mentioned.

602

603 Acknowledgements

604

605 This work is part of a double degree program between Åbo Akademi University (ÅAU) and
606 University of Southern Denmark (SDU). The study was funded by The Maj and Tor Nessling
607 Foundation (Project 201600125: Baltic Sea blue carbon: environmental gradients influencing
608 the carbon sink capacity of seagrass meadows) and University of Southern Denmark. We
609 acknowledge Archipelago Centre Korpoström and University of Southern Denmark for excellent
610 working facilities. Tiina Salo and Karine Gagnon are acknowledged for field assistance and
611 Katrine Kierkegaard and Birthe Christensen for their assistance in the laboratory work.

Field Code Changed

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639
640
641
642
643
644
645
646
647
648

References

- Alongi, D. M.: Carbon Cycling and Storage in Mangrove Forests. *Ann. Rev. Mar. Sci.* 6:195–219, 2014.
- Anderson, M.J., Gorley, R.N. and Clarke, K.R.: PERMANOVA for PRIMER: guide to software and statistical methods. Plymouth, United Kingdom: PRIMER-E Ltd., 2008.
- Atwood, T.B., Connolly, R.M., Euan, G.R., Lovelock, C.E., Heithaus, M.R. and Hays, G.C, Predators help protect carbon stocks in blue carbon ecosystem. *Nat. Clim. Change* 2015.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. and Silliman, B.R.: The value of estuarine and coastal ecosystem services. *Ecol. Monogr.* 81(2):169-193, 2011.
- Boström C., Baden S., Bockelmann A., Dromph K., Fredriksen S. and Gustafsson C.: Distribution, structure and function of Nordic eelgrass (*Zostera marina*) ecosystems: implications for coastal management and conservation. *Aq. Cons. Mar. Freshwater Ecosyst.* 24(3):410-434, 2014
- Boström, C., Törnroos, A. and Bonsdorff, E.: Invertebrate dispersal and habitat heterogeneity: Expression of biological traits in a seagrass landscape. *J. Exp. Mar. Biol. Ecol.*, 390(2):106-117, 2010.
- Boström, C., Bonsdorff, E., Kangas, P. and Norkko, A.: Long-term changes of a brackish-water eelgrass (*Zostera marina* L.) community indicate effects of coastal eutrophication. *Estuar. Coast. and Shelf Sci.* 55(5):795-804, 2002.
- Boström, C., O'Brien, K., Roos, C. and Ekebom, J.: Environmental variables explaining structural and functional diversity of seagrass macrofauna in an archipelago landscape. *J. Exp. Mar. Biol. Ecol.* ,335(1):52-73, 2006
- Bouillon S. and Boschker H.T.S.: Bacterial carbon sources in coastal sediments: a cross-system analysis based on stable isotope data of biomarkers. *Biogeosciences*, 3(2):175-185, 2006.
- Bouillon, S., Dahdouh-Guebas, F., Rao, A.V.V.S., Koedam, N. and Dehairs, F.: Sources of organic carbon in mangrove sediments: variability and possible ecological implications. *Hydrobiologia*, 495(1-3):33-39, 2003.

Bouillon, S., Dehairs, F., Velimirov, B., Abril, G. and Borges, A.V.: Dynamics of organic and inorganic carbon across contiguous mangrove and seagrass systems (Gazi Bay, Kenya). *J. Geophys. Research-Biogeosciences* 112(G2):G02018, 2007.

Burnham, K.P. and Anderson, D.R.: *Model Selection and Multi-Model Inference: A Practical Information-Theoretic Approach* (2nd Edition). Secaucus, NJ, USA: Springer, 2002.

Conley, D.J., Björck, S., Carstensen, J., Bonsdorff, E., Destouni, G. and Gustafsson, B.G.: Hypoxia- related processes in the Baltic Sea. *Env. Sci. Tech.*, 2009.

DMU-report, Ærtebjerg, G., Andersen, J.H. and Hansen, O.S.(Eds.): *Nutrients and Eutrophication in Danish Marine Waters . A Challenge for Science and Management*, National Environmental Research Institute, Denmark, 2003.

Downie, A., von Numers, M. and Boström, C.: Influence of model selection on the predicted distribution of the seagrass *Zostera marina*. *Estuar. Coast. and Shelf S.*, 121:8-19, 2013.

Duarte, C.M. and Cebrian, J.: The fate of marine autotrophic production. *Limnol. Oceanogr.*, 41(8):1758-1766, 1996.

Duarte, C.M., Middelburg, J.J. and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences*, 2(1):1-8, 2005.

Duarte, C.M., Kennedy, H., Marba, N. and Hendriks, I.: Assessing the capacity of seagrass meadows for carbon burial: Current limitations and future strategies. *Ocean Coast Man.*, 83:32-38 2013.

Duarte, C.M., Losada, I.J., Hendriks, I.E., Mazarrasa, I. and Marba, N.: The role of coastal plant communities for climate change mitigation and adaptation. *Nature Clim. Change*, 3 (11):961-968, 2013.

Folk, R.L. and Ward, W.C.: Brazos river bar: a study of significance of grain size parameters. *J. Sediment Petrol* 1957;27:3, 1957

Fonseca and Fisher, J.S.: A comparison of canopy friction and sediment movement between four species of seagrass with reference to their ecology and restoration. *Marine ecology progress series*. Oldendorf, 29(1):15-22, 1986

Fonseca, M.S. and Cahalan, J.A.: A preliminary evaluation of wave attenuation by four species of seagrass. *Estuar. Coast. Shelf Sci.* 1992 12; 35(6):565-576, 1992.

Fourqurean, J.W. and Schrlau, J.E. Changes in nutrient content and stable isotope ratios of C and N during decomposition of seagrasses and mangrove leaves along a nutrient availability gradient in Florida Bay, USA. *Chem. Ecol.* 19:373-390, 2003.

Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marba, N., Holmer, M. and Mateo, A.M.: Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience*, 5(7):505-509, 2012.

Fry B. and Sherr E.B. $\delta^{13}\text{C}$ measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contrib. Mar. Sci.*, 27:13, 1984.

Fry, B., Scalan, R.S., Parker, P.L.: *Stable Carbon Isotope Evidence for 2 Sources of Organic-Matter in*

Coastal Sediments - Seagrasses and Plankton. *Geochim. Cosmochim. Acta* 41(12):1875-1877, 1977.

Gacia, E., Granata, T.C. and Duarte, C.M.: An approach to measurement of particle flux and sediment retention within seagrass (*Posidonia oceanica*) meadows. *Aquat. Bot.*, 65(1-4):255-268, 1999.

Gacia, E. and Duarte, C.M. and Middelburg, J.J.: Carbon and nutrient deposition in a Mediterranean seagrass (*Posidonia oceanica*) meadow. *Limnol. Oceanogr.* 47(1):23-32, 2002.

Gattuso, J.P., Frankignoulle, M. and Wollast R. Carbon and carbonate metabolism in coastal aquatic ecosystems. *Ann. Rev. Ecol. Syst.*, 29:405-434, 1998.

Greiner, J.T., McGlathery, K.J., Gunnell, J. and McKee, B.A. Seagrass Restoration Enhances "Blue Carbon" Sequestration in Coastal Waters. *Plos One*, 8(8):e72469, 2013.

Gustafsson, C. and Boström, C.: Algal mats reduce eelgrass (*Zostera marina* L.) growth in mixed and monospecific meadows. *Journal of Experimental Marine Biology and Ecology*.461:85-92, 2014.

Hendriks, I.E., Sintes, T., Bouma, T.J. and Duarte, C.M.: Experimental assessment and modeling evaluation of the effects of the seagrass *Posidonia oceanica* on flow and particle trapping. *Mar. Ecol. Prog. Ser.* 356:163-173, 2008.

Herr, D., Pidgeon, E., Laffoley, D. (Eds.): Blue Carbon. Policy Framework: Based on the Discussion of the International Blue Carbon Policy Working Group, IUCN and Arlington, Gland, Switzerland, USA: CI. viþ39pp, 2012.

Holmer, M., Duarte, C.M., Boschker, H.T.S. and Barron, C.: Carbon cycling and bacterial carbon sources in pristine and impacted Mediterranean seagrass sediments. *Aquat. Microb. Ecol.*, 36(3):227-237, 2004.

Holmer, M., Baden, S., Bostrom C and Moksnes P.: Regional variation in eelgrass (*Zostera marina*) morphology, production and stable sulfur isotopic composition along the Baltic Sea and Skagerrak coasts. *Aquat. Bot.*, 91(4):303-310, 2009.

Holmer, M. 2009. Productivity and biogeochemical cycling in seagrass ecosystems. In *Coastal Wetlands: An Integrated Ecosystem Approach*, ed. G.M.E. Perillo, E. Wolanski, D.R. Cahoon and M.M. Brinson, 377-401. Amsterdam, The Netherlands.

Holmer, M., Wirachwong, P. and Thomsen, M.S.: Negative effects of stress-resistant drift algae and high temperature on a small ephemeral seagrass species. *Mar. Biol.*, 158(2):297-309, 2011.

Kennedy, H., Gacia, E., Kennedy, D.P., Papadimitriou, S. and Duarte, C.M.: Organic carbon sources to SE Asian coastal sediments. *Estuar. Coast. and Shelf S.*, 60(1):59-68, 2004.

Kennedy, H. and Björk, M.: Seagrasses, in *The Management of Natural Coastal Carbon Sinks in Coastal Ecosystems: Investigating and Realising the Potential*. 2009.

Kennedy, H., Beggins, J., Duarte, C.M., Fourqurean, J.W., Holmer, M. and Marba, N.: Seagrass sediments as a global carbon sink: Isotopic constraints. *Glob. Biogeochem. Cycles*, 24:GB4026, 2010.

Kristensen, E. and Holmer, M.: Decomposition of plant materials in marine sediment exposed to different electron acceptors (O₂, NO₃⁻, and SO₄²⁻), with emphasis on substrate origin, degradation kinetics, and the role of bioturbation. *Geochim. Cosmochim. Acta*, 65(3):419-433, 2001.

Lavery, P.S., Mateo, M., Serrano, O. and Rozaimi, M.: Variability in the Carbon Storage of Seagrass Habitats and Its Implications for Global Estimates of Blue Carbon Ecosystem Service. *Plos One* ,8 (9):e73748, 2013.

Legendre, P. and Anderson, M.J.: Distance-based redundancy analysis: Testing multispecies responses in multifactorial ecological experiments (vol 69, pg 1). *Ecol. Monogr.* , 69 (4):512-512, 1999.

Leipe, T., Tauber, F., Vallius, H., Virtasalo, J., Uścinowicz, S., Kowalski, N., Hille, S., Lindgren, S., and Myllyvirta, T.: Particulate organic carbon (POC) in surface sediments of the Baltic Sea. *Geo-Mar. Lett.* 31:175-188, 2011.

Macreadie, P.I., Baird, M.E., Trevathan-Tackett, S.M., Larkum, A.W.D. and Ralph, P.J.: Quantifying and modelling the carbon sequestration capacity of seagrass meadows - A critical assessment. *Mar. Pollut. Bull.*, 83(2):430-439, 2014.

Marba, N., Arias-Ortiz, A., Masque, P., Kendrick, G. A., Mazarrasa, I., Bastyan, G. R., Garcia-Orellana, J. and Duarte, C. M.: Impact of seagrass loss and subsequent revegetation on carbon sequestration and stocks. *J. of Ecology* 103:296-302, 2015

McLeod, E., Chmura, G.L., Bouillon, S., Salm, R., Bjork, M. and Duarte, C.M.: A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front. Ecol. Env.*, 9(10):552-560, 2011.

Miyajima, T., Hori, M., Hamaguchi, M., Shimabukuro, H., Adachi, H. and Yamano, H.: Geographic variability in organic carbon stock and accumulation rate in sediments of East and Southeast Asian seagrass meadows. *Glob. Biogeochem. Cycles*, 29(4):397-415, 2015.

Moksnes, P.-E. and Cole, S. G.: Valuing multiple eelgrass ecosystem services in Sweden: Fish production and uptake of carbon and nitrogen. *Front. Mar. Sci.* 2(121):1-18, 2015.

Moncreiff, C.A. and Sullivan M.J.: Trophic importance of epiphytic algae in subtropical seagrass beds: evidence from multiple stable isotope analyses. *Mar. Ecol. Prog. Ser.*, 215:93-106, 2001.

Nellemann, C., Corcoran, E., Duarte, C.M., Valdés, L., De Young, C. and Fonseca, L.: Blue Carbon. A Rapid Response Assessment, 2009.

Norkko, A. and Bonsdorff, E.: Rapid zoobenthic community responses to accumulations of drifting algae. *Mar. Ecol. Prog. Ser.*, 131:143-157, 1996.

Olesen, B. and Sandjensen, K.: Patch Dynamics of Eelgrass *Zostera marina*. *Mar. Ecol. Prog. Ser.* 106(1-2):147-156, 1994.

Orth, R.J., Carruthers, T.J.B., Dennison, W.C., Duarte, C.M., Fourqurean, J.W. and Heck, K.L.Jr. : A global crisis for seagrass ecosystems. *Bioscience*, 56(12):987-996, 2006.

Pedersen, M. Ø., Serrano, O., Mateo, M.A, Holmer, M.: Temperature effects on decomposition of a *Posidonia oceanica* mat. *Aquat. Microb. Ecol.* 65: 169-182, 2011.

Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A. and Sifleet, S: Estimating Global "Blue Carbon" Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *Plos One*, 7(9):e43542, 2012.

Petersen, J.K., Maar, M., Ysebaert, T. and Herman, P.M.J.: Near-bed gradients in particles and nutrients above a mussel bed in the Limfjorden: influence of physical mixing and mussel filtration. *Mar. Ecol. Prog. Ser.*, 490:137-146, 2013.

Phillips, D.L. and Gregg, J.W.: Source partitioning using stable isotopes: coping with too many sources. *Oecologia*, 136(2):261-269, 2003.

Rask, N., Pedersen, S.E. and Jensen, M.H.: Response to lowered nutrient discharges in the coastal waters around the island of Funen, Denmark. *Hydrobiologia*, 393:69, 1999.

Rasmussen, E.: The wasting disease of eelgrass *Zostera marina* and its effects on environmental factors and fauna. *Seagrass ecosystems. A scientific perspective*. New York: Marcel Dekker Inc., 1977.

Rasmussen, J.R., Pedersen, M.F., Olesen, B., Nielsen, S.L. and Pedersen, T. M.: Temporal and spatial dynamics of ephemeral drift-algae in eelgrass *Zostera marina*, beds. *Estuar. Coast. and Shelf S.* 119:167-175, 2013.

Salovius, S. and Bonsdorff, E.: Effects of depth, sediment and grazers on the degradation of drifting filamentous algae (*Cladophora glomerata* and *Pilayella littoralis*). *Journal of Experimental Marine Biology and Ecology*. 298:93-109, 2004.

Schlacher, T.A and Conolly, R.M.: Effects of acid treatment on carbon and nitrogen stable isotope ratios in ecological samples: a review and synthesis. *Methods in Ecology and Evolution* 5(6):541-550, 2014.

Serrano, O., Lavery, P.S., Rozaimi, M. and Mateo, A. M.: Influence of water depth on the carbon sequestration capacity of seagrasses. *Global Biogeochem. Cycles* 28(9):950-961, 2014.

Short, F. and Duarte, C.M.: Methods for the measurement of seagrass growth and production. In: Short and Cole, editors. *Global seagrass research methods* Amsterdam: Elsevier; p. 155-182, 2001.

Spalding, M., Taylor, M., Ravilious, C., Short, F., Green, E.: Global overview: The distribution and status of seagrasses. In: Green, E.P. and Short F.T.: editors. *World Atlas of Seagrasses: Present Status and Future Conservation*. Berkeley California: University of California Press, p. 5-26, 2003.

United States Government. Technical Support Document: Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866 (United States Government Interagency Working Group on Social Cost of Carbon, Washington, DC), 2010.

Vichkovitten, T. and Holmer, M. Contribution of plant carbohydrates to sedimentary carbon mineralization. *Org. Geochem.*, 35(9):1053-1066, 2004.

Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C. and Olyarnik, S.: Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. Natl. Acad. Sci. USA*, 106(30):12377-12381, 2009.

Wiles, P.J., van Duren, L.A., Hase, C., Larsen, J. and Simpson, J.H.: Stratification and mixing in the Limfjorden in relation to mussel culture. *J. Mar. Syst.*, 60(1-2):129-143, 2006.




Fig. 1. The study sites in Denmark and Finland. a) study regions, b) Finnish study sites, c) Limfjorden study sites, d) Funen study sites.




Fig. 2. Silt content (%) (a, c) and degree of sediment sorting (b, d) at the study sites in Finland and Denmark, respectively. Lower values in degree of sorting indicate well-sorted sediment types.





Fig. 3. Sediment profiles of organic carbon density (mg C cm^{-3}) in the top 25 cm of the Finnish and Danish eelgrass (*Zostera marina*) meadows. Note the difference in the scale of x-axis between the regions. Numbers below detection limit ($\%OC < 0.01 \%DW$) are not included in the figure. Average (\pm SEM; $n=1-3$)

727
728
729
730
731
732
733
734
735
736
737



738
739
740
741
742
743

Fig. 4. Corg stocks (g C m⁻²) in the top 25 cm of sediment in Finnish and Danish eelgrass (*Zostera marina*) meadows. Note that the value of Thurøbund (grey bar) corresponds to right y-axis.

744
745
746
747
748
749
750




Fig. 5. Relative contribution of different organic matter sources (*Z. marina*, *P. perfoliatus*, *P. pectinatus*, *Ruppia cirrhosa*, phytoplankton and drift algae) to the ^{13}C signal of the sediment surface layer (0-2 cm) in Finnish and Danish eelgrass (*Z. marina*) meadows.




Fig. 6. Distance-based redundancy analysis (DbRDA) plot showing the environmental parameters (percentage of *Z. marina* in sediment carbon pool, above:belowground-ratio, annual eelgrass production, sediment silt content [%], sediment dry density and sediment porosity) fitted to the variation in the Corg stock (g C m^{-2}) at the Finnish and Danish eelgrass (*Z. marina*) sites, respectively. Vectors indicate direction of the parameters' effect. Site codes: Finland; Ko=Kolaviken, Ry=Ryssholmen, Tv=Tvärminne, Få=Fårö, Ly=Lyddaren, Lå=Långören, Hu=Hummelskär, Jä=Jänisholm, Sa=Sackholm, Ån=Ängsö. Site codes: Denmark; Ag12=Agerø12, Ag3=Agerø3, Vi=Visby, Lg=Løgstør, Lo=Lovns, Th=Thurøbund, Lu=Lunkebugt, Da=Dalby, Ke=Kertinge, Ny=Nyborg.

Table 1. Location, silt content (% silt), sediment dry density (Dry dens. g cm⁻³), sediment organic carbon content (SedOC, % DW), sediment organic matter content (SedOM, %DW), $\delta^{13}\text{C}$ sediment surface, $\delta^{13}\text{C}$ *Z. marina* leaves, $\delta^{13}\text{C}$ *Z. marina* rhizomes, seagrass shoot density (Shoot dens., shoots m⁻²), seagrass above and belowground biomass (AB and BB, gDW m⁻²), root:shoot-ratio (R:S), and aboveground production (Production, gDW m⁻²y⁻¹) at the sampling sites. SE (n= 3–4) is given. Annual seagrass production is calculated from pooled values of replicates per site and therefore no SE is shown.

Country/ Location	Silt content (%)	Dry dens. (g cm ⁻³)	SedOM (% DW)	SedOC (% DW)	$\delta^{13}\text{C}$ sediment surface	$\delta^{13}\text{C}$ <i>Z. marina</i> leaves	$\delta^{13}\text{C}$ <i>Z. marina</i> rhizomes	Shoot density (shoots m ⁻²)	AB (gDW m ⁻²)	BB (gDW m ⁻²)	R:S	Production (gDW m ⁻² y ⁻¹)
Finland												
Fårö	5.0±1.4	1.32±0.025	0.66±0.07	0.13±0.001	-20.6±0.3	-9.7±0.4	-9.0±0.20	304±32	138±20	167±28	1.27±0.13	773
Hummelskär	9.0±2.6	1.33±0.009	1.06±0.20	0.35±0.019	-19.4±1.2	-9.3±0.3	-9.8±0.25	364±31	70±11	28±2	0.45±0.06	446
Jänisholm	7.1±2.1	1.37±0.076	0.93±0.20	0.33±0.135	-22.1±0.4	-10.8±0.4	-11.0±0.28	128±17	65±16	46±2	1.44±0.53	270
Kolaviken	1.9±0.2	1.34±0.035	0.75±0.02	0.13±0.011	-19.5±0.2	-10.3±0.3	-11.4±0.34	476±96	74±6	149±16	2.07±0.27	324
Lyddaren	4.9±2.5	1.34±0.171	1.75±0.70	0.45±0.094	-13.5±3.5	-8.8±0.4	-9.6±0.29	228±42	86±7	57±12	0.64±0.09	505
Långören	4.4±2.1	1.42±0.046	2.70±2.10	0.19±0.019	-18.9±0.4	-8.5±0.1	-8.9±0.15	436±53	121±46	68±25	0.58±0.06	788
Ryssholmen	2.7±0.6	1.34±0.054	0.89±0.20	0.16±0.004	-20.7±0.3	-11.5±0.1	-11.5±0.29	756±57	160±3	136±16	0.86±0.11	803
Sackholm	12.4±1.9	1.36±0.042	0.95±0.20	0.26±0.027	-21.1±0.8	-10.3±0.7	-9.9±0.34	774±234	110±18	37±9	0.31±0.04	377
Tvärminne	9.2±1.9	1.33±0.034	0.88±0.20	0.20±0.016	-22.7±0.6	-11.6±0.1	-11.5±0.25	112±11	99±16	38±7	0.37±0.01	436
Ängsö	6.3±0.5	1.36±0.052	0.84±0.02	0.20±0.010	-20.1±0.3	-10.3±0.1	-10.3±0.28	604±98	91±6	63±9	0.67±0.05	521
FIN average	6.3±1	1.35±0.014	1.4±0.3	0.24±0.033	-19.9±0.3	-10.1±0.3	10.3±0.32	417±75	101±3	79±5	0.87±0.06	524
Denmark												
Agero 3	29.4±6.2	1.24±0.085	1.94±0.60	2.30±0.082	-13.0±1.7	-9.2±0.5	-11.1±0.22	448±89	181±33	84±8	0.52±0.07	1075
Agero 12	27.3±7.7	1.35±0.173	1.65±0.80	0.29±0.135	-17.4±0.8	-10.7±0.3	-11.9±0.21	404±90	110±2	46±9	0.40±0.08	576
Dalby	8.1±1.2	1.37±0.034	0.67±0.03	0.12±0.009	-17.3±0.7	-9.7±0.3	-10.5±0.56	400±48	76±7	83±10	1.09±0.11	470
Kertinge	27.1±1.5	1.15±0.045	12.59±1.60	3.23±0.236	-16.6±0.2	-9.2±0.1	-9.8±0.08	328±64	90±17	64±14	0.68±0.02	527
Lovns	17.3±2.7	1.22±0.092	2.90±0.50	1.53±0.088	-16.3±2.4	-11.5±0.4	-12.2±0.37	360±27	141±4	100±11	0.70±0.06	848
Lunkebugt	33.0±7.4	1.23±0.227	4.72±2.40	1.71±0.806	-16.9±0.3	-8.9±0.9	-10.6±0.38	347±81	210±10	382±24	1.82±0.08	1056
Løgstør	4.0±0.4	1.23±0.025	0.75±0.03	0.31±0.089	-17.7±0.4	-9.7±0.4	-10.4±0.51	300±14	149±11	63±13	0.42±0.07	755
Nyborg	0.5±0.3	1.17±0.027	0.42±0.02	0.10±0.006	-17.6±1.1	-9.3±0.2	-10.6±0.34	652±30	203±24	214±50	1.00±0.14	1179
Thurøbund	34.6±2.8	1.27±0.030	14.48±0.80	5.78±0.512	-15.5±0.4	-8.2±0.1	-9.0±0.22	420±98	101±16	398±15	4.54±0.70	619
Visby	21.0±3.6	1.25±0.021	1.17±0.06	2.18±0.201	-13.8±1.2	12.0±0.6	-12.4±0.70	520±21	193±13	49±4	0.25±0.01	2172
DK average	20.2±3.9	1.25±0.022	3.9±1.5	1.75±0.563	-16.20±0.2	-9.8±0.4	-10.9±0.33	418±32	145±5	148±14	1.14±0.13	928

811 Table 2. Estimated average carbon stocks (g C m⁻² and Mt C), annual areal carbon accumulation (Corg acc. t C ha⁻¹y⁻¹) and annual carbon accumulation
812 (Annual Corg (Mt C y⁻¹) in Finnish and Danish eelgrass (*Z. marina*) meadows. Denmark_{lost} = eelgrass area of the region lost since the beginning 1900's.
813 Limfjorden_{lost} = eelgrass area of the region lost since the 1900's. See text for calculations. *) mean carbon density (mg C cm⁻³) calculated for Denmark
814 is used. n.d= no data. For calculations of annual carbon accumulation three different sediment accumulation rates were applied (0.32 mm y⁻¹;
815 Miyayima et al. 2015, 2.02 mm y⁻¹; Duarte et al. 2013b and 4.2 mm y⁻¹; Serrano et al. 2014), for Corg seq the sediment accumulation rate of 2.02 mm
816 y⁻¹ was used.
817

Region	Seagrass area (km ²)	Carbon density (mg C cm ⁻³)	Corg stock (g C m ⁻²)	Corg stock (Mt C)	Corg acc. (t C ha ⁻¹ y ⁻¹)	Annual Corg accumulation (Mt C y ⁻¹)		
						819		
						0.32 mm y ⁻¹	2.02 mm y ⁻¹	4.2 mm y ⁻¹
Finland	30	2.60±0.09	627±25	0.019±< 0.001	0.052	0.002	0.016	0.0328
Limjorden	18	10.57±1.66	2644±207	0.047± 0.007	0.213	0.006	0.038	0.079
Funen	179	24.32±9.15	6005±1127	1.090±0.410	0.491	0.139	0.881	1.832
Denmark _{min}	673	17.45±9.42*	4324±1188*	2.164±0.005	0.352	0.376	2.373	3.636
Denmark _{max}	1345	17.45±9.42*	4324±1188*	5.868±0.014	0.352	0.75	4.741	9.859
Denmark _{lost}	5381-6230	17.45±9.42*	17.45±9.42*	23.478-27.183	n.d	n.d	n.d	n.d

820
821
822
823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

Table 3. Table from DistLm analysis showing variables in the marginal tests and the results for statistical analysis.

MARGINAL TESTS				
Variable	SS	Pseudo-F	P-value	Proportion
1. Root:shoot- ratio	5309	10.64	0.002	0.155
2. Sediment dry density	10704	26.37	0.001	0.313
3. Annual eelgrass production	4959	9.82	0.002	0.145
4. Shoot density	48	0.08	0.911	0.001
5. Porosity	3507	6.61	0.01	0.102
6. Silt content (%)	12653	33.99	0.001	0.369
7. C:N-ratio of plant material	464	0.79	0.397	0.014
8. <i>Z. marina</i> content (%)	12179	32.02	0.001	0.356
9. Degree of sorting	9725	23.01	0.001	0.284

843 Table 4. Table from DistLm analysis showing results from the sequential tests and solution given
844 by the analysis.

845

Variable							
	AICc	Sum of squares	Pseudo-F	P- value	Proportion	Cumulative proportion	Degrees of freedom
6. Silt content (%)	357.4	12653	33.9	0.001	0.369	0.369	58
1. Root:shoot-ratio	346.0	4375	14.5	0.001	0.127	0.497	57
8. <i>Z. marina</i> content (%)	333.6	3745	15.6	0.001	0.109	0.606	56
3. Production	332.2	805	3.5	0.037	0.023	0.630	55
2. Sediment dry density	331.3	700	3.2	0.049	0.020	0.650	54
5. Porosity	330.8	602	2.8	0.056	0.017	0.668	53
BEST SOLUTION	AICc	R ²	RSS	Variables	Selections		
	330.8	0.668	11363	6	1-3;5;6;8		

846

847

848

849

850

851

852

853

854

855

856

857