

1 Blue carbon stocks in Baltic Sea eelgrass (*Zostera marina*) meadows

2

3 Maria Emilia Röhr^{1,2*}, Christoffer Boström¹, Paula Canal- Vergés³, Marianne Holmer²

4

5 ¹Åbo Akademi University, Faculty of Science and Engineering, Environmental and Marine
6 Biology, Artillerigatan 6, 20520 Åbo, Finland

7 ²University of Southern Denmark, Department of Biology, Campusvej 55, 5230 Odense M,
8 Denmark

9 ³Danish Shellfish Centre, DTU Aqua, Technical University of Denmark, Øroddevej 80, 7900
10 Nykøbing Mors, Denmark

11

12 * Corresponding author;

13

14 Emilia Röhr (mrohr@abo.fi)

15 Abstract

16

17 Although seagrasses cover only a minor fraction of the ocean seafloor, their carbon sink capacity
18 accounts for nearly one-fifth of the total oceanic carbon burial and thus play a critical structural
19 and functional role in many coastal ecosystems. We sampled 10 eelgrass (*Zostera marina*)
20 meadows in Finland and 10 in Denmark to explore seagrass carbon stocks (Corg stock) and
21 carbon accumulation rates (Corg accumulation) in the Baltic Sea area. The study sites represent
22 a gradient from sheltered to exposed locations in both regions to reflect expected minimum and
23 maximum stocks and accumulation. The Corg stock integrated over the top 25 cm of the
24 sediment averaged 627g C m⁻² in Finland, while in Denmark the average Corg stock was over six
25 times higher (4324 g C m⁻²). A conservative estimate of the total organic carbon pool in the
26 regions ranged between 6.98-44.9 t C ha⁻¹. Our results suggest that the Finnish eelgrass
27 meadows are minor carbon sinks compared to the Danish meadows, and that majority of the
28 Corg produced in the Finnish meadows is exported. Our analysis further showed that > 40 % of

29 the variation in the Corg stocks was explained by sediment characteristics i.e. dry density,
30 porosity and silt content. In addition, our analysis show, that the root: shoot- ratio of *Z. marina*
31 explained > 12 % and contribution of *Z. marina* detritus to the sediment surface Corg pool
32 explain > 10 % of the variation in the Corg stocks. The mean monetary value for the present
33 carbon storage and carbon sink capacity of eelgrass meadows in Finland and Denmark, were
34 281 and 1809 € ha⁻¹, respectively. For a more comprehensive picture of seagrass carbon storage
35 capacity, we conclude that future Blue Carbon studies should in a more integrative way,
36 investigate the interactions between sediment biogeochemistry, seascape structure, plant
37 species architecture and the hydrodynamic regime.

38 Keywords: Blue Carbon, eelgrass, seagrass, carbon stock, carbon accumulation, sequestration,
39 carbon sink

40

41 Copyright statement

42

43 The article is distributed under the [Creative Commons Attribution 3.0](#) License. Unless otherwise
44 stated, associated published material is distributed under the same licence. Authors
45 grant Copernicus Publications a license to publish the article and identify itself as the original
46 publisher. Authors grant Copernicus Publications commercial rights to produce hardcopy
47 volumes of the journal for purchase by libraries and individuals. Authors grant any third party
48 the right to use the article freely under the stipulation that the original authors are given credit
49 and the appropriate citation details are mentioned.

50

51 Introduction

52

53 The atmospheric carbon dioxide (CO₂) enters the ocean via gas-exchange processes at the ocean-
54 atmosphere interface. In the ocean dissolved inorganic carbon is fixed in photosynthesis by

55 primary producers, and released again through respiration. A large percentage of this fixed
56 carbon is stored and sequestered in the sediments of vegetated coastal ecosystems of which the
57 three globally most significant are saltmarshes, mangrove forests and seagrass meadows (Herr
58 et al. 2012). The carbon stored by these ecosystems is known as Blue Carbon (Duarte et al. 2005
59 Duarte et al. 2013a, Nellemann et al. 2009). Blue Carbon ecosystems function as carbon sinks, in
60 which the rate of carbon sequestered by the ecosystem exceeds the rate of carbon lost through
61 respiration and export.

62 Seagrass meadows play a critical structural and functional role in many coastal ecosystems
63 (Orth et al. 2006). Although seagrass meadows only cover globally about 300 000 - 600 000 km²
64 of the ocean sea floor, corresponding to 0.1 to 0.2 % of the total area, their carbon sink capacity
65 (capacity of seagrasses to absorb and store carbon in living and dead biomass and in the
66 sediments) may account for up to 18 % of the total oceanic carbon burial (Gattuso et al. 1998,
67 Duarte et al. 2005, Kennedy et al. 2010, Fourqurean et al. 2012). A large portion of the carbon
68 sequestered (captured and stored) by seagrasses is stored in sediments with a conservative
69 value of 10 Pg C in the top 1 meter of seagrass sediments (Fourqurean et al. 2012).
70 Consequently, recent global estimates imply that seagrass sediments store almost 25 200 to 84
71 000 t C km² (Fourqurean et al. 2012). More importantly, carbon in submerged sediments is
72 stored for timescales of millennia while terrestrial soils are usually less stable and only
73 sequester carbon up to decades (Hendriks et al. 2008).

74

75 The coasts of Scandinavia and the Baltic Sea are key distribution areas for eelgrass (*Zostera*
76 *marina* L.) meadows (Boström et al. 2002, Boström et al. 2014). The meadows extend from fully
77 saline (>30) along the Norwegian coast to brackish (5-6) archipelago areas of Finland. This
78 region is estimated to support >6 000 individual meadows covering at least 1 500 – 2 000 km²,
79 which is four times more than the combined eelgrass area of Western Europe (Spalding et al.
80 2003, Boström et al. 2014). Consequently, this region plays a key role in the coastal carbon
81 dynamics, but we presently lack estimates of the role of eelgrass for carbon storage in temperate

82 sediments. Seagrasses are lost at accelerating rates and it has been estimated that 29 % of global
83 seagrass area has disappeared since the initial recording of seagrasses in 1879 (Waycott et al.
84 2009). This decline could have severe consequences on the total capacity of marine ecosystems
85 to store and sequester carbon in addition to the other ecosystem services seagrass meadows
86 provide. Little is known about the magnitude of carbon emissions from degraded seagrasses
87 ecosystems, not to mention its economic implications. A recent study points out, that despite the
88 importance of these ecosystems in the global carbon budget, none of the three Blue Carbon
89 ecosystems have been included in global carbon market protocols (Pendleton et al. 2012).

90

91 Seagrasses exhibit marked differences in shoot architecture and grow under variable
92 environmental settings, making direct extrapolations between species and locations difficult.
93 Consequently, there is a pressing need to better understand which factors are causing variability
94 in carbon storage (Corg stocks, carbon stored in living and dead seagrass biomass and
95 sediments) and the capacity of seagrass meadows to sequester carbon (Corg accumulation) in
96 seagrass sediments. Indeed, recent studies show considerable influence of seagrass habitat
97 setting, sediment characteristics and species-specific traits on the variability in carbon storage
98 capacity in seagrass meadows (Duarte et al. 2013a, Lavery et al. 2013, Miyajima et al. 2015).
99 Such differences contribute to uncertainty in local and global estimates of the carbon storage
100 capacity and carbon dynamics in coastal seagrass areas.

101

102 In order to determine seagrass Corg stocks and Corg accumulation, knowledge on the sources of
103 the carbon stored in the sediments is also crucial. The different Corg sources vary in their
104 turnover compared to seagrasses (sources other than seagrasses being typically faster) and
105 volumes of standing stock (typically less) and thus affect the dynamics of the Corg stocks and
106 accumulation (Fry et al. 1977, Kennedy et al. 2004, Kennedy et al. 2010). Seagrasses are known
107 to be enriched in $\delta^{13}\text{C}$ compared to other potentially sources of Corg in the seagrass sediments,
108 such as plankton, macroalgae, allochthonous carbon material, seagrass epiphytes, and benthic

109 microalgae (Kennedy et al. 2004, Kennedy et al. 2010, Fry and Sherr 1984, Moncreiff and
110 Sullivan 2001, Bouillon et al. 2002, Bouillon and Boschker 2006, Macreadie et al. 2014). Thus,
111 the stable isotope signals of seagrasses and other potential Corg sources can be relatively easily
112 and reliably used as a proxy for identification of the origin of Corg in seagrass sediment carbon
113 pool (Kennedy et al. 2010). Unfortunately, the current knowledge base on how these factors
114 interact and influence carbon fluxes and storage is, at best, limited at both local and global
115 scales.

116

117 In this study, we contrast storage, accumulation rates and sources of the accumulated carbon in
118 eelgrass (*Zostera marina*) meadows in two regions differing in salinity, temperature and
119 seagrass productivity, namely Finland and Denmark. Specifically we asked;

120 (1) How large is the carbon storage capacity (Corg stocks) of Baltic Sea eelgrass meadows?

121 (2) Which are the environmental factors determining the variability of carbon storage (Corg
122 stocks) and accumulation (Corg accumulation) at local and regional scales?

123 (3) How do the sediment characteristics influence the carbon storage (Corg stocks) of
124 eelgrass meadows at local and regional scales?

125 (4) How much carbon (Corg stocks) is presently stored in Finnish and Danish eelgrass
126 meadows, respectively.

127 (5) What is the present and historically lost (only in Denmark) monetary carbon value?

128

129 Materials and methods

130 Study area

131

132 Plant and sediment samples were collected in June-September 2014 from 10 sites in Finland
133 (The Archipelago Sea) and 10 sites in Denmark (Funen and Limfjorden) (Fig. 1). The Baltic Sea
134 sediments are typically mineral sediments consisting of glaciofluvial deposits and only a small

135 fraction of the sediment carbon content consists of carbonates (Leipe et al. 2011). The inorganic
136 carbon content in our samples was low and contributed with 0.5-5 % of total carbon content (n=
137 10 sites per region) and therefore carbonates were not removed from the sediment samples
138 prior to the analysis to avoid analytical errors in low organic samples (Schlacher and Connolly
139 2014). However, when interpreting the data it should be kept in mind, that given the %IC
140 variation in the samples for the different sites (range 0.1-5.78 %; average 3.33%), the
141 inorganic carbon could cause bias in the stable isotope signals of the sediment surface
142 samples (range 0.16-1.17‰; average 0.76‰).

143 The study sites in each region spanned a gradient from sheltered to exposed areas. The
144 Archipelago Sea of southwestern Finland is a shallow (mean depth 23 m), brackish (salinity 5-6)
145 coastal area characterized by a complex mosaic of some 30.000 islands and skerries (Boström et
146 al. 2006, Downie et al. 2013). The region is heavily influenced by human pressures, especially
147 eutrophication, and exhibits naturally steep environmental gradients, as well as, strong
148 seasonality in temperature and productivity (Boström et al. 2014).

149
150 Limfjorden is a brackish water area in the Jutland peninsula connected to both North Sea and
151 Kattegat with salinity ranging from 17 to 35. The Fjord has a surface area of ~1500 km² and a
152 mean depth of 4.7 m (Olesen and Sandjensen 1994, Wiles et al. 2006, Petersen et al. 2013).
153 Funen is located between the Belt Seas in the transition zone where waters from Baltic Sea and
154 Kattegat meet. The salinity of the area ranges between 10 and 25 and the annual mean water
155 temperature ranges from 10-15° C (Rask et al. 1999). This study was conducted in shallow (< 10
156 m) fjords around Funen. Also the Danish areas are heavily influenced by human pressures,
157 especially eutrophication from intense agricultural farming. (DMU; Danmarks
158 Miljøundersøgelser, 2003).

159

160 Field sampling

161

162 All samples were collected from depths of 2.5-3 m by scuba diving. At all sites, three replicate
163 sediment cores (corer: length: 50 cm, diameter: 50 mm) were taken randomly at a minimum
164 distance of 15 m from each other. The corer was manually forced to a depth of 30-40 cm and the
165 sediment between 0-25 cm was used for the analysis. The cores were capped in both ends under
166 water, and kept in a vertical position during transport to the laboratory. Eelgrass production and
167 biomass were measured at all sites from four randomly chosen locations within the eelgrass
168 meadow. In the vicinity of each sediment core, shoot density was counted using a 0.25 m² frame,
169 and above- and belowground biomass samples were collected with a corer (diameter 19.7 cm)
170 and bagged underwater. Additionally, when present, samples of plants and algae (drift algae,
171 other angiosperms, phytoplankton and epiphytes) considered likely carbon sources in the
172 eelgrass meadows were collected from each site for identification and analysis of stable isotope
173 composition. Approximately 10 g wet material was collected for each species. Annual eelgrass
174 production was determined from estimates of previous growth by applying the horizontal
175 rhizome elongation technique (Short and Duarte 2001). From each site, five replicate rhizome
176 samples with the longest possible intact rhizome carefully removed, were collected and
177 transported to the laboratory for further analysis.

178

179 Seagrass variables

180

181 In the laboratory, the above- and belowground biomass was separated and eelgrass leaves and
182 rhizomes were cleaned from epiphytes, detritus and fauna with freshwater and gently scrubbed
183 with a scalpel. All plant material was dried to constant weight (48 h in 60° C). The belowground
184 biomass was separated into living and dead rhizomes and dried separately. Only the living
185 rhizomes were used for the belowground biomass measurements while samples of both living
186 and dead rhizomes were used for analysis of organic carbon (OC) and stable isotopic
187 composition of the organic carbon ($\delta^{13}\text{C}$). The root: shoot-ratio was calculated as the ratio
188 between below- and aboveground biomasses of *Z. marina* samples. A pooled sample of 2

189 youngest leaves from 10 randomly selected shoots were collected prior to drying from the
190 aboveground biomass samples and dried separately for analysis of OC and $\delta^{13}\text{C}$. All samples
191 were analyzed by Thermo Scientific, delta V advantage, isotope ratio mass spectrometer. The
192 measured isotope ratios were represented using the δ - notation with Vienna Peedee belemnite
193 as reference material.

194

195 Determination of annual eelgrass production was done by measuring length of each individual
196 internode of the rhizomes to the nearest millimeters. To obtain an estimate of the mean annual
197 production per site, internode length measurements of individual replicates (n= 5) were pooled.
198 Due to lack of two annual production peaks in both regions the annual production was estimated
199 based on the distance between shortest and longest measured internodes, assuming that they
200 represent the time point when the water temperature was at its minimum and maximum
201 average, respectively. The time points for the water temperatures were obtained from databases
202 of the Finnish and Danish Meteorological Institutes, respectively.

203

204 Sediment variables

205

206 In the laboratory, sediment samples were sliced into sections of 2-5 cm, where the upper 10 cm
207 layer was divided into 2 cm layers and the remaining part in 5 cm layers. From each subsample
208 visible plant parts and fauna were removed before the sediment was homogenized. From the 0-2
209 cm section a subsample of 20 ml was taken for grain size analysis by a Malvern Mastersizer 3000
210 particle size analyzer. The sediment silt content was calculated as the fraction with particle size
211 of 2-63 μm from the range of all particle sizes (Folk and Ward 1957). Sediment water content,
212 dry bulk density and porosity were determined from a subsample of 5 ml that was taken using a
213 cut-off 5 ml syringe and weighed before and after drying at 105°C for 6 h from all sediment
214 layers. The dried sediment samples were homogenized in a mortar and divided into two
215 subsamples from which one was used for analysis of sediment organic matter content (OM, as

216 loss of ignition: 4 h in 520°C), and the other for analysis of sediment $\delta^{13}\text{C}$ and OC as described
217 above for the plant materials. Inorganic carbon content was low in sediments from both regions
218 (< 5 %) and considered insignificant compared to the organic fraction (1-2 order of magnitude
219 higher).

220

221 Corg stock and Corg accumulation calculations

222

223 The depth integrated Corg stocks were calculated according to Lavery et al. (2013) by
224 multiplying the OC (OC mg gDW⁻¹) measured from different sections of the sediment core with
225 the corresponding sediment dry density (g cm⁻³). These numbers were then depth integrated
226 over 25 cm in order to estimate the depth integrated carbon density. To estimate sediment Corg
227 stock and Corg accumulation of Finnish and Danish eelgrass area we used averages from 10 sites
228 from each region in our calculations. The Corg (obtained by depth integration of the carbon
229 density (mg C cm⁻³) of the sampled region was multiplied with estimated seagrass area of the
230 region based on the most recent areal estimates (in km²) of seagrass distribution available in the
231 literature (Boström et al. 2014) and given as Corg in g C m⁻². In Finland the estimated areal
232 extent was 30 km², while in Denmark the extrapolations were based on the minimum and
233 maximum estimates of the areal extent, respectively (673 and 1345 km²; Boström et al. 2014).
234 Results for carbon accumulation (applied by multiplying the depth integrated Corg stock,
235 regional seagrass area and sediment accumulation rate estimate from literature) in each area
236 are given as Corg accumulation (t y⁻¹). Due to lack of long term monitoring of sediment
237 accumulation in eelgrass meadows, we used available minimum, average and maximum
238 sediment accumulation rates in seagrass meadows obtained from literature (Duarte et al. 2013b,
239 Serrano et al. 2014, Miyajima et al. 2015).

240

241 To calculate the total Corg pool in Danish and Finnish eelgrass sediments, we summed the
242 following three components: (1) the annual areal eelgrass carbon accumulation rate (Corg

243 accumulation in $\text{t C ha}^{-1}\text{y}^{-1}$, calculated by dividing the measured Corg stocks (C g DW m^{-2}) in each
244 region with the time that it takes to accumulate this stock with a sedimentation rate of 2.02 mm
245 y^{-1} , (2) the total C in the average living aboveground and belowground *Z. marina* tissue (t C ha
246 $^{-1}$), and (3) the mean Corg stocks (t C ha^{-1}) in eelgrass sediments in Denmark and Finland,
247 respectively. To calculate the present and lost economic value of eelgrass carbon stocks, we used
248 the social cost of carbon of 40.3 € t C^{-1} (United States Government 2010) and multiplied this
249 value with the Corg stocks (t C km^{-2}). To estimate the Danish eelgrass losses over the past 100
250 years in economic terms, we used the calculations above, but accounted for the annually lost
251 sequestration value by multiplying the rate by 100. We used the most recent loss estimates for
252 Denmark for the period 1900-2000, assuming that the present coverage constitutes 10 % or 20
253 % of the historical area, respectively (Boström et al. 2014).

254

255 Sediment carbon sources

256

257 The Isosource 1.3 isotope mixing model software (Phillips and Gregg 2003) was used to estimate
258 the contribution of different carbon sources to the sediment surface Corg stock. We ran the
259 Isosource model using the $\delta^{13}\text{C}$ obtained from stable isotope analysis of *Z. marina* leaves, living
260 and dead rhizomes and for samples of other abundant Corg sources within the meadow ($n=1-5$)
261 with increments of 1 % and tolerance of 0.1. Numbers are given as percentage contribution to
262 the sediment surface carbon pool.

263

264 Data analysis

265

266 All statistical analyses were performed using the PRIMER 6 PERMANOVA+ package (Anderson
267 et al. 2008). A 2-factor mixed model was used, where sampling sites and region (FIN, DEN) were
268 used as fixed factors for the biological response variable (sediment organic carbon stock, g C m^{-2}).
269 In addition we ran reduced, countrywise DistLm models in order to better address the

270 possible differences in regional environmental drivers for Corg stock. Prior to analysis, the
271 environmental predictor variables (degree of sorting, sediment dry density, sediment water
272 content, sediment porosity, sediment silt content, sediment organic content, annual production,
273 root: shoot-ratio, shoot density and percentage of *Z. marina* detritus contribution to Corg) were
274 visually inspected for collinearity using Draftsman plots of residuals. Due to autocorrelation
275 between sediment variables (water content, porosity and dry density) sediment water content
276 was removed from the environmental variables. To achieve normality in the retained
277 environmental variables, data was log-transformed ($\log(X+1)$) and Euclidean distance was used
278 to calculate the resemblance matrix. The biological response variable (Corg stock in g C m^{-2}) was
279 square-root transformed and Bray-Curtis similarity was used to calculate the abundance matrix.
280 The relative importance of different environmental variables was determined by use of DistLm,
281 a distance-based linear model procedure (Legendre and Anderson 1999). DistLm model was
282 constructed using a step-wise procedure that allows addition and removal of terms after each
283 step of the model construction. AICc (Akaike Information Criterion with a correction for finite
284 sample sizes) was chosen as information criterion as it enables to fit the best explanatory
285 environmental variables from of relatively small biological dataset compared to number of
286 environmental variables (Burnham and Anderson 2002). An alpha level of significance of 95 %
287 ($p < 0.05$) was used for all the analysis. All means are reported as mean \pm SE (SEM).

288

289 Results

290

291 Seagrass meadow and sediment characteristics

292

293 In general, the Finnish meadows were found on exposed sandy bottoms while the environmental
294 settings of the eelgrass meadows in Denmark were more variable (Fig. 2). Shoot density was
295 nearly equal in both regions, averaging at 417 ± 75 (shoots m^{-2}) in Finland and 418 ± 32 (shoots
296 m^{-2}) in Denmark. In Finland variation between sites (112-773 shoots m^{-2}) was greater than in

297 Denmark (300-652 shoots m⁻²). In Denmark the highest shoot density was found at the most
298 exposed site (Nyborg), while in Finland the highest shoot density was found at Sackholm, a fairly
299 sheltered site. The lowest shoot densities in Finland and Denmark were found in Tvärminne and
300 Løgstør, respectively. The mean aboveground biomasses were 101±3 and 145±5 (g DW m⁻²) and
301 the mean belowground biomasses 79±5 and 148±13 g (DW m⁻²) at Finnish and Danish sites,
302 respectively. In Denmark, the mean fraction of OC in aboveground and belowground *Z. marina*
303 tissue was 35±0.32 % DW and 29±1.10 % DW, respectively, while the corresponding numbers
304 for Finland were 38±0.24 % DW and 36±0.27 % DW, respectively. Given an average total *Z.*
305 *marina* biomass (above- and belowground) of 293± 22.31 (Denmark) and 180±9.60 g DW m⁻²
306 (Finland), we estimate the Corg pool in bound in living seagrass biomass to 0.66±0.005 and
307 0.94±0.014 t C ha⁻¹ in Finland and Denmark, respectively. The root: shoot-ratio was slightly
308 lower in Finland (0.87±0.05) than in Denmark (1.14±0.12), and varied between 0.29 to 3.29 and
309 0.15 to 6.45 in Finland and Denmark, respectively. The annual production of eelgrass for Finland
310 (average 524±62 g DW m⁻² y⁻¹) showed relatively low variation between sites (270-803 g DW m⁻²
311 y⁻¹) being lowest at Jänisholm and highest at Ryssholmen. In Denmark, the mean annual
312 eelgrass production was almost twice as high (928±159 g DW m⁻² y⁻¹) with large variation (470-
313 2172 g DW m⁻² y⁻¹). Production was lowest and highest at Dalby and Visby, respectively (Table
314 1).

315

316 The sediment characteristics varied significantly between Finland and Denmark. There was a
317 significant difference ($F_{1,9} = 14.7$, $p < 0.003$) between regions in terms of silt content, which was
318 generally lower at Finnish (6.3±1 %) sites than at Danish sites (20.2±3.9 %), although in
319 Denmark the variation between sites ranged from 0.8 % at Nyborg to 31.6 % at Thurøbund
320 (Table 1, Fig. 2). In Finland, the variation between sites was lower and ranged from 1.6%
321 (Kolaviken) to 15.5 % (Sackholm). At the Finnish sites the mean sediment dry density was
322 higher (1.35±0.01 g cm⁻³) compared to the Danish sites (1.25±0.02 g cm⁻³), and the Finnish sites
323 exhibited lower within-region variability ranging from 1.1 g cm⁻³ at Lyddaren to 1.5 cm⁻³ at

324 Långören, while the Danish sites varied from 0.3 g cm^{-3} at Thurøbund to 1.5 g cm^{-3} at Visby. The
325 Finnish sites showed consistently lower pools of organic matter (LOI: $1.4 \pm 0.3 \text{ \%DW}$) compared
326 to the average of Danish sites (LOI: $3.9 \pm 1.5 \text{ \% DW}$). Similarly, the mean OC content was lower in
327 Finland (0.24 ± 0.033) than in Denmark (1.75 ± 0.563). Consequently, the mean water content was
328 similarly lower in Finland ($20.9 \pm 0.4 \text{ \%}$: range 16-29 %) than in Denmark ($37.4 \pm 1.8 \text{ \%}$: range 17-
329 76 %) (Table 1). Sediment porosity was similar in both regions, and ranged from 0.25 to 0.30 in
330 Finland, and from 0.20 to 0.40 in Denmark. At the Finnish sites, the proxy (degree of sorting)
331 that was used to estimate exposure, varied from 0.8 to 1.5 (ϕ), with Kolaviken being the most
332 exposed and Ängsö being the most sheltered site. In Denmark degree of sorting varied from 0.6
333 to 2.1 (ϕ), with Nyborg and Visby being the most exposed and sheltered sites, respectively (Fig.
334 2).

335

336 Organic carbon stocks

337 The profiles of carbon densities (g C cm^{-3}) in the upper 25 cm of the sediment showed marked
338 differences both between and within the sampled regions. At the Finnish sites, where eelgrass
339 typically grows at exposed locations, the sediment carbon density was low (mean $2.6 \pm 0.09 \text{ mg C}$
340 cm^{-3}) and declined with depth at most of the 10 study sites (Fig. 3). At the Danish sites, however,
341 the sediment carbon density was more variable (mean $17.45 \pm 9.42 \text{ mg C cm}^{-3}$) both within and
342 between sites (Fig. 3). Depth integrated Corg stocks (0-25 cm, g C m^{-2} , Fig. 4) were particularly
343 high at one sheltered site in Funen, namely Thurøbund. This site is characterized by soft
344 sediments with high organic content, high annual eelgrass production and high belowground
345 biomass (Table 1). The lowest eelgrass Corg stocks in Denmark were found at two relatively
346 exposed and sandy sites, namely Nyborg and Dalby (Fig. 4). The estimate of average total Corg
347 stock in Finland was $0.019 \pm 0.001 \text{ Mt C}$ taking the total area of eelgrass into account (30 km^2 ;
348 Table 2). Using minimum and maximum estimates of the eelgrass area in Denmark the estimates

349 for mean total sediment Corg stock in Denmark were 2.164 ± 0.005 Mt C or 5.868 ± 0.014 Mt C,
350 respectively (673 and 1345 km²; Table 2).

351

352 Using an annual carbon accumulation value of 0.05 and 0.35 t C ha⁻¹ y⁻¹ for Finland and
353 Denmark, respectively, and assuming sediment accumulation of 2.02 mm y⁻¹ on average (Table
354 2), the total pool of Corg in the *Z. marina* meadows (Corg bound in living biomass, sediment Corg
355 stock and Corg accumulation) corresponds to 6.98 t C ha⁻¹ (698t km⁻²) and 44.9 t C ha⁻¹ (4490 t
356 km⁻²) for Finland and Denmark, respectively. Using the social cost of carbon of 40.3 € t C⁻¹
357 (United States Government 2010), the present economic value of eelgrass carbon in Finnish and
358 Danish eelgrass meadows is estimated at 281 and 1809 € ha⁻¹, respectively. Using an average of
359 these values (1045 € ha⁻¹) and a conservative estimate of the eelgrass acreage in the Baltic Sea
360 (2100 km²: Boström et al. 2014), we estimate a total monetary value of the present
361 sequestration by eelgrass meadows to be 219.4 million euro. Given the total eelgrass loss in
362 Denmark for the time period 1900-2000 is between 5381 km² (present area 20 % of historical
363 distribution) and 6053 km² (present area 10 % of historical distribution), this equals to a Corg
364 loss of 0.042 and 0.048 Gt C, respectively. Using the acreage value (1045 € ha⁻¹) these areal loss
365 estimates corresponds to a lost economic value between 562 and 632 million euro, for the
366 minimum and maximum areal loss estimates, respectively.

367

368 Corg accumulation

369

370 The estimates for annual Corg accumulation in the Finnish seagrass meadows (30 km²) were
371 low (0.002, 0.016, 0.033 Mt C y⁻¹), when applying sediment accumulation rates of 0.32, 2.02 and
372 4.20 mm y⁻¹, respectively. The low Corg accumulation in Finnish meadows was a result of low
373 mean Corg stocks and relatively small size of seagrass area compared to Denmark (Table 2). The
374 estimates for annual Corg accumulation for the Danish sites differed between the two sub-
375 regions Limfjorden (18 km²) and Funen (179 km²). At the sampling sites around Funen, the Corg

376 accumulation was 0.139, 0.881 and 1.832 Mt C y⁻¹, while in Limfjorden the Corg accumulation
377 was lower (0.006, 0.038 and 0.079 Mt C y⁻¹) and similar to Corg accumulation for Finland. Using
378 upper and lower eelgrass areal estimates, total Corg accumulation based on 3 sediment
379 accumulation rates in Denmark were more than four orders of magnitude higher (0.376, 2.373,
380 3.636 and 0.75, 4.741 and 9.859 Mt C y⁻¹) than the estimated total Corg accumulation in Finnish
381 eelgrass meadows.

382

383 Carbon sources

384

385 The $\delta^{13}\text{C}$ values of the surface sediment within regions were quite homogenous ranging from -
386 22.8 to -18.9 ‰ and -17.6 to 13.5 ‰, in Finland and Denmark respectively. The analytical error
387 for the sediment $\delta^{13}\text{C}$ values was 2.8 ‰. The $\delta^{13}\text{C}$ in *Z. marina* tissues ranged from -11.4 to -8.5
388 ‰ and from -12.5 to -8.2‰, in Finland and Denmark, respectively. There was no significant
389 difference between living above- and belowground tissue and decomposed belowground tissue
390 and samples were pooled in the isotope mixing model. Although *Z. marina* was the dominant
391 seagrass species in Finland, the study sites included both monospecific and mixed seagrass
392 meadows. Mixed meadows typically contained pondweeds, e.g. *Potamogeton pectinatus* and
393 *Potamogeton perfoliatus*. In particular, *P. pectinatus* ($\delta^{13}\text{C}$ -11.3 to -7.6 ‰) and *P. perfoliatus*
394 ($\delta^{13}\text{C}$ -15.6 to -12.6 ‰) were both present at five of the Finnish study sites (Jänisholm,
395 Sackholm, Hummelskär, Tvärminne and Fårö) and *P. pectinatus* was present at Kolaviken,
396 Ryssholmen and Lyddaren. *Ruppia cirrhosa* (-11.5 to -8.8 ‰) was less abundant and found at
397 three of the Finnish sites (Sackholm, Ängsö, Kolaviken) and at one study site in Denmark
398 (Kertinge). The $\delta^{13}\text{C}$ for phytoplankton ranged from -24.6 to -22.6 ‰ and -18.6 to 16.4 ‰, in
399 Finland and Denmark, respectively. Drift algae was present at all Danish study sites, except
400 Thurøbund, and had $\delta^{13}\text{C}$ values from -17.9 to -13.5 ‰, but only at five Finnish sites (Ängsö,
401 Ryssholmen, Fårö, Långören and Hummelskär) with $\delta^{13}\text{C}$ values ranging from -20.0 to -16.3‰.

402

403 The isotope mixing model indicated that at all Finnish sites, phytoplanktonic material was the
404 major contributor (43-86 %) to the sediment surface Corg pool. In Denmark *Z. marina*
405 contributed with 13-81 % to the sediment surface Corg pool, contribution being lowest at the
406 most exposed site in Nyborg and highest in Visby. The corresponding numbers for Finland were
407 1.5-32 %, being lowest and highest in Tvärminne and Lyddaren, respectively (Fig. 5).

408

409 Environmental factors explaining carbon pools

410

411 The combined (FIN + DK) DistLm analysis showed, that three sediment variables (dry density,
412 silt content, porosity) and three plant variables (annual eelgrass production, the root:shoot-ratio
413 and *Z. marina* contribution to the sediment carbon pool) explained 67 % of the variation in the
414 sediment Corg stock (g C m^{-2}) (Table 3 and 4, Fig. 6). Specifically, sediment silt content alone
415 explained > 36 % of the variation in Corg stocks (Table 3). In both regions, exposed sites
416 characterized by sandy, low organic sediments and low silt content, had low Corg stocks. In
417 contrast, at sheltered sites like Thurøbund in Denmark, we measured the highest sediment Corg
418 stock along with highest silt and water content among all sites. Although sediment porosity and
419 sediment dry density also contributed to the model, they were of minor importance (~2 %
420 each).

421 The combined (FIN+ DK) DistLm analysis also showed that the *Z. marina* contribution to the
422 sediment surface carbon pool explained 10.9 % of the variation in the measured Corg stocks
423 (Fig. 6, Table 3 and Table 4). Drift algae was a significant contributor (72 %) to the sediment
424 surface Corg pool at the Danish sites, while it appeared to play only a minor role (0-21 %) in
425 Finland. The carbon sources were generally more mixed at the Danish study sites compared to
426 the Finnish sites where phytoplankton dominated (Fig. 5).

427

428 While the overall model including all sites explained almost 70 % of the variation in carbon

429 stocks (Table 3, 4), and indicated that the most relevant environmental variables were included
430 in this Baltic scale analysis, reduced, countrywise DistLM models revealed different results. In
431 particular, variability in Finnish carbon stocks were explained up to 50 % by geological variables
432 (porosity, sorting and sediment dry density), while the best sequential model for carbon stock
433 variability at Danish sites explained 75 % of the total variance. In contrast to Finland, the role of
434 eelgrass related variables (relative proportion of *Z. marina* in the sediment and the root:shoot
435 ratio) were most important and explained 40 % and 25 %, respectively of the carbon stock
436 variability.

437

438 Discussion

439

440 Recent studies have shown considerable variation in the global estimates of carbon stocks (Corg
441 stocks) and carbon accumulation rates (Corg accumulation) in seagrass meadows, indicating an
442 incomplete understanding of factors influencing this variability (Fourqurean et al. 2012, Duarte
443 et al. 2013a, Lavery et al. 2013, Miyayima et al. 2015). The Baltic Sea forms a key distribution
444 area for eelgrass in Europe, but similarly to the global data sets, we have so far lacked estimates
445 on seagrass carbon stocks and accumulation.

446

447 In our study, the Finnish eelgrass meadows showed consistently very low Corg stocks and Corg
448 accumulation, and the meadows were minor carbon sinks compared to the Danish meadows.
449 The Danish sites showed more variation in the sediment Corg stock and accumulation and Corg
450 stock was particularly high at one site, Thurøbund ($26138 \pm 385 \text{ g C m}^{-2}$), which is a relatively
451 sheltered site with high organic sediments. Expectedly, due to both larger overall eelgrass
452 acreage and larger Corg stocks in the Danish meadows, the total Corg accumulation ($0.38\text{-}9.86 \text{ t}$
453 C y^{-1}) was three to four orders of magnitude higher than in the Finnish meadows ($0.002\text{-}0.033 \text{ t}$
454 C y^{-1}). As eelgrass in Finland generally grow in more exposed locations potentially due to

455 increased interspecific competition with freshwater plants such as common reed (*Phragmites*
456 *australis*) in sheltered locations (Boström et al. 2006), it is probable that most of the Corg
457 produced in the Finnish meadows is exported, and thus incorporated in detrital food webs in
458 deeper bottoms. This argument is supported when applying sediment accumulation rates from
459 literature, as only 0.15 – 2.0 % of the annual production accumulated in Finnish meadows, while
460 the corresponding numbers for Denmark were 0.6 -7.8 %. Duarte and Cebrian (1996) estimated
461 that on average 25 % of the global seagrass primary production is exported, and seagrass
462 detritus may thus contribute significantly to Corg stocks in other locations, a fact that is often
463 overlooked.

464

465 Extrinsic drivers of carbon sequestration in seagrass meadows

466

467 As proposed in previous work accumulation of fine grained size fractions in seagrass sediments,
468 relative to those accumulated in bare sediments, appears to be one of the major factors
469 influencing the carbon sink capacity of seagrass meadows (Kennedy et al. 2010, Miyajima et al.
470 2015), and may thus be a useful proxy for the sink capacity. In addition, it is well known, that
471 seagrasses modify sediments by reducing water flow and consequently increasing particle
472 trapping and sedimentation and reducing resuspension (Fonseca and Fisher 1986, Fonseca and
473 Cahalan 1992, Gacia et al. 2002, Hendriks et al. 2008, Boström et al. 2010) and also increasing
474 Corg (Kennedy et al. 2010).

475

476 In this study, the DistLm analysis showed, that contribution of *Z. marina* to the sediment surface
477 carbon pool was an important driver (> 10.9 %) of the variation in the sediment Corg stock
478 (Table 3 and 4, Fig. 6) when the model included both regions. Surprisingly, the reduced
479 countrywise analysis revealed different results and showed that *Z. marina* contribution to the
480 sediment surface carbon pool was an important driver for Corg stocks only in Denmark. We

481 believe that the countrywise differences in explanatory variables might relate to the more
482 pronounced influence of eelgrass for carbon stocks at Danish sites. Indeed, these sites exhibited
483 on average 30 % higher aboveground biomasses, 45 % higher belowground biomasses, 24 %
484 higher root:shoot- ratios and 44 % higher productivity compared to the Finnish sites. In Finland
485 and the northern Baltic Sea, eelgrass meadows appears to be primarily physically controlled,
486 and thus sediment variables play a relatively more pronounced role. The results from the model
487 were also supported by our data in which we found increasing Corg stocks at the Danish sites,
488 where *Z. marina* was the major source of organic carbon, contributing with 13-81% to the
489 surface sediment Corg, while in contrast, at the Finnish sites where only a minor fraction of
490 carbon buried in sediments derive from eelgrass detritus (1.5-39.6 %) the Corg stocks were low.
491 Correspondingly, the average $\delta^{13}\text{C}$ value (-16.2 ‰) in the Danish sediment samples was similar
492 to the global median value (-16.3±0.2 ‰) reported by Kennedy et al. (2010) in which on
493 average 51 % of the carbon was derived from seagrass detritus, whereas it was -20±0.6 ‰ in
494 Finland indicative of higher contribution from other more negative carbon sources., such as
495 phytoplankton. The importance of the *Z. marina* contribution to the Corg stocks may be
496 explained by slow decomposition rates of seagrass tissue. Especially, the high proportion of
497 refractory organic compounds in the seagrass belowground parts and high C:N:P-ratios of
498 seagrass tissue in general make seagrasses less biodegradable than most marine plants and
499 algae (Fourqurean and Schrlau 2003, Vichkovitten and Holmer 2004, Kennedy and Björk 2009,
500 Holmer et al. 2011, Röhr et al in prep.). The slow decomposition rates are also a result of
501 reduced sediment conditions commonly encountered in Danish seagrass meadows (Kristensen
502 and Holmer 2001, Holmer et al. 2009, Pedersen et al. 2011). Despite the extensive distribution
503 (2-29 ha), high biomasses (300-800 g DW m⁻²) and major impact of drifting algal mats on coastal
504 ecosystem functioning (Norkko and Bonsdorff 1996, Salovius and Bonsdorff 2004, Rasmussen et
505 al. 2013, Gustafsson and Boström 2014), the stable isotope composition of the sediments
506 suggests that drift algae had a surprisingly minor influence on the sediment surface Corg pool in
507 both regions. Thus, despite present on several sampling sites, drift algae is likely exported and

508 mineralized in deeper sedimentation basins. Furthermore, we found that at all study sites in
509 both regions, there were several other potential sources influencing the sediment surface Corg
510 pool. Similarly, Bouillon et al. (2007) showed that in seagrass sediments adjacent to mangrove
511 forests in Kenya, none of their sites had seagrass material as the sole source of Corg, and instead
512 mangrove-derived detritus contributed significantly to the seagrass sediment Corg pool.

513

514 Similarly to the contribution of *Z. marina* to the sediment surface carbon pool, the root:shoot-
515 ratio explained 12.7 % of the variation in the Corg stocks when both regions were included in
516 the model, but in the reduced countrywise models it was important driver for Corg stocks in
517 only Denmark. Accordingly, the highest Corg stocks, belowground biomass and root: shoot-ratio
518 was found in Thurøbund (Denmark). In Finland, the highest root: shoot-ratio (2.07) was found at
519 Kolaviken, with a relatively low Corg stock (397 g C m⁻²). Due to higher degree of exposure at the
520 site (degree of sorting 0.7 ϕ) compared to Thurøbund (1.4 ϕ) it is likely that large portion of the
521 eelgrass production was exported away from the meadow and not stored in the sediment. The
522 mean shoot densities were almost identical between regions, and shoot density did not
523 contribute to the model explaining Corg.

524

525 The annual eelgrass production explained only 2.3 % of the variation in the Corg stocks in the
526 combined model. The annual production rates were almost twice as high at Danish sites
527 compared to the Finnish sites. Regional differences in seagrass productivity may be caused by
528 differences in e.g. the inorganic carbon concentration in water column and light availability
529 between the regions (with higher values in Denmark), which both affect the photosynthetic
530 capacity of the plant (Hellblom and Björk 1999, Holmer et al. 2009, Boström et al. 2014).
531 Eelgrass production tend to be higher in physically exposed areas compared to more sheltered
532 areas, which can be due to improved sediment oxygen conditions and hydrodynamical effects
533 (Hemminga and Duarte 2000). This finding was not supported by our study, in which we found

534 the highest annual eelgrass production rates at both the most sheltered and exposed sites,
535 namely Visby and Nyborg (DK).

536

537 Geographical comparisons of carbon stocks and accumulation

538

539 Our estimated Corg stocks for the study sites were generally lower (627-4324 t C km⁻²) than
540 estimates (25200-84000 t C km⁻²) found in the literature (Nelleman et al. 2009, , Fourqurean et
541 al. 2012). Several of the studies were conducted in the Mediterranean *P. oceanica* meadows - a
542 habitat with superior carbon sequestration and storage capacity (Duarte et al. 2005, Lavery et al.
543 2013). The average sizes of Corg stocks in Finnish and Danish eelgrass meadows were also
544 considerably lower than the mean values reported by Alongi et al. (2014) for tropical seagrass
545 meadows (14270 t C km⁻²). In contrast, our estimate for the carbon stock in the top 25 cm for
546 Danish and Finnish meadows (627-6005 g C m⁻²) are comparable to Australian (262-4833 g C m⁻²;
547 Lavery et al. 2013) and Asian estimates (3800-12000 g C m⁻²; Miyajima et al. 2015).

548

549 Consequences of seagrass loss for carbon pools

550

551 Despite the importance of seagrasses, their global distribution has decreased by 29 % since
552 1879 primarily due to anthropogenic pressures (Waycott et al. 2009), thus weakening the
553 carbon sink capacity of marine environments to sequester carbon (Duarte et al. 2005). Since the
554 1970s, the Baltic Sea has been subject to strong anthropogenic pressures (Conley et al. 2009)
555 leading to eelgrass declines in several countries (Boström et al. 2014). In the 1930s, the Danish
556 eelgrass meadows were significantly reduced by the wasting disease (Rasmussen 1977). These
557 regime shifts in Denmark have resulted in a 80-90 % decline corresponding to 6726 km² in the
558 beginning of 1900`s to 673-1345 km² in 2005, using the minimum and maximum estimates for
559 the current coverage area, respectively (Boström et al. 2014). Using the mean carbon density

560 (17.45 mg C cm⁻³) measured at the Danish sites, the lost Corg stock is estimated to 23-27 Mt C
561 and these large-scale seagrass declines, which are also found in Sweden and Germany, have
562 eroded the Corg stocks in the Baltic Sea significantly (Table 2). In Finland there is a lack of long-
563 term monitoring, but the meadows appear to be stable and cover at least 30 km² with no
564 significant loss of Corg stocks.

565

566 Using a carbon monetary value of 40.3€ C⁻¹, we calculated the monetary value of the present
567 carbon storage and sequestration capacity of eelgrass meadows in Finland and Denmark to be
568 281 and 1809 € ha⁻¹, respectively. Pendleton et al. (2014) calculated a global estimated
569 economic cost of lost seagrass meadows to be 1.9-13.7 billion USD. This value was derived from
570 the cost of lost carbon sink capacity, ignoring other lost ecosystem services including e.g. coastal
571 protection, water quality management, food provision and the role of seagrasses as fisheries and
572 key habitats for marine species (Barbier et al. 2011, Atwood et al. 2015). Our estimate also only
573 considers lost carbon sink capacity and can be compared directly with Pendleton et al. (2014).
574 The present economic value of carbon storage and sequestration capacity of Baltic Sea eelgrass
575 meadows is thus between 1.7 and 12 % out of the global seagrass Blue Carbon value.

576

577 While useful, our and previous work still remain snap shots of complex processes causing local
578 and regional variability in estimates of seagrass Blue Carbon stocks and accumulation. Clearly, in
579 order to produce more reliable estimates of global seagrass carbon sequestration rates and
580 stocks, there is a need for more studies integrating and modeling the individual and joint role of
581 e.g. sediment biogeochemistry, seascape structure, plant species architecture and hydrodynamic
582 regime. Since seagrasses are lost at accelerating rates (Waycott et al. 2009), there is also an
583 urgent need for a better understanding of the fate of lost seagrass carbon (Macreadie et al. 2014)
584 and the development of the carbon sink capacity in restored seagrass ecosystems (Nellemann et
585 al. 2009, Greiner et al. 2013, Marba et al. 2015). Nelleman et al. (2009) proposed the use of
586 carbon trading programs using financial incentives for forest conservation, such as REDD+

587 (Reduced Emissions from Deforestation and Degradation) and NAMAs (Nationally Appropriate
588 Mitigation Actions), to include the blue carbon ecosystems as part of their environmental
589 protection protocol. Both of these carbon mitigation programs require ongoing monitoring of
590 organic carbon storage and emission in the different Blue Carbon ecosystems. In order to
591 manage seagrass meadows, mitigate climate change and produce information required for the
592 carbon trading programs, it is fundamental to understand factors influencing the capacity of
593 seagrass meadows to capture and store carbon. By solving these uncertainties, the conservation
594 and restoration of seagrass meadows can be implemented in the most beneficial manner by e.g.
595 giving priority to protection of the seagrass meadows and species with the highest carbon sink
596 capacity and foundation of restoration projects in areas most suitable for seagrass growth
597 (Duarte et al. 2013a).

598 Data availability and competing interest

599

600 The data is available from the corresponding author. The authors declare that they have no
601 conflict of interest.

602 Acknowledgements

603

604 This work is part of a double degree program between Åbo Akademi University (ÅAU) and
605 University of Southern Denmark (SDU). The study was funded by The Maj and Tor Nessling
606 Foundation (Project 201600125: Baltic Sea blue carbon: environmental gradients influencing
607 the carbon sink capacity of seagrass meadows) and University of Southern Denmark. We
608 acknowledge Archipelago Centre Korpoström and University of Southern Denmark for excellent
609 working facilities. Tiina Salo and Karine Gagnon are acknowledged for field assistance and
610 Katrine Kierkegaard and Birthe Christensen for their assistance in the laboratory work.

611

612

- Alongi, D. M.: Carbon Cycling and Storage in Mangrove Forests. *Ann. Rev. Mar. Sci.* 6:195–219, 2014.
- Anderson, M.J., Gorley, R.N. and Clarke, K.R.: PERMANOVA for PRIMER: guide to software and statistical methods. Plymouth, United Kingdom: PRIMER-E Ltd., 2008.
- Atwood, T.B., Connolly, R.M., Euan, G.R., Lovelock, C.E., Heithaus, M.R. and Hays, G.C, Predators help protect carbon stocks in blue carbon ecosystem. *Nat. Clim. Change* 2015.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. and Silliman, B.R.: The value of estuarine and coastal ecosystem services. *Ecol. Monogr.*,81(2):169-193, 2011.
- Boström C., Baden S., Bockelmann A., Dromph K., Fredriksen S. and Gustafsson C.: Distribution, structure and function of Nordic eelgrass (*Zostera marina*) ecosystems: implications for coastal management and conservation. *Aq. Cons. Mar. Freshwater Ecosyst.*,24(3):410-434, 2014
- Boström, C., Törnroos, A. and Bonsdorff, E.: Invertebrate dispersal and habitat heterogeneity: Expression of biological traits in a seagrass landscape. *J. Exp. Mar. Biol. Ecol.*, 390(2):106-117, 2010.
- Boström, C., Bonsdorff, E., Kangas, P. and Norkko, A.: Long-term changes of a brackish-water eelgrass (*Zostera marina* L.) community indicate effects of coastal eutrophication. *Estuar. Coast. and Shelf Sci.* 55(5):795-804, 2002.
- Boström, C., O'Brien, K., Roos, C. and Ekeboom, J.: Environmental variables explaining structural and functional diversity of seagrass macrofauna in an archipelago landscape. *J. Exp. Mar. Biol. Ecol.* ,335(1):52-73, 2006
- Bouillon S. and Boschker H.TS.: Bacterial carbon sources in coastal sediments: a cross-system analysis based on stable isotope data of biomarkers. *Biogeosciences*, 3(2):175-185, 2006.
- Bouillon, S., Dahdouh-Guebas, F., Rao, A.V.V.S., Koedam, N. and Dehairs, F.: Sources of organic carbon in mangrove sediments: variability and possible ecological implications. *Hydrobiologia*, 495(1-3):33-39, 2003.
- Bouillon, S., Dehairs, F., Velimirov, B., Abril, G. and Borges, A.V.: Dynamics of organic and inorganic carbon across contiguous mangrove and seagrass systems (Gazi Bay, Kenya). *J. Geophys. Research-Biogeosciences* 112(G2):G02018, 2007.
- Burnham, K.P. and Anderson, D.R.: *Model Selection and Multi-Model Inference: A Practical Information-Theoretic Approach* (2nd Edition). Secaucus, NJ, USA: Springer, 2002.
- Conley, D.J., Björck, S., Carstensen, J., Bonsdorff, E., Destouni, G. and Gustafsson, B.G.: Hypoxia-related processes in the Baltic Sea. *Env. Sci. Tech.*, 2009.
- DMU-report, Ærtebjerg, G., Andersen, J.H. and Hansen, O.S.(Eds.): *Nutrients and Eutrophication in Danish Marine Waters . A Challenge for Science and Management*, National Environmental Research Institute, Denmark, 2003.

- Downie, A., von Numers, M. and Boström, C.: Influence of model selection on the predicted distribution of the seagrass *Zostera marina*. *Estuar. Coast. and Shelf S.*, 121:8-19, 2013.
- Duarte, C.M. and Cebrian, J.: The fate of marine autotrophic production. *Limnol. Oceanogr.*, 41(8):1758-1766, 1996.
- Duarte, C.M., Middelburg, J.J. and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences*, 2(1):1-8, 2005.
- Duarte, C.M., Kennedy, H., Marba, N. and Hendriks, I.: Assessing the capacity of seagrass meadows for carbon burial: Current limitations and future strategies. *Ocean Coast Man.*, 83:32-38 2013.
- Duarte, C.M., Losada, I.J., Hendriks, I.E., Mazarrasa, I. and Marba, N.: The role of coastal plant communities for climate change mitigation and adaptation. *Nature Clim. Change*, 3 (11):961-968, 2013.
- Folk, R.L. and Ward, W.C.: Brazos river bar: a study of significance of grain size parameters. *J. Sediment Petrol* 1957;27:3, 1957
- Fonseca and Fisher, J.S.: A comparison of canopy friction and sediment movement between four species of seagrass with reference to their ecology and restoration. *Marine ecology progress series*. Oldendorf, 29(1):15-22, 1986
- Fonseca, M.S. and Cahalan, J.A.: A preliminary evaluation of wave attenuation by four species of seagrass. *Estuar. Coast. Shelf Sci.* 1992 12; 35(6):565-576, 1992.
- Fourqurean, J.W. and Schrlau, J.E. Changes in nutrient content and stable isotope ratios of C and N during decomposition of seagrasses and mangrove leaves along a nutrient availability gradient in Florida Bay, USA. *Chem. Ecol.* 19:373-390, 2003.
- Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marba, N., Holmer, M. and Mateo, A.M.: Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience*, 5(7):505-509, 2012.
- Fry B. and Sherr E.B. $\delta^{13}C$ measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contrib. Mar. Sci.*, 27:13, 1984.
- Fry, B., Scalan, R.S., Parker, P.L.: Stable Carbon Isotope Evidence for 2 Sources of Organic-Matter in Coastal Sediments - Seagrasses and Plankton. *Geochim. Cosmochim. Acta* 41(12):1875-1877, 1977.
- Gacia, E., Granata, T.C. and Duarte, C.M.: An approach to measurement of particle flux and sediment retention within seagrass (*Posidonia oceanica*) meadows. *Aquat. Bot.*, 65(1-4):255-268, 1999.
- Gacia, E. and Duarte, C.M. and Middelburg, J.J.: Carbon and nutrient deposition in a Mediterranean seagrass (*Posidonia oceanica*) meadow. *Limnol. Oceanogr.* 47(1):23-32, 2002.
- Gattuso, J.P., Frankignoulle, M. and Wollast R. Carbon and carbonate metabolism in coastal aquatic ecosystems. *Ann. Rev. Ecol. Syst.*, 29:405-434, 1998.
- Greiner, J.T., McGlathery, K.J., Gunnell, J. and McKee, B.A. Seagrass Restoration Enhances "Blue Carbon" Sequestration in Coastal Waters. *Plos One*, 8(8):e72469, 2013.
- Gustafsson, C. and Boström, C.: Algal mats reduce eelgrass (*Zostera marina* L.) growth in mixed and monospecific meadows. *Journal of Experimental Marine Biology and Ecology*. 461:85-92, 2014.

- Hendriks, I.E., Sintes, T., Bouma, T.J. and Duarte, C.M.: Experimental assessment and modeling evaluation of the effects of the seagrass *Posidonia oceanica* on flow and particle trapping. *Mar. Ecol. Prog. Ser.* 356:163-173, 2008.
- Herr, D., Pidgeon, E., Laffoley, D. (Eds.): *Blue Carbon. Policy Framework: Based on the Discussion of the International Blue Carbon Policy Working Group*, IUCN and Arlington, Gland, Switzerland, USA: CI. viþ39pp, 2012.
- Holmer, M., Duarte, C.M., Boschker, H.T.S. and Barron, C.: Carbon cycling and bacterial carbon sources in pristine and impacted Mediterranean seagrass sediments. *Aquat. Microb. Ecol.*, 36(3):227-237, 2004.
- Holmer, M., Baden, S., Bostrom C and Moksnes P.: Regional variation in eelgrass (*Zostera marina*) morphology, production and stable sulfur isotopic composition along the Baltic Sea and Skagerrak coasts. *Aquat. Bot.*, 91(4):303-310, 2009.
- Holmer, M. 2009. Productivity and biogeochemical cycling in seagrass ecosystems. In *Coastal Wetlands: An Integrated Ecosystem Approach*, ed. G.M.E. Perillo, E. Wolanski, D.R. Cahoon and M.M. Brinson, 377-401. Amsterdam, The Netherlands.
- Holmer, M., Wirachwong, P. and Thomsen, M.S.: Negative effects of stress-resistant drift algae and high temperature on a small ephemeral seagrass species. *Mar. Biol.*, 158(2):297-309, 2011.
- Kennedy, H., Gacia, E., Kennedy, D.P., Papadimitriou, S. and Duarte, C.M.: Organic carbon sources to SE Asian coastal sediments. *Estuar. Coast. and Shelf S.*, 60(1):59-68, 2004.
- Kennedy, H. and Björk, M.: *Seagrasses, in The Management of Natural Coastal Carbon Sinks in Coastal Ecosystems: Investigating and Realising the Potential.* 2009.
- Kennedy, H., Beggins, J., Duarte, C.M., Fourqurean, J.W., Holmer, M. and Marba, N.: Seagrass sediments as a global carbon sink: Isotopic constraints. *Glob. Biogeochem. Cycles*, 24:GB4026, 2010.
- Kristensen, E. and Holmer, M.: Decomposition of plant materials in marine sediment exposed to different electron acceptors (O₂, NO₃⁻, and SO₄²⁻), with emphasis on substrate origin, degradation kinetics, and the role of bioturbation. *Geochim. Cosmochim. Acta*, 65(3):419-433, 2001.
- Lavery, P.S., Mateo, M., Serrano, O. and Rozaimi, M.: Variability in the Carbon Storage of Seagrass Habitats and Its Implications for Global Estimates of Blue Carbon Ecosystem Service. *Plos One*, 8(9):e73748, 2013.
- Legendre, P. and Anderson, M.J.: Distance-based redundancy analysis: Testing multispecies responses in multifactorial ecological experiments (vol 69, pg 1). *Ecol. Monogr.*, 69 (4):512-512, 1999.
- Leipe, T., Tauber, F., Vallius, H., Virtasalo, J., Uścińowicz, S., Kowalski, N., Hille, S., Lindgren, S., and Myllyvirta, T.: Particulate organic carbon (POC) in surface sediments of the Baltic Sea. *Geo-Mar. Lett.* 31:175-188, 2011.
- Macreadie, P.I., Baird, M.E., Trevathan-Tackett, S.M., Larkum, A.W.D. and Ralph, P.J.: Quantifying and modelling the carbon sequestration capacity of seagrass meadows - A critical assessment. *Mar. Pollut. Bull.*, 83(2):430-439, 2014.

- Marba, N., Arias-Ortiz, A., Masque, P., Kendrick, G. A., Mazarrasa, I., Bastyan, G. R., Garcia-Orellana, J. and Duarte, C. M.: Impact of seagrass loss and subsequent revegetation on carbon sequestration and stocks. *J. of Ecology* 103:296-302, 2015
- Mcleod, E., Chmura, G.L., Bouillon, S., Salm, R., Bjork, M. and Duarte, C.M.: A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front. Ecol. Env.*, 9(10):552-560, 2011.
- Miyajima, T., Hori, M., Hamaguchi, M., Shimabukuro, H., Adachi, H. and Yamano, H.: Geographic variability in organic carbon stock and accumulation rate in sediments of East and Southeast Asian seagrass meadows. *Glob. Biogeochem. Cycles*, 29(4):397-415, 2015.
- Moksnes, P.-E. and Cole, S. G.: Valuing multiple eelgrass ecosystem services in Sweden: Fish production and uptake of carbon and nitrogen. *Front. Mar. Sci.* 2(121):1-18, 2015.
- Moncreiff, C.A. and Sullivan M.J.: Trophic importance of epiphytic algae in subtropical seagrass beds: evidence from multiple stable isotope analyses. *Mar. Ecol. Prog. Ser.*, 215:93-106, 2001.
- Nellemann, C., Corcoran, E., Duarte, C.M., Valdés, L., De Young, C. and Fonseca, L.: Blue Carbon. A Rapid Response Assessment, 2009.
- Norkko, A. and Bonsdorff, E.: Rapid zoobenthic community responses to accumulations of drifting algae. *Mar. Ecol. Prog. Ser.*, 131:143-157, 1996.
- Olesen, B. and Sandjensen, K.: Patch Dynamics of Eelgrass *Zostera marina*. *Mar. Ecol. Prog. Ser.* 106(1-2):147-156, 1994.
- Orth, R.J., Carruthers, T.J.B., Dennison, W.C., Duarte, C.M., Fourqurean, J.W. and Heck, K. L.Jr. : A global crisis for seagrass ecosystems. *Bioscience*, 56(12):987-996, 2006.
- Pedersen, M. Ø., Serrano, O., Mateo, M.A, Holmer, M.: Temperature effects on decomposition of a *Posidonia oceanica* mat. *Aquat. Microb. Ecol.* 65: 169-182, 2011.
- Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A. and Sifleet, S: Estimating Global "Blue Carbon" Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *Plos One*, 7(9):e43542, 2012.
- Petersen, J.K., Maar, M., Ysebaert, T. and Herman, P.M.J.: Near-bed gradients in particles and nutrients above a mussel bed in the Limfjorden: influence of physical mixing and mussel filtration. *Mar. Ecol. Prog. Ser.*, 490:137-146, 2013.
- Phillips, D.L. and Gregg, J.W.: Source partitioning using stable isotopes: coping with too many sources. *Oecologia*, 136(2):261-269, 2003.
- Rask, N., Pedersen, S.E. and Jensen, M.H.: Response to lowered nutrient discharges in the coastal waters around the island of Funen, Denmark. *Hydrobiologia*, 393:69, 1999.
- Rasmussen, E.: The wasting disease of eelgrass *Zostera marina* and its effects on environmental factors and fauna. *Seagrass ecosystems. A scientific perspective*. New York: Marcel Dekker Inc., 1977.
- Rasmussen, J.R., Pedersen, M.F., Olesen, B., Nielsen, S.L. and Pedersen, T. M. : Temporal and spatial dynamics of ephemeral drift-algae in eelgrass *Zostera marina*, beds. *Estuar. Coast. and Shelf S.*

119:167-175, 2013.

Salovius, S. and Bonsdorff, E.: Effects of depth, sediment and grazers on the degradation of drifting filamentous algae (*Cladophora glomerata* and *Pilayella littoralis*). *Journal of Experimental Marine Biology and Ecology*. 298:93-109, 2004.

Schlacher, T.A and Conolly, R.M.: Effects of acid treatment on carbon and nitrogen stable isotope ratios in ecological samples: a review and synthesis. *Methods in Ecology and Evolution* 5(6):541-550, 2014.

Serrano, O., Lavery, P.S., Rozaimi, M. and Mateo, A. M.: Influence of water depth on the carbon sequestration capacity of seagrasses. *Global Biogeochem. Cycles* 28(9):950-961, 2014.

Short, F. and Duarte, C.M.: Methods for the measurement of seagrass growth and production. In: Short and Cole, editors. *Global seagrass research methods* Amsterdam: Elsevier; p. 155-182, 2001.

Spalding, M., Taylor, M., Ravilious, C., Short, F., Green, E.: Global overview: The distribution and status of seagrasses. In: Green, E.P. and Short F.T.: editors. *World Atlas of Seagrasses: Present Status and Future Conservation*. Berkeley California: University of California Press, p. 5-26, 2003.

United States Government. Technical Support Document: Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866 (United States Government Interagency Working Group on Social Cost of Carbon, Washington, DC), 2010.

Vichkovitten, T. and Holmer, M. Contribution of plant carbohydrates to sedimentary carbon mineralization. *Org. Geochem.*, 35(9):1053-1066, 2004.

Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C. and Olyarnik, S.: Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. Natl. Acad. Sci. USA*, 106(30):12377-12381, 2009.

Wiles, P.J., van Duren, L.A., Hase, C., Larsen, J. and Simpson, J.H.: Stratification and mixing in the Limfjorden in relation to mussel culture. *J. Mar. Syst.*, 60(1-2):129-143, 2006.

615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650






Fig. 1. The study sites in Denmark and Finland. a) study regions, b) Finnish study sites, c) Limfjorden study sites, d) Funen study sites.



651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675

Fig. 2. Silt content (%) (a, c) and degree of sediment sorting (b, d) at the study sites in Finland and Denmark, respectively. Lower values in degree of sorting indicate well-sorted sediment types.


676
677



678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697

Fig. 3. Sediment profiles of organic carbon density (mg C cm^{-3}) in the top 25 cm of the Finnish and Danish eelgrass (*Zostera marina*) meadows. Note the difference in the scale of x-axis between the regions. Numbers below detection limit ($\%OC < 0.01 \%DW$) are not included in the figure. Average (\pm SEM; $n=1-3$).


698
699
700
701
702



703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718

Fig. 4. Corg stocks (g C m⁻²) in the top 25 cm of sediment in Finnish and Danish eelgrass (*Zostera marina*) meadows. Note that the value of Thurøbund (grey bar) corresponds to right y-axis.


719
720
721



722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747

Fig. 5. Relative contribution of different organic matter sources (*Z. marina*, *P. perfoliatus*, *P. pectinatus*, *Ruppia cirrhosa*, phytoplankton and drift algae) to the ^{13}C signal of the sediment surface layer (0-2 cm) in Finnish and Danish eelgrass (*Z. marina*) meadows.

748
 749
 750
 751
 752
 753
 754
 755
 756



757
 758
 759
 760
 761
 762
 763
 764
 765
 766

Fig. 6. Distance-based redundancy analysis (DbRDA) plot showing the environmental parameters (percentage of *Z. marina* in sediment carbon pool, above:belowground-ratio, annual eelgrass production, sediment silt content [%], sediment dry density and sediment porosity) fitted to the variation in the Corg stock (g C m^{-2}) at the Finnish and Danish eelgrass (*Z. marina*) sites, respectively. Vectors indicate direction of the parameters effect. Site codes: Finland; Ko=Kolaviken, Ry=Ryssholmen, Tv=Tvärminne, Få=Fårö, Ly=Lyddaren, Lå=Långören, Hu=Hummelskär, Jä=Jänisholm, Sa=Sackholm, An=Ängsö. Site codes: Denmark; Ag12=Agerø12, Ag3=Agerø3, Vi=Visby, Lg=Løgstør, Lo=Lovns, Th=Thurøbund, Lu=Lunkebugt, Da=Dalby, Ke=Kertinge, Ny=Nyborg.

767 Table 1. Location, silt content (% silt), sediment dry density (Dry dens. g cm⁻³), sediment organic carbon content (SedOC, % DW), sediment organic
 768 matter content (SedOM, %DW), δ¹³ C sediment surface, δ¹³ C *Z. marina* leaves, δ¹³ C *Z. marina* rhizomes, seagrass shoot density (Shoot dens., shoots
 769 m⁻²), seagrass above and belowground biomass (AB and BB, gDW m⁻²), root:shoot-ratio (R:S), and aboveground production (Production, gDW m⁻² y⁻¹)
 770 at the sampling sites. SE (n= 3–4) is given. Annual seagrass production is calculated from pooled values of replicates per site and therefore no SE is
 771 shown.

772
 773
 774

Country/ Location	Silt content (%)	Dry dens. (g cm ⁻³)	SedOM (% DW)	SedOC (% DW)	δ ¹³ C sediment surface	δ ¹³ C <i>Z. marina</i> leaves	δ ¹³ C <i>Z. marina</i> rhizomes	Shoot density (shoots m ⁻²)	AB (gDW m ⁻²)	BB (gDW m ⁻²)	R:S	Production (gDW m ⁻² y ⁻¹)
Finland												
Fårö	5.0±1.4	1.32±0.025	0.66±0.07	0.13±0.001	-20.6±0.3	-9.7±0.4	-9.0±0.20	304±32	138±20	167±28	1.27±0.13	773
Hummelskär	9.0±2.6	1.33±0.009	1.06±0.20	0.35±0.019	-19.4±1.2	-9.3±0.3	-9.8±0.25	364±31	70±11	28±2	0.45±0.06	446
Jänisholm	7.1±2.1	1.37±0.076	0.93±0.20	0.33±0.135	-22.1±0.4	-10.8±0.4	-11.0±0.28	128±17	65±16	46±2	1.44±0.53	270
Kolaviken	1.9±0.2	1.34±0.035	0.75±0.02	0.13±0.011	-19.5±0.2	-10.3±0.3	-11.4±0.34	476±96	74±6	149±16	2.07±0.27	324
Lyddaren	4.9±2.5	1.34±0.171	1.75±0.70	0.45±0.094	-13.5±3.5	-8.8±0.4	-9.6±0.29	228±42	86±7	57±12	0.64±0.09	505
Långören	4.4±2.1	1.42±0.046	2.70±2.10	0.19±0.019	-18.9±0.4	-8.5±0.1	-8.9±0.15	436±53	121±46	68±25	0.58±0.06	788
Ryssholmen	2.7±0.6	1.34±0.054	0.89±0.20	0.16±0.004	-20.7±0.3	-11.5±0.1	-11.5±0.29	756±57	160±3	136±16	0.86±0.11	803
Sackholm	12.4±1.9	1.36±0.042	0.95±0.20	0.26±0.027	-21.1±0.8	-10.3±0.7	-9.9±0.34	774±234	110±18	37±9	0.31±0.04	377
Tvärminne	9.2±1.9	1.33±0.034	0.88±0.20	0.20±0.016	-22.7±0.6	-11.6±0.1	-11.5±0.25	112±11	99±16	38±7	0.37±0.01	436
Ångsö	6.3±0.5	1.36±0.052	0.84±0.02	0.20±0.010	-20.1±0.3	-10.3±0.1	-10.3±0.28	604±98	91±6	63±9	0.67±0.05	521
FIN average	6.3±1	1.35±0.014	1.4±0.3	0.24±0.033	-19.9±0.3	-10.1±0.3	10.3±0.32	417±75	101±3	79±5	0.87±0.06	524
Denmark												
Agero 3	29.4±6.2	1.24±0.085	1.94±0.60	2.30±0.082	-13.0±1.7	-9.2±0.5	-11.1±0.22	448±89	181±33	84±8	0.52±0.07	1075
Agero 12	27.3±7.7	1.35±0.173	1.65±0.80	0.29±0.135	-17.4±0.8	-10.7±0.3	-11.9±0.21	404±90	110±2	46±9	0.40±0.08	576
Dalby	8.1±1.2	1.37±0.034	0.67±0.03	0.12±0.009	-17.3±0.7	-9.7±0.3	-10.5±0.56	400±48	76±7	83±10	1.09±0.11	470
Kertinge	27.1±1.5	1.15±0.045	12.59±1.60	3.23±0.236	-16.6±0.2	-9.2±0.1	-9.8±0.08	328±64	90±17	64±14	0.68±0.02	527
Lovns	17.3±2.7	1.22±0.092	2.90±0.50	1.53±0.088	-16.3±2.4	-11.5±0.4	-12.2±0.37	360±27	141±4	100±11	0.70±0.06	848
Lunkebugt	33.0±7.4	1.23±0.227	4.72±2.40	1.71±0.806	-16.9±0.3	-8.9±0.9	-10.6±0.38	347±81	210±10	382±24	1.82±0.08	1056
Løgstør	4.0±0.4	1.23±0.025	0.75±0.03	0.31±0.089	-17.7±0.4	-9.7±0.4	-10.4±0.51	300±14	149±11	63±13	0.42±0.07	755
Nyborg	0.5±0.3	1.17±0.027	0.42±0.02	0.10±0.006	-17.6±1.1	-9.3±0.2	-10.6±0.34	652±30	203±24	214±50	1.00±0.14	1179
Thurøbund	34.6±2.8	1.27±0.030	14.48±0.80	5.78±0.512	-15.5±0.4	-8.2±0.1	-9.0±0.22	420±98	101±16	398±15	4.54±0.70	619
Visby	21.0±3.6	1.25±0.021	1.17±0.06	2.18±0.201	-13.8±1.2	12.0±0.6	-12.4±0.70	520±21	193±13	49±4	0.25±0.01	2172
DK average	20.2±3.9	1.25±0.022	3.9±1.5	1.75±0.563	-16.20±0.2	-9.8±0.4	-10.9±0.33	418±32	145±5	148±14	1.14±0.13	928

775

776 Table 2. Estimated average carbon stocks (g C m⁻² and Mt C), annual areal carbon accumulation (Corg acc. t C ha⁻¹y⁻¹) and annual carbon accumulation
777 (Annual Corg (Mt C y⁻¹) in Finnish and Danish eelgrass (*Z. marina*) meadows. Denmark_{lost} = eelgrass area of the region lost since the beginning 1900`s.
778 Limfjorden_{lost}= eelgrass area of the region lost since the 1900`s. See text for calculations. *) mean carbon density (mg C cm⁻³) calculated for Denmark
779 is used. n.d= no data. For calculations of annual carbon accumulation three different sediment accumulation rates were applied (0.32 mm y⁻¹;
780 Miyayima et al. 2015, 2.02 mm y⁻¹; Duarte et al. 2013b and 4.2 mm y⁻¹; Serrano et al. 2014), for Corg seq the sediment accumulation rate of 2.02 mm
781 y⁻¹ was used.
782

Region	Seagrass area (km ²)	Carbon density (mg C cm ⁻³)	Corg stock (g C m ⁻²)	Corg stock (Mt C)	Corg acc. (t C ha ⁻¹ y ⁻¹)	Annual Corg accumulation (Mt C y ⁻¹)		
						784		
						0.32 mm y ⁻¹	2.02 mm y ⁻¹	4.2 mm y ⁻¹
Finland	30	2.60±0.09	627±25	0.019±< 0.001	0.052	0.002	0.016	0.0328
Limjorden	18	10.57±1.66	2644±207	0.047± 0.007	0.213	0.006	0.038	0.079
Funen	179	24.32±9.15	6005±1127	1.090±0.410	0.491	0.139	0.881	1.832
Denmark _{min}	673	17.45±9.42*	4324±1188*	2.164±0.005	0.352	0.376	2.373	3.636
Denmark _{max}	1345	17.45±9.42*	4324±1188*	5.868±0.014	0.352	0.75	4.741	9.859
Denmark _{lost}	5381-6230	17.45±9.42*	17.45±9.42*	23.478-27.183	n.d	n.d	n.d	n.d

785

786 Table 3. Table from DistLm analysis showing variables in the marginal tests and the results for
787 statistical analysis.
788

MARGINAL TESTS

Variable	SS	Pseudo-F	P-value	Proportion
1. Root: shoot- ratio	5309	10.64	0.002	0.155
2. Sediment dry density	10704	26.37	0.001	0.313
3. Annual eelgrass production	4959	9.82	0.002	0.145
4. Shoot density	48	0.08	0.911	0.001
5. Porosity	3507	6.61	0.010	0.102
6. Silt content (%)	12653	33.99	0.001	0.369
7. C:N-ratio of plant material	464	0.79	0.397	0.014
8. <i>Z. marina</i> content (%)	12179	32.02	0.001	0.356
9. Degree of sorting	9725	23.01	0.001	0.284

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808 Table 4. Table from DistLm analysis showing results from the sequential tests and solution given
 809 by the analysis.

810

Variable	AICc	Sum of squares	Pseudo-F	P- value	Proportion	Cumulative proportion	Degrees of freedom
Silt content (%)	357.4	12653	33.9	0.001	0.369	0.369	58
Root: shoot-ratio	346.0	4375	14.5	0.001	0.127	0.497	57
<i>Z. marina</i> content (%)	333.6	3745	15.6	0.001	0.109	0.606	56
Production	332.2	805	3.5	0.037	0.023	0.630	55
Sediment dry density	331.3	700	3.2	0.049	0.020	0.650	54
Porosity	330.8	602	2.8	0.056	0.017	0.668	53
BEST SOLUTION	AICc	R ²	RSS	Variables	Selections		
	330.8	0.668	11363	6	1-3;5;6;8		

811

812

813

814

815

816

817

818