

Interactive comment on “The status and challenge of global fire modelling” by S. Hantson et al.

E. Chuvieco (Referee)

emilio.chuvieco@uah.es

Received and published: 21 March 2016

General evaluation The paper presents a framework for evaluation of fire models, particularly in the context of analyzing the potential impacts of climate changes in altering fire regimes. The paper is well written and generally well documented (few minor issues are commented later). It does provide a good overview of existing global DGVM including a fire component, with tables summarizing the main assumptions and drivers. I think the paper would benefit from extending the specific benchmark test that will be used to compare the model performance, as the current version only gives insights on potential approaches. I believe the authors should be more specific on what are the planning to do to actually compare model performance, which indicators will they use, on which period and area (including target resolution).

Minor issues: Line 65: a comma is missed after seasonality: “frequency, intensity, seasonality etc”. Line 71: What a significant fraction means? Please quantify Lines 70-

[Printer-friendly version](#)

[Discussion paper](#)



Interactive
comment

86: No reference to N2O emissions from fires is made. Why? Line 92: Johnston et al., 2012 is missed in the references section. Lines 108-109: "Fire risk is not quantitatively related to area burnt, fuel consumption, or fire emissions". It is not clear what you mean here. Most fire risk systems are assessed with fire statistics (Chuvieco et al. 2014; Chuvieco et al. 2010; Padilla and Vega-Garcia 2011; Paz et al. 2011), and some are associated to burned area and fuel consumption (Consume, for instance: see Pettinari and Chuvieco 2015). Line 143: "outcrops can act as natural barriers to fire fronts". Natural barriers is duplicated from previous line. Line 147: "and highest in areas of intermediate water availability", assuming a dry period exists. Line 159: "purpose, for example for forest clearance, agricultural waste burning or fire", please add pasture management, which is the most common factor in many areas of the world. Line 170: "gross domestic product, GDP, that are linked to population density) results from the co-variance of population density with vegetation production and moisture". This sentence may be tinged, as those relations depend on other factors, such as the importance of agricultural sector in regional economy. For a global analysis, you may be interested to read Chuvieco and Justice 2010. Bowman et al. 2011 has also an interesting analysis of human-fire relations. Line 198: the JSBACH acronym is not defined. Lines 368: When citing alternative sources of model assessment, you do not include reference to the GFED dataset (Giglio et al. 2013), which is widely used for fire –emissions analysis. A reference to the synthesis analysis of Mouillot et al (2014) may be relevant in this point. Please, also note that soil moisture is not equivalent to fuel moisture. The CCI Soil moisture product does not really estimate vegetation wetness. Lines 383-388: When analyzing different global burned area products, you may refer to the intercomparison analysis published by Chang and Song 2009 or the most recent validation effort by Padilla et al. 2015. Line 381 and 814: Please note that ESA MERIS burned area product is officially named Fire_cci (see Chuvieco et al. 2016, which also includes an assessment of fire emissions derived from this product). The temporal resolution of the Fire_cci product is Burn Date for the pixel product at approx. 300 m resolution. However, the burned area is accumulated in 15 day periods for a gridded

[Printer-friendly version](#)[Discussion paper](#)

version of product, which has 0.5 d resolution. Line 459: please, include the updated reference to Alonso-Canas and Chuvieco 2015. Line 819. In fig. 1 you may add to Fuel load, Fuel continuity, which is related to fragmentation.

References Alonso-Canas, I., & Chuvieco, E. (2015). Global Burned Area Mapping from ENVISAT-MERIS data. *Remote Sensing of Environment*, 163, 140-152. Bowman, D.M.J.S., Balch, J., Artaxo, P., Bond, W.J., Cochrane, M.A., D'Antonio, C.M., DeFries, R., Johnston, F.H., Keeley, J.E., Krawchuk, M.A., Kull, C.A., Mack, M., Moritz, M.A., Pyne, S., Roos, C.I., Scott, A.C., Sodhi, N.S., & Swetnam, T.W. (2011). The human dimension of fire regimes on Earth. *Journal of Biogeography*, 38, 2223-2236. Chang, D., & Song, Y. (2009). Comparison of L3JRC and MODIS global burned area products from 2000 to 2007. *Journal of Geophysical Research*, 114, 10.1029/2008JD11361. Chuvieco, E., Aguado, I., Jurdao, S., Pettinari, M.L., Yebra, M., Salas, J., Hantson, S., de la Riva, J., Ibarra, P., Rodrigues, M., Echeverría, M., Azqueta, D., Román, M.V., Bastarrika, A., Martínez, S., Recondo, C., Zapico, E., & Martínez-Vega, F.J. (2014). Integrating geospatial information into fire risk assessment. *International Journal of Wildland Fire*, 23, 606-619. Chuvieco, E., Aguado, I., Yebra, M., Nieto, H., Salas, J., Martín, P., Vilar, L., Martínez, J., Martín, S., Ibarra, P., de la Riva, J., Baeza, J., Rodríguez, F., Molina, J.R., Herrera, M.A., & Zamora, R. (2010). Development of a framework for fire risk assessment using remote sensing and geographic information system technologies. *Ecological Modelling*, 221, 46-58. Chuvieco, E., & Justice, C. (2010). Relations between Human Factors and Global Fire Activity. In E. Chuvieco, J. Li & X. Yang (Eds.), *Advances in earth observation of global change* (pp. 187-200). Dordrecht ; London: Springer. Chuvieco, E., Yue, C., Heil, A., Mouillot, F., Alonso-Canas, I., Padilla, M., Pereira, J.M., Oom, D., & Tansey, K. (2016). A new global burned area product for climate assessment of fire impacts. *Global Ecology and Biogeography*, DOI 10.1111/geb.12440. Giglio, L., Randerson, J.T., & Werf, G.R. (2013). Analysis of daily, monthly, and annual burned area using the fourth generation global fire emissions database (GFED4). *Journal of Geophysical Research: Biogeosciences*, 118, 317-328. Mouillot, F., Schultz, M.G., Yue, C., Cadule, P., Tansey, K., Ciais, P.,

Interactive
comment

& Chuvieco, E. (2014). Ten years of global burned area products from spaceborne remote sensing—A review: Analysis of user needs and recommendations for future developments. *International Journal of Applied Earth Observation and Geoinformation*, 26, 64-79. Padilla, M., Stehman, S.V., Hantson, S., Oliva, P., Alonso-Canas, I., Bradley, A., Tansey, K., Mota, B., Pereira, J.M., & Chuvieco, E. (2015). Comparing the Accuracies of Remote Sensing Global Burned Area Products using Stratified Random Sampling and Estimation. *Remote Sensing of Environment*, 160, 114-121. Padilla, M., & Vega-Garcia, C. (2011). On the comparative importance of fire danger rating indices and their integration with spatial and temporal variables for predicting daily human-caused fire occurrences in Spain. *International Journal of Wildland Fire*, 20, 1-13. Paz, S., Carmel, Y., Jahshan, F., & Shoshany, M. (2011). Post-fire analysis of pre-fire mapping of fire-risk: A recent case study from Mt. Carmel (Israel). *Forest Ecology and Management*, 262, 1184-1188. Pettinari, M.L., & Chuvieco, E. (2015). Generation of a global fuel dataset using the Fuel Characteristic Classification System. *Biogeosciences Discuss.*, 12, 17245-17284.

Interactive comment on *Biogeosciences Discuss.*, doi:10.5194/bg-2016-17, 2016.

[Printer-friendly version](#)

[Discussion paper](#)

