

We thank both reviewers for their comprehensive assessment of our paper.

Reviewer 1 (R1)

R1 contended that the ms provided very interesting data from the paired flux tower sites but suggested we did "... not emphasise the strengths of their work, which is the detailed time series that can compare processes for the two sites. Rather they attempt to extrapolate the results across northern Australia in ways that are not transparent and appear to have a number of flaws"

This was a reasonable criticism but this ms is part of the OzFlux Special Issue and here we wanted to highlight the use of flux observations in refining GHG emissions and impacts on national accounts, a management focus rather than a detailed physiological analysis.

The flux observations were used to highlight respiration differences between sites, the magnitude of soil CO₂ emissions during tillage and preparation for cropping, the continued uptake of carbon post-deforestation from grass growth and woody re-sprouting, as well as the net loss of the natural C sink that we observed at the uncleared analogue site.

Other papers in the Special Issue have documented the flux characteristics of north Australian savannas – data from four savanna flux towers were included in the overview paper of Beringer et al. 2016 (doi:10.5194/bg-2016-152).

R1 provided very useful comments aimed at improved descriptions and estimation of fuels and associated emissions from the debris burning post deforestation, an important component of the total emission from this LUC. As a result the Methods section has been re-written and restructured and broken into 8 sections instead of 6. The revised ms also features a new Table, Table 3, that describes in a more transparent manner, the distribution of fuels as measured across the deforested site using fuel classifications as defined by the Australian Government's savanna abatement methodology and latest emissions factors. Table 3 includes data for each fuel type (fine, coarse, heavy) including fuel mass and the associated emissions factors for each GHG (CO₂ CH₄ and N₂O), N:C and %C content with the emissions calculation described by an equation, Equation 1.

The emission calculation is now based on emission factors as recommended by R1, namely those of Meyer et al. 2012 and the Savanna Abatement Methodology Determination, March 2015, a methodology that is now legislated by the Commonwealth government (www.legislation.gov.au/Details/F2015L00344). This is a more robust and transparent reporting of emissions from the debris fire, although the new estimate (121.9 Mg CO₂-e ha⁻¹) differs by only 5% (now smaller) when compared to the original estimate. The new estimate has a higher contribution from non-CO₂ fluxes.

These improvements are in response to R1's comments arising from text on P7 to P12 of the original ms and we believe this has improved the clarity of the Methods section.

We suspect R1 (and R2) assumed we had one or two large stockpiles of heavy fuels that burnt very hot for 10 days or more. This was not the case as burning of the site consisted of ignition of the cured grasses and woody fuels in situ, with no stock-piling as an initial phase of burning. To ensure safety, ignition was done in blocks, which is why the process took 20+ days. After an initial burn, unburnt woody debris was then stockpiled and re-burnt until incinerated in a second phase. As such we had multiple debris piles distributed across the 295 ha block as opposed to one or two large piles burning at very high temperatures. Again, this has been described more fully in the revised text and to aid in the description of the LUC phases, in particular the fire event and aftermath, a colour Plate, Plate 1, has now been included.

Plate 1 consists of 4 images of the site showing the initial deforestation event, the debris fire and stockpiling and the finalised state of the site prior to bed preparation for cropping.

Descriptions of the data sources for savanna-specific deforestation emissions across north Australia have been improved as R1 (and R2) found this hard to follow. All data were sourced from the NGGI in collaboration with staff from the Commonwealth government's departmental reporting team, as was acknowledged in the Acknowledgements. References to the methodology are given. In short, the savanna boundary that was defined by Fox et al. 2001 was applied as a spatial 'mask' to constrain the area of emissions estimates to the savanna region only. These were then compared to emissions from savanna burning from the same area.

R1 queried how the value of 78, 605 ha y⁻¹ in Table 4 (now Table 5) was derived. It is described in text in the Methods section, in Table 4 (5)'s caption and footnote for the table - this value is derived from the savanna constrained deforestation area and is the mean savanna area defrosted per year 1990-2013.

R1 also queried our LUC emission figure for 2013 and suggested our reported value was low. Firstly we report the mean from 1990 to 2013 and secondly we are not reporting emissions from all activities with the LUC sector – only emissions from Activity 2 'Deforestation' are relevant to our study. This was indicated in the original text. Plus our emissions estimate is specifically limited to the savanna land area across WA, NT and Queensland. If R1 looked up reportable emissions for these states in their entirety, there would be a significant difference compared to our reported value as the standard data reported by the Commonwealth includes the non-savanna (non-tropical) areas of each state where there have been significantly higher deforestation rates. Comparing with Cook et al 2010 may be problematic as the area included in each study would need to be identical, especially areas of Queensland, which have experienced significant clearances in southern and central Queensland, outside of our study area. It should also be noted that for the regional savanna estimates, we are simply compiling emissions data for either savanna burning or deforestation as estimated by the Commonwealth, but constrained to the savanna area as defined by Fox et al.

There were a number of other inconsistencies R1 commented on and these have all been addressed: fire frequency data were inconsistent and have been corrected; the citation for biomass allometry is confined to Williams et al. 2005; reference to fire-line intensities has been deleted given our fuel load is a mixture of grass and heavy fuel, with heavy fuel dominating.

We have also improved text in the Methods describing CWD estimation, in particular dealing with hollowing of large CWD fragments – we do not 'add missing biomass' as R1 queried, our

method is designed to estimate the missing *volume* to ensure we do not provide a large overestimate of CWD. This was not entirely clear in the original ms and the text is now improved. We are estimating the volume for each CWD fragment that is then converted to biomass using specific wood densities assigned to our 5 rot classes that we define.

R1 also commented on our text re stand replacement events such as cyclones and/or floods which would take 4 decades to recover the lost carbon. The original text was confusing as given the site locations, neither of these events / scenarios is feasible and this sentence has been modified accordingly. The only agent of stand replacement in the region of our sites would be deforestation and conversion for agricultural production.

We include both CO₂ (not reportable) and non-CO₂ emissions (reportable) for savanna burning for comparison with deforestation emissions.

Reviewer 2 (R2)

Comments by R2 related to improvements in expression and typographical errors throughout the ms as well as an inclusion of a statement of potential errors.

All suggested changes of R2 have been implemented.

Fig 2 on energy balance closure was removed as suggested by R2 and text describing slope statistics from the closure analysis has now been incorporated included in the revised text.

R2 queried the nature of the gap filling approach used – a unique ANN model was developed for each LUC phase given the significant change in canopy and microclimatic characteristics of each phase. Text describing this has been improved. Errors associated with gap filling using the DINGO system were minimal as we had less than 10% of data that was missing.

In this study we used 30 minute covariance data for the calculation of fluxes not the raw data as is inferred from the paper of Isaac et al. in the Special Issue.

We can confirm fire emissions were not included in the NEE measurements and the value of 0.9 (BEF) was derived from an assessment of remaining heavy fuel levels post-fire. This value influences calculations of both CO₂ and non-CO₂ as described in the new methods section and the new Table 5 that gives emissions factors.

R2 queried the value of 2.75 Mg ha⁻¹ – this is the combined emission from the soil tillage phases over the last 6 months of the measurements, as is described in the text.

As requested by R2, a statement on potential errors associated with our emissions estimate from the debris fire has been included, which is based on uncertainty measures as described Russell-Smith et al. (2009) for key parameters used in the Australian savanna emissions methodology. Given a number of key parameters were measured on site in this study, with fuels measured across a well-defined area, our errors will be relatively low when compared with catchment scale to regional scale projects that the methodology has been designed for.

As R2 suggested, we statistically tested for site differences for each LUC phase (1-way ANOVA) with all phases significantly different except the pre-clearing phase, Phase 1. Significantly different mean NEE are identified in Table 3.

Figure 1 has been improved as requested, with latitude and longitude lines marked and a higher resolution coastline used.

We thank both reviews for their very constructive comments on the paper.

**Quantifying the relative importance of greenhouse gas emissions from current and future
savanna land use change across northern Australia**

Bristow M.^{1,2}, Hutley L.B.¹, Beringer J.³, Livesley S.J.⁴, Edwards A.C.¹, Arndt S.K.⁴

¹School of Environment, Research Institute for the Environment and Livelihoods, Charles Darwin
University, NT, Australia, 0909

²Department of Primary Industry and Fisheries, Berrimah, NT, Australia, 0828

³School of Earth and Environment, The University of Western Australia, Crawley, WA, Australia,
6009

⁴School of Ecosystem and Forest Sciences, The University of Melbourne, [Burnley](#), Victoria,
Australia, 3121

Correspondence to: Lindsay B. Hutley (lindsay.hutley@cdu.edu.au)

Keywords: Deforestation, clearing, eddy covariance, savanna burning, fire emissions

1 Abstract

2 Clearing and burning of tropical savanna leads to globally significant emissions of greenhouse
3 gases (GHG) ~~although~~, however there is large uncertainty relating to the magnitude of this flux.
4 Australia's tropical savannas occupy ~~over 25% the northern quarter~~ of the ~~continental continent~~, a
5 region of increasing interest for further exploitation of land mass and water resources. Land use
6 decisions across this vast biome have ~~at~~ the potential to ~~significantly~~ influence the national
7 greenhouse gas budget, ~~particularly because they are the focus of likely agricultural expansion~~. To
8 better quantify emissions from savanna deforestation and investigate the ~~role~~ impact of deforestation
9 on national GHG emissions, we undertook a paired site ~~approach was used~~. ~~The CO₂ exchange was~~
10 ~~measured~~ measurement campaign where emissions were quantified from two tropical savanna
11 woodland sites; one that was ~~cleared~~ deforested and prepared for agricultural land use, and a second
12 analogue site that remained uncleared for the duration of a 22 month ~~observation period campaign~~.
13 At both sites, net ecosystem exchange (~~NEE~~) of CO₂ was measured using the eddy covariance (~~EC~~)
14 method. Observations at the ~~cleared~~ deforested site ~~was were~~ continuous before, during and after the
15 clearing event, providing high resolution data that tracked CO₂ emissions through multiple ~~line~~
16 phases of land use change. At the deforested site, post-clearing debris was allowed to cure for six
17 months and was subsequently burnt, followed by extensive soil preparation for cropping.
18 ~~At~~ During the ~~cleared site, post-clearing debris was allowed to cure for 6 months and was~~
19 ~~subsequently burnt, followed by extensive soil preparation for cropping~~. ~~Emissions burning, fluxes~~
20 of CO₂ as measured by the eddy covariance tower were excluded. For this phase, emissions were
21 estimated ~~from the debris fire~~ by quantifying ~~the~~ on-site biomass prior to ~~clearing~~ deforestation and
22 applying savanna-specific ~~emission~~ emission factors to estimate a fire-derived GHG emission. ~~This~~
23 ~~was added to net CO₂ fluxes as measured by the eddy covariance tower, giving a total GHG~~
24 ~~emission of 154 Mg CO₂-e ha⁻¹ from a savanna woodland with a that included both CO₂ and non-~~
25 CO₂ gases. ~~The~~ total fuel ~~load~~ (mass that was consumed during the debris burning was 40.9 Mg C

1 ha⁻¹ and included above- and below- ground woody ~~debris~~biomass, coarse woody debris, twigs, leaf
2 litter plusand C₄ grass ~~fuel~~) of 40.9 Mg C ha⁻¹. ~~This~~fuels. Emissions from the burning were added to
3 the net CO₂ fluxes as measured by the eddy covariance tower for other post-deforestation phases to
4 provide a total GHG emission ~~was dominated by the combustion of cleared debris which was 83%~~
5 of the ~~from this land use change.~~

6 The total emission from this savanna woodland was 148.3 Mg CO₂-e ha⁻¹ with the ~~remainder~~
7 ~~coming~~debris burning responsible for 121.9 Mg CO₂-e ha⁻¹ or 82% of the total emission. The
8 remaining emission was attributed to CO₂ efflux from soil ~~emissions~~disturbance during site
9 preparation for agriculture (10% of the total emission) and decay of debris during the curing period
10 prior to burning. ~~Soil disturbance from ploughing and site preparation for cropping was responsible~~
11 ~~for almost 10% of the total emission. Fluxes (8%).~~ Over the same period, fluxes at the uncleared
12 savanna woodland site were ~~tracked~~measured using ~~an additional a second~~ flux tower ~~for~~and over
13 the 22 month observation period ~~and over this time the~~ cumulative NEE was ~~-2.1 Mg C ha⁻¹~~, a net
14 carbon sink of -2.1 Mg C ha⁻¹, or -7.7 Mg CO₂-e ha⁻¹.

15 Estimated emissions for this savanna type were then ~~upsealed~~extrapolated to a regional scale to
16 1) provide estimates of the magnitude of GHG emissions from any future deforestation. ~~At current~~
17 ~~rates of deforestation, savanna burning is as significant a source of GHG emissions as deforestation,~~
18 ~~with fire emissions occurring and 2) compare with GHG emissions from prescribed savanna~~
19 burning that occurs across north Australian savanna every year ~~across this.~~ Emissions from current
20 rate of annual savanna ~~biome. However, expanded deforestation could exceed fire emissions and a~~
21 ~~clearing scenario was~~ across north Australia was double that of reportable (non-CO₂ only) savanna
22 burning. However, if the total GHG emission is accounted, CO₂ plus non-CO₂ emissions, burning
23 emissions are an order of magnitude larger than that arising from savanna deforestation. We
24 examined ~~which suggested that clearing a scenario of expanded land use that required additional~~
25 deforestation of savanna woodlands over and above current rates. This analysis suggested that

1 significant expansion of deforestation area across the northern savanna woodlands could add ~~up to~~
2 ~~5~~an additional 3% to Australia's national GHG account for the duration of the ~~clearing~~
3 ~~activities~~land use change. This bottom-up study provides data that can reduce uncertainty associated
4 with land use change for this extensive tropical ecosystem and provide an assessment of the relative
5 magnitude of GHG emissions from savanna burning and deforestation~~as well as~~. Such knowledge
6 can contribute to informing ~~northern~~ land use decision making processes associated with land and
7 water resource development.

8

1.0 Introduction

An increase in greenhouse gas (GHG) emissions through human-related activities is leading to rapid change in the climate system (IPCC 2013) ~~and it~~. It is therefore crucial to obtain data describing the net GHG balance at regional to global scales to better characterise anthropogenic forcing of the atmosphere (Tubiello et al., 2015). Emissions from land-use change (LUC) are the integral of ecosystem transformations that can include emissions from deforestation and conversion to agriculture, logging and harvest activity, shifting cultivation, as well as regrowth sinks following harvest and/or abandonment of previously cleared agriculture lands (Houghton et al. 2012). At present, LUC emits $0.9 \pm 0.5 \text{ Pg C y}^{-1}$ to the atmosphere, which is approximately 10% of anthropogenic carbon emissions (Le Quéré et al., 2014). Data sources and methods used to estimate LUC emissions are diverse. These include census-based historical land use reconstructions and land use statistics, satellite estimates of biomass change through time ~~(Baccini et al., 2012)~~(Baccini et al., 2012), satellite monitored fire activity and burn area estimates associated with deforestation ~~(van der Werf et al., 2010)~~(van der Werf et al., 2010). In addition, there is increasing use of ecosystem models coupled with remote sensing to estimate emissions from LUC (Galford et al. 2011).

Emissions associated with the LUC sector have the highest degree of uncertainty given the complexity of processes involving net emissions and Houghton et al. (2012) assessed this uncertainty at $\sim 0.5 \text{ Pg C y}^{-1}$, which is of the same order of magnitude as the emissions themselves. Uncertainties in estimating GHG emissions arising from savanna clearing, associated debris burning and conversion to agriculture are greater than those for tropical forests (Fearnside et al., 2009). It is important to quantify the emissions and their uncertainties in savannas particularly because tropical savanna woodland and grasslands occupy a large area globally ($27.6 \text{ million km}^2$), greater than tropical forest ($17.5 \text{ million km}^2$, Grace et al., 2006). Deforestation and associated fire from these biomes are the largest contributors to global LUC emissions (Le Quéré et al., 2014). Much of these

GHG emissions are from the Brazilian Amazonia, an agricultural area that has been expanding since the 1990s. However, over the last decade, the rate of tropical forest deforestation in this region has decreased from 16,000 km² in early 2000s to ~6,500 km² by 2010 (Grace et al., 2006)(Lapola et al., 2014), but at the expense of the Brazilian cerrado, a vast savanna biome of some 2.04 million km², where clearing rates have been maintained (Ferreira et al., 2013, 2016; Galford et al., 2013). Given the suitability of the cerrado topography and soils for mechanized agriculture, the Cerrado may become the principal region of LUC in Brazil (Lapola et al., 2014).

~~Tropical savanna woodlands and grasslands occupy a large area globally (27.6 million km²), greater than tropical forest (17.5 million km²; Grace et al., 2006). Deforestation and associated fire from these biomes are the largest contributors to global LUC emissions (Le Quéré et al., 2014). Much of these GHG emissions are from the Brazilian Amazonia, an agricultural area that has been expanding since the 1990s. However, over the last decade, the rate of tropical forest deforestation in this region has decreased from 16,000 km² in early 2000s to ~6,500 km² by 2010 (Lapola et al., 2014), but at the expense of the Brazilian cerrado, a vast savanna biome of some 2.04 million km², where clearing rates have been maintained (Ferreira et al., 2013, 2016; Galford et al., 2013). Given the suitability of the cerrado topography and soils for mechanized agriculture as well as potential leakage pressure resulting from declining deforestation rates of tropical forests in Amazonia, the Cerrado may become the principal region of LUC in Brazil (Lapola et al., 2014).~~

North Australia is one of the world's major tropical savanna regions, extending some 1.93 million km² across north-west Western Australia, the northern half of the Northern Territory and Queensland (Fisher and Edwards, 2015). This biome occupies approximately one ~~third~~quarter of the Australian continent and since European arrival, 5% has been cleared for improved pasture, horticulture and cropping (Landsberg et al. 2011), making it one of least disturbed savanna regions in the world (Woinarski et al., 2007). However, this small percentage equates to a substantial area of 9.2 million hectares and LUC and associated economic development in northern Australia is a

government imperative and this is likely to involve expansion and intensification of grazing, irrigated cropping, horticulture and forestry (~~Northern Australia Committee, 2014~~)(Committee on Northern Australia, 2014). Drivers of this potential expansion in food and fibre production include the exploitation of growing markets of Asia as well as domestic factors such as the perception that land and water resources of north Australia can provide a future agricultural resource base to offset the expected declines in agricultural productivity in southern Australia due to adverse impacts of climate change (Steffan and Hughes, 2013).

Historically, intensive agricultural developments in northern Australia have been implemented based on limited scientific knowledge with dysfunctional policy and market settings, and as a result there has been limited success (~~Cook, 2009~~)(Cook, 2009). Future expansion needs to be underpinned by sound understanding of the consequences of regional scale land transformation on carbon and water budgets and GHG emissions. Any significant expansion ~~northern agricultural production would require significant clearance of native savanna vegetation, with unknown increases in GHG emissions. Accurate and transparent measurement of sinks and sources of GHGs has become an imperative to quantify impacts of LUC, in particular clearing and managed savanna burning, on national GHG accounts (Meyer et al., 2012).~~

~~Most LUC studies occur at catchment, regional or biome scales (Houghton et al. in northern agricultural production would require clearance of native savanna vegetation, with unknown increases in GHG emissions. Most LUC studies occur at catchment, regional or biome scales (Houghton et al., (2012) and are not underpinned by good understanding of underlying processes.2012) and are not underpinned by good understanding of underlying processes.~~ However, there are an increasing number of plot-scale studies using eddy covariance and chamber methods to provide direct measures of net GHG fluxes from contrasting land uses (Lambin et al., 2013). These studies typically compare microclimate and fluxes of GHGs from pastures and/or crops with adjacent forest ecosystems under a range of management conditions (e.g. Anthoni et al. 2004; Zona

et al. 2013) or natural grasslands and different cropping types (e.g. Zenone et al., 2011). In tropical regions, there is a focus on transitions from forest to pasture and from forest to crops for food or bioenergy production (Galford et al., 2010, 2011; Wolf et al. 2011, 2011; Sakai et al. 2004, 2004).

~~There are fewer studies that directly measure GHG emissions and sinks prior to, during and after LUC at one site. Land use change can involve rapid changes in net GHG emissions over varying temporal scales (minutes, hours, and seasonal cycles) and continuous flux measurements are essential to capture these events (Hutley et al. 2005). However, there are no direct observations of emissions from savanna clearing in northern Australia, contributing to the uncertainty associated with the LUC sector in Australia's national GHG accounts (Commonwealth of Australia, 2015).~~

There are few studies that directly measure GHG emissions and sinks prior to, during and after LUC at a single site. Land use change can involve rapid changes in net GHG emissions over varying temporal scales (minutes, hours, and seasonal cycles) and continuous flux measurements are essential to capture the magnitude of these events (Hutley et al. 2005). However, there are no direct observations of emissions from savanna clearing in northern Australia, contributing to the uncertainty associated with the LUC sector in Australia's national GHG accounts (Commonwealth of Australia, 2015a).

Our objective is to provide a comprehensive assessment of GHG emissions associated with savanna clearing. Our aims are to 1) quantify the typical rates of CO₂ exchange of intact tropical savanna and make comparative measurements from an analogue site that was to be cleared, 2) quantify CO₂ fluxes before, during and after a clearing event, 3) estimate both CO₂ and non-CO₂ (CH₄ and N₂O) GHG emissions arising from burning of cleared debris and 4) quantify ecosystem scale GHG balance for this land use conversion. and compare with emissions from savanna fire, a significant source of GHG emissions across north Australia.

2.0 Methods

In this study we used a paired site approach, where concurrent fluxes of CO₂, water vapour and energy were measured using eddy covariance towers from an uncleared savanna woodland site and a similar savanna woodland site on the same soil type that was to be cleared, burnt and prepared for agricultural production. Fluxes of CO₂ were monitored for 161 days prior to clearing at both sites with observations continuing during the clearing event (deforestation) and for another 507 days through phases of woody debris and grass curing, burning and soil preparation through raking and ploughing. The entire observation period was 668 days. Flux observations of net CO₂ exchange were combined with on-site biomass measurements and regionally calibrated pyrogenic emissions factors to estimate emissions of ~~CH₄ and N₂O (Russell-Smith et al., 2009b) and CO₂ (Hurst et al., 1994)~~CO₂, CH₄ and N₂O (Meyers et al. 2012, Commonwealth of Australia, 2015b) from burning of the cleared debris- that was a key component of the land conversion. Fire derived emissions were combined with net CO₂ fluxes from the land conversion phases to provide a total net emission in units of CO₂-e for this ~~land conversion~~LUC. In this paper, we use the term deforestation to describe ‘savanna clearing’. Deforestation is defined under Australia’s National Greenhouse Accounting system, as the loss of forest/woodland cover due to direct human-induced actions ~~and fail~~that fails to regenerate cover via natural regrowth or restoration planting (Commonwealth of Australia, ~~2015~~2015a).

2.1 Study sites

Both savanna woodland sites were located within the Douglas-Daly River catchment approximately 300 km south of Darwin, Northern Territory (Fig. 1). Both sites are OzFlux sites (www.ozflux.org.au), with flux observations ongoing at the uncleared (UC) savanna ~~since 2007~~ (Beringer and Hutley, 2016; Beringer et al., 2011; Hutley et al., 2011)site since 2007 (Beringer et al. 2016; Beringer et al., 2011; Hutley et al., 2011). OzFlux is the regional Australian and New Zealand flux tower network that aims to provide continental-scale monitoring of CO₂ fluxes and surface energy balance to assess trends and improve predictions of Australia’s terrestrial biosphere

1 and climate (Beringer et al., 2016). The UC site is broadly representative of Australian tropical
2 savanna woodland found on deep, well drained sandy loam soils at sites with ~1000 mm ~~annual~~
3 ~~rainfall~~MAP (Table 1). The cleared savanna site (CS) ~~cleared~~ was carefully selected to ensure the
4 vegetation and soils were as similar to the UC site as possible, and with topography suitable for
5 eddy covariance measurements.

6 Both sites were classified as savanna woodland ~~type 4B2 using Northern Territory Government~~
7 ~~vegetation mapping ((type 4B2, Aldrick and Robinson 1972~~1972, 1:50,000 mapping);) with an
8 overstorey cover of 30%, equivalent to the 'Eucalypt woodland' Major Vegetation Group (MVG) of
9 the National Vegetation Information System (NVIS, Commonwealth of Australia, 2003). The sites
10 were dominated by an overstorey of *Eucalyptus tetrodonta* (F. Muell.), *Corymbia latifolia* (F.
11 Muell.) ~~and Erythrophleum chlorostachys (F. Muell.) with a mid-storey of small deciduous trees~~
12 ~~and shrubs and an understorey of C4 grasses (Sorghum, Heteropogon spp. dominated).~~). Soils at
13 both the UC and CS sites were red kandosols of the haplic mesotrophic great group (Isbell, 2002),
14 characterised as deep, sandy-loams (Table 1). The long-term mean annual ~~(\pm sd)~~
15 ~~rainfall~~precipitation (MAP) (\pm SD) at the UC site was estimated at 1180 ± 225 mm (1970-2012,
16 Australian Water Availability Project (AWAP), www.csiro.au/awap), similar to the CS site at 1107
17 ± 342 mm (1985-2013, Bureau of Meteorology station, Tindal, NT).

18 ~~Both UC and CS~~ Slopes at both sites ~~had a flat topography (slopes of were~~ $< 2\%$ ~~)~~% with a
19 ~~consistent savanna woodland~~ fetch of ~1.5 km at the UC site, and ~1 km at the CS site. At both
20 sites, 23 m guyed masts were installed to support eddy covariance instruments at 21.5 m ~~at both~~
21 ~~sites~~above-ground. The tower at the CS site was moved three times to ensure adequate fetch was
22 maintained according to seasonal wind direction during clearing and phases of the land use ~~change~~
23 ~~as well as~~conversion. Instrument height was also adjusted given the height of the ~~canopy~~ surface
24 post-clearing and during the soil tillage phase (Table 2). ~~Instruments were mounted at the top of the~~
25 ~~tower during these phases (7 m and 3 m).~~

~~Remotely sensed fire scar history~~ Satellite-derived burnt area mapping is available across north Australia at 250 m resolution (North Australian Fire Information system (NAFI), www.firenorth.org.au) ~~indicating and indicated~~ that fires had occurred within the flux footprint of the UC flux tower in 5 out of the last 13 years (2000-2013), whereas no fires had occurred within the footprint of the CS site. The average fire return time for the entire Australian savanna biome is ~~one in~~ 3.1 years (Beringer et al., 2015).

2.2 Land-use conversion

~~Conversion of woodland to agricultural land in northern Australia involves pulling trees over using a large chains dragged between two bulldozers, followed by the mechanical stockpiling of woody debris to decay and cure prior to burning. This is followed by raking and stock piling of any remaining debris and re-burning. Finally, there was mechanised ripping of soil to remove remaining coarse root material to 60 cm depth. These processes result in the removal of all above ground and most of the below ground biomass, such that the soil was ready for tillage and cultivation. These phases result in a series of events that may lead to short-term, pulsed GHG emissions that would otherwise be missed or greatly under-estimated by episodic measurements taken at a weekly or monthly frequency after an initial tree felling event (Neill et al., 2006; Weitz et al., 1998). The CS site was cleared between 2 and 6 March 2012, which is towards the end of the wet season; 737 mm of rainfall had fallen since the end of the preceding dry season. Over this five day period, 295 ha of savanna were cleared. The specific sequence and timing of clearing, burning and land preparation phases is given in Table 2.~~

2.3 Flux measurements and data processing

Eddy covariance systems at both sites consisted of CSAT3 3-D ultrasonic anemometers (Campbell Scientific Inc., ~~model CSAT3~~ Logan, USA) and a LI-7500 open-path CO₂ / H₂O analysers (Licor Inc., Lincoln, USA). Flux variables were sampled at 10 Hz and covariances stored every 30 minutes. The LI-7500 gas analysers were calibrated ~~on an at~~ approximately six

~~monthly~~ interval for the duration of the data collection period and were highly stable. ~~Daily~~ Mean daily rainfall, air temperature, relative humidity, soil heat flux (F_g , $W\ m^{-2}$) and volumetric soil moisture (θ_v , $m^3\ m^{-3}$) from surface to 2.5 m depths were measured at both sites. The radiation balance was measured using a CNR4 net radiometer (~~Rn~~ F_n , $W\ m^{-2}$) (~~model CNR4~~, Kipp and Zonen, Zurich).

Thirty minute covariances were stored using data loggers (CR3000, Campbell Scientific, Logan) and data post processing and quality control was undertaken using the OzFluxQC system as described by Isaac et al. ~~(2016)~~.(2016). In this system, data are processed through three levels; Level 1 is the raw data as collected by the data logger, Level 2 are quality-controlled data and Level 3 are post processed and corrected but not gap-filled data. Quality control measures at Level 2 include checks for plausible value ranges, spike detection and removal, manual exclusion of date and time ranges and diagnostic checks for all quantities involved in the calculations to correct the fluxes. Quality checks make use of the diagnostic information provided by the sonic anemometer and the infra-red gas analyser. Level 3 post processing includes 2-dimensional coordinate rotation, low- and high-pass frequency correction, conversion of virtual heat flux to sensible heat flux (F_h , $W\ m^{-2}$) and application of the WPL correction to the latent heat (F_e , $W\ m^{-2}$ and CO_2 fluxes (F_c) ~~(Isaac et al., 2016)~~(Isaac et al., 2016). Level 3 data also include the correction of the ground heat flux for storage in the layer above the heat flux plates (Mayocchi and Bristow, 1995).

Gap filling of meteorology and fluxes along with flux partitioning of net ecosystem exchange (NEE) into gross primary productivity (GPP) and ecosystem respiration (R_e) was performed on the Level 3 data using the Dynamic INtegrated Gap filling and partitioning for Ozflux (DINGO) system as described by ~~Beringer et al., (2016)~~.Beringer et al., (2016b). In summary, DINGO gap fills meteorological variables (air temperature, specific humidity, wind speed and barometric pressure) using nearby Bureau of Meteorology ~~(BoM, www.bom.gov.au)~~ automatic weather stations that were correlated with tower observations. All radiation streams were gap-filled

using a combination of MODIS albedo products (MOD09A1) and ~~Bureau of Meteorology~~ BoM gridded global solar radiation and gridded daily meteorology from the Australian Water Availability Project (AWAP) data set (Jones et al. 2009). Precipitation was gap-filled using either nearby ~~Bureau of Meteorology~~ BoM stations or ~~BoM~~ AWAP data. Soil temperature and moisture were filled using the BIOS2 land surface model (Haverd et al., ~~2013a~~ 2013) run for each site and forced with BoM or AWAP data. Energy balance closure was examined using standard plots of $(F_h + F_e)$ vs $(F_n - F_g)$ using 30 minute flux data. ~~Day time and night time data were~~ from both sites (data not shown). For the CS site, closure was examined using data grouped according to ~~each of the seven~~ nine LUC ~~phase periods, plus a further plot using data pooled across all~~ phases (Fig. as given in Table 2). ~~Data from.~~ For the UC site is also plotted using, all 30 minute data ~~for~~ from 2007-2015 was used.

Gap filling of fluxes was undertaken using DINGO that uses an Artificial Neural Network (ANN) model following Beringer et al. ~~(2007)~~ (2007). Model training uses gradient information in a truncated Newton algorithm. NEE and fluxes of sensible, latent and ground heat fluxes were modelled using the ANN with incoming solar radiation, VPD, soil moisture content, soil temperature, wind speed and MODIS EVI as inputs. The u_{star} threshold for each site was determined following Reichstein *et al.* (2005) and night time observations below the u_{star} threshold were replaced with ANN modelled values of R_e using soil moisture content, soil temperature, air temperature and MODIS EVI as inputs. The ANN R_e model was then applied to daylight periods to estimate daytime respiration and GPP was calculated as the difference between NEE and R_e . For data collected at the CS site, a unique ANN model was developed for each LUC phase given the differing canopy and microclimatology of each phase. At each site, daily NEE, R_e and GPP were calculated for each day of each phase.

2.43 Leaf area index

Canopy leaf area index (LAI) at the CS site in the surrounding intact savanna was measured using a 180° hemispherical lens (Nikon 10.5 mm, f/2.8) after ~~Macfarlane~~ Macarlane et al. (2007).

Three savanna transects were photographed seasonally on 9 occasions over 2.1 years from the pre-clearing phase (October 2011) to December 2013. Along each 100 m transect, 11 hemispherical pictures were taken at 10 m intervals (33 photos for each measure occasion). At both sites the LAI was also estimated using MODIS Collection 5 LAI (MOD15A2) for a 1 km pixel around each tower. The 8-day product was interpolated to daily time series using a spline fit. Only MODIS values with a quality flag of 0 for FparLai_QC were used in the estimate indicating the main algorithm was used (lpdaac.usgs.gov/sites/default/files/public/modis/docs/MODIS-LAI-FPAR-User-Guide.pdf).

2.4 Land use conversion

The specific sequence and timing of clearing, burning and land preparation phases are given in Table 2. Conversion of woodland to agricultural land in northern Australia is typically achieved by pulling trees over using large chains held under tension between two bulldozers. Clearing occurs at the end of the wet season when soil moisture is still high and soil strength low as under these conditions trees are easily pulled over, with a large fraction of the tree root mass extracted when pulled. At the CS site, 295 ha of savanna were deforested between 2 and 6 March 2012 using this technique. A permit for this land conversion had been issued by the regional land management agency following an impact assessment and erosion control planning. Chains were under tension and intercepted tree boles 0.1- 0.2 m height above the ground which assisted in pulling the trees and limited damage to the soil surface. As a result, grasses, woody re-sprouts and shrubs of the understorey remained largely intact following deforestation (Plate 1a). Mechanised ripping of soil to 60 cm depth was also undertaken to remove remaining coarse root material.

A cost-effective method of removing cleared vegetation is curing (drying) and subsequent burning and the land managers at the CS site left debris onsite to for 5 Emissions months through the dry season (March to August, 2011). Burning of debris occurred over a 22 day period in the late dry season, August 2012 (Plate 1b), a period of consistent southerly trade winds of low relative

humidity (10-20%, BoM, Tindal station, NT). Prior to ignition, 100 m fire breaks were installed around the entire 295 ha block and then lit in blocks of ~80 ha in size. There was an initial ignition of the fine and coarse fuels (grasses, litter and twigs, defined below) and woody debris (heavy fuels). Heavy fuels that were not completely consumed following the initial burn were then stockpiled in rows ~1-2 m in height and re-ignited until the fuel was consumed (Plate 1c). Inspection of debris post fire suggested ~5% of fine fuels remained as ash and ~10% of the heavy fuels remained as charcoal, which were subsequently incorporated into the top soil on during soil bed preparation (Plate 1d).

2.5 GHG emissions from debris burning

Emissions of CO₂, CH₄ and N₂O from the debris burning were estimated following the approach as outlined in the IPCC Good Practice Guidelines (IPCC 2003), which uses country or region specific emission factors for fire activity (as indicted by burnt area) and the mass of fuel pyrolised to estimate the emission of each trace gas. This approach is well developed for north Australian savanna (Murphy et al., 2015a) for a range of fuel types (grasses, fine and coarse woody fuels, Russell-Smith et al. 2009). Fuel load (biomass) estimates are essential to this approach and was quantified for four fuel types prior to clearing: 1) above-ground woody biomass, 2) below-ground biomass, 3) surface coarse woody debris (CWD), and 4) grass biomass. Emissions estimates were based on fuel mass per area for each fuel type, carbon content (%), elemental C:N ratios and Australian savanna combustion emissions factors for CH₄ and N₂O (0.0035, 0.0076 respectively, Russell-Smith et al., 2009b) and CO₂ (0.87, Hurst et al., 1994).

To quantify above-ground biomass, pre-clearing, eight 50 x 50 m plots were established within the 295 ha clearing area and all woody plants >1.5 m in height were identified to species with stem diameter at 1.3 m (DBH) and total tree height measured. Savanna specific allometric equations are available (Chen 2002; Williams et al., 2005) and these were used to estimate above-ground biomass for each individual tree and shrub, based on stem DBH and height measurements for all stems in

each plot. Emissions of CO₂, CH₄ and N₂O from the debris burning were estimated following the approach as outlined in the IPCC Good Practice Guidelines (IPCC 2003), which uses country or region specific emission factors for fire activity (indicated by burnt area) and the mass of fuel pyrolised to estimate the emission of each trace gas. This approach is well developed for the fire regime of north Australian savanna and is described by Russell-Smith et al. (2013) and Murphy et al. (2015a). These authors describe a novel GHG emissions abatement methodology for savannas burning that combines indigenous fire practices with an emissions accounting framework, the Emissions Abatement through Savanna Fire Management (Commonwealth of Australia 2015b, www.comlaw.gov.au/Series/F2013L01165). This methodology is a legislative instrument that establishes procedures for abatement projects for prescribed savanna burning and defines emission factors for four fuel classes; fine (grass and litter < 6 mm diameter fragments), coarse (6 mm–5 cm), heavy (>5 cm diameter) and shrubs fuels (Russell-Smith et al., 2013). Emissions of GHGs are estimated based on vegetation type, fuel mass per area for each fuel type, burn area, the burning efficiency (BEF) for each fuel type, defined as the mass of fuel exposed to fire that is pyrolised, the fuel carbon content (%), elemental C:N ratios and emission factors (EF) for each GHG (CO₂, CH₄ and N₂O) and global warming potentials for each gas. Across north Australian savanna, values for BEFs and EFs have been determined for both high (>1000 mm MAP) and low precipitation zones (1000-600 mm MAP) and for both early and late dry season fires, which are fires occurring after 1 August which typically have higher intensity and combustion efficiencies than early dry season fires (Russell-Smith et al. 2013).

We used these definitions of vegetation fuel type (woodland savanna with mixed grass) and associated EF, carbon contents, N:C ratio values as defined in the methodology to estimate GHG emissions from the debris fire using the following equation;

$$E = \sum_i (FL_i \times BEF_i \times CC_i \times N:C_i \times EF_{i,j} \times GWP_i) \quad \text{Equation 1:}$$

where E is the sum of emissions in $\text{Mg CO}_2\text{-e ha}^{-1}$ for each GHG i (CO_2 , CH_4 , and N_2O), FL_j is the fuel load for fuel type j (fine, coarse, heavy) in Mg C ha^{-1} , BEF_j is the burning efficiency factor, CC_j is the fractional carbon content, $N:C_j$ is the fuel nitrogen to carbon ratio (for N_2O emissions), EF_{ij} is the emission factor for GHG i and fuel type j and GWP_i is the global warming potential for each GHG i (after Commonwealth of Australia, 2015b). The debris fire differed from a typical savanna fire in that there was a significantly higher heavy fuel load present and it was of high intensity which consumed the vast majority of fuel (Plate 1c,d), reflected in the assumed BEFs we used. The fire-derived emissions were combined with tower-derived NEE data from the post-clearing phases (Table 3) to give a total emission in $\text{CO}_2\text{-e}$ for this LUC.

2.6 Quantifying fuel loads

To accurately quantify emissions from the debris fire, fine, coarse and heavy fuels were estimated using plots and transects established across the 295 ha deforestation area. For fine fuels, six 100 m transects were randomly located and at 20 m intervals along each transect, all fine (grass, woody litter) and coarse (twigs, sticks) fuels were harvested from 1 m^2 quadrats, dried and weighed to give a mean fine and coarse fuel mass for the site. We assigned on-site coarse woody debris (CWD), above-ground and below-ground biomass estimates to the heavy fuel class (>5 cm diameter fragments). To quantify CWD, an additional six 100 m transects were randomly located across the deforestation area and along each transect the length and diameter of all intersected CWD fragments were recorded to estimate fragment volume. In these savannas, large fragments (>10 cm diameter) are frequently hollowed from the action of termites and fire and the diameter and length of the annulus of such fragments were measured to estimate this missing volume. In addition, large fragments that were tapered were treated as a frustum of a cone and a second diameter was taken at the fragment end to improve volume estimation. Fragment volumes were calculated and converted to mass using rot classes (RC) and associated wood densities (g cm^{-3}). Five rot classes (RC) were defined and assigned to each CWD fragment to capture the decay gradient of fragments. These

~~Below ground biomass was calculated using the root:shoot ratio estimate of Eamus et al. (2002) for these savannas which was 0.38. These savanna trees have no dominant tap root, but large lateral roots in the top 30 cm of soil and up to 90% of root biomass occurs in the top 50 cm (Eamus et al. 2002). As such, we assumed that chaining and bulldozer clearing of all above ground biomass and soil ripping (ploughing) to 60 cm soil depth, plus mechanised removal of root biomass associated with tree bole, resulted in a near complete removal of both above and below ground woody biomass pools. This debris was subsequently stockpiled for curing over the dry season and then burnt (Table 2).~~

~~The CWD pool was defined as any fragment with a diameter >6 mm and was estimated using a line intercept method (Woldendorp et al., 2004). Six 100 m transects, randomly located across the cleared block, were established and along each transect the length and diameter of all intersected CWD fragments were recorded. Large fragments (>100 mm diameter) are frequently hollowed from the action of termites and fire, and the diameter of the annulus and fragment were measured to estimate this missing biomass.~~

~~Five rot classes were adapted from Rice et al. (2004) and assigned to all measured CWD fragments. Rot classes (RC) captures the decay gradient of CWD fragments and were defined as recently fallen, solid wood (RC1), solid wood with or without branches present but with signs of aging (RC2), obvious signs of weathering, still solid wood, bark may or may not be present (RC3), signs of decay with the wood sloughed and friable (RC4) and severe decayseverely decayed fragments with little structural integrity remaining (RC5). A wood density was assigned to each ~~rot~~ classRC and species (where identifiable) after Rose (2006) ~~to provide an accurate estimate of CWD that included hollowing.~~ Grass biomass was estimated using a grassy fuel loads guide developed for north Australian savanna (Johnson, 2001), which was applicable given soil type and annual and perineal grass mixture of both sites. The biomass per area from the four pools was added to give a total fuel load.and Brown (1997) to provide an accurate estimate of CWD mass that included decay~~

and hollowing. For the dominant *Eucalyptus* and *Corymbia* species wood densities ranged from 0.7 g cm⁻³ (RC1) to 0.56 g cm⁻³ (RC 5).

Above-ground biomass was quantified by surveying all woody plants >1.5 m in height or > 2 cm DBH across eight 50 x 50 m plots. All woody individuals were identified to species and stem diameter at 1.3 m height (DBH) and tree height were measured. Region specific allometric equations are available for tree species found at the CS site (Williams et al., 2005) and these were used to estimate above-ground biomass for each individual tree and shrub based on DBH and height. Below-ground biomass was calculated using the root:shoot ratio estimate of Eamus et al. (2002) for these trees. Debris from the March 2012 clearing event was allowed to cure for months through the dry season to ensure a high fire intensity (>5 MW m⁻¹) and combustion efficiency. This period is of similar duration that fuels can naturally cure in this landscape, enabling the application of the regional savanna emission factors as defined by Russell-Smith (2009b). Burning of stockpiled debris lasted approximately 10 days in August 2012, the late dry season (Table 2), a period of high winds and low daytime relative humidity (10-20%). Debris that did not burn was stock-piled for a second time and burnt to ensure all biomass was consumed with ~10% remaining as ash and charcoal, a typical fraction from high severity fires (savanna stands which was 0.38. These trees have large lateral roots in the top 30 cm of soil, with no tap root and 90% of root biomass is found in the top 50 cm of soil (Eamus et al. 2002). As such, we assumed that chaining and bulldozer clearing of all above-ground biomass followed by soil ripping (ploughing) to 60 cm soil depth, plus mechanised removal of root biomass associated with tree boles and subsequent burning, resulted in a near-complete removal of both above- and below-ground woody biomass pools (Plate 1d).

2.7 Deforestation Russell-Smith et al. 2009b). Fine ash blew away and surface charcoal fragments were incorporated into the top soil on ploughing.

Emissions from the debris burning (CO₂ and non-CO₂) were combined with NEE data from the post-clearing period to give a total emission in CO₂-e for this LUC. This estimate included CO₂

~~fluxes as captured by the flux tower plus CO₂ and non-CO₂ (CH₄, N₂O) emissions from debris burning.~~

2.6 Emissions from deforestation and savanna burning emissions at catchment to regional scales

The potential impact of any ~~future expansion of~~expanded deforestation across north Australian savanna landscapes was assessed relative to historic deforestation rates and resultant GHG emissions. ~~These emissions were also compared to~~ and arising from prescribed savanna burning, ~~a~~ This land management activity ~~that~~ contributes ~1.53% to Australia's national GHG emissions (Whitehead et al., 2014) and is 25% of the Northern Territory's annual emissions (~~Commonwealth of Australia, 2015~~); (Commonwealth of Australia, 2015a). Annual emissions from these activities (historic and future savanna deforestation and prescribed burning~~);~~) were estimated at three spatial scales; 1) catchment, 2) state/territory and 3) regional. Emissions estimates from deforestation and savanna burning were compiled for 1) the Douglas-Daly River catchment where the UC and CS sites are located (area 57,571 km²), a catchment with less than 5% of the native vegetation deforested to date (Lawes et al. 2015) but earmarked for future development; 2) the savanna area of Northern Territory (856,000 km²) and ~~for~~3) the savanna region of north Australia (~~1.93 million km²~~) as defined by Fox et al. (~~2001, (2001)~~ with MAP > 600 mm, an area of 1.93 million km² (Fig. 1, insert).

~~Historic deforestation emissions from the Douglas-Daly catchment were estimated using satellite-derived clearing areas (1990–2013) for the catchment as reported by Lawes et al. (2015). These annual deforestation areas were combined with our estimate of GHG emissions from the CS site to give a mean annual estimate in Gg CO₂-e y⁻¹. To estimate GHG emissions from savanna deforestation at state/territory and regional scales, data for the Northern Territory and the north Australian savanna region were taken from the reported emissions under Activity A.2 under the Land Use, Land-Use Change and Forestry sector of State and Territory Greenhouse Gas Inventories~~

(Commonwealth of Australia, 2015). Annual reporting of these GHG emissions is state based, not biome based, and for the regional savanna estimate, data for Western Australia, the Northern Territory and Queensland were used but included only the area within each state that was as defined as savanna after Fox et al. (2001, Fig. 1) and emissions from each state were summed to give the north Australian savanna regional estimate.

To estimate the GHG emissions from future expanded deforestation across north Australia, we upscaled our estimate of deforestation emissions per hectare using the areas identified as having future clearing potential following the land use assessment of north Australian catchments by Petheram et al., (2014). This preliminary assessment identified catchments to be cleared based upon surface water storage potential and proximity of land resources suitable clearing for irrigation development to enable high value farming such as irrigated agriculture, horticulture or improved pasture. Using these criteria, suitable catchments in Western Australia (Fitzroy River, Ord Stage 3; 75 000 ha potential area), the Northern Territory (Victoria, Roper Rivers, Ord Stage 3, Darwin-Wildman River area; 114, 500 ha) and Queensland (Archer, Wenlock, Normanby, Mitchel Rivers; 120 000 ha) were selected. This gives a projected savanna clearing area of 311, 000 ha, equivalent to an additional 16% of cleared land over and above the 1,886,512 ha that has been cleared across the savanna biome since 1990 (Commonwealth of Australia, 2015). Projected emissions calculations included emissions from historic emissions plus additional emissions estimates from the expanded deforestation areas. Emissions from the expanded deforestation areas were calculated assuming any such clearing would occur over a five year period. This filter provided identical areas for comparison of mean annual savanna deforestation and prescribed burning emissions.

For savanna burning, CH₄ and N₂O emissions Emissions of GHG from historic deforestation from the Douglas-Daly catchment were estimated using our estimates for savanna land conversion combined with satellite-derived annual deforestation area (1990-2013) as reported by Lawes et al. (2015) for this catchment to give a catchment scale mean annual estimate of emissions from

1 deforestation in Gg CO₂-e y⁻¹. Annual deforestation emissions data for the Northern Territory and
2 the north Australian savanna region were taken from the National Greenhouse Gas Inventory
3 (NGGI) for the same period 1990-2013. The Department of Environment is responsible for
4 reporting sources of greenhouse gas emissions and removals by sinks in accordance with UNFCCC
5 Reporting Guidelines on Annual Inventories and the supplementary reporting requirements under
6 the Kyoto Protocol. State and Territory GHG Inventories are reported for 1990 to 2013
7 (Commonwealth of Australia, 2015a) and we used data for the Land Use, Land-Use Change
8 and Forestry sector, Activity A.2 Deforestation. These emissions are reported for each State, but are
9 not biome based and for our regional savanna estimate, emissions data for Western Australia, the
10 Northern Territory and Queensland were used but were calculated using the area within each state
11 that was defined as savanna by Fox et al. (2001, Fig. 1). Mean annual deforestation emissions from
12 the savanna area of each state and territory (1990-2013) were summed to calculate a mean (±SD)
13 annual deforestation rate for the north Australian savanna area (1.92 million km²) in Gg CO₂-e y⁻¹.

14 Emissions from savanna burning were calculated using the on-line Savanna Burning Abatement
15 Tool (SAVBat2, www.savbat2.net.au), ~~where an area is defined using~~ using the pre-defined
16 Vegetation Fuel Types (VFTs) mapping for north Australian savanna (Fisher and Edwards, 2015;
17 Thackway, 2014), both components of the Emissions Abatement through Savanna Fire
18 Management methodology. SAVBat2 combines satellite derived burnt area mapping
19 (www.firenorth.org.au) ~~and pre-defined Vegetation Fuel Types (VFTs) mapping (Fisher and~~
20 ~~Edwards, 2015; Thackway, 2014). In accordance with IPCC accounting rules, only non-CO₂ fluxes~~
21 ~~are reported for savanna burning, as it is assumed that CO₂ emissions from dry season burning is~~
22 ~~entirely offset by~~ with fuel load estimates from VFT mapping, GHG emission factors and burn
23 efficiencies to estimate annual emissions from burn areas. In accordance with IPCC accounting
24 rules, only non-CO₂ emissions are reported for savanna burning as it is assumed that CO₂ emissions
25 from dry season burning is offset by re-growth of vegetation (mostly C₄ grasses) in subsequent wet

season(s) (IPCC, 1997). ~~To estimate~~ However, for comparisons with deforestation emissions, we calculated emissions of CO₂ as well as non-CO₂ emissions. SAVBat2 uses satellite-derived burnt area mapping as generated by the North Australia Fire Information (NAFI) system (www.firenorth.org.au), following methods described by. These estimates were compiled for the ~~thesame areas as~~ savanna ~~areas of interest~~ deforestation estimates; the Douglas-Daly River catchment, savanna of the Northern Territory NT and ~~the~~ north Australian savanna ~~region~~. Mean annual burning emissions for 1990-2013 were calculated and are reported as non-CO₂ ~~only~~ (CH₄, N₂O) and total emissions (CO₂, CH₄ and N₂O) in Gg CO₂-e y⁻¹.

2.8 Emissions from expanded deforestation across north Australia

Emissions from expanded deforestation across north Australia was estimated by upscaling our estimate of deforestation emissions per hectare from catchment areas identified as having future clearing potential. These areas were based on the land use assessment of north Australian catchments by Petheram et al. (2014) and identified catchments with development potential based upon surface water storage and proximity of land resources suitable for irrigation development for agriculture, horticulture or improved pastures. Using these criteria, suitable catchments were identified in Western Australia (Fitzroy River, Ord Stage 3; 75 000 ha potential area), the Northern Territory (Victoria, Roper Rivers, Ord Stage 3, Darwin-Wildman River area; 114, 500 ha) and Queensland (Archer, Wenlock, Normanby, Mitchel Rivers; 120 000 ha). This gives a potential savanna deforestation area of 311, 000 ha, equivalent to an additional 16% of cleared land over and above the 1,886,512 ha that has been cleared across the savanna biome since 1990 (Commonwealth of Australia, 2015a). Projected emissions included mean annual emissions from historic deforestation rates plus emissions from this expanded deforestation scenario. Expanded deforestation areas were calculated assuming any such clearing would occur over a five year period and are reported as non-CO₂ (CH₄, N₂O) and total emissions (CO₂, CH₄ and N₂O) in Gg CO₂-e y⁻¹.

3.0 Results

3.1 Pre-clearing site comparisons

Comparisons between the two sites were made using pre-clearing meteorology, flux observations and energy balance closure (Figs. 2, 3) to ensure the CS for UC and UC sites were comparable. Table 3 provides compared (Fig. 2). Mean monthly NEE, R_e and GPP for each LUC phase for both sites are given in Mg ha^{-1} per month. Table 3. Flux measurements prior to clearing (intact canopy phase) were made for 161 days, a period spanning the late dry to early wet season transition (September-December) through to the mid-wet season (January-February, Table 2). Energy balance closure was high. Flux data at the CS site were validated by assessing energy balance closure, with a regression of the between energy balance components giving suggesting closure was high with a slope of 0.91 and an R^2 of 0.95 ($n=4778$). Site differences for this site (Fig. 2, each phase were tested using one-way ANOVA using daily mean NEE with days as replicates. For Phase 1). At the UC site this slope, mean daily NEE was 0.87 using all available data (2007-2015) not significantly different between the two sites during ($P<0.64$, $df=321$). Seasonal patterns of T_{air} , VPD (Fig. 3b2b), LAI (Fig 3e2c) and C fluxes (NEE, GPP, R_e , Fig 3d2d) were similar when both sites were intact, although rainfallprecipitation was 340 mm higher at the UC site (Table 3).

At both sites, NEE shifted from being a weak sink of less than $-1 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ during the late dry season, to a net source of CO_2 during the early wet season (Fig. 3d2d). During this period, R_e increased rapidly from $+2 \mu\text{mol m}^2 \text{ s}^{-1}$ to $+5 \mu\text{mol m}^2 \text{ s}^{-1}$ in early October with the onset of wet season rain, but then remained relatively constant for the remainder of the wet season.

As the wet season progressed, temporal patterns of GPP were similar at both sites and steadily increased to -6 to $-7 \mu\text{mol m}^{-2} \text{ s}^{-1}$ and remained at this rate until cleared (March 2012). R_e was relatively stable during this period and NEE increased to $-2 \mu\text{mol m}^{-2} \text{ s}^{-1}$ through the wet season (December to February). Despite the higher rainfallprecipitation received at the UC site, mean monthly NEE, GPP and R_e differed by $<10\%$ (Table 3, intact canopy phase). Normalising fluxes by

MODIS LAI for each site further reduced differences to 2% (data not shown), suggesting site differences were small and the UC site provides a suitable control for the CS site.

3.2 Fluxes following clearing

Clearing of the 295 ha block commenced on 2 March 2012 and the bulldozers reached the footprint of the flux tower at ~0900h local time on 6 March (Fig. 3a,b). As for Phase 1, energy balance closure of flux tower data for LUC Phases 2 to 4 (post-clearing phases) was high, with a slope >0.9 and $R^2 > 0.92$. Over all phases at the CS site, closure was lower, with a slope of 0.81 ($R^2 = 0.95$, $n=26,395$), similar to that of the UC site at 0.87 ($R^2 = 0.93$, $n=99,998$).

The four day clearing event occurred during relatively high soil moisture conditions, with surface (5 cm depth) θ_v ranging from 0.08 to 0.10 $\text{m}^3 \text{m}^{-3}$ and sub-soil θ_v (50 cm depth) ranging from 0.12 to 0.14 $\text{m}^3 \text{m}^{-3}$. As a result, pre-clearing fluxes were high and NEE reached $-15 \mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$ during the middle of the day (Fig. 43). Mean daily NEE for the week prior to clearing was a net CO_2 sink of $-0.60 \pm 0.63 \mu\text{mol m}^{-2} \text{s}^{-1}$, and was not significantly different to mean daily NEE at the UC site of $-0.80 \pm 0.93 \mu\text{mol m}^{-2} \text{s}^{-1}$ (ANOVA, $P < 0.03$). For the three weeks following clearing, the CS site rapidly became a net source of CO_2 with a mean daily NEE of $+4.38 \pm 0.24 \mu\text{mol m}^{-2} \text{s}^{-1}$, with a much reduced diurnal amplitude and no response to rainfall events (Fig 4a,b). High closure (slope >0.9) was observed during Phases 2 to 4, although this was reduced (slope $=0.75$) for the post-fire and soil preparation, Phases 6-9.

Table 3 provides values of rainfallprecipitation and monthly NEE, Re and GPP for the remaining seven LUC phases following clearing, namely debris decomposition and curing (153 days), burning (22 days), wet season regrowth (80 days), followed by soil tillage and preparation of irrigated raised soil beds (181 days). For each phase, the comparable flux estimate from the UC site is estimated for all post clearing phases and for the entire observation period (Table 3). Following clearing, GPP at the CS site was reduced by a factor of 3.5 when compared to the UC for the same period (March 2012 – January 2013, Table 3). While greatly reduced, GPP still occurred at the CS

site during this 13.7 month period ($-0.38 \text{ Mg C ha}^{-1} \text{ month}^{-1}$), via re-sprouting of felled overstorey and sub-dominant trees and shrubs, as well as grass germination and growth stimulated by early wet season rainfallprecipitation (November 2012-January 2013, 361 mm, Table 3). Ecosystem respiration during this period was higher at the UC site ($+1.12 \text{ Mg C ha}^{-1} \text{ month}^{-1}$) when compared to the CS site ($+0.82 \text{ Mg C ha}^{-1} \text{ month}^{-1}$) and given the large decline in GPP, the CS site was a small net C source at $+0.51 \text{ Mg C ha}^{-1} \text{ month}^{-1}$, as compared to the UC site which was a weak sink of $-0.03 \text{ Mg C ha}^{-1} \text{ month}^{-1}$.

Cumulative NEE over all the post-clearing LUC phases was $+7.2 \text{ Mg C ha}^{-1}$ at the CS site as compared to a net sink of $-0.78 \text{ Mg C ha}^{-1}$ at the UC site (Table 3). The temporal dynamics of cumulative NEE across all LUC phases (note differences in phase duration) is summarised in Fig. 54, which compares fluxes from both sites for the complete observation period. Three significant periods of C emission are evident in Fig. 54. Firstly, the clearing event and the subsequent switch from a C sink to a net source of 1.9 Mg C ha^{-1} due to soil disturbance and the decomposition of biomass. Secondly, this was followed by a reduction in source strength over the dry season of 2012, attributable to a reduction in R_e during the dry season (2012 dry season pre-burn phase, Table 3). Thirdly, there were other major emissions attributed to soil tillage and bed preparation in the wet and dry seasons of 2013, a cumulative net emission of $+2.75 \text{ Mg C ha}^{-1}$ that occurred over the final six months (Fig. 54) in preparation for cropping. Over this phase, the UC site was a net sink of $-0.62 \text{ Mg C ha}^{-1}$.

3.4 Emissions from debris burning

~~At the CS site, the~~ Table 4 gives fuels loads, BEF, EF, carbon content and N:C ratios for each fuel distribution of cleared debris type used to estimate the GHG emission from the debris burning. Fuel load was dominated by heavy fuels (~~$\geq 6 \text{ mm diameter}$, as defined by Russell-Smith et al., 2009~~), with a mean ($\pm \text{SD}$) above-ground biomass of ~~2726.9 ± 7.50~~ $2726.9 \pm 7.50 \text{ Mg C ha}^{-1}$ and a range of 14.4 to $39.3 \text{ Mg C ha}^{-1}$ across the eight biomass plots. The mean ~~coarse root~~ below-ground biomass was

estimated at $10.69.0 \pm 2.9 \text{ Mg C ha}^{-1}$. The CWD pool was assessed using six transects and was spatially variable, with mean CWD of $1.5 \pm 1.84 \text{ Mg C ha}^{-1}$. Grass biomass was assessed at 0.9 Mg C ha^{-1} . The total biomass from the four pools was estimated to be $40.9 \text{ Mg C ha}^{-1}$, which became the fuel load following clearing, soil ripping and lifting of tree stumps. CWD was $1.4 \pm 0.6 \text{ Mg C ha}^{-1}$. Fine and associated root mass, 5 months of dry season curing and stocking piling. We assumed 90% of all coarse fuels were consumed with 10% ash and charcoal remaining as observed by Russell-Smith et al. (2009b) for late dry season fires. 1.4 ± 0.7 and $0.5 \pm 1.0 \text{ Mg C ha}^{-1}$ respectively, giving a total fuel mass of $38.2 \text{ Mg C ha}^{-1}$. Using this biomass data, these fuel loads with savanna emission factors EF and the BEFs estimated for the site gave an emissions of CO_2 , CH_4 and N_2O , for each fuel type and the estimated emission from debris burning totalled $121.9 \text{ Mg CO}_2\text{-e ha}^{-1}$, with 9.5% of this total comprising non- CO_2 emissions (Table 4).

3.5 Total GHG emission

Emissions derived from debris burning needs to be combined with the post-clearing NEE as measured by the EC system to provide a total GHG emissions estimate from this LUC in units of $\text{CO}_2\text{-e}$. The LUC phases following clearing spanned a 502 day period (Table 3), and NEE was $+7.2 \text{ Mg C ha}^{-1}$ (Table 3), which is in units of $\text{CO}_2\text{-e}$ or $+26.4 \text{ Mg CO}_2\text{-e ha}^{-1}$. In comparison, NEE from the UC site over the same period was $-0.78 \text{ Mg C ha}^{-1}$ or $-2.9 \text{ CO}_2\text{-e ha}^{-1}$. Adding emission NEE from post-clearing phases (Phases 2-9, Table 3) to emissions from debris burning to NEE (Table 4) gave a total emission of $+148.3 \text{ Mg CO}_2\text{-e ha}^{-1}$ for the CS site, although the fire-derived non- CO_2 emission was only 3% of this total. The CO_2 -only emission from fire debris burning plus post-clearing NEE was $+136.7 \text{ Mg CO}_2 \text{ ha}^{-1}$, which was a flux 5045 times larger than the observed savanna CO_2 sink at the UC site over this the post-clearing period.

3.6 Upscaled and projected emissions from deforestation and savanna burning

Table 45 provides mean (\pm SD) GHG emissions estimates for the two of the most significant GHG-savanna burning and deforestation for 1990-2013. At all spatial scales, annual mean burnt area dwarfed the mean annual land area deforested. For the Douglas-Daly catchment area, reportable non-CO₂ emissions from savanna burning were 577 \pm 124 Gg CO₂-e y⁻¹, almost four times larger than emissions from the mean annual savanna deforestation rate of 163 \pm 162 Gg CO₂-e y⁻¹. For the Northern Territory savanna, mean annual burning emissions were an order of magnitude larger than mean annual deforestation emissions (Table 4) and two orders of magnitude larger if CO₂ emissions were included. At the regional scale of, the mean annual deforestation rate across the northern Australian savanna, the deforestation rate was 16,161 \pm 5,601 Gg CO₂ y⁻¹, with emissions from Queensland savanna area dominating this amount at 15,762 \pm 5,566 Gg CO₂ y⁻¹. This is an emission-double that of the reportable savanna(non-CO₂ only) emission from prescribed burning at 6,740 \pm 1,740 Gg CO₂ y⁻¹ (Table 45).

Emissions estimates that include future deforestation rates would be equivalent to savanna burning, at least for the duration of the additional clearing-deforestation. For the Douglas-Daly catchment, this future emission is estimated at 781756 Gg CO₂-e y⁻¹ and across the Northern Territory savanna area, this would be 3,574413 Gg CO₂-e y⁻¹, similar rates of emission that are equivalent to current savanna-burning emissions for the catchment (Douglas-Daly catchment, 577 \pm 124) and the state scales (Northern Territory (Table 4) savanna, 3,490 \pm 922 Gg CO₂-e y⁻¹). Emissions that include future deforestation rates for the northern Australian savanna region were estimated at 24,769393 Gg CO₂-e y⁻¹ and would be ~30% in excess of current regional three times the reportable savanna burning annual emissions (Table 5).

4.0 Discussion

Australia has lost approximately 40% of its native forest and woodland since colonisation (Bradshaw, 2012)(Bradshaw, 2012), with most of this clearing for primary production in the eastern and south-eastern coastal region. Attention has now turned to the productivity potential

development of the largely intact northern savanna landscapes, which will involve trade-offs between management of land and water resources for primary production and biodiversity conservation (~~Adams and Pressey, 2014; Grundy et al., 2016~~)(Adams and Pressey, 2014; Grundy et al., 2016). Globally and in Australia, savanna fire ecology and fire derived GHG emissions have been reasonably well researched (Beringer et al., 1995; Cook and Meyer, 2009; Livesley et al., 2011; Meyer et al., 2012; Walsh et al., 2014; van der Werf et al., 2010) and the impacts of fire on the functional ecology of Australian savanna has been recently reviewed by Beringer et al. (2015). In this study, we have focussed ~~upon~~ savanna deforestation and land preparation for agricultural use. These phases result in a series of events that may lead to pulsed GHG emissions that would otherwise be missed or greatly under-estimated by episodic measurements taken at a weekly or monthly frequency after an initial tree felling event (Neill et al., 2006; Weitz et al., 1998).

~~Eddy covariance measurements~~

We used the eddy covariance methodology as it provides a direct and non-destructive measurement of the net exchange of CO₂ (and other GHG gases) at high temporal resolution, ranging from 30-~~min~~ minute intervals to daily, monthly, seasonal and annual estimates. ~~It~~The method is useful as a full carbon accounting tool as all exchanges of CO₂ from autotrophic and heterotrophic components of the ecosystem undergoing change are quantified. (~~Hutley et al., 2005~~). This ~~makes it a useful tool~~approach provides essential data for bottom-up GHG and carbon accounting studies as ~~micrometrological~~micrometeorological conditions and associated fluxes can be tracked through time for the duration of a land use conversion (~~Hutley et al., 2005~~).

~~At the CS site, burning of post-clearing debris comprised 83% of the total emission of 154.4 Mg CO₂-e ha⁻¹ (0.154 Gg CO₂-e ha⁻¹ y⁻¹), with the remainder attributed to the net ecosystem exchange of CO₂ as measured by the flux tower. This comprised significant CO₂ losses via respiration of debris, enhanced soil CO₂ efflux from soil disturbance and tillage, which was partially offset by net uptake of CO₂ from woody re-growth, re-sprouting, grass germination and growth~~

(Fig. 5). Soil disturbance via ripping, tillage and preparation was responsible for almost 10% of the CO₂ emission from the conversion. The EC flux tower was operational during the clearing event, demonstrating the utility of this method as the switch of the ecosystem being a net CO₂ sink to being a net source occurred over a few hours of the clearing event completing (Fig. 3). During the LUC phase changes, there was little evidence of major pulses of CO₂ flux, instead there was a rapid transition to a new diurnal pattern following management events, such as the clearing (Fig. 4) or the commencement of soil preparation (data not shown). This is in contrast to non-CO₂ fluxes emissions, in particular N₂O, with short term emissions often follow disturbance (Grover et al., 2012; Zona et al., 2013).

At the CS site, burning of post-clearing debris comprised 82% of the total emission of 148.4 Mg CO₂-e ha⁻¹, with the remainder attributed to NEE as measured by the flux tower. This flux comprised significant CO₂ losses via respiration of debris, enhanced soil CO₂ efflux from soil disturbance and tillage, which was partially offset by net uptake of CO₂ from woody re-sprouting post-clearing and periods of grass growth following wet season rainfall (Fig. 4). Soil disturbance via ripping, tillage and preparation was responsible for 10% of the CO₂ emission from the conversion. The EC flux tower was operational during the clearing event, demonstrating the utility of this method as the switch of the ecosystem from being a net CO₂ sink to being a net source occurred over a number of hours as the clearing event was completed (Fig. 3). During the LUC phase changes, there was little evidence of major pulses of CO₂ flux, instead there was a rapid transition to a new diurnal pattern following the clearing (Fig. 3) or the commencement of soil preparation (data not shown). This is in contrast to non-CO₂ flux emissions, in particular N₂O, with short term emissions often follow disturbance (Grover et al., 2012; Zona et al., 2013) and can be a significant fraction of annual emissions.

The net CO₂ source as measured by the flux tower represents an emission that would be missed if vegetation biomass density alone was used to estimate LUC emissions, the approach used in

current remote sensing LUC studies ~~for~~at regional and ~~national accounting purposes~~. As ~~such, continental scales, data that is~~ the ~~emission associated complete oxidation~~basis of biomass by ~~fire would be more comparable to reported deforestation~~emissions reporting for the LULUC sector.

The total GHG emission we report in this study is more accurately described as a land conversion, as it includes the oxidation of biomass plus emissions associated with soil disturbance and tillage required for a conversion to a cropping or grazing system.

The emission estimate from this study does not include non-CO₂ soil derived fluxes of CH₄ and N₂O, which can be significant for LUC events in certain ecosystems (Tian et al., 2015). Grover et al. ~~(2012) compared soil CO₂ and non-CO₂ fluxes from native savanna with young pasture and old pastures (5-7 and 25-30 years old) in the Douglas-Daly River catchment. Soil emissions of CO₂-e were 30% greater on the pasture sites as compared with native savanna sites, with this change being dominated by increases in CO₂ emission and soil CH₄ exchange shifting from a small net sink to a small net source at the pasture sites. Non-CO₂ soil fluxes were generally small, especially N₂O emissions, although these measurements were made many years after the LUC event and there is uncertainty as to their relevance for a recently deforested and converted savanna site. An additional pathway for CH₄ and N₂O emissions in these savannas is through termite activity (Jamali et al., 2011a, 2011b), and in our study, termite mounds were abundant across the CS site, but were largely destroyed by clearing and soil preparation, potentially reducing net non-CO₂ emissions. Further work is required to quantify these fluxes and refine our total emission estimate for this LUC event.~~(2012) compared soil CO₂ and non-CO₂ fluxes from native savanna with young pasture and old pastures (5-7 and 25-30 years old) in the Douglas-Daly River catchment. Soil emissions of CO₂-e were 30% greater on the pasture sites as compared with native savanna sites, with this change being dominated by increases in CO₂ emission and soil CH₄ exchange shifting from a small net sink to a small net source at the pasture sites. Non-CO₂ soil fluxes were generally small, especially N₂O emissions, although these measurements were made many years after the LUC event and there is uncertainty as to their relevance for a recently deforested and converted savanna site. An additional

pathway for CH₄ and N₂O emissions in these savannas is via termite activity (Jamali et al., 2011a, 2011b). In our study, termite mounds were abundant across the CS site but were largely destroyed by clearing and soil preparation, potentially reducing the net non-CO₂ emission following conversion. Further work is required to quantify these non-CO₂ fluxes not associated with debris burning to refine our total emission estimate for savanna deforestation.

This land conversion represents the loss of decades of carbon accumulation in this mesic savanna (>1000 mm MAP), ecosystems which are currently thought to ~~be~~ a weak carbon sink (~~Beringer et al., 2015~~). (Beringer et al., 2015). The 8-year ensemble mean NEE for the UC site was -0.11 ± 0.16 Mg C ha⁻¹ y⁻¹ and is representative of a savanna site at ~~or near~~ a near-equilibrium state in terms of carbon balance, given the ~~moderate~~low fire frequency (~~13~~ in 513 years, Table 21) with high severity fires uncommon (1 in 8 years of ~~measurement~~flux measurements). The annual increase in tree biomass at this UC site is 0.6 t C ha⁻¹ y⁻¹ (Rudge, Hutley, Beringer, unpublished data), ~~suggesting and given an above-ground standing biomass of 28 t C ha⁻¹ suggests a~~ regeneration period of approximately four ~~to five~~ decades after stand ~~replacing~~replacement disturbance event (~~extreme fire, cyclone, flood, harvest~~). ~~such as deforestation for this savanna type.~~

Even after the large pool of carbon is lost following oxidation of biomass, carbon loss ~~will~~may ~~continue~~continue on cleared land via continued soil carbon mineralisation, leading to a slow decline in soil carbon storage that is frequently reported for forest to cropping LUC ~~systems~~(Jarecki and Lal, 2003; Lal and Follett, 2009). Conversion of forest or woodland to improved pasture grazing may result in either increases or decreases in soil carbon (Sanderman et al., 2010). Alternatively, it is possible that carbon sequestration may occur post-clearing via woody regrowth if a cleared site is abandoned and not further prepared for cultivation. This has actually been a relatively common transition and a significant sequestration pathway that needs to be included in savanna LUC assessments (Henry et al. 2015). Admittedly, if savanna cleared land does fully transition to a

cropping system, some fraction of the lost carbon could also be replaced or sequestered by ~~the new~~
~~agricultural or~~ horticultural or forestry land ~~uses~~.

~~There are few detailed, plot-scale studies of GHG emissions from savanna clearing in north~~
~~Australia. Several studies (Law and Garnett 2009, 2011) used the Full Carbon Accounting Model~~
~~(FullCAM Ver 3.0, Commonwealth of Australia, 2015a; Richards and Evans, 2004) to generate~~
~~spatial maps of above and below ground biomass and soil organic carbon pools across the NT. The~~
~~FullCAM model uses spatial and temporal soil, climate, rainfall data with NVIS major vegetation~~
~~classes to simulate carbon losses (as GHG emissions) and uptake between the terrestrial biological~~
~~system and the atmosphere. Land use change scenarios can be run within the model and Law and~~
~~Garnett (2009) examined deforestation emissions from the Eucalypt woodland NVIS vegetation~~
~~class, as per UC and CS site classification. Modelled emissions were 136 ± 42 Mg CO₂-e,~~
~~comparable to the deforestation (fire only) estimate of 128 Mg CO₂-e reported in our study.~~

~~Henry et al. (2015) used a life cycle assessment approach to quantifying GHG emissions from~~
~~LUC associated with beef production in eastern Australia. Australia's major beef producing areas~~
~~across central and southern Queensland and northern central New South Wales were classified into~~
~~11 bioregions, with the northern-most bioregion, the northern Brigalow Belt, falling within the~~
~~savanna biome. Vegetation biomass from this bioregion was estimated at 84.7 ± 7.1 Mg ha⁻¹ or ~ 41.4~~
~~Mg C ha⁻¹, with an emission estimated at 129 Mg CO₂-e (Henry et al., 2015), almost identical to the~~
~~woodland biomass density and resultant emission with deforestation (fire derived emission) from~~
~~the CS site of this study.~~

~~Our emissions estimate is robust for this vegetation class and can be upscaled and as such can~~
~~be compared and upscaled with other land sector activities, such as prescribed savanna burning. At~~
~~a regional scale, current levels of savanna burning dominate emissions compared to land clearing~~
~~rates (Table 4). The cumulative deforestation across the savanna region since 1990 (1,886,512 ha) is~~
~~17 times smaller than the annual savanna burn area as approximately 30 to 70% of the savanna area~~

1 is burnt annually (Russell-Smith et al., 2009a). Modelling NEP for savanna biome for 1990–2010
2 (Beringer et al., 2015; Haverd et al., 2013), suggests the north Australia savanna is near carbon
3 neutrality, or is a weak source of CO₂ to the atmosphere once regional scale fire emission are
4 included. As such, the IPCC assumption that CO₂ emissions from the previous year's burning are
5 recovered by the following year's wet season growth may have some validity for regional scale
6 GHG accounting. This assumption at plot to catchment scales may not be valid, as localised
7 interannual variability in rainfall, site history and fire regimes may result in either net accumulation
8 or loss of carbon (Hutley and Beringer, 2011; Murphy et al., 2014, 2015). Assuming year to year
9 CO₂ loss from burning is re-sequestered, assessment of the non-CO₂-only emissions from savanna
10 burning with clearing is useful. This comparison suggests projected deforestation emissions (24,769
11 Gg CO₂-e y⁻¹, Table 4) could be well in excess of current annual burning emissions, at least for the
12 period of enhanced clearing, which in this study, we assume to be five years.

13 There are few detailed, plot scale studies of GHG emissions from savanna clearing in north
14 Australia. Several studies (Law and Garnett 2009, 2011) used the Full Carbon Accounting Model
15 (FullCAM Ver 3.0, Commonwealth of Australia, 2015a; Richards and Evans, 2004) to generate
16 spatial maps of above- and below-ground biomass and soil organic carbon pools across the NT. The
17 FullCAM model uses spatial and temporal soil, climate, precipitation data with NVIS major
18 vegetation classes to simulate carbon losses (as GHG emissions) and uptake between the terrestrial
19 biological system and the atmosphere. Land use change scenarios can be run within the model and
20 Law and Garnett (2009) examined deforestation emissions from the Eucalypt woodland NVIS
21 vegetation class, as per UC and CS site classification. Modelled emissions were 136±42 Mg CO₂-e,
22 comparable to our deforestation estimate of 121.4 Mg CO₂-e. Henry et al. (2015) used a life cycle
23 assessment approach to quantify GHG emissions from LUC associated with beef production in
24 eastern Australia. Australia's major beef producing areas across central and southern Queensland
25 and northern central New South Wales were classified into 11 bioregions, with the northern most

bioregion, the northern Brigalow Belt, falling within the savanna biome. Vegetation biomass from this bioregion was estimated at $84.7 \pm 7.1 \text{ Mg ha}^{-1}$ or $\sim 41.4 \text{ Mg C ha}^{-1}$, with an emission estimated at $129 \text{ Mg CO}_2\text{-e}$ (Henry et al., 2015), similar to the woodland biomass density and resultant emission with deforestation from the CS site of this study.

Our emissions estimate is robust for this vegetation class and can be upscaled and compared with other land sector activities such as prescribed savanna burning. At a regional scale, current levels of savanna burning dominate emissions compared to land clearing rates (Table 5). The cumulative deforestation area across the savanna region since 1990 (1,886,512 ha) is 17 times smaller than the mean annual savanna burn area (32 Mha, Table 5) as approximately 30 to 70% of the savanna area is burnt annually (Russell-Smith et al., 2009). Modelling NEP for savanna biome for 1990-2010 (Beringer et al., 2015; Haverd et al., 2013) suggests the north Australian savanna is near carbon neutrality, or is a weak source of CO_2 to the atmosphere once regional scale fire emissions are included. As such, the IPCC assumption that CO_2 emissions from the previous year's burning are recovered by the following year's wet season growth may have some validity for regional scale GHG accounting. This assumption at plot to catchment scales may not be valid, as localised interannual variability in rainfall, site history and fire management can result in either net accumulation or loss of carbon (Hutley and Beringer, 2011; Murphy et al., 2014, 2015b). Assuming year to year CO_2 emitted from burning is re-sequestered, assessment of the non- CO_2 only emissions from savanna burning with deforestation is useful. This comparison suggests projected deforestation emissions ($24,393 \text{ Gg CO}_2\text{-e y}^{-1}$, Table 5) could be well in excess of current annual burning emissions ($6,740 \text{ Gg CO}_2\text{-e y}^{-1}$, Table 5), at least for the period of enhanced clearing, which in this study we assumed to be five years.

In 2013, Australia's total reported GHG emission was 548,440 Gg $\text{CO}_2\text{-e}$ and the impact of expanded savanna deforestation on the national emission can be estimated using data in Table 4. Table 4 provides the5 which provide estimates of mean annual ~~emission-rate~~emissions from an

~~annual-deforested~~deforestation area, giving a mean annual deforestation emission per ha averaged for the entire savanna area, which is $221 \pm 50.8 \text{ Mg CO}_2\text{-e ha}^{-1}$ using 1990 to 2013 data (Commonwealth of Australia, ~~2015~~2015a). This value represents a spatially averaged emission as it is derived from the full range of savanna vegetation types and above-ground biomass, which across the Northern Territory savanna area ranges from 10 to 70 Mg C ha^{-1} (Law and Garnett, 2011). Assuming this emission per ha, an additional 311, 000 ha of savanna deforestation, cleared over a five year period, adds $12,099 \text{ Gg CO}_2\text{-e y}^{-1}$. For the duration of the expanded deforestation, this is a 2.2% increase to Australia's nation emission over and above the historic savanna LUC emissions, ~~(16,161 Gg CO₂-e y⁻¹)~~, which are 2.9% of national emissions. Using our finding from flux tower measurements that a land conversion (deforestation followed by site tillage and preparation for cultivation) adds an additional ~~17~~18% of GHG emissions to a deforestation event, expansion of northern land development could ~~increase Australia's add an additional 3% or 33, 350 Gg CO₂-e y⁻¹ to the reportable national GHG by -6% per annum~~emissions for the duration of the expanded deforestation period.

This assessment is subject to a number of uncertainties. Firstly, a component of our emissions estimate is based on eddy covariance measurements of CO₂ flux, which typically have an error of 10-20% ~~(Aubinet et al., 2012)~~(Aubinet et al., 2012). In this study, energy balance closure suggested fluxes were underestimated by up to ~~19~~13% across the entire observation period ~~(Fig. 2)~~. Energy balance closure ranged from <10% flux loss during the intact canopy phase to >20% error during the final three LUC phases when the flux instruments were at 3 m height measuring net soil CO₂ emissions from the smoothed, vegetation-free ploughed soil surface during preparation.

Secondly, it is difficult to predict the nature of future deforestation (rate, area ~~and~~, specific location) and ~~development across the savanna region and the emissions comparison~~emission comparisons presented here ~~is~~are indicative only. Catchments selected by Petheram et al. (2014) regarded as suitable or with potential ~~to be cleared in the~~for future ~~and development~~ were based on

1 | biophysical properties ~~and only~~, were unconstrained by the regulatory environment and ~~they~~ did not
2 | account for conservation and cultural values placed on identified land and water resources. In
3 | addition, challenges to agricultural expansion in northern Australia include uncertain land and water
4 | tenure, high development costs and lack of existing water infrastructure, logistics and technical
5 | constraints, lack of human capital and distance to ~~market, are markets~~, all factors that ~~are likely~~
6 | ~~to may~~ restrict land clearing. It is well understood that the availability and cost of water for irrigated,
7 | or irrigation assisted agriculture is critical for viable agriculture in northern Australia (Petheram et
8 | al., ~~2008, 2009~~)2008, 2009). Australian ~~government~~Government policies currently support small-
9 | scale, precinct or project scale approaches, based on well-understood water and soil resources,
10 | where water allocation is capped. The current policy and market instruments are likely to ensure
11 | that development remains measured and restricted, unlike development of previous decades in other
12 | regions of eastern and southern Australia.

13 | As a result we used a conservative estimate of potential land suitability area (311, 000 ha over a
14 | five year clearing period), as estimates of assumed clearable area ~~rangeranging~~ up to 700,000 ha
15 | ~~(e.g., Douglas-Daly catchment, Adams and Pressey, 2014)~~(e.g., Douglas-Daly catchment, Adams
16 | and Pressey, 2014) or over 1 million ha across north Australia (Petheram et al. 2014), areas that
17 | may be unlikely given capital investment requirements as well as conservation and cultural
18 | considerations. Our comparison with burning emissions is also influenced by the
19 | ~~clearing~~deforestation period we assume. This was based on patterns of historic rates of clearing as
20 | there are periods when deforestation rates have easily exceeded 311,000 ha over five ~~years~~year
21 | periods, particularly in Queensland (Commonwealth of Australia, ~~2015~~),~~but~~2015a) and a longer
22 | ~~clearing~~-duration of deforestation reduces the ~~annual totals and~~ impact on ~~the annual~~ national GHG
23 | ~~account~~accounting.

24 | There is also uncertainty arising from our emissions from debris burning. Russell-Smith et al.
25 | (2009) estimated errors associated with emissions estimates from the Western Arnhem Land Fire

Abatement (WALFA) project, a savanna burning based GHG abatement scheme operating in the Northern Territory. This is a project area of the 23,893 km² consisting of a wide range vegetation types including open-forest and woodland savanna and sandstone heaths in escarpment areas. Russell-Smith et al. (2009) estimated the accountable emissions from savanna burning at 272 ±100 Gg CO₂-e y⁻¹ (95% CI), an error of 30–35% of the mean. Uncertainty was ascribed to errors in remotely sensed burn area mapping, fuel load estimation, spatial variation of fire severity, errors in BEF for each fuel class and EFs. At the spatial scale of our study area, there were no uncertainties with the burnt area, vegetation structure or fuel type classification, and we used site-specific fuel load estimations used in our calculations, all of which would reduce the error associated with our fire emissions estimate. Russell-Smith et al. (2009) also reported low coefficients of variability (CV%) of for BEFs across fine, coarse and heavy fuel types for high severity fires, ranging from 0.3 to 11% and 2% CV for EFs for CH₄ and N₂O. Site specific sources of error include high spatial variability of on-site fuel loads which had a CV% of ~70% (Table 4) and uncertainty associated with the BEF we assumed for coarse and heavy fuel loads (0.9), which is higher than that derived for late dry season savanna fires (0.36, 0.31 respectively, Russell-Smith et al. 2009). This value was assumed as repeat burning of coarse and heavy fuels ensured ~10% of biomass remained as ash and charcoal at the CS site. This assumed BEF is also consistent with FullCAM (4.00.3) BEF of 0.98 for forest fire with 100% of trees killed, although this is setting is based on Surawski et al. (2012) who found little empirical evidence for BEF for stand-replacement fires. However, given the detailed on-site measurements of fuel load, error in our fire-derived emissions would be of the order of 20% or less.

5.0 Conclusions

While GHG emissions from savanna deforestation are dominated by debris burning, emissions from soil tillage and soil bed preparation is likely to be 20% of the total emission, suggesting satellite-based emissions based on oxidation of cleared vegetation alone do not capture all phases of

1 | LUC prior to cultivation. Savanna burning, using the area as defined in this study, was 1.5% of
2 | Australia's national GHG emissions and is of similar magnitude to emissions associated with
3 | historic savanna deforestation. However, for the deforestation scenario ~~as described in this study (an~~
4 | ~~addition of 311 000 ha of deforestation over five years), could increase~~ Australia's GHG emissions
5 | ~~could increase~~ by ~~between 5 to 10%~~ at least 3% per annum for the duration of the expansion,
6 | depending on the area ~~deforested~~ and deforestation rate. These are indicative estimates only, but
7 | suggest that the impacts of northern agricultural development will have ~~some~~ an impact on the
8 | national GHG budget and will need to be considered in northern land use decision making
9 | processes. These considerations are also particularly relevant given the emission reduction targets
10 | set by Australia following the 21st Conference of Parties to the UN Framework Convention on
11 | Climate Change (COP21 / CMP11) to reduce GHG emissions by 26 to 28% of 2005 by 2030.

12

1 Acknowledgements

2 Financial support for this study was provided by the Australian Research Council's Linkage
3 Project LP100100073 and Discovery Project DP0772981. Beringer is funded under the Australian
4 Research Council's Future Fellowship program (FT1110602). Support for flux data collection and
5 archiving was provided ~~throughby~~ Dr Peter Isaac of the Australian flux network, OzFlux
6 (www.ozflux.org.au), which is funded by the Australian Terrestrial Ecosystem Research Network
7 (TERN) ~~(http://www.tern.org.au, www.tern.org.au)~~. ~~We thank~~ Chris and Bridget Schulz
8 ~~for provided access to the~~ property ~~access~~ and field assistance throughout all phases of the land use
9 change we monitored. We are grateful for the technical expertise and field assistance of Matthew
10 Northwood and Michael Brand who maintained the eddy covariance tower ~~and soil gas exchange~~
11 ~~instruments. Thanks to,~~ Yan-Shih Lin, Amanda Lilleyman, and Allison O'Keefe ~~for provided~~ field
12 support during the intensive field campaigns. ~~We Thanks~~ also ~~thank Dr Peter Isaac of OzFlux for~~
13 ~~help with flux data processing and to~~ the Department of Environment for provision of savanna
14 specific deforestation GHG emissions data, 1990-2013 and the two reviewers who provided
15 constructive comments on the original manuscript.

16

References

- Adams, V. M. and Pressey, R. L.: Uncertainties around the Implementation of a Clearing-Control Policy in a unique catchment in Northern Australia: exploring equity issues and balancing competing objectives, *PLoS One*, 9(5), e9647, 2014.
- Aldrick, J. M., Robinson, C. S.: Report on the Land Units of the Katherine-Douglas Area, Northern Territory Service, Land Conservation Section of the Northern Territory Administration, Department of the Interior, Canberra, 1972.
- Anthoni, P. M., Knohl, A., Rebmann, C., Freibauer, A., Mund, M., Ziegler, W., Kolle, O. and Schulze, E.-D.: Forest and agricultural land-use-dependent CO₂ exchange in Thuringia, Germany, *Glob. Chang. Biol.*, 10(12), 2005–2019, doi:10.1111/j.1365-2486.2004.00863.x, 2004.
- Aubinet, M., Vesala, T. and Papale, D.: *Eddy Covariance: A Practical Guide to Measurement and Data Analysis*, Springer, Dordrecht., 2012.
- ~~Committee on Northern Australia: PIVOT NORTH Inquiry into the Development of Northern Australia: Final Report, Commonwealth of Australia, Canberra, 2014.~~
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S. and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nat. Clim. Chang.*, 2(3), 182–185, 2012.
- Beringer, J., Hutley, L. B., McHugh, I., Arndt, S. K., Campbell, D., Cleugh, H. A., Cleverly, J., de Dios, V., Eamus, D., Evans, B., Ewenz, C., Grace, P., Griebel, A., Haverd, V., Hinko-Najera, N., Huete, A., Isaac, P., Kanniah, K., Leuning, R., Liddell, M. J., Macfarlane, C., Meyer, W., Moore, C., Pendall, E., Phillips, A., Phillips, R. L., Prober, S., Restrepo-Coupe, N., Rutledge, S., Schroder, I., Silberstein, R., Southall, P., Sun, M., Tapper, N. J., van Gorsel, E., Vote, C., Walker, J. and Wardlaw, T.: An introduction to the Australian and New Zealand flux tower network - OzFlux, *Biogeosciences Discuss.*, 2016, 1–52, doi:10.5194/bg-2016-152, 2016.
- Beringer, J., Hutley, L. B., Tapper, N. J. and Cernusak, L. A.: Savanna fires and their impact on net ecosystem productivity in North Australia, *Glob. Chang. Biol.*, 13(5), 990–1004, doi:10.1111/j.1365-2486.2007.01334.x, 2007.
- Beringer, J., Hacker, J., Hutley, L. B., Leuning, R., Arndt, S. K., Amiri, R., Bannehr, L., Cernusak, L. A., Grover, S., Hensley, C., Hocking, D., Isaac, P., Jamali, H., Kanniah, K., Livesley, S., Neining, B., Paw, U., ~~K. T. P.~~, Sea, W., Straten, D., Tapper, N., Weinmann, R., Wood, S. and

1 Zegelin, S.: Special-Savanna patterns of energy and carbon integrated across the landscape, Bull.
2 Am. Meteorol. Soc., 92(11), 1467–+, doi:10.1175/2011bams2948.1, 2011.

3 Beringer, J., Hutley, L. B., Abramson, D., Arndt, S. K., Briggs, P., Bristow, M., Canadell, J. G.,
4 Cernusak, L. A., Eamus, D., Edwards, A. C., Evans, B. J., Fest, B., Goergen, K., Grover, S. P.,
5 Hacker, J., Haverd, V., Kanniah, K., Livesley, S. J., Lynch, A., Maier, S., Moore, C., Raupach, M.,
6 Russell-Smith, J., Scheiter, S., Tapper, N. J. and Uotila, P.: Fire in Australian savannas: from leaf to
7 landscape., Glob. Chang. Biol., 21(1), 62–81, doi:10.1111/gcb.12686, 2015.

8 Beringer, J., McHugh, I., Hutley, L. B., Isaac, P. and Kljun, N.: Dynamic INtegrated Gap-filling
9 and partitioning for OzFlux (DINGO), Biogeosciences Discuss, 1-36, doi:10.5194/bg-2016-188,
10 2016.

11 Bradshaw, C. J. A.: Little left to lose: deforestation and forest degradation in Australia since
12 European colonization, J. Plant Ecol., 5(1), 109–120, doi:10.1093/jpe/rtr038, 2012.

13 Brown, S.: Estimating biomass and biomass change of tropical forest: a primer, FAO Forestry Paper
14 134, Rome, www.fao.org/docrep/w4095e/w4095e00.htm, 1997.

15 Committee on Northern Australia: PIVOT NORTH Inquiry into the Development of Northern
16 Australia: Final Report, Commonwealth of Australia, Canberra, 2014.

17 Commonwealth of Australia: Australian Vegetation Attribute Manual: National Vegetation
18 Information System, Version 6.0, Canberra, 2003.

19 Commonwealth of Australia: State and Territory Greenhouse Gas Inventories 2013, Canberra,
20 20152015a.

21 Commonwealth of Australia: Carbon Credits (Carbon Farming Initiative—Emissions Abatement
22 through Savanna Fire Management) Methodology Determination
23 (www.legislation.gov.au/Details/F2015L00344), Department of the Environment, Canberra, 2015b.

24 Cook, G. D.: Historical perspectives on land use development in northern Australia, in Northern
25 Australia Land and Water Science Review, Office of North Australia, Commonwealth of Australia,
26 Canberra, 2009.

27 ~~Donohue, R. J., Roderick, M. L., McVicar, T. R. and Farquhar, G. D.: Impact of CO₂ fertilization~~
28 ~~on maximum foliage cover across the globe's warm, arid environments, *Geophys. Res. Lett.*,~~
29 ~~40(12), 3031–3035, doi:10.1002/grl.50563, 2013.~~

30 Eamus, D., Chen, X., Kelley, G. and Hutley, L. B.: Root biomass and root fractal analyses of an
31 open Eucalyptus forest in a savanna of north Australia, Aust. J. Bot., 50(1), 31–41,

doi:10.1071/bt01054, 2002.

Fearnside, P. M., Righi, C. A., Graça, P. M. L. de A., Keizer, E. W. H., Cerri, C. C., Nogueira, E. M. and Barbosa, R. I.: Biomass and greenhouse-gas emissions from land-use change in Brazil's Amazonian 'arc of deforestation': The states of Mato Grosso and Rondônia, *For. Ecol. Manage.*, 258(9), 1968–1978, doi:http://dx.doi.org/10.1016/j.foreco.2009.07.042, 2009.

Ferreira, M. E., Ferreira, L. G., Miziara, F. and Soares-Filho, B. S.: Modeling landscape dynamics in the central Brazilian savanna biome: future scenarios and perspectives for conservation, *J. Land Use Sci.*, 8(4), 403–421, doi:10.1080/1747423X.2012.675363, 2013.

Ferreira, M. E., Ferreira, L. G., Latrubesse, E. M. and Miziara, F.: Considerations about the land use and conversion trends in the savanna environments of Central Brazil under a geomorphological perspective, *J. Land Use Sci.*, 11(1), 33–47, 2016.

Fisher, R. and Edwards, A. C.: Fire extent and mapping: procedures, validation and website application, in *Carbon Accounting and Savanna Fire Management*, edited by B. P. Murphy, A. C. Edwards, C. P. Meyer, and J. Russell-Smith, pp. 57–72, CSIRO Publishing, Melbourne, Australia., 2015.

Fox, I. D., Neldner, V. J., Wilson, G. W. and Bannink, P. J.: *The Vegetation of the Australian Tropical Savannas*, Brisbane, 2001.

Galford, G. L., Melillo, J. M., Kicklighter, D. W., Mustard, J. F., Cronin, T. W., Cerri, C. E. P. and Cerri, C. C.: Historical carbon emissions and uptake from the agricultural frontier of the Brazilian Amazon, *Ecol. Appl.*, 21(3), 750–763, doi:10.1890/09-1957.1, 2011.

Galford, G. L., Soares-Filho, B. and Cerri, C. E. P.: Prospects for land-use sustainability on the agricultural frontier of the Brazilian Amazon, *Philos. Trans. R. Soc. London B Biol. Sci.*, 368(1619), 2013.

Grace, J., San, J. J., Meir, P., Miranda, H. S., Montes, R. A., José, J. S., Meir, P., Miranda, H. S. and Montes, R. A.: Productivity and carbon fluxes of tropical savannas, *J. Biogeogr.*, 33(3), 387–400, doi:10.1111/j.1365-2699.2005.01448.x, 2006.

Grover, S. P. P., Livesley, S. J., Hutley, L. B., Jamali, H., Fest, B., Beringer, J., Butterbach-Bahl, K. and Arndt, S. K.: Land use change and the impact on greenhouse gas exchange in north Australian savanna soils, *Biogeosciences*, 9(1), 423–437, doi:10.5194/bg-9-423-2012, 2012.

Grundy, M. J., Bryan, B. A., Nolan, M., Battaglia, M., Hatfield-Dodds, S., Connor, J. D. and Keating, B. A.: Scenarios for Australian agricultural production and land use to 2050, *Agric. Syst.*,

1 142, 70–83, doi:10.1016/j.agry.2015.11.008, 2016.

2 Haverd, V., Raupach, M. R., Briggs, P. R., J. G. Canadell., Davis, S. J., Law, R. M., Meyer, C. P.,
3 Peters, G. P., Pickett-Heaps, C., and Sherman, B.: The Australian terrestrial carbon budget,
4 Biogeosciences, 10, 851–869, doi:10.5194/bg-10-851-2013, 2013.

5 Henry, B. K., Butler, D. and Wiedemann, S. G.: A life cycle assessment approach to quantifying
6 greenhouse gas emissions from land-use change for beef production in eastern Australia, Rangel. J.,
7 37 (3), 273–283, doi.org/10.1071/RJ14112, 2015.

8 Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le
9 Quéré, C. and Ramankutty, N.: Carbon emissions from land use and land-cover change,
10 Biogeosciences, 9(12), 5125–5142, doi:10.5194/bg-9-5125-2012, 2012.

11 Hurst, D. F., Griffith, D. W. T. and Cook, G. D.: Trace gas emissions from biomass burning in
12 tropical Australian savannas, J. Geophys. Res., 99(D8), 16441, doi:10.1029/94JD00670, 1994.

13 Hutley, L. B. and Beringer, J.: Disturbance and climatic drivers of carbon dynamics of a north
14 Australian tropical savanna, in Ecosystem Function in Savannas: Measurement and Modeling at
15 Landscape to Global, edited by M. J. Hill and N. P. Hanan, CRC Press, Boca Raton., 2011.

16 Hutley, L. B., Leuning, R., Beringer, J. and Cleugh, H. A.: The utility of the eddy covariance
17 techniques as a tool in carbon accounting: tropical savanna as a case study, Aust. J. Bot., 53(7),
18 663–675, doi:10.1071/bt04147, 2005.

19 Hutley, L. B., Beringer, J., Isaac, P. R., Hacker, J. M. and Cernusak, L. A.: A sub-continental scale
20 living laboratory: Spatial patterns of savanna vegetation over a rainfallprecipitation gradient in
21 northern Australia, Agric. For. Meteorol., 151(11), 1417–1428,
22 doi:10.1016/j.agrformet.2011.03.002, 2011.

23 IPCC: Revised 1996 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National
24 Greenhouse Gas Inventories, 3 vols, IPCC/OECD/IEA: Paris., France, 1997.

25 IPCC: Guidelines for National Greenhouse Gas Inventories, Paris., 2003.

26 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the
27 Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Eds. Stocker, T.F., D.
28 Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M.
29 Midgley, Cambridge University Press: Cambridge, United Kingdom, 2013

30 Isaac, P.-R., Cleverly, J., McHugh, I., van Gorsel, E., Ewenz, C. and Beringer, J. and McHugh, I.:
31 The .: OzFlux Data: network ~~data path: integration~~ from collection to curation, Biogeosciences

1 | Discuss., 2016, [1–41, doi:10.5194/bg-2016-189](https://doi.org/10.5194/bg-2016-189), 2016..

2 | Isbell, R.: The Australian Soil Classification. In: Australian Soil and Land Survey Handbook,
3 | CSIRO Publishing, Melbourne, 2002.

4 | Jamali, H., Livesley, S. J., Dawes, T. Z., Hutley, L. B. and Arndt, S. K.: Termite mound emissions
5 | of CH₄ and CO₂ are primarily determined by seasonal changes in termite biomass and behaviour,
6 | *Oecologia*, 167(2), 525–534, doi:10.1007/s00442-011-1991-3, 2011a.

7 | Jamali, H., Livesley, S. J., Grover, S. P., Dawes, T. Z., Hutley, L. B., Cook, G. D. and Arndt, S. K.:
8 | The Importance of Termites to the CH₄ Balance of a Tropical Savanna Woodland of Northern
9 | Australia, *Ecosystems*, 14(5), 698–709, doi:10.1007/s10021-011-9439-5, 2011b.

10 | Jarecki, M. K. and Lal, R.: Crop Management for Soil Carbon Sequestration, *Cr. Rev. Plant Sci.*,
11 | 22, 471–502, doi:10.1080/07352680390253179, 2003.

12 | Johnson, A.: North Australian Grassland Fuel Guide – Sturt Plateau and Victoria River District,
13 | Cooperative Research Centre for Tropical Savannas Management, Charles Darwin University,
14 | Darwin, Australia, 2001.

15 | Jones, D., Wang, W. and Fawcett, R.: High-quality spatial climate data sets for Australia, *Aust.*
16 | *Meteorol. Oceanogr. J.*, 58, 233–248, 2009.

17 | Lal, R. and Follett, R. F.: Soils and climate change, in *Soil Carbon Sequestration and the*
18 | *Greenhouse Effect*, edited by R. Lal and R. F. Follett, pp. 1–23, Soil Science Society of America,
19 | Maddison., 2009.

20 | Lambin, E. F., Gibbs, H. K., Ferreira, L., Grau, R., Mayaux, P., Meyfroidt, P., Morton, D. C., Rudel,
21 | T. K., Gasparri, I. and Munger, J.: Estimating the world’s potentially available cropland using a
22 | bottom-up approach, *Glob. Environ. Chang.*, 23(5), 892–901, doi:10.1016/j.gloenvcha.2013.05.005,
23 | 2013.

24 | Landsberg, J., Gillieson, D. and Salt, D.: Trees in savanna landscapes: finding the balance, David
25 | Gillison, Cooperative Research Centre for Tropical Savannas Management, Charles Darwin
26 | University, Darwin, Australia, 2011.

27 | Lapola, D. M., Martinelli, L. A., Peres, C. A., Ometto, J. P. H. B., Ferreira, M. E., Nobre, C. A.,
28 | Aguiar, A. P. D., Bustamante, M. M. C., Cardoso, M. F., Costa, M. H., Joly, C. A., Leite, C. C.,
29 | Moutinho, P., Sampaio, G., Strassburg, B. B. N. and Vieira, I. C. G.: Pervasive transition of the
30 | Brazilian land-use system, *Nat. Clim. Chang.*, 4(1), 27–35, doi.org/10.1038/nclimate2056, 2014.

31 | Law, R. and Garnett, S. T.: Understanding carbon in the Northern Territory: an analysis of future

land use scenarios using the national carbon accounting tool, Cooperative Research Centre for Tropical Savannas Management, Darwin, [2009](#).

Law, R. and Garnett, S. T.: Mapping carbon in tropical Australia: Estimates of carbon stocks and fluxes in the Northern Territory using the national carbon accounting toolbox, *Ecol. Manag. Restor.*, 12(1), 61–68, doi:10.1111/j.1442-8903.2011.00566.x, 2011.

Lawes, M. J., Greiner, R., Leiper, I. A., Ninnis, R., Pearson, D. and Boggs, G.: The effects of a moratorium on land-clearing in the Douglas-Daly region, Northern Territory, Australia, *Rangel. J.*, 37(4), 399–408, doi.org/10.1071/RJ15014, 2015.

[Le Quéré, C., Moriarty, R., Andrew, R. M., Peters, G. P., Ciais, P., Friedlingstein, P., Jones, S. D., Sitch, S., Tans, P., Arneeth, A., Boden, T. A., Bopp, L., Bozec, Y., Canadell, J. G., Chini, L. P., Chevallier, F., Cosca, C. E., Harris, I., Hoppema, M., Houghton, R. A., House, J. I., Jain, A. K., Johannessen, T., Kato, E., Keeling, R. F., Kitidis, V., Klein Goldewijk, K., Koven, C., Landa, C. S., Landschützer, P., Lenton, A., Lima, I. D., Marland, G., Mathis, J. T., Metzl, N., Nojiri, Y., Olsen, A., Ono, T., Peng, S., Peters, W., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Salisbury, J. E., Schuster, U., Schwinger, J., Séférian, R., Segschneider, J., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Werf, G. Lewis, D.: Stray Creek Catchment Vegetation Survey Northern Territory, Department of Infrastructure Planning and Environment, Northern Territory, Palmerston, 2005.](#)

[R., Viovy, N., Wang, Y.-P., Wanninkhof, R., Wiltshire, A. and Zeng, N.: Global carbon budget 2014, Earth Syst. Sci. Data, 7\(1\), 47–85, doi:10.5194/essd-7-47-2015, 2015.](#)

[Macfarlane, C., Hoffman, M., Eamus, D., Kerp, N., Higginson, S., McMurtrie, R. and Adams, M.: Estimation of leaf area index in eucalypt forest using digital photography, Agric. For. Meteorol., 143\(3–4\), 176–188, doi:10.1016/j.agrformet.2006.10.013, 2007.](#)

Mayocchi, C. L. and Bristow, K. L.: Soil surface heat flux: some general questions and comments on measurements, *Agric. For. Meteorol.*, 75, 43–50, 1995.

[Meyer, C. P., Cook, G. D., Reisen, F., Smith, T. E. L., Tattaris, M., Russell-Smith, J., Maier, S. W., Yates, C. P. and Wooster, M. J.: Direct measurements of the seasonality of emission factors from savanna fires in northern Australia, J. Geophys. Res. Atmos., 117\(D20\), n/a-n/a, doi:10.1029/2012JD017671, 2012.](#)

Murphy, B. P., Lehmann, C. E. R., Russell-Smith, J. and Lawes, M. J.: Fire regimes and woody biomass dynamics in Australian savannas, *J. Biogeogr.*, 41(1), 133–144, doi:10.1111/jbi.12204, 2014.

1 Murphy, B. P., Edwards, A. C., Meyer, C. P. (Mick) and Russell-Smith, J.: Carbon Accounting and
2 Savanna Fire Management, CSIRO Publishing, Melbourne, 2015a.

3 Murphy, B. P., Liedloff, A. C. and Cook, G. D.: Does fire limit tree biomass in Australian
4 savannas? *Int. J. Wildl. Fire*, 24(1), 1, doi:10.1071/WF14092, 2015b.

5 Neill, C., Piccolo, M. C., Cerri, C. C., Steudler, P. a. and Melillo, J. M.: Soil solution nitrogen
6 losses during clearing of lowland Amazon forest for pasture, *Plant Soil*, 281(1–2), 233–245,
7 doi:10.1007/s11104-005-4435-1, 2006.

8 Petheram, C., McMahon, T. A. and Peel, M. C.: Flow characteristics of rivers in northern Australia:
9 Implications for development, *J. Hydrol.*, 357(1-2), 93–111, doi:10.1016/j.jhydrol.2008.05.008,
10 2008.

11 Petheram, C., McMahon, T. A., Peel, M. C. and Smith, C. J.: A continental scale assessment of
12 Australia's potential for irrigation, *Water Resour. Manag.*, 24(9), 1791–1817, doi:10.1007/s11269-
13 009-9525-z, 2009.

14 Petheram, C., Gallant, J., Wilson, P., Stone, P., Eades, G., Roger, L., Read, A., Tickell, S.,
15 Commander, P., Moon, A., McFarlane, D. and Marvanek, S. .: Northern rivers and dams: a
16 preliminary assessment of surface water storage potential for northern Australia, Canberra, 2014.

17 ~~Poulter, B., Frank, D., Ciais, P., Myneni, R. B., Andela, N., Bi, J., Broquet, G., Canadell, J. G.,~~
18 ~~Chevallier, F., Liu, Y. Y., Running, S. W., Sitch, S. and van der Werf, G. R.: Contribution of semi-~~
19 ~~arid ecosystems to interannual variability of the global carbon cycle, *Nature*, 509(7502), 600–603,~~
20 ~~doi:10.1038/nature13376, 2014.~~

21 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C.,
22 Buchmann, N., Gilmanov, T., Granier, A., Grunwald, T., Havrankova, K., Ilvesniemi, H., Janous,
23 D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F.,
24 Ourcival, J.-M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G.,
25 Vaccari, F., Vesala, T., Yakir, D. and Valentini, R.: On the separation of net ecosystem exchange
26 into assimilation and ecosystem respiration: review and improved algorithm, *Glob. Chang. Biol.*,
27 11(9), 1424–1439, doi:10.1111/j.1365-2486.2005.001002.x, 2005.

28 Richards, G. P. and Evans, D. M. W.: Development of a carbon accounting model (FullCAM Vers.
29 1.0) for the Australian continent, *Aust. For.*, 67, 277–283, doi:10.1080/00049158.2004.10674947,
30 2004.

31 Rose, D.: The dynamics of coarse ~~Woody Debris~~woody debris in the mesic tropical savannas of the
32 Darwin region: abundance, decomposition and consumption by fire, Honours Thesis, Charles

1 Darwin University, 2006.

2 Russell-Smith, J., Whitehead, P. J., Cooke, P. M. and Yates, C. P.: Challenges and opportunities for
3 fire management in fire-prone northern Australia, in Culture, Ecology and Economy of Fire
4 Management in North Australian Savannas, edited by J. Russell-Smith, P. J. Whitehead, and P. M.
5 Cooke, pp. 1–23, CSIRO Publishing, Melbourne, 2009a.

6 Russell-Smith, J., Murphy, B. P., Meyer, C. P. (Mick), Cook, G. D., Maier, S., Edwards, A. C.,
7 Schatz, J. and Brocklehurst, P.: Improving estimates of savanna burning emissions for greenhouse
8 accounting in northern Australia: limitations, challenges, applications, *Int. J. Wildl. Fire*, 18(1), 1,
9 doi:10.1071/WF08009, 2009b.

10 [Russell-Smith, J., Cook, G. D., Cooke, P. M., Edwards, A. C., Lendrum, M., Meyer, C. P. \(Mick\)](#)
11 [and Whitehead, P. J.: Managing fire regimes in north Australian savannas: applying Aboriginal](#)
12 [approaches to contemporary global problems, *Front. Ecol. Environ.*, 11\(s1\), e55–e63,](#)
13 [doi:10.1890/120251, 2013.](#)

14 Sakai, R. K., Fitzjarrald, D. R., Moraes, O. L. L., Staebler, R. M., Acevedo, O. C., Czikowsky, M.
15 J., da Silva, R., Brait, E. and Miranda, V.: Land-use change effects on local energy, water, and
16 carbon balances in an Amazonian agricultural field., *Glob. Chang. Biol.*, 10(5), 895–907,
17 doi:10.1111/j.1529-8817.2003.00773.x, 2004.

18 Sanderman, J., Farquharson, R. and Baldock, J. A.: Soil carbon sequestration potential: a review for
19 Australian agriculture. Report to the Australian Department of Climate Change, Adelaide, 2010.

20 Steffan, W. and Hughes, L.: The Critical Decade 2013: Climate change science, risks and response,
21 Climate Commission Secretariat, Department of Industry, Innovation, Climate Change, Science,
22 Research and Tertiary Education, Canberra, 2013.

23 [Surawski, N. C., Sullivan, A. L., Roxburgh, S. H. and Cook, G. D.: Review of FullCAM forest fire](#)
24 [event parameters with recommendations supported by a literature review, *CSIRO Ecosystems*](#)
25 [Science, Canberra, 2012.](#)

26 Thackway, R.A.: A vegetation fuel type map for Australia's northern savannas, Department of the
27 Environment, Canberra, 2014.

28 Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., Banger, K., Tao, B., Pan, S., Liu, M.,
29 Zhang, C., Bruhwiler, L. and Wofsy, S.: Global methane and nitrous oxide emissions from
30 terrestrial ecosystems due to multiple environmental changes, *Ecosyst. Heal. Sustain.*, 1(1), art4,
31 doi:10.1890/EHS14-0015.1, 2015.

[Tubiello, F. N., Salvatore, M., Ferrara, A. F., House, J., Federici, S., Rossi, S., Biancalani, R.,](#)
[Condor Golec, R. D., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo, P., Schmidhuber,](#)
[J., Sanz Sanchez, M. J., Srivastava, N. and Smith, P.: The contribution of agriculture, forestry and](#)
[other land use activities to global warming, 1990–2012, Glob. Chang. Biol., 21\(7\), 2655–2660,](#)
[doi:10.1111/gcb.12865, 2015.](#)

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton,
 D. C., DeFries, R. S., Jin, Y. and van Leeuwen, T. T.: Global fire emissions and the contribution of
 deforestation, savanna, forest, agricultural, and peat fires (1997–2009), Atmos. Chem. Phys.,
 10(23), 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

[Walsh, D., Russell-Smith, J. and Cowley, R.: Fire and carbon management in a diversified](#)
[rangelands economy: research, policy and implementation challenges for northern Australia,](#)
[Rangel. J., 36\(4\), 313–322, doi:http://dx.doi.org/10.1071/RJ13122, 2014.](#)

[Weitz, A. M., Veldkamp, E., Keller, M., Neff, J. and Crill, P. M.: Nitrous oxide, nitric oxide, and](#)
[methane fluxes from soils following clearing and burning of tropical secondary forest, J. Geophys.](#)
[Res., 103\(D21\), 28047, doi:10.1029/98JD02144, 1998.](#)

[Whitehead, P. J., Russell-Smith, J. and Yates, C.: Fire patterns in north Australian savannas:](#)
[extending the reach of incentives for savanna fire emissions abatement, Rangel. J., 36\(4\), 371–388,](#)
[2014.](#)

[Williams, R. J., Zerihun, A., Montagu, K. D., Hoffman, M., Hutley, L. B. and Chen, X.: Allometry](#)
[for estimating aboveground tree biomass in tropical and subtropical eucalypt woodlands: towards](#)
[general predictive equations, Aust. J. Bot., 53\(7\), 607, doi:10.1071/BT04149, 2005.](#)

[Woinarski, J., Mackey, B., Nix, H. and Traill, B.: The nature of Northern Australia: it's natural](#)
[values, ecological processes and future prospects, ANU E Press, Canberra., 2007.](#)

Woldendorp, G., Keenan, R. J., Barry, S. and Spenser, R. D.: Analysis of sampling methods for
 coarse woody debris, For. Ecol. Manage., 198, 133–148, 2004.

Wolf, S., Eugster, W., Potvin, C. and Buchmann, N.: Strong seasonal variations in net ecosystem
 CO₂ exchange of a tropical pasture and afforestation in Panama, Agric. For. Meteorol., 151(8),
 1139–1151, doi:10.1016/j.agrformet.2011.04.002, 2011.

Zenone, T., Chen, J., Deal, M. W., Wilske, B., Jasrotia, P., Xu, J., Bhardwaj, A. K., Hamilton, S. K.
 and Philip Robertson, G.: CO₂ fluxes of transitional bioenergy crops: effect of land conversion
 during the first year of cultivation, GCB Bioenergy, 3(5), 401–412, doi:10.1111/j.1757-
 1707.2011.01098.x, 2011.

1 Zona, D., Janssens, I. A., Aubinet, M., Gioli, B., Vicca, S., Fichot, R. and Ceulemans, R.: Fluxes of
2 the greenhouse gases (CO₂, CH₄ and N₂O) above a short-rotation poplar plantation after
3 conversion from agricultural land, *Agric. For. Meteorol.*, 169, 100–110, 2013.

4 |
5

Table captions

Table 1 Site characteristics for the uncleared savanna (UC) and cleared (CS) sites. Site soil orders are given as per Isbell (2002) with savanna vegetation classified using Fox et al. (2001). Fire frequency was estimated from fire mapping taken from the North Australian Fire Information system (NAFI, www.firenorth.org.au) for 2000-2013-2012. The fire frequency estimate for the CS site excluded the debris fires in August 2012. Basal area and stem density is provided for all woody stems >52 cm DBH at both sites-~~CS sites~~. Mean site LAI for the UC is taken from Hutley et al. 2011 and for the CS site, ~~it~~ was estimated from canopy hemispherical photos, see text for details.

Table 2 Characteristics of land conversion phases during the 668 day observation period at the savanna clearing site (CS). Also given are the canopy heights following LUC phases and flux ~~tower~~instrument heights that were adjusted following clearing, burning and then soil preparation phases.

Table 3 ~~Mean rainfall, Cumulative precipitation and mean~~ NEE, Re and GPP ($\text{Mg C ha}^{-1} \text{ month}^{-1}$) for each of the ~~land-use change~~LUC phases at the CS site as measured by the ~~eddy covariance~~flux tower. ~~Comparative~~These fluxes are given for the UC site for ~~the identical~~these same periods. One-way ANOVA was used to test for differences between mean daily NEE for each LUC phase with significantly different means labelled with an asterisk. On the days of ignition during the debris burning phase, flux data at the CS site were excluded. Integrated fluxes are given for the post-clearing period (507 days) and the entire observation period (668 days) for both sites in Mg C ha^{-1} .

~~Table 4~~Table 4 Measured fuel loads, assumed burning efficiencies (BEF), carbon contents, N:C ratio and emissions factors (EF) used to estimate GHG emissions from the burning of the post-deforestation fine, coarse and heavy fuel debris. Emission factors, carbon content and C:N ratio were assumed for the vegetation fuel type woodland savanna with mixed grass (code hWMI) as given in the Emissions Abatement through Savanna Fire Management methodology (Commonwealth of Australia, 2015b), available at www.legislation.gov.au/Details/F2015L00344 and Meyers et al. 2012.

Table 5 Greenhouse gas emissions for 1990-2013 from prescribed savanna burning and savanna deforestation at catchment (Douglas-Daly River), state/territory (Northern Territory savanna area) and regional scales (north Australian savanna area, Fig. 1). For savanna burning, burnt area and associated mean annual emissions (\pm ~~sd~~SD) are given for both reportable non-CO₂ (CH₄, N₂O) and total emissions (CO₂, CH₄ and N₂O). For the identical areas as used for savanna burning, mean

1 | annual GHG emissions from deforestation (\pm ~~sd~~SD) are given. For the Douglas-Daly River
2 | catchment, deforestation area was taken from Lawes et al. (2015) and combined with deforestation
3 | emissions from the CS site. Deforestation emissions (1990-2013) for the NT and the north
4 | Australian savanna area are taken from the State and Territory Greenhouse Gas Inventories
5 | (Commonwealth of Australia, ~~2015~~-2015a). In bold text are the emissions associated with the
6 | current deforestation rate plus expanded deforestation areas as identified by Petheram et al. (2014),
7 | which are combined with emissions from the CS site to give an upscaled estimate of potential
8 | emissions with agricultural development at the three spatial scales.

9

10 | Table 1

11

<i>Site</i>	<i>UC</i>	<i>CS</i>
Location	14°09'33.12"S, 131°23'17.16"E	14°33'48.71"S, 132°28'39.47"E
Soils	Red Kandosol (Blain)	Red Kandosol (Blain)
	Woodland savanna	Woodland savanna
Vegetation type;	<u>Savanna woodland with mixed grasses</u>	<u>Savanna woodland with mixed grasses</u>
dominant species	Map unit D4 . <i>E. tetradonta</i> , <i>C. latifolia</i> , <i>Terminalia grandiflora</i> , <i>Sorghum spp</i> , <i>Heteropogon triticeus</i>	Map unit D4 . <i>E. tetradonta</i> , <i>Erythrophleum chlorostachys</i> , <i>Corymbia</i> <i>bleeseri</i> , <i>Sorghum spp</i> , <i>H. triticeus</i>
Map unit area (km²)	59,986	59,986
Fire frequency (y⁻¹)	Error!	
Hyperlink reference not valid.)	<u>0.323</u>	<u>0.07</u>
Basal area (m² ha⁻¹)	8.3	6.8
Canopy height (m)	16.4	14.2
Above-ground biomass (Mg C ha⁻¹)	<u>30.6 ± 9.2</u>	<u>26.2 ± 7.0</u>
Stem density (ha⁻¹)	330 ± 58	643 ± 102
Overstorey LAI (wet/dry)	n/a / 0.8	0.9 / 0.5
MODIS LAI (wet/dry)	1.5 / 0.9	1.6 / 1.0
MAP (mm)	1372 ^a / 1180 ^b	1107 ^c
Max T_{air} (°C)	37.5 (Oct) / 31.2 (Jun)	37.5 (Oct) / 29.7 (Jun)
Min T_{air} (°C)	23.8 (Jan) / 12.6 (Jul)	25.0 (Nov) / 13.7 (Jul)

12 | ^aOn-site observations, 2007-2012, ^bgridded ~~rainfall~~precipitation (AWAP, 1970-2012), ^cTindal BoM station (14.52S, 132.38E, data from 1985-2013).

13

14

15 Table 2

16

17

<i>Season</i>	<i>Period</i>	<i>LULUC phases</i>	<i>Canopy height (m)</i>	<i>TowerInstru ment height (m)</i>
Late dry season	Sep - Oct 2011	Intact savanna	16	23 21.5
Wet season pre-clearing	Oct 2011 - Feb 2012	Intact savanna	16	23 21.5
Wet season clearing	Mar - May 2012	Savanna cleared deforested using bulldozers, followed by debris decomposition, understory grass germination	2-4 3	7
Dry season pre- burn	May - Aug 2012	Vegetation debris curing, understorey grass growth	2- 3	7
Debris burning	Aug 2012	Debris and grasses burnt then , soil ripped to 60 cm to remove roots, roots and remaining debris stockpiled, re-burnt	0 2	7
Dry season post--burn	Aug - Nov 2012	Vegetation removed, followed by grass Grass and shrubs germination and resprouting Site cultivation removal Removal remaining below-ground biomass. Rainfall stimulation of Wet season rains stimulates grass, shrubs and trees germination growth, shrub re-sprouting and growth	0 -1	7
Early wet season	Nov 2012 - Jan 2013		1- 3	7
Wet season	Jan - Mar 2013	All regenerated vegetation removed via , soil cultivation bed preparation	0	3
Dry season	Apr - Jul 2013	Soil cultivation in stages	0	3

18

19

20 | Table 3
21
22

<i>LULUC phases</i>	<i>Phase numb er (d)</i>	<i>Period (d)</i>	<i>CS</i>				<i>UC</i>			
			<i>Rainfall</i>	<i>NEE</i>	<i>Re</i>	<i>GPP</i>	<i>Rainfall</i>	<i>NEE</i>	<i>Re</i>	<i>GPP</i>
			<i>all</i>	<i>E</i>			<i>all</i>			
			(mm)	(Mg C ha ⁻¹ month ⁻¹)			(mm)	(Mg C ha ⁻¹ month ⁻¹)		
Intact canopy cover	<u>1</u>	161	736.6	-0.23	1.5 ₇	-1.79	1076.8	-0.25	1.45	-1.70
Clearing event	<u>2</u>	4	59.4	0.23 [*]	1.9 ₅	-1.73	59.8	0.38 [*]	1.80	-1.50
Wet-dry debris curing, decomposition	<u>3</u>	59	143.2	0.98 [*] -	1.3 ₉	-0.41	412.0	0.32 ^{**}	1.53	-1.22
Dry season pre-burn	<u>4</u>	94	0	0.34 [*] -	0.5 ₇	-0.23	2.4	0.15 ^{**}	0.94	-0.79
Fire emissions late dry	<u>5</u>	22	0	0.90 [*] -	0.7 ₆	0.0	0.0	-0.01 ^{**}	0.71	-0.72
Dry season post-burn	<u>6</u>	67	2.2	0.31 [*] -	0.3 ₇	-0.06	64.4	-0.28	0.64	-0.91
Early wet regrowth	<u>7</u>	80	361.0	0.03 [*] -	0.9 ₉	-0.96	345.8	-0.32	1.80	-2.12
Wet season site prep	<u>8</u>	91	701.7	0.62 [*] -	0.9 ₉	-0.37	914.4	-0.20 ^{**}	1.67	-1.88
Dry season final bed prep. and cultivation	<u>9</u>	90	0	0.29 [*] -	0.3 ₂	-0.02	10.8	0.06 ^{**}	0.91	-0.85

				<u>(Mg C ha⁻¹)</u>				<u>(Mg C ha⁻¹)</u>		
	Total post-clearing	507	1267. 5	7.2 [*] -	12. 8	-5.6	1809. 6	-0.78 ^{**}	20.7	-21.5
	Total all phases	668	2004. 1	6.0 [*] -	21. 2	-15.2	2886. 4	-2.1 ^{**}	28.5	-30.6
23										
24										

25 | *Denotes significantly different mean NEE at the 5% level, ** significant at 1%.
26 |

27 Table 4

28

<u>Fuel type</u>	<u>Fuel load</u> <u>(Mg C ha⁻¹)</u>	<u>BEF</u>	<u>Carbon</u> <u>content</u>	<u>N:C ratio</u>	<u>EF CO₂</u>	<u>EF CH₄</u>	<u>EF N₂O</u>	<u>Emissions</u> <u>(Mg CO₂-e ha⁻¹)</u>			
								<u>CO₂</u>	<u>CH₄</u>	<u>N₂O</u>	<u>Total</u>
<u>Fine</u>	<u>1.1 ± 0.70</u>	<u>0.95</u>	<u>0.46</u>	<u>0.0096</u>	<u>0.97</u>	<u>0.0031</u>	<u>0.0075</u>	<u>3.9</u>	<u>0.1</u>	<u>0.04</u>	<u>4.0</u>
<u>Coarse</u>	<u>0.5 ± 1.0</u>	<u>0.9</u>	<u>0.46</u>	<u>0.0081</u>	<u>0.92</u>	<u>0.0031</u>	<u>0.0075</u>	<u>1.5</u>	<u>0.0</u>	<u>0.01</u>	<u>1.6</u>
<u>Heavy - AGB</u>	<u>26.2 ± 7.0</u>	<u>0.9</u>	<u>0.46</u>	<u>0.0081</u>	<u>0.87</u>	<u>0.01</u>	<u>0.0036</u>	<u>75.2</u>	<u>7.9</u>	<u>0.32</u>	<u>83.4</u>
<u>Heavy - CWD</u>	<u>1.4 ± 0.6</u>	<u>0.9</u>	<u>0.46</u>	<u>0.0081</u>	<u>0.87</u>	<u>0.01</u>	<u>0.0036</u>	<u>4.0</u>	<u>2.7</u>	<u>0.11</u>	<u>28.5</u>
<u>Heavy - BGB</u>	<u>9.0 ± 2.4</u>	<u>0.9</u>	<u>0.46</u>	<u>0.0081</u>	<u>0.87</u>	<u>0.01</u>	<u>0.0036</u>	<u>25.7</u>	<u>0.0</u>	<u>0.02</u>	<u>4.4</u>
<u>Total</u>								<u>110.2</u>	<u>11.1</u>	<u>0.50</u>	<u>121.9</u>

29

30

Table 5

<i>Savanna region</i>	<i>Savanna burning</i>			<i>Savanna deforestation</i>			
	Burnt area ^a	Emissions non-CO ₂ ^a	Emissions total ^a	Deforestation area	Emissions total	Expanded deforestation area ^d	Expanded emissions total ^d
	(ha y ⁻¹)	(Gg CO ₂ -e y ⁻¹)	(Gg CO ₂ -e y ⁻¹)	(ha y ⁻¹)	(Gg CO ₂ -e y ⁻¹)	(ha)	(Gg CO ₂ -e y ⁻¹)
Douglas-Daly River catchment	2,482,100 ±490,400	577 ±124	14,270 ±3064	1275 ±454 ^b	163 ±162 ^b	20,000	781756
Northern Territory	13,419,410 ±487,300	3,490 ±922	86,255 ±22,880	1,717 ±611 ^c	398 ±128 ^c	114,500	5,9493,413
North Australian	32,249,254 ±11,176,004	6,740 ±1,729	166,586 ±42,725	78,605 ±34,976 ^c	16,161 ±5601 ^c	311,000	28,26024,393

^aBurnt area and emissions data estimated using the on-line Savanna Burning Abatement Tool (SAVBat2), 1990-2013. These emissions are CH₄ and N₂O only.

^bDeforestation area data taken from Lawes et al. (2015), upscaled using the emissions from the CS site from this study, 1990-2013

^cDeforestation area and emissions data taken from the State and Territory Greenhouse Gas Inventories (Commonwealth of Australia, ~~2015~~2015a), 1990-2013

^dExpanded deforestation area data taken from catchments as identified by Petheram et al. (2014), upscaled using the GHG emissions from the CS site from this study and added to historic emissions

40 **Figure captions**

41

42 Figure 1 Location of the uncleared site (UC) and the cleared savanna (CS) sites south of Darwin,
43 Northern Territory. The inset figure shows the distribution of the savanna biome across northern
44 Australia as defined by Fox et al. (2001).

45 ~~Figure 2 Energy balance closure plots using 30 minute flux data from the CS and UC sites. For the~~
46 ~~CS site, data were grouped according to the LUC phase periods using pooled day time and night~~
47 ~~time data. Plots are arranged clockwise and labelled ‘Phase 1’ through to ‘Phase 7’, with a further~~
48 ~~plot of all data pooled from all phases (‘All Phases’). The final plot is the long term UC site is also~~
49 ~~plotted using 30 minute data for 2007–2015. Statistical data for each regression is given for each~~
50 ~~plot.~~

51 ~~Figure 3~~ Figure 2 Comparative meteorology and fluxes for the uncleared (UC) and cleared savanna
52 CS sites prior to the clearing event. Data spans the late dry season (September 2011) through to the
53 mid-wet season prior to the clearing event of 2–6 March 2012. Plots include a) daily
54 ~~rainfall~~ precipitation (black bars UC site, grey bars CS site), mean daily T_{air} (black lines UC, grey
55 CS), b) mean daily VPD (dashed lines; black UC, grey CS), c) interpolated 8-day MODIS LAI
56 (black UC, grey CS), d) NEE (black UC, grey CS) partitioned into R_e (red UC, pink CS) and GPP
57 (dark green UC, pale green CS).

58 ~~Figure 4-a~~ 3a) Daily ~~rainfall~~ precipitation and b) diurnal patterns of NEE at the CS site for the week
59 prior to the clearing event of 2–6 March 2012 (vertical bar) and three weeks post-clearing.

60 ~~Figure 5~~ 4 Cumulative NEE from the CS (red line) and UC sites (black line) for each land use phase
61 (see Table 2 for details) over the entire observational period, September 2011 to July 2013. The UC
62 site is a long-term savanna site of the Australian flux network (OzFlux, see Beringer et al.
63 ~~2016~~) 2016a) and using the sites’ 8-year flux record (2007–2013), the long-term cumulative mean
64 NEE is plotted for each land use phase of (grey line; \pm 95% CI). The dashed line indicates zero net
65 CO_2 flux.

66

67 **Plate caption**

68


69 Plate 1 Key LUC phases associated with: a) the clearing event, Phase 3; b) debris burning of the
70 cured grass, litter and woody fuels following the 5 month curing period, Phase 5; c) stockpiling and
71 ignition of remaining unburnt debris and d) post-fire site preparation with all biomass consumed,
72 Phase 9.

73

74

75

76 **Figure 1**




77

78

79

80 Figure 1




81

82

83 Figure 2

84

85




86 Figure 3


87

88

89



91 Figure 4



92

93 |

