

# The nitrogen, carbon and greenhouse gas budget of a grazed, cut and fertilised temperate grassland

Stephanie K. Jones<sup>1,2</sup>, Carole Helfter<sup>1</sup>, Margaret Anderson<sup>1</sup>, Mhairi Coyle<sup>1</sup>, Claire Campbell<sup>1</sup>, Daniela Famulari<sup>1</sup>, Chiara Di Marco<sup>1</sup>, Netty van Dijk<sup>1</sup>, Y. Sim Tang<sup>1</sup>, Cairistiona F.E. Topp<sup>2</sup>, Ralf Kiese<sup>3</sup>, Reimo Kindler<sup>4</sup>, Jan Siemens<sup>5</sup>, Marion Schrumpp<sup>6</sup>, Klaus Kaiser<sup>7</sup>, Eiko Nemitz<sup>1</sup>, Peter E. Levy<sup>1</sup>, Robert M. Rees<sup>2</sup>, Mark A. Sutton<sup>1</sup>, Ute M. Skiba<sup>1</sup>

1) Centre for Ecology and Hydrology, Edinburgh, Bush Estate, Penicuik, Midlothian EH26 QB, UK

10 2) Scotland's Rural College, King's Buildings, West Mains Road, Edinburgh, EH9 3JG, UK

3) Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Kreuzackbahnstr. 19, 82467 Garmisch-Partenkirchen, D

4) Chair of Waste Management and Environmental Research, Technische Universität Berlin, Franklinstr. 29, 10587 Berlin, D

15 5) Institute of Crop Science and Resource Conservation, Soil Science, Universität Bonn, Nussallee 13, 53115 Bonn, D

6) Max Plank Institute for Biogeochemistry, Hans-Knöll-Str. 10, 07745 Jena, D

7) Martin Luther University Halle-Wittenberg, Von-Seckendorff-Platz 3, 06120 Halle (Saale), D

20 *Correspondence to:* Stephanie K. Jones (Stephanie.Jones@sruc.ac.uk)

## 30 Abstract

Intensively managed grazed grasslands in temperate climates are globally important environments for the exchange of the greenhouse gases (GHGs) carbon dioxide (CO<sub>2</sub>), nitrous oxide (N<sub>2</sub>O) and methane (CH<sub>4</sub>). We assessed the N and C budget of a mostly grazed, occasionally cut, and fertilized grassland in SE Scotland by measuring or modelling all relevant imports and exports to the field as well as changes in soil C and N stocks over time. The N budget was dominated by import from inorganic and organic fertilisers (21.9 g N m<sup>2</sup> yr<sup>-1</sup>) and losses from leaching (5.3 g N m<sup>2</sup> yr<sup>-1</sup>), N<sub>2</sub> emissions and NO<sub>x</sub> and NH<sub>3</sub> volatilisation (6.4 g N m<sup>2</sup> yr<sup>-1</sup>). The efficiency of N use by animal products (meat and wool) averaged 10% of total N input. On average over nine years (2002-2010) the balance of N fluxes suggested that 6.6 ± 4.4 g N m<sup>-2</sup> y<sup>-1</sup> (mean ± confidence interval at p > 0.95) were stored in the soil. The largest component of the C budget was the net ecosystem exchange of CO<sub>2</sub> (NEE), at an average uptake rate of 218 ± 155 g C m<sup>-2</sup>y<sup>-1</sup> over the nine years. This sink strength was offset by carbon export from the field mainly as harvest (48.9 g C m<sup>2</sup> yr<sup>-1</sup>) and leaching (16.4 g C m<sup>2</sup> yr<sup>-1</sup>). The other export terms, CH<sub>4</sub> emissions from the soil, from manure applications and enteric fermentation were negligible and only contributed to 0.02-4.2 % of the total C losses. Only a small fraction of C was incorporated into the body of the grazing animals. Inclusion of these C losses in the budget resulted in a C sink strength of 163 ± 140 g C m<sup>-2</sup>y<sup>-1</sup>. On the contrary, soil stock measurements taken in May 2004 and May 2011 indicated that the grassland sequestered N in the 0-60 cm soil layer at 4.51 ± 2.64 g N m<sup>-2</sup> y<sup>-1</sup> and lost C at a rate of 29.08 ± 38.19 g C m<sup>-2</sup> y<sup>-1</sup>, respectively. Potential reasons for the discrepancy between these estimates are probably an underestimation of C losses, especially from leaching fluxes as well as from animal respiration. The average greenhouse gas (GHG) balance of the grassland was -366 ± 601 g CO<sub>2</sub> eq m<sup>-2</sup> y<sup>-1</sup> and strongly affected by CH<sub>4</sub> and N<sub>2</sub>O emissions. The GHG sink strength of the NEE was reduced by 54% by CH<sub>4</sub> and N<sub>2</sub>O emissions. Enteric fermentation from the ruminating sheep proved to be an important CH<sub>4</sub> source, exceeding the contribution of N<sub>2</sub>O to the GHG budget in some years.

55 *Keywords:* grassland, carbon stocks, carbon sequestration, nitrogen cycling, budget, greenhouse gases

## Introduction

Managed grasslands cover an estimated 26 % of earth's land surface (FAOstat, 2008). The  
60 impact of reactive nitrogen (Nr) losses, carbon (C) sequestration and greenhouse gas (GHG) emissions  
(CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) from these grasslands is therefore of global importance and will become even  
more relevant in the future as increased standards of living in developed countries are expected to  
result in a rapid growth of livestock farming (Caro et al, 2014). Carbon and N cycles in grasslands are  
intricately linked and tightly coupled in extensively managed low N grasslands, with sinks and sources  
65 in equilibrium. Converting such systems to intensively managed N fertilised grasslands in the short  
term may increase the soil organic carbon (SOC) pool from decomposed plant litter and root material  
as well as through rhizodeposition (Rees et al., 2005) until a new equilibrium is reached (Soussana and  
Lemaire, 2014). In the case of the long term Broadbalk experiment in the UK, this equilibrium was  
achieved after 50 years (Powelson et al, 2011). After the conversion to intensive N management, the  
70 tight coupling of the N and C cycles becomes disrupted, leading to emissions of N<sub>2</sub>O and CH<sub>4</sub> at rates  
which may outweigh the benefits of C sequestration. Several studies indicate that managed grasslands  
can sequester C (Kim et al., 1992; Jones et al., 2006; Soussana et al., 2004; Ammann et al., 2007)  
however, uncertainties are high (Janssens et al., 2003). On the contrary, Smith (2014) concluded from  
long-term experiments and chronosequence studies, that changes in agronomic management may lead  
75 to short-term C sequestration, but in the long-term, under constant management and environmental  
conditions, C stocks are relatively stable. In a grassland ecosystem the C balance is determined by the  
net biome exchange (the difference between total C input and losses). In managed grassland  
ecosystems exports through biomass harvesting, the addition of organic manures (from organic  
fertiliser additions and animal excretion) as well as CO<sub>2</sub> and CH<sub>4</sub> losses from animal respiration and  
80 enteric fermentation can make significant contributions to the C budget.

Nutrient budgets are a valuable tool to summarise and understand nutrient cycling in  
agroecosystems and to assess their impact on the environment. As imbalances are not sustainable in the  
long term, N and C budgets can be used as indicators and regulatory policy instruments for nutrient  
management in order to reduce losses and increase efficiency. So far, different Nr species have been  
85 looked at in separate studies according to their form and impact. Few studies have attempted to

calculate N budgets from managed grasslands (e.g. Chen et al., 2004; Ammann et al., 2009; Kramberger et al., 2015), whereas C budgets have been assessed more often and are available for various ecosystems (e.g. Aubinet et al., 2000; Soussana et al., 2007; Ammann et al., 2007, Rytter et al. 2015). To calculate the total C and N budget of an ecosystem all import and export processes have to be assessed by measuring or estimating all imports and exports to an ecosystem. The other method is to measure differences in N and C stocks in the soil over time. This approach has the advantage that it requires the measurement of only a single component of the system. However, a large number of samples are needed at an interval of more than 5 years before detectable changes may be statistically significant (Smith, 2004). Moreover this approach does not provide any information about the different processes leading to the final budget.

In this study we assessed the C and N budget from an intensively managed grassland in Southern Scotland using both approaches. Here we report one of the most detailed analyses of C and N fluxes from a grassland ecosystem over 9 years (2002-2010). This study allowed an analysis of the importance of common grassland management practices such as cutting for silage, grazing of cattle and sheep at different stocking densities, N input by inorganic and organic fertiliser applications, as well as different weather conditions on the N, C and GHG balance. The data provide a unique overview of research undertaken within three European projects GREENGRASS (Soussana et al., 2007), CarboEurope (Schulze et al., 2009) and NitroEurope (Sutton et al., 2007; Skiba et al., 2009).

## 2. Methods

### 2.1 Site description

The experimental site, Easter Bush, is located in South East Scotland, 10 km South of Edinburgh (03°02'W, 55°52' N, 190 m a.s.l) with a mean annual rainfall of  $947 \pm 234$  mm and mean annual temperature of  $9.0 \pm 0.4$  °C (2002-2010). The field (5.424 ha) has been under permanent grassland management for more than 20 years with a species composition of >99% perennial ryegrass (*Lolium perenne*) and < 0.5% with clover (*Trifolium repens*). The soil type is an imperfectly drained Macmerry soil series, Rowanhill soil association (Eutric Cambisol) with a pH of 5.1 (in H<sub>2</sub>O) and a clay fraction of

20-26%. The ground water table was assumed to be at 0.85 m depth on average and the main rooting zone extends down to 0.31 m below soil surface.

115

## 2.2 Grassland management

The grassland was grazed rotationally throughout the experimental period by heifers in calf, ewes and lambs at different stocking densities (Table 1 and Figure 1a). . Livestock units used for heifers, ewes and lambs were 0.75, 0.10 and 0.04, respectively (1 livestock unit has a standard live weight of 600 kg head<sup>-1</sup>; SAC, 1995). Lambs were present on the field from April to September only. The grass was cut for silage only in the first two years, on the 1<sup>st</sup> of June and 8<sup>th</sup> of August 2002 and on the 29<sup>th</sup> of May 2003. Ammonium nitrate fertiliser was applied to the field 3-4 times per year, usually between March and July (56 kg N ha<sup>-1</sup> application<sup>-1</sup> on average). In 2008 an additional fifth mineral N application was added, using urea instead of ammonium nitrate fertiliser. Organic manure was applied on the 28<sup>th</sup> of September 2004 and 27<sup>th</sup> of March 2005 as cattle slurry, using a vacuum slurry spreader, ( Table 4 and 5 and in Fig. 1 a) and b).

125

## 2.3. Annual budget calculations

We assessed the N and C budget by measuring or estimating the import and export of all relevant fluxes to and from the grassland field on an annual basis. Throughout the manuscript all fluxes are presented following the sign convention used in micrometeorology; fluxes from the ecosystem to the atmosphere are positive (exported from the field), while negative values indicate fluxes from the atmosphere to the ecosystem (imported to the field). We set the system boundary for inputs and exports of N and C by the field perimeters (covering an area of 5.4 ha). The balance of all imports and exports results in the observed changes of N and C on this field over time.

130

135

The change in the N balance ( $\Delta N$ ) over time ( $\Delta t$ ) of our grassland ecosystem can be written as:

$$\begin{aligned} \Delta N / \Delta t = & \text{FN}_{\text{org fert.}} + \text{FN}_{\text{synt fert.}} + \text{FN}_{\text{N}_2 \text{ (biol. fixation)}} + \text{FN}_{\text{dep.}} + \\ & \text{FN}_{\text{harvest}} + \text{FN}_{\text{animal}} + \text{FN}_{\text{leaching}} + \text{FN}_{\text{NH}_3/\text{NO}_x \text{ (fert., manure, animal)}} + \\ & \text{FN}_{\text{NO}_x \text{ (soil)}} + \text{FN}_{\text{N}_2\text{O}} + \text{FN}_{\text{N}_2 \text{ (denitr.)}} \end{aligned} \quad (1)$$

140

N imports include the addition of N from organic and inorganic fertiliser ( $FN_{\text{org fert.}} + FN_{\text{synt fert.}}$ ), the fixation of  $N_2$  through biological fixation ( $FN_{N_2 \text{ (biol. fixation)}}$ ) and the deposition of  $NH_3$ ,  $HNO_3$ ,  $NH_4^+$ ,  $NO_3^-$  from dry, and  $NH_4^+$  and  $NO_3^-$  from wet deposition (summarised as  $FN_{\text{dep.}}$ ). Exports include the N lost from plant biomass at cuts for silage ( $FN_{\text{harvest}}$ ), the off-take of N in meat (including bones) and wool from animals ( $FN_{\text{animal}}$ ), the loss of organic and inorganic dissolved N through leaching ( $FN_{\text{leaching}}$ ), the  $NH_3$  and  $NO_x$  emissions from volatilisation of inorganic and organic fertiliser spreading as well as from animal excretion ( $FN_{NH_3/NO_x(\text{fert., manure, animal})}$ ), the emission of  $NO_x$  from the soil ( $FN_{NO_x(\text{soil})}$ ), the emission of  $N_2O$  from the soil ( $FN_{N_2O}$ ) and the loss of  $N_2$  from total denitrification ( $FN_{N_2(\text{denitr.})}$ ).

The change in the C balance ( $\Delta C$ ) over time equals the net biome production (NBP) and can be written for our site as:

$$\Delta C/\Delta t = NBP = FC_{CO_2} + FC_{\text{org fert}} + FC_{\text{animal}} + FC_{CH_4} + FC_{\text{leaching}} + FC_{\text{harvest}} \quad (2)$$

$FC_{CO_2}$  represents the net ecosystem exchange (NEE) of  $CO_2$  and  $FC_{\text{org fert}}$  is the C input through manure application. Carbon input from animal excretion was not included in the budget as it was assumed to be recycled C from plant and soil uptake.  $FC_{\text{animal}}$  includes the C off-take through animal weight increase and wool production. As grazing cows were heifers in calf and ewes milk was consumed by their lambs, there was no C off-take through milk to be considered. Methane emissions from enteric fermentation by ruminants, animal excretion and manure application as well as  $CH_4$  fluxes from the soil are included in  $FC_{CH_4}$ .  $FC_{\text{leaching}}$  is the C lost through dissolved organic and inorganic C and dissolved  $CH_4$  leaching and  $FC_{\text{harvest}}$  represents the C lost from the system through plant biomass export from harvests (cut for silage). Carbon emissions from farm operations (i.e. tractor emissions) or off farm emissions (i.e. fertiliser manufacture) are not included in the C budget.

Details of methods to quantify each N and C budget component, as listed in Eq. (1) and (2), are described under Sect. 2.4 to 2.11 and in the Appendix. Some budget components were measured

170 throughout the 9 years presented, while others were only measured in some years or not at all. Missing data were derived from the literature, models or averages from available data from other years.

The annual net GHG exchange (NGHGE) was calculated from annual NEE ( $FC_{CO_2}$ ),  $CH_4$  ( $FC_{CH_4}$ ) and  $N_2O$  ( $FN_{N_2O}$ ) fluxes using global warming potentials (GWPs) at the 100-year time horizon  
175 (1 for  $CO_2$ , 298 for  $N_2O$  and 25 for  $CH_4$ , IPCC, 2013):

$$NGHGE = (FC_{CO_2}) + FC_{CH_4} * k_{CH_4} + FN_{N_2O} * k_{N_2O} \quad (3)$$

Where;

180  $k_{CH_4} = 9.09$ , since 1 kg  $CH_4$ -C corresponds to 9.09 kg  $CO_2$ -C

$k_{N_2O} = 127$ , since 1 kg  $N_2O$ -N corresponds to 127 kg  $CO_2$ -C

185 In addition the net annual greenhouse gas balance (NGHGB) was calculated by including the loss of C through animal meat and wool production, harvest off take, C leaching and input by organic fertiliser application:

$$NGHGB = NGHGE + FC_{org\ fert} + FC_{animal} + FC_{leach} + FC_{harvest} \quad (4)$$

190

#### **2.4 Nitrogen and carbon import by fertiliser and manure ( $FN_{synt\ fert} + FN_{org\ fert} + FC_{org\ fert}$ )**

Mineral fertiliser was applied by a spreader as either ammonium nitrate or urea. Six month old cattle slurry was spread by a vacuum slurry tanker. Three samples from the slurry tank were taken at each application and analysed for ammoniacal nitrogen ( $NH_3$  and  $NH_4^+$ ), dry matter content, total N, total C,  
195 pH and nitrate. The total N and C import to the field by the slurry was calculated by the volume of the slurry applied and the N and C analyses of the slurry.

#### **2.5 Nitrogen and carbon export by harvest ( $FN_{harvest} + FC_{harvest}$ )**

The farmer estimated a forage harvest of 15 t fresh weight (FW)  $ha^{-1} y^{-1}$  at the first cut and 10 t FW  $ha^{-1}$   
200  $y^{-1}$  at the second cut of a year, based on the plant height at the field at the time of cutting and

information from harvested plot experiments. As there were two cuts in 2002 and one cut in 2003 the estimated harvest was 25 t FW ha<sup>-1</sup> y<sup>-1</sup> for 2002 and 10 t FW ha<sup>-1</sup> y<sup>-1</sup> for 2003. A subsample of harvested vegetation was collected and dried at 80°C for plant N and C analysis using a Carbo-Erba/400 automated N and C analyser.

205

## **2.6 Nitrogen and carbon export by meat and wool ( $FN_{\text{animal}} + FC_{\text{animal}}$ )**

It was estimated by the farmer that heifers increased in weight by 0.8kg per day (starting weight of 450 kg). The ewe weight was assumed to be constant (60 kg). They were fed extra protein (standard cake concentrate) to reduce weight loss during lactation., whereas lambs were brought to the field at a weight of 5 kg and removed when they reached a weight of 45 kg. All animals were weight before they came onto the field at the beginning of the season and again at the end of the season. The total meat export, which includes bones, was calculated from the daily weight increase of heifers and lambs multiplied by the animal number per day. To calculate the N and C export from meat we assumed a N content of meat of 3.5 % and a C content of meat of 21 % (Flindt, 2002), a N content of bones of 7 % and a C content of bones of 20 % (Marchand, 1842), assuming a total bone content of 20 % for sheep (Lambe et al., 2002) and 14 % for heifers (Navajas et al., 2010) . Ewes were sheared annually in June, yielding an estimated 2.5 kg of wool per sheep. Wool N and C export was calculated from wool production multiplied by the average sheep number in June, assuming a N and C content of wool of 16.5 and 50 %, respectively (Roche, 1995).

210

215

220

## **2.7 Nitrogen and carbon leaching ( $FN_{\text{leaching}} + FC_{\text{leaching}}$ )**

Two sets of ten glass suction cups (pore size <1 µm, ecoTech, Bonn, Germany) for soil water and four Teflon suction cups (ecoTech, Bonn, Germany) for soil gas collection were installed in August 2006. One set was located on a slope, another on a hollow. For the budget calculations we only used results from the slope location as the hollow location was frequently water logged. Suction cups were installed horizontally from a soil pit beneath the A horizon (30 cm depth) and at 90cm depth and connected to 2-l glass bottles in an insulated aluminium box placed into the soil pit. Samples were collected every two to three weeks. For further details and description of dissolved organic and inorganic C (DIC, DOC) and

225

dissolved CH<sub>4</sub> analysis see Kindler et al. (2011). Dissolved inorganic and organic N (DIN, DON) and total N (TN) concentrations in leachate water were analysed by colorimetric analysis (San<sup>++</sup>, Automated Wet Chemistry Analyzer - Continuous Flow Analyzer (CFA), Skalar, The Netherlands). Leachate C and N concentrations were measured from October 1<sup>st</sup> 2006 - March 30<sup>th</sup> 2008. Dissolved C and N were calculated by multiplying concentrations of DIC, DOC and dissolved CH<sub>4</sub> or DIN and DON respectively, with leachate volume. The latter was derived from a soil water model based on balancing daily precipitation and evaporation considering the water holding capacity of the soil (Kindler et al., 2011). For the remaining years N was simulated using the LandscapeDNDC model (Haas et al., 2013), with the model tested and validated with comprehensive measured data. For C leaching linear regression models describing the relationship between calculated C leaching fluxes and leachate volume for the measurement period (DOC;  $y = 0.0186x - 0.0695$ ,  $R^2 = 0.8663$ , DIC;  $y = 0.021x - 0.0008$ ,  $R^2 = 0.8056$  and dissolved CH<sub>4</sub>:  $y = 0.0019x - 0.0135$ ,  $R^2 = 0.7623$ ) were used to extrapolate to the remaining years.

## 2.8 Gaseous N fluxes

### 2.8.1 N deposition ( $FN_{dep}$ )

#### Wet N deposition

Wet N deposition was determined from daily samples collected by an automatic precipitation sampler (Eigenbrodt<sup>®</sup> precipitation collector 181/KS, Königsmoor, D) at Auchencorth Moss (3°14'35W, 55°47'34 N), 17 km south west of Easter Bush (Skiba et al., 2013, McKenzie et al., 2015). Precipitation samples were analysed for NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> by ion chromatography (Methrom AG, Switzerland). Annual inorganic N deposition at this site was then adjusted to annual rainfall amounts measured at Easter Bush.

#### Dry N deposition

Cumulative monthly concentrations of gaseous and aerosol N species (NH<sub>3</sub>, HNO<sub>3</sub>, particulate NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>) were collected from another field, about 300m distance from our study field, using a DELTA

system (DENuder for Long Term Atmospheric) (Sutton et al., 2001). N dry deposition fluxes were calculated using the average flux from four different inferential models, as described in detail by Flechard et al. (2011).

260

### **2.8.2 N<sub>2</sub>O fluxes (FN<sub>N2O</sub>)**

From June 2002 to July 2003 N<sub>2</sub>O fluxes were measured continuously by eddy covariance (EC) using an ultra-sonic anemometer coupled with a Tunable Diode Laser absorption spectrometer (TDL) at a frequency of 10 Hz. Details for the gap filling method of the N<sub>2</sub>O–EC data are described in Jones et al., 2011. From August 2006 to November 2010 N<sub>2</sub>O fluxes were measured using manual closed static chambers (Clayton et al., 1994, Skiba et al., 2013). Samples were analysed for N<sub>2</sub>O using a Hewlett Packard 5890 series II gas chromatograph (Agilent Technologies, Stockport, UK), fitted with an electron capture detector (detection limit: N<sub>2</sub>O < 33 ppbV). Fluxes were measured weekly and more frequently during fertilisation. Cumulative fluxes were calculated by gapfilling data for missing days using linear interpolation and summing up all gapfilled data over each calendar year. For the periods where no N<sub>2</sub>O fluxes were measured (January -May 2002, July 2003-March 2004, May 2004-July 2006) fluxes were simulated by LandscapeDNDC (Haas et al., 2013). LandscapeDNDC was tested in detail with available data on plant growth, soil temperature, moisture, inorganic soil N concentration, NO and N<sub>2</sub>O, which resulted in general good agreement of simulations and measurements.

265

270

275

### **2.8.3 NO<sub>x</sub> fluxes (FN<sub>NO<sub>x</sub>(soil)</sub>)**

NO<sub>x</sub> fluxes from the soil were only measured for a short period (June 2009-August 2010). The NO<sub>x</sub> fluxes were measured using an autochamber system described in detail by Butterbach-Bahl et al. (1997). Measurements were made 4 times per day, every 6 hours for an 8 min period per chamber. We used simulated data from Landscape DNDC for years where no NO<sub>x</sub> fluxes were measured.

280

### **2.8.4 NH<sub>3</sub> + NO<sub>x</sub> volatilisation (FN<sub>NH3/NO<sub>x</sub> (fert.,manure, animal)</sub>)**

The fraction of nitrogen that volatilises as NH<sub>3</sub> and NO<sub>x</sub> from applied synthetic fertiliser or cattle slurry application and animal excretion was estimated to be 10% and 20% of total N applied, respectively

285 (IPCC, 2006b). The animal excretion amount was estimated in accordance with the IPCC Guidelines (IPCC, 2006a), for details, see Appendix.

### 2.8.5 N<sub>2</sub> emission by total denitrification ( $FN_{N_2(\text{denitr.})}$ )

290 Di-nitrogen (N<sub>2</sub>) emissions resulting from total denitrification in the soil was not measured in our experiment. We therefore used the N<sub>2</sub> emission rates from LandscapeDNDC simulations.

### 2.8.6 Biological N<sub>2</sub> fixation ( $FN_{N_2(\text{biol. fixation})}$ )

295 The species composition was measured once in 2002 and at monthly intervals in 2003 by the visual estimation method (Braun-Blanquet, 1964), where 50 quadrates of 0.25 m<sup>2</sup> were randomly thrown into the field. As the legume fraction (*Trifolium repens*) was smaller than 0.5% at each measuring point we assumed the nitrogen fixation through plants to be zero.

### 2.9 Exchange of CO<sub>2</sub> ( $FC_{CO_2}$ )

300 NEE was measured continuously from 1.January 2002 till 31.December 2010 by an eddy covariance system consisting of a fast response 3D ultrasonic anemometer (Metek USA-1, Metek GmbH, Elsmhorn, Germany) and a fast closed path CO<sub>2</sub>-H<sub>2</sub>O analyser (LI-COR 7000 infra-red gas analyzer (IRGA), LI-COR, Lincoln, NE, USA). NEE is the arithmetic sum of the gross primary production (GPP) and total ecosystem respiration (TER). Flux partitioning of measured NEE into GPP and TER was calculated by the same online tool used for gapfilling. In this flux partitioning approach, daytime 305 TER is obtained by extrapolation of a night time parameterisation of NEE on air temperature and GPP is the difference between ecosystem respiration and NEE. Contrarily to unmanaged ecosystem, TER at our site also includes the respiratory loss of CO<sub>2</sub> by grazing animals. Net primary production (NPP), which represents the annual plant growth (difference between GPP and autotrophic respiration) was calculated as 50% of GPP (Amthor 2000, Zhang et al., 2009).

310

### 2.10 Methane fluxes ( $FC_{CH_4}$ )

Methane fluxes from the soil were measured with closed static chambers simultaneously with the N<sub>2</sub>O measurements. The same GC was fitted with a flame injection detector (detection limit: CH<sub>4</sub> < 70 ppbV). The minimal detectable flux was 17 ng CH<sub>4</sub>-C m<sup>-2</sup> s<sup>-1</sup>. Fluxes were measured weekly and more frequently at fertiliser events. As measured soil CH<sub>4</sub> fluxes were close to zero and did not vary significantly between months, we calculated CH<sub>4</sub> for months where no CH<sub>4</sub> fluxes were measured (January-May 2002, July 2003-March 2004, May 2004-July 2006), as an average monthly cumulative flux from other years.

Methane emissions from grazing animals, i.e. animal excretion and enteric fermentation, were estimated following the IPCC Tier 2 methodology (IPCC, 2006a: Stewart et al., 2009). For details, see Appendix. Methane emissions from slurry applications were assumed to be small. As no chamber measurements were conducted at the time of slurry spreading, the emissions were estimated as 0.07 % of the applied assuming that emissions were comparable to those in a related study (Jones et al., 2006), where CH<sub>4</sub> was measured from chambers after slurry application on a nearby field in 2002 and 2003.

## 2.11 VOC

Fluxes of non-methane volatile organic compounds (VOC) were not measured. We assumed similar VOC emissions to those reported by Davison et al. (2008) for an intensively managed grassland in Switzerland, where the daily average flux of methanol, acetaldehyde and acetone over 3 days after cutting were 21.1, 5.1. and 2.6 nmol m<sup>-2</sup> s<sup>-1</sup>, respectively. Based on those values, annual VOC emissions from our field were estimated to be in the order of 0.03% of the annual C offtake in harvest and 0.08 % of annual C off-take by grazing animals. We therefore assumed VOC emissions to be negligible and did not account for them in the C balance.

## 2.12 Soil N and C measurements

Total N and C content of the soil were measured in May 2004 and May 2011. One hundred soil cores with an inner diameter of 8.7 (2004) and 8.3 cm (2009), both corers from Eijkelkamp Agrisearch Equipment BV, Giesbeek, The Netherlands) were collected along a regular grid with a distance of 10 m between sampling points on both occasions. The soil sampling grid covered the main footprint area of

340 the site, not the entire field. Cores were separated into layers of 0-5, 5-10, 10-20, 20-30, 30-40, 40-50  
and 50-60 cm. Coarse stones of a diameter > 4 mm and roots of a diameter >1mm were removed from  
the samples prior to drying at 40 °C. Stone and root samples were air-dried separately. Then, soil  
samples were sieved to < 2 mm. Particles > 2 mm were combined with the coarse stones. Dry weights  
of roots and combined stone fractions were determined. Total N and C concentrations in < 2 mm soil  
345 separates were determined using dry combustion (VarioMax, Elementar Analysensysteme GmbH,  
Hanau, Germany). As the site contains no carbonates, total C was assumed to equal organic C. As bulk  
density varies spatially and over time (e.g. through compaction by livestock), the soil N and C content  
per unit ground area to a fixed depth will also change, without any change in the mass fraction of N and  
C in dry soil. Therefore, total N and C stocks were calculated on an equivalent soil mass (ESM) basis,  
350 so that comparisons between years were valid (see Gifford and Roderick, 2003, Wendt and Hauser,  
2013). A cubic polynomial was fitted to the data, to predict cumulative N and C with cumulative soil  
mass in the profile. A soil mass of 800 kg m<sup>-2</sup> was used (Table 7), which corresponds to approximately  
60-cm depth, which was the depth of the corer. Uncertainty in the estimates of stock change was based  
on the prediction intervals in the cubic polynomial at a soil mass of 800 kg m<sup>-2</sup>.

355

### **2.13 Ancillary measurements**

Soil temperature and volumetric soil moisture were continuously recorded at four depths (3.5, 7.5, 15  
and 30 cm) by temperature probes (temperature probe 107, Campbell Scientific, Loughborough, UK)  
and TDR probes (TDR 100, Campbell Scientific, Loughborough, UK), respectively, the latter installed  
360 in June 2002. Rain was measured by a tipping bucket rain gauge, while air temperature and relative  
humidity were measured by an integrated humidity and temperature transmitter (HUMITTER<sup>®</sup>, Vaisala  
Ltd, Suffolk, UK).

### **2.14 Statistical and uncertainty analysis**

365 Random error was determined as 2σ-standard error (95% confidence) of the overall mean according to  
Gaussian statistics. The confidence intervals for group means were used to establish whether or not  
differences were significantly different from zero. Linear correlations between C and N inputs and

outputs were calculated by calendar year. For systematic errors the uncertainty range of measurements as well as of parameterisations and literature based estimates was estimated according to expert judgment. To calculate the combined effect of systematic uncertainties of each flux component on the C and N budget simple Gaussian error propagation rules were used, details are provided in Table S1 (Supplementary material). Confidence intervals are given at the 95% confidence level.

A more detailed description of the methods are provided in the Appendix.

### 3. Results

#### 3.1 Climate and management

The meteorological conditions exhibited substantial inter-annual variability in the study period 2002-2010 (Table 2 and Fig. 2). Annual rainfall ranged from 575 mm to 1238 mm with highest monthly rainfalls of 280 mm month<sup>-1</sup> in September 2002. Lowest annual reported rainfall was in 2010; this low value was caused by a gap in data from January-March, due to snowfall. Average annual air temperature ranged from 8.3 to 9.6 °C with highest daily air temperatures of 30.4 °C in August 2005 and lowest in December 2010 at -10.3 °C. Highest average monthly air temperatures were measured in July 2006 at 17°C and lowest monthly average air temperatures at 2°C in November 2009. In 2003 the highest average annual temperature (9.6° C) and lowest annual rainfall (680 mm) were measured with a correspondingly low annual soil water content of 31 %. The duration of the growing season was defined per calendar year as the period bounded by the first and last 5 consecutive days with mean daily air temperature  $\geq 5$  °C. The length of the growing season (LGS) varied between 151 days (2006) and 242 days (2009) (Table 2).

Livestock stocking density exhibited both intra- and inter-annual variability. The average annual stocking density was lowest in 2002 and 2003 at 0.27 LSU ha<sup>-1</sup> y<sup>-1</sup> and 0.54 LSU ha<sup>-1</sup> y<sup>-1</sup>, respectively (Table 1), which were the years where the grass was cut for silage and no lambs were present in the field. In 2007, 2008, 2009 and 2010 no heifers were present in the field. Highest annual average stocking density occurred in 2004 and 2007 at 0.99 LSU ha<sup>-1</sup> y<sup>-1</sup> and 0.91 LSU ha<sup>-1</sup> y<sup>-1</sup>, respectively. Maximum monthly stocking density occurred in September 2006 at 13.8 LSU ha<sup>-1</sup>, while interim

periods with no grazing at all were observed in all years (Fig. 1a). Mineral N fertiliser was applied split into 3 to 5 applications per year, ranging from 2.5 to 9.6 g N m<sup>-2</sup> application<sup>-1</sup> (Fig. 1b). Organic manure was applied in 2004 and 2005 as cattle slurry, spread at a rate of 6.9 and 15.8 g N m<sup>-2</sup> application<sup>-1</sup>, respectively, which resulted in a C input of 55.4 and 171.8 g C m<sup>-2</sup> application<sup>-1</sup>, respectively (Fig. 1b and c). The grass was only cut in 2002 and 2003. Harvested biomass in 2002 and 2003 ranged from 2.60 to 3.75 t DW ha<sup>-1</sup> cut<sup>-1</sup> which resulted in an N off-take ranging from 1.7 to 4.7 g N m<sup>-2</sup> cut<sup>-1</sup> and a C removal from the field ranging from 113.1 to 169.5 g C m<sup>-2</sup> cut<sup>-1</sup> (Fig. 1c).

### 3.2. N budget

In our grassland system the N balance is the difference between the N input through fertiliser and atmospheric deposition and the N output through harvest, animal export, leaching and gaseous emissions. The total resulting balance over the nine years, derived from flux calculations and estimations, showed that N was stored at an average rate of  $-6.6 \pm 4.4$  g N m<sup>-2</sup> y<sup>-1</sup> ( $p < 0.05$ ). From 2003 to 2010, N was stored at a rate of  $-1.9$  to  $-17.2$  g N m<sup>-2</sup> y<sup>-1</sup>, whilst in 2002 N was lost at a rate of 6.4 g N m<sup>-2</sup> y<sup>-1</sup> (Table 4). The major N input consisted of inorganic fertiliser, ranging from  $-11$  to  $-25.9$  g N m<sup>-2</sup> y<sup>-1</sup>, averaging at  $-19.4$  g N m<sup>-2</sup> y<sup>-1</sup>, while N deposition represented only between 1.9 and 5.9% of the total N input. During the years where N was stored, a significant positive correlation between total N input from fertiliser and N storage was observed ( $R^2 = 0.55$ ). Largest losses resulted from leaching at an average rate of  $5.34 \pm 3.4$  g N m<sup>-2</sup> y<sup>-1</sup> and were estimated to be highest in 2002 at 14.9 g N m<sup>-2</sup> y<sup>-1</sup> and lowest in 2003 at 0.09 g N m<sup>-2</sup> y<sup>-1</sup>. The total N off take through meat and wool ranged from 0.15-3.12 g N m<sup>-2</sup> y<sup>-1</sup>, while the total annual N offtake from harvest was 5.0 g N m<sup>-2</sup> y<sup>-1</sup> in 2002 and 4.68 g N m<sup>-2</sup> y<sup>-1</sup> in 2003. Amongst gaseous exchanges, highest losses were estimated from N<sub>2</sub> emissions, averaging at 2.76 g N m<sup>-2</sup> y<sup>-1</sup> with maximum losses of 4.12 g N m<sup>-2</sup> y<sup>-1</sup> in 2009, although in 2004 and 2005 losses from NO<sub>x</sub>/NH<sub>3</sub> volatilisation from excretion and organic fertilisation exceeded losses from N<sub>2</sub> emissions. Losses through NO<sub>x</sub> from the soil were always less than 1% of the total N exchange (0.2 g N m<sup>-2</sup> y<sup>-1</sup>). Nitrous oxide emissions ranged from 0.11 to 1.27 g N m<sup>-2</sup> y<sup>-1</sup>, representing 1.3-8.4 % of the total N export. Annual N<sub>2</sub>O emissions showed no correlation with precipitation, livestock density or

total N input. N<sub>2</sub>O emission factors (percentage of N lost from total N inputs by mineral and organic fertiliser), ranged between 0.6 and 7.5 % (Table 6).

425 To investigate the influence of different managements on the N and C budget, we separated  
experimental years into harvested and grazed (2002 and 2003) and grazed only years (2004-2010 Fig. 2  
and 3). During the harvested years, the main loss of N from the system occurred through leaching  
(39.2% of total N inputs), followed by the export through harvest (25.2%), while the export from  
animals (meat and wool) accounted for less than 2 % of total losses (Fig. 2a). The main loss to the  
430 atmosphere resulted from total denitrification (N<sub>2</sub>; 15.4%), followed by NO<sub>x</sub>/NH<sub>3</sub> volatilisation from  
inorganic N fertiliser applications (9.5%), while N<sub>2</sub>O emissions accounted for 3.3%, NO<sub>x</sub>/NH<sub>3</sub>  
volatilisation from excretion for 2.7% and NO<sub>x</sub> from soil for less than 1%. The residual 2% represents  
the N storage in the soil and the uncertainty of the budget. When grazed-only years were considered  
(Fig. 2b), the residual part was the highest at 38.6%. Losses through leaching (19.9%) and N<sub>2</sub> (11.4%)  
435 were lower in grazed years compared to harvested years, while the export through grazing animals were  
considerably higher at 15.8% (sum of N loss through meat, wool and NO<sub>x</sub>/NH<sub>3</sub> volatilisation from  
excretion). An additional loss occurred in grazed years through the volatilisation of NO<sub>x</sub>/NH<sub>3</sub> from  
organic fertiliser applications in 2004 and 2005 (3%). Losses through N<sub>2</sub>O and NO<sub>x</sub>/NH<sub>3</sub> from inorganic  
fertiliser were comparable to harvested years at 2.5% and 8.3%, respectively.

440 Cumulative soil N stocks were derived from soil core measurements taken in May 2004 and  
May 2011. In 2004 N stocks were 840.86 (±11.89) g N m<sup>-2</sup> and in 2011 they were 870.02 (±14.14) g N  
m<sup>-2</sup>. Nitrogen storage over the 7 years was calculated from the cumulative equivalent soil mass (ESM)  
for the soil mass increment of 800 kg m<sup>-2</sup>, which corresponds to approximate 60 cm depth. The  
estimated N storage over the 7 years was -4.51 ± 2.64 g N m<sup>-2</sup> y<sup>-1</sup> (Table 7) and was a significant N  
445 accumulation to the soil (p < 0.01). The estimated N storage derived from flux calculations between  
2004 and 2010 was -8.44 ± 4.21 g N m<sup>-2</sup> y<sup>-1</sup>, which is almost 2 times more than that calculated by  
sequential soil analysis, however, values were not significantly different from each other.

### 3.3. C budget

450 Annual C inputs through photosynthesis (GPP) varied between -982.1 and -2162.9 g C m<sup>-2</sup>, and losses through autotrophic and heterotrophic respiration (TER) varied between 972.1 and 2183.2 g C m<sup>-2</sup>, both considerably larger than any other C fluxes (Table 5). If only the NEE was considered (difference between GPP and TER), the grassland acted as a sink for CO<sub>2</sub> at an average of 218 ± 155 g C g C m<sup>-2</sup> y<sup>-1</sup>, and the CO<sub>2</sub> uptake was significantly different from zero (p < 0.05). The sink strength ranged from -  
455 10 g C m<sup>-2</sup> y<sup>-1</sup> (2006) to -606 g C m<sup>-2</sup> y<sup>-1</sup> (2009), only in 2004, the grassland was a small source of CO<sub>2</sub> (72 g C m<sup>-2</sup> y<sup>-1</sup>). Taking into account all C inputs and outputs (NBP), C was sequestered on average at 163 ± 140 g C m<sup>-2</sup> y<sup>-1</sup> over the nine years, although the storage was not significantly different from zero (p<0.05). In 2004 and 2006 C was lost from the ecosystem. The major C import resulted from NEE in all years apart from 2005, when the C input from manure application was larger. Highest C export  
460 occurred from harvest in 2002 and 2003 (270.6 and 169.5 g C m<sup>-2</sup>y<sup>-1</sup> respectively), while second largest export in 2002 and 2003 and largest exports in other years was leaching (6.8 to 25.1 g C m<sup>-2</sup>y<sup>-1</sup>). The measured C leaching value for 2007 (15.4 g C m<sup>-2</sup>y<sup>-1</sup>, table 5) differs from the leaching value published for Easter Bush by Kindler et al. (2011), as we only used values of one of the two measured sites in this manuscript (slope, not hollow, as the hollow location was frequently water logged). The third largest C  
465 loss consisted of C export from meat in 2004-2010, ranging from 6.4-15.8 g C m<sup>-2</sup> y<sup>-1</sup>. In 2002 and 2003, when no lambs were present in the field, C export from meat was exceeded by CH<sub>4</sub> losses from enteric fermentation. Carbon export from wool ranged from 0.5 to 2.1 g C m<sup>-2</sup> y<sup>-1</sup>. CH<sub>4</sub> emissions from organic fertilisation, soil processes and animal excretion were always less than 1 % of the total C losses. CH<sub>4</sub> losses from enteric fermentation ranged from 1.5 to 5.7 g C m<sup>-2</sup> y<sup>-1</sup>, corresponding to 0.5-22.5 % of  
470 all C losses from the ecosystem. The NBP was dominated by the NEE. A high livestock density tended to reduce the net sink strength. A significant negative correlation of NEE with stocking density could be seen (R<sup>2</sup>=0.47). The NBP correlated positively with rainfall (R<sup>2</sup>=0.48) and there was only a weak correlation between NEE and rainfall (R<sup>2</sup>=0.38).

The net primary production (NPP) in years when grass was harvested and grazed (2002 and  
475 2003) and grazed only (2004– 2010) are presented in Figure 3. In both management types most C was lost through ecosystem respiration, (67% and 71% of NPP, respectively). Harvest export represented 21% of NPP. Leaching accounted for 1.5% of NPP during harvested years and 2.2% in grazed only

years. Animal export (meat and wool) consisted of 1.5% of NPP in grazed only years and was 0.2% of NPP in grazed and harvested years. The sum of all CH<sub>4</sub> emissions (from organic fertilisation, excretion, enteric fermentation and soil) was less than 1% of the NPP. The residual part, which includes the C storage in the soil as well as the uncertainty of the budget, was estimated at 10% and 24% of NPP in harvested and grazed or grazed years, respectively.

The C content for the cumulative soil mass increment 0-800 kg m<sup>-2</sup> (~ 0-60 cm) was lower in 2011 (12026.05 ± 190.19 g C m<sup>-2</sup>) compared to 2004 (11824.87 ± 187.84 g C m<sup>-2</sup>), resulting in a C loss of 29.08 ± 38.19 g C m<sup>-2</sup> (Table 7). In comparison, based on flux calculations C was stored at 180 ± 180 g C m<sup>-2</sup> y<sup>-1</sup> over the 7 years. However, neither C loss calculated by sequential soil analysis, nor C storage estimated from flux calculations were significantly different from zero.

### 3.4. Greenhouse gas budget

Average greenhouse gas fluxes, net GHG exchange (NGHGE) and attributed net GHG balance (NGHGB) for 2002-2010 are shown in Figure 4. The CO<sub>2</sub> storage from the NEE provided the largest term in the annual GHG budget. Carbon dioxide (NEE) was sequestered over the 9 years at a rate of -799 ± 567 g CO<sub>2</sub> m<sup>-2</sup> y<sup>-1</sup>. This storage was significantly different from zero (p < 0.05). On average, the annual net GHG exchange (NGHGE) was highly correlated with annual NEE (R<sup>2</sup>=0.96). On average the grassland was a source of the GHGs CH<sub>4</sub> and N<sub>2</sub>O at a rate of 148 ± 30 and 285 ± 131 g CO<sub>2</sub> m<sup>-2</sup> y<sup>-1</sup>, respectively, both being significantly different from zero (p < 0.001 and p < 0.01, respectively). Nitrous oxide losses ranged from 52 g CO<sub>2</sub> eq. m<sup>-2</sup> y<sup>-1</sup> (2004) to 588 g CO<sub>2</sub> eq. m<sup>-2</sup> y<sup>-1</sup> (2007) (data for each year not shown). Methane from soil processes, manure input as well as animal excretion, accounted for less than 5% of total CH<sub>4</sub> emissions. Methane emissions from enteric fermentation ranged from 53 g CO<sub>2</sub> eq. m<sup>-2</sup> y<sup>-1</sup> (2002) to 199 g CO<sub>2</sub> eq. m<sup>-2</sup> y<sup>-1</sup> (2004). Annual total CH<sub>4</sub> emissions correlated positively with annual live stock density (R<sup>2</sup>=0.99). The CH<sub>4</sub> emissions, which were predominately (> 97%) of ruminant origin weakened the sink strength of NEE by 18 %. If both CH<sub>4</sub> and N<sub>2</sub>O were considered the total trade-off of NEE was a substantial 54% and increased to a total of 67 %, if only grazed years were considered. On average the grassland represented a GHG sink of -366 ± 601 g CO<sub>2</sub> m<sup>-2</sup> y<sup>-1</sup>, if only NEE, CH<sub>4</sub> and N<sub>2</sub>O were included (NGHGE). If all C components (FC<sub>org.fert</sub>, FC<sub>animal</sub>, FC<sub>leaching</sub>,

FC<sub>harvest</sub>) are included, the sink strength of the grassland decreased to  $-182 \pm 560 \text{ g CO}_2 \text{ m}^{-2} \text{ y}^{-1}$  (NGHGB). This represents a weakening of the sink strength of the NGHGE by 50 %, mainly due to the export of harvest. However, it has to be noted that in harvested years the return of the manure, resulting from the grass fed to livestock off -site, would reduce the GHG balance. If only grazed years were considered the sink strength increased slightly by 5.4 %, due to the C input from manure in 2004 and 2005. Both, NGHGE and NGHGB were not significantly different from zero.

## 4. Discussion

### 4.1. Nitrogen balance

The main N inputs in our study were from inorganic and organic fertilizer additions. The amount of N added through fertilizer was determined by national recommendations (SAC, 2013) and lies within the range of N applied in other European studies with similar management (e.g. Laws et al., 2000; Allard et al., 2007; Ammann et al., 2009). Atmospheric N deposition (wet and dry) accounted only for a small fraction of the total N input to our managed grassland. This is in contrast to semi natural systems, where atmospheric N deposition and biological fixation represents the main N input (Pheonix et al., 2006, Bleeker et al., 2011). As our experimental field was sown as a grass mixture (without clover) the legume fraction was less than 1% and biological N<sub>2</sub> fixation therefore a negligible source of N.

The data obtained from our budget were used to calculate the Nitrogen Use Efficiency (NEU) expressed as the ratio between N in crop and animal products (in this case either the crop harvest or the sum of meat, wool and milk) to the total N inputs to the system (fertilizer, imported manure). The NUE of herbage in cut years (2002 and 2003) of 25% (Figure 2a) seems low compared to reported N efficiencies of 55-80% in harvested herbage from managed temperate grasslands (Ball and Ryden 1984; Ammann et al., 2009). It has been shown that the NUE in crops is significantly higher compared to the NUE in animal production (Galloway and Cowling, 2002). The inclusion of grazing ruminants introduces an additional trophic level altering the NUE of herbage as the nitrogen in the grazed grass is consumed and converted to meat, milk, wool, or is excreted. The lower NUE in the grass production in our study is therefore partly due to grazing. Furthermore, it has been shown that the proportion partitioned to plant uptake decreases as the total amount of soil inorganic N increases (Scholefield et al.,

1991), which is a further explanation for a low NUE in herbage in our high N input system. The NUE of  
535 animal products on our grassland system ranged from 6 to 21% in grazed only years (2004-2010), with  
an average of 9.9 %. This is in agreement with the NUE reported for sheep of 6.2 % by Van der Hoek  
(1998) and beef production systems, which reported N efficiencies range from 6 to 12% (Whitehead et  
al., 1986; Tyson et al., 1992) and 5-20% (Ball and Ryden, 1984). Approximately 85% of total harvested  
N is used to feed livestock (Sutton 2011). A measure to reduce N pollution could therefore be the  
540 reduction of meat consumption or a larger fraction of meat produced from grassland only (Smith et al.  
2013).

Nitrogen was lost from our grassland to the environment through different pathways. Nitrogen  
leaches from grassland soils in the form of nitrate ( $\text{NO}_3^-$ ), ammonium ( $\text{NH}_4^+$ ) and dissolved organic N  
(DON). Overall, leaching from our field ( $5.3 \pm 3.4 \text{ g N m}^{-2} \text{ y}^{-1}$ ) was comparable to values measured at  
545 intensively grazed pastures in Ireland ( $1.8\text{-}6.4 \text{ g N m}^{-2} \text{ y}^{-1}$ , Watson et al., 2007) and England ( $3.8\text{-}13.3 \text{ g}$   
 $\text{N m}^{-2} \text{ y}^{-1}$ , Scholefield et al., 1993) or croplands (e.g. Bechmann et al., 1998), max. leaching losses of  
 $10.4 \text{ g N m}^{-2} \text{ y}^{-1}$ ). However, leaching from our study was high compared to the Swiss NitroEurope site,  
where a maximum loss of  $0.35 \text{ g N m}^{-2} \text{ y}^{-1}$  was estimated from an ungrazed grass/clover sward, despite  
comparable annual rainfall and N inputs (Ammann et al., 2009). This difference can be explained by the  
550 different plant cover and management. It has been shown that clover introduction can reduce leaching  
(Owens et al., 1994), whereas grazing tends to increase leaching (Cuttle and Scholefield, 1995). Grazed  
grasslands tend to have higher N leaching rates than cut grasslands since highly concentrated N  
deposited in urine is inefficiently recovered by herbage and prone to leaching. The uneven distribution  
of excreted organic N further enhances leaching due to the formation of N hotspots, which has been  
555 observed at outdoor pig farms (e.g. (Eriksen, 2001). Ryden et al., (1984a) showed a 5.6 times higher  
leaching loss from grazed compared to cut grassland with 36% of total N inputs lost from grazed  
compared to 6% lost from cut grassland. On our site leaching represented about 20 % of total inputs in  
grazed years, compared to 39% in the cut years. However, the higher value in cut years was due to the  
high rainfall in 2002.

560 Due to its high background in the atmosphere,  $\text{N}_2$  fluxes cannot be measured directly in the  
field. There are different methods to measure  $\text{N}_2$  fluxes indirectly, which have been summarized by

Groffman et al. (2006). In our study, we estimated N<sub>2</sub> losses using the process based biogeochemical model LandscapeDNDC (Haas et al., 2013, Molina-Herrera et al., 2016). These losses represented the highest gaseous N losses from our grassland in most years, with an average of 12.6 % of total N inputs and 14 % of inorganic fertilizer N inputs. This is comparable with an average N<sub>2</sub> loss of 12.5 % from inorganic N applications measured by the acetylene inhibition method from a fertilized and cut, but ungrazed grassland in Switzerland (Rudaz et al., 1999). Using the same method, Van der Salm et al. (2007) reported a higher loss of 22% of total N input from a cattle grazed pasture on a heavy clay soil in the Netherlands. In addition to the impact of the heavy clay soil, which could have enhanced denitrification due to reduced oxygen concentrations, grazing is likely to have enhanced denitrification rates in Van der Salm's study. Grazing not only enhances denitrification through soil compaction caused by trampling animals but also due to the formation of N hot spots resulting from unevenly distributed soil N from excretion. In our study N<sub>2</sub> losses simulated by LandscapeDNDC were based on average (per ha<sup>-1</sup>) changes of the soil N pool instead of the more uneven distribution of soil N in hot spots like urine patches. Therefore it is likely that N<sub>2</sub> losses in our study have been underestimated and better estimation would contribute to a significant reduction in the uncertainties associated with the overall N budget.

Annual N<sub>2</sub>O emissions measured in our study (0.1 to 1.3 g N m<sup>-2</sup> y<sup>-1</sup>) are within the range of literature values from reported grazed as well as un-grazed European grasslands (Leahy et al., 2004; Flechard et al., 2007). Generally N<sub>2</sub>O losses are higher from grazed grassland compared to cut, ungrazed pasture (Velthof and Oenema, 1995; Luo et al., 1999) due to a more anaerobic environment as a consequence of soil compaction caused by animal treading and the influence of N and C from the deposition of animal excreta to the soil. We did not observe any correlations between annual N<sub>2</sub>O emissions and stocking density, rainfall or total N input. This demonstrates that N<sub>2</sub>O emissions are not simply a uniform fraction of N applied, as suggested by the Tier 1 IPPC methodology, but are also influenced by the type of N applied, by stocking density, and by the rainfall at the time of fertilization (Jones et al., 2007; Flechard et al., 2007). We found a relationship between the cumulative precipitation one week before and three weeks after fertilization with N<sub>2</sub>O emissions (R<sup>2</sup>=0.53) (Skiba et al., 2013). This relationship, together with the influence of stocking density and type of N applied needs to be

590 considered when developing Tier 2 N<sub>2</sub>O emission factors. In our study EFs were above the uncertainty range (0.3 - 3 %) given by IPCC Tier 1 guidelines (IPCC, 2006b) in four out of nine years.. However, it has been shown that the N<sub>2</sub>O emission factor from managed grassland can be higher, especially under wet conditions and with a high soil C content as this is the case for Scottish soils (Jones et al., 2007; Dobbie et al., 1999; Buckingham et al., 2013).

595 In grazed pastures NH<sub>3</sub> volatilizes from urine patches, decomposing dung as well as from fertilizers containing urea and NH<sub>4</sub><sup>+</sup> (Twigg et al 2011). Increased rates of NH<sub>3</sub> losses have been associated with high stocking densities under a rotational grazing system by Ryden and Mc Neill (1984). In our study, N volatilized as NH<sub>3</sub> and NO<sub>x</sub> from inorganic and organic fertiliser and animal excretion, accounted for a considerable amount of total N, with losses of 13 % in cut and grazed years  
600 (2002, 2003) and 17 % in grazed only years. In contrast, soil NO<sub>x</sub> emissions from our grassland were estimated to be negligible, accounting for less than 1% of the total budget. Soil NO<sub>x</sub> emissions result predominantly from microbial nitrification of either added N fertilizers or following the mineralization of soil organic matter, animal excretions or added manure. Emissions tend to be linked with aerobic soil conditions (Davidson, 1991).

605 Results from soil analysis taken in May 2004 and May 2011 indicate that our field has stored N (-4.51 ± 2.64 g N m<sup>-2</sup> y<sup>-1</sup>). The N budget assessed from the net N flux balance showed that N was stored in the soil of our grassland over the same 7 years at a higher rate (-8.44 ± 4.21 g N m<sup>-2</sup> y<sup>-1</sup>), although values were not significantly different from each other. The slight shifts in measurement periods (May 2004 – May 2011) for the soil stock calculations and the period for flux budget calculations (Jan 2004 –  
610 Dec 2010), is presumed to be insignificant in this comparison. Results from both methods are within the range of literature values. Neeteson and Hassink (1997) found a N accumulation in SOM of 0-25 g N m<sup>-2</sup> y<sup>-1</sup> from two cattle-grazed farms in the Netherlands, while Watson et al. (2007) reported a N storage in grazed Irish grasslands ranging from 10-15.2 g N m<sup>-2</sup> y<sup>-1</sup>, depending on N inputs. Soil N storage assessed from soil measurements from a cut grassland close to our field, where plots were treated with  
615 cattle slurry, stored N over 6 years at a rate of -2.17 g N m<sup>-2</sup> y<sup>-1</sup> in the top 10 cm, while, in the same experiment, a N loss was observed from mineral N and urea treatments (4.5 and 8.3 g N m<sup>-1</sup>y<sup>-1</sup>, respectively) (Jones et al., 2007). In contrast, Schipper et al. (2007) reported an average loss of 9.1 g N

m<sup>-2</sup> y<sup>-1</sup> in the top 100 cm from managed grasslands over 20 years in New Zealand. The reason for the small difference between methods (flux measurements vs sequential soil sampling) in our study might  
620 lie in a possible underestimation of losses from flux measurements. Uncertainties of our estimates are high, especially those for N losses. The largest absolute systematic uncertainty for the N balance was attributed to N leaching as for most years values were modelled using data to validate the model from only one location. The uncertainty of the leaching estimate would therefore be reduced if the model could be validated with data measured from several locations. The second highest systematic  
625 uncertainty was attributed to losses through N<sub>2</sub>, followed by NO<sub>x</sub>/NH<sub>3</sub> emission from excretion, NO<sub>x</sub>/NH<sub>3</sub> emission from inorganic fertilization and inputs from organic fertilization. Combined uncertainties from all components lead to a total systematic uncertainty in the N balance of 2.1 g N m<sup>-2</sup> y<sup>-1</sup> (2004-2010).

#### 630 **4.2. Carbon balance**

On an annual basis our grassland site was a sink for atmospheric CO<sub>2</sub> in most years. NEE was only positive in 2004, which was likely to be due to a high livestock density. Generally, grazing causes a very gradual impact on the CO<sub>2</sub> uptake as a part of the field is defoliated each day. The reduced leaf area index (LAI) then leads to a reduced CO<sub>2</sub> uptake by plants. In addition to the reduced LAI, grazing  
635 presents a source of CO<sub>2</sub> from animal respiration, thereby reducing the CO<sub>2</sub> sink of the grassland within the field (Levy et al., submitted). The maximum uptake of CO<sub>2</sub> measured in our study is close to the upper range of NEE reported for temperate grasslands (100 to 600 g C m<sup>-2</sup> y<sup>-1</sup>, (IPCC, 1996). On average over the 9 years the magnitude of the NEE on our grassland (-218.0 ± 154.5 g C m<sup>-2</sup> y<sup>-1</sup>) was close to the average NEE measured in a comparison of nine European grasslands over two years (240 ±  
640 70 g C m<sup>-2</sup> y<sup>-1</sup>) by Soussana et al. (2007) and comparable to the CO<sub>2</sub> sink capacity of managed Irish grasslands measured by Byrne et al. (2007) (290 ± 50 g C m<sup>-2</sup> y<sup>-1</sup>) or Leahy et al. (2004) (257 g C m<sup>-2</sup> y<sup>-1</sup>). Despite high variability over the 9 years, the average NEE value was significantly different from zero (p < 0.05). The range of the calculated annual gross primary production (GPP) (-982 to -2163 g C m<sup>-2</sup> y<sup>-1</sup>) and terrestrial ecosystem respiration (TER) (972 to 2183 g C m<sup>-2</sup> y<sup>-1</sup>) from our field were within  
645 reported values for other managed grasslands. Gilmanov et al. (2007) reported the GPP of 18

intensively managed European grasslands ranging from 467 to 1874 g C m<sup>-2</sup> y<sup>-1</sup> and TER ranging from 493 to 1541 g C m<sup>-2</sup> y<sup>-1</sup>, while Mudge et al. (2011) reported values of 2000 g C m<sup>-2</sup> y<sup>-1</sup> for GPP and TER from a intensively grazed dairy pasture in New Zealand.

When all components of C import and export were included in addition to the CO<sub>2</sub> exchange (NBP) C was stored in our grassland over the 9 years. However, due to the high variability between years, NBP was not significantly different from zero (p = 0.05), suggesting that our site is carbon neutral. The average C storage value on our site (164 ± 140 g C m<sup>-2</sup> y<sup>-1</sup>) is higher than most estimates reported in literature, but due to the high annual variation, still within the range of reported values; Soussana et al. (2007) reported C storage estimates from European grazed and cut grasslands of 104 ± 73 g C m<sup>-2</sup> y<sup>-1</sup>, and Mudge et al. (2011) reported for a grazed and cut grassland in New Zealand fluxes of 59 ± 56 g C m<sup>-2</sup> y<sup>-1</sup> and 90 ± 56 g C m<sup>-2</sup> y<sup>-1</sup> in two consecutive years. NBP estimates from a Swiss grassland cut for silage was shown to sequester C at a rate of 147 ± 130 g C m<sup>-2</sup> y<sup>-1</sup> (Ammann et al., 2007), while estimates from a cut grassland in Germany was shown to vary from being a sink (-28 g C m<sup>-2</sup> y<sup>-1</sup>) to being a source of C (+25 g C m<sup>-2</sup> y<sup>-1</sup>), depending on years (Prescher et al., 2010). The inclusion of all C imports and exports lead to a weakening of the C sink strength assessed from NEE measurements in five years and even changed the grassland from being a sink to being a source in 2006. Due to the C export from harvest, C sequestration tends to be lower in cut systems. This is represented in our study in the lower residual value of NPP in cut years compared to the residual value from grazed only years (Figure 3), where the residual value represents the C storage in the soil as well as the uncertainty of the budget. However, it has to be kept in mind that the herbage yielded from cuts will end up as animal feed; C will be digested and respired off-site, releasing CO<sub>2</sub> and CH<sub>4</sub> to the atmosphere as well as being returned to the grassland as manure.

Results from soil analysis indicate that our grassland has lost C from 2004-2010 (29±38 g C m<sup>-2</sup> y<sup>-1</sup>, Table 7). In the literature, losses as well as storage of C at various rates have been reported from managed grasslands assessed from soil stock measurements. Soil stock change measurements from our field are comparable with values found in the literature. Depending on the study, managed grasslands in Belgium were shown to either loose (90 g C m<sup>-2</sup> y<sup>-1</sup>, Lettens et al., 2005a) or sequester carbon (4.4 g C m<sup>-2</sup> y<sup>-1</sup> in 0-30 cm, Goidts and Van Wesemael, 2007; 22.5 g C m<sup>-2</sup> y<sup>-1</sup> in 0-30 cm, Lettens et al. 2005b).

Schipper et al. (2007) reported losses of C from pastures in New Zealand over 20 years at an average  
675 rate of  $106 \text{ g C m}^{-2} \text{ y}^{-1}$  (top 100 cm), whereas these losses were a result of an earlier land use change  
from forestry. Schuman, et al., (2002) measured a C sequestration of  $10\text{-}30 \text{ g C m}^{-2} \text{ y}^{-1}$  from US  
rangelands (0-60 cm)), while Watson et al. (2007) measured a C storage at  $112\text{-}145 \text{ g C m}^{-2} \text{ y}^{-1}$  in the  
top 15 cm soil layer from a grazed Irish grassland. Bellamy et al. (2005) showed no evidence of  
680 increased C in the topsoil of grasslands in England and Wales and Hopkins et al. (2009) found no  
significant change of SOC over time in two UK long term experiments. The above mentioned results  
are contrasting and inconclusive, because observed C sinks in grasslands are the effect of land  
management or land use change prior to the beginning of the C stock change measurement. Soussana et  
al (2014) concluded in a theoretical study that grassland SOC sequestration has a strong potential to  
partly mitigate the GHG balance of ruminant production systems at low grazing intensities, but not with  
685 intensive systems. Smith (2014) examined evidence from repeated soil surveys, long term grassland  
experiments and simple mass balance calculations and concluded that, although grasslands can act as C  
sinks, they cannot act as a perpetual C sink and thus could not be used as an offset for GHG emissions.

The comparison of the C storage calculated from the net C flux balance with soil C stock  
changes show that, the flux balance estimated a C sequestration, while based on C stock changes, C was  
690 lost, although neither value was significantly different from zero (Table 7). A literature search by  
Soussana et al. (2010) showed that generally C sequestration calculations on grassland were lower if  
derived from SOC stock changes (average  $-5\pm 30 \text{ g C m}^{-2} \text{ y}^{-1}$ ) compared to C flux balances (average  
 $-22\pm 56 \text{ g C m}^{-2} \text{ y}^{-1}$ ), although these estimates were not significantly different from each other. However,  
in none of those reviewed studies were C flux and C stock change measured in the same field  
695 experiment. A reason for the discrepancy between estimation methods in our study might lie in a  
possible underestimation of C exports in the flux balance calculation, leading to an overestimation of C  
storage in the soil. One underestimated flux could be the export of DIC and DOC. Carbon leaching from  
managed grasslands has not been reported in many studies. Kindler et al. (2011) reported C leaching  
from various European ecosystems, where the measured data (2007) from our experimental field was  
700 part of the study. Our data ( $30.0 \text{ g C m}^{-1} \text{ y}^{-1}$ , average of two locations as published in Kindler et al.  
(2011)) were close to the average value ( $29.4 \text{ g C m}^{-1} \text{ y}^{-1}$ ) of the reported European grasslands, which

showed a range of C losses of 6.5-42.5 g C m<sup>-1</sup> y<sup>-1</sup>. Higher losses have been observed by McTiernan et al. (2001), who measured DOC export from grassland lysimeter plots treated with N fertilizer and slurry over two months. Up-scaled to one year, they measured DOC loss between 25.2 and 70.8 g C m<sup>-2</sup> y<sup>-1</sup>, all above what we measured in our study. Important factors controlling the magnitude of C leaching have been shown to be drainage, the topsoil C/N ratio and the saturation of the subsoil's sorption capacity for organic C (Kindler et al., 2011; McTiernan et al., 2001). In waterlogged soils the soil organic matter (SOM) decomposition and groundwater recharge tend to be reduced and thus the amount of C prone to leaching compared to that under more aerobic conditions associated with drainage. Although our field was drained more than 50 years ago, the drainage system does not operate very well, resulting in large puddles of standing water during prolonged periods of rain. The measured data used for the budget were taken at one sampling point, which was not in a waterlogged area. The spatial heterogeneity within the grassland field caused by uneven water management as well as faeces and urine patches requires to sample at more points in order to obtain a representative leaching value. Therefore our leaching estimates are highly uncertain and could be significantly lower and C exports overestimated. Furthermore, leaching was only measured in one year (2008), while values for remaining years were estimated using a simple regression model with an attributed high uncertainty of 32 % (5.3 g C m<sup>-2</sup> y<sup>-1</sup> of average fluxes). Indeed, Siemens (2003) hypothesized that the underestimation of C leaching from soils can explain a large part of the difference between atmosphere- and land-based estimates of the C uptake of European terrestrial ecosystems. Another underestimated flux could be the loss of CO<sub>2</sub> in the NEE measurements. Gapfilling can introduce uncertainties in the NEE data especially for years with low data capture. Furthermore, CO<sub>2</sub> losses from animal respiration could be underestimated at times due to the animals moving out of the footprint of the EC mast. Using animal respiration values from chamber experiments of 12.1 g CO<sub>2</sub> kg<sup>-1</sup> live weight d<sup>-1</sup> for cows and 11.7 g CO<sub>2</sub> kg<sup>-1</sup> live weight d<sup>-1</sup> for sheep and lambs (Shane Troy, SRUC, personal communication), we estimated a maximum CO<sub>2</sub> loss from animal respiration of 53 g C m<sup>-2</sup> y<sup>-1</sup> (2002-2010) or 59 g C m<sup>-2</sup> y<sup>-1</sup> (2004-2010). So if we assume that all animal respiration has been missed by eddy covariance measurements then the C sink estimated from NEE measurements would be reduced by 24 % (2002-2010) or 33 % (2004-2010). This theoretical maximum 33% reduction would reduce the net carbon balance to ~ 122 g C m<sup>-2</sup> y<sup>-1</sup> (2004-2010).

In addition to uncertainties in the flux budget calculations, uncertainties are also attributed to soil C and N stock measurements. Soil inventory data in our study indicated a loss of C and a storage of N over 7 years, which seems contradictory, although C storage was not significantly different from zero. The uncertainty of soil C and N stock measurements arise from the variability of soil C and N concentrations due to errors from laboratory and to their high spatial variability as well as from the variability in the rock fragment content (Goidts et al., 2009).

### 4.3 Greenhouse gas budget

In the overall N and C budget  $\text{N}_2\text{O}$  and  $\text{CH}_4$  emissions were negligible in terms of N and C losses from the system (1 – 8 % of total N losses and 0.6 - 4.5 % of total C losses, respectively). However, in terms of  $\text{CO}_2$  equivalents,  $\text{N}_2\text{O}$  emissions as well as  $\text{CH}_4$  emissions strongly affected the GHG budget. Indeed, the sink strength of the NEE was weakened by  $\text{N}_2\text{O}$  emissions by 29 % over all years. Methane emissions from soil processes, manure input and animal excretion were negligible in terms of the C budget as well as in the GHG budget. In contrast, enteric fermentation proved to be an important GHG source. Methane emissions were also measured by eddy covariance technique over several months in 2010 on the same field (Dengel et al., 2011). By dividing  $\text{CH}_4$  fluxes by the number of sheep in the field each day, Dengel et al. calculated  $\text{CH}_4$  emissions per head of livestock as  $7.4 \text{ kg CH}_4 \text{ head}^{-1} \text{ y}^{-1}$  for sheep, which is close to the emission factor used in our budget of  $7.6 \text{ kg CH}_4 \text{ head}^{-1} \text{ y}^{-1}$  for ewes, showing that our estimates were realistic. The positive correlation of  $\text{CH}_4$  emissions with the stock density indicates that any changes in animal production will have a major impact on the global  $\text{CH}_4$  budget. The weakening of the GHG sink strength of the NEE by  $\text{N}_2\text{O}$  and  $\text{CH}_4$  emissions, show the importance of those two gases in terms of global warming. Thus, adapting the management of grasslands by adding fertilizer or manure to increase plant growth and thus improve C sequestration in soils may increase  $\text{N}_2\text{O}$  emissions, while changing land use from cropland to pasture in the attempt to reduce C losses from soils might lead to increased  $\text{CH}_4$  losses from grazing animals if the total number of animals increases rather than animals are fed in a different way.

## 5. Conclusion

In our study only a small proportion of the N inputs from inorganic fertilizer and organic manure were converted to animal outputs or stored in the soil, while the main part was lost through leaching and gaseous emissions. An improvement of the NUE would mean both an economic profit for the farmer as well as an environmental benefit. Estimates from flux budget calculations indicated that our grassland was sequestering C. However, although grasslands can act as C sinks, they cannot act as a perpetual C sink and thus could not be used as an offset for GHG emissions (Smith et al., 2014). Instead, as it is easier and faster for soils to lose than to gain carbon, care must be taken to preserve C loss by management options, rather than trying to increase carbon stocks in grasslands. There was a discrepancy between soil stock measurements and flux budget calculations for the C as well as the N budget. The reason for the discrepancy between C budget estimates might lie in a possible underestimation of C exports such as leaching and animal respiration as well as the uncertainty due to gapfilling in the NEE data. The N accumulation might have been overestimated by the flux calculations through a possible overestimation of N losses, mainly through leaching as well as through N<sub>2</sub> and NO<sub>x</sub>NH<sub>4</sub> emissions. Furthermore, uncertainties are also attributed to soil C and N stock measurements. Our data have shown that the information about the potential of managed grasslands to act as sinks or sources for GHG is important for mitigation and adaption purposes. High plant productivity, stimulated by fertilisation, resulted in high plant CO<sub>2</sub> fixation. However, increased N losses through N<sub>2</sub>O emissions counteracted the benefits of C sequestration in terms of GHG emissions. Furthermore, CH<sub>4</sub> emissions from enteric fermentation largely reduced the positive effect of CO<sub>2</sub> uptake, especially in years where NEE rates were small. We therefore conclude that CO<sub>2</sub> exchange alone is not sufficient for the estimation of the GWP of a managed grassland ecosystem. Only a comprehensive approach, combining C and N cycling will help us to better understand functionalities of ecosystems and to improve modelling by integrating this knowledge.

## Acknowledgments

We would like to thank Chris Flechard for help with N deposition modelling, Patricia Ricci for help with IPCC methodology used for CH<sub>4</sub> animal emissions and NO<sub>x</sub>/NH<sub>3</sub> volatilisation calculations and

Sue Owen for help with VOC emission calculations. We are grateful to Ivan Simmons, Julia Drewer and Ian Leith for their help with field infrastructure. The authors gratefully acknowledge funding from the EU project GREENGRASS (EC EVK2-CT2001-00105), the NitroEurope Integrated Project (contract 017841) and CarboEurope (Contract No. GOCE-CT-2003-505572).

790

## Appendix A. Supplementary information

### Reference List

- 795 Allard, V., Soussana, J. F., Falcimagne, R., Berbigier, P., Bonnefond, J. M., Ceschia, E.,  
D'hour, P., Henault, C., Laville, P., Martin, C., and Pinares-Patino, C.: The role of grazing management  
for the net biome productivity and greenhouse gas budget (CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>) of semi-natural  
grassland, *Agr. Ecosyst. Environ.*, 121.1-2, 47-58, 2007.
- 800 Ammann, C., Flechard, C. R., Leifeld, J., Neftel, A., and Fuhrer, J.: The carbon budget of  
newly established temperate grassland depends on management intensity, *Agr. Ecosyst. Environ.*,  
121.1-2, 5-20, 2007.
- 805 Ammann, C., Neftel, A., Spirig, C., Leifeld, and J., Fuhrer, J.: Nitrogen balance of hay  
meadows with and without fertilization, *Agrarforschung*, 16.9, 348-53, 2009.
- Amthor, J.S.: The McCree–de Wit–Penning de Vries–Thornley Respiration Paradigms: 30 Years Later,  
*Annals of Botany* 86 (1): 1–20, 2000.
- 810 Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A. S.,  
Martin, P. H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grunwald, T.,  
Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., and Vesala, T.: Estimates of  
the annual net carbon and water exchange of forests: The EUROFLUX methodology, *Adv. Ecol. Res.*,  
30, 113-75, 2000.
- 815 Ball, P. R. and Ryden, J. C.: Nitrogen Relationships in Intensively Managed Temperate  
Grasslands, *Plant Soil*, 76.1-3, 23-33, 1984.
- 820 Bechmann, M., Eggestad, H. O., and Vagstad, N.: Nitrogen balances and leaching in four  
agricultural catchments in southeastern Norway, *Environ. Pollut.*, 102, 493-99, 1998.
- Bellamy, P. H., Loveland, P. J., Bradley, R. I., Lark R. M., and Kirk, G. J. D.: Carbon  
losses from all soils across England and Wales 1978–2003, *Nature*, Vol 437, 245-48., 2005.

- 825 Bleeker, A., Hicks, W. K., Dentener, F., Galloway, J., Erisman, J. W.: N deposition as a threat to the World's protected areas under the Convention on Biological Diversity. *Environ. Pollut.* 159, 2280e2288, 2011.
- Braun-Blanquet J.: *Pflanzensoziologie, Grundzüge der Vegetationskunde*, 3. Aufl. Springer, 1964, Verlag, Wien and New York.
- 830
- Byrne, K. A., Kiely, G., and Leahy, P: Carbon sequestration determined using farm scale carbon balance and eddy covariance, *Agr. Ecosyst. Environ.*, 121.4, 357-64, 2007.
- 835 Buckingham, S., Rees, R.M. & Watson, C.A. 2013. Issues and pressures facing the future of soil carbon stocks with particular emphasis on Scottish soils. *The Journal of Agricultural Science*, 152, 699-715.
- Butterbach-Bahl, K., Gasche, R., Breuer, L., and Papen, H.: Fluxes of NO and N<sub>2</sub>O from temperate forest soils: impact of forest type, N deposition and of liming on the NO and N<sub>2</sub>O emissions, *Nutr. Cycl. Agroecosys.*, 48, 79-90, 1997.
- 840
- Caro, D., Davis S. J., Bastianoni, S., Caldeira K.: Global and regional trends in greenhouse gas emissions from livestock. *Climatic Change*, 126:203–216, 2014
- 845 Chen, W., McCaughey, W. P., and Grant, C. A.: Pasture type and fertilization effects on N-2 fixation, N budgets and external energy inputs in western Canada, *Soil. Biol. Biochem.*, 36.8, 1205-12, 2004.
- Clayton, H., Arah, J. R. M., and Smith, K. A.: Measurement of Nitrous-Oxide Emissions from Fertilized Grassland Using Closed Chambers. *J. Geophys. Res.-atmos.*, 99. D8, 16599-607, 1994.
- 850
- Cuttle, S. P. and Scholefield, D.: Management options to limit nitrate leaching from grassland. *J. Cont. Hydrol.*, 20, 299-312, 1995.
- 855
- Davidson, E. A., 1991. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In: Rogers, J. E., Whitman, W. B. (Eds.), *Microbial Production and Consumption of Greenhouse Gases: Methane, Nitrogen Oxides and Halomethanes*. American Society of Microbiology, Washington, DC, pp. 219–236.
- 860
- Davison, B., Brunner, A., Ammann, C., Spirig, C., Jocher, M., and Nefel, A.: Cut-induced VOC emissions from agricultural grasslands, *Plant Biol.*, 10.1, 76-85, 2008.

- 865 Dengel, S., Levy, P. E., Grace, J., Jones, S. K., and Skiba, U. M.: Methane emissions from sheep pasture, measured with an open-path eddy covariance system, *Glob. Change Biol.*, 17.12, 3524-33, 2011.
- Di Marco, C., Skiba, U., Weston, K., Hargreaves, K., and Fowler, D.: Field scale N<sub>2</sub>O flux measurements from grassland using eddy covariance, *Water Air Soil Poll.: Focus*, 4.6, 143-49, 2004.
- 870 Dobbie, K. E., McTaggart, I. P., and Smith, K. A.: Nitrous oxide emissions from intensive agricultural systems: Variations between crops and seasons, key driving variables, and mean emission factors, *J. Geophys. Res.-atmos.*, 104.D21, 26891-99, 1999.
- 875 Eriksen, J.: Nitrate leaching and growth of cereal crops following cultivation of contrasting temporary grasslands, *J. Agr. Sci.*, 136, 271-81, 2001.
- Flechard, C. R., Ambus, P., Skiba, U., Rees, R.M., Hensen, A., van Amstel, A., Pol-van Dassel, A.V., Soussana, J. F., Jones, M., Clifton-Brown, J., Raschi, A., Horvath, L., Neftel, A.; Jocher, M., Ammann, C., Leifeld, J., Fuhrer, J., Calanca, P., Thalman, E., Pilegaard, K., Di Marco, C., 880 Campbell, C., Nemitz, E., Hargreaves, K. J., Levy, P. E., Ball, B. C., Jones, S. K., van de Bulk, W. C. M., Groot, T., Blom, M., Domingues, R., Kasper, G., Allard, V., Ceschia, E., Cellier, P., Laville, P., Henault, C., Bizouard, F., Abdalla, M., Williams, M., Baronti, S., Berretti, F., and Grosz, B.: Effects of climate and management intensity on nitrous oxide emissions in grassland systems across Europe, *Agr. Ecosyst. Environ.*, 121.1-2, 135-52, 2007.
- 885 Flechard, C. R., Nemitz, E., Smith, R. I., Fowler, D., Vermeulen, A. T., Bleeker, A., Erismann, J. W., Simpson, D., Zhang, L., Tang, Y. S., and Sutton, M. A.: Dry deposition of reactive nitrogen to European ecosystems: a comparison of inferential models across the NitroEurope network, *Atmos. Chem. Phys.*, 11.6, 2703-28, 2011.
- 890 Flindt, R.: *Biologie in Zahlen: Eine Datensammlung in Tabellen mit ueber 10000 Einzelwerten.* Spektrum akademischer Verlag. Gustav Fischer, 249, 2002.
- 895 Galloway, J.N. and Cowling, E.B.: Reactive Nitrogen and the World: 200 Years of Change. 2002. *Ambio*, Vol. 31, No. 2, Optimizing Nitrogen Management in Food and Energy Productions, and Environmental Change (Mar., 2002), pp. 64-71.
- 900 Gilmanov, T. G., Aires, L., Barcza, Z., Baron, V. S., Beileli, L., Beringer, J., Billesbach, D., Bonal, D., Bradford, J., Ceschia, E., Cook, D., Corradi, C., Frank, A., Gianelle, D., Gimeno, C., Gruenwald, T., Guo, H.Q., Hanan, N., Haszpra, L., Heilman, J., Jacobs, A., Jones, M. B., Johnson, D. A., Kiely, G., Li, S. G., Magliulo, V., Moors, E., Nagy, Z., Nasyrov, M., Owensby, C., Pinter, K., Pio, C., Reichstein, M., Sanz, M. J., Scott, R., Soussana, J. F., Stoy, P. C., Svejcar, T., Tuba, Z., and Zhou, G. S.: Partitioning European grassland net ecosystem CO<sub>2</sub> exchange into gross primary productivity and

- ecosystem respiration using light response function analysis, *Agr. Ecosyst. Environ.*, 121.1-2, 93-120, 2007.
- 905
- Goidts, E. and Van Wesemael, B.: Regional assessment of soil organic carbon changes under agriculture in Southern Belgium (1955-2005), *Geoderma*, 141.3-4, 341-54, 2007.
- 910
- Goidts, E., Van Wesemael, B. and Crucifix, M.: Magnitude and sources of uncertainties in soil organic carbon (SOC) stock assessments at various scales. *European Journal of Soil Science*, October 2009, 60, 723–739, 2009.
- 915
- Groffman, P. M., Altabet, M. A., Bohlke, J. K., Butterbach-Bahl, K., David, M. B., Firestone, M. K., Giblin, A. E., Kana, T. M., Nielsen, L. P., and Voytek, M. A.: Methods for measuring denitrification: Diverse approaches to a difficult problem, *Ecol. Appl.*, 16.6, 2091-122, 2006.
- 920
- Haas, E., Klatt, S., Frohlich, A., Kraft, P., Werner, C., Kiese, R., Grote, R., Breuer, L., Butterbach-Bahl, K.: LandscapeDNDC: a process model for simulation of biosphere–atmosphere–hydrosphere exchange processes at site and regional scale. *Landscape Ecol*, 28, 615–636, 2013. DOI 10.1007/s10980-012-9772-
- 925
- Hopkins, D. W., Waite, I. S., McNicol, J. W., Poulton, P. R., Macdonald, A. J. and O'Donnell, A. G.: Soil organic carbon contents in long-term experimental grassland plots in the UK (Palace Leas and Park Grass) have not changed consistently in recent decades, *Glob. Change Biol.*, 15, 1739-1754, 2009.
- 930
- IPCC 1996. Climate change 1995. Impacts, adaptation and mitigation of climate change: scientific, technical analysis, contribution of working group II to the 2nd assessment reports of the IPCC. 1996. Intergovernmental Panel on Climate Change and Cambridge University Press Cambridge UK.
- 935
- IPCC 2006a . Guidelines for National Greenhouse Gas Inventories, Agriculture, Forestry and Other Land Use. Chapter 10: Emissions from Livestock and Manure Management. Intergovernmental Panel on Climate Change (IPCC), Volume 4. 2006. Institute for Global Environmental Strategies, Tokyo, Japan.
- 940
- IPCC 2006b. Guidelines for National Greenhouse Gas Inventories, Agriculture, Forestry and Other Land Use. Chapter 11: N<sub>2</sub>O Emissions from Managed Soils, and CO<sub>2</sub> Emissions from Lime and Urea Application. Intergovernmental Panel on Climate Change (IPCC) Volume 4. 2013. Institute for Global Environmental Strategies, Tokyo, Japan.
- Janssens, I. A., Freibauer, A., Ciais, P., Smith, P., Nabuurs, G. J., Folberth, G.,

- 945 Schlamadinger, B., Hutjes, R. W. A., Ceulemans, R., Schulze, E. D., Valentini, R., and Dolman, A.J.: Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic CO<sub>2</sub> emissions, *Science*, 300.5625, 1538-42, 2003.
- 950 Jones, S. K., Rees, R. M., Kosmas, D., Ball, B. C., and Skiba, U. M.: Carbon sequestration in a temperate grassland; management and climatic controls, *Soil Use Manage.*, 22.2, 132-42, 2006.
- Jones, S. K., Rees, R. M., Skiba, U. M., and Ball, B. C.: Influence of organic and mineral N fertiliser on N<sub>2</sub>O fluxes from a temperate grassland, *Agr. Ecosyst. Environ.*, 121.1-2, 74-83, 2007.
- 955 Jones, S. K., Famulari, D., Di Marco, C. F., Nemitz, E., Skiba, U. M., Rees, R. M., and Sutton, M.A.: Nitrous oxide emissions from managed grassland: a comparison of eddy covariance and static chamber measurements, *Atmospheric Measurement Techniques*, 4.10, 2179-94, 2011.
- 960 Kim, J., Verma, S. B., and Clement, R. J.: Carbon-Dioxide Budget in A Temperate Grassland Ecosystem. *J. Geophys. Res-Atmos.*, 97.D5, 6057-63, 1992.
- 965 Kindler, R., Siemens, J., Kaiser, K., Walmsley, D. C., Bernhofer, C., Buchmann, N., Cellier, P., Eugster, W., Gleixner, G., Grunwald, T., Heim, A., Ibrom, A., Jones, S. K., Jones, M., Klumpp, K., Kutsch, W., Larsen, K. S., Lehuger, S., Loubet, B., McKenzie, R., Moors, E., Osborne, B., Pilegaard, K., Rebmann, C., Saunders, M., Schmidt, M. W. I., Schrupf, M., Seyfferth, J., Skiba, U., Soussana, J. F., Sutton, M.A., Tefs, C., Vowinkel, B., Zeeman, M. J., and Kaupenjohann, M.: Dissolved carbon leaching from soil is a crucial component of the net ecosystem carbon balance, *Glob. Change Biol.*, 17.2, 1167-85, 2011.
- 970 Kramberger, M., Podvršnik, A., Gselman, V., Šuštar, J., Kristl, M., Muršec, M., Lešnik, D., Škorjanc: The effects of cutting frequencies at equal fertiliser rates on bio-diverse permanent grassland: Soil organic C and apparent N budget. *Agriculture, Ecosystems & Environment*, Volume 212, Pages 13-20B, 2015.
- 975 Lambe, N.R., Navajas, E.A, McLean, K.A., Simm, G., Bünger, L.: Changes in carcass traits during growth in lambs of two contrasting breeds, measured using computer tomography, *Livestock Science*, 107, 37-52, 2007.
- 980 Laws, J. A., Falge, E., Gu, L., Baldocchi, D. D., Bakwin, P., Berbigier, P., Davis, K., Dolman, A. J., Falk, M., Fuentes, J. D., Goldstein, A., Granier, A., Grelle, A., Hollinger, D., Janssens, I. A., Jarvis, P., Jensen, N. O., Katul, G., Mahli, Y., Matteucci, G., Meyers, T., Monson, R., Munger, W., Oechel, W., Olson, R., Pilegaard, K., Paw, K. T., Thorgeirsson, H., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Comparison of grassland management systems for beef cattle using self-contained farmlets: effects of contrasting nitrogen inputs and management strategies on nitrogen budgets, and herbage and animal production, *Agr. Ecosyst. Environ.*, 80.3, 243-54, 2000.
- 985

Leahy, P., Kiely, G., and Scanlon, T. M.: Managed grasslands: A greenhouse gas sink or source? *Geophys. Res. Lett.*, 31.20, 2004.

990 Lettens, S., Van Orshoven, J., Van Wesemael, B., De Vos, B., and Muys, B.: Stocks and fluxes of soil organic carbon for landscape units in Belgium derived from heterogeneous data sets for 1990 and 2000, *Geoderma*, 127.1-2, 11-23, 2005 a.

995 Lettens, S., Van Orshoven, J., Van Wesemael, B., Muys, B., and Perrin, D.: Soil organic carbon changes in landscape units of Belgium between 1960 and 2000 with reference to 1990, *Glob. Change Biol.*, 11.12, 2128-40, 2005b.

Luo, J., Tillman, R. W., and Ball, P. R.: Grazing effects on denitrification in a soil under pasture during two contrasting seasons, *Soil. Biol. Biochem.*, 31.6, 903-12, 1999.

1000 Marchand, R.F.: Ueber die chemische Zusammensetzung der Knochen. *Journal für praktische Chemie* 27, 83–97, 1842. DOI: 10.1002/prac.18420270117

1005 McKenzie, R. M., Özel, M. Z., Cape, J. N., Drewer, J., Dinsmore, K. J., Nemitz, E., Hamilton, J. F., Sutton, M. A., Gallagher, M. W. and Skiba, U.: The import and export of organic nitrogen species at a Scottish ombrotrophic peatland, *Biogeosciences Discussions*, 12, 515-554, 2015.

McTiernan, K. B., Jarvis, S.C., Scholefield, D., and Hayes, M. H. B.: Dissolved organic carbon losses from grazed grasslands under different management regimes, *Water Res.*, 35.10, 2565-69, 2001.

1010 Molina-Herrera S., Haas, E., Klatt, S., Kraus, D., Augustin, J., Magliulo, V., Tallec, T., Ceschia, E., Ammann C., Loubet, B., Skiba, U., Jones, S., Brümmer, C., Butterbach-Bahl, K., Kiese, R.: A modelling study on mitigation of N<sub>2</sub>O emissions and NO<sub>3</sub> leaching at different agricultural sites across Europe using LandscapeDNDC, *Science of the Total Environment*, 553, 128–140, 2016.

1015 Mudge, P. L., Wallace, D. F., Rutledge, S., Campbell, D. I., Schipper, L. A., Hosking, C. L.: Carbon balance of an intensively grazed temperate pasture in two climatically contrasting years, *Agr. Ecosyst. Environ.*, 144.1, 271-80, 2011.

1020 Navajas, E. A., Richardson, R. I., Fisher, A. V., Hyslop, J. J., Ross, D. W., Prieto, N., Simm, G. and Roehe, R.: Predicting beef carcass composition using tissue weights of a primal cut assessed by computed tomography, *Animal*, 4:11, pp 1810–1817, 2010. doi:10.1017/S1751731110001096

1025 Neeteson, J. J. and Hassink, J.: Nitrogen Budgets of Three Experimental and Two Commercial Dairy Farms in the Netherlands. In: M.K. van Ittersum & S.C. van de Geijn (Eds.), *Perspectives for Agronomy. Adopting ecological principles and managing resource use.* Elsevier Science

BV, Amsterdam. *Developments in Crop Science* 25, pp. 171-178. 1997.

1030 Owens, L. B., Edwards, W. M., and Vankeuren, R. W.: Groundwater Nitrate Levels Under Fertilized Grass and Grass-Legume Pastures. *J. Environ. Qual.*, 23.4, 752-58, 1994.

1035 Phoenix, G. K., Hicks, W.K., Cinderby, S., Kuylenstierna, J.C.I., Stock, W. D., Dentener, F. J., Giller, K. E., Austin, A. T., Lefroy, R. D. B., Gimeno, B. S., Ashmore, M. R., Ineson, P.: Atmospheric nitrogen deposition in world biodiversity hotspots: the need for a greater global perspective in assessing N deposition impacts. *Global Change Biology*, 12, pp. 470–476, 2006.

1040 Powlson, D.S., Whitmore, A.P. and Goulding, K.W.T.: Soil carbon sequestration to mitigate climate change: A critical re-examination to identify the true and the false. *European Journal of Soil Science*, 62, 42-55, 2011.

Prescher, A. K., Grunwald, T. and Bernhofer, C.: Land use regulates carbon budgets in eastern Germany: From NEE to NBP, *Agr. Forest Meteorol.*, 150.7-8, 1016-25, 2010.

1045 Rees, R. M., Bingham, I. J., Baddeley, J. A., and Watson, C.A.: The role of plants and land management in sequestering soil carbon in temperate arable and grassland ecosystems. *Geoderma*, 128.1-2, 130-54, 2005.

1050 Roche J.: *The International Wool Trade*. Woodhead Publishing, 1995.

Rudaz, A. O., Walti, E., Kyburz, G., Lehmann, P., and Fuhrer, J.: Temporal variation in N<sub>2</sub>O and N-2 fluxes from a permanent pasture in Switzerland in relation to management, soil water content and soil temperature, *Agr. Ecosyst. Environ.*, 73.1, 83-91, 1999.

1055 Ryden, J. C., Ball, P. R., and Garwood, E. A.: Nitrate Leaching from Grassland, *Nature* 311.5981, 50-53, 1984a.

1060 Rytter, R. M., Rytter, L., Hogbom, L.: Carbon sequestration in willow (*Salix* spp.) plantations on former arable land estimated by repeated field sampling and C budget calculation. *Biomass and Bioenergy*, 83, 483-492, 2015.

1065 Schipper, L. A., Baisden, W. T., Parfitt, R. L., Ross, C., Claydon, J. J., and Arnold, G.: Large losses of soil C and N from soil profiles under pasture in New Zealand during the past 20 years, *Glob. Change Biol*, 13.6, 1138-44, 2007.

Scholefield, D., Lockyer, D. R., Whitehead, D. C., and Tyson, K. C. A.: A Model to Predict Transformations and Losses of Nitrogen in Uk Pastures Grazed by Beef-Cattle, *Plant Soil*, 132.2, 165-77, 1991.

- 1070 Scholefield, D., Tyson, K. C., Garwood, E.A., Armstrong, A.C., Hawkins, J., and Stone, A. C.: Nitrate Leaching from Grazed Grassland Lysimeters - Effects of Fertilizer Input, Field Drainage, Age of Sward and Patterns of Weather, *J. Soil Sci.*, 44.4, 601-13, 1993.
- Schuman, G. E., Janzen, H. H., and Herrick, J. E.: Soil carbon dynamics and potential carbon  
1075 sequestration by rangelands, *Environ. Pollut.*, 116.3, 391-96, 2002.
- Schrumpf, M., Schulze, E. D., Kaiser, K., and Schumacher, J.: How accurately can soil organic carbon stocks and stock changes be quantified by soil inventories? *Biogeosciences*, 8, 1193-1212, 2011.
- 1080 Schulze, E.D., Luysaert, S., Ciais, P., Freibauer, A., Janssens, I.A., Soussana, J.F., Grace, J., Levin, I., Thiruchittampalam, B., Heimann, M., Dolman, A.J., Valentini, R., Bousquet, P., Peylin, P., Peters, W., Rödenbeck, C., Etiope, G., Vuichard, N., Wattenbach, M., Nabuurs, G.J., Poussi, Z., Nieschulze, J., Gach, J.H.: Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance.  
1085 *Nature Geoscience* 2: 842-850, 2009.
- Scottish Agricultural College, Farm management handbook SAC; ed. Linda Chadwick, Edinburgh, 1995.
- 1090 Scottish Agricultural College, Technical Note TN652: Fertiliser recommendations for grasslands. A., Sinclair, P.A., Shipway, B., Crooks, 2013.
- Siemens, J.: The European carbon budget: A gap, *Science*, 302.5651, 1681, 2003.
- 1095 Skiba, U., Drewer, J., Tang, Y. S., van Dijk, N., Helfter, C., Nemitz, E., Famulari, D., Cape, J. N., Jones, S. K., Twigg, M., Pihlatie, M., Vesala, T., Larsen, K. S., Carter, M. S., Ambus, P., Ibrom, A., Beier, C., Hensen, A., Frumau, A., Erisman, J. W., Br'uggemann, N., Gasche, R., Butterbach-Bahl, K., Neftel, A., Spirig, C., Horvath, L., Freibauer, A., Cellier, P., Laville, P., Loubet, B., Magliulo, E., Bertolini, T., Seufert, G., Andersson, M., Manca, G., Laurila, T., Aurela, M., Lohila A., Zechmeister-Boltenstern, S., Kitzler, B., Schaufler, G., Siemens, J., Kindler, R., Flechard, C., and Sutton, M. A.:  
1100 Biosphere atmosphere exchange of reactive nitrogen and greenhouse gases at the NitroEurope core flux measurement sites: Measurement strategy and first annual data set, *Agr. Ecosys. Environ.*, 133, 139–149, 2009
- 1105 Skiba U, Drewer J, Tang Y, van Dijk N, Helfter C, Nemitz E, Famulari D, Cape J, Jones S, Twigg M.: Biosphere-atmosphere exchange of reactive nitrogen and greenhouse gases at the NitroEurope core flux measurement sites: Measurement strategy and first data sets. *Agriculture, Ecosystems & Environment* 133 (3- 4): 139-149, 2009. doi: 10.1016/j.agee.2009.05.018
- 1110 Skiba, U., Jones, S. K., Drewer, J., Helfter, C., Anderson, M., Dinsmore, K., McKenzie, R., Nemitz, E., and Sutton, M. A.: Comparison of soil greenhouse gas fluxes from extensive and

intensive grazing in a temperate maritime climate, *Biogeosciences* 10, 1231-1241, 2013.

Smith, P. : How long before a change in soil organic carbon can be detected? *Glob. Change Biol.*, 10.11, 1878-83, 2004

1115

Smith, P., Haberl, H., Popp, A., Erb, K.-H., Lauk, C., Harper, R., et al.: How much land based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? *Global Change Biology*, 19, 2285–2302, 2013.

1120

Smith, P.: Do grasslands act as a perpetual sink for carbon? *Glob. Change Biol.*, 20, 2708–2711, 2014.

Soussana, J. F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T., and Arrouays, D.: Carbon cycling and sequestration opportunities in temperate grasslands, *Soil Use Manage.*, 20, 219-30, 2004.

1125

Soussana, J. F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E., Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A., Horvath, L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A., Raschi, A., Baronti, S., Rees, R.M., Skiba, U., Stefani, P., Manca, G., Sutton, M., Tubaf, Z., and Valentini, R.: Full accounting of the greenhouse gas (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>) budget of nine European grassland sites, *Agr. Ecosyst. Environ.*, 121.1-2, 121-34, 2007.

1130

Soussana, J. F., Tallec, T., and Blanfort, V.: Mitigating the greenhouse gas balance of ruminant production systems through carbon sequestration in grasslands, *Animal*, 4.3, 334-50, 2010.

1135

Soussana, J. F. and Lemaire, G.: Coupling carbon and nitrogen cycles for environmentally sustainable intensification of grasslands and crop-livestock systems, *Agr. Ecosyst. Environ.*, 190 9–17, 2014.

1140

Stewart, A. A., Little, S. M., Ominski, K.H., Wittenberg, K. M., and Janzen, H. H.: Evaluating greenhouse gas mitigation practices in livestock systems: an illustration of a whole-farm approach, *J. Agr. Jci.*, 147, 367-82, 2009.

1145

Sutton, M. A., Tang, Y. S., Miners, B., and Fowler, D.: A New Diffusion Denuder System for Long-Term, Regional Monitoring of Atmospheric Ammonia and Ammonium, *Water Air Soil Poll.: Focus*, 1.5-6, 145-56, 2001.

1150

Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach-Bahl, K., Cellier, P., de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S., Laville, P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C. R., Herrmann, B., Calanca, P. L., Schjoerring, J. K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B. A., Tietema, A., Peñuelas, J., Kesik, M., Brueggemann, N., Pilegaard, K., Vesala, T., Campbell, C.L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P.,

Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, *Environ. Pollut.*, 150, 125–139, 2007.

1155

Sutton M.A., Oenema, O., Erisman, J.W., Leip, A., van Grinsven, H. and Winiwarten, W.: Too much of a good thing. *Nature* 472, 159-161, 2011. doi:10.1038/472159a

1160

Tyson, K. C., Garwood, E. A., Armstrong, A.C., and Scholefield, D.: Effects of Field Drainage on the Growth of Herbage and the Liveweight Gain of Grazing Beef-Cattle, *Grass Forage Sci.*, 47.3, 290-301, 1992.

1165

Twigg, M. M., House, E., Thomas, R., Whitehead, J., Phillips, G.J., Famulari, D., Fowler, D., Gallagher, M.W., Cape, J.N., Sutton, M.A., Nemitz, E.: Surface/atmosphere exchange and chemical interactions of reactive nitrogen compounds above a manured grassland *Agricultural and Forest Meteorology*, Volume 151, Issue 12, , Pages 1488-1503, 2011.

1170

Van der Hoek, K. W.: Nitrogen efficiency in global animal production, *Environ. Pollut.*, 102, 127-32, 1998.

Van der Salm, C., Doling, J., Heinen, M., and Velthof, G. L.: Estimation of nitrogen losses via denitrification from a heavy clay soil under grass, *Agr. Ecosyst. Environ.*, 119, 311-319, 2007.

1175

Velthof, G. L. and Oenema, O.: Nitrous oxide fluxes from grassland in the Netherlands .2. Effects of soil type, nitrogen fertilizer application and grazing, *Eur. J. Soil Sci.*, 46.4, 541-49, 1995.

1180

Watson, C. J., Jordan, C., Kilpatrick, D., McCarney, B., and Stewart, R.: Impact of grazed grassland management on total N accumulation in soil receiving different levels of N inputs. *Soil Use Manage.*, 23.2, 121-128, 2007.

Whitehead, D. C., Pain, B. F., and Ryden J. C.: Nitrogen in UK grassland agriculture, *J. R. Agric. Soc.*, 147, 190-201. 1986.

1185

Zhang, Y., Xu, M., Chen, H. and Adams, J.: Global Pattern of NPP to GPP Ratio Derived from MODIS Data: Effects of Ecosystem Type, Geographical Location and Climate. *Global Ecology and Biogeography* 18 (3): 280–90, 2009. doi:10.1111/j.1466-8238.2008.00442.x.

**Table 1.** Average annual stocking rates [LSU\* ha<sup>-1</sup> y<sup>-1</sup>].

	2002	2003	2004	2005	2006	2007	2008	2009	2010
Heifers	0.12	0.38	0.05	0.15	0.27	0	0	0	0
Ewes	0.14	0.16	0.82	0.56	0.51	0.68	0.68	0.61	0.53
Lambs	0	0	0.12	0.12	0.12	0.23	0.14	0.11	0.12
all animals	0.27	0.54	0.99	0.83	0.90	0.91	0.83	0.72	0.65

\* LSU stands for Live stock units

**Table 2.** Weather characteristics of each measurement year.

	2002	2003	2004	2005	2006	2007	2008	2009	2010
Annual mean temperature [°C]	9.2	9.6	8.9	8.8	9.3	9.1	8.6	8.9	8.3
Maximum temperature [°C]	23.6	29.5	27.4	30.4	26.6	21.4	23.5	28.0	24.0
Minimum temperature [°C]	-5.1	-8.4	-4.9	-6.6	-5.5	-7.5	-5.8	-7.8	-10.3
Annual rainfall [mm]	1238	680	1169	1028	1120	904	1065	744	575
Soil water content [% by volume]	36.9	31.0	40.3	45.2	36.6	37.7	41.5	39.4	-
Water filled pore space [%]	68.0	57.2	74.3	83.3	67.5	69.5	76.5	72.6	-
Length of growing season* [days]	180	196	156	177	151	186	193	242	226

\* The plant growing season begins and ends with periods of consecutive days, where daily temperatures average more than 5 °C without any five-day spells of temperatures below 5 °C.

**Table 3.** Nitrogen budget and balance for each measurement year and average values, confidence intervals at  $p > 0.95$  (CI) and systematic uncertainties (uncert.) for 2002-2010 [ $\text{g N m}^{-2} \text{y}^{-1}$ ]. Negative numbers represent uptake while positive numbers represent loss of N from this grassland ecosystem. Letters indicate data published in previous publications.

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2002-2010		
										average	CI	uncert.
Organic fertilisation	0	0	-6.9	-15.8	0	0	0	0	0	-2.5	3.6	0.2
Inorganic fertilisation	-20.6	-16.0	-11.0	-17.3	-22.4	-17.3	-25.9	-25.0	-19.0	-19.4	3.1	0.2
Wet deposition	-0.4	-0.6	-0.6	-0.7	-0.6	-0.6	-0.5	-0.4	-0.4	-0.5	0.1	0.2
<sup>a</sup> Dry deposition	-0.5	-0.4	-0.3	-0.3	-0.2	-0.3	-0.2	-0.2	-0.2	-0.3	0.1	0.2
Harvest	5.0	4.7	0	0	0	0	0	0	0	1.1	1.4	0.2
Meat (incl. bones)	0.2	0.6	2.3	3.1	2.6	2.9	1.8	1.3	1.5	1.80	0.7	0.2
Wool	0	0	0.6	0.5	0.4	0.7	0.7	0.2	0.2	0.4	0.2	0.0
<sup>b</sup> Leaching	14.9.0	0.1	0.1	4.6	10.6	4.2	5.6	2.6	5.0	5.3	3.1	1.71
N <sub>2</sub>	3.7	2.2	1.3	1.7	2.8	3.0	3.3	4.1	3.6	2.9	0.6	0.8
<sup>c</sup> N <sub>2</sub> O	1.1	0.1	0.1	0.4	0.9	1.3	0.8	0.4	0.4	0.6	0.3	0.2
NO <sub>x</sub> (soil)	0.3	0.1	0	0.1	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.1
NO <sub>x</sub> ,NH <sub>3</sub> (inorg.fert.)	2.1	1.6	1.1	1.7	2.2	1.7	2.6	2.5	1.9	1.9	0.3	0.6
NO <sub>x</sub> ,NH <sub>3</sub> (org.fert.)	0	0	1.4	3.2	0	0	0	0	0	0.5	0.7	0.3
NO <sub>x</sub> ,NH <sub>3</sub> (excretion)	0.5	0.8	1.5	1.6	1.5	2.4	2.1	1.6	1.5	1.5	0.4	0.7
N balance	6.4	-6.9	-10.4	-17.2	-1.9	-1.9	-9.5	-12.7	-5.6	-6.6	4.4	2.2

<sup>a</sup>Flecharad et al. 2011: Dry deposition, modelled average value of the two years 2007/2008

<sup>b</sup>Molina-Herrera et al. 2016: N leaching modelled 2005-2010

<sup>c</sup>Molina-Herrera et al. 2016: N<sub>2</sub>O fluxes modelled 2005-2010.

<sup>d</sup>Di Marco et al. 2004: N<sub>2</sub>O fluxes measured by eddy covariance (half hourly) June 2002 to June 2003

<sup>e</sup>Jones et al. 2011: N<sub>2</sub>O fluxes measured by eddy covariance (half hourly) and chambers (hourly) during measurement campaigns in June 2003, March/May/July 2007 and May/July 2008.

<sup>f</sup>Flecharad et al. 2007: annual N<sub>2</sub>O fluxes measured by eddy covariance in 2002/2003 and by chambers in 2004

<sup>g</sup>Skiba et al. 2013: annual N<sub>2</sub>O fluxes measured by chambers from Jan. 2007 – Sept. 2010

**Table 4.** Carbon budget and balance for each measurement year and average values, confidence intervals at  $p > 0.95$  (CI) and systematic uncertainties (uncert.) for 2002-2010 [ $\text{g C m}^{-2} \text{y}^{-1}$ ]. Negative numbers represent uptake, while positive numbers represent loss of C from the grassland ecosystem. Letters indicate data published in previous publications.

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2002-2010		
											average	CI
GPP	-2162.9	-1982.0	-2111.4	-1662.4	-982.1	-1722.7	-1441.2	-1722.4	-2015.4	-1755.8	244.4	105.3
TER	1726.9	1725.9	2183.2	1638.5	972.1	1606.7	1324.0	1116.7	1547.0	1537.9	236.2	92.3
NPP	-1081.5	-991.0	-1055.7	-831.2	-491.1	-861.3	-720.6	-861.2	-1007.7	-877.9	122.2	-52.8
<sup>a</sup> CO <sub>2</sub> (NEE)	-436.0	-256.1	71.8	-24.0	-10.0	-115.9	-117.1	-605.7	-468.4	-217.9	154.5	80.0
Organic fert.	0.0	0.0	-55.4	-171.8	0.0	0.0	0.0	0.0	0.0	-25.2	37.8	5.0
Harvest	270.6	169.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	48.9	65.5	5.4
Meat (incl. bones)	0.9	2.9	11.4	15.6	12.9	14.3	9.0	6.3	7.3	9.0	3.3	10
Wool	0.0	0.0	1.7	1.5	1.3	2.1	2.0	0.7	0.5	1.1	0.5	0.1
<sup>b</sup> Leaching	25.1	7.0	22.1	18.7	19.4	15.4	17.0	6.8	14.3	16.4	4.3	5.26
CH <sub>4</sub> (organic fert.)	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<sup>c</sup> CH <sub>4</sub> (soil)	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1
CH <sub>4</sub> (excretion)	0.0	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.0	0.0
CH <sub>4</sub> (enteric ferm.)	1.5	3.2	5.7	4.8	5.2	5.2	4.8	4.1	3.8	4.3	0.8	0.9
C balance (NBP)	-137.8	-73.3	57.6	-154.9	29.1	-78.7	-84.1	-587.7	-440.3	-163.3	139.5	16.0

<sup>a</sup>Soussana et al. 2007: NEE July 2002- Dec 2004

<sup>a</sup>Skiba et al. 2013: NEE 2007-2010

<sup>a</sup>Kindler et al. 2011: NEE average multiyear value 2004-2007

<sup>b</sup>Kindler et al. 2011: C leaching losses October 2006- Sept 2008 (Slope value corresponds to data used in this publication).


<sup>c</sup>Skiba et al. 2013: CH<sub>4</sub> (soil) 2007-2010

**Table 5.** Annual N<sub>2</sub>O exchange, total N input by fertiliser (mineral and organic) and N<sub>2</sub>O emission factors, expressed as percentage of total N inputs in 2002-2010.

	N <sub>2</sub> O flux [g N m <sup>-2</sup> y <sup>-1</sup> ]	Total N input [g N m <sup>-2</sup> y <sup>-1</sup> ]	EF [%]
2002	1.14	20.60	5.5
2003	0.14	15.98	0.9
2004	0.11	11.00	0.6
2005	0.36	17.25	1.1
2006	0.88	22.43	3.9
2007	1.25	17.25	7.2
2008	0.84	25.93	3.2
2009	0.41	24.95	1.6
2010	0.35	18.98	1.9


**Table 6.** N and C soil stocks (g N or C m<sup>-2</sup>) in May 2004 and May 2011 and budgets (g N or C m<sup>-2</sup> y<sup>-1</sup>) over 7 years based on repeated soil N and C stock inventories and flux budget calculations (January 2004 - December 2010). Soil stock changes are based on a soil mass of 800 kg m<sup>-2</sup>, which corresponds to approximately 60 cm depth. The flux budgets are averages for the years 2004 – 2010 (Table 4 & 5). Numbers in brackets represent confidence intervals. Negative numbers are sinks.

	Nitrogen	Carbon
soil stocks in 2004	840.68 (11.89)	12026.05 (190.19)
soil stocks in 2011	870.02 (14.14)	11824.87 (187.84)
soil stock change	-4.51 (2.64)	29.08 (38.19)
flux budget	-8.44 (4.21)	-179.7 (180)




**Figure 1.** Stocking density (a), nitrogen (b) and carbon (c) input and export from inorganic and organic fertiliser and harvest from 2002-2010. LSU stands for livestock unit, where 1 livestock unit has a standard live weight of 600 kg head<sup>-1</sup>.


a) N budget (% total N input); harvested and grazed




b) N budget (% total N input); grazed only



**Figure 2.** Mean annual nitrogen budget for Easter Bush, showing the fate of total N input (fertiliser and deposition) in (a) years when harvested for silage (2002 and 2003) and (b) in years when only grazing took place (2004–2010). The residual term includes all the error in the budget calculation, as well as any net accumulation of soil organic nitrogen.



**Figure 3.** Mean annual carbon budget for Easter Bush, showing the fate of net primary productivity (NPP) in (a) years when harvested for silage (2002 and 2003) and (b) in years when only grazing took place (2004–2010). Heterotrophic respiration includes the respiration of soil microbes, cows and sheep. The residual term includes all the error in the budget calculation, as well as any net accumulation of soil organic carbon.



**Figure 4.** Average greenhouse gas fluxes, net GHG exchange (NGHGE) and attributed net GHG balance (NGHGB, includes  $FC_{org\ fert}$ ,  $FC_{animal}$ ,  $FC_{leaching}$ ,  $FC_{harvest}$ ) for 2002-2010. Positive values correspond to losses and negative values to storage of greenhouse gases to and from the grassland system, respectively. The CH<sub>4</sub> component comprises CH<sub>4</sub> fluxes from enteric fermentation, animal excretion, slurry application and soil exchange, while the N<sub>2</sub>O component is the N<sub>2</sub>O flux from the soil. CO<sub>2</sub> represent the Net Ecosystem Exchange (NEE). Global warming potentials of 298 and 25 were used for N<sub>2</sub>O and CH<sub>4</sub> respectively, using a time horizon of 100 yrs (IPCC, 2013). Thin error bars represent variations (confidence intervals at  $p > 0.95$ ) between years, while thick error bars represent the systematic uncertainty of each value.