

Interactive comment on “Long-term nutrient fertilization and the carbon balance of permanent grassland: any evidence for sustainable intensification?” by Dario A. Fornara et al.

L. Schipper (Referee)

Schipper@waikato.ac.nz

Received and published: 9 July 2016

Thanks for the responses.

In conclusions I dont agree that the previous work that I did suggests in the long-term pasture soils lose carbon. I think there needs to be a broader discussion here 'some' pasture systems will gain C and some will lose with time and many/most will likely be at steady state. Our previous work (Schipper et al 2014) demonstrated that the majority of soils in long-term pasture on flat land in New Zealand appear at steady state except for a couple of soil orders Gley (losses attributed to drainage and enhanced C respiration, as was suggested in Belgium) and Allophanic soils (we do not have an explanation for this loss).

[Printer-friendly version](#)

[Discussion paper](#)



In contrast pasture on hill country was gaining C, possibly due to recovery after past sheet wash erosion but not known.

So in the end we we need to understand the variety of pasture management systems that may increase or decrease soil C but a general statement that carbon increases or decreases in pastures is not really supported, it depends strongly on past land use and current management.

Interactive comment on Biogeosciences Discuss., doi:10.5194/bg-2016-224, 2016.

BGD

Interactive
comment

Printer-friendly version

Discussion paper

