

1 **Hydrologically transported dissolved organic carbon influences soil respiration in a tropical**
2 **rainforest**

3 **Running title: DOC influences soil respiration**

4 Author: W.-J. Zhou ^{1,2,3}, H.-Z. Lu ^{1,2,3}, Y.-P. Zhang ^{1,2*}, L.-Q. Sha ^{1,2*}, D. Schaefer ^{1,2}, Q.-H. Song ^{1,2,3},
5 Y. Deng ^{1,2}, X.-B. Deng^{1,2}

6 Affiliation:

7 ^{1.} Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden,
8 Chinese Academy of Sciences, Mengla, Yunnan 666303, China

9 ^{2.} Xishuangbanna Station for Tropical Rain Forest Ecosystem Studies, Chinese Ecosystem
10 Research Net, Mengla, Yunnan 666303, China

11 ^{3.} Graduate University of the Chinese Academy of Sciences, Beijing 100039, China

12 **Corresponding author:**

13 Y.-P. Zhang, Tel: +86-871-65160904, Fax: +86-871-65160916, E-mail: yipingzh@xtbg.ac.cn

14 L.-Q.Sha, Tel: +86-871-65160904, Fax: +86-871-65160916, E-mail: shalq@xtbg.ac.cn

15 **Keywords:**

16 Dissolved organic carbon (DOC), Soil temperature, Soil water content, Soil respiration, Tropical
17 rainforest

18 **Paper type:**

19 Primary research articles

Abstract

To better understand the effect of the dissolved organic carbon (DOC) transported by hydrological processes (rainfall, throughfall, litter leachate, and surface soil water (0–20 cm)) on soil respiration in tropical rainforests, we detected the DOC flux in rainfall, throughfall, litter leachate, and surface soil water (0–20 cm), compared the seasonality of $\delta^{13}\text{C}_{\text{DOC}}$ in each hydrological process, and $\delta^{13}\text{C}$ in leaves, litter, and surface soil, and analyzed throughfall, litter leachate, and surface soil water (0–20 cm) effect on soil respiration in a tropical rainforest in Xishuangbanna, southwest China. Results showed: The surface soil intercepted $94.4 \pm 1.2\%$ of the annual litter leachate DOC flux and is a sink for DOC. The throughfall and litter leachate DOC fluxes amounted to 6.81% and 7.23% of the net ecosystem exchange, respectively, indicating that the DOC flux through hydrological processes is an important component of the carbon budget, and may be an important link between hydrological processes and soil respiration in a tropical rainforest. Even the variability in soil respiration is more dependent on the hydrologically transported water than DOC flux insignificantly, soil temperature and soil water content (at 0–20 cm). The difference in $\delta^{13}\text{C}$ between the soil, soil water (at 0–20 cm), throughfall, and litter leachate indicated that DOC is transformed in the surface soil and decreased the sensitivity indices of soil respiration of DOC flux to water flux, which suggests that soil respiration is more sensitive to the DOC flux in hydrological processes, especially the soil water DOC flux, than to soil temperature or soil moisture.

38

39 **1. Introduction**

40 Dissolved organic carbon (DOC), the most active form of fresh carbon, stimulates microbial activity
41 and affects CO₂ emissions from the surface soil (Bianchi, 2011, Chantigny, 2003, Cleveland *et al.*,
42 2006). This indicates that the proportion of DOC that leaches from the soil is a crucial component of
43 the carbon balance (Kindler *et al.*, 2011, Stephan *et al.*, 2001), which is also estimated as the high ratio
44 of DOC flux to net ecosystem exchange (NEE) in forests, grasslands, and croplands (Sowerby *et al.*,
45 2010). The DOC from water-extractable soil carbon is regenerated quickly and functions as an
46 important source of substrate for soil respiration (SR), especially microbial heterotrophic respiration
47 (HR) (Cleveland *et al.*, 2004, Jandl and Sollins, 1997, Schwendenmann and Veldkamp, 2005), which
48 contributes more to SR than does autotrophic respiration. Laboratory studies have shown that DOC
49 also plays an important role in SR in the surface soil (De Troyer *et al.*, 2011, Fröberg *et al.*, 2005, Qiao
50 *et al.*, 2013). However the mechanisms underlying the effects of DOC on the carbon budget and SR in
51 the field remain unclear.

52 Hydrological processes that transport DOC, such as throughfall and litter leachate, are important
53 sources of DOC in surface soil water (De Troyer *et al.*, 2011, Kalbitz *et al.*, 2000, Kalbitz *et al.*, 2007,
54 Kindler *et al.*, 2011). The soil retains most of the DOC that reaches the soil surface from the throughfall
55 and litter leachate (Chuyong *et al.*, 2004, Dezzio and Chacón, 2006, Liu and Sheu, 2003, Liu *et al.*,
56 2008, McJannet *et al.*, 2007, Schrumpf *et al.*, 2006, Zimmermann *et al.*, 2007). Qiao *et al.* (2013)
57 suggested that the addition of labile organic carbon increases the decomposition of the native soil
58 organic carbon (SOC) by exerting a priming effect, and augments the CO₂ emissions in subtropical

forests. Because of the massive rainfall in tropical rainforests, more DOC flux is transported to the soil by throughfall and litter leachate than in other forests. The high temperature and leaching in tropical forests may mean that the fresh DOC from hydrological processes affects SR differently in tropical rainforests than in boreal and temperate forests (De Troyer *et al.*, 2011, Fröberg *et al.*, 2005, Qiao *et al.*, 2013). For this reason, research into the role of hydrologically transported DOC in the SR in tropical rainforest is essential.

The fate of DOC intercepted by the surface soil can be determined from variations in the DOC flux and $\delta^{13}\text{C}_{\text{DOC}}$ among soil water, soil, litter leachate, and throughfall. Based on the seasonal and source (canopy leaf, litter, or soil) differences in $\delta^{13}\text{C}$ (De Troyer *et al.*, 2011), $\delta^{13}\text{C}_{\text{DOC}}$ studies have shown that DOC transported from aboveground water and from the desorption of soil aggregates is retained in the surface soil by soil absorption or is involved in surface carbon biochemical dynamics through soil water leaching and microbiological activity (Comstedt *et al.*, 2007, De Troyer *et al.*, 2011, Fang *et al.*, 2009, Kindler *et al.*, 2011). This proposal has been confirmed in a laboratory leaching experiment simulating a temperate forest, as performed by Park *et al.* (2002), who reported that the cumulative amount of CO_2 evolved is positively related to the availability of carbon (Park *et al.*, 2002). Furthermore, fresh DOC fed to the surface soil influences soil CO_2 emissions in both the short term (3–14 days) and long term (month to years) (Davidson *et al.*, 2012). Therefore, several models of the surface soil carbon efflux indicate that DOC is a factor that influences CO_2 emissions (Blagodatskaya *et al.*, 2011, Guenet *et al.*, 2010, Yakov, 2010) based on recent research with controlled experiments. However, the natural mechanism underlying the effects of the hydrologically transported DOC flux on CO_2 emissions remains unclear. The precipitation rate, NEE, and litterfall are all high in tropical forests (Tan *et al.*, 2010, Zhang *et al.*, 2010), and several studies have shown that DOC plays an important role

in the carbon balance in these settings (Fontaine *et al.*, 2007, McClain *et al.*, 1997, Monteith *et al.*, 2007). Here, we investigate the relative contribution of hydrologically transported DOC to SR in a rainforest compared with the contributions of soil temperature and moisture, which has not been extensively studied until now.

Our study was performed in a tropical rainforest at Xishuangbanna in southwest China, on the northern edge of a tropical region. This forest has less annual rainfall (1557 mm), a smaller carbon sink (1667 kg C ha⁻¹) (Tan *et al.*, 2010, Zhang *et al.*, 2010), lower SR (5.34 kg CO₂ m⁻² yr⁻¹) (Sha *et al.*, 2005), and less litterfall (9.47 ± 1.65 Mg C ha⁻¹ yr⁻¹) (Tang *et al.*, 2010) than typical rainforests of the Amazon and around the equator. We hypothesized that throughfall and litter leachate DOC flux are important in carbon budget, and that hydrologically transported DOC significantly affects SR in the tropical rainforest at Xishuangbanna. To test these hypotheses, we determined the SR, HR, and DOC fluxes in the rainfall, throughfall, litter leachate, and surface soil water (0–20 cm depth), the seasonal variability in $\delta^{13}\text{C}$ (isotopic abundance ratio of ¹³C) in DOC ($\delta^{13}\text{C}_{\text{DOC}}$) and in the carbon pools in the soil, litter, and canopy leaves in this tropical forest.

2 Materials and methods

2.1 Study site

The study site is located at the center of the National Forest Reserve in Menglun, Mengla County, Yunnan Province, China (21°56'N, 101°15'E), and has suffered relatively little human disturbance. The weather in the study area is dominated by the north tropical monsoon and is influenced by the southwest monsoon, with an annual average temperature of 21.5 °C, annual average rainfall of 1557 mm, and average relative humidity of 86%. Based on the precipitation dynamics, the rainy season

occurs between May and October (with 84.1% of the total annual precipitation) and the dry season between November and April.

The dominant trees are *Terminalia myriocarpa* and *Pometia tomentosa*, which are typical tropical forest trees. Canopy height is about 45m, the land cover ratio is 100%, there are 311 species that diameter at breast height (DBH) is larger than 2cm (Cao *et al.*, 1996). The topographic slope is 12 °–18 °, and the soil type is oxisol, formed from Cretaceous yellow sandstone, with a pH of 4.5–5.5 and a clay content ($d < 0.002$ mm) of 29.5% in the surface soil (0–20 cm) (Tang *et al.*, 2007).

2.2 Experimental set-up

At the study plot (a 23.4 ha catchment), three rainfall collectors were set above the canopy on a 70 m eddy flux tower to collect rain samples. Each collector had a polytetrafluoroethylene (PTFE) funnel (2.5 cm diameter) connected to a brown glass bottle, which was rinsed with distilled water before each collection. There were four replicates of throughfall, litter leachate, and soil water (20 cm depth) respectively. All the collectors were set around the eddy flux tower randomly. All the collectors were distributed randomly around the eddy flux tower. The throughfall collectors were 200×40 cm² V-shaped tanks made of stainless steel. A PTFE tube connected the collector to a polyethylene sampling barrel. The litter leachate was collected in 40 cm × 30 cm × 2 cm PTFE plates. In the plate, we layered 100-, 20-, and 1-mesh silica sand from the bottom to the upper edge, to a depth of 2 cm, to ensure that the litterfall fragments did not reach the bottom of the plate and to filter the leachate. The bottom of the plate was curved into an arc shape, causing the leachate to flow together at the bottom funnel. The funnel was connected by a PTFE tube to a 10 L bottle further down the slope. The soil water collector was designed like the litter leachate collector. The collection system was buried in soil

at a depth of 20 cm along the surface slope. To reduce the disturbance from digging as much as possible, all the soil collectors were placed in holes that were approximately the size of the PTFE collector, and all soil was added from the bottom to the surface, layer by layer. All the soil water and litter leachate collectors were set in place 3 months before the samples were collected, to minimize the influence of their installation.

The water fluxes from rainfall and throughfall were estimated with an installed water-level recorder. The recorder was set to measure the average discharge at 30 min intervals. The daily and weekly water fluxes from rainfall and throughfall were calculated from the data recorded automatically between 08:00 and 08:00 on the following day (local time). The water fluxes from the litter leachate and soil water were determined daily by manual observation.

We set four 5 m × 5 m plots around the eddy flux tower to measure SR and HR using the trenching method. In each plot, three paired trenches and control treatments were used to detect both HR and SR. Each treatment covered an area of 50 × 50 cm². Most fine roots occur in the first 0–20 cm of soil and few occur below a depth of 50 cm in the soil of tropical rainforests. In each trenched treatment, a polyvinyl chloride panel was installed, and a 50-cm-deep trench was filled with *in situ* soil to protect root respiration during the trenching treatment.

The soil respiration was measured using a Li-820 system (Li-Cor Inc., Lincoln, NE, USA), which consisted of an infrared gas analyzer with a polyvinyl chloride chamber (diameter of 15 cm and height of 15.0 cm). A polyvinyl chloride collar (diameter of 15 cm and height of 5 cm) was installed in the forest floor to a depth of ~3 cm. All the leaf litter and small branches were left in the collar. Soil respirations were detected from 09:00 to 14:00 local time when was taken to represent respiration in that day (Sha et al. 2005, Yao et al. 2011) biweekly from February 2008 to February

2009.

Soil temperature and moisture

From 2008 to 2011, soil temperature and moisture at a depth of 5 cm were measured every 15 min with a Campbell Scientific data logger (Campbell Scientific, North Longan, Utah, USA) which was fixed to the eddy flux tower. The daily average soil temperature and moisture were calculated as the daily means of the data collected every 15 min.

During soil respiration observation period between February 2008 and January 2009, soil water content (0–12 cm) was detected by time-domain reflectometry (TDR100, Campbell Scientific, USA) in the soil close to every chamber. At the same time, the soil temperature (0–10 cm) and the air temperature were recorded with a needle thermometer.

Soil, leaf, and litter sampling

Soil (0–20 cm depth) near the soil water collectors, and leaf samples and litter samples from around the water collector were collected in August and October, 2010, and in January, March, and May, 2011.

The leaves of the dominant species were randomly picked from the canopy around the plots, and litter samples were collected from around the plots. Soil samples were collected with a steel soil sampler (diameter = 5 cm, height = 20 cm). All the leaf and litter samples were oven-dried to constant weight at 60 °C. After drying, the leaf and litter were ground and passed through a 1 mm screen. Wind-dried soil was manually broken by hand and sieved (100 mesh) to remove larger particles, roots, and visible soil fauna. Plant and soil samples were analyzed for total C and $\delta^{13}\text{C}$ values with an elemental analyzer (Elementar vario PYRO cube, Germany) coupled to an continuous flow system isotope ratio mass spectrometer (IsoPrime 100 Isotope Ratio Mass Spectrometer, Germany, EA-MS). Samples (1.00–3.00 mg plant samples and 10–40 mg soil sample dried and sieved through 100 mesh size) were wrapped in

a tin boat and loaded into the auto-sampler (EA3000, Eurovector, Milan, Italy) coupled to the EA-IRMS. The sample was flash combusted in a combustion reactor held at 1120°C. The produced CO₂ was separated by the CO₂ absorption column, and carried by helium to ion source for measurements. The reference CO₂ (>99.999%) flowed in at 420 seconds and lasted for 30 seconds. The isotopic results are expressed in standard notation ($\delta^{13}\text{C}$) in parts per thousand (‰) relative to the standard Pee Dee Belemnite:

$$\delta^{13}\text{C} = \left[\frac{{}^{13}\text{R}_{\text{sample}}}{{}^{13}\text{R}_{\text{standard}}} - 1 \right] \times 1000 \quad (1)$$

where R is the molar ratio $^{13}\text{C}/^{12}\text{C}$.

2.3 Water sampling and analysis

All the 24 h cumulative water samples were collected at the sampling sites between 08:00 and 10:00 (local time), following the procedure outlined by Zhou et al. (2013), using high-density polyethylene bottles. The sampling bottles were completely filled, allowing no headspace. After the bottles were washed with 3% HCl solution, they were rinsed with distilled water. Before sample collection, the bottles were pre-rinsed three times with the sample water. The study was performed over three full calendar years, from January 1, 2009, to December 31, 2011. The water samples were collected on the day following a rain event during the dry season and once a week during the rainy season in 2009, and once a week in 2010 and 2011. All the water samples were immediately transported to the laboratory in insulated bags to prevent DOC decomposition.

Based on the analytical method of Zhou *et al.* (2013), all the samples were vacuum-filtered through a 0.45 μm glass fiber filter (Tianjinshi Dongfang Changtai Environmental Protection Technology, Tianjin, China) and were pre-rinsed with deionized water and the sample water under vacuum. The

filtered samples were analyzed for DOC within 24 h of collection using a total organic carbon/total nitrogen (TOC/TN) analyzer (LiquiTOC II, Elemental Analyses System GmbH, Germany).

To analyze the water DOC isotopic $\delta^{13}\text{C}$ -DOC ($\delta^{13}\text{C}_{\text{DOC}}$), the samples were collected on the same day as the leaves, litter, and 0–20 cm soil samples were collected. Subsamples (500 mL) of the rain, throughfall, litter leachate, and soil water samples were passed through a 0.45 μm glass fiber filter and transferred to another 500 mL polyethylene terephthalate bottle. All the filtered water was frozen and placed in a freeze dryer until it was reduced to a fine powder. The $\delta^{13}\text{C}$ of the freeze-dried DOC was analyzed with a method similar to that for the plant and soil samples. Considering the lower C content, more sample amount (20–60mg) were weighted, the combustion temperature was set at 920°C, and the reference CO_2 flowed in at 475 seconds, later than for the soil and plant samples. The sample $\delta^{13}\text{C}$ abundance were calculated according to Eq (1).

2.4 Calculations and statistics

The correlations among the daily water flux and DOC concentration, SR, HR, soil moisture, and soil temperature from February 2008 to January 2009, and the weekly SR and HR rates and the amounts of DOC and water in 2009–2011 were tested with Pearson's correlation (two-tailed) and nonlinear regression tests. One-way analysis of variance (ANOVA) was used to compare the hydrological DOC fluxes among different hydrological processes. The seasonal difference of hydrological DOC fluxes, $\delta^{13}\text{C}_{\text{DOC}}$ was tested by independent sample t test. The SPSS 15.0 software was used for all calculations.

Because the individual correlations between the water flux and the DOC concentration in the throughfall, litter leachate, and soil water were significant (Fig. S1), the regression equations used for the water flux and DOC concentration ($Y = ae^{bx}$) were as follows:

$$C_{TF} = 48.69e^{-0.097x} \quad \text{adjusted } r^2 = 0.3883, p = 0.002 \quad (2)$$

$$C_{LL} = 60.93e^{-0.048x} \quad \text{adjusted } r^2 = 0.4131, p < 0.001 \quad (3)$$

$$C_{sw20} = 6.78e^{-0.02048x} \quad \text{adjusted } r^2 = 0.5840, p < 0.001 \quad (4)$$

where C_{TF} , C_{LL} , and C_{sw20} are the DOC concentrations (mg L^{-1}) in the throughfall, litter leachate, and soil water (0–20 cm), respectively, and x is the water flux per day (mm d^{-1}).

We did not collect all the individual rainfall events, throughfall, litter leachate, and soil water samples to analyze the DOC concentrations, but interpolated all the DOC concentrations and water fluxes according to eq(2)–(4).

The daily DOC flux was calculated as

$$F = CV/100 \quad (5)$$

where F is the daily DOC flux ($\text{kg C ha}^{-1} \text{d}^{-1}$), C is the DOC concentration (mg L^{-1}), and V is the water flux (mm d^{-1}) per day.

The weekly carbon flux ($\text{kg C ha}^{-1} \text{week}^{-1}$) was calculated as the sum of the daily DOC fluxes.

Soil temperature and soil water content of eddy flux tower explained 89.96% and 80.57% dynamic of that of soil respiration observation plot from Feb. 2008 to Jan. 2009 respectively, and the correlations between soil temperature at a depth of 5 cm and both SR and HR were strong (Fig. S2) between February 2008 and January 2009. SR and HR during the period from January 1, 2009 to December 31, 2011 were calculated based on the equation $Y = ae^{bx}$ from the data collected between February 2008 and January 2009, as follows:

$$SR = 46.37e^{(0.11T5)} \quad r^2 = 0.8966, p < 0.0001 \quad (6)$$

$$HR = 18.90e^{(0.14T5)} \quad r^2 = 0.8372, p < 0.0001 \quad (7)$$

where SR is total soil respiration ($\text{mg CO}_2 \text{ m}^{-2} \text{h}^{-1}$), HR is heterotrophic respiration ($\text{mg CO}_2 \text{ m}^{-2} \text{h}^{-1}$),

and T5 is soil temperature at 5 cm depth.

Sensitivity indices calculations

Firstly, weekly soil respirations fluxes, weekly average of soil temperature and soil water content, weekly water and DOC fluxes were standardized by the ratio of measured value to the mean value during the observation period. Secondly, linear regression equation was used between the standardized soil respirations values and T, SWC, water and DOC fluxes respectively. Thirdly, we considered the slope of the linear regression as the sensitivity indices which showed the soil respirations variation rate with soil temperature, soil water content, water and DOC fluxes changing.

3 Results

3.1 Water and DOC fluxes in a tropical rainforest

The seasonal and annual water fluxes decreased from the rainfall to the surface soil (Fig. 1a). The interception rate of the water between hydrological processes was higher in the dry season than in the rainy season (Fig. 1a, Table 1). The highest annual interception rate was between the litter leachate and the surface soil ($63.85 \pm 7.98\%$), which was $62.19 \pm 15.07\%$ in the rainy season and $81.64 \pm 23.38\%$ in the dry season.

The seasonal dynamics of the DOC flux were similar to those of the water flux (Fig. 1, Table 1). The annual DOC flux increased from rainfall ($41.9 \pm 3.8 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) to throughfall ($113.5 \pm 8.5 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) and to litter leachate ($127.7 \pm 8.5 \text{ kg C ha}^{-1} \text{ yr}^{-1}$), and then decreased sharply to the surface soil at 0–20 cm ($7.07 \pm 1.4 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) (Fig. 1b). The surface soil intercepted most of the DOC coming from the previous layer (annual: $94.4 \pm 1.2\%$, dry season: $96.7 \pm 4.4\%$, rainy season: $93.9 \pm 2.6\%$). That the interception rates for water and DOC were greatest in the surface soil indicates the surface soil

is the most important water and DOC sink in this tropical rainforest (Table 1).

3.2 Isotopic characteristics of DOC in the hydrological processes of a tropical rainforest

During the transfer of rainfall to soil water (0–20 cm), $\delta^{13}\text{C}_{\text{DOC}}$ was highest in the rainfall DOC and lowest in the throughfall DOC in both the rainy and dry seasons (Table 2). The seasonal difference in $\delta^{13}\text{C}_{\text{DOC}}$ was highest in the surface soil water (3.25‰) and lowest in the litter leachate (0.11‰). From the litter leachate to the surface soil water, $\delta^{13}\text{C}_{\text{DOC}}$ increased significantly by 4.26‰ ($p = 0.05$) in the rainy season, but increased by only 1.12‰ (not significant, $p = 0.39$) in the dry season. $\delta^{13}\text{C}$ increased from the canopy leaves to the soil and did not differ significantly between seasons (Table 2).

In both the dry and rainy seasons, $\delta^{13}\text{C}_{\text{DOC}}$ in water was higher than $\delta^{13}\text{C}$ in the corresponding element (comparing throughfall with leaves, litter leachate with litter, and soil water with soil at 20 cm depth) (Table 2). The smallest difference between $\delta^{13}\text{C}_{\text{DOC}}$ and $\delta^{13}\text{C}$ in each compartment occurred between soil water DOC and soil carbon in the dry season, which was only 0.23‰. The greater difference between $\delta^{13}\text{C}_{\text{DOC}}$ and $\delta^{13}\text{C}$ in the rainy season than in the dry season for soil water and soil (Table 2) indicates that the biogeochemical dynamics of DOC are more active in the rainy season than in the dry season in soil.

3.3 Surface soil CO_2 flux dynamics in a tropical rainforest

In the tropical rainforest at Xishuangbanna, SR was dominated by HR (Fig. 2). HR contributed more to SR during the rainy season ($76.8 \pm 0.8\%$) than during the dry season ($66.5 \pm 0.5\%$), and the annual contribution of HR to SR was $71.7 \pm 0.7\%$. SR and HR were higher in the rainy season than in the dry season, similar to the dynamics of the hydrological and DOC fluxes (Fig. 1).

Standardized soil temperature explained 98.7% and 98.2% of the variation in standardized SR and HR,

respectively, and standardized soil moisture explained 55.8% and 56.8% of the variation in standardized SR and HR, respectively (Table 3). The sensitivity indices of SR and HR for soil temperature at a depth of 10 cm were 0.56 and 0.46, respectively, whereas their sensitivity indices for soil moisture were 0.65 and 0.53, respectively, based on observational data (Table 3, Fig. S2).

3.4 Influence of DOC flux on soil CO₂ flux in a tropical rainforest

There were significant correlations between the standardized weekly SR and HR and the standardized weekly water fluxes and DOC fluxes through the hydrological processes (Table 3). Based on the definition of the temperature-dependent sensitivity index for soil respirations, which is the slope of standardized soil respirations caused by increase in standardized temperature, we also defined a soil-water-content-dependent sensitivity index, a DOC-flux-dependent sensitivity index, and a water-flux-dependent sensitivity index in this study, analogous to the temperature-dependent sensitivity index for SR (Table 3). An independent *t* test showed that the DOC-flux-dependent sensitivity indices for SR (2.72 ± 0.51) and HR (2.21 ± 0.42) were significantly lower than the water-flux-dependent sensitivity indices for SR (2.87 ± 0.52 , $t = -2.68$, $p = 0.06$) and HR (2.33 ± 0.43 , $t = -2.57$, $p = 0.06$), respectively, which indicates that SR and HR were more sensitive to the water flux than to the DOC flux through the hydrological processes. The significant difference was observed between the water-flux-dependent indices ($t = 13.78$, $p < 0.001$) for SR (2.87 ± 0.52) and HR (2.33 ± 0.43), or between the DOC-flux-dependent indices ($t = 13.12$, $p < 0.001$) for SR (2.72 ± 0.51) and HR (2.21 ± 0.42).

The soil-water-content-dependent sensitivity indices for HR (0.53) and SR (0.65) were than the soil-temperature-dependent sensitivity indices (HR, 0.46; SR, 0.56), but less than all the

water-flux-dependent and DOC-flux-dependent sensitivity indices for SR and HR (Table 3). This indicates that SR and HR are more sensitive to the hydrological water flux and DOC flux than to the soil water content and soil temperature. A comparison of the sensitivity indices for water flux, DOC flux, soil temperature, and soil moisture in all the hydrological processes reveals that SR and HR were most sensitive to the water flux (3.70) dynamics which is a little higher than DOC flux (3.57) in the soil water (0–20 cm depth) when weekly variations in the Xishuangbanna tropical rainforest were considered.

4 Discussion

Our results showed that the throughfall carried most of the DOC ($113.5 \pm 8.5 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) through the hydrological processes in the Xishuangbanna tropical rainforest, which amounted to 6.81% of the NEE ($1.67 \times 10^3 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) (Tan *et al.*, 2010) in this tropical rainforest in southwest China. The litter leachate DOC ($127.7 \pm 8.5 \text{ kg}$) accounted for 7.23% of the NEE in this forest. This result indicates that the throughfall DOC is an important component of the tropical rainforest carbon budget. The litter leachate fed a great deal of DOC to the soil, but the surface soil intercepted $94.4 \pm 1.2\%$ ($127.7 \pm 8.0 \text{ kg}$) of the DOC, and the surface soil water DOC flux was only $7.1 \pm 1.4 \text{ kg C ha}^{-1} \text{ yr}^{-1}$, which was slightly less than that at the headwater stream outlet ($10.31 \text{ kg C ha}^{-1} \text{ yr}^{-1}$) (Zhou *et al.*, 2013). The surface soil intercepted the bulk of the litter leachate DOC and transported little DOC to the deep layer, indicating that the surface soil is the DOC sink in the tropical rainforest in Xishuangbanna. The small seasonal differences in $\delta^{13}\text{C}_{\text{DOC}}$ in the rainfall, throughfall, and litter leachate indicate that the DOC in the aboveground water is seasonally stable (Table 2). However, $\delta^{13}\text{C}_{\text{DOC}}$ in the soil water (at 0–20 cm) was higher in the rainy season (3.25‰) than in the dry season, indicating that the DOC

reaction in the surface soil is seasonal. In the dry season, $\delta^{13}\text{C}_{\text{DOC}}$ in the surface soil water ($-27.1 \pm 2.2\text{‰}$) was similar to $\delta^{13}\text{C}_{\text{DOC}}$ in the soil ($-27.3 \pm 0.1\text{‰}$), indicating that the soil is the major source of soil water DOC. This is attributable to the combined absorption effects of the high clay content (Fr berg, 2004, Lemma *et al.*, 2007, Sanderman and Amundson, 2008, Tang *et al.*, 2007) and the lack of water carrying DOC through the different compartments in the dry season. Therefore, most DOC is locally produced rather than transported. Less water and the lower DOC input from litter leachate and throughfall to the surface soil (Fig. 1) also contribute to a reduction in microbial activity, which contributes negligibly to the soil DOC when the soil moisture and soil temperature are low in the dry season (Wu *et al.*, 2009). In the rainy season, the soil water content and soil temperature are higher, so there is more vigorous biogeochemical activity in the surface soil (Bengtson and Bengtsson, 2007). Therefore, more DOC is released from the soil to be mineralized by microorganisms, and there is more ^{13}C in the soil water DOC ($\delta^{13}\text{C} = -23.9 \pm 2.2\text{‰}$) than in the soil ($\delta^{13}\text{C}$, $-27.3 \pm 0.1\text{‰}$). The relatively low $\delta^{13}\text{C}_{\text{DOC}}$ in the litter leachate ($\delta^{13}\text{C}_{\text{DOC}} = -28.1 \pm 2.7\text{‰}$) compared with the soil water indicates that the DOC from the litter leachate has attended in the carbon cycle in the surface soil (Cleveland *et al.*, 2006, De Troyer *et al.*, 2011). Furthermore, most of the DOC from the throughfall, litter leachate, and litter was fed to the surface soil, and the soil water $\delta^{13}\text{C}_{\text{DOC}}$ value was higher than that of the throughfall, litter leachate DOC, and $\delta^{13}\text{C}$ soil (0–20 cm) values (Table 2). These data indicate that all the DOC transported by the throughfall and litter leachate was ultimately involved in the surface soil carbon cycle (Fr berg *et al.*, 2003, 2005, Kammer *et al.*, 2012), and has also contributed to the SR because it is an important part of the surface soil carbon cycle in the tropical rainforest at Xishuangbanna. Laboratory-based studies of tropical forests have shown that DOC primes the soil CO_2 flux (Qiao *et al.*,

2013). A study of a temperate forest showed that the rate of DOC production is one of the rate-limiting steps for SR (Bengtson and Bengtsson, 2007). Comparative studies of ^{13}C and ^{14}C in DOC and SOC have also shown that fresh organic carbon stimulates the activity of old carbon, and increases the emission of CO_2 because DOC is the substrate of microbial activity (Cleveland *et al.*, 2004, 2006, Hagedorn and Machwitz, 2007, Hagedorn *et al.*, 2004, Qiao *et al.*, 2013). Because the microbial biomass and potential carbon mineralization rates are higher in soils with higher DOC contents than in soils with lower DOC contents (Montaño *et al.*, 2007), the DOC turnover rate (Bengtson and Bengtsson, 2007) is rapid and the transformation period is short (3–14 days) (Cleveland *et al.*, 2006, De Troyer *et al.*, 2011). This indicates that DOC is involved in the surface soil carbon cycle in the short term by affecting SR (Cleveland *et al.*, 2004, 2006). Although we did not determine the period of the DOC turnover cycle, the weekly DOC flux passing through the hydrological processes (throughfall, litter leachate, soil water, and interception by the surface soil) significantly explained SR and HR, with higher sensitivity indices than the indices for the soil water content and soil temperature (Table 3), predicting that DOC has a significant impact on soil CO_2 emissions in this tropical rainforest.

It is important to consider which part of the DOC flux in the hydrological processes of this tropical rainforest most strongly influences SR. Previous studies have shown that of all the factors affecting SR, it is most sensitive to soil temperature (Bekku *et al.*, 2003, Reichstein *et al.*, 2003, Zheng *et al.*, 2009), as in the tropical forest at Xishuangbanna (Sha *et al.*, 2005). Although soil temperature better explained SR and HR than the DOC flux, the sensitivity indices for the soil water DOC fluxes were higher than the sensitivity indices for soil temperature, although temperature explained the rate of SR better than the DOC flux (Table 3). At this study site, HR, which depends predominantly on microbial activity and substrates, contributed the major fraction of SR (Fig. S2), so not only HR, but also SR depends most

strongly on the microbial and respiratory substrates in this tropical rainforest. Therefore, the DOC transported by the forest hydrological processes, from litter decomposition, root exudates, and the soil itself, will contribute to SR (Table 2). The bioavailability of the DOC transported by hydrological processes is greater than that of SOC (De Troyer *et al.*, 2011, Kindler *et al.*, 2011). The DOC from throughfall and litter leachate is also an important contributor because $\delta^{13}\text{C}_{\text{DOC}}$ differs between the surface soil water and the litter leachate and throughfall (Table 2). Although ectotrophic mycorrhizae contribute significantly to SR in the rhizospheres of some temperate and boreal forests (Neumann *et al.*, 2014; Tomè *et al.*, 2016), in this tropical rainforest, EMF: *Paraglomus*, a kind of endomycorrhiza, occupies more than 90% of the mycorrhizal community (Shi, 2014). Together with roots, and root exudate, it contributes to the autotrophic SR, which is only 28.9% of the total SR, so the mycorrhiza is not the dominant contributor to SR in this tropical rainforest. The other details of the biogeochemical processes affecting DOC in the surface soil are not obvious in this study. However, according to both laboratory and field studies, the DOC intercepted by the surface soil clearly affects HR (Table 3), together with the DOC from litter decomposition and the soil itself (Cleveland *et al.*, 2004, 2006, Hagedorn and Machwitz, 2007, Hagedorn *et al.*, 2004, Jandl and Sollins, 1997, Keiluweit *et al.*, 2015; Montañó *et al.*, 2007, Qiao *et al.*, 2013, Schwendenmann and Veldkamp, 2005;). Considering the effect of DOC on SR, the surface soil water DOC is the most sensitive index of HR and SR (Table 3). The DOC-flux-dependent sensitivity indices for the different parts of the hydrological processes in this tropical rainforest were a little less but insignificant than the amount-of-water-dependent sensitivity indices, which shows that the DOC flux affects SR less than the amount of water passing through the system, because of the combined effects of water and DOC on SR. According the DOC significant contribution of soil respirations (Cleveland *et al.*, 2004, 2006, Hagedorn and Machwitz, 2007,

Hagedorn *et al.*, 2004, Qiao *et al.*, 2013), the little difference mechanisms between DOC and water flux of tropical rainforest should be declared in the future study.

This study demonstrates that the surface soil is a sink for the DOC transported by hydrological processes (Fig. 1), and that HR and SR are sensitive to the DOC flux through these processes. The most sensitive indicator of SR is the soil water flux (at 0–20 cm) and followed by soil water DOC flux, both exceeding the sensitivity of the soil temperature, soil water content, and other water flux, and DOC flux along all the hydrological processes (Table 3). The variations in $\delta^{13}\text{C}$ in DOC, soil, and plants also partly support the notion that the soil water DOC flux is the more sensitive index of SR in this tropical rainforest. The results suggest that the DOC transported by hydrological processes plays the more important role in the SR processes. In the context of global climate change, more attention must be paid to the contribution of hydrologically transported DOC in future studies of the mechanisms of SR.

Author contribution

W.-J. Zhou and D. Schaefer, H.-Z. Lu, Sha L-Q, Y.-P Zhang designed the experiments and W.-J. Zhou, H.-Z. Lu, Q.-H. Song, Y. Deng, X.-B. Deng carried them out. W-J Zhou prepared the manuscript with contributions from all co-authors.

Acknowledgments

This work was supported by the Natural Science Foundation of Yunnan Province (2016FB073, 2015FB188), the National Natural Science Foundation of China (41271056, U1202234, 40801035), the Strategic Priority Research Program of the Chinese Academy of Sciences (No. XDA05020302 and XDA05050601), the CAS 135 Project (XTBG-F01), and the Science and Technology Service Network

Initiative of the Chinese Academy of Sciences (No. KFJ-EW-ST5-084).

We thank the staff and technicians of the Xishuangbanna Station for Tropical Rain Forest Ecosystems who assisted with field measurements and the Public Technology Service Center of Xishuangbanna Tropical Botanical Garden, CAS, who contributed to ¹³C analyses. We also thank Zhi-Gang Chen and Zhi-Hua Zhou for assistance with sampling, Zhi-Ling Chen and Li-Fang OU for laboratory work, and Jan Mulder and Jing Zhu for reviewing the manuscript.

References

- Bekku Y. S., Nakatsubo T., Kume, A., Adachi M., Koizumi, H.: Effect of warming on the temperature dependence of soil, respiration rate in arctic, temperate and tropical soils, *Appl. Soil. ecol.*, 22, 205–210, (2003)
- Bengtson P., Bengtsson G. R.: Rapid turnover of DOC in temperate forests accounts for increased CO₂ production at elevated temperatures, *Eco.Lett*, 10, 783–790, 2007.
- Bianchi T. S.: The role of terrestrially derived organic carbon in the coastal ocean: A changing paradigm and the priming effect, *P. Natl. Acad. Sci. USA.*, 108, 19473–19481, 2011.
- Blagodatskaya E., Yuyukina T., Blagodatsky S., Kuzyakov Y.: Three-source-partitioning of microbial biomass and of CO₂ efflux from soil to evaluate mechanisms of priming effects, *Soil Biol. Biochem.*, 43, 778–786, 2011.
- Cao M., Zhang J., Feng Z., Deng J., Deng X.: Tree species composition of a seasonal rain forest in Xishuangbanna, Southwest China, *Trop. Ecol.*, **37**, 183–192, 1996.
- Chantigny M.: Dissolved and water-extractable organic matter in soils: a review on the influence of land use and management practices, *Geoderma*, 113, 357–380, 2003.

424 Chuyong G. B., Newbery D. M., Songwe N. C.: Rainfall input, throughfall and stemflow of nutrients in
 425 a central African rain forest dominated by ectomycorrhizal trees, *Biogeochemistry*, **67**, 73–91,
 426 2004.

427 Cleveland C. C., Neff J. C., Townsend A. R., Hood E.: Composition, dynamics, and fate of leached
 428 dissolved organic matter in terrestrial ecosystems: results from a decomposition experiment,
 429 *Ecosystems*, **7**: 175–285, 2004.

430 Cleveland C. C., Nemergut D. R., Schmidt S. K., Townsend A. R.: Increases in soil respiration
 431 following labile carbon additions linked to rapid shifts in soil microbial community composition,
 432 *Biogeochemistry*, **82**, 229–240, 2006.

433 Comstedt D., Boström B., Marshall J.D., Holm A., Slaney M., Linder S., Ekblad A.: Effects of elevated
 434 atmospheric carbon dioxide and temperature on soil respiration in a boreal forest using $\delta^{13}\text{C}$ as a
 435 labeling tool, *Ecosystems*, **9**, 1266–1277, 2007.

436 Davidson E.A., Samanta, S., Caramori, S.S., Savage, K.: The Dual Arrhenius and Michaelis-Menten
 437 kinetics model for decomposition of soil organic matter at hourly to seasonal time scales, *Global*
 438 *Change Biol.*, **18**, 371–384, 2012.

439 De Troyer, I., Amery F., Van Moorleghe C., Smolders E., Merckx R.: Tracing the source and fate of
 440 dissolved organic matter in soil after incorporation of a ^{13}C labelled residue: A batch incubation
 441 study, *Soil Biol. Biochem.*, **43**, 513–519, 2011.

442 Dezzeo N., Chacón N.: Nutrient fluxes in incident rainfall, throughfall, and stemflow in adjacent
 443 primary and secondary forests of the Gran Sabana, southern Venezuela, *Forest Eco. Manag.*, **234**,
 444 218–226, 2006.

445 Fang H.-J., Yu G.-R., Cheng S.-L., Mo J.-M., Yan J.-H., Li S.: ^{13}C abundance, water-soluble and

446 microbial biomass carbon as potential indicators of soil organic carbon dynamics in subtropical
 447 forests at different successional stages and subject to different nitrogen loads, *Plant Soil*, 320,
 448 243–254, 2009.

449 Fang Q.L., Sha L.: Fine roots turnover of tropical seasonal rain forest in Xishuangbanna, Yunnan, SW
 450 China, *Journal of Mountain Science*, 23 (4):488-494, 2005.

451 Fontaine S., Barot S., Barré P., Bdioui N., Mary B., Rumpel C.: Stability of organic carbon in deep soil
 452 layers controlled by fresh carbon supply, *Nature*, 450, 277–280, 2007.

453 Fröberg M.: Processes controlling production and transport of dissolved organic carbon in forest soils,
 454 Doctoral thesis, 2004.

455 Fröberg, M., Berggren, D., Bergkvist, B., Bryant, C., Knicker, H.: Contributions of O_i, O_e and O_a
 456 horizons to dissolved organic matter in forest floor leachates, *Geoderma*, 113, 311–322, 2003.

457 Fröberg, M., Kleja, D.B., Bergkvist, B., Tipping, E., Mulder, J.: Dissolved organic carbon leaching
 458 from a coniferous forest floor – a field manipulation experiment, *Biogeochemistry*, 75, 271–287,
 459 2005.

460 Guenet, B., Neill, C., Bardoux, G., Abbadie, L.: Is there a linear relationship between priming effect
 461 intensity and the amount of organic matter input? *Appl. Soil Ecol.*, 46, 436–442, 2010.

462 Hagedorn, F., Machwitz, M., 2007. Controls on dissolved organic matter leaching from forest litter
 463 grown under elevated atmospheric CO₂, *Soil Bio. Bioch.*, 39, 1759–1769, 2007.

464 Hagedorn, F., Saurer, M., Blaser, P.: A ¹³C tracer study to identify the origin of dissolved organic
 465 carbon in forested mineral soils, *Eur. J. Soil Sci.*, 55, 91–100, 2004.

466 Jandl, R., Sollins, P.: Water extractable soil carbon in relation to the belowground carbon cycle, *Bio.*
 467 *Fer. Soils*, 25, 196–201, 1997.

468 Kalbitz, K., Meyer, A., Yang, R., Gerstberger, P.: Response of dissolved organic matter in the forest
 469 floor to long-term manipulation of litter and throughfall inputs, *Biogeochemistry*, 86, 301–318,
 470 2007.

471 Kalbitz, K., Solinger, S., Park, J.-H., Michalzik, B., Matzner, E.: Controls on the dynamics of dissolved
 472 organic matter in soils: a review, *Soil Sci.*, 165, 277–304, 2000.

473 Kammer, A., Schmidt, M.W.I., Hagedorn, F.: Decomposition pathways of ¹³C-depleted leaf litter in
 474 forest soils of the Swiss Jura, *Biogeochemistry*, 108, 395–411, 2012.

475 Keiluweit, M., Bougoure, J. J., Nico, P. S., Pett-Ridge, J., Weber, P. K., Kleber, M.: Mineral protection
 476 of soil carbon counteracted by root exudates, *Nature Climate Change*, 5(6), 588–595, 2015.

477 Kindler, R., Siemens, J.A.N., Kaiser, K., Walmsley, D.C., Bernhofer, C., Buchmann, N., Cellier, P.,
 478 Eugster, W., Gleixner, G., GrÜnwald, T., Heim, A., Ibrom, A., Jones, S.K., Jones, M., Klumpp, K.,
 479 Kutsch, W., Larsen, K.S., Lehuger, S., Loubet, B., McKenzie, R., Moors, E., Osborne, B.,
 480 Pilegaard, K.I.M., Rebmann, C., Saunders, M., Schmidt, M.W.I., Schrumpf, M., Seyfferth, J.,
 481 Skiba, U.T.E., Soussana, J.-F., Sutton, M.A., Tefs, C., Vowinckel, B., Zeeman, M.J.,
 482 Kaupenjohann, M.: Dissolved carbon leaching from soil is a crucial component of the net
 483 ecosystem carbon balance, *Global Change Bio.* 17, 1167–1185, 2011.

484 Lemma, B., Nilsson, I., Kleja, D.B., Olsson, M., Knicker, H.: Decomposition and substrate quality of
 485 leaf litters and fine roots from three exotic plantations and a native forest in the southwestern
 486 highlands of Ethiopia, *Soil Bio. Bioch.*, 39, 2317–2328, 2007.

487 Liu, C.P., Sheu, B.H., 2003. Dissolved organic carbon in precipitation, throughfall, stemflow, soil
 488 solution, and stream water at the Guandaushi subtropical forest in Taiwan, *Forest Eco. Manag.*,
 489 172, 315–325, 2003.

490 Liu, W.J., Liu, W.Y., Li, J.T., Wu, Z.W., Li, H.M.: 2008. Isotope variations of throughfall, stemflow and
 491 soil water in a tropical rain forest and a rubber plantation in Xishuangbanna, SW China, *Hydrol.*
 492 *Res.*, 39, 437–449, 2008.

493 McClain, M.E., Richey, J.E., Brandes, J.A., Pimentel, T.P.: Dissolved organic matter and
 494 terrestrial-lotic linkages in the central Amazon basin of Brazil, *Global Biogeochem. Cy.*, 11,
 495 295–311, 1997.

496 McJannet, D., Wallace, J., Reddell, P.: Precipitation interception in Australian tropical rainforests: I.
 497 Measurement of stemflow, throughfall and cloud interception, *Hydrol. Process.*, 21, 1692–1702,
 498 2007.

499 Montañó, N.M., García-Oliva, F., Jaramillo, V.J.: Dissolved organic carbon affects soil microbial
 500 activity and nitrogen dynamics in a Mexican tropical deciduous forest, *Plant Soil*, 295, 265–277,
 501 2007.

502 Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, H.A., Forsius, M., Høgåsen, T., Wilander, A.,
 503 Skjelkvåle, B.L., Jeffries, D.S., Vuorenmaa, J., Keller, B., Kopáček, J., Vesely, J.: Dissolved
 504 organic carbon trends resulting from changes in atmospheric deposition chemistry, *Nature*, 450,
 505 537–540. 2007.

506 Neumann J., Matzner E.: Contribution of newly grown extramatrical ectomycorrhizal mycelium and
 507 fine roots to soil respiration in a young Norway spruce site, *Plant soil*, 378(1–2): 73–82, 2014.

508 Park, J.H., Kalbitz, K., Matzner, E.: Resource control on the production of dissolved organic carbon
 509 and nitrogen in a deciduous forest floor, *Soil Bio. Bioche.*, 34, 1391–1391, 2002..

510 Qiao, N., Schaefer, D., Blagodatskaya, E., Zou, X., Xu, X., Kuzyakov, Y.: Labile carbon retention
 511 compensates for CO₂ released by priming in forest soils, *Global Change Biol.*, 20(6): 1943–1954,

2014..

Reichstein, M., Rey, A., Freibauer, A., Tenhunen, J., Valentini, R., Banza, J., Casals, P., Cheng, Y.F., Grunzweig, J.M., Irvine, J., Joffre, R., Law, B.E., Loustau, D., Miglietta, F., Oechel, W., Ourcival, J.M., Pereira, J.S., Peressotti, A., Ponti, F., Qi, Y., Rambal, S., Rayment, M., Romanya, J., Rossi, F., Tedeschi, V., Tirone, G., Xu, M., Yakir, D.: Modeling temporal and large-scale spatial variability of soil respiration from soil water availability, temperature and vegetation productivity indices, *Global Biogeochem. Cy.*, 17, DOI: 10.1029/2003GB002035, 2003.

Sanderman, J., Amundson, R.: A comparative study of dissolved organic carbon transport and stabilization in California forest and grassland soils, *Biogeochemistry*, 89, 309–327, 2008.

Schrumpf, M., Zech, W., Lehmann, J., Lyaruu, H.V.C.: TOC, TON, TOS and TOP in rainfall, throughfall, litter percolate and soil solution of a montane rainforest succession at Mt. Kilimanjaro, Tanzania, *Biogeochemistry*, 78, 361–387, 2006.

Schwendenmann, L., Veldkamp, E.: The role of dissolved organic carbon, dissolved organic nitrogen, and dissolved inorganic nitrogen in a tropical wet forest ecosystem, *Ecosystems*, 8, 339–351, 2005..

Sha, L.Q., Zheng, Z., Tang, J.W., Wang, Y.H., Zhang, Y.P., Cao, M., Wang, R., Liu, G.G., Wang, Y.S., Sun, Y.: Soil respiration in tropical seasonal rain forest in Xishuangbanna, SW China, *Science in China Series D-Earth Sciences*, 48, 189–197, 2005(In Chinese).

Shi L.L.: Soil Microbial Community in Forest Ecosystem as Revealed by Molecular Techniques —diversity pattern, maintenance mechanism, and the response to disturbance, A Dissertation Submitted to University of Chinese Academy of Sciences, In partial fulfillment of the requirement Doctor of Philosophy, 2014 (In Chinese).

534 Sowerby, A., Emmett, B.A., Williams, D., Beier, C., Evans, C.D.: The response of dissolved organic
535 carbon (DOC) and the ecosystem carbon balance to experimental drought in a temperate
536 shrubland, *Eur. J. Soil Sci.*, 61, 697–709, 2010.

537 Stephan, S., Karsten, K., Egbert, M.: Controls on the dynamics of dissolved organic carbon and
538 nitrogen in a Central European deciduous forest, *Biogeochemistry*, 55, 327–349, 2001.

539 Tan, Z., Zhang, Y., Yu, G., Sha, L., Tang, J., Deng, X., Song, Q.: Carbon balance of a primary tropical
540 seasonal rain forest, *J. Geophys. Res.*, 115 (D4), DOI: 10.1029/2009JD012913, 2010.

541 Tang, J.W., Cao, M., Zhang, J.H., Li, M.H., Litterfall production, decomposition and nutrient use
542 efficiency varies with tropical forest types in Xishuangbanna, SW China: a 10-year study, *Plant*
543 *soil*, 335, 271–288, 2010.

544 Tang, Y.L., Deng, X.B., Li, Y.W., Zhang, S.B.: Research on the difference of soil fertility in the
545 different forest types in Xishuangbanna, *Journal of Anhui Agricultural Sciences*, 35,
546 779–781, 2007 (In Chinese).

547 Tomè E., Ventura, M., Folegot, S., Zanutelli, D., Montagnani, L., Mimmo, T., Tonon, G., Tagliavini, M.,
548 Scandellari, F.: Mycorrhizal contribution to soil respiration in an apple orchard. *Appl. Soil*
549 *ecol.*, 101, 165–173, 2016.

550 Wu, Y., Yang, X., Yu, G.: Seasonal fluctuation of soil microbial biomass carbon and its influence factors
551 in two types of tropical rainforests, *Ecology and Environmental Sciences*, 18, 658–663, 2009 (In
552 Chinese).

553 Yakov, K.: Priming effects: Interactions between living and dead organic matter. *Soil Bio. Biochem.* 42,
554 1363–1371, 2010.

555 Zhang, Y., Tan, Z., Song, Q., Yu, G., Sun, X.: Respiration controls the unexpected seasonal pattern of

556 carbon flux in an Asian tropical rain forest, *Atmos. Environ.* 44, 3886–3893, 2010.

557 Zheng, Z.M., Yu, G.R., Fu, Y.L., Wang, Y.S., Sun, X.M., Wang, Y.H.: Temperature sensitivity of soil

558 respiration is affected by prevailing climatic conditions and soil organic carbon content: A

559 trans-China based case study, *Soil Bio. Biochem.*, 41, 1531–1540, 2009.

560 Zimmermann, A., Wilcke, W., Elsenbeer, H.: Spatial and temporal patterns of throughfall quantity and

561 quality in a tropical montane forest in Ecuador, *J. Hydrol.* 343, 80–96, 2007.

562

Table legends

Table 1 The interception rate of the water between hydrological processes in the tropical rainforest at Xishuangbanna southwest China

Table 2 DOC $\delta^{13}\text{C}$ dynamics along the hydrological processes (R, rainfall, TF, throughfall, LL, litter leachate) and the $\delta^{13}\text{C}$ in leaves, litter, and surface soil in the tropical rainforest at Xishuangbanna, southwest China

Table 3 Results of a regression analysis of the weekly water flux, DOC flux, soil respiration (SR), and heterotrophic respiration (HR) along the hydrological processes (TF, throughfall, LL, litter leachate) in the tropical rainforest in Xishuangbanna, southwest China.

Figure captions

Figure 1 Amount of water (A) and DOC flux along the hydrological processes in the tropical rainforest at Xishuangbanna, southwest China.

Figure 2 Dynamics of soil respiration (SR) and heterotrophic respiration (HR) (a) and soil temperature at 5cm and soil water content at 10cm (b) in the tropical rainforest at Xishuangbanna, southwest China. The shaded area indicates the rainy season.

579

580 Table 1

Interception (%)		Annual	Rainy season (%)	Dry season
Water flux	Between TF and P	53.9±11.7	43.1±2.7	41.3±14.8
	Between LL and TF	33.9±6.6	33.9±9.8	34.1±27.6
	Between SW20cm and LL	63.8±8.0	62.2±15.1	81.6±23.3
DOC flux	Between TF and R	137.0±19.9	182.0±16.0	170.8±7.8
	Between LL and TF	1.1±17.0	16.1±9.4	12.7±4.3
	Between SW20cm and LL	-96.7±4.4	-93.9±2.6	-94.4±1.2

581 P indicates rainfall, TF indicates throughfall, LL indicates litter leachate, SW20 indicates soil water at a
582 depth of 20 cm.

583 Table2

584

Season	P	TF	LL	Soil water (0–20 cm) ‰	Leaves	Litter	Soil (0–20 cm)
Rainy season	-23.9 ± 3.3^a	-28.7 ± 1.7^{bc}	-28.1 ± 2.7^{bc}	$-23.9 \pm 1.6^a *$	-32.4 ± 0.6^d	-30.4 ± 0.2^{cd}	-27.3 ± 0.1^b
Dry season	-23.8 ± 1.3^a	-29.1 ± 1.6^{bc}	-28.1 ± 1.5^{bc}	-27.1 ± 2.2^b	-32.5 ± 0.5^d	-30.2 ± 0.1^{cd}	-27.3 ± 0.1^{bc}

585 P indicates rainfall, TF indicates throughfall, LL indicates litter leachate, SW20 indicates soil water at a
586 depth of 20 cm.

587 Different superior letters indicate significant differences between the treatments according to Lsd test
588 ($P < 0.05$).

589 *indicates the significant seasonal difference according to independent sample t test ($p < 0.1$)

Table 3

		SR				HR			
		a	b	R^2	p	a	b	R^2	p
DOC flux	T	0.56	0.54	0.987	<0.001	0.46	0.64	0.982	<0.001
	SWC	0.65	0.41	0.558	<0.001	0.53	0.52	0.568	<0.001
	P	2.31	-1.17	0.423	<0.001	1.86	-0.74	0.425	<0.001
	TF	2.36	-1.25	0.429	<0.001	1.91	-0.83	0.413	<0.001
	LL	2.71	-1.57	0.355	<0.001	2.21	-1.10	0.366	<0.001
	SW20	3.57	-2.23	0.227	<0.001	2.91	-1.62	0.240	<0.001
Water flux	LL-SW20	2.66	-1.53	0.352	<0.001	2.17	-1.07	0.363	<0.001
	P	2.42	-1.35	0.323	<0.001	1.96	-0.92	0.331	<0.001
	TF	2.55	-1.44	0.316	<0.001	2.06	-0.99	0.323	<0.001
	LL	3.02	-1.83	0.301	<0.001	2.46	-1.31	0.312	<0.001
	SW20	3.70	-2.34	0.166	<0.001	3.02	-1.71	0.178	<0.001
	LL-SW20	2.64	-1.54	0.257	<0.001	2.14	-1.08	0.267	<0.001

Equations used to calculate the sensitivity indices: sensitivity index = a, where b is the constant of the regression equation for standardized soil respirations, and soil temperature, soil water content, water flux, and DOC flux: $Y = aX + b$, where Y is the standardized soil respiration rate, and X is standardized soil temperature, soil water content, water flux, or DOC flux.

T indicates soil temperature at a depth of 5 cm, SWC indicates soil water content, TF indicates throughfall, LL indicates litter leachate, SW20 indicates soil water at a depth of 20 cm, LL- SW20 indicates the difference between litter leachate and soil water at a depth of 20 cm.



