Abiotic versus biotic controls on soil nitrogen cycling in drylands along a 3200 km transect

Dongwei Liu^{1*}, Weixing Zhu^{1, 2}, Xiaobo Wang^{1*}, Yuepeng Pan³, Chao Wang¹, Dan Xi^{1, 5}, Edith Bai¹, Yuesi Wang³, Xingguo Han¹, Yunting Fang^{1, 4}

5

¹ Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese
 Academy of Sciences, Shenyang, 110016, China

- ² Department of Biological Sciences, Binghamton University, State University of New York,
 Binghamton, NY 13902
- ³ State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry

(LAPC), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China
 ⁴ Qingyuan Forest CERN, Chinese Academy of Sciences, Shenyang 110016, China

- Gingydan Forest CERRY, Chinese Academy of Sciences, Shenyang 110010, China
 ⁵ College of Resources and Environment, University of Chinese Academy of Sciences, Beijing,
 100040, China
- 14 100049, China 15
- *These authors contributed equally to this work.
- 17

18 Corresponding Author:

- 19 Weixing Zhu
- 20 Department of Biological Sciences, Binghamton University State University of New York,
- 21 Binghamton, NY 13902-6000
- 22 Phone: (607) 777-3218
- **23** Fax: (607)-777-6521
- 24 Email: wxzhu@binghamton.edu
- 25
- 26 Yunting Fang
- 27 Institute of Applied Ecology, the Chinese Academy of Science, No.72, Wenhua Road, Shenyang, P.
- 28 R. China, 110016
- 29 Phone: +86-24-83970541
- 30 Fax: +86-24-83970300
- 31 Email: fangyt@iae.ac.cn
- 32

33	Supplementary Information
34	
35	Supplementary Figure legends:
36	
37	Figure S1 Climate variables. The positive correlation (b) between aridity index (AI) and precipitation
38	demonstrated that the precipitation gradient here can also be considered as an AI gradient. In each site,
39	n=1.
40	
41	Figure S2 Soil moisture along the precipitation gradient. Soil moisture was expressed as a percent of dry
42	soil on a mass basis. In each site, n=5.
43	
44	
45	
46	



48 Figure S1



