

1 **Multi-gas and multi-source comparisons of six land use emission datasets**  
2 **and AFOLU estimates in the Fifth Assessment Report, for the tropics for**  
3 **2000-2005.**

4  
5 **Short title:** AFOLU dataset comparisons  
6

7 **Authors affiliation:**

8 Rosa Maria Roman-Cuesta<sup>1,2\*</sup>, Martin Herold<sup>2</sup>, Mariana C. Rufino<sup>1</sup>, Todd S. Rosenstock<sup>3</sup>,  
9 Richard A. Houghton<sup>4</sup>, Simone Rossi<sup>5</sup>, Klaus Butterbach-Bahl<sup>6,7</sup>, Stephen Ogle<sup>8</sup>, Benjamin  
10 Poulter<sup>9</sup>, Louis Verchot<sup>10,11</sup>, Christopher Martius<sup>1</sup>, Sytze de Bruin<sup>2</sup>.

11  
12 <sup>1</sup> Center for International Forestry Research (CIFOR), P.O Box 0113 BOCBD, Bogor 16000,  
13 Indonesia.

14 <sup>2</sup> Laboratory of Geo-Information Science and Remote Sensing - Wageningen University &  
15 Research. Droevendaalsesteeg 3, 6708PB. Wageningen. The Netherlands.

16 <sup>3</sup> World Agroforestry Centre (ICRAF). PO Box 30677-00100, Nairobi. Kenya.

17 <sup>4</sup> Woods Hole Research Center. 149 Woods Hole Road Falmouth, MA, 02540-1644, US.

18 <sup>5</sup> Global Environmental Monitoring Unit, Institute for Environment and Sustainability,  
19 European Commission, Joint Research Centre, TP. 440 21020 Ispra, Varese 21027, Italy,

20 <sup>6</sup> International Livestock Research Institute (ILRI) P.O. Box 30709. Nairobi 00100, Kenya

21 <sup>7</sup> Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research (IMK-  
22 IFU), Garmisch-Partenkirchen, Germany

23 <sup>8</sup> Natural Resource Ecology Laboratory, Campus Delivery 1499, Colorado State University,  
24 Fort Collins, Colorado 80523-1499, USA.

25 <sup>9</sup> Ecosystem Dynamics Laboratory. Montana State University. P.O. Box 172000. Bozeman,  
26 MT 59717-2000. USA.

27 <sup>10</sup> International Center for Tropical Agriculture, Km17 Recta Cali-Palmira, Apartado Aéreo  
28 6713, Cali, Colombia.

29 <sup>11</sup> Earth Institute Center for Environmental Sustainability, Columbia University, New York,  
30 USA.

31 \* Corresponding author. Telephone:+31317485919, Fax: Email: rosa.roman@wur.nl

32

33 **Keywords:** AFOLU, Land use greenhouse gas emissions, Land Use Land Cover Change and  
34 Forestry, LULUCF, mitigation, Fifth Assessment Report, gross emissions flux.

35

## 36 **ABSTRACT**

37 The Agriculture, Forestry and Other Land Use (AFOLU) sector contributes with ca. 20-25%  
38 of global anthropogenic emissions (2010), making it a key component of any climate change  
39 mitigation strategy. AFOLU estimates remain, however, highly uncertain, jeopardizing the  
40 mitigation effectiveness of this sector. Comparisons of global AFOLU emissions have shown  
41 divergences of up to 25%, urging for improved understanding on the reasons behind these  
42 differences. Here we compare a diversity of AFOLU emission datasets (i.e. FAOSTAT,  
43 EDGAR, the newly developed AFOLU “Hotspots”, “Houghton”, “Baccini”, and EPA) and  
44 estimates given in the Fifth Assessment Report, for the tropics (2000-2005), to identify  
45 plausible explanations for the differences in: i) aggregated gross AFOLU emissions, and ii)

46 disaggregated emissions by sources, and by gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O). We also aim to iii)  
47 identify countries with low agreement among AFOLU datasets, to navigate research efforts.  
48 Aggregated gross emissions were similar for all databases for the AFOLU: 8.2 (5.5-12.2), 8.4  
49 and 8.0 Pg CO<sub>2</sub>e.yr<sup>-1</sup> (Hotspots, FAOSTAT and EDGAR respectively), Forests: 6.0 (3.8-10),  
50 5.9, 5.9 and 5.4 PgCO<sub>2</sub>e.yr<sup>-1</sup> (Hotspots, FAOSTAT, EDGAR, and Houghton), and  
51 Agricultural sectors: 1.9 (1.5-2.5), 2.5, 2.1, and 2.0 PgCO<sub>2</sub>e.yr<sup>-1</sup> (Hotspots, FAOSTAT,  
52 EDGAR, and EPA). However, this agreement was lost when disaggregating by sources,  
53 continents, and gases, particularly for the forest sector, with fire leading the differences.  
54 Agricultural emissions were more homogeneous, especially livestock, while croplands were  
55 the most diverse. CO<sub>2</sub> showed the largest differences among datasets. Cropland soils and  
56 enteric fermentation led the smaller N<sub>2</sub>O and CH<sub>4</sub> differences. Disagreements are explained  
57 by differences in conceptual frameworks (i.e. carbon-only vs multi-gas assessments,  
58 definitions, land use versus land cover, etc), in methods (Tiers, scales, compliance with  
59 Intergovernmental Panel on Climate Change (IPCC) guidelines, legacies, etc) and in  
60 assumptions (i.e. carbon neutrality of certain emissions, instantaneous emissions release, etc)  
61 that call for more complete and transparent documentation for all the available datasets.  
62 Enhanced dialogue between the carbon (CO<sub>2</sub>) and the AFOLU (multi-gas) communities is  
63 needed to reduce discrepancies of land use estimates.

64

## 65 **1. INTRODUCTION**

66 Modelling studies suggest that to keep the global mean temperature increase to less than 2°C  
67 and to remain under 450 ppm of CO<sub>2</sub> by 2100, CO<sub>2</sub> emissions must be cut 41-72% below  
68 2010 levels by 2050 (IPCC, 2014), and global emissions levels must be reduced to zero (a  
69 balance between sources and sinks) before 2070 and below zero, through removal processes,  
70 after that (Anderson, 2015; UNEP, 2015). To reach these ambitious goals, tremendously rapid

71 improvements in energy efficiency and nearly a quadrupling of the share of zero and low  
72 carbon energy supply (i.e. renewables, nuclear energy, and carbon dioxide capture and storage  
73 (CCS), including bioenergy (BECCS)) would be needed by 2050 (IPCC, 2014; Friedlingstein  
74 et al., 2014; UNEP, 2015). Since there is no scientific evidence on the feasibility of CCS  
75 technologies (Anderson, 2015), renewables and the land use sector are among the most  
76 plausible options (Canadell and Schulze, 2014). Optimistic estimates suggest that the AFOLU  
77 sector (here indistinctively also called land use sector) could contribute from 20 to 60% of the  
78 total cumulative abatement to 2030 including bioenergy (Smith et al., 2014).

79

80 The Agriculture, Forestry, and other Land Use (AFOLU) sector roughly contributes with a  
81 quarter (10-12 PgCO<sub>2</sub>e.yr<sup>-1</sup>) of the total anthropogenic greenhouse gas (GHG) emissions (50  
82 PgCO<sub>2</sub>e.yr<sup>-1</sup>) (Smith et al., 2014) through a few human activities: deforestation, forest  
83 degradation, and agriculture including cropland soils, paddy rice, and livestock (Smith et al.,  
84 2014). Despite the acknowledged importance of the emissions from the land use sector in  
85 global mitigation strategies, assessing GHG emissions and removals from this sector remains  
86 technically and conceptually challenging (Abad-Viñas et al., 2014; Ciais et al., 2014). This  
87 challenge relates to an incomplete understanding of the processes that control the emissions  
88 from the land use sector (Houghton et al., 2012), especially post-disturbance dynamics (Frank  
89 et al., 2015; Poorter et al., 2016) and to various sources of error that range from inconsistent  
90 definitions, methods, and technical capacities (Romijn et al., 2012, 2015; Abad-Viñas et al.,  
91 2014), to special features of the land use sector such as legacy and reversibility/non-  
92 permanence effects (Estrada et al., 2014), or to the difficulty to separate anthropogenic from  
93 natural emissions (Estrada et al., 2014; Smith et al., 2014). As a result, the AFOLU emissions  
94 are the most uncertain of the all the sectors in the global budget, reaching up to 50 percent of  
95 the emissions mean (Houghton et al., 2012; Smith et al., 2014; Tubiello et al., 2015). This is

96 important since uncertainties jeopardize the effectiveness of the AFOLU sector to contribute  
97 to climate change mitigation. Thus, country compliances to their mitigation targets are likely  
98 to be controversial when the uncertainty is equal to or greater than the pledged emission  
99 reductions (Grassi et al., 2008; Pelletier et al., 2015).

100

101 Currently, data on AFOLU emissions are available through national greenhouse gas  
102 inventories, which are submitted to the United Nations Framework Convention on Climate  
103 Change (UNFCCC), but these national estimates cannot be objectively compared due to  
104 differences in definitions, methods, and data completeness (Houghton et al., 2012; Abad-  
105 Viñas et al., 2014). More comparable AFOLU data are offered in global emission databases  
106 such as EDGAR or FAOSTAT (Smith et al., 2014; Tubiello et al., 2015), or more sectorial  
107 datasets such as the Houghton's Forestry and other Land Use (FOLU) data (Houghton et al.,  
108 2012), and the US Environmental Protection Agency non-CO<sub>2</sub> emissions for agriculture -  
109 including livestock (USEPA, 2013). While national inventories and global databases are  
110 currently the best bottom up emissions data we count on, their utility to inform on what the  
111 atmosphere receives has been contested. Late research shows disagreements between the  
112 trends of reported emissions and atmospheric growth since 1990 for CO<sub>2</sub> (Francey et al.,  
113 2010, 2013a, 2013b), for CH<sub>4</sub> (Montzka et al., 2011), and for N<sub>2</sub>O (Francey et al., 2013b). In  
114 the case of CO<sub>2</sub>, Francey *et al.* conclude that the differences between atmospheric and  
115 emission trends for CO<sub>2</sub> might be more related to under-reported emissions (~9 PgC = 33  
116 PgCO<sub>2</sub> for the period 1994-2005), than to adjustments in the terrestrial sinks (i.e. increased  
117 CO<sub>2</sub> removals in oceans and forests). On the other hand, global AFOLU databases suffer from  
118 inconsistencies that lead to global CO<sub>2</sub>e emissions differences of up to 25% (2000-2009)  
119 (Tubiello et al., 2015): 12.7 vs 9.9 PgCO<sub>2</sub>e.yr<sup>-1</sup> for EDGAR and FAOSTAT, respectively.  
120 These datasets also disagreed in the contribution of the AFOLU sector to the total

121 anthropogenic budget in 2010 (i.e. 21% and 24% for FAOSTAT vs EDGAR) (Tubiello et al.,  
122 2015), and on the relative share of the emissions from agriculture versus FOLU since 2010  
123 Tubiello et al., 2015). Thus, while EDGAR implies a relatively equal contribution (IPCC,  
124 2014), FAOSTAT reports agricultural emissions being larger contributors to the total  
125 anthropogenic budget ( $11.2\pm 0.4\%$ ) than forestry and other land uses ( $10\pm 1.2\%$ ) (Tubiello et  
126 al., 2015), with a steady growth trend of 1% since 2010.

127

128 Understanding the inconsistencies among AFOLU datasets is an urgent task since they  
129 preclude our accurate understanding of land-atmosphere interactions, GHG effects on climate  
130 forcing and, consequently, the utility of modelling exercises and policies to mitigate climate  
131 change (Houghton et al., 2012; Grace et al., 2014; Smith et al., 2014; Sitch et al., 2015; Tian  
132 et al., 2016). The land use sector plays a prominent role in the Paris Agreement (Art.5), with  
133 many countries including it as mitigation targets in their Nationally Determined Contributions  
134 (NDCs) (Grassi and Dentener, 2015; Richards et al., 2015; Streck, 2015). It is then urgent to  
135 understand how much and why different AFOLU datasets differ in their emissions estimates,  
136 so that we can better navigate countries' land-based mitigation efforts, and help to validate  
137 their proposed claims under the UNFCCC.

138

139 Here we compare gross AFOLU emissions estimates for the tropics, for 2000-2005, from six  
140 datasets: FAOSTAT, EDGAR, "Houghton", "Baccini", the US Environmental Protection  
141 Agency data (EPA), and a recently produced, spatially explicit AFOLU dataset, that we will  
142 hereon call "Hotspots" (Roman-Cuesta et al., 2016). We aim to identify differences and  
143 plausible explanations behind: i) aggregated AFOLU, FOLU and Agricultural gross  
144 emissions, ii) disaggregated contributions of the emission sources for the different datasets,

145 iii) disaggregated contribution of the different gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O), and iv) national scale  
146 disagreements among datasets.

147

## 148 **2. METHODS**

### 149 *2.1 Study area*

150 Our study area covers the tropics and the subtropics, including the more temperate regions of  
151 South America (33° N to 54° S, 161° E to 117° W). Land use change occurs nowhere more  
152 rapidly than in the tropics (Poorter et al., 2016) so its study has global importance. Moreover,  
153 the tropics suffer from the largest data and capacity gaps (Romijn et al., 2012; 2015) and their  
154 need to access AFOLU data and understand their differences is more crucial. We selected the  
155 period 2000-2005 for being the common temporal range for all the datasets. This period is not  
156 for the recent past but that does not affect the comparative nature of this research. Our study  
157 area focuses at the country level and includes eighty countries, following Harris et al., (2012).  
158 We ran the comparisons on gross emissions. While gross and net emissions are equally  
159 important, they offer different information (Richter and Houghton, 2011; Houghton et al.,  
160 2012). Net land use emissions consider the emissions by the sources and the removals by the  
161 sinks (i.e. forest growth, forest regrowth after disturbances, organic matter stored in soils) in a  
162 final emission balance where the removals are discounted from the emissions. Gross  
163 assessments can consider both the emissions produced by the sources (gross emissions) and  
164 the removals absorbed by the sinks (gross removals), but they are not balanced out. Gross  
165 emissions are useful to navigate mitigation implementation since they offer direct information  
166 on the sources and sinks that need to be acted upon through policies and measures to enhance  
167 and promote mitigation (see further information on net and gross alternatives in Roman-  
168 Cuesta et al., 2016).

169

170 **2.2 AFOLU datasets**

171 *Hotspots* (Roman-Cuesta et al., 2016): this is a multi-gas (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) spatially explicit  
172 (0.5°) database on gross AFOLU emissions and associated uncertainties for the tropics and  
173 subtropics for the period 2000-2005, at Tier 2 and Tier 3 levels (see Supplement for the  
174 definition of Tiers). This database locates the hotspots of tropical AFOLU emissions, which  
175 should help to estimate mitigation potentials, and prioritize the areas and the land activities  
176 that require most urgent mitigation action. It combines available published GHG datasets for  
177 the key sources of emissions in the AFOLU sector as identified by the Fifth Assessment  
178 Report of the Intergovernmental Panel on Climate Change (AR5, Smith et al., 2014):  
179 deforestation, forest degradation (fire, wood harvesting), crop soils, paddy rice, and livestock  
180 (enteric fermentation and manure management). It also includes agricultural peatland  
181 decomposition using Tier 1 emission factors (see details in Roman-Cuesta et al., 2016). .  
182 Forest emissions focus on aboveground biomass, with the exception of peat fires. More  
183 detailed methodological information is available in Roman-Cuesta et al., (2016).

184

185 *FAOSTAT*: covers agriculture, forestry and other land uses and their associated emissions of  
186 CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, following IPCC, 2006 Guidelines at Tier 1 (Tubiello et al., 2013, 2014).  
187 Emissions are estimated for nearly 200 countries, annually, for the reference period of 1961–  
188 2012 (agriculture) and 1990–2012 (FOLU), based on national activity data submitted by  
189 countries and further collated by FAO. Projected emission data are available for 2030 and  
190 2050. FAOSTAT includes estimates of emissions from biomass fires, peatland drainage and  
191 fires, based on geo-spatial information, as well as on forest carbon stock changes (both  
192 emissions and removals) based on national-level FAO Forest Resources Assessment data  
193 (FRA 2010).

194



195 *EDGAR*: The Emissions Database for Global Atmospheric Research (EDGAR) provides  
196 global GHG emissions from multiple gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs and SF<sub>6</sub>) at 0.1°  
197 and country levels. The EDGAR database covers all IPCC sectors (energy, industry, waste  
198 management, and AFOLU), mostly applying IPCC 2006 guidelines for emission estimations  
199 (EDGAR 2012). We downloaded the EDGAR's 4.2 Fast Track 2010 (FT 2010). FT 2010  
200 emissions cover the period 2000-2010 in an annual basis, at the country level.

201

202 "*Houghton*": Houghton's bookkeeping model calculates the net and gross fluxes of carbon  
203 (CO<sub>2</sub> only) between land and atmosphere that result from land management (Houghton, 1999,  
204 2012; Houghton and Hackler, 2001; Houghton et al., 2012). The net estimate includes  
205 emissions of CO<sub>2</sub> from deforestation, shifting cultivation, wood harvesting, wood debris  
206 decay, biomass burning (for deforestation fires only, peatland fires were not included in our  
207 version of their data), and soil organic matter from cultivated soils. It also includes sinks of  
208 carbon in forests recovering from harvest and agricultural abandonment under shifting  
209 cultivation.. The model does, however, not include forests that are not logged, cleared or  
210 cultivated. Rates of growth and decomposition are ecosystem specific and do not vary in  
211 response to changes in climate, CO<sub>2</sub> concentrations, or other elements of environmental  
212 change. Therefore, forests grow (and wood decays) at the same rates in 1850 and 2015.  
213 Unlike other databases all carbon in the ecosystem considered is accounted for: live  
214 vegetation, soil, slash (woody debris produced during disturbance), and wood products. We  
215 downloaded regional annual emissions from the TRENDS (1850-2005) dataset for the tropics:  
216 Central and South (CS) America, tropical Africa and South and South East Asia. Only net  
217 emissions were available. No spatially disaggregated data were offered (i.e. countries).  
218 Houghton's data are, unlike all the other datasets, net aggregated FOLU estimates, for CO<sub>2</sub>-  
219 only.

220

221 *"Baccini"*: These are gross FOLU tropical emissions published by Baccini et al., (2012). Data  
222 are gross emissions for the period 2000-2010 disaggregated into: deforestation (4.18  
223 PgCO<sub>2</sub>.yr<sup>-1</sup>), wood harvesting (1.69 PgCO<sub>2</sub>.yr<sup>-1</sup>), biomass burning (2.86 PgCO<sub>2</sub>.yr<sup>-1</sup>) and,  
224 wood debris decay (3.04 PgCO<sub>2</sub>.yr<sup>-1</sup>). We excluded this last variable to make it more  
225 comparable to the other datasets, where CWD is frequently excluded (Table 1). Baccini's  
226 estimates refer to a tropical area slightly smaller than our study region and they are offered as  
227 an aggregated value (no continental or country data available)

228

229 *The US Environmental Protection Agency (EPA)*: global non-CO<sub>2</sub> projected emissions for the  
230 period 1990-2030 for the Agriculture, Energy, Industrial Processes and Waste sectors, for  
231 more than twenty gases. EPA uses future net emissions projections of non-CO<sub>2</sub> GHGs as a  
232 basis for understanding how future policy and short-term, cost-effective mitigation options  
233 can affect these emissions. EPA follows the Global Emissions Report, which uses a  
234 combination of country-prepared, publicly-available reports consistent with IPCC guidelines  
235 and guidance (USEPA, 2013). When national emissions estimates were unavailable, EPA  
236 produced its own non-CO<sub>2</sub> emissions using IPCC methodologies (i.e., international statistics  
237 for activity data, and the default IPCC Tier 1 emission factors). Deviations to this  
238 methodology are discussed in each of the source-specific methodology sections of USEPA  
239 (2012). No FOLU estimates are included in this dataset. We downloaded agricultural  
240 emissions offered as 5-year intervals at country level, disaggregated by gas (N<sub>2</sub>O and CH<sub>4</sub>),  
241 and by emission sources.

242

243 *IPCC AR5*: The AR5 is a synthesis report, not a repository of global data. However, new  
244 AFOLU data are produced by the merging of peer-reviewed data such as Figures 11.2, 11.4,

245 11.5 and 11.8 in chapter 11 of the AR5 (Smith et al., 2014). We will contrast our six datasets  
246 against the data from these newly produced figures.

247

248 Table 1 shows a summary of key similarities and differences of the assessed AFOLU datasets  
249 and the data from the AR5. The exact variables used for each database are described in Table  
250 S1 in the Supplement.

251

### 252 *2.3 Estimating comparable gross AFOLU emissions for all datasets*

253 We focus on human-induced gross emissions only, excluding fluxes from unmanaged land  
254 (i.e. natural wetlands). We focus on direct emissions excluding indirect emissions whenever  
255 possible (i.e. nitrate leaching and surface runoff from croplands). Delayed fluxes (legacies)  
256 are important (i.e. underestimations of up to 62% of the total emissions when recent legacy  
257 fluxes are excluded) (Houghton et al., 2012) but are frequently omitted in GHG assessments  
258 that derive from remote sensing, such as the deforestation emissions used in the Hotspots  
259 database, which relies on Harris et al., 2012). Wood harvesting emissions also excluded  
260 legacy fluxes. We assumed instantaneous emissions of all carbon that is lost from the land  
261 after human action (Tier 1, IPCC 2006) (i.e. deforested and harvested wood), with no  
262 transboundary considerations (i.e. the emissions are assigned wherever the disturbance takes  
263 place, particularly important for Harvested Wood Products). Life-cycle substitution effects  
264 were neither considered for harvested wood (Peters et al., 2012). Some exceptions were  
265 allowed when data were already aggregated (i.e. for Houghton's and EPA's datasets we could  
266 not exclude indirect emissions linked to forest decay and agriculture, respectively), or because  
267 their legacy (past decay) estimates corresponded to an important source (i.e. EDGAR's post  
268 burned decay and decomposition emissions represent deforestation) (Tubiello et al., 2015).  
269 Databases include a diversity of emission sources and gases under AFOLU, not always

270 following IPCC requirements (some exclude peatland emissions, some include energy into the  
271 AFOLU emissions, some exclude non CO<sub>2</sub> emissions, etc). However, to compare the AFOLU  
272 emission estimates among databases we choose exactly the same sources: deforestation, wood  
273 harvesting, fire, livestock (enteric fermentation + manure management), cropland soil  
274 emissions, rice emissions, emissions from drained histosols), and the same gases CO<sub>2</sub>, CH<sub>4</sub>,  
275 and N<sub>2</sub>O, and documented what was included in each case (See Table S1 the Supplement).  
276 For the case of fire, for all the databases, we excluded CO<sub>2</sub> emissions that came from biomass  
277 burning in non-woody vegetation such as savannas and agriculture, since they are assumed to  
278 be in equilibrium with annual regrowth processes (for CO<sub>2</sub> gases only) (IPCC 2003, 2006).

279

#### 280 ***2.4 Correcting known differences among datasets estimates***

281 Tubiello *et al.* (2015) identified four main differences that resulted in larger estimates for the  
282 EDGAR data than for FAOSTAT, under the AFOLU estimates of the AR5 (Smith *et al.*,  
283 2014): 1. The inclusion of energy emissions under the agriculture budget, 2. Inclusion of  
284 savannah burning, 3. Higher rice emissions due to the use of the IPCC 1996 guidelines instead  
285 of the IPCC 2006 guidance, 4. FOLU's unresolved differences due to unclear metadata on  
286 EDGAR's proxy for deforestation (post burned decay and decomposition). We have corrected  
287 for the first two in our data comparison. No energy, and no CO<sub>2</sub> for savannah burning have  
288 been included in the AFOLU estimates in any of our analyses.

289

#### 290 ***2.5 Country emissions***

291 To characterize countries emissions' variability we estimated the standard deviations for the  
292 different emission sectors: i. forest (deforestation + fire + wood harvesting), ii. agriculture  
293 (cropland soils + paddy rice), iii. livestock, and the aggregated AFOLU emissions, for the  
294 three most complete datasets (Hotspots, FAOSTAT, EDGAR), per country. We grouped the

295 standard deviations into four percentiles to aggregate countries into levels of emission  
296 variability: high agreement (=low variability, low standard deviations, <25<sup>th</sup>percentile),  
297 moderate agreement (25<sup>th</sup>-50<sup>th</sup> percentiles), low agreement (25<sup>th</sup>-50<sup>th</sup> percentiles), and very  
298 low agreement (= very high variability, very high standard deviations, >75<sup>th</sup> percentile). See  
299 Supplement for a further discussion on issues regarding countries' emissions variability.

300

### 301 **3. RESULTS AND DISCUSSION**

#### 302 ***3.1 Aggregated AFOLU, FOLU and Agricultural emissions***

303 We found good agreement among datasets for the aggregated tropical scales with AFOLU  
304 values of 8.0 (5.5-12.2) (5<sup>th</sup>-95<sup>th</sup> percentiles), 8.4 and 8.0 PgCO<sub>2</sub>e.yr<sup>-1</sup> (for the Hotspots,  
305 FAOSTAT and EDGAR, respectively). FOLU (deforestation and forest degradation)  
306 contributed with 6.0 (3.8-10), 5.9, 5.9 and 5.4 PgCO<sub>2</sub>e.yr<sup>-1</sup> for the Hotspots, FAOSTAT,  
307 EDGAR, and Houghton datasets respectively. Agriculture (livestock, cropland soils and rice  
308 emissions) reached 1.9 (1.5-2.5), 2.5, 2.1, and 2.0 PgCO<sub>2</sub>e.yr<sup>-1</sup> for the Hotspots, FAOSTAT,  
309 EDGAR, and EPA datasets respectively (Figure 1, Table 2). Forest emissions represented  
310 ≥70% of the tropical AFOLU gross mean annual budget for 2000-2005 (the Hotspots database  
311 and Houghton showing the highest and the lowest estimates), and agriculture represented the  
312 remaining 25-30% AFOLU emissions (FAOSTAT and Hotspots showing the highest and the  
313 lowest values). Houghton's FOLU value (5.4 PgCO<sub>2</sub>.yr<sup>-1</sup>) is a net estimate that includes  
314 carbon dynamics associated to forest land use changes, and forest removals from areas under  
315 logging and shifting cultivation and it is, as expected, lower than the forest gross emissions.  
316 Its value for the tropics was, however, higher than the net FOLU value used in the IPCC AR5  
317 (4.03 PgCO<sub>2</sub>e.yr<sup>-1</sup> for 2000-2009) (Houghton *et al.* 2012). Since boreal and temperate forest  
318 sinks are reported to be quasi-neutral (Houghton et al., 2012), these differences are unclear.  
319 There is a variety of Houghton' net FOLU estimates in current bibliography (i.e. 4.03

320 PgCO<sub>2e</sub>.yr<sup>-1</sup> for 2000-2009 in Smith *et al.* (2012), 4.9 for 2000 and 4.2 for 2010 (Tubiello et  
321 al., 2015) that likely correspond to different updates of the same dataset, but create confusion  
322 and would call for verified official values that could be consistently used.

323

324 The IPCC AR5 offers a FOLU gross value for the tropics of ca. 8.4 PgCO<sub>2</sub>.yr<sup>-1</sup> (2000-2007)  
325 (Fig 11.8 in AR5, Smith et al., 2014) (Fig S1, Supplement) which corresponds to Baccini's  
326 estimates using Houghton's bookkeeping model. This value is in the upper range of the  
327 Hotspots gross FOLU emissions: 6 (3.8-10) PgCO<sub>2e</sub>.yr<sup>-1</sup> (2000-2005), and higher than the  
328 mean gross FOLU emissions from all the other datasets (approx. 6 PgCO<sub>2e</sub>.yr<sup>-1</sup>) (Table 2).  
329 The time periods are not identical and we do not compare the same gases (i.e. the  
330 bookkeeping model focuses on CO<sub>2</sub> only, while we run a multi-gas assessment). However, the  
331 differences mainly relate to unreported choices behind the inclusion/exclusion of emission  
332 sources and the description of their methods, in the AR5. Thus, the 8.4 PgCO<sub>2</sub>.yr<sup>-1</sup> gross  
333 estimate does not include fire, and has larger contributions from shifting cultivation (2.35  
334 PgCO<sub>2</sub>.yr<sup>-1</sup>) and wood-harvesting (2.49 PgCO<sub>2</sub>.yr<sup>-1</sup>), than the deforestation and wood-  
335 harvesting emissions in the Hotspots selected datasets (Figure 2). Numbers used in Figure  
336 11.8 also exclude other gross emissions offered in Baccini et al. (2012), which is the citation  
337 used in Fig. 11.8. Explicit, complete, and transparent documentation is encouraged for the  
338 next AFOLU figures in the IPCC Assessment Reports. Another consideration of AFOLU  
339 estimates in the Assessment Reports relates to the use of the bookkeeping model to estimate  
340 land use, land use change and forest (LULUCF) emissions. As useful as this model is, its  
341 framework does not follow the IPCC AFOLU guidelines (IPCC, 2006), particularly regarding  
342 the concept of managed land. Thus, forests that are on managed land but are not suffering  
343 from direct human activities are considered carbon neutral (Houghton *pers. comm.*). Partly  
344 because of that, the net emission estimates of LULUCF from Houghton et al. (2012) used in

345 the AR5 (4.03 PgCO<sub>2</sub>.yr<sup>-1</sup> 2000-2009) differ from LULUCF country reports for the same  
346 period, which are close to zero (Grassi and Dentener, 2015; Federici et al., 2016). The use of  
347 IPCC compliant models for the IPCC Assessment Reports, or/and some documentation that  
348 warned about these inconsistencies, would be useful in future assessments.

349

350 Emissions in the agricultural sector are mostly net, since sink effects in the soils are small and  
351 frequently temporal (USEPA, 2013; Smith et al., 2014). Comparisons against global  
352 agricultural emissions show that for the year 2000, global estimates more than doubled the  
353 Hotspots values (i.e. 5 and 5.5 PgCO<sub>2</sub>e.yr<sup>-1</sup> vs ca. 2 PgCO<sub>2</sub>e.yr<sup>-1</sup> in all datasets) (Tubiello et  
354 al., 2015) (Table 2), suggesting larger contributions of agricultural emissions from non-  
355 tropical countries. Unexplained methodological differences such as the inclusion or not of  
356 indirect emissions and the lack of an exhaustive list of the variables included in the  
357 agricultural emissions, difficult further comparisons.

358

### 359 ***3.2 Disaggregated gross emissions: contributions of the emission sources***

360 While the gross aggregated estimates suggested a good level of agreement among datasets  
361 (Figure 1), differences occur when comparing the emissions sources leading the AFOLU  
362 budgets (Figure 2). The FOLU sector showed the largest differences, mainly due to the  
363 estimates of forest degradation, and particularly fire (FAOSTAT and EDGAR showed the  
364 lowest and highest values). The forest sector is the most uncertain term in the AFOLU  
365 emissions due to both uncertainties in areas affected by land use changes and other  
366 disturbances, and by uncertain forest carbon densities (Houghton et al., 2012; Grace et al.,  
367 2014; Smith et al., 2014). Agricultural sources were more homogeneous (ca. 2 PgCO<sub>2</sub>e.yr<sup>-1</sup>  
368 for all datasets) (Figure 1), with livestock and cropland soil emissions as the most and least  
369 similar (Figure 2). The homogeneity in livestock emissions was expected since most datasets

370 use common statistics (FAO) to derive herd numbers per country.

371

### 372 3.2.1 Deforestation

373 Deforestation emissions were 2.9 (1.0-10.1), 3.7, and 2.5 and 4.2 PgCO<sub>2</sub>.yr<sup>-1</sup> (Hotspots,  
374 FAOSTAT, EDGAR, and Baccini, respectively), with Baccini and EDGAR showing the  
375 highest and the lowest values. Their values represent, however, very different scenarios: gross  
376 deforestation for the Hotspots and Baccini datasets (forest losses only), net deforestation for  
377 FAOSTAT (forest losses minus forest gains), and forest fire and post-burn decay for EDGAR  
378 (Table 3). The Hotspots (Harris et al., 2012) and Baccini et al., (2012) datasets offer gross  
379 deforestation estimates that rely on Hansen et al., (2010)'s forest cover loss areas. However,  
380 they report different tropical emissions (0.81 and 1.14 PgC.yr<sup>-1</sup>) because they use different  
381 carbon density maps: Harris *et al.* (2012) rely on Saatchi *et al.*(2011) and Baccini rely on  
382 Baccini *et al.*(2010). EDGAR does not provide a category for deforestation, and their Forest  
383 Fire and Decay category (5F) (Table 3, and Table S1 in the Supplement) is used as a proxy for  
384 deforestation (Tubiello et al., 2015). Such an approximation leads to underestimations since  
385 not all carbon losses from deforestation are necessarily associated with the use of fire  
386 (Tubiello et al., 2015). In spite of being net emissions, the deforestation estimates for  
387 FAOSTAT were higher than the gross estimates of Hotspots and Baccini. This is partly due to  
388 FAOSTAT's inclusion of fire emissions from humid tropical forests (see section 3.2.3), which  
389 the other datasets did not. Baccini's larger estimates of gross deforestation included more  
390 carbon pools than the other datasets (i.e. soil, Coarse Woody Debris (CWD), litter). Baccini *et*  
391 *al.* (2012) reported that their estimated gross and net emissions from tropical deforestation  
392 were the same value (4.2 Pg CO<sub>2</sub>.yr<sup>-1</sup>). The difference with Houghton's net emissions (5.4  
393 PgCO<sub>2</sub>.yr<sup>-1</sup>) (Figure 2) corresponds, then, to non-offset carbon emissions from other land uses  
394 and activities included in the bookkeeping model: degradation by logging and shifting



395 cultivation, decomposition and decay, and cultivated soils. Houghton's tropical net emissions  
396 for 2000-2005 are high, but lower than Houghton's reported net estimates in the 80's (7  
397 PgCO<sub>2</sub>.yr<sup>-1</sup>) (Houghton, 1999).

398

### 399 *3.2.2 Forest degradation: wood harvesting and fire emissions*

400 Forest degradation can be defined in many ways (Simula, 2009), but no single operational  
401 definition has been agreed upon by the international community (Herold et al., 2011a). It  
402 typically refers to a sustained human-induced loss of carbon stocks within forest land that  
403 remains forest land. In this study, similarly to Federici et al., (2015), we consider degradation  
404 any annual removal of carbon stocks that does not account for deforestation, without temporal  
405 scale considerations (i.e. time needed for disturbance recovery, or time to guarantee a  
406 sustained reduction of the biomass). We assessed two major degradation sources: wood  
407 harvesting and fire. Soil degradation is poorly captured in many datasets, and mainly focuses  
408 on fire in equatorial Asian peatland forests and drained peatlands (Hooijer et al., 2010).  
409 Better understanding of the processes and emissions behind forest degradation would be key  
410 for climate mitigation efforts not only because forest degradation is a wide spread  
411 phenomenon (i.e. affects much larger areas than deforestation (Herold et al., 2011b)) but also  
412 because the lack of knowledge of net carbon effects frequently results in assumptions of  
413 carbon neutrality of the affected standing forests, particularly for fire (Houghton et al., 2012;  
414 Le Quéré et al., 2014), which is likely leading to an underestimation of forest and AFOLU  
415 emissions (Brando et al., 2014; Turetsky et al., 2015; Roman-Cuesta et al., 2016).

416

417 Gross emissions from forest degradation were larger than deforestation for the Hotspots,  
418 EDGAR and Baccini's datasets, with degradation-to-deforestation ratios of 108%, 120%, and  
419 128%, respectively. FAOSTAT had degradation emissions of 60% of the deforestation, partly

420 due to its anomalously low fire contribution (see next section). Houghton et al., (2012)  
421 pointed out that global FOLU net fluxes were led by deforestation with a smaller fraction  
422 attributable to forest degradation, while the opposite was true for gross emissions (degradation  
423 being 267% of deforestation emissions). This large ratio relates to their inclusion of shifting  
424 cultivation under degradation. This is a definition issue, which would not fit the definition of  
425 degradation chosen in this study, where a complete forest cover loss would represent  
426 deforestation and not degradation.

427

### 428 3.2.2.1 Fire

429 Fire led the gross forest degradation emissions in the tropics in 2000-2005 (Figure 2): 2 (1.1-  
430 2.7), 0.2, 3.4, 2.9 PgCO<sub>2</sub>e.yr<sup>-1</sup> for the Hotspots, FAOSTAT, EDGAR, and Baccini datasets,  
431 respectively) (Figure 2). The Hotspots estimates are conservative compared to Van der Werf  
432 et al., (2010)'s global emissions of 7.7 PgCO<sub>2</sub>e.yr<sup>-1</sup> for 2002-2007, due to the removal of CO<sub>2</sub>  
433 from deforestation fires (to avoid double counting with deforestation emissions), to the  
434 exclusion of fires in grasslands and agricultural residues, and to the Hotspots' smaller study  
435 area. FAOSTAT and EDGAR had the lowest and the highest fire values. FAOSTAT lowest  
436 values relate to omissions that are currently in the process of being corrected (Rossi *pers.*  
437 *comm.*): 1. the complete exclusion of CO<sub>2</sub> from fire in humid tropical forests and other forests  
438 (Table 3, Table S1), which FAOSTAT relocated as net forest conversion emissions, partly  
439 explains their larger deforestation values (FAOSTAT kept, however, CH<sub>4</sub> and N<sub>2</sub>O for fire in  
440 humid tropical forests and other forests), and 2. The use of default parameters for fuel in peats  
441 from the IPCC 2006 Guidelines instead of the new IPCC Wetland supplement which offer  
442 considerable higher values (Rossi et al., 2016). Moreover, FAOSTAT uses GFED3.0-burned  
443 area (Giglio et al., 2010) in their estimates while the other datasets use GFED3.0-emissions  
444 (Van der Werf et al., 2010). EDGAR fire emissions were the largest most likely because they

445 included decay. Their dataset considers some undefined “forest fires” (5A) and  
446 “wetland/peatland fires and decay” (5D) (Table 3; Table S1 in the Supplement). Peatland  
447 decay probably explains EDGAR’s larger emissions in Asia, while we assume that EDGAR’s  
448 highest fire emissions for CS America might respond to deforestation fires which were not  
449 included in the Hotspots to avoid double counting with deforestation, and relocated in  
450 FAOSTAT to deforestation emissions (Figure 3, Table 3). The Hotspots dataset showed  
451 higher gross fire emissions for Africa due to the inclusion of woodland fire, which EDGAR  
452 and FAOSTAT probably excluded. Baccini et al., (2012)’s fire emissions:  $2.9 \text{ PgCO}_2\text{e.yr}^{-1}$   
453 (2000-2010) derive from Houghton’s bookkeeping but it is unclear how these emissions were  
454 estimated.

455 In spite of the importance of fire as a degradation source, this variable is frequently  
456 incompletely included, either through unaccounted gases (i.e.  $\text{CH}_4$  and  $\text{N}_2\text{O}$  are excluded in  
457 the carbon community but their omission represent 17-34% of the gross  $\text{CO}_2$  fire emissions)  
458 (Valentini et al., 2014; Roman-Cuesta et al., 2016), or to unaccounted components (i.e fires in  
459 tropical temperate forests such as conifers or dry forests such as woodlands, are frequently  
460 excluded) (Houghton et al., 2012). Unaccounted fire emissions also derive from  
461 methodological choices (i.e. only inter-annual fire anomalies being considered) (Le Quéré et  
462 al., 2014), from poor satellite observations such as understory fires in humid closed canopy  
463 forests) (Alencar et al., 2006; 2012, Morton et al., 2013), or satellite fire omissions in certain  
464 regions (i.e. high Andean fires) (Bradley and Millington, 2006; Oliveras et al., 2014). Other  
465 omissions relate to the current exclusion of non-Asian peatland fires (i.e American tropical  
466 montane cloud forest peatland fires) (Asbjornsen et al., 2005; Roman-Cuesta et al., 2011;  
467 Oliveras et al., 2013; Turetsky et al., 2015).

468

469 Fire suffers, moreover, from a series of assumptions that do not apply so easily to other types  
470 of degradation: 1. Assuming a non-human nature of the fires (deforestation fire vs wildfires),  
471 which in tropical areas contrasts with multiple citations referring to the 90% human causality  
472 of fires (Cochrane et al., 1999; Roman-Cuesta et al., 2003; Alencar et al., 2006; Van der Werf  
473 et al., 2010). 2. Assuming *force-majeure* conditions that lead to non-controllable fires due to  
474 extreme climate conditions, which frequently results in incomplete assessment and reporting  
475 of emissions. This assumption contrasts with research on how human activities have seriously  
476 increased fire risk and spread in the tropics (Uhl and Kauffman, 1990; Laurance and  
477 Williamson, 2001; Roman-Cuesta et al., 2003; Hooijer et al., 2010), and clearly expose how  
478 most of the fires in the humid tropics would not occur in the absence of human influences  
479 over the landscape (Roman-Cuesta et al., 2003). 3. Assuming carbon neutrality and full  
480 biomass recovery after fire in standing forests. This is a generous assumption that contrasts  
481 with numerous studies on tropical forest die-back following fire events in non-fire adapted  
482 humid tropical forests (Cochrane et al., 1999; Barlow et al., 2008; Roman-Cuesta et al., 2011;  
483 Brando et al., 2012; Oliveras et al., 2013; Balch et al., 2015). All these phenomena casts  
484 doubts on the robustness of these assumptions and call for a much more comprehensive  
485 inclusion of fire emissions into forest degradation budgets.

486

#### 487 3.2.2.2 *Wood harvesting*

488 There is not a unique way to estimate wood harvesting emissions as exposed in the guidelines  
489 for harvested wood products of the IPCC (IPCC 2006). Assumptions regarding the final use  
490 of the wood products, decay times, substitution effects, international destination of the  
491 products and time needed for forests to recover their lost wood, can fully change the emission  
492 budgets. In our study, wood harvesting emissions were 1.2 (0.7-1.6), 2.0, 1.7 PgCO<sub>2</sub>.yr<sup>-1</sup> for  
493 the Hotspots, FAOSTAT and Baccini data, respectively (Tables 3, Table S1 in the

494 Supplement). Harvested wood products derive from FAO's country reports (i.e. FAOSTAT  
495 forest products). All datasets included fuel wood and industrial roundwood (Tables 3, Table  
496 S1). EDGAR excluded fuelwood from the AFOLU budget and placed it instead into the  
497 energy budget (EDGAR, 2012), which explains its absence in Figure 2. Wood harvesting  
498 emissions were larger in FAOSTAT than in the Hotspot data (Figure 2) partly due to the  
499 inclusion of some extra categories of fuels (i.e. charcoal and residues) that were not included  
500 in the Hotspot database (Table 3, Table S1 in the Supplement). Charcoal represents 26% of  
501 the total wood-harvesting emissions in FAOSTAT. Differences on wood harvesting affected  
502 more Asia and CS America (where the Hotspot data were half of FAOSTAT's), whilst Africa  
503 presented almost identical values (Figure 3), reasons for these continental differences are  
504 unclear. Baccini's high emissions on wood harvesting could partly relate to their inclusion of  
505 extra biomass due to felling damages (i.e. 20-67% of the AGB is damaged, and 20% is left  
506 dead in BGB) (Houghton, 1999).

507

### 508 *3.2.3 Livestock*

509 Livestock emissions were the most homogeneous among the emissions sources (Figure 2)  
510 with estimates of 1.2 (0.8-1.5), 1.1, 1.2, 1.1 PgCO<sub>2e</sub>.yr<sup>-1</sup> for the Hotspots, FAOSTAT,  
511 EDGAR and EPA respectively, in range with the estimates in the AR5 (Fig 11.5 in Smith et  
512 al., 2014). Values were similar in spite of deriving from different Tiers (i.e. Tier 3 for Herrero  
513 et al., (2013), Tier 1 for FAOSTAT and EDGAR. EPA used Tier 3 but for incomplete data  
514 series, otherwise Tier 1 was applied (USEPA, 2013)). All datasets included enteric  
515 fermentation (CH<sub>4</sub>) and manure management (N<sub>2</sub>O, CH<sub>4</sub>). All of them relied on FAO data for  
516 livestock heads, although they used different years (i.e. 2000 for Herrero et al., (2013) data in  
517 the Hotspots, and 2007-2010 for EDGAR). From a continental perspective, FAOSTAT and  
518 EDGAR estimates were the closest while the Hotspots and EPA's were less similar. The

519 Hotspots showed higher emissions for Africa and Asia and lower for CS America, than the  
520 other three datasets. Divergences likely relate to different Tiers. CS America and Asia showed  
521 the highest values, with Africa following closely (Figure 3), similar to what is reported in the  
522 AR5 (Smith et al., 2014). Globally, livestock is the largest source of CH<sub>4</sub> emissions, with  
523 three-fourth of the emissions coming from developing countries, particularly Asia (USEPA,  
524 2013, Tubiello et al., 2014). Three out of the top-5 emitting countries are in the tropics:  
525 Pakistan, India and Brazil (USEPA, 2013) and while Asia hosts the largest livestock  
526 emissions, the fastest growing trends in 2011 correspond to Africa (Tubiello et al., 2014).

527

#### 528 *3.2.4 Cropland emissions*

529 The estimates of cropland emissions reached values of 0.18 (0.16-0.19), 0.56, 0.6 and 0.64  
530 PgCO<sub>2e</sub>.yr<sup>-1</sup> for the Hotspots, FAO, EDGAR and EPA datasets respectively, for N<sub>2</sub>O and CO<sub>2</sub>  
531 from changes in soil organic carbon content. Cropland soil emissions (N<sub>2</sub>O and soil organic  
532 carbon stocks (CO<sub>2</sub>) heavily depend on land management practices (i.e. tillage, fertilization  
533 and irrigation practices) and climate (Crowther et al., 2015). We chose exactly the same land  
534 practices in all datasets to allow comparisons (Table 3, S1 in the Supplement). For this reason,  
535 we excluded N<sub>2</sub>O emissions from grassland soils, drainage of organic soils, and restoration of  
536 degraded lands (Table 3). This restrictions resulted in lower emissions than those estimated  
537 for cropland soils in the AR5 (Fig. 11.5 in Smith et al., 2014). The Hotspots and EPA showed  
538 the lowest and the highest estimates (Figures 2, 3). With the exception of the Hotspots, the  
539 other datasets agreed well at the tropical scale, with FAOSTAT and EDGAR being almost  
540 identical, also at continental scales. EPA disagreed more than the other datasets at the  
541 continental scales, with underestimations for Asia, probably related to the parameterization of  
542 their emission model. All three datasets used FAO's activity data, and for EDGAR and  
543 FAOSTAT the same emission factors must have been used. The Hotspot showed anomalously

544 low emissions partly because it only included six major crop types (maize, soya, sorghum,  
545 wheat, barley, millet) for which the emission model (DAYCENT) counted on reliable  
546 parametrization (*Ogle pers. comm*). Emissions from other important crops in the tropics (i.e.  
547 sugar cane, tobacco, tea, etc) were excluded, as well as emissions from croplands in organic  
548 soils, due to model constraints.

549

### 550 *3.2.5 Peatland drainage for agriculture*

551 Estimates of drained peatlands (mainly for agricultural purposes) suggest large omissions in  
552 the Hotspots database with emissions one order of magnitude lower ( $28 \text{ TgCO}_2\text{e.yr}^{-1}$ ) than  
553 FAOSTAT (ca.  $500 \text{ TgCO}_2\text{e.yr}^{-1}$ ) and one order of magnitude lower than the values reported  
554 for peatland drainage in Asia alone (Hooijer et al. 2010) ( $355\text{-}855 \text{ TgCO}_2\text{e.yr}^{-1}$ ). The lower  
555 values in the Hotspots dataset relate to much smaller agricultural areas with histosols (0.4 mill  
556 ha) than those reported by FAOSTAT for the same countries (7mill ha). This area difference  
557 is partly due to the methodological approach used by Ogle et al. (2013) where only six major  
558 crop covers are considered: maize, wheat, sorghum, soya beans, millet and barley, and partly  
559 to the unmatching spatial scales of histosols and croplands (i.e. 1km for histosols and 50km  
560 for croplands) which result in underestimations of the final area.

561

### 562 *3.2.6 Paddy rice*

563 When paddy fields are flooded, decomposition of organic material gradually depletes the  
564 oxygen present in the soil and floodwater, causing anaerobic conditions in the soil that favour  
565 methanogenic bacteria that produce  $\text{CH}_4$ . Some of this  $\text{CH}_4$  is dissolved in the floodwater, but  
566 the remainder is released to the atmosphere, primarily through the rice plants themselves. Net  
567 emission estimates for paddy rice were 0.55 (0.4-0.833), 0.33, 0.37,  $0.30 \text{ PgCO}_2\text{e.yr}^{-1}$  for the  
568 Hotspots, FAOSTAT, EDGAR and EPA datasets, respectively. The Hotspots showed the

569 highest emissions (Figure 2), but only in Asia (Figures 3). Part of the reason behind these  
570 differences refers to the final gases estimated in Li et al., (2013)'s which included CH<sub>4</sub>, N<sub>2</sub>O  
571 and decomposition of Soil Organic Carbon (SOC) (CO<sub>2</sub>) (Table 3, S1), while the others only  
572 focused on CH<sub>4</sub>. In Li et al., (2013)'s estimates, N<sub>2</sub>O were 48% of the CH<sub>4</sub> emissions,  
573 explaining the doubled emissions in the Hotspots database. SOC was a sink, with -0.076  
574 PgCO<sub>2</sub>.yr<sup>-1</sup>.

575  
576 Based on the explanations above, Table 4 points out the likely least reliable emission sources  
577 for each dataset considering disagreements among emission estimates due to  
578 biased/divergent/incomplete definitions and methods. Houghton's sinks are suggested as least  
579 reliable since they suffer from compatibility issued with IPCC guidance and exclude sinks  
580 from non-disturbed areas and sinks from forests undergoing disturbances other than wood  
581 harvesting or recovery from shifting cultivation (Grassi and Dentener 2015; Federici et al.,  
582 2016).

583

### 584 ***3.3 Differences in the relative contribution of greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O)***

585 GHG emissions (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) showed good agreement at the sectoral level (FOLU and  
586 agriculture) (Figure 5), that disappeared at the disaggregated level (Figure 6). CO<sub>2</sub> showed the  
587 largest disagreements among datasets and gases, led by forests emissions and particularly fire.  
588 SOC accumulation was reported in the Hotspots data (Li et al., 2013) but it is uncertain if it is  
589 included in the other datasets.

590

591 Non-CO<sub>2</sub> emissions showed lower variability than CO<sub>2</sub> (Figure 6). Livestock led CH<sub>4</sub>  
592 emissions and showed the largest differences among datasets, with the Hotspot data (Herrero  
593 et al., (2013) having the lowest CH<sub>4</sub> emissions, which were compensated with larger N<sub>2</sub>O than



594 the other datasets (Figure 6 b,c). At a global level, wetlands dominates natural CH<sub>4</sub> emissions,  
595 while agriculture and fossil fuels represent 2/3 of all human emissions, with smaller  
596 contributions coming from biomass burning, the oceans, and termites (Montzka et al., 2011).  
597 Fire non-CO<sub>2</sub> emissions were quite similar among datasets, confirming that FAOSTAT  
598 omissions were CO<sub>2</sub> related (see section 3.2.3). Thus, as exposed in FAOSTAT's metadata,  
599 only N<sub>2</sub>O and CH<sub>4</sub> are considered in forest fires, excluding CO<sub>2</sub> from aboveground biomass.  
600 As expected, N<sub>2</sub>O emissions in crops showed large differences, with the Hotspots having the  
601 lowest values (3 times lower). Rice N<sub>2</sub>O emissions were omitted in all datasets except the  
602 Hotspots (Li et al., 2013), which also included SOC.

603

604 The importance of multigas assessments relates to their role in climate change mitigation due  
605 to their radiative forcing (RF), understood as a measure of the warming strength of different  
606 agents (gases and not gases) in causing global warming (W.m<sup>-2</sup>). CO<sub>2</sub> is the most (400ppm in  
607 2015) and longest living gas which makes it the leading force of global warming (Anderson  
608 2012). Non-CO<sub>2</sub> GHG are less abundant in the atmosphere (1,774 ppb and 319 ppb for CH<sub>4</sub>  
609 and N<sub>2</sub>O in 2005 respectively) but have larger warming potentials (x 28 for CH<sub>4</sub>) and (x 265  
610 for N<sub>2</sub>O) (AR4) but shorter lifetimes than CO<sub>2</sub> (~9 and ~120 years, respectively). In spite of  
611 their shorter lifespans they offer an additional opportunity to mitigate climate change  
612 (Montzka et al., 2011) partly because they play a role in atmospheric chemistry that  
613 contributes to short-term warming (Montzka et al., 2011) and partly because their presence  
614 counteracts CO<sub>2</sub> terrestrial sinks (Tian et al., 2016),

615

### 616 ***3.4 Country level emissions***

617 Country comparisons showed poor agreement among datasets for all the emission sectors,  
618 particularly for the largest emitters (i.e. Brazil, Argentina, India, Indonesia) (Figures 7, 8).

619 Forests led the AFOLU disagreements (as observed by the similarity of Figure 7 a,b). From a  
620 continental perspective, Central and South America showed more countries with high levels  
621 of disagreement, suggesting the need for further data research.

622

### 623 *3.5 Some reflections on the datasets*

#### 624 *3.5.1 Original goals*

625 Different datasets were developed for different purposes that have influenced the methods and  
626 approaches chosen to estimate their land use GHGs. Thus, EDGAR was created with an air  
627 pollution focus making its land emissions weaker. Contrastingly, FAOSTAT carries FAO's  
628 focus on land particularly agriculture (data available since the 60s), with forest data added  
629 later through the FRA assessments (1990, 2005, 2010, 2015). The 'Hotspot' database was  
630 created to identify the areas with the largest land use emissions in the tropics (emissions  
631 hotspots), while Houghton's accent is on historical LULUCF emission trends (since 1850).  
632 EPA concentrates on industrial, energy, and agricultural emissions -forests are excluded- with  
633 an interest on human health and mitigation. Moreover, due to its long existence, several  
634 datasets rely on FAOSTAT's long-term agricultural data, which is probably the reason behind  
635 the higher homogeneity of agricultural emission estimates (i.e. crops, rice, and livestock)  
636 among datasets). FAOSTAT's forest emissions use FRA data, which get updated every 5  
637 years. Different FRA versions strongly influence forest emission estimates which makes it  
638 important to acknowledge the FRA version used when contrasting FAOSTAT emissions.  
639 when comparing estimates (i.e. differences up to 22% between the forest sink estimates using  
640 FRA2015 and FRA2010 have been reported by Federici et al., 2015). Similarly, official  
641 updates of Houghton's bookkeeping TRENDS data, as well as researchers' self-tuned  
642 versions of his model, result in emission differences that are difficult to track.

643

644 *4.2 IPCC guidelines and guidance:* Under the UNFCCC, countries are requested to use the  
645 latest IPCC AFOLU guidelines to estimate their GHG emissions (i.e. IPCC 2006 and 2003 for  
646 developed and developing countries, respectively). The use of different guidelines, Tiers, and  
647 approaches influences the final emission estimates. Compliance with IPCC has two main  
648 consequences: 1. the total area selected to report emissions, and 2. the choice of *land use* over  
649 *land cover*. In the first case, under IPCC guidance, the total area selected to report emissions  
650 would include all the land under human influence (the *managed land* concept, which includes  
651 areas under active and non-active management). Houghton’s bookkeeping model (and the  
652 carbon modelling community in general) do not comply well with the *managed land* concept,  
653 resulting in different net emissions from forest land uses and land use changes (LULUCF)  
654 than IPCC compliant country emissions (Grassi and Dentener, 2015; Federici et al., 2016). In  
655 the second case, the selection of *land uses* instead of *land covers* has partly been behind the  
656 recent controversy between FAO and the Global Forest Watch’s reported estimates on  
657 deforestation trends (Holmgren 2015). Estimates of deforestation that rely on *land cover* are  
658 higher than those using *land use*, since forest losses under forest land uses -that remain forest  
659 land use- are not considered deforestation (i.e. logged areas will regrow). In our analysis,  
660 FAO and Houghton relies on *land use* for deforestation, while the ‘Hotspots’ and EDGAR  
661 rely on *land cover*. FAOSTAT and the ‘Hotspots’ rely on the 2006 IPCC Guidelines for  
662 National Greenhouse Gas Inventories (IPCC, 2006). FAOSTAT uses Tier 1 and standard  
663 emission factors, while the ‘Hotspots’ use a combination of Tiers (Tier 3 for all emissions  
664 except wood harvesting and cropland emissions over histosols that rely on Tier 1). EDGAR  
665 reports the use of 2006 IPCC Guidelines for the selection of the emission factors but some of  
666 their methodological approaches are not always consistent with IPCC guidelines (i.e.  
667 deforestation expressed as the decay of burned forests, wood-harvesting is part of the energy  
668 sector, agricultural energy balances are included in the AFOLU budget). EPA methods are

669 reported to be consistent with IPCC guidelines and guidance, with Tier 1 methodologies used  
670 to fill in missing or unavailable data (USEPA, 2013).

671

#### 672 **4. CONCLUSIONS**

673 The Paris Agreement (COP21) counts on the Nationally Determined Contributions (NDCs) as  
674 the core of its negotiations to fight climate change. As March 2016, 188 countries had  
675 submitted their NDCs under the UNFCCC (FAO, 2016) with agriculture (crops, livestock,  
676 fishery and aquaculture) and forests as prominent features in meeting the countries' mitigation  
677 and adaptation goals (86% percent of the countries include AFOLU measures in their NDCs,  
678 placing it second after the energy sector) (FAO, 2016). However, there exists large variability  
679 in the way countries present their mitigation goals, and quantified sector-specific targets are  
680 rare (FAO, 2016). Variability relates not only to the lack of a standardized way to report  
681 mitigation commitments under the NDCs, but also to uncertainties and gaps in the AFOLU  
682 data. The Paris Agreement relies on a 5-year cycle stock-taking process to enhance mitigation  
683 ambition, and to keep close to the 2°C target. To be effective and efficient, stock-taking needs  
684 robust, transparent and certain numbers (at least with known uncertainties). This is true both  
685 for national emission reports and NDCs, but also for the global datasets that can be used to  
686 review the feasibility of countries' mitigation claims, and the real space for further mitigation  
687 commitments. We have here compared the gross AFOLU emissions of six datasets to search  
688 for disagreements, gaps, and uncertainties, focusing on the tropical region. Conclusions  
689 depend on the spatial scale. For the tropical scale:

- 690 • Data aggregation offers more homogeneous emission estimates than disaggregated  
691 data (i.e.continental level, gas level, emission source level).
- 692 • Forest emissions are the most uncertain of the AFOLU sector, with deforestation  
693 having the highest uncertainties.

- 694       • Agricultural emissions, particularly livestock, are the most homogeneous of the  
695       AFOLU emissions.
- 696       • Forest degradation, both fire and wood harvesting, show the largest variabilities  
697       among databases.
- 698       • CO<sub>2</sub> is the gas with longer-term influence in climate change trends, but it remains the  
699       most uncertain among the AFOLU gases and the most variable, in absolute value,  
700       among datasets (Figure 5) Fire leads this variability (Figure 6).
- 701       • Among the non-CO<sub>2</sub> gases, N<sub>2</sub>O showed the most variable emission estimates, in  
702       absolute value, in all the emission sources and for all the datasets (Figure 6).
- 703       • Emissions from histosols/peatlands remain incomplete or fully omitted in most  
704       datasets.

705 For the country and continental scales:

- 706       • Large emitters show the highest levels of data disagreement in the tropics, enhancing  
707       the need for data improvement to guarantee effective mitigation action.
- 708       • Forest lead the emission disagreement in the total AFOLU emissions.
- 709       • Central and South America showed the largest continental disagreements on emission  
710       data for all the land sectors.

711

## 712 ***4.1 Next steps***

### 713 *4.1.1 Enhancing dialogue between the carbon and the AFOLU research communities*

714 Research ran by the carbon community is pivotal for AFOLU assessments and while these  
715 two research communities overlap, they do not focus on exactly the same topics. The carbon  
716 community works with CO<sub>2</sub> emissions-only, fully excluding non-CO<sub>2</sub> gases, particularly N<sub>2</sub>O.  
717 It moreover rather focuses on forests and associated land use changes, excluding emissions  
718 from agriculture. The AFOLU community has, contrarily, a multi-gas approach (CO<sub>2</sub>, CH<sub>4</sub>,

719 N<sub>2</sub>O) and includes emissions from both forests and agriculture. For these reasons, estimates of  
720 the carbon community cannot be considered as AFOLU estimates, and certain confusion  
721 appears in the IPCC's AR5 with an incorrect AFOLU labelling (Table 11.1, Fig S2 in the  
722 Supplement). There is great space for these two communities to cooperate but further  
723 dialogue is needed to promote closer and more coordinated action. Future steps might include  
724 the adoption of the *managed land* concept by the carbon community; and ways to include  
725 legacy emissions by the AFOLU community.

726

#### 727 4.2.2 Improving data quality

728 The quality of the reported AFOLU emissions can be assessed through the UNFCCC  
729 principles: completeness, comparability, consistency, accuracy and transparency, which can  
730 help navigate the improvements of national monitoring systems. From these principles, the  
731 reviewed datasets performed well in *consistency* (they applied similar methods and  
732 assumptions over time, with the exception of 'Hotspots' that did not include temporal data).  
733 *Transparency* was excellent for FAOSTAT with well elaborated and publicly available  
734 metadata linked to their offered data, while EDGAR performed poorly due to insufficient  
735 metadata. Improving transparency is an urgent call for future action. *Accuracy and*  
736 *uncertainty* are also urgent calls. Thus, in spite of their importance to fully understand the  
737 emission trends and dynamics, only Houghton and the 'Hotspots' provided uncertainties.  
738 FAO offered uncertainties as a percent value for each emission source. *Completeness and*  
739 *omissions* are also urgent tasks because all datasets are incomplete (Table 1) (i.e. missing  
740 pools, missing gases) and omissions affect all datasets. Complete emission reporting should  
741 consider the importance of:

- 742 • Forest soil CO<sub>2</sub> and N<sub>2</sub>O emissions (Werner et al., 2007) (i.e. N<sub>2</sub>O tropical forest soil  
743 emissions of 0.7 PgCO<sub>2</sub>e.yr<sup>-1</sup>).

- 744 • Emissions from CH<sub>4</sub> and N<sub>2</sub>O from drained peatland soils, and from wetlands over  
745 managed land (i.e. conservation).
- 746 • All forest fire types (i.e. temperate conifers and woodlands; understory fires over  
747 humid closed canopy forests (Alencar et al., 2006; Morton et al., 2013) (i.e. 85,500  
748 km<sup>2</sup>, 1999-2010 in southern Brazilian Amazon); fire emissions over peatland soils and  
749 peatland forests out of Asia (Román-Cuesta et al., 2011; Oliveras et al., 2014) (i.e. 4-8  
750 TgCO<sub>2</sub>e, 1982-1999, for the tropical high Andes from Venezuela to Bolivia)
- 751 • CO<sub>2</sub> emissions from other components of wood harvesting other than fuel and  
752 industrial roundwood (i.e. charcoal, residues).
- 753 • CO<sub>2</sub> emissions from tree biomass loss due to fragmentation (Numata et al., 2010; Pütz  
754 et al., 2014) (i.e 0.2 Pg C y<sup>-1</sup>)
- 755 • CO<sub>2</sub> due to decomposition and decay of forests under extreme events: hurricanes  
756 (Read and Lawrence, 2003; Negron-Juarez et al., 2010) (i.e the 2005 convective  
757 storm, the Amazon basin suffered from an estimated tree mortality of 542±121 million  
758 trees); intense droughts (Phillips et al., 2009, 2010; Brienen et al., 2015) (i.e. the 2005  
759 Amazonian drought resulted in 1.2-1.6 PgC emissions and the atmosphere has yet to  
760 see 13.9 PgCO<sub>2</sub> (3.8 PgC) of the Amazon necromass carbon produced since 1983);

761

762 Further suggestions on improving data gaps and knowledge for the AFOLU sector have been  
763 reported by Smith *et al.* (2014); Houghton *et al.* (2012); USEPA (2013) and Sist *et al.* (2015),  
764 with a focus on soil data and crop production systems, as well as an improved understanding  
765 of the mitigation potentials, costs and consequences of land use mitigation options.

766

## 767 **5. REFERENCES**

768 Abad-Viñas, R., Blujdea, V., Federici, S., Hiederer, R., Pilli, R., Grassi, G.: Analysis and

769 proposals for enhancing Monitoring, Reporting, and Verification of greenhouse gases from  
770 Land Use, Land Use Change and Forestry in the EU. Technical Report  
771 071201/2011/211111/CLIMA.A2. Joint Research Centre, Ispra, Italy, 2015. Available at:  
772 <http://publications.jrc.ec.europa.eu/repository/bitstream/JRC91414/lb-na-26813-en-n.pdf>  
773 Achard, F., Beuchle, R., Mayaux, P., Stibig, H. J., Bodart, C., Brink, A., Carboni, S., Desclée,  
774 B., Donnay, F., Eva, H. D., Lupi, A., Raši, R., Seliger, R. and Simonetti, D.: Determination  
775 of tropical deforestation rates and related carbon losses from 1990 to 2010. *Global Change*  
776 *Biology*, 20, 2540-2554, 2014.

777 Alencar, A., Nepstad, D., Vera-Diaz, MC.: Forest Understory Fire in the Brazilian Amazon in  
778 ENS and non-ENSO years: Area Burned and Committed Carbon Emissions. *Earth*  
779 *Interactions*, 10, 1-17, 2006.

780 Anderson, K.: The inconvenient truth of carbon offsets. *Nature News*, 484, 7, 2012.

781 Anderson, K.: Duality in climate science. *Nature Geoscience*, 8, 898–900, 2015.

782 Asbjornsen, H., Gallardo-Hernández, C., Velázquez-Rosas, N., García-Soriano, R.: Deep  
783 ground fires cause massive above- and below-ground biomass losses in tropical montane  
784 cloud forests in Oaxaca, Mexico. *Journal of Tropical Ecology*, 21, 427-434, 2005.

785 Baccini, A., Goetz, S.J., Walker, W.S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J.,  
786 Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., Houghton, R.A.: Estimated carbon  
787 dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature*  
788 *Climate Change*, 2, 182-185, 2012.

789 Balch, J., Brando, P., Nepstad, D. Coe, M., Silverio, D., Massad, T., Davidson, E., Lefebvre,  
790 P., Oliveira-Santos, C., Rocha, W., Cury, R., Parsons, A., Carvalho, K.: The susceptibility of  
791 Southeastern Amazon Forests to Fire: Insights from a Large Scale Burn Experiment.  
792 *BioScience*, 65, 893-905, 2015.



793 Barlow, J., Peres, C.: Fire-mediated dieback and compositional cascade in an Amazonian  
794 forest. *Philosophical Transactions of the Royal Society*, 363, 1787-1794, 2008.

795 Bellassen, V., Luysaert, S.: Managing forests in uncertain times. *Nature*, 506, 153-156, 2014.

796 Brando, PM., Nepstad, DC., Balch, JK., Bolker, B., Christman, MC., Coe, M., Putz, F.: Fire-  
797 induced tree mortality in a neotropical forest: the roles of bark traits, tree size, wood density,  
798 and fire behaviour. *Global Change Biology*, 18, 630-641, 2012.

799 Brando, PM., Balch, JK., Nepstad, DC., Morton, D., Putz, F., Coe, M., Silverio, D., Macedo,  
800 M., Davidson, E., Nobrega, C., Alencar, A., Soares-Filho, B.: Abrupt increases in  
801 Amazonian tree mortality due to drought-fire interactions. *Proceedings of the National*  
802 *Academy of Sciences*, 11, 6347-6352, 2014.

803 Brienen, R., Phillips, O., Feldspausch, T., Gloor, E., Lloyd, J., Lopez-Gonzalez, G.,  
804 Morteagudo-Mendoza, A., Malhi, Y., Lewis, S., Vasquez Martinez, R., Alexiades, M.,  
805 Alvarez, E., Alvarez-Loayzada, P., Zagt, R.: Long term decline of the Amazon carbon sink.  
806 *Nature*, 519, 344-361, 2015.

807 Canadell, J., Schulze, D. Global potential of biospheric carbon management for climate  
808 mitigation. *Nature Communications*, 5, 5282-5293, 2014.

809 Ciais, P., Dolman, J., Bombelli, A., Duren, A., Peregon, A., Rayner, P., Miller, C., Gobron,  
810 N., et al.: Current systematic carbon-cycle observations and the need for implementing a  
811 policy-relevant carbon observing system. *Biogeosciences*, 11, 3547-3602, 2014.

812 Cochrane, M., Alencar, A., Schulze, M., Souza, C., Nepstad, D., Lefebvre, P., Davidson, E.:  
813 Positive feedbacks in the fire dynamics of closed canopy tropical forests. *Science*, 284,  
814 1832-1835, 1999.

815 Crowther, T., Thomas, S., Maynard, D., Baldrian, P., Covey, K., Frey, S., van Diepen, L.,  
816 Bradford, M.: Biotic interactions mediate soil microbial feedbacks to climate change.  
817 *Proceedings of the National Academy of Science*, 112, 7033-7038, 2015.

818 EDGAR. The Emissions Database for Global Atmospheric Research (2012) Part III:  
819 Greenhouse gas emissions. [http://edgar.jrc.ec.europa.eu/docs/IEA\\_PARTIII.pdf](http://edgar.jrc.ec.europa.eu/docs/IEA_PARTIII.pdf)

820 Estrada, M., Lee, D., Murray, B., O'Sullivan, R., Penman, J., Streck, C.: Land Use in a Future  
821 Climate Agreement. # S-LMAQM-13-CA-1128 U.S. Department of State, 2014. Available  
822 at: <http://merid.org/land-use-in-ADP/>

823 FAO. The Agriculture sectors in the intended nationally determined contributions. Analysis.  
824 Environmental and natural resource management working paper 61. Rome, Italy. 2016.  
825 Available at: <http://www.fao.org/3/a-i5687e.pdf>

826 Federici, S., Tubiello, F., Salvatore, M., Jacobs, H., Schmidhuber, J.: New estimates of CO<sub>2</sub>  
827 forest emissions and removals: 1990-2015. *Forest Ecol.Manag.*, 3, 89-98, 2015.

828 Federici, S., Grassi, G., Harris, N., Lee, D., Neeff, T., Penman, J., Sanz-Sanchez, M., Wolosin  
829 M.: GHG fluxes from forests: an assessment of national reporting and independent science  
830 in the context of the Paris Agreement. Working Paper. UCLA, San Francisco, 2016.  
831 Available at: [http://www.climateandlandusealliance.org/wp-content/uploads/2016/06/](http://www.climateandlandusealliance.org/wp-content/uploads/2016/06/GHG_Fluxes_From_Forests_Working_Paper.pdf)  
832 [GHG\\_Fluxes\\_From\\_Forests\\_Working\\_Paper.pdf](http://www.climateandlandusealliance.org/wp-content/uploads/2016/06/GHG_Fluxes_From_Forests_Working_Paper.pdf)

833 Friedlingstein, P., Andrew, R., Rogelj, J., Peters G., Canadell J., Knutti, R., Luderer, G.,  
834 Raupach, M., Schaeffer, M., van Vuuren, D., Le Quéré, C.: Persistent growth of CO<sub>2</sub>  
835 emissions and implications for reaching climate targets. *Nature geoscience*, 7, 709-715,  
836 2014.

837 Forest Resources Assessment (FRA) (2005) <http://www.fao.org/forestry/fra/fra2005/en/>

838 Forest Resources Assessment (FRA) (2010). <http://www.fao.org/forestry/fra/fra2010/en/>

839 Frank, D., Reichstein, M., Bahn, M., Thonicke, K., Frank. D., Mahecha, M., Smith, P., van  
840 der Velde, M. et al. Effects of climate extremes on the terrestrial carbon cycle: concepts,  
841 processes and potential future impacts. *Global Change Biology*, 21, 2861-2880, 2015.

842 Francey, R., Trudinger, C., Van der Schoot, M., Krummel, P., Steele, L., Langenfelds, L.:  
843 Differences between trends in atmospheric CO<sub>2</sub> and the reported trends in anthropogenic  
844 CO<sub>2</sub> emissions. *Tellus*, 62B, 316-328, 2010.

845 Francey, R., Trudinger, C., Van der Schoot, M., Law, M., Krummel, P., Langenfelds, R.,  
846 Steele, P., Allison, C., Stavert, A., Andres, R., Rödenbeck, C.: Atmospheric verification of  
847 anthropogenic CO<sub>2</sub> emission trends. *Nature Climate Change*, 3, 520-525, 2013a.

848 Francey, R., Trudinger, C., Van der Schoot, M., Law, M., Krummel, P., Langenfelds, R.,  
849 Steele, P., Allison, C., Stavert, A., Andres, R., Rödenbeck, C.: Reply to Anthropogenic CO<sub>2</sub>  
850 emissions. *Nature Climate Change*, 3, 603-604, 2013b.

851 Giglio, L., Randerson, J., van der Werf, G., Kasibhatla, P., Collatz, G., Morton D., DeFries,  
852 R.: Assessing variability and long-term trends in burned area by merging multiple satellite  
853 fire products. *Biogeosciences*, 7, 1171–1186, 2010.

854 Grace, J., Mitchard, E., Gloor, E.: Perturbations in the carbon budget of the tropics. *Global*  
855 *Change Biology*, 20, 3238-3255, 2014.

856 Grassi, G., Dentener, F.: Quantifying the contribution of the Land Use Sector to the Paris  
857 Climate Agreement. The LULUCF sector within the Intended Nationally Determined  
858 Contributions. EUR 27561. JRC Science for Policy Report. Ispra, Italy, 2015. Available at:  
859 <http://publications.jrc.ec.europa.eu/repository/bitstream/JRC98451/jrc%20lulucf->  
860 [indc%20report.pdf](http://publications.jrc.ec.europa.eu/repository/bitstream/JRC98451/jrc%20lulucf-indc%20report.pdf)

861 Grassi, G., Monni, S., Federici, S., Achard, F., Mollicone, D.: Applying the conservativeness  
862 principle to REDD to deal with uncertainties of the estimates. *Environmental Research*  
863 *Letters*, 3, 035005, 2008.

864 Hansen, M., Stehman, S., Potapov, P.: Quantification of global gross forest cover loss.  
865 *Proceedings of the National Academy of Sciences*, 107, 8650-8655, 2010.

866 Harris, N., Brown, S., Hagen, S., Saatchi, S., Petrova, S., Salas, W., Hansen, M., Potapov, P.,  
867 Lotsch, A.: Baseline Map of Carbon Emissions from Deforestation in Tropical Regions.  
868 Science, 336, 1576-1578, 2012.

869 Herrero, M., Havlik, P., Valin, H., Notenbaert, A., Rufino, M., Thornton, P., Blümmel, M.,  
870 Weiss, F., Grace, D., Obesteiner, M.: Biomass use, production, feed efficiencies, and  
871 greenhouse gas emissions from global livestock systems. Proc. Natl. Acad. Sci., 110, 20888-  
872 20893, 2013.

873 Herold, M., Roman-Cuesta, RM., Heymell, V., Hirata, Y., Van Laake, P., Asner, G., Souza,  
874 C., Avitabile, V., MacDicken, K.: A review of methods to measure and monitor historical  
875 carbon emissions from forest degradation. FAO. Unasylva, 238, 16-24, 2011a. Available at:  
876 <http://www.fao.org/docrep/015/i2560e/i2560e04.pdf>

877 Herold, M., Roman-Cuesta, RM., Mollicone, D., Hirata, Y., Van Laake, P., Asner, G., Souza,  
878 C., Skutch, M., Avitabile, V., MacDicken, K.: Options for monitoring and estimating historic  
879 carbon emissions from forest degradation in the context of REDD+. Carbon Balance and  
880 Management, 6, 13-20, 2011b.

881 Holmgren, P. Can we trust country-level data from global forest assessments? Available at:  
882 [http://blog.cifor.org/34669/can-we-trust-country-level-data-from-global-forest-](http://blog.cifor.org/34669/can-we-trust-country-level-data-from-global-forest-assessments?fnl=en)  
883 [assessments?fnl=en](http://blog.cifor.org/34669/can-we-trust-country-level-data-from-global-forest-assessments?fnl=en) Last view September 2016.

884 Hooijer, A., Page, S., Canadell, J., Silvius, M., Kwadijk, J., Wosten, H., Jauhiainen, J.:  
885 Current and future CO<sub>2</sub> emissions from drained peatlands in Southeast Asia.  
886 Biogeosciences, 7, 1505-1514, 2010.

887 Houghton, RA.: The annual net flux of carbon to the atmosphere from changes in land use  
888 1850-1990. Tellus B, 51, 298-313, 1999.

889 Houghton, RA.: How well do we know the flux of CO<sub>2</sub> from land-use change? Tellus B, 62,  
890 337-351, 2010.

891 Houghton, RA.: Carbon emissions and the drivers of deforestation and forest degradation in  
892 the tropics. *Current Opinion in Environmental Sustainability*, 4, 597-603, 2012.

893 Houghton, RA., Hackler, JL.: Carbon Flux to the Atmosphere from Land-Use Changes: 1850  
894 to 1990. ORNL/CDIAC-131, NDP-050/R1. Carbon Dioxide Information Analysis Center,  
895 U.S. Department of Energy, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A,  
896 2001.

897 Houghton, RA., House, JL., Pongratz, J., van der Werf, G., DeFries, R., Hansen, M., Le  
898 Quere, C., Ramankutty, R.: Carbon emissions from land use and land-cover change.  
899 *Biogeosciences*, 9, 5125-5142, 2012.

900 IPCC. Intergovernmental Panel on Climate Change.: Good Practice Guidance for Land Use,  
901 Land Use and Forestry.. IPCC National Greenhouse Gas Inventory Programme. (ed. Penman  
902 J, Gytarsky M, Hiraishi T, Krug T, Kruger D, Ppatti R, Buendia L, Miwa K, Ngara T,  
903 Tanabe K, Wagner F) IGES. Kanagawa, Japan, 2003 [http://www.ipcc-](http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf_files/GPG_LULUCF_FULL.pdf)  
904 [nggip.iges.or.jp/public/gpglulucf/gpglulucf\\_files/GPG\\_LULUCF\\_](http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf_files/GPG_LULUCF_FULL.pdf)FULL.pdf

905 IPCC. Intergovernmental Panel on Climate Change.: AFOLU Guidelines for National  
906 Greenhouse gas Inventories. Vol. 4: Agriculture, Forestry and Other Land Use (eds  
907 Eggleston S, Buendia L, Miwa K, Ngara T, Tanabe K). IGES, Kanagawa, Japan. 2006.  
908 <http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.htm>

909 IPCC. Intergovernmental Panel on Climate Change.: Summary for Policymakers. In: Climate  
910 Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth  
911 Assessment Report of the Intergovernmental Panel on Climate Change (eds Edenhofer O,  
912 Pichs-Madruga R, Sokona Y, E, Farahani E, Kadner S, Seyboth K, Adler A, Baum I,  
913 Brunner S, Eickemeier P, Kriemann B, Savolainen J, Schlomer S, von Stechow C, Zwickel  
914 T, Minx JC). Cambridge University Press, Cambridge, United Kingdom and New York,  
915 NY, USA, 2014.

916 Laurance, W., Williamson, G.: Positive feedbacks among forest fragmentation, drought and  
917 climate change in the Amazon. *Conservation Biology*, **15**, 1529–1535, 2001.

918 Le Quéré, C., Peters, G.P., Andres, R. J., Andrew, R. M., Boden, T. A., Ciais, P.,  
919 Friedlingstein, P., Houghton, R.A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneeth,  
920 A., Arvanitis, A., Bakker, D. C.E., Bopp, L., Canadell, J. G., Chini, L. P., Doney, S. C.,  
921 Harper, A., Harris, I., House, J.I., Jain, A. K., Jones, S. D., Kato, E., Keeling, R. F., Klein  
922 Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Maignan, F., Omar, A., Ono, T.,  
923 Park, G.-H., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S.,  
924 Schwinger, J., Segschneider, J., Stocker, B. D., Takahashi, T., Tilbrook, B., van Heuven, S.,  
925 Viovy, N., Wanninkhof, R., Wiltshire, A., and Zaehle, S. Global carbon budget 2013, *Earth  
926 System Science Data*, **6**, 235-263, 2014.

927 Li, C., Salas, W., DeAngelo, B., Rose, S.: DNDC9.5 in EPA (2013) Global Mitigation of non-  
928 CO<sub>2</sub> Greenhouse Gases: 2010-2030. EPA Technical Report-430-R-13-011, US, 2013.  
929 Country data available at: [http://www.epa.gov/climatechange/EPAactivities/economics/  
930 nonco2projections.html](http://www.epa.gov/climatechange/EPAactivities/economics/nonco2projections.html)

931 Montzka, S.A., Dlugokencky, E.J., Butler, J.H.: Non-CO<sub>2</sub> greenhouse gases and climate  
932 change. *Nature*, **476**, 43-51, 2011.

933 Morton, D.C., Le Page, Y., DeFries, R., Collatz, G.J., Hurtt, G.C.: Understory fire frequency  
934 and the fate of burned forests in southern Amazonia. *Philosophical Transactions of the  
935 Royal Society B*, **368**, 20120163, 2013.

936 Negron-Juarez, R.I., Chambers, J., Guimaraes, G., Zeng, H., Raupp, C., Marra, D., Ribeiro,  
937 G., Saatchi, S., Nelson, B., Higuchi, N.: Widespread Amazon forest tree mortality from a  
938 single cross-basin squall line event. *Geophysical Research Letters*, **37**, L16701, 2010.

939 Numata, I., Cochrane, M., Roberts, D., Soares, J., Souza, C., Sales, M.: Biomass collapse and  
940 carbon emissions from forest fragmentation in the Brazilian Amazon. *Journal of*  
941 *Geophysical Research*, 115, G03027, 2010.

942 Ogle, S. et al. in EPA (2013) *Global Mitigation of non-CO2 Greenhouse Gases: 2010-2030*.  
943 EPA Technical Report-430-R-13-011, 2013.(data available upon request) Country data  
944 available at:[http://www.epa.gov/climatechange/EPAactivities/economics/nonco2projections.](http://www.epa.gov/climatechange/EPAactivities/economics/nonco2projections.html)  
945 [html](http://www.epa.gov/climatechange/EPAactivities/economics/nonco2projections.html)

946 Oliveras, I., Malhi, Y., Salinas, N., Huaman, V., Urquiaga-Flores, E., Kala-Mamani, J.,  
947 Quintano-Loaiza, JA., Cuba-Torres, I., Lizarraga-Morales, N., Roman-Cuesta, RM.:  
948 Changes in forest structure and composition after fire in tropical montane cloud forests near  
949 the Andean treeline. *Plant Ecology and Diversity*,7, 329-340, 2013.

950 Oliveras, I. Anderson, D., Malhi, Y.: Application of remote sensing to understanding fire  
951 regimes and biomass burning emissions of the tropical Andes. *Global Biogeochemical Cy*  
952 *cles*, 28, 480-496, 2014.

953 Pelletier, J., Busch, J., Potvin, C.: Addressing uncertainty upstream or downstream of  
954 accounting for emissions reductions from deforestation and forest degradation. *Climatic*  
955 *Change*, 130, 635-648, 2015.

956 Peters, G., Davis, J., Andrew, R.: A synthesis of carbon in international trade. *Biogeosciences*,  
957 9, 3247-3276, 2012.

958 Phillips, O., Aragao, L., Lewis, S., et al.: Drought Sensitivity of the Amazon Rainforest.  
959 *Science*, 323, 1344-1347, 2009.

960 Phillips, O., van der Heijden, G., Lewis, S., et al.: Drought-mortality relationships for tropical  
961 forests. *New Phitologist*, 187, 631-646, 2010.

962 Phillips, O., Lewis, S.:Evaluating the tropical forest carbon sink. *Global Change Biology*, 20,  
963 2039-2041, 2014.

964 Poorter, L., Bongers, F., Aide et al.: Biomass resilience of Neotropical secondary forests.  
965 Nature, 530, 211-214, 2016.

966 Pütz, S., Groeneveld, J., Henle, K., Knogge, C., Martensen, A., Metz, M.: Long-term carbon  
967 loss in fragmented Neotropical forests. Nature communications, 5, 5037-5045, 2014.

968 Read, L., Lawrence, D.: Recovery of biomass following shifting cultivation in dry tropical  
969 forests of the Yucatan. Ecological Applications, 13, 85–97, 2003.

970 Richards, M., Bruun, T., Campbell, BM., Gregersen, L., Huyer, S., Kuntze, V., Madsen, T.,  
971 Oldvig, M., Vasileiou, I.: How countries plan to address agricultural adaptation and  
972 mitigation: An analysis of Intended Nationally Determined Contributions. CGIAR Research  
973 Program on Climate Change, Agriculture and Food Security (CCAFS), Copenhagen, 2015.

974 Richter, D., Houghton, RA.: Gross CO<sub>2</sub> fluxes from land-use change: implications for  
975 reducing global emissions and increasing sinks. Carbon management, 2, 41-47, 2011.

976 Roman-Cuesta, RM., Gracia, M., Retana, J.: Environmental and human factors influencing  
977 fire trends in Enso and non-Enso years in tropical Mexico. Ecological Applications, 13,  
978 1177–1192, 2003.

979 Roman-Cuesta, RM., Salinas, N., Asbjornsen, H. et al.: Implications of fires on carbon  
980 budgets in Andean cloud montane forest: The importance of peat soils and tree resprouting.  
981 Forest Ecology and Management, 261, 1987–1997, 2011

982 Roman-Cuesta, RM., Rufino, M., Herold, M., Butterbach-Bahl, K., Rosenstock, T., Ogle, S.,  
983 Li, S., Herrero, M., Poulter, B., Verchot, L., Martius, C., Stuiver, J., De Bruin, S.: Hotspots  
984 of tropical land use emissions: patterns, uncertainties, and leading emission sources for the  
985 period 2000-2005. Biogeosciences, 13, 4253–4269, 2016.

986 Romijn, E., Herold, M., Koistra, L., Murdiyarso, D., Verchot, L.: Assessing capacities of non-  
987 Annex I countries for national forest monitoring in the context of REDD+. Environmental  
988 Science Policy, 19–20, 33–48, 2012.



989 Romijn, E., Lantican, C., Herold, M., Lindquist, E., Ochieng, R., Wijaya, A., Murdiyarso, D.,  
990 Verchot, L.: Assessing change in national forest monitoring capacities of 99 tropical  
991 countries. *Forest Ecology and Management*, 352, 109–123, 2015.

992 Saatchi, S., Harris, N., Brown, S., et al.: Benchmark map of forest carbon stocks in tropical  
993 regions across three continents. *Proceedings of the National Academy of Sciences*, 108,  
994 9899-9904, 2012.

995 Simula, M.: Towards defining forest degradation: comparative analysis of existing definitions.  
996 *Forest Resources Assessment. Working Paper 154*. FAO. Rome, Italy, 2009. Available at:  
997 <ftp://ftp.fao.org/docrep/fao/012/k6217e/k6217e00.pdf>

998 Sist, P., Rutishauser, E., Peña-Claros, M., et al.: The Tropical Managed Forests Observatory:  
999 a research network addressing the future of logged forests. *Applied Vegetation Science*, 18,  
1000 171-174, 2015.

1001 Smith, P., Martino, D., Cai, Z., et al.: Greenhouse gas mitigation in agriculture. *Philosophical*  
1002 *Transactions of the Royal Society B: Biological Sciences*, 363, 789-813, 2008.

1003 Smith, P., Bustamante, M., Ahammad, H., et al.: Agriculture, Forestry and Other Land Use  
1004 (AFOLU). In: *Climate Change 2014: Mitigation of Climate Change. Contribution of*  
1005 *Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on*  
1006 *Climate Change* (eds Edenhofer O, Pichs-Madruga R, Sokona Y, E, Farahani E, Kadner S,  
1007 Seyboth K, Adler A, Baum I, Brunner S, Eickemeier P, Kriemann B, Savolainen J,  
1008 Schlomer S, von Stechow C, Zwickel T, Minx JC). Cambridge University Press, Cambridge,  
1009 United Kingdom and New York, NY, USA, 2014.

1010 Streck, C.: Forests and Land Use in the Paris Agreement. *Climate Focus*. 2015. Available at:  
1011 [http://www.climatefocus.com/sites/default/files/20151223%20Land%20Use%20and%20the](http://www.climatefocus.com/sites/default/files/20151223%20Land%20Use%20and%20the%20Paris%20Agreement%20FIN.pdf)  
1012 [%20Paris%20Agreement%20FIN.pdf](http://www.climatefocus.com/sites/default/files/20151223%20Land%20Use%20and%20the%20Paris%20Agreement%20FIN.pdf)

1013 Tian, H., Lu, C., Ciais, P., Michalak, A., Canadell, J., Saikawa, E., Huntzinger, D., Gurney,  
1014 K., Sitch, S., Zhang, B., Yang, J., Bousquet, P., Bruhwiler, L., Chen, G., Dlugokencky, E.,  
1015 Wofsy, S.: The terrestrial biosphere as a net source of greenhouse gases to the atmosphere,  
1016 *Nature*, 531, 225–228, 2016.

1017 Tubiello, F., Salvatore, M., Rossi, S., Ferrara, A., Fitton, N., Smith, P.: The FAOSTAT  
1018 database of greenhouse gas emissions from agriculture. *Environmental Research Letters*, 8,  
1019 015009-19, 2013.

1020 Tubiello, F., Salvatore, M., Córdor Golec, R., Ferrara, A., Rossi, S., Biancalani, R., Federici,  
1021 S., Jacobs, H., Flammini, A.: Agriculture, Forestry and Other Land Use Emissions by  
1022 Sources and Removals by Sinks 1990 – 2011 Analysis. Working Paper Series ESS/14-  
1023 02.FAO Statistical Division. Rome, Italy, 2014. Available at:  
1024 <http://www.fao.org/docrep/019/i3671e/i3671e.pdf>

1025 Tubiello, F., Salvatore, M., Ferrara, A., House, J., Federici, S., Rossi, S., Biancalani, R.,  
1026 Condor Golec, R., Jacobs, H., Flammini, A., Prospero, P., Cardenas-Galindo, P.,  
1027 Schmidhuber, J., Sanz Sanchez, MJ., Srivastava, N., Smith, P. The contribution of  
1028 Agriculture, Forestry and other Land Use Activities to Global Warming, 1990-2012. *Global*  
1029 *Change Biol.*, 21, 2655–2660, 2015.

1030 Turetsky, M., Benscoter, B., Page, S., Rein, G., Van der Werf, G., Watts, A.: Global  
1031 vulnerability of peatlands to fire and carbon loss. *Nature Geosciences*, 8, 11-14, 2015.

1032 Uhl, C., Kauffman, J.: Deforestation effects on fire susceptibility and the potential response of  
1033 the tree species to fire in the rainforest of the eastern Amazon. *Ecology*, 71, 437–449, 1990.

1034 USEPA. United States Environmental Protection Agency. Global Anthropogenic Non-CO<sub>2</sub>  
1035 Greenhouse Gas Emissions: 1990-2030. EPA 430-R-12-006. Washington, DC. 2012.  
1036 Available at: <http://www.epa.gov/climatechange/economics/international.html>

1037 USEPA. United States Environmental Protection Agency. Global Mitigation of non-CO2  
1038 Greenhouse Gases: 2010-2030. Technical Report-430-R-13-011, 2013. Available at:  
1039 [http://www.epa.gov/climate change/Downloads/EPAactivities/MAC\\_Report\\_2013.pdf](http://www.epa.gov/climate%20change/Downloads/EPAactivities/MAC_Report_2013.pdf)  
1040 Valentini, R., Arneeth, A., Bombelli, A. et al.: A full greenhouse gases budget of Africa:  
1041 synthesis, uncertainties and vulnerabilities. *Biogeosciences*, 11, 381-407, 2014.  
1042 Van der Werf, G., Randerson, J., Giglio, L., Collatz, G., Mu, M., Kasibhatla, P., Morton, D.,  
1043 DeFries, R., Jin, Y., van Leeuwen, T.: Global fire emissions and the contribution of  
1044 deforestation, savannah, forest, agricultural, and peat fires (1997–2009). *Atmospheric*  
1045 *Chemistry and Physics*, 10, 11707–11735, 2010.  
1046 Werner, C., Butterbach-Bahl, K., Haas, E., Hickler, T., Kiese, R.: A global inventory of N<sub>2</sub>O  
1047 emissions from tropical rainforest soils using a detailed biogeochemical model. *Global*  
1048 *biogeochemical Cycles*, 21, GB3010, 2007.

1049

## 1050 **6. CONTRIBUTIONS**


1051 RMRC, MR, MH designed the study. SO, BP provided data and ran quality controls of the  
1052 data. RMRC, MR, MH, KBB, TR, LV, CM, SR, RH, SO, BP,SdB discussed the results and  
1053 contributed to writing. SdB advised on statistical choices.

1054

## 1055 **7. ACKNOWLEDGEMENTS**

1056 This research was generously funded by the Standard Assessment of Mitigation Potential and  
1057 Livelihoods in Smallholder Systems (SAMPLES) Project as part of the CGIAR Research  
1058 Program Climate Change, Agriculture, and Food Security (CCAFS). Funding also came from  
1059 two European Union FP7 projects: GEOCarbon (283080) and Independent Monitoring of  
1060 GHG Emissions-N° CLIMA.A.2/ETU/2014/0008. Partial funds came through CIFOR from  
1061 the governments of Australia (Grant Agreement # 46167) and Norway (Grant Agreement


1062 #QZA-10/0468). In the memory of Changsheng Li.



1064




1065

1066 **Figure 1:** AFOLU (Agriculture, Forestry and Other Land Use) emissions estimates  
 1067 (PgCO<sub>2</sub>e.yr<sup>-1</sup>) for the period 2000-2005 for the tropics, for six datasets (Hotspots, FAO  
 1068 (FAOSTAT), EDGAR, EPA, Baccini and Houghton), disaggregated into FOLU (Forestry and  
 1069 Other Land Use) and Agricultural emissions. Uncertainties are only provided in the Hotspot  
 1070 dataset. EPA data do not include a FOLU sector. Houghton and Baccini's are FOLU, CO<sub>2</sub>-  
 1071 only, datasets and do not include agricultural emissions. Houghton offers net emissions while  
 1072 Baccini's data are gross emissions for deforestation + fire + wood harvesting (Baccini et al.,  
 1073 2012).



1075

1076 **Figure 2:** Tropical gross annual emissions (2000-2005) comparisons, for the leading emission  
 1077 sources in the AFOLU sector, for the Hotspots, FAOSTAT, EDGAR, Baccini, EPA and  
 1078 Houghton datasets). Bars indicate uncertainty estimates ( $1\sigma$  from mean). No uncertainty  
 1079 estimates are available for the other datasets. Houghton’s data are net land use emissions  
 1080 rather than deforestation and are offered for visual comparisons against Baccini’s gross  
 1081 deforestation estimate which includes gross deforestation + fire + wood harvesting.  
 1082 Uncertainties are only provided in the Hotspot dataset. EPA data do not include a FOLU  
 1083 sector. Forest degradation would be the sum of fire and harvesting emissions.




1085 **Figure 3:** Continental disaggregated emissions for the individual emission sources  
1086 ( $\text{PgCO}_2\text{e.yr}^{-1}$ ). Bars indicate uncertainty estimates ( $1\sigma$  from mean). No uncertainty estimates  
1087 are available for the other datasets.

1088


1089








1090

1091 **Figure 4:** Disaggregation of cropland soil emissions from drained peatlands for datasets with  
 1092 available data: FAOSTAT and Hotspots. Organic soils are excluded in EPA’s cropland  
 1093 emissions.




1094

1095 **Figure 5:** Contribution of the different AFOLU GHGs (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) for the different  
 1096 datasets. Bars indicate uncertainty estimates (1σ from mean). No uncertainty estimates are  
 1097 available for the other datasets.



1099 **Figure 6:** GHG emission contribution (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) of the leading AFOLU emission  
1100 sources. Bars indicate uncertainty estimates (1σ from mean). No uncertainty estimates are  
1101 available for the other datasets.

1102



1103

1104

1105


1106

1107

1108

1109

**Figure 7:** Country level agreement for (a) AFOLU and (b) forest emissions for the FAOSTAT, EDGAR and ‘Hotspots’ databases. The categories of agreement are percentiles of the standard deviations and represent data dispersion = data variability (i.e. High agreement = low data variability = low data dispersion  $\leq 25^{\text{th}}$  percentile, Moderate agreement:  $25^{\text{th}}$ - $50^{\text{th}}$  percentiles, Low agreement:  $50^{\text{th}}$ - $75^{\text{th}}$  percentiles, and Very Low agreement = high data variability  $\geq 75^{\text{th}}$  percentile)



1110

1111 **Figure 8:** Country level agreement for (a) Cropland and (b) livestock emissions for the  
 1112 FAOSTAT, EDGAR and ‘Hotspots’ databases. The categories of agreement are percentiles of  
 1113 the standard deviations and represent data dispersion = data variability (i.e. High agreement =  
 1114 low data variability = low data dispersion  $\leq 25^{\text{th}}$  percentile, Moderate agreement:  $25^{\text{th}}$ - $50^{\text{th}}$   
 1115 percentiles, Low agreement:  $50^{\text{th}}$ - $75^{\text{th}}$  percentiles, and Very Low agreement = high data  
 1116 variability  $\geq 75^{\text{th}}$  percentile)

1117

1118

	<b>Hotspots</b>	<b>FAOS TAT</b>	<b>EDGAR</b>	<b>Houghton</b>	<b>Baccini</b>	<b>EPA</b>	<b>AR5</b>
<b>Gross/Net emissions</b>	Gross	Gross	Gross	Net	Gross	Gross	Net
<b>Uncertainty<sup>a</sup></b>	√	No	No	No	No	No	√
<b>Transparency</b>	High	High	Low <sup>b</sup>	Low	Low	Intermediate	Low
<b>IPCC compliant</b>	√	√	√	Not fully <sup>c</sup>	Not fully <sup>d</sup>	√	Not fully <sup>e</sup>
<b>Forest carbon Pools</b>	AGB + BGB	AGB + BGB	AGB	AGB+BGB +Soil +CWD+Litter	AGB+BGB +Soil +CWD+Litter	Soil	AGB+BGB +Soil +CWD+Litter
<b>Gases</b>	CO <sub>2</sub> ,C H <sub>4</sub> , N <sub>2</sub> O	CO <sub>2</sub> ,C H <sub>4</sub> , N <sub>2</sub> O	CO <sub>2</sub> ,C H <sub>4</sub> , N <sub>2</sub> O	CO <sub>2</sub>	CO <sub>2</sub>	CO <sub>2</sub> ,CH <sub>4</sub> , N <sub>2</sub> O	CO <sub>2</sub> for forests. CO <sub>2</sub> ,CH <sub>4</sub> , N <sub>2</sub> O for agriculture and peatlands.
<b>Tier 1</b>	√	√	√			√	-
<b>Tier 2, 3</b>	√			√	√	√	-
<b>Spatial Disaggregation<sup>f</sup></b>	Pixel (0.5°)	Country	Country <sup>g</sup>	Region	Region	Country	Region
<b>Peatlands</b>	√	√	√	No	No	No	√

1120 **Table 1:** differences and similarities of the assessed AFOLU datasets.

1121 a Uncertainty at the level of disaggregation at which data are available to download.

1122 b Low means there is no metadata available, or metadata does not properly document the  
1123 processes followed to estimate the emissions.

1124 c EDGAR data on deforestation emissions does not follow IPCC guidelines.

1125 d The bookkeeping approach does not follow the concept of managed land, and does not  
1126 include the sink of forests remaining forests in managed land other than logged forests and  
1127 those regrowing after shifting cultivation.

1128 e Based on Houghton et al. (2012).

1129 f Available disaggregated data.

1130 g We selected data at the country scale to favour comparability with other datasets (i.e.  
1131 FAOSTAT) even though data are available at pixel level (0.1°).

1132



	Gross Tropical (PgCO <sub>2</sub> e.yr <sup>-1</sup> )						
(a)	2000-2005					2000-2007	
	Hotspots	FAOSTAT	EDGAR-JRC	Houghton	EPA	Baccini	AR5
<b>Agriculture</b>	1.9 (1.5-2.5)	2.5	2.1	-	2.0	-	
<b>FOLU</b>	6 (3.8-10)	5.9	5.9	5.4*	-	12.3**	8.2***
<b>AFOLU</b>	8 (5.5-12.2)	8.4	8	-	-	-	
	Net Global PgCO <sub>2</sub> e.yr <sup>-1</sup>						
(b)	2000			2010			2000/09
	FAOSTAT	EDGAR-JRC	Houghton	FAOSTAT	EDGAR-JRC	Houghton	AR5
<b>Agriculture</b>	5	5.5	-	5.2	5.8	-	5
<b>FOLU</b>	4.9	6.5	4.9	4.9	5.5	4.2	5
<b>AFOLU</b>	9.9	12	-	10.1	11.3	-	10

1133

1134 **Table 2:** Summary of (a) tropical gross emissions estimates for agriculture, FOLU (Forestry and Other Land Use) and AFOLU (Agriculture,  
1135 Forestry and Other Land Use) for all the datasets (Hotspots, FAOSTAT, EDGAR, EPA, Houghton) (2000-2005) and published data (Baccini et  
1136 al. (2012), AR5 (Smith et al., 2014)) (2000-2007), and of (b) net global estimates as reported by Tubiello et al., (2015). Houghton and EPA offer  
1137 FOLU and agricultural data only, respectively, and therefore estimates for AFOLU are not complete.

1138 \*Data exposed in Figure 11.2 in Chapter 11 Smith et al. (2014), it corresponds to a net FOLU estimate without agriculture.

1139 \*\* Baccini et al., (2012) reported gross estimates for the FOLU components.

1140 \*\*\* Baccini et al., (2012) estimates selected for the AR5 FOLU values in Figure 11.8, Chapter 11, WG-III.

1141

	Deforestation	Wood Harvesting	Fire	Enteric Fermentation	Manure management	Agricultural soils	Cropland over histosols	Rice	Others
<b>CO<sub>2</sub></b>	Hotspots <sup>1</sup> FAOSTAT <sup>2</sup> Houghton <sup>3</sup> Baccini <sup>1</sup>	Hotspots <sup>4</sup> FAOSTAT <sup>5</sup> Houghton <sup>4</sup> Baccini <sup>4</sup>	Hotspots <sup>6</sup> FAOSTAT <sup>7</sup> EDGAR <sup>8</sup> Houghton <sup>9</sup> Baccini <sup>9</sup>				Hotspots <sup>10</sup> FAOSTAT <sup>11</sup>		EDGAR <sup>12</sup>
<b>CH<sub>4</sub></b>			Hotspots <sup>13</sup> FAOSTAT <sup>14</sup> EDGAR <sup>15</sup>	Hotspots FAOSTAT EDGAR EPA	Hotspots FAOSTAT EDGAR EPA			Hotspots FAOSTAT EDGAR EPA	
<b>N<sub>2</sub>O</b>			Hotspots <sup>13</sup> FAOSTAT <sup>14</sup> EDGAR <sup>15</sup>		Hotspots FAOSTAT EDGAR EPA	Hotspots <sup>16,17</sup> FAOSTAT <sup>16,18</sup> EDGAR <sup>16,17,19</sup> EPA <sup>16,19</sup>	Hotspots FAOSTAT	Hotspots	
<b>dSOC</b>						Hotspots		Hotspots	

1142 **Table 3:** Characteristics of the emission sources used in this comparative assessment disaggregated by greenhouse gases for the period 2000-  
1143 2005, for the Hotspots, FAOSTAT, EDGAR, EPA, Houghton, and Baccini’s datasets (based in gross emissions from Baccini et al. (2012)).  
1144 Superindices specify differences among datasets and/or indicate the exact data included in our database comparisons. EPA offers only non-CO<sub>2</sub>  
1145 emissions for agriculture. Houghton offers only CO<sub>2</sub> FOLU emissions. Baccini’s gross emissions include deforestation, fire and wood harvesting  
1146 only. dSOC refers to changes in Soil Carbon stocks. Wood harvesting and fire are considered as forest degradation.

1147 <sup>1</sup> Gross deforestation.

1148 <sup>2</sup> Net deforestation. Forest fire emissions included in deforestation

1149 <sup>3</sup> Houghton net CO<sub>2</sub>-only estimates are not deforestation emissions, but land use and land use change fluxes including deforestation, forest  
1150 degradation, and cropland, abandoned land, and agricultural soil organic carbon (SOC).

1151 <sup>4</sup> Nationally reported fuel wood and industrial roundwood.

1152 <sup>5</sup> Nationally reported fuel wood, charcoal, fuel residues and industrial roundwood.

1153 <sup>6</sup> Long-term CO<sub>2</sub> emissions only (i.e savannas/agricultural fires excluded). Peat, forests and woodland fires are included (as defined by Van der  
1154 Werf et al., 2010). Deforestation fires excluded

1155 <sup>7</sup> CO<sub>2</sub> from the combustion of organic soils. Forest fire emissions excluded.

1156 <sup>8</sup> CO<sub>2</sub> Forest fires + wetland/peatland fires and decay (5A, and 5D classes).

1157 <sup>9</sup> Humid forest deforestation fires, and peatland fires + decay.

1158 <sup>10</sup> CO<sub>2</sub> emissions from organic soils. Tier 1 approach. EF=20 tC.ha<sup>-1</sup>.yr<sup>-1</sup> (IPCC 2006). Only for the six crop types reported by the agricultural  
1159 soils (maize, soya, sorghum, wheat, barley, millet). N<sub>2</sub>O emissions not included.

- 1160 <sup>11</sup> CO<sub>2</sub> emissions from organic soils. Tier 1 approach. EF=20 tC.ha<sup>-1</sup>.yr<sup>-1</sup> (IPCC 2006). N<sub>2</sub>O emissions not included.
- 1161 <sup>12</sup> CO<sub>2</sub> for fuelwood is part of the energy balance.
- 1162 <sup>13</sup> CH<sub>4</sub> and N<sub>2</sub>O emissions for peat, forests and woodland, savannahs and agriculture fires.
- 1163 <sup>14</sup> CH<sub>4</sub>, N<sub>2</sub>O emissions from fire in humid tropical forests and other forests, as well as CH<sub>4</sub>, N<sub>2</sub>O from the combustion of organic soils.
- 1164 <sup>15</sup> CH<sub>4</sub>, N<sub>2</sub>O for forest fires + wetland/peatland fires and decay (5A, and 5D classes).
- 1165 <sup>16</sup> Direct agricultural emissions only
- 1166 <sup>17</sup> Fertilizers, manure, crop residues
- 1167 <sup>18</sup> Synthetic fertilizers + Manure applied to soils + Crop residues + Manure applied to pastures.
- 1168 <sup>19</sup> Indirect emissions
- 1169

1170

	Hotspots	FAOSTAT	EDGAR	Houghton*	Baccini	EPA	AR5*
Deforestation							
Fire							
Wood-harvesting							
Livestock							
Cropland							
Paddy Rice							
Peatland							
Forest sinks							

1171 **Table 4:** Identification of the least reliable emission sources for each dataset considering  
 1172 disagreements among emission estimates due to biased/divergent/incomplete definitions and  
 1173 methods.