

Manganese in the West Atlantic Ocean in context of the first global ocean circulation model of manganese

Marco van Hulten^{1,6}, Rob Middag^{2,3,4}, Jean-Claude Dutay¹, Hein de Baar^{4,5}, Matthieu Roy-Barman¹, Marion Gehlen¹, Alessandro Tagliabue⁷, and Andreas Sterl⁶

¹Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL, CEA–Orme des Merisiers, 91191 Gif-sur-Yvette, France

²University of Otago, 364 Leith Walk, Dunedin, 9016, New Zealand

³University of California Santa Cruz (UCSC), 1156 High Street, Santa Cruz, CA 95064, USA

⁴Royal Netherlands Institute for Sea Research (NIOZ), Landsdiep 4, 1797 SZ 't Horntje, Texel, the Netherlands

⁵University of Groningen (RUG), Postbus 72, 9700 AB Groningen, the Netherlands

⁶Royal Netherlands Meteorological Institute (KNMI), Utrechtseweg 297, 3731 GA De Bilt, the Netherlands

⁷University of Liverpool, 4 Brownlow Street, Liverpool L69 3GP, UK

Correspondence to: M. M. P. van Hulten (mvhulten@lsce.ipsl.fr)*

Abstract. Dissolved manganese (Mn) is a biologically essential element. Moreover, its oxidised form is involved in removing itself and several other trace elements from ocean waters. Here we report the longest thus far 17 500 km length full-depth ocean section of dissolved Mn in the West Atlantic Ocean, comprising 1320 data values of high accuracy. The goal of this study is to combine these new observations with state-of-the-art modelling to give a first assessment of the main sources and redistribution of Mn throughout the ocean. To this end, we simulate the distribution of dissolved Mn using a global-scale circulation model. This first model includes simple parameterisations to account for the sources, processes and sinks of Mn in the ocean. Oxidation and (photo)reduction, aggregation, settling, as well as biological uptake and remineralisation by plankton, are included in the model. Our model provides, together with the observations, the following insights:

- The high surface concentrations of manganese are caused by the combination of photoreduction and sources to the upper ocean. The most important sources are sediments, dust, and, more locally, rivers.
- Observations and model simulations suggest that surface Mn in the Atlantic Ocean moves downwards into the southward flowing North Atlantic Deep Water (NADW), but because of strong removal rates there is no elevated concentration of Mn visible anymore in the NADW south of 40° N.
- The biological cycle is not well understood: when incorporating Mn in (and release from) plankton, similarly to phosphorus, Mn becomes extremely low in large parts of the surface of the Pacific Ocean. This is inconsistent with observed dissolved Mn concentrations in these Pacific surface waters.

*Postprints may be found at [arXiv:1606.07128](https://arxiv.org/abs/1606.07128).

- There is a mainly homogeneous background concentration of dissolved Mn of about 0.10 nM to 0.15 nM throughout most of the deep ocean. The model reproduces this by means of a threshold on particulate manganese oxides of 25 pM, suggesting that a minimal concentration of particulate Mn is needed before aggregation and removal become efficient.
- The observed sharp hydrothermal signals are produced by assuming both a high source and a strong removal of Mn near hydrothermal vents.

1 Introduction

Dissolved manganese (Mn_{diss}) is taken up by phytoplankton, because Mn is crucial for photosynthesis and other biological functions (Raven, 1990). Furthermore, its oxidised form (Mn_{ox}) plays an important role in the removal of several other trace metals from seawater (Yamagata and Iwashima, 1963). While in the open ocean manganese (Mn) exists in small concentrations, it is the twelfth most plentiful element in the Earth's crust (Wedepohl, 1995). In seawater, Mn occurs in many forms, among which the bioavailable dissolved form. After phytoplankton death, incorporated Mn sinks downwards together with the dead material, but most of the organic material is remineralised before reaching the sea floor (Sarmiento and Gruber, 2006), releasing incorporated Mn back to the water.

Another important mechanism of storing Mn in particles, besides biological incorporation, is the removal of dissolved Mn via larger colloids on which the Mn is oxidised to insoluble Mn(IV) (and possibly the other Mn(III) oxidation state), and the subsequent aggregation by particulate matter. Oxidation occurs everywhere in the ocean where oxygen is available. This process can be strongly accelerated by Mn(II)-oxidising microorganisms, primarily bacteria and fungi (Sunda and Huntsman, 1988, 1994; Tebo et al., 2005), but we do not understand the role of these organisms quite well (Nealson, 2006). The reverse process is the reduction of Mn oxides to bioavailable dissolved Mn(II), i.e. $Mn^{2+}(aq)$. The full oxidation/reduction (redox) equilibrium reaction, in its most simple form, is given by (Froelich et al., 1979):



Reduction is significantly faster under the influence of sunlight. It is hence referred to as *photoreduction* when irradiance is the major contributor. The relative rate of reduction compared to oxidation is important for Mn(II) availability. An overall net higher oxidation rate implies more particle formation, hence more Mn export.

Manganese enters the open ocean through lithogenic dust deposition (Baker et al., 2016) and lateral advection from reducing sediments (Homoky et al., 2016). Sediments along the relatively shallow (<1000 m depth) ocean margins tend to receive more organic deposition, hence by enhanced microbial decomposition are more anoxic and a stronger source of reduced Mn than deep sea sediments. Surface $[Mn_{diss}]$ is especially high in the central Atlantic Ocean and up to at least 30° N because of high dust input from the Sahara in combination with photoreduction (Landing and Bruland, 1987; Jickells, 1995; Guieu et al., 1994; Baker et al., 2006; De Jong et al., 2007; Wu et al., 2014). Similarly, $[Mn_{diss}]$ is high in the northern Indian Ocean (Vu and Sohrin, 2013). Dissolved Mn diffuses out of anoxic sediments, because sediment microorganisms reduce Mn_{ox} if there is no more oxygen (or nitrate) left (Li et al., 1969; Landing and Bruland, 1980; Sundby and Silverberg, 1985; Middag et al., 2012).

As long as there is oxygen in the sediment, the organic carbon is remineralised by using this oxygen as an electron acceptor (Froelich et al., 1979).

Rivers are another source of Mn to the ocean (Elderfield, 1976; Aguilar-Islas and Bruland, 2006). While much of the fluvial Mn is removed within the estuaries into the sediments by scavenging and aggregation, a large part may finally be transported to the ocean by diffusion from and resuspension of the sediments (Jeandel, 2016; Charette et al., 2016). Typically, the smaller sediment particles (0.5 μm to 4 μm) have a high Mn content, and, because of their small size, are able to reach the open ocean (Yeats et al., 1979; Sundby et al., 1981; Trefry and Presley, 1982). Manganese may also flux into the ocean by melting sea ice (Middag et al., 2011b). Finally, overwhelming evidence is found of manganese fluxing out of hydrothermal vents (Klinkhammer et al., 1977, 1985; Hydes et al., 1986; Klinkhammer et al., 2001; Middag et al., 2011a, b; German et al., 2016).

Downward fluxes of settling particles that have been collected in sediment traps show a strong correlation between lithogenic particles and authigenic Mn (Roy-Barman et al., 2005). Therefore, lithogenic particles are likely to play a significant role in the removal (oxidation, scavenging and aggregation) of Mn (Dutay et al., 2015). Here the ballast effect of lithogenic particles, which typically have a density of about twice of that of seawater, is likely playing a major role for rapid settling of agglomerates of biogenic and lithogenic particles including Mn-oxides coatings (Balistrieri et al., 1981). This is consistent with the fact that most suspended particles are small (less than 2 μm), but the larger aggregates are deemed to be the significant contributors to the vertical flux (McCave, 1975; Bishop and Fleisher, 1987). The complete process may be more complicated than described above (Boyle et al., 2005), e.g. because Mn binds to dissolved ligands such that more of it may stay in solution (Sander and Koschinsky, 2011; Madison et al., 2013; Luther et al., 2015). Manganese oxides are important scavengers of other trace metals like iron, cobalt, nickel and zinc (Yamagata and Iwashima, 1963; Murray, 1975; Means et al., 1978; Saito et al., 2004; Tonkin et al., 2004), as well as insoluble radionuclides such as thorium and protactinium (e.g Reid et al., 1979; Hayes et al., 2015; Jeandel et al., 2015). Therefore, Mn availability does not only directly impact primary production but may also play a role in removing other elements from the surface ocean. These elements include biologically essential trace metals as well as many more trace elements.

Published observational studies, like the excellent results of Wu et al. (2014), show low, relatively constant Mn_{diss} concentrations of around 0.15 nM away from the boundaries. Towards the surface there is a sharp increase of $[\text{Mn}_{\text{diss}}]$. Other notable elevations are near oceanic ridges. These elevations of $[\text{Mn}_{\text{diss}}]$ at mid-depths can be ascribed to hydrothermal activity near those regions. The hydrothermal plumes typically extend in the order of 1000 km (Middag et al., 2011a, b), or reach even up to 3000 km (Resing et al., 2015).

What makes the distribution of Mn_{diss} relatively homogeneous in the interior of the ocean? How can this be reconciled with localised features where $[\text{Mn}_{\text{diss}}]$ is very elevated compared to that stable “background concentration”? Until now only local model simulations of the Mn ocean cycle have been performed that focus on the processes most relevant for the respective regions. In other words, the different features of the Mn_{diss} distribution have not been brought together in a unified model.

The case of hydrothermal activity has been studied by Lavelle et al. (1992) who modelled Mn in the deep ocean near hydrothermal vents. They included four Mn tracers, namely, a dissolved form, small particles associated with bacteria, larger aggregate particles, and one in sediments as the model includes benthic fluxes. They found that “more than 80 % of the

hydrothermal Mn is deposited within several hundred kilometres of the ridge crest though dissolved Mn concentrations beyond that distance exceed background levels by many times”. This illustrates the high Mn input and removal from hydrothermal vents, as well as the already mentioned large plume extend.

5 The Mn_{diss} distribution in the North Pacific Ocean has been modelled by Johnson et al. (1996), in a 1-D vertical model neglecting horizontal transport. Their oxidation model depends on the oxygen concentration and the hydroxide activity. Their region of interest was on the upper and intermediate depth ocean, and their goal was to reproduce the $[Mn_{diss}]$ maximum in the Oxygen Minimum Zone (OMZ). Except for the North Pacific, the OMZ is also present in other basins, amongst which the northwestern Indian Ocean (Saager et al., 1989; Lewis and Luther, 2000). Johnson et al. (1996) found that the combination
10 of remineralisation rates and decreased oxidation in the OMZ explained the $[Mn_{diss}]$ subsurface maximum in their profiles, whereas a flux from the continental margin sediments appeared not responsible. In the euphotic zone Mn_{diss} was incorporated in phytoplankton, while remineralised Mn_{diss} in the aphotic (and disphotic) zone was lost by oxidation and scavenging.

While these modelling efforts are useful for their purposes, no studies exist in which the global ocean Mn_{diss} distribution is modelled. To arrive at a more integrated understanding of the behaviour of pelagic Mn, we include a Mn model in a global
15 ocean general circulation model. Specifically, we will test a simple mechanism that should be able to give insight into the apparently stable Mn_{diss} concentration, and its contrast to strong Mn sources. This is the first time that a global ocean model for manganese has been written and assessed. It is a basic model that should give a starting point for further studies. At the moment there is no mechanistic evidence for typical uptake-remineralisation processes as is the case for, e.g., iron. However, as there is of course uptake of Mn by phytoplankton, and sometimes it is even a limiting element (Middag et al., 2013), we
20 include a biological cycle of Mn in the model. While we perform a model simulation on a global scale, we give more attention to the Mn_{diss} distribution in the Atlantic Ocean. Specifically, two dense and high-accuracy datasets as part of the GEOTRACES programme were available at the time of this study, namely the here reported GA02 section in the West Atlantic and the GIPY5 at the Zero Meridian in the Southern Ocean, which will be presented in this paper. Furthermore, in those regions we can study important properties of the ocean geochemistry of manganese and the interaction with circulation, among which the Atlantic
25 overturning circulation and hydrothermal activity.

In this study our goal is to assess the fundamental processes that are the most important to accurately simulate $[Mn_{diss}]$, including the aforementioned properties of the dissolved Mn distribution. To this end, we will first introduce our Mn model with its processes, sources and sinks. We will show the results of a reference simulation, which will be compared with recent high-accuracy observations. Also several sensitivity simulations will be presented, studying the effects of the biological cycle,
30 the intense nature of hydrothermal vents and the strong removal of manganese from the seawater.

2 Methods

2.1 Observations in the West Atlantic Ocean

2.1.1 Sample collection

For the determination of tracer concentrations, notably dissolved aluminium and dissolved manganese, samples were collected along the GEOTRACES Atlantic Meridional GA02 transect of the Netherlands (Fig. 5). Sampling was done with an all-titanium ultraclean CTD sampling system for trace metals (De Baar et al., 2008) with novel PVDF samplers (Rijkenberg et al., 2015; 5 Middag et al., 2015b, a). Immediately upon recovery, the complete titanium frame with its 24 PVDF samplers was placed inside a clean room environment sampling from where the sub-samples for trace metal analysis were collected. The water was filtered from the PVDF samplers over a 0.2 μm filter cartridge (Sartobran-300, Sartorius) under pressure (1.5 atm) of (inline prefiltered) nitrogen gas. Sub-samples for dissolved metals were taken in cleaned (Middag et al., 2009, for cleaning procedure) LDPE sample bottles. All sample bottles were rinsed five times with the sample seawater. Seawater samples were acidified 10 with HCl to a concentration of 0.024 M HCl which results in a pH of 1.7 to 1.8 with Baseline® Hydrochloric Acid (Seastar Chemicals Inc.).

2.1.2 Analysis of dissolved Mn

Analyses of dissolved manganese were performed on shipboard with the method developed by Doi et al. (2004) with some slight modifications in the preparation and brands of the chemicals used. Furthermore, samples were buffered in-line with an 15 ammonium borate sample buffer to a pH of 8.5 ± 0.2 . This buffer was produced by dissolving 30.9 g of boric acid (Suprapure, Merck) in 1 L MQ water (Millipore Milli-Q) deionised water $R > 18.2 \text{ M}\Omega \text{ cm}^{-1}$ and adjusting the pH to 9.4 with ammonium hydroxide (Suprapure, Merck).

The buffered sample was pre-concentrated during 150 s on a Toyopearl AF-Chelate 650M (TosoHaas, Germany) column. Hereafter the column was rinsed for 60 s with MQ water to remove interfering salts. The Mn was subsequently eluted from 20 the column for 200 seconds with a solution of 0.1 M three times quartz distilled formic acid (reagent grade, Merck) containing 0.1 M hydrogen peroxide (Suprapure, Merck) and 12 mM ammonium hydroxide (Suprapure, Merck). The pH of this carrier solution was adjusted to 2.9 ± 0.05 . The eluate with the dissolved Mn passed a second column of immobilised 8-hydroxyquinoline (Landing et al., 1986) to remove interfering iron ions in the carrier solution (Doi et al., 2004). Hereafter the carrier mixed with 0.7 M ammonium hydroxide (Suprapure Merck) and a luminol solution. The latter luminol solution was made by diluting 25 600 μl luminol stock solution and 10 μl TETA (triethylenetetramine, Merck) in 1 dm^3 MQ. The luminol stock solution was made by diluting 270 mg luminol (3-aminophthalhydrazide, Aldrich) and 500 mg potassium carbonate in 15 ml MQ. The resulting mixture of carrier solution, ammonium hydroxide and luminol solution had a pH of 10.2 ± 0.05 and entered a 3 m length mixing coil placed in a water bath of 25 °C. Hereafter the chemiluminescence was detected with a Hamamatsu HC135 Photon counter. Concentrations of dissolved Mn were calculated in nanomole per litre (nM) from the photon emission peak 30 height of triplicate measurements.

| Mn _{diss} (nmol kg ⁻¹) | Consensus | Middag et al. (2015b) |
|---|-------------|------------------------------|
| SAFe S | 0.79 ± 0.06 | 0.82 ± 0.02 |
| SAFe D2 | 0.35 ± 0.05 | 0.33 ± 0.01 (<i>n</i> = 24) |
| GEOTRACES S | 1.46 ± 0.14 | 1.47 ± 0.03 (<i>n</i> = 10) |
| GEOTRACES D | 0.21 ± 0.03 | 0.18 ± 0.01 (<i>n</i> = 5) |

Table 1. Compliance with the international GEOTRACES reference samples program. Left column are the international consensus values. Right column are the values reported here as part of the GA02 dataset.

The system was calibrated using standard additions from a 5000 nM Mn stock solution (Fluka) to filtered acidified seawater of low Mn concentration that was collected in the sampling region. A five-point calibration line (0, 0.1, 0.2, 0.6 and 1.2 nM standard additions) and blank determination were made every day. The three lowest points (0, 0.1 and 0.2 nM) of the calibration line were measured in triplicate and the two highest points (0.6 and 1.2 nM) in duplicate in order to add more weight to the lower part of the calibration line. The blank was determined by measuring acidified MQ which was below the detection limit and subsequently no blank was subtracted. The limit of detection defined as three times the standard deviation of the lowest value observed was < 0.01 nM. The flow injection system was rinsed every day with a 0.5 M HCl solution.

An internal reference sample was measured in triplicate every day. This was a sub-sample of a 25 dm³ volume of filtered seawater that was taken at the beginning of Leg 1 (also used during Leg 2, i.e. Iceland to the equator) and Leg 3 (Punta Arenas, Chile, to the equator). The relative standard deviation (i.e. the precision) of this replicate analysis seawater sample that was analysed 40 times on different days in triplicate was 2.57 % (Leg 1 and 2) and 1.21 % for 17 analyses during Leg 3. The relative standard deviation on single days was on average 1.37 % and the absolute values were 0.45 and 0.61 nM for the first two legs and Leg 3, respectively.

As external comparison, the international reference samples collected on the GEOTRACES Intercalibration Cruise (www.geotraces.org) as well as from the SAFe cruise (Johnson et al., 2007) were analysed for Mn. At different locations, large volumes of seawater were sampled of which subsamples were analysed by various laboratories, resulting in different reference values. These labs include the Royal Netherlands Institute for Sea Research (NIOZ), who also analysed the GA02 transect data. An independent referee removed outliers and averaged the reference values, resulting in a consensus value, which is here considered as the true value. These consensus values are listed in Table 1, together with the values determined by NIOZ. The distributions of the measurements from NIOZ lie within the precision of the consensus values.

The vertical profiles of the BATS station for the Netherlands sampling at 13 June 2010 and the US sampling at 19 November 2011 are presented in Fig. 1. For both occupations the samples were analysed by ICP-MS by Rob Middag at UCSC. Moreover, Rob Middag had done shipboard FIA analyses during the 2010 cruise. There is excellent agreement between the three datasets in the deep waters. Small differences within the upper ~500 m between the Netherlands sampling and the US sampling are deemed to be real due to the seasonal variability over the more than one year time difference of sampling. Furthermore, these Mn profiles are also consistent with the profiles determined by Landing et al. (1995).

Besides the shipboard Flow Injection Analysis (FIA) analysis, samples were also analysed on shore, using mass spectrometer analysis. Figure 2 shows the correlation between the two methods of analysis for the determination of $[\text{Mn}_{\text{diss}}]$, shipboard and laboratory measurements. There is a very good agreement between the shipboard and mass spectrometer analyses, which strongly suggests a high observational accuracy. In this study we used the observational data obtained from the shipboard FIA for the comparison with the model simulations.

2.2 Model description

5 In order to simulate the three-dimensional (3-D) distribution of dissolved Mn, we use the general circulation model *Océan Parallélisé* (OPA) that is part of NEMO, a framework for ocean models (Madec and the NEMO team, 2016). We use the ORCA2-LIM configuration of NEMO. The spatial resolution is 2° by $2^\circ \cos(\phi)$ (where ϕ is the latitude) with an increased meridional resolution to 0.5° in the equatorial domain (Madec and Imbard, 1996). The model has 30 vertical layers, with an increased vertical thickness from 10 m at the surface to 500 m at 5000 m depth. Representation of the topography is based
 10 on the partial-step thickness (Barnier et al., 2006). Lateral mixing along isopycnal surfaces is performed both on tracers and momentum (Lengaigne et al., 2003). The parameterisation of Gent and McWilliams (1990) is applied from 10° poleward to represent the effects of non-resolved mesoscale eddies. Vertical mixing is modelled using the turbulent kinetic energy (TKE) scheme of Gaspar et al. (1990), as modified by Madec and the NEMO team (2016). The fluid dynamics used to drive our model is identical to that used in Aumont et al. (2015).

15 The model contains two tracers of Mn, referred to as dissolved (Mn_{diss}) and oxidised (Mn_{ox}) manganese. The dissolved phase is operationally defined. These tracers are driven by the equations set out in this section, as well as the velocity fields obtained from OPA. Instead of calculating the dynamical variables of the model (velocity and mixing), we run it off-line, using a climatology with a resolution of five days.

Figure 3 presents the conceptual scheme of our manganese model. The internal processes are biological uptake and rem-
 ineralisation (green arrows), reduction, oxidation, aggregation and burial (blue arrows in the figure), and are described in the following subsections. Manganese sources from rivers, the atmosphere, sediments and hydrothermal vents are presented as
 5 arrows at the top and bottom of the figure. These four Mn sources are presented in Fig. 4, and Table 2 lists the absolute contributions to the different basins by each of these sources, as well as the relative contribution of every source to the world ocean. The model parameters are summarised in Table 3. In the following subsections we will describe how the different sources and processes are included in the model.

2.2.1 Atmospheric source

Manganese is added to the pool of Mn_{diss} in the upper model layer, according to

$$10 \left. \frac{\partial \mathcal{D}}{\partial t} \right|_{\text{dust}} = \frac{\alpha \cdot f_{\text{Mn,dust}}}{m \cdot \Delta z_1} \cdot \Phi_{\text{dust}}, \quad (1)$$

where \mathcal{D} is the dissolved Mn concentration, α is the solubility of Mn in dust, $f_{\text{Mn,dust}}$ is the mass fraction of Mn in dust, m is the molar mass of Mn, and $\Delta z_1 = 10$ m is the upper model layer thickness. The lithogenic dust deposition flux, Φ_{dust} , is

| Basin | Dust | Rivers | Sediment | Hydrothermal |
|---------------------------------------|------|--------|----------|--------------|
| Atlantic Ocean | 2200 | 127 | 924 | 13917 |
| Pacific Ocean | 1183 | 94 | 1237 | 59846 |
| Indian Ocean | 1506 | 25 | 442 | 18653 |
| Southern Ocean | 14 | 0 | 206 | 7601 |
| Arctic Ocean | 23 | 19 | 463 | 2269 |
| Mediterranean Sea | 673 | 12 | 91 | 0 |
| total amount (Mmol yr ⁻¹) | 5598 | 277 | 3363 | 102286 |
| relative amount (%) | 5.0 | 0.2 | 3.0 | 91.7 |

Table 2. Absolute amount of effective annual input by an imposed Mn flux from sediments, rivers and hydrothermal vents into each basin (Mmol yr⁻¹). The Southern Ocean is defined as the ocean south of 58.7° S. The line “total amount” denotes how much world-wide Mn is added to the ocean due to a specific flux; “relative amount” is normalised to the total Mn input flux.

| Parameter | Symbol | Value used | Known range | Reference |
|--|------------------------------------|---------------------------------------|---|--|
| Mass fraction of Mn in dust | $f_{\text{Mn,dust}}$ | 880 ppm | 696 to 880 ppm | Wedepohl (1995); Mendez et al. (2010) |
| Dust Mn solubility | α | 40 % | 10 to 70 % | Baker et al. (2006) |
| Sediment source Mn/Fe ratio | $r_{\text{Mn:Fe, sed}}$ | 0.2 | uncertain | Bortleson and Lee (1974); Slomp et al. (1997) |
| River source Mn/Fe ratio | $r_{\text{Mn:Fe, riv}}$ | 0.214 | uncertain | Sarmiento and Gruber (2006, p. 2) |
| Hydrothermal Mn/ ³ He ratio | $r_{\text{Mn:}^3\text{He, hydro}}$ | 0.10×10^9 | uncertain | - |
| Settling speed of Mn _{ox} | w_{ox} | 1–10 m d ⁻¹ | 0.9–1.4 m d ⁻¹ | Roy-Barman (2009) |
| Oxidation rate constant | k_{ox} | $0.341 \times 10^{-3} \text{ h}^{-1}$ | uncertain | Bruland et al. (1994) |
| Photoreduction rate constant | $k_{\text{red, light}}$ | $98 \times 10^{-3} \text{ h}^{-1}$ | $50\text{--}150 \times 10^{-3} \text{ h}^{-1}$ | Sunda and Huntsman (1994) |
| Aphotic reduction rate | $k_{\text{red, dark}}$ | $1.70 \times 10^{-3} \text{ h}^{-1}$ | $0.98\text{--}14.3 \times 10^{-3} \text{ h}^{-1}$ | Bruland et al. (1994) |
| Aggregation threshold | \mathcal{X}_{thr} | 25 pM | hypothetical | - |
| Mn/P incorporation ratio | $R_{\text{Mn:P}}$ | 0.36×10^{-3} | 0.33×10^{-3} to 5×10^{-3} | Twining and Baines (2013) |

Table 3. Mn model parameters for the *Reference* simulation. The settling velocity w_{ox} is given by Eq. 11. In *LowHydro* the hydrothermal flux and the maximum settling speed are both reduced by a factor of 10 ($r_{\text{Mn:}^3\text{He, hydro}} = 0.01 \times 10^9$; and $w_{\text{ox}} \equiv 1 \text{ m d}^{-1}$ or, equivalently, $\mathcal{X}_{\text{thr}} \rightarrow \infty$). In the sensitivity simulation *NoThreshold* the aggregation threshold \mathcal{X}_{thr} is set to zero.

derived from the Interaction with Chemistry and Aerosols (INCA) model (Hauglustaine et al., 2004). Here we use a 12 month climatology of INCA’s output as a forcing.

15 The average mass fraction of Mn in the Earth's upper crust is 527 ppm (Wedepohl, 1995). However, the fraction measured in Saharan dust is 880 ppm (Mendez et al., 2010), consistent with Guieu et al. (1994) and Statham et al. (1998). Since most of the dust deposited on the Atlantic Ocean originates from the Sahara and our focus is the Atlantic Ocean, the value of 880 ppm is used for $f_{\text{Mn,dust}}$.

The solubility of Mn from dust is uncertain and relatively high compared to most other trace metals. Here $\alpha = 40\%$ of the Mn in dust is assumed to dissolve, largely consistent with the values reported by Guieu et al. (1994); Jickells (1995); Baker et al. (2006); De Jong et al. (2007); Buck et al. (2010). Several studies report even higher values ($> 50\%$), which are, however, mainly from anthropogenic or otherwise processed dust, while the lower reported values ($< 50\%$) are from natural dust, mainly of Saharan origin.

Figure 4a presents the average Mn dissolution flux of the 12 months climatology. Globally this is 5.6 Gmol yr^{-1} of Mn.

10 2.2.2 River source

The manganese river source is modelled analogously to iron, which is part of the biogeochemical model PISCES (Aumont et al., 2015). This means that our manganese influx is proportional to the total dissolved (organic and inorganic) carbon flux, just like for iron (Aumont et al., 2015). Hence, the modelled concentration change of Mn_{diss} caused by river input is given by

$$\frac{\partial \mathcal{D}}{\partial t} \Big|_{\text{rivers}} = r_{\text{Mn:Fe,river}} \cdot \frac{\partial [\text{Fe}_{\text{diss}}]}{\partial t} \Big|_{\text{rivers}}, \quad (2)$$

15 where $[\text{Fe}_{\text{diss}}]$ is the dissolved iron concentration, and $r_{\text{Mn:Fe,river}}$ is the effective manganese/iron flux ratio, here set to 0.214, based on river dissolved concentrations (Sarmiento and Gruber, 2006, p. 2). For comparison, the Mn/Fe flux ratio is much higher than the crustal ratio, which is around 0.02 (Wedepohl, 1995), but probably underestimated as we do not consider external sources of particulate manganese. The effective Mn_{diss} input into the ocean by rivers is presented in Fig. 4b. The global, effective river flux of Mn is $0.28 \text{ Gmol yr}^{-1}$.

20 2.2.3 Sediment source

The largest contribution of Mn to the upper ocean is dust deposition, but over large shelf and slope regions (e.g. polar oceans) the flux of Mn_{diss} from the sediment can be of the same order of magnitude as, or higher than, the dust deposition flux (e.g. Middag et al., 2013, for the Southern Ocean). The (redox) reactions in the sediment are not explicitly modelled, since the sediment is not part of our model domain. Therefore, Mn addition from the sediment is modelled as a prescribed source. Since we do not have global maps of Mn_{diss} sediment-seawater flux, we parameterise the flux based on existing parameterisations of nitrate and iron.

Middelburg et al. (1996) derived an empirical model for calculating the denitrification rate as a function of the seafloor depth:

$$\hat{C}_{\text{Fsed}} = -0.9543 + 0.7662 \cdot \ln(z_{\text{Fsed}}) - 0.235 \cdot \ln(z_{\text{Fsed}})^2, \quad (3)$$

where ζ_{Fsed} is the natural logarithm of the denitrification flux ($\mu\text{mol cm}^{-2} \text{d}^{-1}$ of carbon) and z_{Fsed} is a function of bathymetry. Aumont et al. (2015) used this model for their sediment source of dissolved Fe in the PISCES model. They used a high-resolution bathymetric map to account for the shallow shelves, and modulated the seafloor depth ($z \rightarrow z_{\text{Fsed}}$) as follows:

$$z_{\text{Fsed}} = \min\left(8, \left(\frac{z}{500 \text{ m}}\right)^{-1.5}\right). \quad (4)$$

- 5 We set the maximum iron flux to $1 \mu\text{mol m}^{-2} \text{d}^{-1}$, following Aumont and Bopp (2006). We assume a porewater ratio $r_{\text{Mn:Fe, sed}} = 0.2$, and follow the same method as Aumont et al. (2015). Our final sediment addition of Mn_{diss} into bottom water is given by:

$$\left.\frac{\partial \mathcal{D}}{\partial t}\right|_{\text{sediments}} = \frac{0.2 \mu\text{mol m}^{-2} \text{d}^{-1}}{\Delta z_{\text{sed}}} \cdot \min(1, 2e^{\zeta_{\text{Fsed}}}), \quad (5)$$

- where z_{sed} is the gridbox thickness of the bottom gridbox (just above the seafloor) in metre. With this prescribed source, the Mn flux is limited to a maximum of about $75 \mu\text{mol m}^{-2} \text{yr}^{-1}$. This can be seen in Fig. 4c: the higher-than- $70 \mu\text{mol m}^{-2} \text{yr}^{-1}$ regions on the shelves is especially notable as it falls in the upper (red) part of the colour scale. The global sediment flux of Mn is 3.4 Gmol yr^{-1} .

2.2.4 Hydrothermal source

- Hydrothermal vent Mn_{diss} fluxes are modelled proportional to that of ^3He in Ruth et al. (2000) and Dutay et al. (2004). This approach is shown to have worked for iron (Bowers et al., 1988; Douville et al., 2002; Tagliabue et al., 2010; Resing et al., 2015). The basic equation for the change of Mn_{diss} from hydrothermal vent influx is

$$\left.\frac{\partial \mathcal{D}}{\partial t}\right|_{\text{hydrothermal}} = r_{\text{Mn:}^3\text{He, hydro}} \cdot \frac{\partial [^3\text{He}]}{\partial t}, \quad (6)$$

- where $r_{\text{Mn:}^3\text{He, hydro}} = 0.1 \times 10^9$ is the ratio between the Mn_{diss} and ^3He effective inflow from hydrothermal vents into the model domain. Recent observational studies found $\text{Mn}_{\text{diss}}/^3\text{He}$ concentration ratios of 4.0×10^6 (Kawagucci et al., 2008) and 3.5×10^6 (Resing et al., 2015) at observational sites close to hydrothermal outflux regions. To satisfy this, we assume that a dissolved fraction of 4% is left when the hydrothermal plume reaches these two observational sites.

- This high ‘‘solubility fraction’’ means that the hydrothermal vents are a large source of Mn_{diss} compared to Fe (Tagliabue et al., 2010; Resing et al., 2015). The integrated Mn flux is 102 Gmol yr^{-1} . In our simulations we will show that we need to assume such a large flux to explain the observations. This choice also relates to the fast modelled removal rate of Mn near the depth of hydrothermal vents (explained later). The hydrothermal Mn_{diss} source is presented in Fig. 4d.

2.2.5 Redox processes

Reduction and oxidation of Mn within the water column is a combination of several processes. Here we assume that Mn_{ox} is subject to reduction with a constant rate, but significantly stimulated by sunlight (e.g. Sunda and Huntsman, 1994). This is taken into account in the model by using different k_{red} for the euphotic and aphotic zones of the ocean. However, there

are other processes playing a role that can locally be important. Those include the microbial enhancement of the rate of oxidation in regions where Mn_{diss} supply is high (Sunda and Huntsman, 1994; Tebo et al., 2005), and the dependence on the O_2 concentration and pH (Johnson et al., 1996; Rijkenberg et al., 2014, for Fe at GA02). However, at this stage we decide to not include a dependency on $[\text{O}_2]$ to the model. Hence, here we choose to model Mn following a pseudo-first-order reaction where k_{ox} and k_{red} are pseudo-first-order rate constants (conventional primes omitted, e.g. Stone and Morgan, 1990):

$$\left. \frac{\partial \mathcal{D}}{\partial t} \right|_{\text{redox}} = -k_{\text{ox}} \mathcal{D} + k_{\text{red}} \mathcal{X} \quad (7)$$

$$\left. \frac{\partial \mathcal{X}}{\partial t} \right|_{\text{redox}} = k_{\text{ox}} \mathcal{D} - k_{\text{red}} \mathcal{X}, \quad (8)$$

where \mathcal{X} is the particulate oxidised Mn concentration, and

$$k_{\text{red}} = \begin{cases} k_{\text{red,light}} & \text{in the euphotic zone} \\ k_{\text{red,dark}} & \text{elsewhere,} \end{cases} \quad (9)$$

where $k_{\text{red,light}}$ is the reduction rate in the euphotic zone and $k_{\text{red,dark}}$ in the aphotic zone. The euphotic zone is defined as the depths where the sunlight intensity is at least 1%. The value for $k_{\text{red,light}}$ is taken from Sunda and Huntsman (1994) and has an observational uncertainty of a factor of three. The $k_{\text{red,dark}}$ is much smaller and much more uncertain (varying a factor of fifteen in Sunda and Huntsman (1994)). As we have both dissolved and particulate Mn in our model, we chose to fit the $k_{\text{ox}}/k_{\text{red,light}}$ to $[\text{Mn}_{\text{ox}}]/[\text{Mn}_{\text{diss}}]$ from observations, resulting in a value within the observational range of Sunda and Huntsman (1994). At the time of this study only Bruland et al. (1994) was known to us as reporting accurate dissolved and particulate Mn for the same samples (at the VERTEX-IV station). From this k_{ox} was calculated, and as Bruland et al. (1994) sampled the deep ocean as well, $k_{\text{red,dark}}$ was derived from that study as well (Table 3).

For one simulation, we will introduce a threshold on the oxidation process of Mn_{diss} . In that case, oxidation only takes place when higher than a certain threshold value \mathcal{D}_{thr} ; in other words, k_{ox} is multiplied by $H(\mathcal{D} - \mathcal{D}_{\text{thr}})$. The Heaviside step function $H(x)$ equals zero where $x \leq 0$ and one where $x > 0$. Based on observations, we have estimated $\mathcal{D}_{\text{thr}} = 0.125$ nM, corresponding to the observed deep ocean background value away from the influence of hydrothermal sources.²

2.2.6 Settling and burial

Manganese oxides settle, resulting in a concentration change according to

$$\left. \frac{\partial \mathcal{X}}{\partial t} \right|_{\text{settling}} = -w_{\text{ox}} \cdot \frac{\partial \mathcal{X}}{\partial z}, \quad (10)$$

where w_{ox} is the settling velocity, set to a constant 1 m d^{-1} as long as $[\text{Mn}_{\text{ox}}] \leq 25$ pM. If $[\text{Mn}_{\text{ox}}] > 25$ pM, the settling velocity is not a constant any more but a function of depth. Still, in the mixed layer $w_{\text{ox}} = 1 \text{ m d}^{-1}$, but if $[\text{Mn}_{\text{ox}}] > 25$ pM the sinking

²The value can be reproduced by first sorting the Mn_{diss} concentrations. Then we cut of the third quartile to remove the high values (above 0.42 nM), and chose a low value for a typical background concentration (0.13 nM is the first quartile). The reason for not choosing simply the minimum value is that values close to that can be lower than the ‘background value’ because of local removal processes.

speed increases linearly such that it reaches a value of 10 m d^{-1} at 2.5 km depth. The manganese oxide is buried when arriving
 30 at the ocean floor, which means that it is removed from the model domain.

The rationale for increasing the settling velocity above a certain threshold is that if Mn_{ox} is large enough, dense aggregates
 form that have faster sinking rates. This is not unlike increasing velocities of detritus in some models (Kriest and Oschlies,
 2011; Aumont et al., 2015). Notably mineral particles (sand, clay, carbonate) of high density in the order of 2–3 times that
 of seawater are responsible for this (McCave, 1975). Particulate organic carbon may play an important role as well (Passow
 5 and De La Rocha, 2006). In this way, $[\text{Mn}_{\text{diss}}]$ does not go to zero while still providing a deep ocean sink. As long as this
 critical particulate Mn concentration of 25 pM is not reached, aggregation of small Mn_{ox} particles does not yet occur. The
 choice of the critical concentration of 25 pM, also referred to as the *aggregation threshold*, is derived from the redox rate
 constants in combination with a typical, low value of the observed dissolved Mn concentration (here chosen as 0.125 nM):
 $\mathcal{X}_{\text{thr}} = (k_{\text{ox}}/k_{\text{red, dark}}) \cdot 0.125 \text{ nM} = 25 \text{ pM}$. More precisely, settling of Mn_{ox} follows Eq. (10) with the settling velocity

$$10 \quad w_{\text{ox}}/(\text{m d}^{-1}) = 1 + 9 \cdot \max\left(0, \frac{z - \text{MLD}}{2.5 \text{ km}}\right) \cdot H(\mathcal{X} - \mathcal{X}_{\text{thr}}), \quad (11)$$

where z is the depth, MLD is the mixed layer depth, and \mathcal{X}_{thr} is the aggregation threshold set to 25 pM.

2.2.7 Biological cycle

Manganese is incorporated into phytoplankton during growth. To this end, we run the PISCES-v2 biogeochemical ocean model
 (Aumont et al., 2015) together with our manganese model. The biological processes where Mn is involved are modelled
 15 proportionally to the change in phosphate concentration, thus it is given by:

$$\frac{\partial \mathcal{D}}{\partial t} \Big|_{\text{biology}} = R_{\text{Mn:P}} \cdot \frac{\partial [\text{PO}_4^{3-}]}{\partial t} \Big|_{\text{biology}}. \quad (12)$$

Here $R_{\text{Mn:P}}$ is the extended Redfield ratio for Mn and $[\text{PO}_4^{3-}]$ is the phosphate concentration. The ratio is set to 0.36×10^{-3} ,
 an average value for the manganese to phosphorus ratio in phytoplankton (Twining and Baines, 2013). This is close to the value
 0.4 derived by Middag et al. (2013) from dissolved distributions.

20 There are four types of plankton in PISCES: nanophytoplankton, diatoms, microzooplankton and mesozooplankton (pre-
 sented in Fig. 3 as the box “living”). There are two detrital pools, namely small particles settling with 2 m d^{-1} and large
 particles settling in our model configuration with 50 m d^{-1} (box “detrital”). Manganese is incorporated in the four living pools
 according to the PISCES equations for phosphorus. Through other processes they become part of detrital material or particulate
 organic matter. These pools are, however, not explicitly followed. Only the biological sources and sinks of Mn_{diss} are modelled,
 entailing the conversion of Mn_{diss} to the two phytoplankton pools, and from all particle pools to Mn_{diss} .

2.3 Simulations

The *Reference* simulation was spun up for 600 yr to reach a steady state. From year 100 onwards two sensitivity simulations
 5 were forked off to run in parallel with *Reference* for another 500 yr. These simulations are variations of the *Reference* simu-
 lation that uses the parameters listed in Table 3. Table 4 lists the simulations and their key parameters.

| Simulation name | $r_{\text{Mn:}^3\text{He,hydro}}$ | $w_{\text{ox}}/(\text{m d}^{-1})$ | $\mathcal{X}_{\text{thr}}/\text{pM}$ | $\mathcal{D}_{\text{thr}}/\text{nM}$ | $R_{\text{Mn:P}}$ |
|----------------------|--------------------------------------|-----------------------------------|--------------------------------------|--------------------------------------|-----------------------|
| <i>Reference</i> | 0.10×10^9 | 1–10 | 25 | 0 | 0.36×10^{-3} |
| <i>NoBio</i> | 0.10×10^9 | 1–10 | 25 | 0 | 0 |
| <i>LowHydro</i> | 0.01×10^9 | 1 | – | 0 | 0.36×10^{-3} |
| <i>NoThreshold</i> | 0.10×10^9 | 1–10 | 0 | 0 | 0.36×10^{-3} |
| <i>OxidThreshold</i> | 0.10×10^9 | 1–10 | 0 | 0.125 | 0.36×10^{-3} |

Table 4. List of simulations with the parameters changed compared to the reference simulation in boldface.

Our second simulation is without a biological cycle of Mn (*NoBio*). In this simulation Eq. 12 is removed from the model, or, equivalently, $R_{\text{Mn:P}}$ is set to zero. This simulation should illustrate the consequences of the (lack of) biological incorporation and subsequent remineralisation of Mn.

10 The goal of *LowHydro* is to investigate whether the combination of the high hydrothermal input of Mn_{diss} , and strong aggregation (modelled as a high settling velocity), is needed to get an accurate distribution of Mn_{diss} , i.e. one where its concentration compares well with the observations. Specifically, we want to explain the sharp observed Mn plumes. To this end, we first decreased the settling velocity to a constant 1 m d^{-1} (or, equivalently, $\mathcal{X}_{\text{thr}} \rightarrow \infty$), which, as expected, resulted in a wide spreading of Mn_{diss} and a too high $[\text{Mn}_{\text{diss}}]$ almost everywhere in the deep ocean (not presented). Decreasing only hydrothermal
5 input would trivially result in a proportionally smaller $[\text{Mn}_{\text{diss}}]$ near the vents. For this reason *LowHydro* contains two changes compared to the *Reference* simulation: a tenfold decrease in both the hydrothermal flux and the maximum settling velocity.

In our fourth simulation, *NoThreshold*, we want to see if an aggregation threshold is needed for an accurate $[\text{Mn}_{\text{diss}}]$ simulation. The threshold is removed by setting \mathcal{X}_{thr} in Eq. 11 to zero.

The fifth and final simulation, *OxidThreshold* is one that has a threshold in the oxidation process instead of the particle
10 sinking, setting effectively a minimum concentration of dissolved Mn.

2.4 Model validation

In this study we will mainly use data from the GEOTRACES programme, but for a worldwide global ocean comparison one has to rely also on data that were collected in the era before the reference samples of SAFe and GEOTRACES were available. Table 5 lists the datasets from GEOTRACES expeditions, as well as several other datasets; Fig. 5 shows the coordinates of the
15 stations.

Most of these GEOTRACES datasets are part of the publicly available GEOTRACES Intermediate Data Product 2014 (IDP) (Mawji et al., 2015), except for GP16 (Resing et al., 2015) that will be released as the GEOTRACES Intermediate Data Product 2017.

20 These observations are used for a visual global ocean data-model comparison. Details on the statistical and visual model–data comparison are presented in Appendix A. Of these observations, only the West Atlantic Ocean data (the GEOTRACES GA02 transect, consisting of 64 PE 319, PE 321 and JC 057; 1440 points) and the Zero-Meridian Southern Ocean data (GIPY5; 570

| Transect | Year | Expedition | Ocean basin | Citation | # |
|--|-----------|-----------------------------|--------------------------|--------------------------------------|------|
| GEOTRACES transects | | | | | |
| GIPY11 | 2007 | ARK XXII/2 | Arctic Ocean | Middag et al. (2011b) | 773 |
| GIPY4 | 2008 | MD166 BONUS-GoodHope | Southern Ocean | Boye et al. (2012) | 233 |
| GIPY5 | 2008 | ANT XXIV/3 | Southern Ocean | | |
| | | | a) Zero Meridian | Middag et al. (2011a) | 468 |
| | | | b) Weddell Sea | Middag et al. (2013) | 176 |
| | | | c) Drake Passage | Middag et al. (2012) | 221 |
| GI04 | 2009/2010 | KH-09-5 | Indian Ocean | Vu and Sohrin (2013) | 233 |
| GA02 | 2010 | 64 PE 319 | Northwest Atlantic Ocean | <i>this study</i> | 384 |
| GA02 | 2010 | 64 PE 321 | Northwest Atlantic Ocean | <i>this study</i> | 504 |
| GA02 | 2011 | JC 057 | Southwest Atlantic Ocean | <i>this study</i> | 432 |
| GA03 | 2010 | US GT10 | North Atlantic | Wu et al. (2014) | 91 |
| GA03 | 2011 | US GT11 | North Atlantic | Wu et al. (2014) | 578 |
| GP16 | 2013 | US EPZT | South Pacific | Resing et al. (2015) | 874 |
| Other expeditions and datasets | | | | | |
| - | 1983 | VERTEX-IV | North Pacific Ocean | Landing and Bruland (1987) | 27 |
| - | 2006 | EUCFe (<i>Kilo Moana</i>) | Pacific Ocean | Slemons et al. (2010) | 349 |
| - | 2005/2006 | CLIVAR P16 | Pacific Ocean | Milne and Landing (unpublished data) | 174 |
| - | 2007 | CoFeMUG | South Atlantic | Noble et al. (2012) | 429 |
| Total number of dissolved Mn measurements: | | | | | 5742 |

Table 5. Observational $[Mn_{diss}]$ used for comparison with the model simulations. GEOTRACES datasets are indicated by GEOTRACES transect codes. Their accuracies are approved by GEOTRACES on the basis of results of reference samples and cross-over stations. At the time there were no reference samples available for the other datasets, or they were not used for Mn.

points) have been used for statistical comparison with the model. Moreover, only the data of the shipboard FIA have been used, which have five stations less than the on-shore mass spectrometry determinations. The focus of this study is the West Atlantic Ocean for several reasons. Firstly, recently measurements have been carried out in that region, resulting in a large consistent (one method) dataset. Other regions generally contain fewer measurements and are mainly based on different methods by different analysts. Secondly, the West Atlantic Ocean is of importance to the Atlantic meridional overturning circulation, and hence the deep ocean cycling of for example the major nutrients. Therefore the West Atlantic Ocean was chosen as a key site for the ~18000 km long GEOTRACES GA02 transect for the collection of data for dissolved Mn and a suite of other trace elements and isotopes. The GEOTRACES GIPY5 transect at the Zero Meridian in the Southern Ocean complements the GA02 transect up to Antarctica, giving a more complete picture of the ocean circulation. For these reasons in this study we focus on the GA02 and GIPY5 transects.

The particulate $[\text{Mn}_{\text{ox}}]$ measurements from Landing and Bruland (1987) were used to tune the redox model. The data from the CLIVAR P16 cruise are unpublished; sampling is from the surface up to 1000 m depths; its methods of analysis are described by Milne et al. (2010).

3 Results

3.1 Observations in the West Atlantic Ocean

Figure 6 shows the West Atlantic GEOTRACES transect at the top 500 m. The concentration of Mn_{diss} near the ocean surface is high, going up to 4 nM and beyond. The highest surface concentrations are observed in the regions of high dust deposition as previously observed for aluminium (Middag et al., 2015a) and consistent with dust as the main source of Mn to the surface open ocean. The Mn_{diss} distribution is mainly homogeneous in the intermediate and deep ocean, where dissolved Mn has a concentration of about 0.10 nM to 0.15 nM.

North of 40° N high concentrations of dissolved Mn reach deeper than the top couple of hundred metres apparently because of vertical mixing. As can be seen in Fig. 7, some of it seems to go all the way down to about 3000 m. However, other sources may be (partly) responsible as well, among which advection from diffusive sediments and hydrothermal vents. By plotting all deep waters at enhanced resolution of concentration (Fig. 7), the Mn maxima of hydrothermal plumes and the O_2 minimum zones are better discernible. Whereas Mn and Fe have a very similar redox chemistry in the oceans, not surprisingly the maxima of Mn and Fe (Rijkenberg et al., 2014) overlap very nicely (the red dashed and black solid ellipses in Fig. 7). Some of the Mn may be transported southward by North Atlantic Deep Water (NADW), but this cannot be deduced from this transect: very soon the $[\text{Mn}_{\text{diss}}]$ reaches the typical background concentration. Similarly, slightly elevated concentrations in the subsurface are also observed in the northward advecting Antarctic Intermediate Water (AAIW around 1000 m), but once again concentrations reach the typical background concentration. In the Northern Hemisphere, the subsurface waters underlying the high dust deposition region near the equator have relatively elevated Mn_{diss} concentrations down to 750–1000 m depth, implying some influence of the dust deposition and particle export on the subsurface Mn concentrations, but yet again, the typical background concentration is reached further down. Elevated local features are located at almost 2 km depth on the Zero Meridian at 50° S (Middag et al., 2011a, Fig. 2; also presented as dots from 52° S on our Fig. 9) and almost 3 km depth in the West Atlantic transect at, and just south of, the equator, and at the Denmark Strait overflow (Fig. 7). The elevated Mn at the Denmark Strait overflow for a small part coincides with elevated aluminium attributed to sediment resuspension (Van Hulst et al., 2014; Middag et al., 2015a). However, there is a mismatch between the highest Al concentrations observed around 45° N and the highest Mn in the northernmost part of the transect (Middag et al., 2015a). This implies the source of Mn is related to the DSOW rather than the sediment resuspension occurring while the DSOW advects into the North Atlantic Ocean. The elevated deep Mn at the other locations match with elevated Fe and is most likely of hydrothermal origin (Klunder et al., 2011; Gerringa et al., 2015). Finally, at low latitudes, in the very deep ocean, $[\text{Mn}_{\text{diss}}]$ decreases with depth, suggesting a slower circulation in that region or an additional/enforced export of Mn to the sediment.

3.2 Reference simulation

Figure 8 shows the modelled and measured dissolved Mn concentrations at four depths; observations as coloured dots (same scale). The dissolved Mn concentration is high in the surface of the Atlantic, Indian and Arctic oceans, mostly consistent with the observations (Fig. 8a). The model also reproduces the latitudinal gradient of $[\text{Mn}_{\text{diss}}]$ in the Atlantic and Indian oceans, reflecting dust deposition patterns (Fig. 4a).

Figure 9 shows the dissolved Mn concentrations at full depth in the Zero-Meridian Southern Ocean and the West Atlantic Ocean from the *Reference* simulation; observations as coloured dots. Lower concentrations in the deep ocean are reproduced by the model (Figs 8b–d and 9). Both the model and observations present a mainly homogeneous distribution of just over 0.1 nM, though there are a number of sub-0.1 nM measurements in the polar oceans. The Mn_{diss} concentrations are generally higher near the surface compared to the deep ocean, both in the observations and the model. This is caused by a combination of dust deposition and photoreduction. Also the penetration of Mn_{diss} from Mn-rich surface waters into the deep ocean at around 50° N is reproduced by the model, but this is scavenged quickly before traversing southward in the NADW, which is consistent with early studies (e.g. Bender et al., 1977). Finally, the *Reference* simulation reproduces the measurements near hydrothermal vents in the Atlantic and Indian oceans (Figs 8c and 9).

Still, at many places, the Mn_{diss} concentration is underestimated by the model, with the most notable exceptions of the Southern and Arctic oceans. The underestimation is especially pronounced at the surface of the Pacific Ocean (Fig. 8a). Furthermore, in the model the Mn-rich water from the Amazon does not reach the GA02 transect as opposed to the observations. This potentially explains the underestimated concentration of Mn_{diss} in that region. The simulation overestimates measured $[\text{Mn}_{\text{diss}}]$ in the Arctic Ocean, except for some coordinates like near the Gakkel Ridge (the dark red dot in Fig. 8c at 90° E) (Middag et al., 2011b). The Gakkel Ridge is not in our forcing field (Fig. 4d).

Figure 10 presents $[\text{Mn}_{\text{diss}}]$ of the GA03 west to east transect. Observations from Wu et al. (2014) are presented as coloured dots. Again, the general patterns are captured, but the concentration of Mn_{diss} very close to the Mid-Atlantic Ridge is underestimated, while above the ridge (at about 2–3 km depth) it is overestimated. In other words, the modelled hydrothermal Mn_{diss} signal is not as sharp as in the observations. It is difficult to improve this feature because of the low vertical model resolution at that depth.

To more objectively compare the different simulations to each other, we list several goodness-of-fit statistics in Table 6. They compare the model simulations to observational data from the GEOTRACES GA02 transect. The model–data $[\text{Mn}_{\text{diss}}]$ correlation coefficient has a value of 0.77 for the *Reference* simulation. The reliability index, the average factor by which model predictions differ from observations, shows that on average the model differs by a factor of 1.67 from observations (Stow et al., 2009, and Appendix A).

| Simulation | Correlation | RMS deviation (nM) | Reliability index | Mn (Gmol) | Percentage Mn _{ox} |
|----------------------|---------------------|---------------------|---------------------|-----------|-----------------------------|
| <i>Reference</i> | 0.77 (± 0.06) | 0.46 (± 0.06) | 1.67 (± 0.07) | 440 | 14.1 |
| <i>NoBio</i> | 0.78 | 0.45 | 1.66 | 409 | 14.0 |
| <i>LowHydro</i> | 0.63 | 0.61 | 2.77 | 1023 | 15.8 |
| <i>NoThreshold</i> | 0.77 | 0.46 | 1.88 | 373 | 13.7 |
| <i>OxidThreshold</i> | 0.72 | 0.46 | 1.72 | 405 | 12.6 |

Table 6. Statistical model–data comparison for $[\text{Mn}_{\text{diss}}]$ at the GEOTRACES West Atlantic GA02 transect. The significance errors in the entries of *Reference* are $\pm 2\sigma$ from Monte Carlo samplings (Appendix A). Bold face values denote significant worsening compared with *Reference*; deviations of other values from the *Reference* value are insignificant. The last two columns are the total modelled Mn in seawater, and the Mn oxides as a portion of the total.

3.3 Biological cycle disabled

As Mn plays an important role as a trace nutrient, it is likely that biology has an impact on $[\text{Mn}_{\text{diss}}]$ at the ocean surface. Therefore, a simulation has been performed where biological incorporation and remineralisation of Mn_{diss} was not included, henceforth named *NoBio*.

10 Figure 11 shows that across most of the surface ocean the biological cycle of Mn in *Reference* causes a notable decrease of $[\text{Mn}_{\text{diss}}]$ compared to *NoBio*, though there is an increase at around 20° N/S in the Pacific Ocean. The decrease is up to 100 % near the equator and up to 80 % at 40° S. In the Atlantic Ocean $[\text{Mn}_{\text{diss}}]$ has not changed much between the two model simulations, explaining the insignificant change in the model–data comparison (Table 6). However, deeper in the Pacific Ocean, $[\text{Mn}_{\text{diss}}]$ is higher in *Reference* compared to *NoBio*.

15 Figure 12 shows $[\text{Mn}_{\text{diss}}]$ of the model compared with the observations at two depths for the *NoBio* simulation. The changes between the simulations are only big in the Pacific Ocean (Fig. 12). Elsewhere biology does not appear to have a big effect on the Mn_{diss} concentration. *NoBio* generally compares better with the observations in the Pacific Ocean, especially with the US EPZT transect (GP16).

3.4 Hydrothermal flux and export reduced

20 We have shown that the *Reference* simulation gives a reasonably accurate distribution of the effects of hydrothermal vents and the background concentration in the deep ocean. This is achieved by setting a large Mn_{diss} flux from hydrothermal vents and a big maximum settling velocity at the depth of the vents to remove the hydrothermal Mn away from the source. The simulation presented in this section, *LowHydro*, is meant to test if the high flux is necessary for an accurate Mn_{diss} distribution.

25 Figure 13 shows the relative change in $[\text{Mn}_{\text{diss}}]$ at four depths when both the hydrothermal input and the maximum settling velocity are decreased by a factor of 10 (*LowHydro*). In the surface ocean (Fig. 13a) there are both moderate increases and decreases. However, there are very large increases in places with deep convection that connect with the deep ocean (over 500 % in the Weddell Sea). In the deep ocean below about 1 km depth, such a large increase is not limited to the Weddell Sea but

stretches over much of the Atlantic and Indian oceans (Fig. 13c and d). Exceptions are the locations near the oceanic ridges, where there is hydrothermal activity. At those locations $[\text{Mn}_{\text{diss}}]$ decreased by up to 80 %.

30 The $[\text{Mn}_{\text{diss}}]$ transects in Fig. 14 show that *LowHydro* performs worse than the *Reference* simulation. For instance, the high $[\text{Mn}_{\text{diss}}]$ with a clear hydrothermal origin at 54° S has disappeared, while at the same time the deep ocean filled up with Mn_{diss} resulting in a consistent overestimation of about a factor of five in most of the deep West Atlantic Ocean. In the South Pacific Ocean and the Indian Ocean the signature of hydrothermal input of Mn_{diss} is – though smaller and worse than in *Reference* – still clearly present, with $[\text{Mn}_{\text{diss}}]$ values near the ridges distinctly different from the “background” concentration (results not shown). Furthermore, the statistics of *LowHydro* compared to *Reference* unambiguously show that $[\text{Mn}_{\text{diss}}]$ worsened (Table 6). The *LowHydro* simulation is significantly worse in all three statistics: the gradients from hydrothermal vents have disappeared,
5 and $[\text{Mn}_{\text{diss}}]$ is much too high throughout the ocean compared to *Reference*.

3.5 Aggregation threshold disabled

The simulation *NoThreshold* does not impose the aggregation threshold, meaning that settling is unconstrained in this simulation. Figure 15 presents $[\text{Mn}_{\text{diss}}]$ from *NoThreshold*. In the intermediate and deep ocean south of 40° N the concentration of Mn_{diss} is generally more than 50 % smaller than that in *Reference* (Fig. 9).

10 The homogeneous, already low background concentration of Mn_{diss} is reduced to close to zero (at least in the deep South Atlantic Ocean), while the hydrothermal signals are still correctly represented in the *NoThreshold* simulation. This explains that neither the correlation coefficient nor the root-mean-square deviation of $[\text{Mn}_{\text{diss}}]$ in *NoThreshold* differ significantly from those of *Reference*. This means that away from ocean ridges the spatial variation of the modelled $[\text{Mn}_{\text{diss}}]$ is similar to that of the observations (Stow et al., 2009). However, the reliability index, the average factor by which model predictions differ from
15 observations, has changed from 1.67 to 2.77, a change that is significant by over 30 standard deviations (Table 6). Therefore, the *NoThreshold* simulation is much worse than the *Reference* simulation.

3.6 Oxidation threshold enabled

As a final simulation, we limited the oxidation, rather than the settling of the manganese oxides. The resulting dissolved Mn concentrations are similar to those of *Reference* that has an aggregation threshold. Figure 16 shows the $\text{Mn}_{\text{ox}}/\text{Mn}_{\text{diss}}$
20 concentration ratio of the two simulations as well as observations at (155° W, 28° N). The blue squares represent the ratios between the particulate and dissolved Mn measurements taken during the VERTEX-IV cruise (Bruland et al., 1994). The left panel shows that this ratio lies near 0.005 in the upper 100 m, and the right panel shows the deep ocean that has higher values. The green line is the $\text{Mn}_{\text{ox}}/\text{Mn}_{\text{diss}}$ concentration ratio from the *Reference* simulation, and the red, dashed line is the *OxidThreshold* simulation. For the *Reference* simulation, the modelled \mathcal{X}/\mathcal{D} generally lies at the lower end of the observed
25 ratio, while only between 200 and 400 m particles are overestimated compared to dissolved Mn. Whereas in the upper 100 m of the ocean both simulations are consistent with those from the VERTEX-IV cruise, in the deep ocean only the *Reference* simulation is accurate. In *OxidThreshold* the ratio is underestimated.

4 Discussion

Many of the properties of the Mn_{diss} distribution in the world ocean are reproduced by our *Reference* simulation (Sect. 3.2).

30 However, some discussion of the assumptions of the underlying processes in the model is required.

4.1 Margin sediments

We have chosen to use a simple sediment flux parameterisation for this study. In our model, Mn_{diss} is added to bottom water from anoxic sediments analogously to the iron flux in the model PISCES (Aumont et al., 2015). We think that coupling the model to the sediment would be difficult and maybe not that useful, because the sediment source is large at shallow sediments, while most of the Mn burial occurs near the hydrothermal vents. This first-order approach works mostly well, but shortcomings can be identified.

On the one hand, the flux may be overall underestimated, for instance because Mn reduction releases a higher free energy than Fe reduction, so that Mn oxides reduce more easily (Sarmiento and Gruber, 2006, pp. 232–236). Furthermore, high benthic Mn fluxes have been observed from eastern North Pacific marine sediments (McManus et al., 2012). On the other hand, at some large-shelf regions, like in parts of the Arctic Ocean, the Mn flux is overestimated, because the low model resolution does not handle large shelf regions well.

Slomp et al. (1997) measured pore water concentrations of dissolved Fe and Mn. The Mn/Fe ratios based on their maximum pore water concentrations range from 0.2 to 1.2. Balzer (1982) reports fluxes from sediment to bottom water of dissolved Mn and Fe that yield an average ratio of 0.5. The value that we used, 0.2, is at the lowest end of the reported values. This choice is mainly due to the fact that the crude iron flux parameterisation results in increases of the Mn flux that are out of proportion in some regions of the ocean. This is a known limitation of this parameterisation that can also be seen in the modelled iron distribution (Tagliabue and Völker, 2011). In the Arctic Ocean, a higher sedimentary Mn input would induce an even stronger overestimation than it already does with $Mn/Fe=0.2$ (Fig. 4c). This had also been concluded by Van Hulten et al. (2013) who presented a sensitivity study that used the same parameterisation for aluminium sediment input as for Mn here.

20 Nevertheless, a higher Mn_{diss} input would be beneficial for the simulation in the Atlantic and especially the Pacific Ocean where $[Mn_{diss}]$ is underestimated everywhere but south of 50° S. Underestimation may be due to upwelling from effluxes from OMZ sediments in several regions of the ocean. In the East Pacific Ocean the California Current induces Ekman transport, and hence equatorial upwelling. High $[Mn_{diss}]$ in the North Pacific Ocean are suggested by upwelling from OMZ sediments in the East Pacific (Landing and Bruland, 1987). This is partly captured by our model (Fig. 8a, East Pacific around 20° N). In the South Pacific Ocean this effect is more clear in the data of Resing et al. (2015) (Fig. 8a,b, East Pacific around 20° S).

To summarise, we chose to use a simple sediment flux parameterisation, which is not adequate for representing Mn_{diss} effluxes from OMZ sediments. Albeit out of the scope of this study, a better parameterisation, or even an explicit sediment model, could make for a more realistic simulation, both in terms of accuracy as well as reliability in a methodological sense (Petersen, 2006, § 3.6). This should be pursued in future studies.

30 4.2 Biological cycle

4.2.1 General discussion and the GIPY5 transect

Our model includes biological processes involving manganese in a way similar to that of phosphorus where the rate variables are multiplied by a typical Mn/P ratio based on the measured plankton content ratio. We had chosen to follow all phosphorus cycling as described by Aumont et al. (2015) for consistency reasons. We did not want to evaluate all potential details of biology but rather take a simple approach.

At this stage, we lack observational constraints allowing us to develop a more complex representation of the biological Mn cycle. Specifically, even though in the Southern Ocean upper waters the removal and remineralisation cycle of Mn_{diss} correlates well with the nutrients PO_4^{3-} and NO_3^- (Middag et al., 2011a), there is no mechanistic evidence for how the uptake-remineralisation process exactly should work for Mn. Of course, Mn plays an important role in biology. It even limits primary production in parts of the Southern Ocean (Coale, 1991; Middag et al., 2011a, 2013; Browning et al., 2014), although the trace nutrient iron often appears more significant (Buma et al., 1991). This is at least suggested by the observational data: at the Zero-Meridian GIPY5_e transect south of 58° S, the dissolved Mn concentration is very low at the surface, but from ~ 50 m to 250 m there is a maximum of $[Mn_{diss}]$ (Fig. 9, left transect). This is probably due to remineralised particulate organic matter (Middag et al., 2011a). The maximum is not found in the model. For instance, near 66° S at 1000 m upwards the modelled $[Mn_{diss}]$ decreases monotonically, which suggests that biology does not explain the subsurface maximum. However our model may not be adequate. Firstly, it may be unreasonable to treat Mn biologically identical to P. Secondly, the subsurface maximum can be a direct consequence of a low oxygen concentration, because that would decrease the oxidation rate k_{ox} , and our redox model does not depend on $[O_2]$. This is not the most probable hypothesis, because at least during the ANT XXIV/3 expedition only a slight oxygen minimum occurred around 300–400 m depth which lies well below the Mn maximum (Middag et al., 2011a). Thirdly, there could be problems with the underlying model. For instance, if the vertical mixing in the underlying circulation model is too strong, low- $[Mn_{diss}]$ waters mix from the deep Southern Ocean upwards through the 100 m to 200 m region, possibly removing the whole signal.

The model suggests that a Mn biological cycle following P may have an important contribution to the Southern Hemisphere, and especially at low latitudes. This should be tested further through field experiments and more sophisticated model simulations.

By far the most settling Mn is in the biologically incorporated portion. The oxidised fraction has only a significant, though still smaller, contribution near hydrothermal vents compared to incorporated Mn (not presented). It seems that, especially for low latitudes, biological incorporation is more important than oxidation.

4.2.2 Relation to redox and settling

Our model yields small $[Mn_{diss}]$ values at the surface of the eastern Pacific Ocean (equator and 40° S, Fig. 8a). This appears to be due to the biological cycle (Fig. 11a). Moreover, the model does not include the effects of oxygen minima, especially

that comes from reductive sediments and upwells in the eastern Pacific Ocean. The remineralisation signal generates a more accurate simulation at intermediate depths (same figures, panels b; Fig. 12b).

Overall, the amount of manganese ($Mn_{diss} + Mn_{ox}$) in the world ocean is slightly larger with biology (440 Gmol) compared to without (409 Gmol, Table 6). This is not expected, because biological incorporation would give an extra sink of Mn incorporated in particulate organic matter, referred to as POMn. The explanation of this paradox lies in the interaction between the biological cycle and the settling process of Mn_{ox} in the model.

We apply a concentration-dependent sinking speed, which generates an approximately homogeneous distribution of dissolved manganese of around $0.125 \mu M$ in much of the deep ocean. There is a finite range of input fluxes where this works. If the input flux gets above a certain value, $[Mn_{diss}]$ exceeds this background value, as it occurs in the Atlantic Ocean north of $35^\circ N$. However, if the input flux is low enough, $[Mn_{diss}]$ decreases below this background value because of slow Mn_{ox} settling, like it does in the East Pacific Ocean. Inflow of higher $[Mn_{diss}]$ waters by large-scale circulation also counts as a source that can elevate Mn, but the currents in the deep Pacific Ocean are slow.

The biological processes modify this behaviour. In PISCES there are two types of particulate carbon: one of large particles that sinks with $50 m d^{-1}$ (in our version) and one of small particles that sinks with $2 m d^{-1}$. Most carbon, and so manganese, is present in the pool of small particles. Even though, at least in the model, the Mn sources in the Pacific Ocean are small, there is a lot of Mn_{diss} in the surface ocean that would ultimately be Mn_{ox} if it were mixed down below the photic zone depth. However, with the biological cycle enabled, phytoplankton incorporates a part of the Mn_{diss} . POMn (just like POC) sinks down into the deep ocean. During the downward propagation of the organic Mn particles, they remineralise, releasing Mn_{diss} in the deep ocean. A part of this Mn_{diss} is converted to Mn_{ox} that sinks down inefficiently ($1 m d^{-1}$) when Mn_{ox} has not yet reached the equilibrium concentration \mathcal{X}_{thr} . The remineralised POMn fills up, through oxidation, the Mn_{ox} pool in the deep ocean.

Biological processes store Mn in plankton, which is preserved from strong Mn oxide settling (in the model). This new pool of manganese fills the pools of dissolved and oxidised Mn in the deep ocean. This new pool is provided by the plankton that took it from the Mn_{diss} pool in the photic zone. Without biology this would be available for oxidation to Mn_{ox} during night. Adding this extra Mn to the integrated total yields a higher Mn quantity in *Reference* than in *NoBio*.

4.3 Redox rates

While we chose to model Mn reduction and oxidation as first-order-reaction kinetics, redox of Mn within the water column is a combination of several processes. Firstly, Mn_{ox} is subject to non-biological reduction, significantly stimulated by sunlight (e.g. Sunda and Huntsman, 1994). This is taken into account by using different k_{red} for the euphotic and aphotic zones of the ocean. Secondly, the rate of oxidation is enhanced by microbes (bacteria and fungi) in regions where Mn_{diss} supply is high (Sunda and Huntsman, 1994; Tebo et al., 2005), and it depends on the O_2 concentration and pH. This has been observed in the North Pacific Ocean (Johnson et al., 1996, and references therein), as well as in some GEOTRACES transects in the Atlantic Ocean. Measurements from the West Atlantic Ocean and Zero-Meridian Southern Ocean GEOTRACES cruises show some variations with depth, e.g. near $10^\circ N$ and near $40-47^\circ N$ (top of Fig. 7). This corresponds to the maximum of dissolved Fe associated with the oxygen minimum as reported for the same cruise by Rijkenberg et al. (2014). Our model does not reproduce such a

30 maximum of dissolved Mn, because Mn redox does not depend on $[O_2]$. Furthermore, fluxes from reducing sediments are not expected to reproduce the feature either (Johnson et al., 1996). This means that at least for those regions it would be logical to include a dependence on $[O_2]$. However, this is not the pattern that attracts the most attention when looking at the full transect. The striking patterns are rather the high concentration at the surface and near the equator around 2.5 km to 3 km depth (Fig. 7). For this reason we have not included a dependency on $[O_2]$ to the model. In other words, while k_{ox} depends on oxygen, we assumed here that the oxygen concentration is generally so high that its variation does not affect k_{ox} .

In our standard model, oxidation takes place everywhere with the same $k_{ox} = 0.341 \times 10^{-3} \text{ h}^{-1}$. Based on the underestimation of the concentrations in the surface of the Atlantic and Pacific oceans, one may think that $[Mn_{ox}]$ is too high compared to $[Mn_{diss}]$. In other words, assuming approximate equilibrium, the ratio k_{ox}/k_{red} could be too high. However, the k values in our model are chosen to get values consistent with the ratio of concentration measurements of Mn_{ox} and Mn_{diss} . To this end we
5 set $k_{red,light}$ to the mean value found by Sunda and Huntsman (1994), then used the dissolved and particulate profiles in Fig. 4 of Bruland et al. (1994) to derive k_{ox} and $k_{red,dark}$. If we were to ignore that ratio, and increase the first-order rate constants for reduction with a factor of five (a factor of three higher than the upper value of the range found by Sunda and Huntsman (1994)), the model would not yield sufficiently high $[Mn_{diss}]$ (results not presented).

Still, one may introduce a threshold on the oxidation process (instead of the settling of the particles). The purpose would
10 be to find out if this would result in higher surface $[Mn_{diss}]$, without having a significant effect on the ratio between the dissolved manganese and the oxides. In such a model simulation, oxidation only takes place when $[Mn_{diss}]$ is higher than a certain threshold \mathcal{D}_{thr} , equal to 0.125 nM in *OxidThreshold*. Analogous to Eq. 10, k_{ox} is multiplied by $H(\mathcal{D} - \mathcal{D}_{thr})$. While this yields a distribution of Mn_{diss} not significantly worse in the West Atlantic Ocean than that of *Reference*, the Mn_{ox}/Mn_{diss} concentration ratio around 500 m and below is then strongly underestimated compared to the VERTEX data (Fig. 16, right
15 panel, red dashed line). This pleads for using an aggregation rather than an oxidation threshold.

4.4 Export dynamics


The very homogeneous low dissolved concentration of Mn at ~ 0.1 nM in the deep waters (Fig. 8) of the West Atlantic GA02 section is remarkable. Such deep water values of ~ 0.1 nM were also found in the deep waters of the Antarctic Ocean (Middag et al., 2011a, 2012, 2013), as well as deep waters of the Mediterranean Sea (2013 cruise, data will be included in
20 the GEOTRACES Intermediate Data Product 2017). For the Atlantic Ocean, these deep concentrations are much lower than the dissolved Mn values of ~ 0.5 nM to 0.6 nM reported for deep samples at several Atlantic stations in the 1990s (Saager et al., 1997; Statham and Burton, 1986; Statham et al., 1998). The sampling resolution is too coarse to exclude spatial variation, but it appears the values produced by pioneering efforts to study the oceanic Mn distributions are overestimations.

The new ultraclean sampling methods (Rijkenberg et al., 2015), rigorous calibrations and excellent accuracy at the Bermuda
25 crossover station for twelve trace metals (Middag et al., 2015b) now yield a much larger database of uniformly very low Mn ~ 0.1 nM in deep waters. Given these very uniform and much lower background Mn concentrations, the hydrothermal Mn plumes are better discernible and much larger widespread than realised in the 1990s. Exactly over or very near the ridge crest this is recently shown dramatically in a plume of more than 500 km wide over the Gakkel Ridge in the high-latitude Arctic

Ocean (Middag et al., 2011b), and in the Antarctic Ocean with a plume that extends more than 1000 km, at a site near the Bouvet triple junction where three ocean ridges meet (Middag et al., 2011a). Similar long range extent of the Mn hydrothermal plume has recently been discerned in the Pacific Ocean (Resing et al., 2015).

The West Atlantic GA02 section is quite far west from the Mid-Atlantic Ridge, nevertheless the hydrothermal plume extent still is seen in a dissolved Mn maximum of 0.2 nM to 0.3 nM in the 2500 m to 3000 m depth range just south of the Equator. This is consistent with observations of the more complete hydrothermal plumes along two meridional (east-west) sections across the Mid Atlantic Ridge. These are (i) at a station (USGT11-16 at 26.14° N, 44.83° W) over the TAG hydrothermal site (discovered in 1972 by the NOAA Trans-Atlantic Geotraverse Project), and adjacent stations of the GEOTRACES GA03 section in the North Atlantic Ocean (Wu et al., 2014), as well as (ii) at ~12° S in the South Atlantic Ocean where also distinct plumes of hydrothermal Mn were observed (Noble et al., 2012, their Fig. 1).

Remarkably, the background concentration of dissolved Mn is quite uniform in the 0.10 to 0.15 nM range. Preliminary equilibrium calculations predict an equilibrium of 0.19 nM total dissolved Mn with the solid phase pyrochroite $\text{Mn}(\text{OH})_2(s)$. This total dissolved equilibrium concentration is quite close to the observed 0.10 to 0.15 nM background range measured in the oceans. Please notice that in pyrochroite the Mn is divalent Mn(II) just as the dissolved Mn forms. Disregarding organic complexes, the latter dissolved equilibrium Mn(II) forms would comprise 0.148 nM (78 %) free Mn^{2+} ion, 0.027 nM (~ 14 %) MnCl^+ ion, 0.011 nM (~ 5.8%) $\text{MnCl}_2^0(aq)$ and a very low abundance 0.000028 nM (~0.0015 %) MnOH^+ ion. The above most simple Reaction R1 for oxidation removal from seawater, may perhaps be envisioned to be split into several reaction steps as follows:



where the dominant (78 %) dissolved Mn^{2+} species is via the minor dissolved MnOH^+ species converted to solid $\text{Mn}(\text{OH})_2$, and next this is oxidised to one or several of various oxide forms, for example hausmannite Mn_3O_4 that is a mixed valency state (III, IV) Mn-oxide, or $\text{Mn}(\text{IV})\text{O}_2$ pyrolusite or birnessite. Please notice that in the Reference simulation an aggregation threshold for particulate Mn at 25 pM was invoked, that via redox rate constants is based on typical background dissolved Mn concentrations that was chosen at 0.125 nM. Latter threshold value of 0.125 nM was required to have the model simulate the observed data, and is quite close to the equilibrium concentration of 0.19 nM versus pyrochroite. The notion of perhaps an equilibrium control is not new. Previous pioneering measurements led to an apparent background concentration of Mn of 0.4 nM or higher, that is now shown to be too high, yet otherwise attempts were made to compare these values in terms of equilibrium versus hausmannite Mn_3O_4 (Klinkhammer and Bender, 1980). Von Langen et al. (1997) did propose a reaction mechanism also involving conversion of Mn^{2+} via $\text{Mn}(\text{OH})^+$ as in above (R2). Further research is needed to verify and confirm the here discussed hypothesis of equilibrium of dissolved Mn at ~ 0.19 nM versus pyrochroite.

In the *NoThreshold* simulation $[\text{Mn}_{\text{diss}}]$ is underestimated in much of the ocean. When removing the threshold, the RMSD significantly worsens: it appears that the threshold is needed to reasonably simulate $[\text{Mn}_{\text{diss}}]$. But there are several alternatives to the aggregation threshold that may keep $[\text{Mn}_{\text{diss}}]$ in check.

30 Firstly, instead of an aggregation threshold, a criterion on oxidation may be imposed, because, hypothetically, there may be a lower limit below which oxidising microbes would not proliferate and the pseudo-first-order rate constant would then be lower. The simulation with an oxidation threshold based on this idea gives similar results as the aggregation threshold, but as we saw in the previous section, it would influence significantly the Mn_{ox}/Mn_{diss} concentration ratio in the deep ocean, making for a less accurate simulation. Therefore, this alternative should probably not be pursued.

Secondly, the process of oxidation is strongly mediated by adsorption onto particles, after which it forms larger aggregates that settle faster. This is now parameterised by increasing the settling velocity of particles with depth. However, a more explicit model for adsorption/desorption could be useful. Possibly, much of the Mn_{ox} does not settle efficiently. In that case, another tracer of adsorbed Mn is needed that would export oxidised Mn (in adsorbed or aggregated form) but not too fast to account 5 for the background $[Mn_{diss}]$. The adsorption should not take place onto a homogeneous pool of particles as is effectively done in our model, but rather onto e.g. calcium carbonate ($CaCO_3$) (Martin and Knauer, 1983; Fraústo da Silva and Williams, 2001) or lithogenic particles (Roy-Barman, 2009). Sediment samples show a strong correlation between authigenic manganese and lithogenic particles, while Mn does not show any correlation with biogenic silica, POC or $CaCO_3$ according to recent observations and modelling (Roy-Barman, 2009). On the one hand, this suggests that only lithogenic particles are a scavenger 10 of Mn. On the other hand, we do not know whether the correlation comes from Mn adsorption onto lithogenic particles, or if it is lithogenic in itself.

Thirdly, oxidation is mediated by microbes (mainly bacteria) (Sunda and Huntsman, 1994; Von Langen et al., 1997; Tebo et al., 2005; Fraústo da Silva and Williams, 2001). Possibly they prolong the time of Mn_{ox} particles spent in the euphotic zone, but this has not been tested. In the model we assume that these processes are included in the redox rate constant. If we 15 want the redox rate constants to depend on inhomogeneously distributed quantities like the bacterial distribution or the oxygen concentration, the equations need to be modified, and our model may need to be extended. Subsequently, more Mn would stay suspended.

Fourthly, it has been suggested that dissolved Mn ligands keep Mn in solution (Sander and Koschinsky, 2011; Madison et al., 2013; Luther et al., 2015). That would result in less conversion to manganese oxide aggregates, and hence may solve 20 the removal problem, but this may yield wrong Mn_{ox}/Mn_{diss} concentration ratios like we saw for *OxidThreshold*. Similarly, nanoparticles, which have a size that falls within the operationally defined dissolved Mn, keep Mn afloat as long as they do not aggregate.

4.4.1 Hydrothermal activity

In the model, hydrothermal Mn_{diss} is added to the ocean based on a model proxy of 3He (Dutay et al., 2004). For the *Reference* 25 simulation the hydrothermal Mn_{diss} flux was set to 0.10×10^9 mole Mn for each mole of 3He . There is no reference to literature values in Table 3 since these are very uncertain. For instance, Klinkhammer and Bender (1980) note that “Corliss and Dymond (1979) reported that the Mn : 3He ratios vary by over a factor of 2 in the Galapagos vents, and as we discuss below, similar variations are found at 21° N”. We had chosen a value to yield an acceptable distribution of Mn_{diss} .

The goal of *LowHydro* is to investigate whether the combination of the high hydrothermal input of Mn_{diss} , and strong aggregation is needed to get a more accurate distribution of Mn_{diss} , namely that of *Reference*. To this end, the choice was made to decrease hydrothermal input and also to decrease the settling velocity in *LowHydro*. In *LowHydro* the spatial distribution of $[Mn_{diss}]$ becomes more homogeneous. Neither the low background concentration nor the high hydrothermal vent concentrations are reproduced anymore (Fig. 14). Decreasing hydrothermal input and the settling velocity by a smaller factor than 10 results in a Mn_{diss} distribution that is better than *LowHydro* but not as good as in *Reference* (not presented). In any case, the main result is that a high hydrothermal input very similar to or higher than that used in the *Reference* simulation is needed for an accurate simulation of $[Mn_{diss}]$.

According to observations a negligible amount of Mn from hydrothermal vents reaches the surface ocean. As Bruland and Lohan (2006) put it: “The hydrothermal input of iron and manganese [...] is essentially all scavenged and removed in the deep sea prior to having a chance to mix back into the surface waters.” However, this is what we have thought about iron while there are now doubts about this (Tagliabue et al., 2010). Similarly, hydrothermal Mn may not be completely removed from the ocean before reaching the surface by currents and vertical mixing. There are different potential reasons for this. First, the oxidation of Mn(II) is thermodynamically favoured, but the large activation energy of Mn(II) oxidation renders Mn(II) stable in aquatic environments (Nealson, 2006). In our model we assumed that the activation energy is not an issue. Furthermore, the background “dissolved” Mn particles are not Mn^{2+} ions but rather suspended colloids or complexes. Candidates include colloidal Mn_{ox} and Mn bound to ligands, respectively the second and third potential reasons for the observed background “ Mn_{diss} ” concentration. The low Mn_{diss} concentrations away from hydrothermal vents are established in our model by removing Mn_{ox} from the high hydrothermal Mn input down to 25 pM.

Interestingly, in this way, in *Reference*, hydrothermal flux accounts for 92 % of the total Mn input. Dust deposition is 18 times as small as hydrothermal input, and still $[Mn_{diss}]$ in the upper 1000 m of the ocean is dominated by Mn_{diss} release from marginal sediments and dissolution from deposited dust. Nonetheless, some of the hydrothermally derived Mn_{diss} reaches the ocean surface. In our model, 8 % of Mn in the surface is hydrothermal of origin. The reason for the importance of sediment and dust is that in the model the settling velocity of Mn_{ox} is much higher near hydrothermal vents (up to 10 m d^{-1}) than in the surface ocean (1 m d^{-1}). The modelling study by Lavelle et al. (1992) suggests a settling velocity much higher than our w_{ox} . They used a model with an additional tracer of large particles (aggregate products) that had settling speeds of up to 175 m d^{-1} (near hydrothermal vents, as they model those regions). This shows that using an increasing settling velocity is justified. Moreover, it suggests that at least two particle tracers are needed in the model. It is worth investigating in future studies if this would make for a more accurate and reliable model.

25 5 Conclusions

This is the first study in which the 3-D distribution of Mn_{diss} has been modelled and compared to the recent observations from the GEOTRACES programme. A combination of photoreduction and Mn sources to the upper ocean yields high surface concentrations of dissolved manganese. However, the concentration is somewhat underestimated by the model. The Mn sources

to the ocean surface and the sinks are uncertain and could therefore be adjusted for a more accurate simulation. The most important sources for the upper ocean are sediments, dust, and, more locally, rivers, whereas hydrothermal vents are the most important in the deep ocean. The observed sharp hydrothermal signals are produced by assuming both a high source and a strong removal of Mn near hydrothermal vents. Our model further shows that the Mn at the surface in the Atlantic Ocean moves downwards into the North Atlantic Deep Water, but because of strong removal the Mn signal does not propagate southwards.

There is a mainly homogeneous background concentration of dissolved Mn of about 0.10 nM to 0.15 nM throughout most of the deep ocean. Our model reproduces this by means of a threshold on solid manganese oxides of 25 pM, suggesting that a minimal concentration of particulate Mn is needed before aggregation and removal become efficient. An aggregation threshold, as applied in our model, appears reasonable, and does not affect the modelled Mn_{ox}/Mn_{diss} concentration ratio. An oxidation threshold is more troublesome as it affects this ratio in an undesirable way. Since the settling condition as a sole threshold already appears to mitigate most of the removal, it is reasonable to further develop the model with the aggregation threshold and without the oxidation threshold. Still, the simplified redox and subsequent settling of Mn_{ox} is incomplete and could be too imprecise, as future studies might show.

Our model includes biology, and this has a big impact in the Pacific Ocean. However, we have at the moment no clear evidence for typical uptake-remineralisation processes as for iron. Biological incorporation should be improved in future versions of the model. Besides processes in the photic zone, it may also be needed to model microbial activity throughout the ocean, as that is not homogeneously distributed. Our chemical first-order reaction may be insufficient for many modelling purposes.

Hydrothermal fluxes of Mn were set to such a high rate that we must assume 96 % is scavenged near the outflow of the vents. This choice was made to account for the local high Mn_{diss} near the oceanic ridges. This was combined with a high w_{ox} in the deep ocean to prevent hydrothermal Mn_{diss} spreading far away from the oceanic ridges in a too high concentration. As an alternative, settling may not need to be set as high as 10 m d^{-1} in the deep ocean, but then it is unclear how to simulate the local nature of $[Mn_{diss}]$ anomalies near the ocean ridges. One possibility is to include extra species of Mn, which may include a very fast sinking particle and a very reactive Mn species.

Process studies of Mn are necessary to determine the rate constants, and possibly thresholds, for redox, scavenging and aggregation. More measurements of particulate Mn_{ox} concentrations would be useful as well. When measuring particulate Mn, it needs to be clear what is measured exactly. It would for instance be useful if (1) a range of particle sizes were measured, and (2) a structural analysis of the particles were performed, such that one can unambiguously say onto which particle Mn were adsorbed or into which particle it were incorporated.

25 Appendix A: Data–model comparison

To compare quantitatively the model results of this study with the observations, the focus is on the West Atlantic Ocean. The data of the GA02 GEOTRACES transect in that region are used. First the model output is horizontally interpolated onto the station coordinates, keeping the vertical model grid (10 m at the surface up to 500 m thickness near the bottom). Then the model output that lies closest to each of the observations are associated with each other. To be precise, for each observation, we take the shallowest gridbox whose upper bound lies deeper than the observation, after which residuals can be defined as $P_i - O_i$, where O_i is the observed and P_i the modelled $[Mn_{diss}]$, for each

$i \in \{1, \dots, N\}$, with N the number of data points. The interpolation introduces a representative error that is taken to be part of the residual (Van Hulten, 2014). Then several statistics are determined, namely the Root Mean Square Deviation (RMSD), the Reliability Index (RI) and the correlation coefficient r .

In addition to classical statistical indices (r , RMSD), another additional performance indicator has been used as suggested in previous skill assessment studies (Stow et al., 2009; Vichi and Masina, 2009): the reliability index (Leggett and Williams, 1981). The reliability index “quantifies the average factor by which model predictions differ from observations” (Stow et al., 2009). It is in essence the root-mean-square deviation, but it uses the logarithm of the residual. This is useful when both large and small values need to be considered (as for this case). The RI is given by

$$RI = \exp \sqrt{\frac{1}{N} \sum_{i=1}^N \left(\log \frac{O_i}{P_i} \right)^2}. \quad (\text{A1})$$

where O_i is the measured concentration with index i , P_i is the model prediction associated to the respective observation i , and N is the number of observations.

Furthermore, for each sensitivity simulation the significance of the change in each goodness-of-fit statistic compared with the corresponding reference simulation is calculated. This is determined by means of a Monte Carlo simulation on the reference simulation for which a subsample of 400 is randomly selected from the original set of 1800 or 1650 data–model points. They are the pairs of observations and model output, both on the model grid. This is done 50 000 times, and from this the 2σ confidence interval is calculated (the mean \pm two times the standard deviation). Suppose that we want to simulate q , and assume q is in steady state. Given a simulation X , for each model simulation Y resulting in $q_Y(\mathbf{x})$, the average RMSD of the Monte Carlo simulation of $q_Y(\mathbf{x})$ must be outside the 2σ confidence range of the RMSD distribution of $q_X(\mathbf{x})$ to say that Y is a significant improvement or worsening compared to X .

The statistics tell us how accurate our simulations are, but it does not tell how reliable they are in a wider sense (Petersen, 2006). For this reason, these statistics are amended with comparison plots and an extensive discussion of those. For the visual comparison between model and observations, horizontal and vertical cross-sections of the model data are presented. Using the same colour scale, observations are plotted as coloured dots to directly compare the model with the observations. Horizontal $[\text{Mn}_{\text{diss}}]$ sections are presented for four different depths, where “surface” signifies the average over the upper 45 m, “500 m” is 400–600 m averaged, ‘2500 m’ is 2100–2900 m averaged and ‘4500 m’ is 4000–5000 m averaged. The colour scale is not linear to better show the main features at both low and high concentrations of Mn_{diss} . The vertical $[\text{Mn}_{\text{diss}}]$ sections are calculated from the 3-D model data by converting the ORCA2 gridded model data to a rectilinear mapping and interpolating the rectilinear data onto the cruise track coordinates.

20 Appendix B: Code availability

We used NEMO 3.6 svn 7036 that can be checked out through this command:

```
$ svn checkout -r 7036 http://forge.ipsl.jussieu.fr/nemo/svn/branches/2015/nemo_v3_6_STABLE
```

However, the latest stable version of NEMO is usually advisable. A login will be asked that you can create on the NEMO website.

The manganese module comes as part of different Fortran 90/95 source code files that are to be found in the electronic supplement. Those should be put in `NEMOGCM/CONFIG/*/MY_SRC/` that adds and overrides symlinks in `NEMOGCM/CONFIG/*/WORK/`. The asterisk must be replaced with your configuration name that should be based on `ORCA2_OFF_PISCES`.

Both NEMO and the manganese model are available under the CeCILL licence.

Acknowledgements. We would like to thank several people in particular who helped in different ways in this study. Angela Milne, William Landing and Joseph Resing kindly provided their manganese data from the Pacific Ocean. We thank Caroline Slomp, Catherine Jeandel and Micha Rijkenberg for the useful discussion. Loes Gerringa has kindly provided speciation calculations on manganese. Loes Gerringa and Micha Rijkenberg organised and served as expedition leaders of the three cruises of the GA02 section. This research was funded by the NWO (grant 839.08.410; GEOTRACES, Global Change and Microbial Oceanography in the West Atlantic Ocean and grant 820.01.014 GEOTRACES Netherlands-USA Joint Effort on Trace Metals in the Atlantic Ocean). This study was partly supported by a Swedish Research Council grant (349-2012-6287) in the framework of the French-Swedish cooperation in the common research training programme in the climate, environment and energy agreement between VR and LSCE, for the project “Particle transport derived from isotope tracers and its impact on ocean biogeochemistry: a GEOTRACES project in the Arctic Ocean”. The sampling and analysis of the data of Milne and Landing (unpublished data) were supported by NSF grants OCE-0223378, OCE-0550317 and OCE-0649639.

References

- Aguilar-Islas, A. and Bruland, K.: Dissolved manganese and silicic acid in the Columbia River plume, *Mar. Chem.*, 101, 233–247, doi:10.1016/j.marchem.2006.03.005, 2006.
- 5 Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochem. Cy.*, 20, GB2017, doi:10.1029/2005GB002591, 2006.
- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies, *Geosci. Model Dev.*, 8, 2465–2513, doi:10.5194/gmd-8-2465-2015, 2015.
- Baker, A., Jickells, T., Witt, M., and Linge, K.: Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean, *Mar. Chem.*, 98, 43–58, doi:10.1016/j.marchem.2005.06.004, 2006.
- 10 Baker, A. R., Landing, W. M., Bucciarelli, E., Cheize, M., Fietz, S., Hayes, C. T., Kadko, D., Morton, P. L., Rogan, N., Sarthou, G., Shelley, R. U., Shi, Z., Shiller, A., and van Hulst, M. M. P.: Trace element and isotope deposition across the air–sea interface: progress and research needs, *Phil. Trans. R. Soc. A: Mathematical, Physical and Engineering Sciences*, 374, doi:10.1098/rsta.2016.0190, part of the special issue *Biological and climatic impacts of ocean trace element chemistry*, 2016.
- 15 Balistrieri, L., Brewer, P., and Murray, J.: Scavenging residence times of trace metals and surface chemistry of sinking particles in the deep ocean, *Deep-Sea Res. Pt A*, 28, 101–121, 1981.
- Balzer, W.: On the distribution of iron and manganese at the sediment/water interface: thermodynamic versus kinetic control, *Geochim. Cosmochim. Ac.*, 46, 1153–1161, doi:10.1016/0016-7037(82)90001-1, 1982.
- Barnier, B., Madec, G., Penduff, T., Molines, J.-M., Tréguier, A.-M., Sommer, J. I., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., 20 Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and Cuevas, B. D.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy-permitting resolution, *Ocean Dynamics*, 56, 543–567, doi:10.1007/s10236-006-0082-1, <http://archimer.ifremer.fr/doc/2006/publication-3514.pdf>, 2006.
- Bender, M. L., Klinkhammer, G. P., and Spencer, D. W.: Manganese in seawater and the marine manganese balance, *Deep-Sea Res.*, 24, 799–812, doi:10.1016/0146-6291(77)90473-8, 1977.
- 25 Bishop, J. K. and Fleisher, M. Q.: Particulate manganese dynamics in Gulf Stream warm-core rings and surrounding waters of the N.W. Atlantic, *Geochim. Cosmochim. Ac.*, 51, 2807–2825, doi:10.1016/0016-7037(87)90160-8, 1987.
- Bortleson, G. and Lee, G.: Phosphorus, iron, and manganese distribution in sediment cores of six Wisconsin lakes, *Limnol. Oceanogr.*, 19, 794–801, 1974.
- Bowers, T., Campbell, A., Measures, C., Spivack, A., Khadem, M., and Edmond, J.: Chemical controls on the composition of vent fluids at 30 13°–11° N and 21° N, East Pacific Rise, *J. Geophys. Res.*, 93, 4522–4536, doi:10.1029/JB093iB05p04522, 1988.
- Boye, M., Wake, B. D., Garcia, P. L., Bown, J., Baker, A. R., and Acherberg, E. P.: Distributions of dissolved trace metals (Cd, Cu, Mn, Pb, Ag) in the southeastern Atlantic and the Southern Ocean, *Biogeosci.*, 9, 3231–3246, doi:10.5194/bg-9-3231-2012, 2012.
- Boyle, E. A., Bergquist, B. A., Kayser, R. A., and Mahowald, N.: Iron, manganese, and lead at Hawaii Ocean Time-series station ALOHA, *Geochim. Cosmochim. Ac.*, 69, 933–952, doi:10.1016/j.gca.2004.07.034, 2005.
- 35 Browning, T. J., Bouman, H. A., Henderson, G. M., Mather, T. A., Pyle, D. M., Schlosser, C., Woodward, E. M. S., and Moore, C. M.: Strong responses of Southern Ocean phytoplankton communities to volcanic ash, *Geophys. Res. Lett.*, 41, 2851–2857, doi:10.1002/2014GL059364, 2014.

- Bruland, K. and Lohan, M.: Controls of Trace Metals in Seawater, in: *The Oceans and Marine Geochemistry*, edited by Elderfield, H., vol. 6, chap. 6.02, pp. 23–47, Elsevier, http://es.ucsc.edu/~kbruland/Manuscripts/BRULAND/BrulandChpt6_02_TraceMetals.pdf, 2006.
- Bruland, K., Orians, K., and J.P., C.: Reactive trace metals in the stratified central North Pacific, *Geochim. Cosmochim. Ac.*, 58, 3171–3182, doi:10.1016/0016-7037(94)90044-2, 1994.
- 5 Buck, C., Landing, W., Resing, J., and Measures, C.: The solubility and deposition of aerosol Fe and other trace elements in the North Atlantic Ocean, *Mar. Chem.*, 120, 57–70, doi:10.1016/j.marchem.2008.08.003, 2010.
- Buma, A. G. J., de Baar, H. J. W., Nolting, R. F., and van Bennekom, A. J.: Metal enrichment experiments in the Weddell-Scotia Seas: Effects of iron and manganese on various plankton communities, *Limnol. Oceanogr.*, 36, 1865–1878, doi:10.4319/lo.1991.36.8.1865, 1991.
- Charette, M. A., Lam, P. J., Lohan, M. C., Kwon, E. Y., Hatje, V., Jeandel, C., Shiller, A. M., Cutter, G. A., Thomas, A., Boyd, P. W., Homoky, W. B., Milne, A., Thomas, H., Andersson, P. S., Porcelli, D., Tanaka, T., Geibert, W., Dehairs, F., and Garcia-Orellana, J.: Coastal ocean and shelf-sea biogeochemical cycling of trace elements and isotopes: lessons learned from GEOTRACES, *Phil. Trans. R. Soc. A*, 374, doi:10.1098/rsta.2016.0076, part of the special issue *Biological and climatic impacts of ocean trace element chemistry*, 2016.
- 10 Coale, K. H.: Effects of iron, manganese, copper, and zinc enrichments on productivity and biomass in the subarctic Pacific, *Limnol. Oceanogr.*, 36, 1851–1864, doi:10.4319/lo.1991.36.8.1851, 1991.
- 15 Corliss, J. and Dymond, J.: on the Galapagos Rift, *Science*, 203, 16, doi:10.1126/science.203.4385.1073, 1979.
- Doi, T., Obata, H., and Maruo, M.: Shipboard analysis of picomolar levels of manganese in seawater by chelating resin concentration and chemiluminescence detection, *Anal. Bioanal. Chem.*, 378, 1288–1293, doi:10.1007/s00216-003-2483-z, 2004.
- Douville, E., Charlou, J., Oelkers, E., Biennu, P., Jove Colon, C., Donval, J., Fouquet, Y., Prieur, D., and Appriou, P.: The rainbow vent fluids (36° 14' N, MAR), *Chemical Geology*, 184, 37–48, doi:10.1016/S0009-2541(01)00351-5, 2002.
- 20 Dutay, J., Jean-Baptiste, P., Campin, J., Ishida, A., Maier-Reimer, E., Matear, R., Mouchet, A., Totterdell, I., Yamanaka, Y., Rodgers, K., et al.: Evaluation of OCMIP-2 ocean models' deep circulation with mantle helium-3, *J. Mar. Syst.*, 48, 15–36, doi:10.1016/j.jmarsys.2003.05.010, 2004.
- Dutay, J.-C., Tagliabue, A., Kriest, I., and Hulten, M.M.P. van: Modelling the role of marine particle on large scale ²³¹Pa, ²³⁰Th, Iron and Aluminium distributions, *Prog. Oceanogr.*, doi:10.1016/j.pocean.2015.01.010, 2015.
- 25 Baar, H.J.W. de, Timmermans, K., Laan, P., De Porto, H., Ober, S., Blom, J., Bakker, M., Schilling, J., Sarthou, G., Smit, M., et al.: Titan: A new facility for ultraclean sampling of trace elements and isotopes in the deep oceans in the international GEOTRACES program, *Mar. Chem.*, 111, 4–21, doi:10.1016/j.marchem.2007.07.009, 2008.
- Langen, P.J. von, Johnson, K., Coale, K., and Elrod, V.: Oxidation kinetics of manganese (II) in seawater at nanomolar concentrations, *Geochim. Cosmochim. Ac.*, 61, 4945–4954, doi:10.1016/S0016-7037(97)00355-4, 1997.
- 30 Elderfield, H.: Manganese fluxes to the oceans, *Mar. Chem.*, 4, 103–132, 1976.
- Fraústo da Silva, J. and Williams, R.: *The biological chemistry of the elements: the inorganic chemistry of life*, Oxford University Press, 2 edn., 2001.
- Froelich, P., Klinkhammer, G., Bender, M., Luedtke, N., Heath, G., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., and Maynard, V.: Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis, *Geochim. Cosmochim. Ac.*, 43, 1075–1090, doi:10.1016/0016-7037(79)90095-4, 1979.
- 35 Gaspar, P., Gregoris, Y., and Lefevre, J. M.: A simple eddy kinetic energy model for simulations of the ocean vertical mixing: Tests at station Papa and Long-Term Upper Ocean Study Site site, *J. Geophys. Res.*, 95, 16,179–16,193, doi:10.1029/JC095iC09p16179, 1990.

- Gent, P. and McWilliams, J.: Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.*, 20, 150–155, doi:10.1175/1520-0485(1990)020<0150:MIOCM>2.0.CO;2, 1990.
- German, C. R., Casciotti, K. A., Dutay, J.-C., Heimbürger, L. E., Jenkins, W. J., Measures, C. I., Mills, R. A., Obata, H., Schlitzer, R., Tagliabue, A., Turner, D. R., and Whitby, H.: Hydrothermal impacts on trace element and isotope ocean biogeochemistry, 374, doi:10.1098/rsta.2016.0035, part of the special issue *Biological and climatic impacts of ocean trace element chemistry*, 2016.
- 5 Gerringa, L., Rijkenberg, M., Schoemann, V., Laan, P., and de Baar, H.: Organic complexation of iron in the West Atlantic Ocean, *Mar. Chem.*, 177, Part 3, 434–446, doi:10.1016/j.marchem.2015.04.007, cycles of metals and carbon in the oceans—A tribute to the work stimulated by Hein de Baar, 2015.
- Guieu, C., Duce, R., and Arimoto, R.: Dissolved input of manganese to the ocean: Aerosol source, *J. Geophys. Res. Atmos.*, 99, 18 789–18, doi:10.1029/94JD01120, 1994.
- 10 Hauglustaine, D., Hourdin, F., Jourdain, L., Filiberti, M.-A., Walters, S., Lamarque, J.-F., and Holland, E.: Interactive chemistry in the Laboratoire de Météorologie Dynamique general circulation model, *J. Geophys. Res.: Atmospheres*, 109, D04 314, doi:10.1029/2003JD003957, 2004.
- Hayes, C. T., Anderson, R. F., Fleisher, M. Q., Lam, P. J., Ohnemus, D. C., Huang, K.-F., Robinson, L. F., Lu, Y., Cheng, H., Edwards, R. L., and Moran, S. B.: Intensity of Th and Pa scavenging partitioned by particle chemistry in the North Atlantic Ocean, *Mar. Chem.*, 15 doi:10.1016/j.marchem.2015.01.006, 2015.
- Homoky, W. B., Weber, T., Berelson, W. M., Conway, T. M., Henderson, G. M., van Hulten, M., Jeandel, C., Severmann, S., and Tagliabue, A.: Quantifying trace element and isotope fluxes at the ocean–sediment boundary: a review, 374, doi:10.1098/rsta.2016.0246, part of the special issue *Biological and climatic impacts of ocean trace element chemistry*, 2016.
- Hulten, M.M.P. van: Aluminium and Manganese in the West Atlantic Ocean, Ph.D. thesis, University of Groningen, <http://irs.ub.rug.nl/ppn/384091636>, printed version available on request to the author, 2014.
- 20 Hulten, M.M.P. van, Sterl, A., Dutay, J.-C., Tagliabue, A., Gehlen, M., Baar, H.J.W. de, and Middag, R.: Aluminium in an ocean general circulation model compared with the West Atlantic GEOTRACES cruises, *J. Mar. Syst.*, 126, 3–23, doi:10.1016/j.jmarsys.2012.05.005, traces and Tracers: Selected papers from the Joint Liège Colloquium on Ocean Dynamics – Bonus-GoodHope – GEOTRACES meeting, 2013.
- 25 Hulten, M.M.P. van, Sterl, A., Middag, R., de Baar, H., Gehlen, M., Dutay, J.-C., and Tagliabue, A.: On the effects of circulation, sediment resuspension and biological incorporation by diatoms in an ocean model of aluminium, *Biogeosci.*, 11, 3757–3779, doi:10.5194/bg-11-3757-2014, 2014.
- Hydes, D., Statham, P., and Burton, J.: A vertical profile of dissolved trace metals (Al, Cd, Cu, Mn, Ni) over the median valley of the mid Atlantic ridge, 43° N, *Sci. Total Environ.*, 49, 133–145, doi:10.1016/0048-9697(86)90236-6, 1986.
- 30 Jeandel, C.: Overview of the mechanisms that could explain the “Boundary Exchange” at the land–ocean contact, *Phil. Trans. R. Soc. A: Mathematical, Physical and Engineering Sciences*, 374, doi:10.1098/rsta.2015.0287, <http://rsta.royalsocietypublishing.org/content/374/2081/20150287>, part of the special issue *Biological and climatic impacts of ocean trace element chemistry*, 2016.
- Jeandel, C., van der Loeff, M. R., Lam, P. J., Roy-Barman, M., Sherrell, R. M., Kretschmer, S., German, C., and Dehairs, F.: What did we learn about ocean particle dynamics in the GEOSECS–JGOFS era?, *Progress in Oceanography*, 133, 6–16, doi:10.1016/j.pocean.2014.12.018,
- 35 GEOTRACES Synthesis and Modeling: The role of particles in the marine biogeochemical cycles of trace elements and their isotopes, 2015.
- Jickells, T.: Atmospheric inputs of metals and nutrients to the oceans, *Mar. Chem.*, 48, 199–214, doi:10.1016/0304-4203(95)92784-P, 1995.

- Johnson, K., Coale, K., Berelson, W., and Michael Gordon, R.: On the formation of the manganese maximum in the oxygen minimum, *Geochim. Cosmochim. Ac.*, 60, 1291–1299, doi:10.1016/0016-7037(96)00005-1, 1996.
- Johnson, K., Elrod, V., Fitzwater, S., Plant, J., Boyle, E., Bergquist, B., Bruland, K., Aguilar-Islas, A., Buck, K., Lohan, M., et al.: Developing standards for dissolved iron in seawater, *Eos, Transactions American Geophysical Union*, 88, 131–132, doi:10.1029/2007EO110003, 5 2007.
- Jong, J. de, Boyé, M., Gelado-Caballero, M., Timmermans, K., Veldhuis, M., Nolting, R., van den Berg, C., and de Baar, H.: Inputs of iron, manganese and aluminium to surface waters of the Northeast Atlantic Ocean and the European continental shelf, *Mar. Chem.*, 107, 120–142, doi:10.1016/j.marchem.2007.05.007, 2007.
- Kawagucci, S., Okamura, K., Kiyota, K., Tsunogai, U., Sano, Y., Tamaki, K., and Gamo, T.: Methane, manganese, and helium-3 in newly discovered hydrothermal plumes over the Central Indian Ridge, 18°–20° S, *Geochemistry, Geophysics, Geosystems*, 9, 10 10.1029/2008GC002082, 2008.
- Klinkhammer, G. and Bender, M.: The distribution of manganese in the Pacific Ocean, *Earth Planet. Sc. Lett.*, 46, 361–384, doi:10.1016/0012-821X(80)90051-5, 1980.
- Klinkhammer, G., Bender, M., and Weiss, R.: Hydrothermal manganese in the Galapagos Rift, *Nature*, 269, 319–320, doi:10.1038/269319a0, 15 1977.
- Klinkhammer, G., Rona, P., Greaves, M., and Elderfield, H.: Hydrothermal manganese plumes in the Mid-Atlantic Ridge rift valley, *Nature*, 314, 727–731, doi:10.1038/314727a0, 1985.
- Klinkhammer, G., Chin, C., Keller, R., Daehlmann, A., Sahling, H., Sarthou, G., Petersen, S., Smith, F., and Wilson, C.: Discovery of new hydrothermal vent sites in Bransfield Strait, Antarctica, *Earth Planet. Sc. Lett.*, 193, 395–407, doi:10.1016/S0012-821X(01)00536-2, 2001.
- 20 Klunder, M., Laan, P., Middag, R., de Baar, H., and van Ooijen, J.: Dissolved iron in the Southern Ocean (Atlantic sector), *Deep-Sea Res. Pt II*, 58, 2678–2694, doi:10.1016/j.dsr2.2010.10.042, 2011.
- Kriest, I. and Oschlies, A.: Numerical effects on organic-matter sedimentation and remineralization in biogeochemical ocean models, *Ocean Model.*, 39, 275–283, doi:10.1016/j.ocemod.2011.05.001, 2011.
- Landing, W. and Bruland, K.: Manganese in the north Pacific, *Earth Planet. Sc. Lett.*, 49, 45–56, doi:10.1016/0012-821X(80)90149-1, 1980.
- 25 Landing, W. and Bruland, K.: The contrasting biogeochemistry of iron and manganese in the Pacific Ocean, *Geochim. Cosmochim. Ac.*, 51, 29–43, doi:10.1016/0016-7037(87)90004-4, 1987.
- Landing, W., Haraldsson, C., and Paxeus, N.: Vinyl polymer agglomerate based transition metal cation-chelating ion-exchange resin containing the 8-hydroxyquinoline functional group, *Anal. Chem.*, 58, 3031–3035, doi:10.1021/ac00127a029, 1986.
- Landing, W., Cutter, G., Dalziel, J., Flegal, A., Powell, R., Schmidt, D., Shiller, A., Statham, P., Westerlund, S., and Resing, J.: Analytical intercomparison results from the 1990 Intergovernmental Oceanographic Commission open-ocean baseline survey for trace metals: Atlantic Ocean, *Mar. Chem.*, 49, 253–265, 1995.
- 30 Lavelle, J. W., Cowen, J. P., and Massoth, G. J.: A model for the deposition of hydrothermal manganese near ridge crests, *J. Geophys. Res.: Oceans*, 97, 7413–7427, doi:10.1029/92JC00406, 1992.
- Leggett, R. W. and Williams, L. R.: A reliability index for models, *Ecological Modelling*, 13, 303–312, doi:10.1016/0304-3800(81)90034-X, 35 1981.
- Lengaigne, M., Madec, G., Menkes, C., and Alory, G.: Effect of isopycnal diffusion in the tropical Pacific Ocean, *J. Geophys. Res.*, 108, 10.1029/2002JC001704, doi:10.1029/2002JC001704, 2003.

- Lewis, B. and Luther, G.: Processes controlling the distribution and cycling of manganese in the oxygen minimum zone of the Arabian Sea, *Deep-Sea Res. Pt II*, 47, 1541–1561, doi:10.1016/S0967-0645(99)00153-8, 2000.
- Li, Y.-H., Bischoff, J., and Mathieu, G.: The migration of manganese in the Arctic basin sediment, *Earth Planet. Sc. Lett.*, 7, 265–270, doi:10.1016/0012-821X(69)90063-6, 1969.
- Luther, G. W., Madison, A. S., Mucci, A., Sundby, B., and Oldham, V. E.: A kinetic approach to assess the strengths of ligands bound to soluble Mn(III), *Mar. Chem.*, 173, 93–99, doi:10.1016/j.marchem.2014.09.006, SCOR WG 139: Organic Ligands – A Key Control on Trace Metal Biogeochemistry in the Ocean, 2015.
- Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, *Climate Dynamics*, 12, 381–388, doi:10.1007/BF00211684, 1996.
- Madec, G. and the NEMO team: NEMO ocean engine, Note du Pole de Modélisation, Institut Pierre-Simon Laplace, 2016.
- 10 Madison, A., Tebo, B., Mucci, A., Sundby, B., and Luther, G.: Abundant Porewater Mn(III) Is a Major Component of the Sedimentary Redox System, *Science*, 341, 875–878, doi:10.1126/science.1241396, 2013.
- Martin, J. and Knauer, G.: VERTEX, *Deep-Sea Res. Pt A*, 30, 411–425, 1983.
- Mawji, E., Schlitzer, R., Dodas, E. M., Abadie, C., Abouchami, W., Anderson, R. F., Baars, O., Bakker, K., Baskaran, M., Bates, N. R., Bluhm, K., Bowie, A., Bown, J., Boye, M., Boyle, E. A., Branell, P., Bruland, K. W., Brzezinski, M. A., Bucciarelli, E., Buesseler, K., 15 Butler, E., Cai, P., Cardinal, D., Casciotti, K., Chaves, J., Cheng, H., Chever, F., Church, T. M., Colman, A. S., Conway, T. M., Croot, P. L., Cutter, G. A., de Baar, H. J., de Souza, G. F., Dehairs, F., Deng, F., Dieu, H. T., Dulaquais, G., Echevoyen-Sanz, Y., Edwards, R. L., Fahrbach, E., Fitzsimmons, J., Fleisher, M., Frank, M., Friedrich, J., Fripiat, F., Galer, S. J., Gamo, T., Solsona, E. G., Gerringa, L. J., Godoy, J. M., Gonzalez, S., Grosstefan, E., Hatta, M., Hayes, C. T., Heller, M. I., Henderson, G., Huang, K.-F., Jeandel, C., Jenkins, W. J., John, S., Kenna, T. C., Klunder, M., Kretschmer, S., Kumamoto, Y., Laan, P., Labatut, M., Lacan, F., Lam, P. J., Lannuzel, D., 20 le Moigne, F., Lechtenfeld, O. J., Lohan, M. C., Lu, Y., Masqué, P., McClain, C. R., Measures, C., Middag, R., Moffett, J., Navidad, A., Nishioka, J., Noble, A., Obata, H., Ohnemus, D. C., Owens, S., Planchon, F., Pradoux, C., Puigcorb , V., Quay, P., Radic, A., Rehk mper, M., Remenyi, T., Rijkenberg, M. J., Rintoul, S., Robinson, L. F., Roeske, T., Rosenberg, M., van der Loeff, M. R., Ryabenko, E., Saito, M. A., Roshan, S., Salt, L., Sarthou, G., Schauer, U., Scott, P., Sedwick, P. N., Sha, L., Shiller, A. M., Sigman, D. M., Smethie, W., Smith, G. J., Sohrin, Y., Speich, S., Stichel, T., Stutsman, J., Swift, J. H., Tagliabue, A., Thomas, A., Tsunogai, U., Twining, B. S., van Aken, 25 H. M., van Heuven, S., van Ooijen, J., van Weerlee, E., Venchiarutti, C., Voelker, A. H., Wake, B., Warner, M. J., Woodward, E. M. S., Wu, J., Wyatt, N., Yoshikawa, H., Zheng, X.-Y., Xue, Z., Zieringer, M., and Zimmer, L. A.: The GEOTRACES Intermediate Data Product 2014, *Mar. Chem.*, 177, doi:10.1016/j.marchem.2015.04.005, 2015.
- McCave, I.: Vertical flux of particles in the ocean, *Deep-Sea Res. and Oceanographic Abstracts*, 22, 491–502, doi:10.1016/0011-7471(75)90022-4, 1975.
- 30 McManus, J., Berelson, W. M., Severmann, S., Johnson, K. S., Hammond, D. E., Roy, M., and Coale, K. H.: Benthic manganese fluxes along the Oregon–California continental shelf and slope, *Cont. Shelf Res.*, 43, 71–85, doi:10.1016/j.csr.2012.04.016, 2012.
- Means, J. L., Crerar, D. A., Borcsik, M. P., and Duguid, J. O.: Radionuclide adsorption by manganese oxides and implications for radioactive waste disposal, *Nature*, 274, 44–47, doi:10.1038/274044a0, 1978.
- Mendez, J., Guieu, C., and Adkins, J.: Atmospheric input of manganese and iron to the ocean: Seawater dissolution experiments with Saharan and North American dusts, *Mar. Chem.*, 120, 34–43, doi:10.1016/j.marchem.2008.08.006, 2010.
- 35 Middag, R., de Baar, H., Laan, P., and Bakker, K.: Dissolved aluminium and the silicon cycle in the Arctic Ocean, *Mar. Chem.*, 115, 176–195, doi:10.1016/j.marchem.2009.08.002, 2009.

- Middag, R., de Baar, H., Laan, P., Cai, P., and van Ooijen, J.: Dissolved manganese in the Atlantic sector of the Southern Ocean, *Deep-Sea Res. Pt II*, 58, 2661–2677, doi:10.1016/j.dsr2.2010.10.043, 2011a.
- Middag, R., de Baar, H., Laan, P., and Klunder, M.: Fluvial and hydrothermal input of manganese into the Arctic Ocean, *Geochim. Cosmochim. Ac.*, 75, 2393–2408, doi:10.1016/j.gca.2011.02.011, 2011b.
- 5 Middag, R., de Baar, H., Laan, P., and Huhn, O.: The effects of continental margins and water mass circulation on the distribution of dissolved aluminum and manganese in Drake Passage, *J. Geophys. Res.*, 117, C01 019, doi:10.1029/2011JC007434, 2012.
- Middag, R., de Baar, H., Klunder, M., and Laan, P.: Fluxes of dissolved aluminum and manganese to the Weddell Sea and indications for manganese co-limitation, *Limnol. Oceanogr.*, 58, 287–300, doi:10.4319/lo.2013.58.1.0287, 2013.
- Middag, R., Hulthen, M.M.P. van, Aken, H.M. van, Rijkenberg, M., Gerringa, L., Laan, P., and Baar, H.J.W. de: Dissolved aluminium in the
10 ocean conveyor of the West Atlantic Ocean: Effects of the biological cycle, scavenging, sediment resuspension and hydrography, *Mar. Chem.*, 177, Part 1, 69–86, doi:10.1016/j.marchem.2015.02.015, biogeochemistry of trace elements and their isotopes, 2015a.
- Middag, R., Séférian, R., Conway, T., John, S., Bruland, K., and de Baar, H.: Intercomparison of dissolved trace elements at the Bermuda Atlantic Time Series station, *Mar. Chem.*, 177, Part 3, 476–489, doi:10.1016/j.marchem.2015.06.014, cycles of metals and carbon in the oceans—A tribute to the work stimulated by Hein de Baar, 2015b.
- 15 Middelburg, J. J., Soetaert, K., Herman, P. M. J., and Heip, C. H. R.: Denitrification in marine sediments: A model study, *Global Biogeochem. Cy.*, 10, 661–673, doi:10.1029/96GB02562, 1996.
- Milne, A. and Landing, W.: CLIVAR P16: Dissolved manganese in the upper 1000 m of the Pacific Ocean, unpublished data.
- Milne, A., Landing, W., Bizimis, M., and Morton, P.: Determination of Mn, Fe, Co, Ni, Cu, Zn, Cd and Pb in seawater using high resolution magnetic sector inductively coupled mass spectrometry (HR-ICP-MS), *Analytica Chimica Acta*, 665, 200–207,
20 doi:10.1016/j.aca.2010.03.027, 2010.
- Murray, J. W.: The interaction of metal ions at the manganese dioxide-solution interface, *Geochim. Cosmochim. Ac.*, 39, 505–519, doi:10.1016/0016-7037(75)90103-9, 1975.
- Nealson, K. H.: The Manganese-Oxidizing Bacteria, in: *The Prokaryotes: Volume 5: Proteobacteria: Alpha and Beta Subclasses*, edited by Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., pp. 222–231, Springer New York, New York, NY,
25 doi:10.1007/0-387-30745-1_11, 2006.
- Noble, A. E., Lamborg, C. H., Ohnemus, D. C., Lam, P. J., Goepfert, T. J., Frame, C. H., Casciotti, K. L., DiTullio, G. R., Jennings, J. C., Saito, M. A., et al.: Basin-scale inputs of cobalt, iron, and manganese from the Benguela-Angola front to the South Atlantic Ocean, *Limnol. Oceanogr.*, 2012.
- Passow, U. and De La Rocha, C. L.: Accumulation of mineral ballast on organic aggregates, *Global Biogeochem. Cy.*, 20,
30 doi:10.1029/2005GB002579, 2006.
- Petersen, A.: *Simulating nature*, Ph.D. thesis, 2006.
- Raven, J. A.: Predictions of Mn and Fe use efficiencies of phototrophic growth as a function of light availability for growth and of C assimilation pathway, *New Phytologist*, 116, 1–18, doi:10.1111/j.1469-8137.1990.tb00505.x, 1990.
- Reid, D. F., Key, R. M., and Schink, D. R.: Radium, thorium, and actinium extraction from seawater using an improved manganese-oxide-coated fiber, *Earth Planet. Sc. Lett.*, 43, 223–226, doi:10.1016/0012-821X(79)90205-X, 1979.
- 35 Resing, J. A., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., and Tagliabue, A.: Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean, *Nature*, 523, 200–203, doi:10.1038/nature14577, 2015.

- Rijkenberg, M., Middag, R., Laan, P., Gerringa, L., Aken, H.M. van, Schoemann, V., Jong, J.T.M. de, Jong, J. de, and Baar, H.J.W. de: The Distribution of Dissolved Iron in the West Atlantic Ocean, *PloS one*, 9, e101323, doi:10.1371/journal.pone.0101323, 2014.
- Rijkenberg, M. J., de Baar, H. J., Bakker, K., Gerringa, L. J., Keijzer, E., Laan, M., Laan, P., Middag, R., Ober, S., van Ooijen, J., Ossebaar, S., van Weerlee, E. M., and Smit, M. G.: "PRISTINE", a new high volume sampler for ultraclean sampling of trace metals and isotopes, *Mar. Chem.*, 177, Part 3, 501–509, doi:10.1016/j.marchem.2015.07.001, cycles of metals and carbon in the oceans—A tribute to the work stimulated by Hein de Baar, 2015.
- Roy-Barman, M.: Modelling the effect of boundary scavenging on Thorium and Protactinium profiles in the ocean, *Biogeosci.*, 6, 3091–3107, doi:10.5194/bg-6-3091-2009, 2009.
- Roy-Barman, M., Jeandel, C., Souhaut, M., Rutgers van der Loeff, M., Voege, I., Leblond, N., and Freydier, R.: The influence of particle composition on thorium scavenging in the NE Atlantic ocean (POMME experiment), *Earth Planet. Sc. Lett.*, 240, 681–693, doi:10.1016/j.epsl.2005.09.059, 2005.
- Rüth, C., Well, R., and Roether, W.: Primordial ^3He in South Atlantic deep waters from sources on the Mid-Atlantic Ridge, *Deep-Sea Res. Pt I*, 47, 1059–1075, doi:10.1016/S0967-0637(99)00077-1, 2000.
- Saager, P. M., de Baar, H. J., and Burkill, P. H.: Manganese and iron in Indian Ocean waters, *Geochim. Cosmochim. Ac.*, 53, 2259–2267, doi:10.1016/0016-7037(89)90348-7, 1989.
- Saager, P. M., de Baar, H. J., de Jong, J. T., Nolting, R. F., and Schijf, J.: Hydrography and local sources of dissolved trace metals Mn, Ni, Cu, and Cd in the northeast Atlantic Ocean, *Mar. Chem.*, 57, 195–216, doi:10.1016/S0304-4203(97)00038-8, 1997.
- Saito, M. A., Moffett, J. W., and DiTullio, G. R.: Cobalt and nickel in the Peru upwelling region: A major flux of labile cobalt utilized as a micronutrient, *Global Biogeochem. Cy.*, 18, doi:10.1029/2003GB002216, 2004.
- Sander, S. and Koschinsky, A.: Metal flux from hydrothermal vents increased by organic complexation, *Nat. Geosci.*, 4, 145–150, doi:10.1038/ngeo1088, 2011.
- Sarmiento, J. and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton University Press, 2006.
- Slemons, L., Murray, J., Resing, J., Paul, B., and Dutrieux, P.: Western Pacific coastal sources of iron, manganese, and aluminum to the Equatorial Undercurrent, *Global Biogeochem. Cy.*, 24, doi:10.1029/2009GB003693, 2010.
- Slomp, C., Malschaert, J., Lohse, L., and Van Raaphorst, W.: Iron and manganese cycling in different sedimentary environments on the North Sea continental margin, *Cont. Shelf Res.*, 17, 1083–1117, doi:10.1016/S0278-4343(97)00005-8, 1997.
- Statham, P. and Burton, J.: Dissolved manganese in the North Atlantic Ocean, 0–35‰N, *Earth Planet. Sc. Lett.*, 79, 55–65, doi:10.1016/0012-821X(86)90040-3, 1986.
- Statham, P., Yeats, P., and Landing, W.: Manganese in the eastern Atlantic Ocean, *Mar. Chem.*, 61, 55–68, 1998.
- Stone, A. and Morgan, J.: Kinetics of Chemical Transformation in the Environment, in: *Aquatic chemical kinetics*, edited by Stumm, W., vol. 2, chap. 1, pp. 1–42, John Wiley and Sons, 1990.
- Stow, C., Jolliff, J., Jr., D. M., Doney, S., Allen, J., Friedrichs, M., Rose, K., and Wallhead, P.: Skill assessment for coupled biological/physical models of marine systems, *J. Mar. Syst.*, 76, 4–15, doi:10.1016/j.jmarsys.2008.03.011, 2009.
- Sunda, W. and Huntsman, S.: Effect of sunlight on redox cycles of manganese in the southwestern Sargasso Sea, *Deep-Sea Res. Pt A*, 35, 1297–1317, doi:10.1016/0198-0149(88)90084-2, 1988.
- Sunda, W. and Huntsman, S.: Photoreduction of manganese oxides in seawater, *Mar. Chem.*, 46, 133–152, doi:10.1016/0304-4203(94)90051-5, 1994.
- Sundby, B. and Silverberg, N.: Manganese fluxes in the benthic boundary layer, *Limnol. Oceanogr.*, pp. 372–381, 1985.

- Sundby, B., Silverberg, N., and Chesselet, R.: Pathways of manganese in an open estuarine system, *Geochim. Cosmochim. Ac.*, 45, 293–307, doi:10.1016/0016-7037(81)90240-4, 1981.
- 1045 Tagliabue, A. and Völker, C.: Towards accounting for dissolved iron speciation in global ocean models, *Biogeosci.*, 8, 3025–3039, doi:10.5194/bg-8-3025-2011, 2011.
- Tagliabue, A., Bopp, L., Dutay, J.-C., Bowie, A., Chever, F., Jean-Baptiste, P., Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G., Aumont, O., Gehlen, M., and Jeandel, C.: Hydrothermal contribution to the oceanic dissolved iron inventory, *Nat. Geosci.*, 3, 252–256, doi:10.1038/NGEO818, 2010.
- 1050 Tebo, B., Johnson, H., McCarthy, J., and Templeton, A.: Geomicrobiology of manganese (II) oxidation, *Trends in Microbiology*, 13, 421–428, doi:10.1016/j.tim.2005.07.009, 2005.
- Tonkin, J. W., Balistrieri, L. S., and Murray, J. W.: Modeling sorption of divalent metal cations on hydrous manganese oxide using the diffuse double layer model, *Applied Geochemistry*, 19, 29–53, doi:10.1016/S0883-2927(03)00115-X, 2004.
- Trefry, J. H. and Presley, B. J.: Manganese fluxes from Mississippi Delta sediments, *Geochim. Cosmochim. Ac.*, 46, 1715–1726, doi:10.1016/0016-7037(82)90112-0, 1982.
- 1055 Twining, B. and Baines, S.: The Trace Metal Composition of Marine Phytoplankton, *Annu. Rev. Mar. Sci.*, 5, doi:10.1146/annurev-marine-121211-172322, 2013.
- Vichi, M. and Masina, S.: Skill assessment of the PELAGOS global ocean biogeochemistry model over the period 1980–2000, *Biogeosci.*, 6, 2333–2353, doi:10.5194/bg-6-2333-2009, 2009.
- Vu, H. and Sohrin, Y.: Diverse stoichiometry of dissolved trace metals in the Indian Ocean, *Sci. Rep.*, 3, doi:10.1038/srep01745, 2013.
- 1060 Wedepohl, K.: The composition of the continental crust, *Geochim. Cosmochim. Ac.*, 59, 1217–1232, doi:10.1016/0016-7037(95)00038-2, 1995.
- Wu, J., Roshan, S., and Chen, G.: The distribution of dissolved manganese in the tropical-subtropical North Atlantic during US GEOTRACES 2010 and 2011 cruises, *Mar. Chem.*, doi:10.1016/j.marchem.2014.08.007, 2014.
- Yamagata, N. and Iwashima, K.: Monitoring of Sea-Water for Important Radioisotopes Released by Nuclear Reactors, *Nature*, 200, 52–52, doi:10.1038/200052a0, 1963.
- 1065 Yeats, P., Sundby, B., and Bewers, J.: Manganese recycling in coastal waters, *Mar. Chem.*, 8, 43–55, doi:10.1016/0304-4203(79)90031-8, 1979.






Figure 1. $[Mn_{diss}]$ at the 16 June 2010 (NL) and the 19 November 2011 (US) occupations of the crossover site of the Bermuda Atlantic Time Series (BATS) station. There is good agreement between the two sets of samples, as well as the two different analytical methods, the shipboard FIA by NIOZ for the NL2010 samples, and the lab ICP-MS at UCSC for both the NL2010 and the US2011 samples. Each lab had its own independently prepared lab standards, further confirming the overall accuracy. Analyses by Rob Middag at NIOZ in 2010 and in 2011 at UCSC.



(a)



(b)

Figure 2. The correlation between the two methods of analysis for the determination of $[Mn_{diss}]$ at GA02 (shipboard and laboratory mass spectrometry): (a) at Bermuda; (b) all 55 West Atlantic stations.




Figure 3. Model scheme: biology in green; redox and scavenging in blue; circulation and mixing (dynamics) in grey; and the sources dust, rivers, hydrothermal and sediment are in light brown, light blue, dark brown and red, respectively. Mn_{diss} is the dissolved and Mn_{ox} the oxidised Mn.




Figure 4. Sources of Mn to the ocean: effective Mn_{diss} input flux Φ . Three-dimensional fields are vertically integrated, such that dimensions are molar fluxes. Ranges vary between the different sources.




Figure 5. Transect (or expedition) names corresponding to station colours: **GIPY11** in the Arctic Ocean; **GIPY4** and **GIPY5** in the Atlantic sector of the Southern Ocean; **GI04** in the Indian Ocean; **GA02** in the West Atlantic Ocean; **GA03** in the North Atlantic Ocean; **VERTEX-4** in the North Pacific Ocean; **EUCFe** in the equatorial Pacific Ocean; **CLIVAR P16** in the Pacific Ocean; **GP16** in the South Pacific; **CoFeMUG** in the South Atlantic Ocean. See Table 5 for an overview with references and the number of observations.




Figure 6. Observations of $[Mn_{diss}]$ (nM) along the GEOTRACES GA02 transect in the West Atlantic Ocean. Dots are the locations of the measurements. Upper 500 m.




Figure 7. Observations of $[Mn_{diss}]$ (nM) along the GEOTRACES GA02 transect in the West Atlantic Ocean. Dots are the locations of the measurements. Below 500 m. The by red dashed lines denoted high dissolved iron concentrations are inferred from Rijkenberg et al. (2014, their Figure 3A). The by blue dashed lines denoted high dissolved aluminium concentrations are inferred from Middag et al. (2015a, their Fig. 3a). Note that the concentrations are generally much smaller than in the surface, and that the colour scale is different from that of Fig. 6.




Figure 8. $[Mn_{diss}]$ (nM) at four depth layers in the world ocean for the reference simulation (*Reference*) after 500 yr (annual average). Observations are presented as coloured dots; white is the land mask of the model grid.




Figure 9. $[Mn_{diss}]$ (nM) at the Zero-Meridian GIPY5_e and the West Atlantic GA02 GEOTRACES transects for *Reference* after 500 yr (annual average). Observations are presented as coloured dots.




Figure 10. $[Mn_{diss}]$ (nM) at the North Atlantic GA03 GEOTRACES transect for *Reference* after 600 yr (annual average). Observations are presented as coloured dots.




Figure 11. The effect of biological incorporation on $[Mn_{diss}]$ at four depths: $(Reference - NoBio)/NoBio$. Grey is the land mask of the model domain; the black contour is that of the real continental coasts.




Figure 12. Annual average of $[Mn_{diss}]$ (nM) at two depth in the world ocean for the simulation without a Mn biological cycle (*NoBio*) of year 500 after forking from *Reference*.




Figure 13. Result from the simulation with hydrothermal vent Mn input decreased and settling velocity decreased (*LowHydro*). Relative difference in $[\text{Mn}_{\text{diss}}]$ (%) between *Reference* and *LowHydro*. To represent the changes of much larger than 100 %, the scale is increased from +50 % upwards.




Figure 14. $[\text{Mn}_{\text{diss}}]$ (nM) from the simulation with hydrothermal vent Mn input decreased and settling velocity decreased (*LowHydro*) at the West Atlantic GA02 and Zero-Meridian Southern Ocean GIPY5_e GEOTRACES transects.




Figure 15. $[\text{Mn}_{\text{diss}}]$ (nM) at the West Atlantic GA02 and Zero-Meridian Southern Ocean GIPY5_e GEOTRACES transects. Simulation without threshold (*NoThreshold*).




Figure 16. $[Mn_{ox}]/[Mn_{diss}]$ in the Pacific Ocean at the VERTEX-IV station in the upper 110 m (left) and at full depth (right). Blue squares are observations, the green line is the *Reference* simulation, and the red dashed line is the ratio from *OxidThreshold*.