

## ***Interactive comment on “Yedoma Ice Complex of the Buor Khaya Peninsula (southern Laptev Sea)” by Lutz Schirrmeister et al.***

**Lutz Schirrmeister et al.**

lutz.schirrmeister@awi.de

Received and published: 3 August 2016

Due to a copying error at the completion of this manuscript, the conclusions have been replaced by Acknowledgements. The correct versions of conclusion and acknowledgment are the following.

**Conclusion** The studied Yedoma IC permafrost exposed at the western coast of the Buor Khaya Peninsula in the central Laptev Sea accumulated from about  $54.1 \pm 3.4$  kyr BP (IRSL age  $51.1 \pm 4.9$  kyr) until the onset of the Lateglacial to Holocene warming. The timing of IC formation as well as its cryolithological properties and temperature are similar to known Yedoma IC sequences in the Laptev Sea region and beyond. Three robust end-members (rEMs) represent fine silt, medium silt, and fine sand fractions in grain-size distribution data and support multiple transport and depositional processes

C1

during IC formation. Post-depositional refreezing processes during the MIS3 interstadial with relatively warm and moist summer conditions created unusually high solute concentrations in pore water. The OM characteristics of the Yedoma IC reflect typical continental tundra-steppe environments of western Beringia during the late Pleistocene. Eroding Yedoma IC deposits are one of the OM sources in the marine realm of the Laptev Shelf.

**Acknowledgments** The study presented here is part of the Helmholtz-Russia Joint Research Group. We gratefully acknowledge the German Federal Ministry of Education and Research (BMBF) for funding this study as part of the joint German-Russian research project CARBOPERM (03G0836B, 03G0836F). Moreover, we acknowledge the support by the European Research Council (Starting Grant #338335) and the Initiative and Networking Fund of the Helmholtz Association (#ERC-0013). We thank our colleagues who helped during fieldwork in 2014, especially Dmitry Mel'nichenko and colleagues from the Hydrobase Tiksi, and Waldemar Schneider from AWI Potsdam who greatly supported the logistics of this expedition. The analytical work in the AWI laboratories was expertly conducted by Ute Bastian, Dyke Scheidemann, Antje Eulenburg, Hanno Meyer, 10 and Lutz Schöncke. We are grateful to Marcus Richter (TU Dresden) and to Ingrid Stein (TU Bergakademie Freiberg) for IRSL analyses and to Jörg Erzinger, Sabine Schumann (GFZ Potsdam), and Torben Windirsch (University of Potsdam) for TS measurements. Finally, the paper benefited by constructive comments and English language correction from Candace O'Connor (Fairbanks Alaska).

---

Interactive comment on Biogeosciences Discuss., doi:10.5194/bg-2016-283, 2016.

C2