

Response to Reviewers (bg-2016-286)

Associate Editor Decision:

Publish subject to technical corrections (28 Feb 2017) by Akihiko Ito Comments to the Author: Dear Authors: Thank you for uploading the revised manuscript. I studied the manuscript and found that it was considerably improved in comparison with the previous one. Specifically, you replaced the word “long-term” by “multi-seasonal” and considered the random effect in the statistical analysis. I with this study carries valuable information to research community on the methane emission from a temperate, human-affected lake. Only one technical correction is needed: in the supplement, please add “S” to the table number (e.g., Table S1).

Answer: Changed as suggested.

Associate Editor Comments

Thank you for sending your detailed replies to the referee comments. The three referees gave insightful comments and in general evaluated your study positively. Although one referee did not find ‘strong relevance to Biogeosciences’ in your results and discussion, others concluded that the study falls within the journal scope. Importantly, the three referees claimed that this is not a ‘long-term’ study, because this study was conducted only for four years. Also, they made many critical comments on the spatial and temporal representativeness of measurements (e.g., monthly interval at three sites). Therefore, you should revise the manuscript very carefully, such that any overstatement on observational representativeness is avoided and possible uncertainties are fully discussed. Additionally, referees made a recommendation about the statistical analysis, such that inter-site difference should be treated as a random effect in mixed effects models. As you replied, although the site effect was not a critical factor, you should update the manuscript by including the

statistical re-analysis results. I'm looking forward to receiving a revised manuscript with major revision.

Answer: Thank you very much! We have considered all the suggestions carefully in the revised manuscript. We changed the word “long-term” to “multi-seasonal”, explained the sampling representativeness, and added the random effect in the statistical analysis result in the revised version.

Reviewer 1

General comments

The manuscript reports on the temporal patterns of methane (CH₄) efflux in the largest lake in China and the various factors that influence these fluxes over different timescales. CH₄ efflux was slightly greater than other lakes with an area greater than 1km, but was comparable to that found in tropical lakes. The variables best explaining variation on CH₄ efflux was timescale dependent but, overall, temperature was important over seasonal scales and wind speed on a bihourly scale. The paper is well written despite a few grammatical errors. As the authors point out, there is a lack of data explaining CH₄ effluxes in this region and also in larger, nonalpine lakes more generally. As such, the results from this paper will add to the limited understanding of CH₄ dynamics in these lake types. However, I cannot recommend that this paper be published in its current form. I have major issues with a) the premise of the paper, b) some overreaching statements that are made, and c) the statistical approaches used - all have major implications for the generalisation of the results. It is difficult for me to assess the results and technical aspects of this study until statistical changes are made.

Answer: We thank the reviewer so much for the constructive comments and suggestions. We have considered all the comments and suggestions carefully in revising the manuscript.

1. One major concern is that the study was undertaken in a very small area (three sites with 20km of each other) even though the lake is the largest in China by area (3283sq km). Further, the study sites are situated in a section of the lake that appears to be relatively confined. There is nothing wrong with the site selection. However, the authors cannot make statements about the whole lake because they don't know if the spatial and temporal patterns of CH₄ vary the same way across the lake. They need to qualify in all statements that the research was undertaken in one small section of the lake. It is not a study of CH₄ effluxes from Poyang Lake, but it is a study of CH₄ effluxes from one section of Poyang Lake.

Answer: We agree with the Reviewer that the CH₄ efflux in the Poyang Lake has a large spatial variation as evidenced in our previous study which examined the spatial variations of greenhouse gas effluxes (including CH₄) over the lake with 44 sampling locations. The current study focuses on the temporal dynamics of CH₄ efflux. We chose the 3 sites to roughly represent the average CH₄ efflux of the whole lake based on the results of our previous study (Liu et al., 2013).

Therefore, our results reflect the general situation of the lake.

2. Another major concern is a statistical one. The authors use average values from three different locations in Poyang Lake for all analyses. The justification for this was to 'minimize the effect of the spatial variation of CH₄ efflux on the temporal dynamics of the efflux'. However I suspect the main motivation for doing this was because the environmental variables were only collected at one location (it is not clear where the environmental variables were collected). Was this the case? Given that CH₄ was only

measured in three locations of the lake, surely the degree of variation between them is very important to a) understand and/or b) account for in statistical analyses. The authors should re-analyse their results in one of the following ways: – treat each study site as a random effect in mixed effects models so that variation among the three sites is taken into account when investigating the annual, seasonal, and diurnal variation, as well as the relationships with measured explanatory variables. Including site as a random effect would enable the researchers to make more general statements about CH₄ fluxes from Poyang Lake – this is just common practice these days and should be incorporated into the study design / statistical analyses. A random effect for site effectively means that these study sites are a random sample of all potential sites in the lake – this is where the generalisability comes in. Please see Section 8.1.1 (Types of predictor variables (factors)) in Quinn & Keough (2002; Experimental Design & Data Analysis for Biologists) or another similar book for information about mixed effects models and random and fixed factors. – split the analyses into two parts. The first analysis will not average the three study sites prior to the analysis and investigate the spatial and temporal patterns in CH₄ among them. The second analysis could average the study sites (still preferably treat study site as a random factor) and relate this to the measured explanatory variables.

Answer: We actually collected environmental variables at each site except water level which was monitored at the Xingzi Hydrological Station. We appreciate the Reviewers' suggestion (also see Reviewer 2's comments) and re-analyzed the data by treating the site as a random effect. We found that the site effect was not statistically significant including seasonal and diel CH₄ effluxes over the 4-year period (Table S2, S3 in the supplementary material). We also re-analyzed our data for each site and found that the differences among the 3 sites were minor

with the 4-year mean of $0.53 \text{ mmol m}^{-2} \text{ day}^{-1}$, $0.55 \text{ mmol m}^{-2} \text{ day}^{-1}$, and $0.54 \text{ mmol m}^{-2} \text{ day}^{-1}$ respectively. In addition, we found that the seasonal patterns of CH_4 effluxes at three sites were similar and also in line with the seasonal pattern averaged over the 3 sites. Nevertheless, in the stepwise multiple regressions analyses, the same environmental variables were selected in the final model for each individual site as for the average of the 3 sites with the regression coefficients slightly different (Table S5), but not statistically significant ($p > 0.20$). So we have focused on presenting the site-averaged CH_4 efflux and its dynamics due to the length limitation of the paper. But we explained the site effect on CH_4 effluxes in the revised version.

3. One more major concern is the notion that this is a long-term study. 4 years is not long term. Remove all reference to this study being long term, including the second sentence of the Abstract which introduces the idea that this research is filling the knowledge gap around the lack of long term research on CH_4 fluxes. Instead, the authors should frame this ‘knowledge gap’ around the lack of multi-seasonal investigations into CH_4 effluxes – this is exactly what this paper addresses.

Answer: We agree and thank you so much for the constructive suggestion. We removed the phrase “long term” and changed the tones accordingly in the text during the version. In addition, we have focused on multi-seasonal investigations of CH_4 effluxes as suggested.

Specific comments

1. Line 18. It is stated continuous measurements of CH_4 efflux was measured, but measurements were not continuous. Monthly measurements were made. Change all reference to continuous measurements in the manuscript to monthly measurements.

Answer: Changed as suggested in the revised version.

2. Line 121-124. Are these parameters an average of the entire lake or for a specific location? Please specify.

Answer: These parameters are averages of the entire lake. We added the information in the revised version (Page 7/lines 135-138).

3. Section 2.3. Environmental variables. Where were the environmental variables collected from? Where samples collected at each of the three study sites and then averaged or from just one site? This information is very important.

Answer: The environmental variables were measured at each of the three study sites and then averaged over the sites except water level which was monitored at a single hydrological station (national class station). We added more details of the environmental variables in the revised version (Page 11/lines 221-222).

4. Line 331-332. This concluding sentence only relates to the first sentence of this paragraph and does not relate or link to the remaining text in the paragraph. This sentence should only be left if a re-working of the paragraph better supports this argument.

Answer: We deleted the sentence because it is not the main point of the paragraph.

5. Line 337-341. An argument is made that this study has lower diurnal variation in CH₄ efflux than other studies and this may be due to differences in sample size in other studies. I would think that more frequent sampling would in-fact lead to more variation. The authors need to report on how much diurnal variability in CH₄ efflux there was among the study sites.

Answer: We agree with the Reviewer that the diurnal range (maximum – minimum) of CH₄ efflux depends on sample size and sampling frequency, which

makes the comparison with other lakes less meaningful. Therefore, we deleted the discussion on comparing the ranges of CH₄ effluxes in different lakes, which are not the main focus of the current study (Also see the reviewer 2' comments).

Technical comments

1. Line 163-182. The description of how CH₄ efflux due to ebullition is very confusing and long.

Answer: We rewrote this part to clarify the confusion in the revised version (Page 10/lines 190-200).

2. From Line 125, where the ebullition and diffusive fluxes are introduced, I would suggest briefly describing how, or how not, the chambers can be used to differentiate these two fluxes.

Answer: Chambers cannot be used to differentiate ebullitive and diffusive fluxes. In the current study, the chambers can give the total flux including ebullitive and diffusive fluxes. We rewrote this section as suggested in the revised version.

3. Line 312. Remove 'obviously'.

Answer: Removed as suggested (Page 18/line 380).

Reviewer 2

This manuscript presents 4 years of CH₄ flux patterns in the largest lake in China and environmental factors that influence CH₄ flux rates. It falls well within the scope of Biogeosciences, but several aspects need to be improved for publication. Some suggestions: 1) How do you define "long-term"? To me, 4-year observations can be short-term. Also, all the statements related to seasonal or inter-annual variability need to be justified because CH₄ flux rates measured on one day may not represent flux rates of one month. Furthermore, daily CH₄ flux rates could have been overestimated,

considering that CH₄ flux rates are measured during the day each month, when CH₄ flux rates were higher than those at night according to diel cycle measurements. 2) All the assumptions are met for regression models? Did you consider any interactions among variables? In addition, did you also carry out the analysis before averaging the flux rates, with replicates as random effects? If so, how did the results differ from those after averaging? 3) In the discussion section some results were described, which did not appear in the result section. Results and discussion need to be better separated. In addition, the interpretation of the results needs to be better supported in the discussion section, focusing clearly on the core messages, i.e., what the results mean and what we can learn from this study.

Answer: We thank the reviewer so much for the constructive comments and suggestions. We have considered all the comments and suggestions carefully in revising the manuscript. Firstly, we avoided using “long-term” as suggested and focused on multi-seasonal dynamics of CH₄ effluxes. We totally agree with the Reviewer that the measured CH₄ effluxes on one day did not represent the mean efflux rate of the month. We used the daily measurements as sampling points to explore the relationships between the CH₄ efflux and environmental variables. We calculated the monthly, seasonal and annual mean CH₄ effluxes using interpolation method (e.g. regression or the random forest model). It is true that most of our measurements were taken during the daytime. However, the daytime and nighttime average CH₄ effluxes were not statistically different ($p = 0.19$). Moreover, we built our statistical models based on the daytime mean efflux and daytime averages of environmental variables and the nighttime efflux was calculated based on the nighttime averages of the same environmental variables. This avoided the overestimation of daily CH₄ efflux. Secondly, we re-analyzed

our data for each site and also treated site as a random effect as suggested. As a result, we found that site had no significant effect on the measured seasonal and diel CH₄ effluxes over the 4-year period (Table S2, S3). In the stepwise multiple regressions analyses, the same environmental variables were selected in the final model for each site as for the 3-site average though the coefficients of each variable were slightly different (Table S5), but not statistically significant ($p > 0.20$). The seasonal patterns of CH₄ effluxes at individual sites were very similar to the seasonal pattern by averaging CH₄ effluxes over the 3 sites. Therefore, we used average values of the 3 sites in our analyses, but we added those information to the result section. Thirdly, we included the interactions among environmental variables in the revised version as suggested (Table S4). Finally, we rewrote the result and discussion sections as suggested to clarify relevant issues.

1. Line#47-51, there are too few references to represent the minimum and maximum flux rates in lakes, especially given that those references are from lakes in China and Norway only. Also, if such values can be presented with more studies, how would seasonal variations look like in comparison to diurnal ones?

Answer: We agree with the Reviewer that there are too few studies measuring lake CH₄ efflux in the literature and the sampling size and frequency was also different among the limited number of studies (Also see Reviewer #1's comments, specific question 5). Therefore, we deleted the range (maximum and minimum) comparison among lakes and focused on comparing the mean efflux of various lakes in the revised version.

2. Line#75-78, can you add references for each variable? Line#64-72 well covered the references for each variable, but this section lacks it.

Answer: We added related references in the method section as suggested in the revised version (Page 4/lines 81-84).

3. Line#78-82, it sounds like investigating in large lakes is not important. Please rephrase or add some more sentences to justify the importance of this research.

Answer: We added some sentences and references to emphasize the importance of CH₄ emissions from large lakes as suggested (Page 5/lines 88-91).

4. Line#86-87, I suggest adding references that describe the previous studies, e.g. Liu et al. (2013).

Answer: Thanks for your suggestion. We added references to describe the previous studies in the revised version (Page 5/ lines 97-102).

5. Line#109, what are the species names of Carex?

Answer: The species name of Carex in Poyang Lake is Carex cinerascens Kükenth and Carex argyi Lev. et Vant. We added the species scientific names in the revised version (Page 7/ lines 123-124).

6. Line#128-146, this section can be written more concisely.

Answer: We rewrote this section as suggested in the revised version.

7. Line#166, can decreases in CH₄ concentrations right after ebullition events be solely explained by diffusion back to lake water? If CH₄ molecules were diffused back to the lake water, partial pressure of CH₄ inside the chamber should be very high, inhibiting further emission from lake water to chamber. Can they be partially from irregular air mixing inside the chamber, which results in errors in CH₄ concentrations? Then, the current method for calculating flux rates needs to be reconsidered.

Answer: We speculate that the short-term decrease or leveling-off of CH₄ concentration inside the chamber after ebullition was mainly caused by the back diffusion of CH₄ to surface water due to the high CH₄ concentration in the

bubbles. This back-diffusion phenomenon has been evidenced for CH₄ efflux over water surfaces (Varadharajan et al., 2010; Wik et al., 2013). The ebullition suddenly increased CH₄ concentration, and thus partial pressure of CH₄, in the chamber headspace, which reversed the normal CH₄ diffusion gradient between surface water and chamber space. We do not think irregular mixing is the main cause in the current study because we had a mixing fan running in each chamber during the whole period of measurement.

8. Line#167-182, this section is confusing. It can be written clearly and concisely.

Answer: We rewrote this section more clearly and concisely as suggested.

9. Line#200, were water and sediment samples collected at three sampling points for flux measurements? The paragraph from line#229 can be given in a Table.

Answer: Yes, we collected water and sediment samples at each of the three sampling sites when taking flux measurements. We added a table (Table S1) to the supplementary material section in the revised version as suggested.

10. Line#241, T test → t-test

Answer: Thank you for pointing out the typo. We changed “T test” to t-test as suggested.

11. Line#242, flux rates are measured three times per season and they may not well represent flux rates of one season of the year. Then, can deviation of these three values be used to quantify inter-annual variability?

Answer: We agree that 3 measurements in a season for a given year are not enough to represent the seasonal mean CH₄ efflux due to the high temporal variation of the efflux. In the current study, we used 4-year data to compare the seasonal variations, which means 12 data points for each season. We changed the values in Table 1 accordingly by using 12 data points to calculate the seasonal

mean effluxes in the revised version (Page 40). For quantifying inter-annual variability we have to interpolate the measured CH₄ effluxes to annual efflux through modeling approach. The details of the modeling work were presented in another paper (Liu et al. 2016, in revision). We used the model results to compare the inter-annual, seasonal, and diurnal variabilities of CH₄ efflux in the Poyang Lake.

12. Line#247, please write what b represents in the equation.

Answer: Thank you for your suggestion. Here b is the exponent of the exponential function between CH₄ efflux and sediment temperature. We added it to the text in the revised version (Page 13/lines 275-276).

13. Line#278, what do you mean by “inconsistent and obvious”?

Answer: This is a typo. We fixed it in the revised version (Page 15/line 312).

14. Line#309-331, this part can be written more concisely.

Answer: Rewritten as suggested.

15. Line#331-332, sentences of this paragraph do not support this conclusion.

Answer: We deleted the concluding sentence.

16. Line#335, here again, can the absolute values be compared with a few references, which are probably based on different observation periods?

Answer: We agree that comparing the extreme values (minimum and maximum) among different lakes is not much meaningful. So, we deleted the relevant text and focused on comparing diurnal patterns.

17. Line#338-342, a larger number of data points can produce wider range of values.

Answer: See answers to question #16.

18. Line#345-356, possible explanations can be added, such as potential drivers that can affect diel CH₄ flux patterns and their variations (if measured).

Answer: Wind speed strongly influenced diel CH₄ efflux variations in our study.

We discussed this point in the 4.3 section.

19. 4.2 CH₄ effluxes in summer, this section contains a lot of new results, which were not presented in the result section. Also, some sentences describe very detailed information from other studies, which hinders the main focus of the paragraph.

Answer: We moved them to the “Results” section and rewrote the discussion by focusing on our own results.

20. 4.3 Timescale dependence of wind, substrate availability, and temperature effects on CH₄ effluxes, here again, a lot of new results are reported, such as line#410-414, line#436-451, line#457-461 (repetition from result section), and line#462-468.

Answer: Again, we moved the results to the “Results” section and rewrote the discussion accordingly.

21. Line#473-475, considering uncertainties related to infrequent measurements (CH₄ efflux rates measured on one day may not represent the mean rates of that month), this kind of statement needs to be corrected.

Answer: According to our model-based interpolation we found that July had the maximum monthly efflux, while January had the minimum. This conclusion is coincidently in line with the 4-year measurements though we had only 4-day measurements in each month. Therefore, we think that the conclusion still holds.

22. Table 3, can you add the observation period of each study for better comparison? Also, sorting the rows by lake size and climate would make this Table easier to read.

Answer: Great idea! We added the observation period of each study and sorted the rows by lake size in the revised version.

23. Figure 3 and 4, can you add error bars from spatial variability?

Answer: We added errors bars from spatial variability for Figure 3 and 4 as

suggested in the revised version.

Reviewer 3

Specific Comments:

1. Most of the results and discussions were built on the environmental variables and methane flux data. However, there are no data of biogeochemical related environmental variables shown in the figures and tables except Table 2. I would suggest to present the raw data of measured environmental variables in the supplementary material.

Answer: Thank you for the suggestion. We added a table (Table S1) to the Supplementary Material section to present the raw data of measured environmental variables, such as sediment total nitrogen content, water level, DOC content in the water, and pH in the sediment , in the revised version.

2. Substrate availability (Line, 432), biological (e.g., microbial activities) and biochemical (e.g., sediment carbon and nitrogen contents processes) (Lines 454-455) are very important factors to link methane efflux to the biogeochemical cycles and understand methane source and sink. Unfortunately, no comprehensive data or evidence to support the role of substrates and microbial activities on methane efflux in this manuscript which could be an important contribution to this journal.

Answer: We agree with the Reviewer that substrates and microbial activities are important to understanding methane sources and sinks in lakes. In our earlier studies we found that sediment carbon and nitrogen ratio were highly correlated with microbial biomass and community structure (Liu et al., 2015) which was also highly associated with greenhouse gas (CO₂, CH₄, and N₂O) fluxes in the Poyang Lake (Liu and Xu, 2016). In the current study, we focus on examining environmental variables (e.g. climate) that may affect the temporal patterns and

variations of CH₄ effluxes in the Poyang Lake. We have added the related information and references to the discussion section in the revised version. Further investigation on the mechanisms of biological and biochemical controls on CH₄ production and oxidation requires lab-based experiments with isotope and microbial DNA sequencing techniques which are beyond the scope of the current study.

3. It might be a risk to use the data from three sampling sites measured from one day (1 hr? Line 148) to represent methane efflux in that month. For example, it appears a contradiction between high methane efflux measured in July 2011 in Figure 3 and low methane efflux measured in July 2011 in Figure 4a.

Answer: We measured CH₄ effluxes at monthly interval to examine the seasonal dynamics of the efflux and the value does not necessarily represent the monthly average of CH₄ efflux. We measured CH₄ effluxes from early morning to late afternoon with about 6 cycles of measurements during the day (Pages 10-11/lines 213-219). The values of methane efflux measured in July 2011 in Figure 3 and Figure 4a are different because of different units. CH₄ efflux in Figure 3 was measured on a daily scale, but CH₄ efflux in Figure 4a was based on the hourly scale. So we used different units to present seasonal and diel patterns of CH₄ effluxes.

4. How long and what time did the authors deploy the floating chambers in the three sampling sites within a day for the study at the large temporal scales (Fig. 3)? I feel 4-year measurements are not a very large temporal scale especially there are no continues measurements/monitering such as deploying floating chamber within a short interval (every week or every two to three days). Since high methane efflux was shown in the early mornings in Fig. 4a, b and d, were the floating chambers deployed

at the same time at three different sites for the data shown in Fig. 3?

Answer: We measured CH₄ fluxes from early morning to late afternoon with about 6 cycles of measurements during the day for the 4-year study. For sampling frequency we measured every monthly. We agree with the Reviewer that 4 year is not “long-term” given the relatively low sampling frequency. So we deleted “long-term” and focused on the multi-seasonal investigations of CH₄ effluxes as suggested by Reviewer 1 in the revised manuscript. We used three boats to monitor CH₄ fluxes at the three sites, so the floating chambers were deployed at about the same time at the sites as shown in Fig. 3.

5. The area and water table of Poyang Lake fluctuate dramatically between the wet and dry seasons. The authors only have short but not clear descriptions of the effect of water level on methane efflux, e.g., in Lines 404-405 and Line 432. Methane efflux might be high in dry seasons instead of summer, since methane efflux is expected to be high under lower water level due to decreasing of the hydrostatic pressure (e.g. Chanton et al. 1989). Are there any difference in water level between three sampling sites in different seasons (The mean water depth at three sites should not be always 3m through the whole year; Line 186)? The authors might consider a simple calculation of methane solubility changes due to water level fluctuations to strengthen the role of water level on methane efflux, e.g., Line 432.

Answer: It is true that the Poyang Lake features a large seasonal variation of water level, high water level in summer and low in winter. However, the water level at the 3 sites was very similar at a given time of the year. We agree that hydrostatic pressure affects CH₄ efflux as reported in Chanton et al. (1989), but our data showed that CH₄ efflux was positively correlated with water level. This is because the water level in the Poyang Lake co-varies with other factors,

such as temperature and NH_4^+ content in the water, which also affect the CH_4 efflux throughout the year. For example, we found that the CH_4 efflux was highly correlated with sediment temperature at an annual scale. Our results suggest that the CH_4 efflux in the Poyang Lake was dominated by temperature rather than water level. The high CH_4 efflux in summer was contributed to strong microbial activities induced by warmer temperature and high substrate availability from the flooding water in summer. Therefore, we think the positive correlation between CH_4 efflux and water level in the Poyang Lake is a pseudo relation which does not reflect the hydrostatic pressure effect on CH_4 efflux as evidenced by Chanton et al. (1989). It is possible to examine the water level effect by calculating CH_4 solubility change due to water level fluctuation. However, given the large seasonal variation of temperature in the study area it is very difficult to separate the water level effect based on the CH_4 efflux measurements on the water surface. In addition, water level induced CH_4 solubility change may affect short-term (minutes to hours) CH_4 diffusion gradient and thus CH_4 efflux and it should have little impact on CH_4 efflux as long as a new diffusion equilibrium has established. Thus, we did not calculate methane solubility changes to further investigate the water level effect on CH_4 efflux in revised the manuscript.

6. As the authors stated in the introduction that methane is driven by three major mechanisms such as molecular diffusion, bubble ebullition and plant-mediated transportation, bubble ebullition is not the only pathway for methane to transport from water to the air. However, data for dissolved methane concentrations in lake water and sediments are lack in this study. No bubble ebullition doesn't mean no methane efflux. I would suggest to include diffusive methane flux to the air for comparison in the

future by analyzing surface water methane concentrations and using the equation from the gas-transfer model e.g., Wanninkhof (1992).

Answer: Great idea! We will take this suggestion in our future study.

7. Since many environmental factors and methane fluxes collected in October 2010 in Poyang Lake have been shown in Liu et al., (2013) for spatial studies, the authors may include Liu et al. (2013) in the introduction and discussions to emphasize why the three sampling sites were chosen in this timescale study and the relations between different environmental factors and methane effluxes in Autumn (October).

Answer: Based on our previous study which examined the spatial variation of CH₄ efflux in the Poyang Lake (Liu et al., 2013), we chose the 3 sites which gave CH₄ effluxes close to the average efflux of the lake. We provided detailed information of Liu et al. (2013) in the introduction and discussion sections as suggested in the revised version.

Minor Comments:

1. Lines 57-59: Please add references for the studies in high-latitude, tropical and subtropical lakes.

Answer: We added references for the studies in high-latitude, tropical and subtropical lakes in the revised version (Page 4/lines 61-63).

2. Line 129: What fluxes did the floating chamber measured while inserting 20 cm above the water surface?

Answer: The chamber measured the total CH₄ efflux including diffusive and ebullitive fluxes as described in the method section. The plant-mediated CH₄ transportation was negligible because no vascular plants grew above water surface at our study sites.

3. Line 150: the air samples ==> the gas samples

Answer: Changed as suggested.

4. Line 159-160; Fig. 4: Since methane efflux was calculated by using a linear regression model to the methane concentration data, should the minimum value be zero instead of a negative value? There should be no negative methane value detected by GC.

Answer: The negative efflux means CH₄-uptake by the lake water due probably to the short-time change in air pressure.

Reference:

- Chanton, J. P.: Gas transport from methane-saturated, tidal freshwater and wetland sediments, *Limnol.Oceanogr.*, 34, 807-819, 1989.**
- Liu, L. X., Xu, M., Lin, M., Zhang, X.: Spatial variability of greenhouse gas effluxes and their controlling factors in the Poyang Lake in China, *Pol. J. Environ. Stud.*, 22, 749-758, 2013.**
- Liu, L. X., Xu, M., Qiu, S., Shen, R. C.: Spatial patterns of benthic bacteria communities in a large lake, *Int. Rev. Hydrobiol.*, 100, 97-105, 2015.**
- Liu, L. X., Xu, M.: Microbial biomass in sediments affects greenhouse gas effluxes in Poyang Lake in China, *J. Freshw. Ecol.*, 31, 109-121, 2016.**
- Liu, L. X., Xu, M.: Modeling temporal patterns of methane effluxes using multiple regression and random forest in Poyang Lake, China, *Wetland Ecology and Management (In Revision)*.**
- Varadharajan, C., Hermosillo, R., Hemond, H. F.: A low-cost automated trap to measure bubbling gas fluxes. *Limnol.Oceanogr.*, 8, 363-375, 2010.**
- Wik, M., Crill, P. M., Varner, R. K., Bastviken, D.: Multiyear measurements of ebullitive methane flux from three subarctic lakes. *J. Geophys. Res.*, 118, 1307-1321, 2013.**

Timescale dependence of environmental controls on methane efflux in Poyang Lake, China

Lixiang Liu¹, Ming Xu^{1,2*}, Renqiang Li¹, Rui Shao^{1,3}

¹Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

²Center for Remote Sensing and Spatial Analysis, Department of Ecology, Evolution and Natural Resources, Rutgers University, New Brunswick, NJ 08901, USA

³University of Chinese Academy of Sciences, Beijing 100049, China

Correspondence to: Ming Xu (mingxu@ igsnrr.ac.cn)

Abstract

Lakes are an important natural source of CH₄ to the atmosphere. However, the ~~long-term-multi-seasonal~~ CH₄ efflux in lakes has been rarely studied. In this study, the CH₄ efflux in Poyang Lake, the largest freshwater lake in China, was measured ~~continuousmonthly~~ly over a 4-year period by using the floating chamber technique. The mean annual CH₄ efflux throughout the 4 years was 0.54 mmol m⁻² day⁻¹, ranging from 0.47 to 0.60 mmol m⁻² day⁻¹. The CH₄ efflux had a high seasonal variation with an average summer (June to August) efflux of 1.34 mmol m⁻² day⁻¹ and winter (December to February) efflux of merely 0.18 mmol m⁻² day⁻¹. The efflux showed no apparent diel pattern, although most of the peak effluxes appeared in the late morning, from 10:00 h to 12:00 h. Multivariate stepwise regression on a seasonal scale showed that environmental factors, such as sediment temperature, sediment total nitrogen content, dissolved oxygen, and total phosphorus content in the water, mainly regulated the CH₄ efflux. However, the CH₄ efflux only showed a strong positive linear correlation with wind speed within a day on a bihourly scale in the multivariate regression analyses but almost no correlation with wind speed on diurnal and seasonal scales.

Keywords: Methane, Sediment temperature, Temperature sensitivity, Substrate availability, Wind speed

带格式的

1. Introduction

Methane (CH₄) contributes to about 20% of global warming in terms of radiative forcing, and its concentration in the atmosphere increased at a rate of 0.5 ppb year⁻¹ in 1999–2006; this rate rapidly increased to 6 ppb year⁻¹ from 2007 to 2011 (IPCC, 2013). Although the total global lake area accounts for approximately 3.7% of the Earth's nonglaciated land area (Verpoorter et al., 2014), CH₄ emissions from global lakes account for up to 14.9% of natural CH₄ emissions (IPCC, 2013). However, this estimate has been associated with large uncertainties because of the high spatial and temporal variations of CH₄ emissions and the insufficient ~~multi-seasonal long-term~~ measurements of CH₄ effluxes, especially in tropical and subtropical lakes (Yang et al., 2011; Ortiz-Llorente and Alvarez-Cobelas, 2012; Bastviken et al., 2015; Li and Bush, 2015).

CH₄ effluxes in lakes feature high temporal variations (Käki, 2001; Xing et al., 2004; Duan et al., 2005; Xing et al., 2005, 2006; Palma-Silva et al., 2013). For example, previous studies found that the ~~mean-minimum-and-maximum~~ CH₄ effluxes over a day were ~~0.018–1.36~~ and ~~12868.85–27~~ mmol m⁻² day⁻¹, respectively (Xing et al., 2004; Duan et al., 2005; Chen et al., 2007; Podgrajsek et al., 2014a, ~~2014b~~); even larger variations were found on a seasonal scale (Xing et al., 2005, 2006; Duan et al., 2005; Ortiz-Llorente and Alvarez-Cobelas, 2012; Wik et al., 2014). These large variations in CH₄ effluxes highlight the importance of frequent and ~~long-term~~ ~~multi-seasonal~~ measurements (Bastviken et al., 2008; Chen et al., 2013; Bastviken et al., 2015). Unfortunately, most earlier studies on CH₄ emissions were based on short-term measurements, ranging from daily to seasonal scales, and were conducted during the day time (Xing et al., 2004; Duan et al., 2005; Xing et al., 2005; Schrier-Uijl et al., 2011; Rõõm et al., 2014). To our knowledge, multi-~~year-seasonal~~

measurements of CH₄ effluxes have only been conducted in high-latitude lakes (Utsumi et al., 1998a, b; Huttunen et al., 2003; Rødm et al., 2014; Wik et al., 2014), and few studies on tropical and subtropical lakes (Xing et al., 2005, 2006; Ortiz-Llorente and Alvarez-Cobelas, 2012), especially large ones, had measurement durations longer than one year.

The magnitude of CH₄ emission mainly depends on the dynamic balance between the microbial processes of CH₄ production, oxidation, physical transportation from the anaerobic zone to the atmosphere in lakes, and regulation by multiple, interconnected physical, chemical, and biological variables (Sun et al., 2012; Liu et al., 2013; Serrano-Silva et al., 2014; Rasilo et al., 2015). CH₄ production and oxidation are microbial processes regulated by organic carbon loading, dissolved organic matter, lake nutrient status, and N availability (Bridgham et al., 2013; Liu et al., 2013; Hershey et al., 2014; Rasilo et al., 2015); temperature (Liikanen et al., 2003; Marotta et al., 2014; Yvon-Durocher et al., 2014); lake depth and size (Juutinen et al., 2009; Rasilo et al., 2015); pH, O₂, NO₃²⁻, Fe³⁺, and SO₄²⁻ in the sediment and water column (van Bodegom and Scholten 2001; Schrier-Uijl et al., 2011; Bridgham et al., 2013); and populations and potential activities of methanogens and methanotrophs (Segers, 1998; van Bodegom and Scholten, 2001; Liu et al., 2015, 2016). CH₄ transportation is driven by three major mechanisms, namely, molecular diffusion, bubble ebullition, and plant-mediated transportation (Bridgham et al., 2013; Chen et al., 2013; Zhu et al., 2016). These mechanisms are affected by water stratification and seasonal overturns of the water mass, which are determined by temperature (Palma-Silva et al., 2013; Rødm et al., 2014), wind-forced mixing (Wanninkhof, 1992; Palma-Silva et al., 2013), water depth (Liu et al., 2013), boundary layer dynamics (Poindexter et al., 2015; Anthony and Macintyre, 2016), hydrostatic pressure (Chanton et al., 1989), and

different vascular plants (Juutinen et al., 2009; Zhu et al., 2016). Most studies examined CH₄ emissions and their influencing factors in small lakes because of their large contribution to the global CH₄ budget (Bastviken et al., 2004; Downing et al., 2010; Bartosiewicz et al., 2015; Holgerson et al., 2016). Although small lakes are a large source of atmospheric CH₄, CH₄ emissions from large lakes was not neglected due to their large areas (Bastviken et al., 2010; Rasilo et al., 2015; Townsend-Small et al., 2016). However, few studies reported temporal CH₄ emissions and their key regulating factors at different temporal scales in large lakes. Therefore, investigating the impacts of physical and biological factors on temporal CH₄ effluxes based on multi-seasonal long-term measurements in a large lake is also important to estimate lake CH₄ emissions.

Poyang Lake, a subtropical lake, is the largest freshwater lake in China, but its annual-multi-seasonal CH₄ emissions have not been adequately measured. In our previous study, we have explored the spatial variations of CH₄ efflux over the lake with 44 sampling locations (Liu et al., 2013). In addition, we also found that microbial biomass and community structure highly influenced CH₄ efflux in Poyang Lake (Liu et al., 2016). In this study, we measured the CH₄ efflux in three sites which we chose on the basis of our previous result over the course of 4 years in Poyang Lake to (1) examine the annual-multi-seasonal mean CH₄ efflux; (2) explore the CH₄ efflux dynamics, including diel and seasonal and inter-annual variations; and (3) quantify the relationships between the CH₄ efflux and environmental factors, and identify the possible factors driving CH₄ effluxes at different temporal scales.

2. Materials and methods

2.1. Site description

Poyang Lake (28°22'–29°45'N, 115°47'–116°45'E) is located in Southern China in Jiangxi Province, with a surface area of 3283 km² and a total catchment area of 162,000 km², which is separated to the northern and southern parts by the Songmen Mountain. Poyang Lake receives water input from five main tributaries, namely, the Raohe River, Xinjiang River, Fuhe River, Ganjiang River, and Xiushui River. The climate is humid subtropical with a mean annual temperature of 17.5 °C and an annual precipitation of 1680 mm (Ye et al., 2011). Vegetation in the lake is composed of macrophytes, including *Carex* sp. (*Carex cinerascens* Kükenth and *Carex argyi* Lev. et Vant.) and *Artemisia selengensis* in the hydrophyte zone, and the main submerged aquatic macrophytes, including *Ceratophyllum demersum*, *Potamogeton malaianus*, *Potamogeton crispus*, and *Hydrilla verticillata* (Wang et al., 2011).

This study was conducted near the Poyang Lake Laboratory of the Wetland Ecosystem Research Station (operated by the Chinese Academy of Sciences), which is located in the northern sub-basin of Poyang Lake in Xingzi County, Jiangxi Province (Fig. 1). The five tributaries flow into the lake in the southeast of Xingzi County, which then joins with the Yangtze River. The water level fluctuated dramatically from 7.78 m to 18.57 m above sea level (Wu Song) between the wet (April to September) and dry seasons (October to March) during the study period because of rainfall and Three Gorge management. Poyang Lake is not stratified (Zhu and Zhang, 1997), with mean and maximum depths of 8 and 23 m, respectively. The mean concentrations of total nitrogen (TN), total phosphorous (TP), suspended solids (SS), and chlorophyll *a* (Chl *a*) in the Poyang lake were 3.45, 0.11, 39.98, and 9.04 mg L⁻¹, respectively (Yao et al., 2015).

2.2. CH₄ efflux measurements

The CH₄ efflux was measured using floating chambers, including both ebullition and diffusive fluxes (Bastviken et al., 2004, 2010). The floating chamber was fabricated using a PVC pipe 100 cm in length and 20 cm in diameter with Styrofoam floats attached to the sides. The floating chambers were inserted 80 cm into the water and 20 cm above the water surface to minimize the perturbation of the surface water flow to the pressure inside the chambers. We tested the chamber system with different insertion depths in the laboratory and field, and found that the current depth of about 80 cm could effectively prevent the impacts of the surrounding Styrofoam floats while maintaining the chamber balance in moderate winds. A similar design of floating chambers was used in previous studies (Lorke et al., 2015; Zhao et al., 2015). Zhao et al. (2015) have recently conducted a systematic comparison of the effects of chamber shape, dimension, and insertion depth into the water on CH₄ effluxes and found that insertion depth only slightly affects the CH₄ efflux measured in the Three Gorges Reservoir when wind speed is relatively low. In the current study, the insertion depth was deeper than those of previous studies to avoid the impact of waves in Poyang Lake on the chamber body. Earlier studies also found that floating chambers should be seated at the water surface with minimal insertion into the water in a flowing-water system to minimize the “drag” effect of flowing water on chamber pressure (Bastviken et al., 2010; Vachon et al., 2013; McGinnis et al., 2015). Except for some waves, However, the water in Poyang Lake did not have an apparent directional flow except for some waves during the measurement period. So In the current study, the insertion depth was deeper than those of previous studies to avoid the impact of waves in Poyang Lake on the chamber body in the current study. A detailed description of the floating chamber system can be found in Liu et al. (2013). So we measured the

total CH₄ efflux including both ebullition and diffusive effluxes and cannot differentiate ebullitive and diffusive fluxes by our chamber.

We collected a gas sample (ambient concentration) immediately after the chamber was closed and three other samples at a 20 min interval for 1 h. The gas was extracted into a 12 mL evacuated glass vial by a 2 mL syringe needle with an air pump, which enhanced the pressure in the vial to 3 bars. Subsequently, the gas-air samples were transported immediately to a laboratory for CH₄ concentration analysis. The CH₄ concentration was measured using a gas chromatograph equipped with a flame ionization detector (GC7890A, Agilent Technologies, Inc., Santa Clara, CA, USA). We used nitrogen (N₂) as the carrier gas, which ran at a flow rate of 30 mL min⁻¹. We calibrated the gas chromatograph for every four samples with a calibration gas of 2.03 ppm at 99.92% precision (China National Research Center for Certified Reference Materials, China). The oven and detector temperatures of the GC were set to 55 °C and 250 °C, respectively.

Calculation of the CH₄ efflux was based on the CH₄ concentration of the four samples using a linear regression model $y = ax + b$, which was calculated on the basis of the slope of the concentration change during the whole period when the chamber was closed. Data quality control was conducted following the method of Rasilo et al. (2015) before the regression models were fitted. As a result, most of the models performed satisfactorily, with a coefficient of determination (R^2) greater than 0.95. In case of ebullition, the CH₄ concentration inside the chamber would deviate from the normal trend. Most of the CH₄ concentrations measured immediately after the ebullition point slightly decreased mainly because of the CH₄ diffusion back to water when the CH₄ concentration inside the chamber space increased suddenly from bubbling. To include the ebullition-induced CH₄ emissions, we only used two

175 measured concentrations, the first measurement (ambient concentration) and an
ebullition-adjusted concentration that was obtained by adding the diffusion-induced
concentration increment when ebullition occurred (Fig. 2). ~~in calculating the CH₄~~
~~efflux when ebullition occurred inside the chamber. The ebullition-adjusted~~
~~concentration was obtained by adding the diffusion-induced concentration increment,~~
180 ~~which is a correction term, to the measured concentration when ebullition occurred.~~
~~The total CH₄ efflux, which includes both ebullition and diffusive effluxes, was~~
~~calculated on the basis of the slope of the concentration change during the whole~~
~~period when the chamber was closed (Fig. 2).~~ Specifically, when ebullition occurred
during the first 20 min, we obtained the ebullition-adjusted concentration by summing
185 up concentration on 20 min and the 2-fold incremental concentration which was from
concentration between the third and fourth sampling times. When the ebullition
occurred at the third sampling, we summed up the concentration at 40 min and the
incremental concentration between the first and second sampling times. When the
ebullition occurred at the fourth sampling, we used the first and fourth sampling
190 concentrations directly to calculate the slope of the total efflux.

Samplings took place at a monthly interval from January 2011 to December 2014
at three sites in Poyang Lake (Fig. 1): site A (Luoxingdun: 29°3'29"N, 116°16'49"E),
site B (Mantianxing: 29°34'25"N, 116°13'29"E), and site C (Huoyanshan: 29°39'0"N,
116°16'11"E). The mean water depth in our sampling sites was 3 m. The sampling
195 sites lacked aquatic plants. Our previous study examined the spatial pattern of the CH₄
efflux in the lake (Liu et al., 2013). Therefore, we focused on the long-term
multi-seasonal dynamics of CH₄ efflux in the current study. At each site, four
chambers were placed approximately 10 m away from a small boat to minimize
disturbance. Measurements were conducted from early morning to late afternoon with

about 6 cycles of measurements for each chamber, except for days when the diel-cycle measurements were taken. We conducted four 24 h measurements at the three sites in 24–25 July 2011, 5–6 September 2012, 13–14 January 2013, and 14–15 January 2015 to examine the diel variations of CH₄ effluxes. These measurements were conducted every 2 h from 8:00 am to 8:00 am the next day, providing 12 cycles of measurements for each chamber per 24 h.

2.3. Environmental variables

Various environmental variables were also measured in the lake sediment, surface water, and atmosphere at each individual site and then averaged when we used.

We collected surface water and sediment samples (0–15 cm) using a plexiglass water grab and a stainless steel sediment sampler (3 cm in diameter) after obtaining gas samples. The water and sediment samples were immediately stored in plastic bottles and bags, respectively. Then, all the samples were stored in ice coolers and transported to a laboratory for analysis within a week. In addition, we measured the wind speed at about 1.5 m above the water surface using a portable anemometer (Testo 410-1, Testo, Germany) and the surface sediment (0–15 cm) temperature using a mercury thermometer. We used a multi-parametric probe (556 MPS, YSI, USA) to measure the water quality factors in situ, such as electrical conductivity and dissolved oxygen (DO) content, at each sampling site from June 2013 to June 2014. The water levels in the lake were obtained from the Xingzi Hydrological Station, about 20 km from our sampling sites.

In the laboratory, the pH values of the water and sediment samples were measured using a pH meter (Delta 320, Mettler–Toledo, Switzerland). Chemical oxygen demand (COD) was measured using the spectrophotometric detection method based on Griess reaction (Jirka and Carter, 1975; Yao et al., 2015). Chl *a* concentration was measured

via spectrophotometry (Rasilo et al., 2015; Yao et al., 2015), which was extracted in 90% ethanol and then analyzed spectrophotometrically at 750 and 665 nm in accordance with ISO 10 260 (1992). The SS level in the lake water was measured by a gravimetric procedure, where the solids from the water sample were filtered, dried, and weighed to determine the total non-filterable residue of the sample (Fishman and Friedman, 1989). TP concentration was measured using the molybdenum blue method after persulfate digestion (Karl and Tien, 1992; Yao et al., 2015). In addition, the nitrate (NO_3^-), ammonium (NH_4^+), TN, and dissolved organic carbon (DOC) contents in the water were measured using a total carbon and nitrogen analyzer using filtered water (Shimadzu TOC-VCSH + TN module, Shimadzu, Japan). The sediment TN and organic carbon contents after total sediment acidification with HCl 1N were determined using a vario MAX CN element analyzer (NA Series 2, CE Instruments, Germany). The values of measured environmental variables in our study were given in Table S1.

Considering the different sampling periods, we classified the environmental variables into three groups (Table S1). The first group included sediment temperature, sediment total nitrogen content, water level, DOC content in the water, pH in the sediment, NH_4^+ and NO_3^- concentrations in the water and sediment, sediment organic carbon content, the ratio of carbon and nitrogen, and the mean daily wind speed over a 48-month period. The second group included TN, TP, COD, and Chl *a* contents in the water, which were sampled between June 2011 and December 2014. We sampled the third group variables from June 2013 to June 2014, including DO content, conductivity, and pH in the water.

2.4. Data analysis

We averaged the CH_4 effluxes of the three sites to minimize the effect of the

250 spatial variation of CH₄ efflux on the temporal dynamics of the efflux. One-way ANOVA followed by post-hoc Tukey's test and paired ~~T-t~~ test were used to analyze the seasonal ~~and inter-annual~~ differences in the CH₄ effluxes. ~~The coefficient of variation (CV) was used to quantify the inter-annual variation of CH₄ efflux.~~ We employed stepwise multiple regressions to identify the environmental factors driving 255 the CH₄ effluxes at different temporal scales. We also used regression and correlation analyses to determine the relationships between independent variables and CH₄ effluxes. In addition, we considered each study site as a random effect in linear mixed effects models in order to take into account CH₄ efflux variations among three sites when we investigated seasonal and diurnal variations as well as the relationships between CH₄ efflux and environmental variables. We used the Vant' Hoff equation to 260 calculate the temperature sensitivity ($Q_{10} = e^{10b}$, where b is the exponent of the exponential function between CH₄ efflux and sediment temperature) of CH₄ efflux (Xu and Qi, 2001; Wei et al., 2015). All statistical analyses were performed using the SPSS 17.0 statistical software (SPSS Inc., Chicago, IL, USA), and graphs were 265 created using the Sigma Plot 11.0 program (Systat Software Inc., San Jose, CA, USA).

3. Results

3.1. CH₄ effluxes in Poyang Lake

3.1.1. Annual CH₄ effluxes

The mean CH₄ efflux was $0.54 \pm 0.053 \text{ mmol m}^{-2} \text{ day}^{-1}$ in Poyang Lake over the 4 year period, with annual mean effluxes of 0.47 ± 0.54 , 0.56 ± 0.41 , 0.52 ± 0.55 , and $0.60 \pm 0.56 \text{ mmol m}^{-2} \text{ day}^{-1}$ in 2011, 2012, 2013, and 2014, respectively (Table 1). The inter annual variation of CH₄ efflux was moderately high with a CV of 9.8% over the 4 years. The mean CH₄ efflux in 2014 was 25.7% greater than that in 2011, justifying the necessity for long term measurements.

3.1.2. Seasonal CH₄ effluxes

The seasonal variations of CH₄ effluxes in Poyang Lake were prominent, demonstrating a similar pattern to that of seasonal temperature (Fig. 3). In general, the annual maximum CH₄ effluxes occurred in summers and the minimum in winters. The CH₄ efflux increased slowly in early spring and then rapidly in May, reaching its maximum in July. After reaching the maximum, the CH₄ efflux decreased sharply in August and September and then slowly before reaching its minimum in January (Fig. 3). Significant differences in the mean CH₄ effluxes existed between summers and the other three seasons throughout the 4 years ($p < 0.05$), whereas the differences in the CH₄ effluxes among the spring, autumn, and winter seasons were not statistically significant ($p > 0.05$) (Table 1). Additionally, the site effect was not statistically significant over the 4-year period (Table S2). The differences among the three sites were minor with the 4-year mean of $0.53 \text{ mmol m}^{-2} \text{ day}^{-1}$, $0.55 \text{ mmol m}^{-2} \text{ day}^{-1}$, and $0.54 \text{ mmol m}^{-2} \text{ day}^{-1}$ respectively. In particular, the seasonal patterns of CH₄ effluxes at the three sites were similar and also in line with the seasonal pattern averaged over the three sites.

3.1. ³². Diel CH₄ effluxes

The CH₄ effluxes in Poyang Lake also exhibited apparent variations within a day because the daily maximum appeared late in the morning (10:00–12:00 h) and the minimum early in the morning the next day (4:00–6:00 h). The diel pattern of the CH₄ efflux was asymmetric, fast increasing in the morning from 8:00 h to 12:00 h and slowly decreasing in the afternoon and during the night, especially in the summer (Fig.

4). However, the diel pattern of the CH₄ efflux was inconsistent ~~and obvious~~. For example, the diel pattern on January 13–14, 2013 was an exception, when the

maximum efflux occurred around 6:00 h on January 14th and a severe cold front with heavy fogs enveloped the Poyang Lake area in the early morning of January 14th. The

CH₄ efflux magnitudes were significantly larger during summer compared to winter.

The diel pattern of CH₄ efflux was vague with an average difference between the daily maximum and minimum of only 0.073 mmol m⁻² h⁻¹. The CH₄ efflux could also

change abruptly throughout a day. For example, the efflux sharply dropped from 0.068 to -0.012 mmol m⁻² h⁻¹ within barely 2 h, as observed on July 23, 2011, indicating that the lake switched from a CH₄ source to sink within a short period of time (Fig. 4a). This abrupt change was also observed in the afternoon of August 28,

2012 (Fig. 4b). Furthermore, we compared the differences of diurnal patterns at each

sites for the four diel sampling. Our results showed that the diel patterns of CH₄

effluxes were similar in the three sites for each diel investigation (Fig. 5) and the site

effect was not statistically significant (Table S3). The diel pattern of the CH₄ efflux

was somewhat different during certain hours such as from 22:00 h to 00:00h on July

24th in 2011. Further analysis showed that the diel pattern of CH₄ effluxes followed the

diel pattern of wind speed (Figs. 5a–5d).

3.2. Relationships between CH₄ efflux and environmental variables

3.2.1 Simple regression relationships between CH₄ efflux and environmental variables

In our study, CH₄ effluxes increased exponentially with sediment temperature for both in the summer and in other seasons (Fig. 6). The CH₄ effluxes were more sensitive to temperature in the summers than in other seasons. The temperature sensitivity, indicated by the Q₁₀ values, was 2.04 and 1.67 in the summer and other three seasons, respectively (Fig. 6).

We found that CH₄ effluxes were also highly associated with other climate and environmental variables in both lake water and sediments. We found that CH₄ effluxes were negatively correlated with NH₄⁺, TN and DO concentrations in the lake water, but positively with Chl *a* content in the water and TN content in the sediment (Table 2). Furthermore, we found that other environmental factors, such as DOC content in the water, pH in the water and in the sediment, NO₃⁻ concentrations in the water and in the sediment, COD, and TP in the water, had insignificant ($p > 0.05$) relationships with CH₄ effluxes in Poyang Lake.

In the current study, we also found that the relationship between CH₄ effluxes and wind speed was scale-dependent. At the diel scale, wind speed was significantly correlated ($p < 0.03$) with CH₄ effluxes for the average of the 3 sites at the diel scale (Fig. 7), but was weakly correlated with CH₄ effluxes at the diurnal and seasonal scales (Fig. 7). In addition, the relationships between wind speed and CH₄ effluxes for each individual site were similar with the relationships for the average of the 3 sites though the regression coefficients for each individual site were slightly different, but not statistically significant ($p > 0.25$).

3.2.2 Multiple regression relationships between CH₄ efflux and environmental variables

In the current study, environmental factors differed in importance depending on

the timescale in the stepwise multiple regressions analyses. The results of stepwise multiple regressions on a seasonal scale showed that the sediment temperature, sediment TN content, DO, and TP content in the water were significant predictors of

CH₄ effluxes (Table 23). It should be noted that multicollinearity didn't occurred among these significant variables (Table S4). In specific, sediment temperature and sediment TN content explained 65% of the variation in CH₄ effluxes for 4 years when we used the first group of factors. The sediment temperature and TN content explained 73% of the CH₄ efflux variations when the second group of variables was added to the first group. The sediment temperature, sediment TN content, DO, and TP contents in the water explained 89% of the CH₄ efflux variation when the three groups of variables were used together. Wind speed was the only significant variable for the CH₄ efflux variations on a diel scale. Wind speed explained 58%, 56%, 84% and 86% of the CH₄ efflux variations in 24–25 July 2011, 5–6 September 2012, 13–14 January 2013 and 14–15 January 2015, respectively (Figs. 5a7a-5d7d). In addition, the same environmental variables were selected in the final model for each individual site as for the average of the 3 sites though the regression coefficients were slightly different (Table S5), but not statistically significant ($p > 0.20$).

4. Discussion

4.1. CH₄ effluxes in Poyang Lake

The mean CH₄ emission in Poyang Lake was moderately higher than those in other large lakes of more than 1 km² in the world. The mean CH₄ emission (0.54 mmol m⁻² day⁻¹) was within the reported range of approximately 0.022–5.85 mmol m⁻² day⁻¹ in boreal and temperate lakes over 1 km² but was obviously lower than diffusive effluxes in subtropical lakes and total effluxes (including diffusion and ebullition) in tropical lakes (Table 34). In addition, the mean CH₄ emission in Poyang Lake was comparable with the diffusive effluxes in tropical lakes (Table 34). For example, previous studies reported that the diffusive CH₄ efflux was 0.65 mmol CH₄ m⁻² day⁻¹ in the TR Lake and 0.50 mmol m⁻² day⁻¹ in the BB Lake in the Pantanal region (Bastviken et al., 2010). However, the mean CH₄ efflux in Poyang Lake was only higher than those in other lakes over 100 km² (except the Värtsjäv Lake). The low CH₄ efflux in the current study was unlikely caused by our floating chamber system because the CH₄ efflux would have increased if the insertion of chambers considerably disturbed the water profiles. The lower CH₄ emissions in our study may be attributed to the low concentration of carbon substrates in the water and sediments in Poyang Lake. The DOC concentration in Poyang Lake was merely 3.3 mg L⁻¹, which was much lower than that of the 5.8 mg L⁻¹ in Biandantang Lake and 7.4 mg L⁻¹ in Donghu Lake, which are two subtropical lakes in China (Xing et al., 2005, 2006). Poyang Lake also has a lower organic carbon content in its sediments than most other lakes. The average organic carbon content in the sediments in Poyang Lake was 0.89%, which was much lower than that of 30.76% averaged over five temperate lakes (Schrier–Uijl et al., 2011) and slightly higher than that of nearly 0.75% in tropical lakes in the Pantanal region (Bastviken et al., 2010). Therefore, the CH₄

emissions in large lakes cannot be ignored when estimating the global CH₄ budget because of their area.

CH₄ effluxes at the air-water interface showed high fluctuations in the four cycles, but showed no significant diurnal differences. The diurnal CH₄ efflux ranged from 0.019 to 0.13 mmol m⁻² h⁻¹, which was within the reported range of other lakes (0.057 to 5.37 mmol m⁻² h⁻¹) over a diurnal cycle (Xing et al., 2004; Duan et al., 2005; Chen et al., 2007; Podgrajsek et al., 2014a, 2014b). The wide range of diurnal CH₄ efflux in previous results may be due to differences in sample size in different studies. For example, CH₄ efflux was measured at 2h intervals with 12 data points over a diurnal cycle in this study, but CH₄ efflux was measured at 3-6 h intervals with only 4-8 data points in previous studies (Käki et al., 2001; Xing et al., 2004; Duan et al., 2005). Another possible reason for these discrepancies was that vegetation might have played an important role in CH₄ efflux in other studies (Käki et al., 2001; Duan et al., 2005), while there was no vegetation in the water where we sampled in Poyang Lake. In addition, our study showed that there were no significant differences in CH₄ effluxes between the nighttime and the daytime, which was inconsistent with other studies (Keller and Stallard, 1994; Bastviken et al., 2010). This inconsistency may be due to incomplete measurement of the diurnal cycle in other studies. For example, CH₄ efflux was only measured three times at sunrise, daytime and sunset to represent a diel cycle in Bastviken et al. (2010). In Keller and Stallard (1994) study, the daytime and nighttime CH₄ efflux measurements were not conducted on the same day. In particular, two studies reported a new finding that hydrodynamic transport contributed more to nighttime CH₄ effluxes than daytime CH₄ effluxes (Poindexter et al. 2015; Anthony and Macintyre 2016). However, we cannot estimate CH₄ effluxes by hydrodynamic transport because we did not measure CH₄ concentration in the water

~~in this study. Further studies are needed to address this issue in the lake.~~

4.2. CH₄ effluxes in summer

The CH₄ effluxes in Poyang Lake were substantially greater in summer than in the other seasons, accounting for more than 63% of the annual total emissions. This finding suggests that summer is the critical season in managing the CH₄ emissions from Poyang Lake. The high effluxes in summer may be attributed to the higher temperature, higher substrate availability, and greater temperature sensitivity during this season than the other seasons.

~~The high summer CH₄ effluxes may due to high temperature in summer. Poyang Lake features a typical monsoon climate with hot summers. During the study period, the mean (June–August) air temperature in summer was 28.5 °C, whereas that in winter was only 5.9 °C.~~ The CH₄ effluxes were highly correlated with the sediment temperature through an exponential function ~~in our study. During the study period, the mean (June–August) air temperature in summer was 28.5 °C, whereas that in winter was only 5.9 °C.~~ Our results confirmed the findings of previous studies that lake CH₄ effluxes are driven by temperature (Bastviken et al., 2008; Marinho et al., 2009; Palma–Silva et al., 2013; Rõm et al., 2014). This is supported by the fact that a warm temperature provides a high optimal temperature for methanogen growth ~~which increases methane production~~ (Nozhevnikova et al., 2007; Rooney–Varga et al., 2007; Duc et al., 2010) ~~and. Moreover, recent studies have reported that high temperatures could increase~~ the proportion of hydrogenotrophic methanogenesis ~~which is an important pathway for CH₄ production~~ (Borrel et al., 2011; Marotta et al., 2014). The high summer CH₄ effluxes might also be because of the ample substrate supply in this season ~~because the decomposition rate of new organic matter was much faster than that of old organic matter (Davidson and Janssens, 2006; Gudas et al., 2010).~~ In the

present study, CH₄ efflux positively correlated with the Chl *a* content (Table 2r = 0.46, data not shown) that was not correlated with other environmental factors (Table S4) and acted as an indicator of primary production. Earlier studies discovered a high amount of labile organic matter, including allochthonous inputs of terrestrial organic matter, during the summer flooding and autochthonous production within-lake by phytoplankton and benthic algae in summer (Crump et al., 2003; Xing et al., 2005, 2006; Bade et al., 2007). ~~The decomposition rate of new organic matter was much faster than that of old organic matter (Davidson and Janssens, 2006; Gudas et al., 2010).~~ Previous studies showed that fresh organic carbon from dead algae stimulates CH₄ emissions in lakes (Huttunen et al., 2002; Xing et al., 2005) because the degradation of dead alga and algal exudates, ~~such as methylated compounds,~~ are the precursors for CH₄ production (Ferrón et al., 2012; Xiao et al., 2015; Liang et al., 2016). However, we did not find any correlation between the CH₄ efflux and DOC content in the water ($p > 0.05$). The algal bloom in summer probably masked the DOC effect on stimulating CH₄ production. Earlier studies demonstrated that 70%–80% of DOC molecules in lakes are recalcitrant carbon, ~~which are composed of humic substances in the lake from the partial degradation of terrestrial lignin in vegetation~~ (Tranvik and Kokalj, 1998; Wetzel, 2001).

The high summer CH₄ effluxes were also driven by the greater temperature sensitivity during summer. The apparent Q₁₀ value in Poyang Lake was 2.04 in summer, which was much greater than the value of 1.67 in the other seasons (Fig. 6). This finding is inconsistent with previous studies in terrestrial and freshwater ecosystems (Davidson and Janssens, 2006; Gudas et al., 2010; Yvon-Durocher et al., 2014), where the Q₁₀ values decreased apparently with the increase in temperature (Xu and Qi 2001a; Chen et al., 2010; Corkrey et al., 2012; Schipper et al., 2014).

However, our result was supported by a recent finding that the temperature sensitivities (Q_{10}) of CH_4 effluxes from lake sediments are greater in the tropics than in boreal regions (Marotta et al., 2014). We speculate that the temperature effect on Q_{10} was confounded by other factors, such as water level and substrate availability.

The addition of a large amount of fresh carbon from summer floods could dramatically boost CH_4 production and thus the apparent Q_{10} values during summer.

4.3. Timescale dependence of wind, substrate availability, and temperature effects on CH_4 effluxes

In this study, the effects of wind, substrate availability, and sediment temperature on CH_4 effluxes were highly timescale dependent. The CH_4 effluxes measured at bihourly intervals positively correlated with wind speed in both simple and multiple regressions (Figs. 5a7a–d, Table 2) but showed no correlation ($p > 0.05$) when the diurnal or seasonal average CH_4 efflux and wind speed were applied (Figs. 5e7e–f).

The effect of wind on CH_4 effluxes was mainly through its effects on the transport, air pressure and storage of CH_4 from the bottom to the surface water (Abril et al., 2005; Hahm et al., 2006; Guérin et al., 2007). Gas diffusion in water is sensitive to pressure changes at the water–air interface (Paganelli et al., 1975; Massmann and Farrier, 1992; Striegl et al., 2001; Nachshon et al., 2012). High wind speed mechanically induces turbulences through friction in the water and brings CH_4 -rich water from the bottom to the surface in lakes (Wanninkhof, 1992; Palma–Silva et al., 2013; Xiao et al., 2013). The CH_4 efflux rapidly decreases or even becomes negative (indicating CH_4 absorption) to compensate for the deficits in the water profile caused by earlier winds when the wind declines or comes to a halt. Our results also confirmed that the CH_4 efflux sharply declined to a negative value after strong wind events (Fig. 4). This wind

effect only worked at short timescales, such as bihourly, when temperature only slightly changed and other biological processes, such as microbial community variation, were relatively stable. At a longer temporal scale, such as seasonal scale as observed in the current study, the wind effect disappeared because the

490 wind-stimulated CH₄ effluxes and the post-wind (or between-gusts) negative effluxes (absorptions) were compensated. Our results suggest that wind exerts minor effects on CH₄ effluxes at large temporal scales when temperature, water level, and substrate availability dominate. Our results also suggest that caution must be taken when one applies the empirical wind speed-driven models developed based on short-term

495 measurements to estimate CH₄ effluxes over long periods, such as months or years.

Meanwhile, the CH₄ effluxes measured at monthly intervals positively correlated with sediment temperature (Fig. 6, [Table 2](#)), but the correlation disappeared when applied at bihourly intervals ($p > 0.05$). The lack of correlation between the CH₄ efflux and sediment temperature as measured on a bihourly scale within a day can be

500 explained by the small variation of sediment temperature within a day, ranging from 0.95 °C to 1.85 °C. Other factors, such as wind and atmospheric pressure, might shadow the weak temperature effect within a day. Instead, we found a high correlation between the bihourly measured CH₄ effluxes and sediment temperature during the diel measurement period in January 14 to 15, 2015 ($r = 0.88$, $p < 0.0001$). Further analyses

505 showed that this temperature effect might be apparent and mainly caused by wind speed because the bihourly measured CH₄ effluxes and wind speed were highly correlated only in January 14 to 15, 2015 and not in the other days ($r = 0.90$, $p < 0.0001$). However, sediment temperature became the dominant factor on a seasonal scale when the temperature ranged from about 4.4 °C in winter to 30.8 °C in summer

510 (Fig. 3). The sediment temperature and CH₄ effluxes averaged over the diurnal period

significantly correlated in the 4-year study period (Fig. 6, [Table 2](#)). Our results suggest that the short-term CH₄ efflux in Poyang Lake was regulated by wind speed, but the ~~long-term~~ [multi-seasonal](#) CH₄ efflux was ultimately controlled by sediment temperature and other biological (e.g., microbial activities) and biochemical (e.g., sediment carbon and nitrogen contents) processes. Therefore, understanding and modeling the dynamics of CH₄ effluxes on lake surfaces require the [multi-seasonal](#) ~~long-term~~ measurements of effluxes and related biotic and abiotic factors in lake water and sediments. Finally, substrate availability, such as sediment TN content, TP, and Chl *a* contents in the water, also influenced CH₄ effluxes on a seasonal scale in the current study ([Table 2](#), [3](#)). However, the effects disappeared when applied at bihourly intervals because the substrate did not change significantly within a day.

In addition to the above-mentioned factors, the DO concentration in the water influenced the CH₄ effluxes in the multivariate regression analysis. In specific, the CH₄ efflux closely correlated with the DO concentration in the water (~~$r = -0.65$~~ [Table 2](#)). This close correlation can be explained by the aerobic CH₄ oxidation in the water. Our result was supported by the previous finding that a high DO concentration in the water results in low CH₄ emission (Rõm et al., 2014; McNicol and Silver, 2015; Yang et al., 2015).

5. Conclusion

The average CH₄ efflux in Poyang Lake during the 4-year study period was $0.54 \pm 0.053 \text{ mmol m}^{-2} \text{ day}^{-1}$, which was moderately higher than that of the other lakes in the world. The CH₄ efflux in Poyang Lake also featured high multi-seasonal variations with the maximum efflux in July and the minimum in January. About 63% of the annual emissions occurred in summer, from June to August. On a seasonal scale, multivariate regression analyses revealed that sediment temperature sediment TN content, TP, and DO contents in the water mainly regulated the CH₄ effluxes. Simple and multivariate regression analyses showed that wind speed influenced the diel CH₄ efflux variations. The effects of sediment temperature, substrate availability, and wind speed on CH₄ effluxes were temporal scale dependent. The CH₄ effluxes increased with the sediment temperature, sediment TN content, Chl *a*, and TP contents in the water on a seasonal scale but were not correlated with sediment temperature on a bihourly scale. In contrast to the temperature and substrate, the CH₄ efflux positively and significantly correlated with wind speed within a day on a bihourly scale but was not correlated with wind speed at larger temporal scales, such as daily and seasonal scales. The timescale dependence of environmental controls on CH₄ effluxes has important implications in modeling CH₄ emissions.

Acknowledgements

550 This research was supported by the National Basic Research Program of China (973
Program, 2012CB417103), Assessment and Valuation of Ecosystem Services in
Qinghai Provincem, China (No. 2013-N-556). We gratefully acknowledge the Poyang
Lake Laboratory Wetland Ecosystem Research, CAS for permission to access the
study site and assistance with field work.

555

References

- Abril, G., Gu érin, F., Richard, S., Delmas, R., Galy-Lacaux, C., Gosse, P., Tremblay, A., Varfalvy, L., Dos Santos, M. A., Matvienko, B.: Carbon dioxide and methane emissions and the carbon budget of a 10-year old tropical reservoir (Petit Saut, French Guiana), *Glob. Biogeochem. Cy.*, 19, GB4007, 2005. <http://dx.doi.org/10.1029/2005GB002457>.
- Anthony, K. W., Macintyre, S.: Nocturnal escape route for marsh gas, *Nature*, 535, 363–365, 2016.
- Bade, D. L., Carpenter, S. R., Cole, J. J., Pace, M. L., Kritzberg, E., Van de Bogert, M. C., Cory, R. M., McKnight, D. M.: Sources and fates of dissolved organic carbon in lakes as determined by whole-lake carbon isotope additions, *Biogeochemistry*, 84, 115-129, 2007.
- Bartosiewicz, M., Laurion, I., MacIntyre, S.: Greenhouse gas emission and storage in a small shallow lake, *Hydrobiologia*, 757, 101-115, 2015.
- Bastviken, D., Cole, J., Pace, M., Tranvik, L.: Methane emissions from lakes: dependence of lake characteristics, two regional assessments, and a global estimate, *Glob. Biogeochem. Cy.*, 18, 1-12, 2004.
- Bastviken, D., Cole, J. J., Pace, M. L., Van de Bogert, M. C.: Fates of methane from different lake habitats: connecting whole-lake budgets and CH₄ emissions, *J. Geophys. Res.*, 113, G02024, 2008. <http://dx.doi.org/10.1029/2007JG000608>.
- Bastviken, D., Natchimuthu, S., Panneer Selvam, B.: Response: inland water greenhouse gas emissions: when to model and when to measure?, *Glob. Change Biol.*, 21, 1379–1380, 2015.

- Bastviken, D., Santoro, A. L., Marotta, H., Pinho, L., Calheiros, D., Crill, P.: Methane
580 emissions from Pantanal, South America, during the low water Season: Toward
more comprehensive sampling, *Environ. Sci. Technol.*, 44, 5450-5455, 2010.
- Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., Enrich-Prast, A.:
Freshwater methane emissions offset the continental carbon sink, *Science* 331,
50-50, 2011.
- 585 Borrel, G., Jézéquel, D., Biderre-Petit, C., Morel-Desrosiers, N., Morel, J. P., Peyret,
P.: Production and consumption of methane in freshwater lake ecosystems, *Res.*
Microbiol. 162, 832-847, 2011.
- Bridgman, S. D., Cadillo-Quiroz, H., Keller, J. K., Zhuang, Q. L.: Methane emissions
from wetlands: biogeochemical, microbial, and modeling perspectives from local
590 to global scales, *Glob. Change Biol.*, 19, 1325-1346, 2013.
- Chanton, J. P.: Gas transport from methane-saturated, tidal freshwater and wetland
sediments, *Limnol. Oceanogr.*, 34, 807-819, 1989.**
- Chen, Y. G., Bai, X. H., Li, X. H., Hu, Z. X., Liu, W. L.: Primary study of the
methane flux on the water-air interface of eight lakes in winter, China. *J. Lake Sci.*
595 19, 11-17, 2007.
- Chen, B. Y., Liu, S. R., Ge, J. P., Chu, J. X.: Annual and seasonal variations of Q_{10}
soil respiration in the sub-alpine forests of the Eastern Qinghai-Tibet Plateau,
China, *Soil Biol. Biochem.*, 42, 1735-1742, 2010.
- Chen, H., Zhu, Q. A., Peng, C. H., Wu, N., Wang, Y. F., Fang, X. Q.: Methane
600 emissions from rice paddies natural wetlands, lakes in China: synthesis new
estimate, *Glob. Change Biol.*, 19, 19-32, 2013.

Corkrey, R., Olley, J., Ratkowsky, D., McMeekin, T., Ross, T.: Universality of thermodynamic constants governing biological growth rates, *Plos One*, 7, e32003, 2012.

605 Crump, B. C., Kling, G. W., Bahr, M., Hobbie, J. E.: Bacterioplankton community shifts in an Arctic Lake correlate with seasonal changes in organic matter source, *Appl. Environ. Microbiol.*, 69, 2253-2268, 2003.

Davidson, E. A., Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165-173, 2006.

610 Downing, J. A.: Emerging global role of small lakes and ponds: little things mean a lot, *Limnetica*, 29, 9-24, 2010.

Duan, X. N., Wang, X. K., Mu, Y. J., Ouyang, Z.Y.: Seasonal and diurnal variations in methane emissions from Wuliangsu Lake in arid regions of China, *Atmos. Environ.*, 39, 4479-4487, 2005.

615 Duc, N. T., Crill, P., Bastviken, D.: Implications of temperature and sediment characteristics on methane formation and oxidation in lake sediments, *Biogeochemistry*, 100, 185-196, 2010.

Fallon, R. D., Harris, S., Hanson, R. S., Brock, T. D.: The role of methane in internal carbon cycling in Lake Mendota during summer stratification, *Limnol. Oceanogr.*, 25, 357-360, 1980.

620 Ferrón, S., Ho, D.T., Johnson, Z. I., Huntley, M. E.: Air-water fluxes of N₂O and CH₄ during microalgae (*Staurosira* sp.) cultivation in an open raceway pond, *Environ. Sci. Technol.*, 46, 10842-10848, 2012.

Fishman, M. J., Friedman, L. C.: Methods for determination of inorganic substances in water and fluvial sediments. US Department of the Interior, Geological Survey, 1989.

Gudas, C., Bastviken, D., Steger, K., Premke, K., Sobek, S., Tranvik, L. J.:

Temperature-controlled organic carbon mineralization in lake sediments, *Nature*, 466, 478-481, 2010.

630 Gu érin, F., Abril, G., Serca, D., Delon, C., Richard, S., Delmas, R., Tremblay, A.,

Varfalvy, L.: Gas transfer velocities of CO₂ and CH₄ in a tropical reservoir and its river downstream, *J. Mar. Syst.*, 66, 161-172, 2007.

Hahm, D., Kim, G., Lee, Y.W., Nam, S.Y., Kim, K.R., Kim, K.: Tidal influence on the sea-to-air transfer of CH₄ in the coastal ocean. *Tellus*, 58B, 88-94, 2006.

635 Hershey, A. E., Northington, R. M., Whalen, S. C.: Substrate limitation of sediment methane flux, methane oxidation and use of stable isotopes for assessing methanogenesis pathways in a small arctic lake, *Biogeochemistry*, 117, 325-336, 2014.

Holgerson, M. A., Raymond, P. A.: Large contribution to inland water CO₂ and CH₄ emissions from very small ponds. *Nature Geosci.*, 9, 222–U150, 2016.

Howard, D. L., Frear, J. I., Pfister, R. M.: The potential for methane carbon cycling in Lake Erie, paper presented at 14th Conference on Great Lakes Research, Int. Assoc. of Great Lakes Res., Ann Arbor, Mich, 1971.

Huttunen, J. T., Alm, J., Liikanen, A., Juutinen, S., Larmola, T., Hammar, T., Silvola, L., Martikainen, P. J.: Fluxes of methane, carbon dioxide and nitrous oxide in boreal lakes and potential anthropogenic effects on the aquatic greenhouse gas emissions, *Chemosphere*, 52, 609–621, 2003.

645 Huttunen, J. T., Väisänen, T. S., Hellsten, S. K., Heikkinen, M., Nykänen, H., Jungner, H., Niskanen, A., Virtanen, M. O., Lindqvist, O.V., Nenonen, O. S., Martikainen, P. J.: Fluxes of CH₄, CO₂, and N₂O in hydroelectric reservoirs Lokka and

650

Porttipahta in the northern boreal zone in Finland, Glob. Biogeochem. Cy., 16, 1-17, 2002.

IPCC, 2013. Climate change 2013: the physical science basis. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, p. 507-507.

ISO 10260: Water quality-Measurement of biochemical parameters-Spectrometric determination of the chlorophyll-a concentration, International Organization for Standardization, Geneva, Switzerland, 1992.

Jirka, A. M., Carter, M. J.: Micro semi-automated analysis of surface and waste waters for chemical oxygen demand, Anal. Chem., 47, 1397-1402, 1975.

Juutinen, S., Rantakari, M., Kortelainen, P., Huttunen, J. T., Larmola, T., Alm, J., Silvola, J., Martikainen, P. J.: Methane dynamics in different boreal lake types, Biogeosciences, 6, 209-223, 2009.

Käki, T., Ojala, A., Kankaala, P.: Diel variation in methane emissions from stands of *Phragmites australis* (Cav.) Trin. ex Steud. and *Typha latifolia* L. in a boreal lake, Aquat. Bot., 71, 259-271, 2001.

Karl, D. M., Tien, G.: MAGIC: A sensitive and precise method for measuring dissolved phosphorus in aquatic environments, Limnol. Oceanogr., 37, 105-116, 1992.

Keller, M., Stallard, F.: Methane emission by bubbling from Gatun Lake, Panama, J. Geophys. Res., 99, 8307-8319, 1994.

- 675 Li, S.Y., Bush, R.T.: Revision of methane and carbon dioxide emissions from inland waters in India, *Glob. Change Biol.*, 21, 6–8, 2015.
- Liang, X., Zhang, X.Y., Sun, Q., He, C. Q., Chen, X. P., Liu, X.Y., Chen, Z. L.: The role of filamentous algae *Spirogyra* spp. in methane production and emissions in streams, *Aquat. Sci.*, 78, 227-239, 2016.
- 680 Liikanen, A., Huttunen, J. T., Murtoniemi, T., Tanskanen, H., Väisänen, T., Silvola, J., Alm, J., Martikainen, P. J.: Spatial and seasonal variation in greenhouse gas and nutrient dynamics and their interactions in the sediments of a boreal eutrophic lake, *Biogeochemistry*, 65, 83–103, 2003.
- Liu, L. X., Xu, M., Lin, M., Zhang, X.: Spatial variability of greenhouse gas effluxes and their controlling factors in the Poyang Lake in China, *Pol. J. Environ. Stud.*, 685 22, 749-758, 2013.
- Liu, L. X., Xu, M., Qiu, S., Shen, R. C.: Spatial patterns of benthic bacteria communities in a large lake, *Int. Rev. Hydrobiol.*, 100, 97-105, 2015.
- Liu, L. X., Xu, M.: Microbial biomass in sediments affects greenhouse gas effluxes in Poyang Lake in China, *J. Freshw. Ecol.*, 31, 109-121, 2016.
- 690 Lorke, A., Bodmer, P., Noss, C., Alshboul, Z., Koschorreck, M.: Technical note: drifting vs. anchored flux chambers for measuring greenhouse gas emissions from running waters, *Biogeosciences*, 12, 14619-14645, 2015.
- Marinho, C. C., Palma-Silva, C., Albertoni, E. F., Trindade, C. R., Esteves, F. A.: Seasonal dynamics of methane in the water column of two subtropical lakes differing in trophic status, *Braz. J. Biol.*, 69, 631-637, 2009.
- 695 McNicol, G., Silver, W. L.: Non-linear response of carbon dioxide and methane emissions to oxygen availability in a drained Histosol, *Biogeochemistry*, 123, 299-306, 2015.

- 700 Marotta, H., Pinho, L., Gudas, C., Bastviken, D., Tranvik, L. J., Enrich-Prast, A.:
Greenhouse gas production in low-latitude lake sediments responds strongly to
warming, *Nat. Clim. Chang.*, 4, 467-470, 2014.
- Massmann, J., Farrier, D. F.: Effects of atmospheric pressures on gas transport in the
vadose zone, *Water Resour. Res.*, 28, 777-791, 1992.
- 705 McGinnis, D. F., Kirillin, G., Tang, K.W., Flury, S., Bodmer, P., Engelhardt, C.,
Casper, P., Grossart, H. P.: Enhancing surface methane fluxes from an
oligotrophic lake: exploring the microbubble hypothesis, *Environ. Sci. Technol.*,
49, 873-880, 2015.
- Miyajima, T., Yamada, Y., Wada, E., Nakajima, T., Koitabashi, T., Hanba, Y. T., Yoshi,
710 K.: Distribution of greenhouse gases, nitrite, and $\delta^{13}\text{C}$ of dissolved inorganic
carbon in Lake Biwa: Implications for hypolimnetic metabolism,
Biogeochemistry, 36, 205–211, 1997.
- Nachshon, U., Dragila, M., Weisbrod, N.: From atmospheric winds to fracture
ventilation: Cause and effect, *J. Geophys. Res.*, 117, 2012.
- 715 Nozhevnikova, A. N., Nekrasova, V., Ammann, A., Zehnder, A. J. B., Wehrli, B.,
Holliger, C.: Influence of temperature and high acetate concentrations on
methanogenesis in lake sediment slurries, *FEMS Microbiol. Ecol.*, 62, 336-344,
2007.
- Ortiz-Llorente, M. J., Alvarez-Cobelas, M.: Comparison of biogenic methane
720 emissions from unmanaged estuaries, lakes, oceans, rivers and wetlands, *Atmos.*
Environ. 59, 328-337, 2012.
- Paganelli, C.V., Rahn, A. A., Wangenstein, O. D.: Diffusion in the gas phase: the
effects of ambient pressure and gas composition, *Respir. Physiol.*, 25, 247-258,
1975.

- 725 Palma-Silva, C., Marinho, C. C., Albertoni, E. F., Giacomini, L. B., Barros, M. P. F.,
Furlanetto, L. M., Trindade, C. R.T., de Assis Esteves, F.: Methane emissions in
two small shallow neotropical lakes: the role of temperature and trophic level,
Atmos. Environ., 81, 373-379, 2013.
- Podgrajsek, E., Sahlström, E., Bastviken, D., Holst, J., Lindroth, A., Tranvik, L.,
730 Rutgersson, A.: Comparison of floating chamber and eddy covariance
measurements of lake greenhouse gas fluxes, Biogeosciences, 11, 4225-4233,
2014a.
- Podgrajsek, E., Sahlström, E., Rutgersson, A.: Diurnal cycle of lake methane flux, J.
Geophys. Res., 119, 236-248, 2014b.
- 735 Poindexter, C. M., Baldocchi, D. D., Matthes, J. H., Knox, S. H., Variano, E. A.: The
contribution of an overlooked transport process to a wetland's methane emissions,
Geophys. Res. Lett., 43, 6276-6284, 2016.
- Rasilo, T., Prairie, Y.T., del Giorgio, P. A.: Large-scale patterns in summer diffusive
CH₄ fluxes across boreal lakes, and contribution to diffusive C emissions,
740 Glob.Change Biol., 21, 1124-1139, 2015.
- Rõõm, E. I., Nõges, P., Feldmann, T., Tuvikene, L., Kisand, A., Teearu, H., Nõges, T.:
Years are not brothers: two-year comparison of greenhouse gas fluxes in large
shallow Lake Võrtsjärv, Estonia, J. Hydrol., 519, 1594-1606, 2014.
- Rooney-Varga, J. N., Giewat, M.W., Duddleston, K. N., Chanton, J. P., Hines, M. E.:
745 Links between archaeal community structure, vegetation type and methanogenic
pathway in Alaskan peatlands, FEMS Microbiol. Ecol., 60, 240-251, 2007.
- Schipper, L. A., Hobbs, J. K., Rutledge, S., Arcus, V. L.: Thermodynamic theory
explains the temperature optima of soil microbial processes and high Q₁₀ values
at low temperatures, Glob. Change Biol., 20, 3578-3586, 2014.

- 750 Schrier-Uijl, A. P., Veraart, A. J., Leffelaar, P. A., Berendse, F., Veenendaal, E. M.:
Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands,
Biogeochemistry, 102, 265-279, 2011.
- 755 [Schultz, M., Faber, E., Hollerbach, A., Schroder, H. G., Guede, H.: The methane
cycling in the epilimnion of Lake Constance, Arch. Hydrobiol., 151, 157–176,
2001.](#)
- Segers, R.: Methane production and methane consumption: a review of process
underlying wetland methane fluxes, Biogeochemistry, 41, 23-51, 1998.
- Serrano-Silva, N., Sarria-Guzmán Y., Dendooven, L., Luna-Guido, M.:
Methanogenesis and methanotrophy in soil: a review, Pedosphere, 24, 291-307,
760 2014.
- Smith, L. K., Lewis, W. M.: Seasonality of methane emissions from five lakes and
associated wetlands of the Colorado Rockies, Global Biogeochem. Cycles, 6,
323- 338, 1992.
- Striegl, R. G., Kortelainen, P., Chanton, J. P., Wickland, K. P., Bugna, G. C.,
765 Rantakari, M.: Carbon dioxide partial pressure and ¹³C content of north
temperate and boreal lakes at spring ice melt, Limnol. Oceanogr., 46, 941-945,
2001.
- Sun, X., Song, C., Guo, Y., Wang, X., Yang, G., Li, Y., Mao, R., Lu, Y.: Effect of
plants on methane emissions from a temperate marsh in different seasons, Atmos.
770 Environ., 60, 277-282, 2012.
- [Townsend-Small, A., Disbennett, D., Fernandez, J. M., Ransohoff, R. W., Mackay, R.,
Bourbonniere, R. A.: Quantifying emissions of methane derived from anaerobic
organic matter respiration and natural gas extraction in Lake Erie. Limnol.
Oceanogr., 61, S356-S366, 2016.](#)

- 775 Tranvik, L., Kokalj, S.: Decreased biodegradability of algal DOC due to interactive effects of UV radiation and humic matter, *Aquat. Microb. Ecol.*, 14, 301-307, 1998.
- Utsumi, M., Nojiri, Y., Nakamura, T., Nozawa, T., Otsuki, A., Seki, H.: Oxidation of dissolved methane in a eutrophic, shallow lake: Lake Kasumigaura, Japan, 780 *Limnol. Oceanogr.*, 43, 471-480, 1998a.
- Utsumi, M., Nojiri, Y., Nakamura, T., Nozawa, T., A Otsuki, A., Takamura, N., Watanabe, M., Seki, H.: Dynamics of dissolved methane and methane oxidation in dimictic Lake Nojiri during winter, *Limnol. Oceanogr.*, 43, 10-17, 1998b.
- Vachon, D., Prairie, Y.T.: The ecosystem size and shape dependence of gas transfer 785 velocity versus wind speed relationships in lakes, *Can. J. Fish. Aquat. Sci.*, 70, 1757-1764, 2013.
- van Bodegom, P. M., Scholten, J. C. M.: Microbial processes of CH₄ production in a rice paddy soil: Model and experimental validation, *Geochimica et Cosmochimica Acta*, 65, 2055-2066, 2001.
- 790 Verpoorter, C., Kutser, T., Seekell, D. A., Tranvik, L. J.: A global inventory of lakes based on high-resolution satellite imagery, *Geophys. Res. Lett.*, 41, 6396-6402, 2014.
- Wang, Y. Y., Yu, X. B., Li, W. H., Xu, J., Chen, Y. W., Fan, N.: Potential influence of water level changes on energy flows in a lake food web, *Chin. Sci. Bull.*, 56, 795 2794-2802, 2011.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373-7382, 1992.

- Wei, D., Xu, R., Tarchen, T., Wang, Y. S., Wang, Y. H.: Considerable methane uptake by alpine grasslands despite the cold climate: *in situ* measurements on the central Tibetan Plateau, 2008-2013, *Glob. Change Biol.*, 21, 777-788, 2015.
- Wik, M., Thornton, B. F., Bastviken, D., MacIntyre, S., Varner, R. K., Crill, P. M.: Energy input is primary controller of methane bubbling in subarctic lakes, *Geophys. Res. Lett.*, 41, 555-560, 2014.
- Wetzel, R.G.: Limnology: Lake and River ecosystems, In: Academic Press. A Harcourt Science and Technology Company, pp: 24570-24577, 2001.
- Xiao, S. B., Liu, W. G., Yang, H., Liu, D., Wang, Y., Peng, F.: Extreme methane bubbling emissions from a subtropical shallow eutrophic pond, *Austin Biometrics and Biostatistics*, 1, 1-6, 2015.
- Xiao, S. B., Wang, Y. C., Liu, D. F., Yang, Z. J., Lei, D., Zhang, C.: Diel and seasonal variation of methane and carbon dioxide fluxes at Site Guojiaba, the Three Gorges Reservoir, *J. Environ. Sci.*, 25, 2065-2071, 2013.
- Xing, Y. P., Xie, P., Yang, H., Ni, L.Y., Wang, Y. S., Tang, W. H.: Diel variation of methane fluxes in summer in a eutrophic subtropical Lake in China, *J. Freshw. Ecol.*, 19, 639-644, 2004.
- Xing, Y. P., Xie, P., Yang, H., Ni, L.Y., Wang, Y. S., Rong, K.W.: Methane and carbon dioxide fluxes from a shallow hypereutrophic subtropical lake in China, *Atmos. Environ.*, 39, 5532-5540, 2005.
- Xing, Y. P., Xie, P., Yang, H., Wu, A. P., Ni, L.Y.: The change of gaseous carbon fluxes following the switch of dominant producers from macrophytes to algae in a shallow subtropical lake of China, *Atmos. Environ.*, 40, 8034-8043 2006,.

Xu, M., Qi, Y.: Spatial and seasonal variations of Q_{10} determined by soil respiration measurements at a Sierra Nevadan forest, *Glob. Biogeochem. Cy.*, 15, 687-696, 2001.

825 Yang, H., Xie, P., Ni, L.Y., Flower, R. J.: Underestimation of CH_4 Emission from Freshwater Lakes in China. *Environ. Sci. Technol.*, 45, 4203-4204, 2011.

Yang, S. S., Chen, I. C., Liu, C. P., Liu, L.Y., Chang, C. H.: Carbon dioxide and methane emissions from Tanswei River in Northern Taiwan, *Atmos. Pollut. Res.*, 6, 52-61, 2015.

830 Yao, X., Wang, S. R., Ni, Z. K., Jiao, L. X.: The response of water quality variation in Poyang Lake (Jiangxi, People's Republic of China) to hydrological changes using historical data and DOM fluorescence, *Environ. Sci. Pollut. Res.*, 22, 3032-3042, 2015.

Ye, X. C., Zhang, Q., Bai, L., Hu, Q.: A modeling study of catchment discharge to Poyang Lake under future climate in China, *Quatern. Int.*, 244, 221-229, 2011.

835 Yvon-Durocher, G., Allen, A. P., Bastviken, D., Conrad, R., Gudas, C., St-Pierre, A., Thanh-Duc, N., del Giorgio, P. A.: Methane fluxes show consistent temperature dependence across microbial to ecosystem scales, *Nature*, 507, 488-491, 2014.

Zhao, Y., Sherman, B., Ford, P., Demarty, M., DelSontro, T., Harby, A., Tremblay, A., Øverjordet, I.B., Zhao, X. F., Hansen, B. H., Wu, B.: A comparison of methods for the measurement of CO_2 and CH_4 emissions from surface water reservoirs: Results from an international workshop held at Three Gorges Dam, June 2012, *Limnol. Oceanogr.: Methods*, 13, 15-29, 2015.

840 Zhu, H. H., Zhang, B.: Poyang Lake-hydrology、biology、sediment、wetland、exploitation andrenovation, In: University of science and technology of China Press. Hefei, pp. 97-99, 1997.

Zhu, D., Wu, Y., Chen, H., He, Y. X., Wu, N.: Intense methane ebullition from open water area of a shallow peatland lake on the eastern Tibetan Plateau, *Sci. Total Environ.*, 542, 57-64, 2016.

Table 1 Seasonal and annual means of CH₄ effluxes with the chamber measurements in Poyang Lake

CH ₄ efflux (mmol m ⁻² day ⁻¹)	2011	2012	2013	2014
Spring (Mar–May)	0.22 ± 0.035b	0.36 ± 0.092bc	0.23 ± 0.16b	0.37 ± 0.084b
Summer (Jun–Aug)	1.34 ± 0.31a	1.21 ± 0.16a	1.36 ± 0.44a	1.44 ± 0.16a
Autumn (Sep–Nov)	0.23 ± 0.12b	0.43 ± 0.14b	0.33 ± 0.12b	0.34 ± 0.16b
Winter (Dec–Feb)	0.11 ± 0.014b	0.23 ± 0.036b	0.14 ± 0.047b	0.23 ± 0.10b
Mean	0.47 ± 0.54a	0.56 ± 0.41ac	0.52 ± 0.55a	0.60 ± 0.56bc

Season	CH ₄ efflux (mmol m ⁻² day ⁻¹)
Spring (Mar–May)	0.30 ± 0.11bd
Summer (Jun–Aug)	1.34 ± 0.32a
Autumn (Sep–Nov)	0.33 ± 0.14b
Winter (Dec–Feb)	0.18 ± 0.077cd

Note: Means with different letters are significantly different as determined by multiple

comparisons on a seasonal scale (one-way ANOVA, post hoc Tukey test, $p < 0.05$) and a pair T-test ($p < 0.05$) on an annual scale.

Table 2 Correlation relationship between seasonal CH₄ efflux and environmental factors

Environmental factors	Correlation coefficient	Environmental factors	Correlation coefficient
Dissolved oxygen	-0.74**	Sediment- NO ₃ ⁻	-0.2
Sediment nitrogen	0.37	Sediment-pH	-0.13
Sediment carbon	0.24	Water-COD	-0.016
pH in the water	-0.29	Water-NO ₃ ⁻	-0.24
Sediment C/N	-0.064	Water-NH ₄ ⁺	-0.36*
Conductivity	-0.37	Water-chla	0.46*
Wind speed	0.008	Water-TN	-0.35*
DOC	-0.015	Water-TP	0.11

Note: Asterisks indicate statistically significant differences between CH₄ efflux and environmental factors (one asterisk, $p < 0.05$; two asterisks, $p < 0.01$).

Table 2.3. Multivariate regressions between seasonal CH₄ efflux and environment factors

No.	Number of variables	Regression Equation	n	R ²	p
Group 1	12	$\text{EffluxCH}_4 = -10.48 + 110.57 \text{ ST} + 65.06 \text{ SNC}$	48	0.65	0.004
Group1 + Group 2	16	$\text{EffluxCH}_4 = -12.66 + 0.57 \text{ ST} + 90.81 \text{ SNC}$	43	0.73	0
Group 1 + Group 2 + Group 3	19	$\text{EffluxCH}_4 = -3.89 + 0.56 \text{ ST} + 102.88 \text{ SNC} - 35.56 \text{ TP} - 0.74 \text{ DO}$	19	0.89	0

Note: Nd means that no variable input to the stepwise regression exists. Variables in group 1 included sediment temperature (ST), sediment total nitrogen content (SNC), water level, DOC content in the water, pH in the sediment, NH₄⁺ and NO₃⁻ concentrations in the water and in the sediment, sediment organic carbon content, the ratio of carbon and nitrogen, and the mean daily wind speed. Variables in group 2 included TN, TP, COD, and Chl *a* contents in the water. Variables in group 3 included DO content, conductivity, and pH in the water.

870

Table 34 Mean CH₄ effluxes in Poyang Lake in comparison with other large lakes

Lake	Lake size (km ²)	Region	Climate	CH ₄ efflux (mmol·m ⁻² ·day ⁻¹)	References
11 lakes	4	Laurentians, Canada	Boreal	4.08 ⁻	Rasilo et al., 2015
26 lakes	47	Chicoutimi, Canada	Boreal	4.08 ⁻	Rasilo et al., 2015
27 lakes	41	Abitibi, Canada	Boreal	4.67 ⁻	Rasilo et al., 2015
46 lakes	474	Chibougamau, Canada	Boreal	0.17 ⁻	Rasilo et al., 2015
20 lakes	7	James Bay, Canada	Boreal	4.08 ⁻	Rasilo et al., 2015
45 lakes	5	Côte-Nord, Canada	Boreal	4.17 ⁻	Rasilo et al., 2015
31 lakes	2	Eastmain, Canada	Boreal	0.58 ⁻	Rasilo et al., 2015
48 lakes	242	Schefferville, Canada	Boreal	0.42 ⁻	Rasilo et al., 2015
Lake Mendota	39.4	North America	Boreal	0.50 ⁻	Fallon et al., 1980
Dillon	43	North America	Boreal	0.61 ⁻	Smith and Lewis, 1992
-Finlen	4.5	Sweden	Boreal	0.02 ⁻	Bastviken et al., 2004
Kevätkä	4	Finland	Boreal	0.22 ⁻	Huttunen et al., 2003
Biwa	674	Japan	Temperate	0.27 ⁻	Miyajima et al., 1997
Kasumigauro	468	Japan	Temperate	0.26 ⁻	Utsunomiya et al., 1998a
Nojiri	4.4	Japan	Temperate	0.06 ⁻	Utsunomiya et al., 1998b
5 lakes	range 1–11, 3436 ^a	Netherlands	Temperate	5.85 ⁻	Schrier-Uijl et al., 2011
-Donghu	27.9	China	Subtropical	4.46 ⁻	Xing et al., 2005
TR lake	71.4	Pantanal, South America	Tropical	0.65 ^a /5.74 ^c	Bastviken et al., 2010
BB lake	36.3	Pantanal, South America	Tropical	0.50 ^a /5.63 ^c	Bastviken et al., 2010
Biantang	3.2	China	Subtropical	4.32 ⁻	Xing et al., 2006
Võrtsjärv	270	Estonia	Boreal	4.28 ^b /2.00 ^f	Räsänen et al., 2014
43 lakes	range 1–10, 782073.8 ^a	worldwide	Mainly boreal	0.12 ⁻	Holgerson and Raymond, 2016
48 lakes	range 10–400, 597789.3 ^a	worldwide	Mainly boreal	0.10 ⁻	Holgerson and Raymond, 2016
6 lakes	>100, 2024015.8 ^a	worldwide	Mainly boreal	0.06 ⁻	Holgerson and Raymond, 2016
Poyang Lake	3283	China	Subtropical	0.54 ⁻	Present study

Lake	Lake size (km ²)	Region	Climate	CH ₄ efflux (mmol m ⁻² day ⁻¹)	References	Sampling period
11 lakes	1	Laurentians, Canada	Boreal	4.08	Rasilo et al., 2015	11/07
Fiolen	1.5	Sweden	Boreal	0.02	Bastviken et al., 2004	Once
31 lakes	2	Eastmain, Canada	Boreal	0.58	Rasilo et al., 2015	14/17 ^a
Keväron	4	Finland	Boreal	0.22	Huttunen et al., 2003	12 times
45 lakes	5	Côte-Nord, Canada	Boreal	1.17	Rasilo et al., 2015	45/07
20 lakes	7	James Bay, Canada	Boreal	1.08	Rasilo et al., 2015	14/67
Dillon	13	North America	Boreal	0.61	Smith and Lewis, 1992	9 times
Lake Mendota	39.4	North America	Boreal	0.5	Fallon et al., 1980	6 times
27 lakes	41	Abitibi, Canada	Boreal	1.67	Rasilo et al., 2015	21/67
26 lakes	47	Chicoutimi, Canada	Boreal	1.08	Rasilo et al., 2015	19/77
16 lakes	171	Chibougamau, Canada	Boreal	0.17	Rasilo et al., 2015	14/27
48 lakes	242	Schefferville, Canada	Boreal	0.42	Rasilo et al., 2015	48/07
Värtsjärv	270	Estonia	Boreal	1.28 ^b /2.09 ^c	Rõm et al., 2014	21 times
6 lake	>100, 2024015.8 ^a	worldwide	Mainly boreal	0.06	Holgerson and Raymond, 2016	Multiple times
18 lakes	range10-100, 597789.3 ^a	worldwide	Mainly boreal	0.1	Holgerson and Raymond, 2016	Multiple times
43 lakes	range1-10, 782073.8 ^a	worldwide	Mainly boreal	0.12	Holgerson and Raymond, 2016	Multiple times
Nojiri	4.4	Japan	Temperate	0.06	Utsunomi et al., 1998 ^b	6 times
5 lakes	range 1-11, 3436 ^c	Netherlands	Temperate	5.85	Schrner-Uijl et al., 2011	twice
Kasumigaura	168	Japan	Temperate	0.26	Utsunomi et al., 1998 ^b	72 times
Biwa	674	Japan	Temperate	0.27	Miyajima et al., 1997	3 times
Biantangang	3.3	China	Subtropical	1.32	Xing et al., 2006	12 times
Donghui	27.9	China	Subtropical	1.46	Xing et al., 2005	48 times
Poyang Lake	3283	China	Subtropical	0.54	Present study	48 times
BB lake	36.3	Pantanal, South America	Tropical	0.50 ^b /5.63 ^c	Bastviken et al., 2010	Once
TR lake	71.4	Pantanal, South America	Tropical	0.65 ^b /5.74 ^c	Bastviken et al., 2010	Once

Note: A means total areas in the given lake size. B means diffusive effluxes and C means total effluxes, including diffusion and ebullition. **D means number of lakes**

measured once/twice. * means 24/0 in 2006, 8/11 in 2007, 0/13 in 2008, 2/10 in 2009, respectively.

Figure Captions

Figure 1. Location of sampling sites in Poyang Lake.

Figure 2. Examples of calculating the slope of total effluxes, including diffusive and ebullitive effluxes. All the concentrations are presented in original (volumetric parts per million-units).

White circles represent the CH_4 concentrations at different sampling times. Grey circles represent the adjusted concentration. Black trendlines represent the data used for the total efflux calculation. The different letters in the figure panels mean different occurrence times for ebullition: no ebullition (a), occurrence of ebullition at 20 min (b), 40 min (c), and 60 min (d), respectively.

Figure 3. Seasonal variations of CH_4 effluxes and sediment temperatures in Poyang Lake.

White circles represent the variation of CH_4 effluxes, and black circles describe the variation of sediment temperature in the 4-year period.

Figure 4. Diel variations of CH_4 effluxes in Poyang Lake.

Different panels present the diel variations of the CH_4 effluxes in 24–25 July 2011 (a), 5–6 September 2012 (b), 13–14 January 2013 (c), and 14–15 January 2015 (d). White circles describe the diel variations of the CH_4 effluxes. Horizontal short dashed lines mean the average value of the diel CH_4 effluxes.

Figure 5. Diel variations of CH_4 effluxes among three sites.

Different panels present the diel variations of the CH_4 effluxes in 24–25 July 2011 (a), 5–6 September 2012 (b), 13–14 January 2013 (c), and 14–15 January 2015 (d).

Relationships between CH_4 effluxes and wind speed in Poyang Lake.

White circles represent the observed values of CH_4 effluxes and wind speed. Different panels mean the variations of CH_4 effluxes at a bihourly interval within a day, including in 24–25 July 2011 (a), 5–6 September 2012 (b), 13–14 January 2013 (c), and 14–15 January 2015 (d), on a diurnal scale (e), and on a seasonal scale (f). Panels e and f include all the measurements during the observation period. We excluded the white crossed circle in figure e in the regression analysis because of a severe cold front.

Figure 6. Relationship between sediment temperature and CH₄ effluxes in Poyang Lake.

White circles represent the observed values of the diurnal mean CH₄ effluxes and sediment temperature in summer, and black circles represent the observed values of the diurnal mean CH₄ effluxes and sediment temperature in the other seasons in the 4-year period. Black lines represent the fitting curves of the relationship between CH₄ effluxes and sediment temperature.

Figure 7. Relationships between CH₄ effluxes and wind speed in Poyang Lake.

White circles represent the observed values of CH₄ effluxes and wind speed. Different panels mean the variations of CH₄ effluxes at a bihourly interval within a day, including in 24–25 July 2011 (a), 5–6 September 2012 (b), 13–14 January 2013 (c), and 14–15 January 2015 (d), on a diurnal scale (e), and on a seasonal scale (f). Panels e and f include all the measurements during the observation period. We excluded the white-crossed circle in figure c in the regression analysis because of a severe cold front.

Fig. 1





Fig. 2



955


960

965

970

|


Fig. 3



带格式的: 字体: 小四

带格式的: 缩进: 左侧: 0 厘米, 悬挂缩进: 2 字符

带格式的: 字体: 小四




975

980


985

Fig. 4



带格式的: 字体: 小四
带格式的: 缩进: 左侧: 0 厘米, 悬挂缩进: 2 字符

带格式的: 字体: 小四




995

1000


1005

Fig.5



带格式的：字体：小四

Fig. 56




带格式的: 字体: 小四

带格式的: 缩进: 左侧: 0 厘米, 悬挂缩进: 2 字符


1015

带格式的: 字体: 小四



1020

Fig.67



带格式的: 字体: 小四

带格式的: 缩进: 左侧: 0 厘米, 悬挂缩进: 2 字符

