

Title:

Biogeochemical cycling and phyto- and bacterio-plankton communities in a large and shallow tropical lagoon (Terminos Lagoon, Mexico) under 2009-2010 El Niño Modoki drought conditions

5

Authors:

Pascal Conan¹, Mireille Pujo-Pay¹, Marina Agab¹, Laura Calva-Benítez², Sandrine Chifflet³,
10 Pascal Douillet³, Claire Dussud¹, Renaud Fichez³, Christian Grenz³, Francisco Gutierrez
Mendieta², Montserrat Origel-Moreno^{2,3}, Arturo Rodríguez-Blanco¹, Caroline Sauret¹,
Tatiana Severin¹, Marc Tedetti³, Rocío Torres Alvarado², Jean-François Ghiglione¹

⁽¹⁾ Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d'Océanographie

15 Microbienne (LOMIC), Observatoire Océanologique, F-66650, Banyuls/mer, France

⁽²⁾ Universidad Autonoma Metropolitana, Departamento de Hidrobiología, México D.F.,
Mexico

⁽³⁾ Aix Marseille Université, CNRS/INSU, Université de Toulon, IRD, Mediterranean Institute
of Oceanography (MIO) UM 110, 13288, Marseille, France

20

Correspondence to: Pascal Conan (pascal.conan@obs-banyuls.fr)

Abstract

25 A large set of biogeochemical [nutrients, dissolved and particulate organic matter, dissolved polycyclic aromatic
hydrocarbons (PAHs)], phytoplanktonic (biomass and photosynthetic activity) and bacterial (diversity and
abundance, including PAH-degrading bacteria, and ectoenzymatic activities) parameters were determined to
understand how the severe drought period relative to the 2009-2010 El Niño Modoki episode influenced
biogeochemical cycling and phyto- and bacterio-plankton communities in Terminos Lagoon (Mexico) under
30 such conditions. During the studied period, the water column of Terminos Lagoon functioned globally as a sink,
and especially as a "nitrogen assimilator", because of high production of particulate and dissolved organic matter
although exportation of autochthonous matter to the Gulf of Mexico was weak. We found that "bottom-up"
control accounted for a large part in the variability of phytoplankton productivity. Nitrogen and phosphorus
stoichiometry mostly accounted for the heterogeneity in phytoplankton and free-living prokaryotes distribution
in the lagoon. In the Eastern part, we found a clear decoupling between areas enriched in dissolved inorganic
35 nitrogen in the North close to Puerto real coastal inlet and areas enriched in phosphate (PO_4) in the South close
to the Candelaria estuary. Such a decoupling limited the potential for primary production resulting in an
accumulation of dissolved organic carbon and nitrogen (DOC and DON, respectively) close to the river mouths.
In the Western part of the lagoon, maximal phytoplankton development resulted from the coupling between
Palizada river inputs of nitrate (NO_3) and particulate organic phosphorus -PP- (but depleted in PO_4) and bacterial
40 activity transforming PP and dissolved organic phosphorus (DOP) to available PO_4 . The Chumpan River only
marginally contributed to PO_4 inputs due to its very low contribution to overall river inputs. The highest
dissolved total PAH concentrations were measured in El Carmen inlet, suggesting an anthropogenic pollution of
the zone probably related to the oil platform exploitation activities in the shallow waters of the South of the Gulf
of Mexico. We also found that a complex array of biogeochemical and phytoplanktonic parameters were the
45 driving force behind the geographical distribution of bacterial community structure and activities. Finally, we
showed that nutrients brought by the Palizada River supported an abundant bacterial community of PAH-
degraders, which are of significance in this important oil production zone.

50

Keywords: biogeochemistry in coastal lagoon, microbial ecology and ecotoxicology, El Niño, lagoon
pollution, Gulf of Mexico, Terminos Lagoon

1. Introduction

55 Coastal lagoons are complex environments, combining features of shallow inland water bodies wholly or partly sealed off from the adjacent coastal oceans, influenced by tide, river input, precipitation *versus* evaporation balance and surface heat balance. Interactions between freshwater and marine sources generate strong gradients of salinity, light and nutrient availability (Hauenstein and Ramírez, 1986). Biological diversity is generally high in these environments (Milessi et al., 2010). Located in the Southern Gulf of Mexico near
60 Campeche sound, Terminos Lagoon is one of the largest tropical coastal lagoon worldwide and its recognised environmental importance and protected status are potentially threatened by petroleum-related industrial activities inshore and offshore (García-Ríos et al., 2013). A first tentative budget of salt and nutrient concluded that Terminos Lagoon was slightly autotrophic on a yearly basis (David, 1999), but this assessment was clearly based on scarce environmental data. Chlorophyll-*a* (CHL) concentration and phytoplankton net production have
65 been reported to respectively range from 1 to 17 $\mu\text{g L}^{-1}$ and from 20 to 300 $\text{gC m}^{-2} \text{a}^{-1}$ (Day et al., 1982), suggesting potential shift from oligotrophic to eutrophic conditions.

In aquatic ecosystems, bacteria utilize a large fraction (up to 90 %) of primary production, since algal carbon exudates might be the principal source for bacterial production (Cole et al., 1988; Conan et al., 1999). Beside the utilization of a considerable part of the available organic matter, bacterioplankton communities also
70 absorb inorganic nutrients thus competing with phytoplankton communities (Conan et al., 2007; Hobbie, 1988). The bulk of organic matter is a highly heterogeneous matrix among which are labile substrates such as proteins or peptides, oligosaccharides, and fatty acids, but most of it is accounted for by complex and refractory substrates (Hoppe et al., 2002). Extracellular enzymes hence are essential to aquatic microorganisms as they allow for the partitioning of complex organic substrates, including high molecular weight compounds which
75 cannot pass through the cell membrane (Arnosti and Steen, 2013). As a function of genetic diversity, the capacity to produce extracellular enzymes is differently distributed in the bacterial community, directly impacting the range of substrates metabolized (Zimmerman et al., 2013). This phenomenon has global-scale implications, since several meta-analysis clearly evidenced differences in the metabolic capacities of microorganisms from temperate, tropical or high latitude waters (Amado et al., 2013; Arnosti et al., 2011). At a
80 local scale, alteration of the evaporation/precipitation balance due to climate change can be challenging especially in the case of coastal lagoon, as it is well known that changes in salinity may alter bacterial diversity and activities (Pedrós-Alió et al., 2000). Local anthropogenic inputs of organic pollutants such as polycyclic aromatic hydrocarbons (PAHs) may also affect bacterial diversity and activities (Aguayo et al., 2014; Jiménez et al., 2011; Rodríguez-Blanco et al., 2010). Indeed, PAHs, which can comprise as much as 25–35 % of total

85 hydrocarbon content in crude oils (Head et al., 2006), are among the most abundant and ubiquitous pollutants in
the coastal environment (González-Gaya et al., 2016). These compounds are recognized by the European and US
environmental agencies as priority pollutants for the aquatic medium due to their toxicity, persistence and ability
to accumulate in the biota (Kennish, 1992). Hence, the presence of PAHs in the marine environment may induce
90 an increase in the indigenous populations of marine bacteria that can break down and utilize these chemicals as
carbon source, the so called “PAH-degrading bacteria” or “PAH degraders”. These bacteria are generally
strongly selected in oil-impacted ecosystems, where they may account for 70 to 90% of the total bacterial
community (Gutierrez et al., 2014; Head et al., 2006).

Despite their importance, few studies consider the bacterial community in tropical inland aquatic
ecosystems (Roland et al., 2010) or coastal lagoons (Abreu et al., 1992; Hsieh et al., 2012; MacCord et al., 2013;
95 They et al., 2013) and almost none dealt with tropical coastal lagoons (Scofield et al., 2015). Though, the present
challenge is to really understand the functioning of such continental to marine environments interface in a
context of growing anthropization. In comparison to the numerous coastal lagoons fringing the Gulf of Mexico,
Terminos Lagoon received moderate scientific attention (Grenz et al., in rev). Among the existing studies, very
few have been conducted on bacterial communities and most of the latter have been based on culture-dependent
100 methods (Lizárraga-Partida et al., 1987; Lizárraga-Partida et al., 1986). However, cultivable bacteria represent a
very small fraction of total present bacteria (<0.1 %; Ferguson et al., 1984) and culture-independent methods are
requested to more accurately assess the diversity and activity of whole bacterial communities in such a vast and
understudied system. Also, Terminos Lagoon is potentially impacted by PAHs, which may come from a
diversity of sources including sea-based activities (spills from ships, platforms and pipelines, ballast water
105 discharge, drilling...) but also rivers, surface runoffs and the atmosphere that carry various urban and industrial
wastes (fuel combustion, traffic exhaust emissions...). Nevertheless, to our knowledge, little is known about the
PAH content in this ecosystem. Even though Noreña-Barroso et al. (1999) have reported PAH concentrations in
the American Oysters *Crassostrea virginica* and Rendon-Von Osten et al. (2007), PAH concentrations in surface
sediments, no data are currently available about dissolved PAH concentrations in surface waters of the Terminos
110 Lagoon.

Our study aims at evaluating the links between: i) biogeochemical (nutrients, dissolved and particulate
organic matter), ii) phytoplanktonic (biomass and photosynthetic activity) and iii) Free-living prokaryotes
(diversity, including PAH-degrading bacteria, and ectoenzymatic activities) parameters in the water column of
Terminos Lagoon (Mexico) after a sustained period of minimum river discharge relative to the 2009-2010 El
115 Niño Modoki episode. After having identified the main sources of nutrients in the lagoon (focused on nitrogen

and phosphorus), we propose a geographical organization of the ecosystem to explain the distribution of the microbial pelagic communities across the lagoon.

120 2. Materials and methods

2.1 Study site and sampling

Terminos Lagoon is a large (1,936 km², volume 4.65 km³) and shallow (average depth 2.4 m) coastal lagoon located in the Mexican state of Campeche (Fig. 1), 18°20' to 19°00' N and 91°10' to 92°00' W. Temperature shows low seasonal variation (27 to 33 °C), but salinity oscillates from brackish to marine waters due to high variability in river runoff (Fichez et al., 2016; Gullian-Klanian et al., 2008). River discharge, precipitation, and groundwater seepage account for 95.44, 4.53 and 0.03 %, respectively. The Chumpan, Candelaria/Mamantel (hereafter Candelaria), and Palizada estuaries account for 5, 19, and 76 %, respectively of freshwater delivered yearly (~12 10⁹ m³ a⁻¹; i.e. about 2.6 times the lagoon volume) to the lagoon (Fichez et al., 2016). The lagoon is connected to the coastal Sea by 2 inlets: El Carmen on the North Western side (4 km long) and Puerto Real on the North Eastern side (3.3 km). About half of the water volume is renewed every 9 days, mostly as a result of tidal exchange. The tide is mainly diurnal, with a mean range of 0.3 m (David and Kjerfve, 1998). Recent results on tidal current modelling (Contreras Ruiz Esparza et al., 2014) revealed a dynamic inshore current entering the lagoon through Carmen inlet, flowing through the southern half of the lagoon and coming out through Puerto Real and a much slower inverse water current flooding the northern central part of the lagoon. That tidally induced hydrodynamic trend generated a counter clock wise circulation gyre located in the centre of the lagoon leeward from Carmen Island.

Samples were collected at 0.2 m depth at 35 stations distributed over the whole lagoon (Fig. 1) from the 21st to the 27th of October, 2009. In 2009, a yearly cumulated discharge of $4.83 \pm 1.71 \cdot 10^9 \text{ m}^3$ broke a historical deficit record over the 1992-2011 period for the Palizada River (average yearly cumulative discharge of $7.19 \pm 4.22 \cdot 10^9 \text{ m}^3 \text{ s}^{-1}$) (Fichez et al., 2016). That exceptional drought period impacted the whole Mesoamerican region during the 2009-2010 El Niño Modoki episode, and resulted in a salinity positive anomaly in Terminos Lagoon that most strongly developed during the post wet season period (Fichez et al., 2016), at the time of our sampling.

A vertical profile of temperature, salinity and fluorescence was carried out at each of the 35 stations with a SeaBird CTD probe (SBE 19) with a precision of 0.01°C for temperature and 0.001 for salinity. Once the

profile completed, water was sampled using a 5L Niskin bottle maintained horizontally at 0.2 m below the surface.

2.2 Nutrients and Dissolved organic matter

150 As soon as the sampling Niskin was retrieved on board, a 40 mL Schott® glass vial previously acid washed was rinsed with sampled water, filled, immediately injected with the fluorometric detection reagent for ammonia determination (as described in Holmes et al., 1999), sealed, and stored in the dark for later analysis at the laboratory. Then, two 30 mL and one 150 mL plastic acid washed vials were then rinsed with sampled water, filled, stored in a specifically dedicated and refrigerated ice cooler, to be later deep-frozen at the laboratory
155 awaiting analysis of dissolved inorganic and organic nutrients as follow.

Nitrate ($\text{NO}_3 \pm 0.02 \mu\text{M}$), nitrite ($\text{NO}_2 \pm 0.01 \mu\text{M}$), phosphate ($\text{PO}_4 \pm 0.01 \mu\text{M}$) and silicate ($\text{Si}(\text{OH})_4 \pm 0.05 \mu\text{M}$) concentrations were measured on a continuous flow autoanalyzer Technicon® AutoAnalyzer II (Aminot and K rouel, 2007) as previously described in Severin et al. (2014). Ammonium ($\text{NH}_4 \pm 10 \text{ nM}$) was detected at nanomolar concentrations by fluorometric detection (Holmes et al., 1999) on a Turner Design Trilogy
160 fluorometer.

Samples for dissolved organic matter (DOM) were filtered through 2 precombusted (24h, 450°C) glass fiber filters (Whatman GF/F, 25 mm). 20 mL were collected for dissolved organic carbon (DOC), into precombusted glass tubes, acidified with orthophosphoric acid (H_3PO_4), and analyzed by high temperature catalytic oxidation (HTCO) (Cauwet, 1999) on a Shimadzu TOCV analyzer. Typical analytical precision is
165 $\pm 0.1\text{--}0.5$ (SD) or 0.2–1 % (CV). 20 mL of samples were collected in Teflon vials for dissolved organic nitrogen (DON) and phosphorus (DOP), and were analyzed by Persulfate wet-oxidation according to Pujo-Pay and Raimbault (1994) and Pujo-Pay et al. (1997).

2.3 Particulate organic matter, chlorophyll and phaeopigment

170 A 4 L plastic acid washed container was also sub-sampling from the Niskin bottle. It was rinsed with sampled water, filled, and stored in a dedicated ice cooler awaiting filtration back at the laboratory as follow. 250 mL of seawater were filtered through a precombusted (24h, 450°C) Whatman GF/F glass filters (25 mm), placed into a Teflon vial and oxidized for particulate organic nitrogen (PON) and phosphorus (PP) measurements (according to Pujo-Pay and Raimbault, 1994). ~1 L was filtered on precombusted (24 h, 450°C) glass fiber filters
175 (Whatman GF/F, 25mm) for particulate organic carbon (POC) and PON measurements. Filters were dried in an

oven at 50°C and stored, in ashed glass vial and in a dessicator until analyses when return from the cruise, on a CHN Perkin Elmer 2400.

For chlorophyll (CHL), 250 mL were filtered on 25 mm diameter Whatman® GF/F filters and immediately stored in liquid nitrogen. CHL and phaeopigment (Phaeo) were later extracted from filters by 100 % methanol (Marker, 1972) and concentrations were determined by the fluorometric technique (Lorenzen, 1966) on a Turner Design Trilogy fluorometer.

2.4 Photosynthetic parameters

Photosynthetic-irradiance parameters (α , P_m^b and I_k) were measured using radioactive ^{14}C -tracer technique (Fitzwater et al., 1982) in a homemade incubator specifically design. 10x60mL Nunc® culture vials were cautiously filled and inoculated with $\text{Na}_2\text{H}^{14}\text{CO}_3$ (final activity of $\sim 0.2 \mu\text{Ci mL}^{-1}$), incubated for 45 min in a 10 light levels irradiance gradient (from 0 to 1327 W m^{-2}), then filtered on Whatman GF/F 25 mm filters, rinsed with 10 % HCl, dried at 45°C for 12 h, and placed into scintillation vials. 10 ml of a liquid scintillation cocktail (Ultima Gold uLLT) were added to the set of scintillation vials 6 h before processing in a Beckman Scintillation Counter. The photosynthetic parameters were determined by fitting each obtained curve with the 'hyperbolic tangent model without photoinhibition' proposed by Jassby and Platt (1976).

2.5 Measurements of dissolved total PAH concentrations

Dissolved total PAH concentrations were determined by using the EnviroFlu-HC submersible UV fluorometer (TriOS Optical Sensors, Germany), a commercially available instrument dedicated to the *in situ* and real time quantification of PAHs in water. The sensor was calibrated in the laboratory before the cruises according to Tedetti et al. (2010) and Sauret et al. (2016). In this work, the mean dissolved total PAH concentrations derived from the sensor are given in ng L^{-1} with a mean coefficient of variation of 10 %.

2.6 Abundance of prokaryotes

Free-living prokaryotes were determined by flow cytometry (Mével et al., 2008). 2 mL seawater samples were fixed with 2 % formaldehyde for 1 h at 4°C. 1 mL sub-sample was incubated with SYBR Green I (Sigma Aldrich, final conc. 0.05% [v/v] of the commercial solution) for 15 min at 20°C in the dark and analysed with a FACS Calibur flow cytometer (Becton Dickinson, San Jose, CA) equipped with an air-cooled argon laser (488 nm, 15 mW). Data acquisition and analysis were done with Cell-Quest software (Becton Dickinson). The

discrimination of the heterotrophs is done by the use of sybr green which induces a green fluorescence and then prokaryotes are separated using the SSC diffraction parameter.

2.7 Total and metabolically active bacterial community structure

210 Nucleic acids were extracted on 0.2 µm-pore-size filter (47 mm, PC, Nucleopore) by filtration of 1 L of pre-filtered (3 µm) water. Co-extraction of DNA and RNA was performed after chemical cell lysis (Ghiglione et al., 1999) with the Qiagen Allprep DNA/RNA extraction kit using manufacturer instructions. DNA and cDNA (by M-MLV reverse transcription of 16S rRNA, Promega) were used as a template for PCR amplification of the variable V3 region of the 16S rRNA gene (*Escherichia coli* gene positions 329–533; Brosius et al., 1981). The primer w34 was fluorescently labelled at the 5'-end position with phosphoramidite (TET, Applied Biosystems). 215 CE-SSCP analysis was performed using the 310 Genetic Analyzer and Genescan analysis software (Applied Biosystems), as previously described (Ortega-Retuerta et al., 2012).

2.8 Extracellular enzymatic activities

220 Aminopeptidase, β-glucosidase and lipase were measured using a VICTOR3 spectrofluorometer (Perkin Elmer) after incubations of 2 h at *in situ* temperature with L-leucine-7-amido-4-methyl coumarin (LL, 5 µM final), MUF-β-D-glucoside (β-Glc, 0.25 µM final) or MUF-palmitate (Lip, 0.25 µM final). These saturated concentrations and optimized time incubations were determined prior to the extracellular enzymatic activities measurement, as previously described (Van Wambeke et al., 2009).

225

2.9 Quantification of PAH-degrading bacteria by Most-Probable-Number

The quantification of PAH-degrading bacteria was performed by the most-probable-number (MPN). A total of 100 µL of each sample was introduced in triplicate in a 48-well microplate with 900 µL of sterile minimum medium, as previously described (Rodríguez-Blanco et al., 2010; Sauret et al., 2016). A mixture of 6 230 PAHs from 2 to 5 rings (naphthalene, fluorene, phenanthrene, fluoranthrene, pyrene and benzo[a]pyrene) prepared in dichloromethane in equimolar concentration was introduced into each well at a final concentration of 10 µg mL⁻¹, as previously described by Sauret et al. (2016). It corresponds to very high concentration of PAH in nature, i.e. 50 times higher than the values found in the harbour of Leghorn (Cincinelli et al., 2001). After 2 weeks of incubation, the change from blue to pink, indicating oxidation of the resazurin contained in the medium 235 was checked and each sample was analysed by flow cytometry. Classical MPN table gave the most probable number of bacteria able to degrade the mixture of six PAHs (Alexander, 1982).

2.10 Statistical analysis

Comparative analysis of 16S rDNA- or 16S rRNA-based CE-SSCP fingerprints was carried out with the
240 PRIMER 6 software (PRIMER-E, Ltd., UK) using Bray-Curtis similarities. We used the similarity profile test
SIMPROF (PRIMER 6) to test the null hypothesis of randomly that a specific sub-cluster can be recreated by
permuting the entry ribotypes and samples, when using hierarchical agglomerative clustering. The significant
branch (SIMPROF, $p < 0.05$) was used as a prerequisite for defining bacterial clusters, and clusters were reported
on non-metric multidimensional scaling (MDS) representation.

245 Canonical correspondence analysis (CCA) was used to investigate the variations in the CE-SSCP
profiles under the constraint of our set of environmental variables, using CANOCO software (version 5.0), as
previously described in Berjeb et al. (2011). Significant variables (i.e. variables that significantly explained
changes in 16S rDNA- and 16S rRNA-based fingerprintings) in our data set were chosen using a forward-
selection procedure. Explanatory variables were added until further addition of variables failed to contribute
250 significantly ($p < 0.05$) to a substantial improvement to the model's explanatory power. Environmental
parameters were previously transformed according to their pairwise distributions, and Spearman's rank pairwise
correlations between the transformed environmental variables were used to determine their significance with
Statel v2.7.

255

3. Results

3.1 Distribution of physical parameters

At the studied period, Terminos Lagoon was characterized by a North West-South East positive
gradient of temperature from >30 to about 27°C (Fig. 2A). Salinity was maximal at Puerto Real inlet (37.50) and
260 along the southern limits of El Carmen Island, intermediate at Candelaria and Chumpan River mouths, and
minimal (21.57) close to the Palizada River (Fig. 2B).

3.2 Distribution of biogeochemical parameters

Nitrate and ammonium concentrations (Fig. 2C and 2D) were maximum close to the Palizada
265 embouchure (16.6 and $0.3 \mu\text{M}$, respectively) and to the Puerto Real inlet ($2.5 \mu\text{M}$ in NO_3 and the highest NH_4
concentration of $1 \mu\text{M}$). In the rest of the lagoon, NO_3 concentrations were quite low and homogeneous (close to

the detection limit of 0.01 μM). NH_4 concentrations were more variable with minimum values on the northern side of the lagoon, and concentration in the range 0.1 to 0.3 μM on the southern inshore side.

270 The distribution pattern for PO_4 (Fig. 2E) significantly differed from N-nutrients. Minimum concentrations ($<0.05 \mu\text{M}$) were measured in the western part of the lagoon under the influence of the Palizada River, indicating very low PO_4 inputs from that river as opposed to nitrogen-nutrients. PO_4 concentrations were also low ($<0.10 \mu\text{M}$) in the centre of the lagoon. The highest PO_4 concentration was measured in front of the Chumpan River (0.17 μM), whereas significant inputs in the Eastern part came from Candelaria River (0.13 μM) and Puerto Real inlet (0.12 μM).

275 The distributions of dissolved organic carbon (DOC; Fig. 2F), nitrogen (DON; Fig. 2G) and phosphorus (DOP; Fig. 2H) concentrations followed a pattern comparable to the one of PO_4 . The highest concentrations of 400, 20 and 1 μM for DOC, DON and DOP, respectively, were measured in the South Eastern part of the lagoon, either in front of the Chumpan or Candelaria estuaries. Maximal DOC and DON concentrations (82% and 95%, respectively) were measured in front of Candelaria River, whereas the maximal DOP concentrations were observed in front of the Chumpan River. Lowest concentrations <200 , 5 and 0.1 μM of DOC, DON and DOP were measured in front of the Palizada mouth and in the case of DOC it even spread along the northern shore of Carmen Island. Significant Spearman's rank correlations ($n=35$, $p<0.05$) were found between DON and DOC ($\rho=0.64$), DOP ($\rho=0.64$) and temperature ($\rho=-0.32$).

285 The 3 rivers were clearly the main sources of particulate organic nitrogen (PON) and phosphorus (PP) in the lagoon (Fig. 3). PON reached a maximum concentration of 9.3 μM in front of the Chumpan estuary and progressively decreased while spreading to the North (Fig. 3A). Concerning PP, the Palizada River was the main source with concentrations close to 0.9 μM , progressively decreasing to 0.6 μM while spreading along the southern shore toward the Chumpan Estuary and 0.5 μM in the north-eastern drift toward Puerto Real passage (Fig. 3B). Significant Spearman's rank correlations ($n=35$, $p<0.05$) were found between PP and PON ($\rho=0.73$), NO_3 ($\rho=0.57$) and salinity ($\rho=-0.56$).

3.3 Photosynthetic pigment and activity

295 Chlorophyll (CHL) and phaeopigment (Phaeo) followed a convergent distribution pattern (Fig. 3C and 3D) with maximum concentrations close or in the vicinity to the Palizada mouth ($>6 \mu\text{gCHL L}^{-1}$ and $\sim 2 \mu\text{gPhaeo L}^{-1}$). A range of 1-6 $\mu\text{gCHL L}^{-1}$ and 1-2 $\mu\text{gPhaeo L}^{-1}$ was encountered in the western part of the lagoon. Concentrations $<1 \mu\text{gCHL L}^{-1}$ and 1-2 $\mu\text{gPhaeo L}^{-1}$ were mostly confined to the eastern part. On a global view, Phaeo accounted for $28 \pm 8 \%$ of CHL on average, hence attesting of rather active phytoplankton communities.

Significant Spearman's rank correlations ($n=35$, $p<0.05$) were found between CHL and Phaeo ($\rho=0.82$) or PP ($\rho=0.74$).

300 The maximum rate of carbon production per unit of chlorophyll at light saturation (P_m^b , Fig. 3E) was minimal ($<0.5 \text{ mgC mgCHL}^{-1} \text{ h}^{-1}$) in the Palizada plume in association with the maximum Phaeo:CHL ratio measured ($>44 \%$). Maximum P_m^b values in excess of $8.0 \text{ mgC mgCHL}^{-1} \text{ h}^{-1}$ were measured close the Chumpan estuary in an area of low Phaeo:CHL ratio ($<25 \%$).

305 3.4 Bacterial abundance and extracellular enzymatic activities

Free-living prokaryotes abundance ranged from 1.0 to $4.8 \cdot 10^6 \text{ cell mL}^{-1}$ (mean= $2.8 \cdot 10^6 \text{ cell mL}^{-1}$, $SD=0.9 \cdot 10^6 \text{ cell mL}^{-1}$, $n=35$), with maximum values observed in the Puerto Real passage and close to the river mouths (Candelaria and Chumpan rivers), except for the Palizada river which showed the highest river-lagoon gradient from maximum to minimal values cited above (Fig. 3F).

310 Cell specific aminopeptidase (Leu-MCA), and phosphatase (MUF-P) activities reached maximum values close to Palizada and Chumpan rivers mouths (33 , and $131.9 \text{ fmol L}^{-1} \text{ h}^{-1} \text{ cell}^{-1}$, respectively (Fig. 4A, and 4B). Cell specific lipase activity (MUF-Lip) was maximum ($10.9 \text{ fmol L}^{-1} \text{ h}^{-1} \text{ cell}^{-1}$; Fig. 4C) from Chumpan river mouth northward towards El Carmen Island, crossing the lagoon approximately in its middle following the isotherms (Fig. 2A). Much lower activities were found over most of the lagoon for all the activities (mean values
315 in $\text{fmol L}^{-1} \text{ h}^{-1} \text{ cell}^{-1}$ are 12.6 ± 8.4 for Leu-MCA, 12.1 ± 24.2 for MUF-P and 2.4 ± 2.6 for MUF-Lip). Significant Spearman's rank correlations ($n=35$, $p<0.01$) were found between aminopeptidase activities and DOC ($\rho=-0.27$), PON ($\rho=0.33$) and to a lesser extent, between phosphatase activities and PO_4 ($\rho=-0.46$), PP ($\rho=0.60$), NO_3 ($\rho=0.69$), CHL ($\rho=0.53$).

320 3.5 Dissolved PAH concentrations and estimated abundance of bacterial PAH-degraders

Dissolved total PAH concentrations (Fig. 5A) were higher close to the El Carmen inlet (332 ng L^{-1}) and relatively lower close to Palizada river (187 ng L^{-1}) and to the Chumpan mouth (166 ng L^{-1}). They were generally lower in the rest of the lagoon ($<130 \text{ ng L}^{-1}$). Quantification by MPN counts showed high enrichment of PAH-degraders close to Palizada river (estimated at $4.6 \cdot 10^4 \text{ cells mL}^{-1}$, equivalent to 4.4% of free-living
325 prokaryotes) (Fig. 5B). Lower values were found close to the Chumpan mouth (estimated at $4.7 \cdot 10^3 \text{ cells mL}^{-1}$, equivalent to 0.2% of free-living prokaryotes), and commonly represented less than 0.1% of the free-living prokaryote abundance in the rest of the lagoon. Quantification by MPN counts showed significant even if low

Spearman's rank correlation with dissolved total PAH concentrations ($\rho=0.37$, $p<0.05$, $n=35$). MPN counts correlation ($p<0.05$, $n=35$) were stronger with PP ($\rho=0.65$) and CHL ($\rho=0.53$).

330

3.6 Spatial distribution of total and metabolically active bacteria by CE-SSCP fingerprints.

Bacterial community structure defined as a function of 16S rDNA-based fingerprints from each sample singled out 3 individual stations (Palizada river, El Carmen inlet and Candelaria river) and aggregated 5 groups of stations (Fig. 6A). Three of those groups included a large number of samples: cluster I grouped 9 stations located in the North-eastern part of the lagoon close to Puerto Real inlet; cluster II grouped 9 stations positioned in the middle of the lagoon up North from Chumpan river to the Carmen Island; cluster III grouped 8 stations situated in the South western of the Carmen Island. Two other groups with fewer stations identified intermediated communities found between the El Carmen inlet and the Palizada River in the western part of the lagoon (cluster V; stations 2, 4, 6) and in the middle of the lagoon, close to the Candelaria river (cluster IV; stations 22, 24, 27).

340

Metabolically active bacterial communities as a function of 16S rRNA-based fingerprints singled out 2 stations (Palizada river and El Carmen inlet) and aggregated 5 groups of stations which are slightly different from the DNA-based clusters (Fig. 6B). Three of those groups included a large number of samples: cluster I formed the largest cluster with 15 stations located in the Eastern part of the lagoon; cluster II grouped 9 stations in the middle of the lagoon up North from Chumpan river to the Carmen Island; cluster III grouped 5 stations in the North Western part of the lagoon, close to El Carmen inlet. Two other groups with fewer stations showed intermediate communities found close to the Palizada river mouth (cluster IV; stations 6 and 8) and further east (cluster V; stations 9 and 12).

345

3.7 Environmental drivers of the total and active prokaryote community structures

To analyse the main environmental factors controlling the spatial distribution of total (Fig. 7A) and active (Fig. 7B) prokaryote communities, we performed a canonical correspondence analysis (CCA). In both DNA- and RNA- based analysis, the cumulative percentage of variance of the species-environment relationship indicated that the first and second canonical axis explained 48 % and 24 % of the total variance, respectively for DNA and 45 % and 31 % for RNA. The remaining axes accounted for less than 14 % of the total variance each, and thus were not considered as significant enough.

355

In the DNA-based CCA, the first canonical axis was positively correlated with NO_3^- and CHL and negatively correlated with concentration of DOC, DOP, DON and oxygen. In the RNA-based CCA, the first

canonical axis was positively correlated with NO_3^- and PAHs and negatively correlated with the concentration of
360 POC, PON, oxygen, salinity, PO_4 and CHL. The concomitant effect of those parameters explained 27 % and
40 % (ratio between the sum of all canonical eigenvalues and the sum of all eigenvalues) of the changes in
bacterial community structure found in the DNA- and RNA-based fractions, respectively (Figure 7).

365 4. Discussion

4.1 Biogeochemical characteristics of Terminos Lagoon under low river discharge conditions

With a contribution of about 76 % to river inputs in the lagoon (Fichez et al., 2016; Jensen et al., 1989),
Palizada River delivers most of the new nitrogen inputs as nitrate and ammonium. High concentrations in
nitrogen were also measured in the Puerto Real inlet, suggesting a second nitrogen source from coastal seawater.
370 These two sources have clearly different impact on primary producer development and activity as shown by the
Phaeo:CHL ratio (<20 % in the vicinity of the Palizada River, but >30 % close to the Puerto Real inlet) and P_m^b
values (low in the Palizada area and higher close to the inlet). So, despite greater chlorophyll degradation
(indicated by high Phaeo concentrations), phytoplanktonic cells were more productive under the influence of
waters from the Gulf of Mexico when compared to those under the river influence. Specifically, in terms of
375 average values for the two zones, with similar nutrient, DOM and POM concentrations, we measured a similar
potential primary production per unit volume (27.5 and 30.2 $\text{mgC m}^{-3} \text{h}^{-1}$ for Palizada River and Puerto Real
inlet respectively), but for about 2-fold lower stock of chlorophyll in the area of the inlet (6.3 and 3.9 mgCHL m^{-3}
for Palizada River and Puerto Real inlet respectively). This is in apparent contradiction with what has been
classically reported on the influence of rivers inputs in coastal areas that generally largely enhanced primary
380 productivity (see for example the Rhone River in the Mediterranean Sea; Pujo-Pay et al., 2006). Decreasing
turbidity along the estuarine to inlet transect is a first factor explaining the seaward offset of phytoplankton
productivity. But higher grazing activity by herbivores in the coastal waters or in the vicinity of the Inlet could
be an explanation to further justify the conjunction of higher Phaeo concentrations together with active
phytoplankton physiology (this should be further study).

385 Moreover, Day *et al.* (1982) demonstrated that small additions of filtered mangrove water had a
stimulatory effect on pelagic primary production in Terminos Lagoon. This observation was later confirmed by
Rivera-Monroy *et al.* (1998), who also evidenced a large temporal variability in stimulating effect, and a rapid
inhibition due to variable humic substance concentrations. The relative decrease of productivity close to the
Palizada plume could be due to humic matter as we also found relatively high concentrations in dissolved PAHs

390 (see hereafter §4.4). Finally, it is clear that bottom-up (nutrients and humic substances) drove the differential responses of phytoplankton productivity in the eastern and western part of the lagoon, certainly in conjunction with grazing activity (top-down control).

At the time of our study, Palizada River and Puerto Real inlet were major sources of nitrogen to the lagoon. Sediments are generally considered to be a significant internal source of nutrients in shallow coastal ecosystems but they may also be a net sink of dissolved nitrogen at least during certain times of the year (Sundbäck et al., 2000; Tyler et al., 2003). Rivera-Monroy et al. (1995a) measured nitrogen fluxes between Estero Pargo (an unpolluted tidal creek), and a fringe mangrove forest in Terminos Lagoon. They reported that mangrove sediments were a sink of NO_3 and NH_4 throughout the year. Denitrification, the dissimilatory reduction of NO_3 to produce N_2O and N_2 , was considered as the main process that contributed to NO_3 loss. However, direct measurements of denitrification rates in the fringe and basin mangroves of Terminos Lagoon indicated low sink of NO_3 (Rivera-Monroy et al., 1995b) on the contrary to what has been evidenced in other mangrove forests (i.e. Twilley, 2013). This was confirmed later by Rivera-Monroy *et al.* (Rivera-Monroy et al., 2007) who hypothesised that most of the inorganic nitrogen was retained in the sediments and not lost via denitrification. They also measured a decoupling between sources of nitrogen and phosphorus and because P is a limiting nutrient, they assumed that the dominant source was tidal flooding as opposed to remineralization from organic matter in the soil. During our study, Origel Moreno (2015) found that benthic carbon mineralization consumed a large proportion (between 67 and 86 %) of the pelagic carbon production. These values are in the higher part of the range calculated for sub-tropical lagoons (Grenz et al., 2010; Machado and Knoppers, 1988) and indicate high biological activity in the sediments. Futhermore, Origel Moreno (2015) estimated that 50 to 95 % of nitrogen was mineralized in the sediment through various N-consuming processes but also that nitrogen was more efficiently mineralized than phosphorus.

Our large scale study considering the whole lagoon brings some information about the potential origin of phosphorus in the water column. It is clear from our measurements that phosphate distribution in the lagoon was disconnected from nitrogen. This impacted the stoichiometry of particulate organic matter (N:P ratio) through the whole lagoon as shown by the surprising and relative low values of PON:PP ratio (<13) at all stations (indicating a particulate nitrogen deficit), except for those located in the southwest part of the lagoon where a canonical Redfield ratio of 16 was measured (Fig. 8). To sustain their growth requirement, primary producers have the ability to decouple their consumption of phosphorus and nitrogen in respect to a variable metabolic plasticity (Conan et al., 2007). In comparison to the previously discussed 2 main sources of NO_3 and NH_4 (Palizada River and Puerto Real inlet) located in the West and North part of the lagoon, we identified two

distinct main sources of PO₄ in Terminos Lagoon during the sampled period: (i) river inputs from Candelaria and Chumpan in the South part, even though their contribution to the overall river discharge is low, and (ii) mineralization of organic phosphorus (PP and DOP) by prokaryotes activity (coherent with ectoenzymatic activities; see hereafter §4.2). Note that the major source of PP in the lagoon was the Palizada River, whereas accumulation of DOP was measured between the Palizada and Chumpan Rivers in the South West of the lagoon. In this area, distribution of dissolved oxygen was minimal compared to the rest of the lagoon, which was coherent with high rates of organic matter mineralization in the water column. Finally during our study, the dominant source of PO₄ was not tidal flooding as hypothesized by Rivera-Monroy *et al.* (2007), but the mineralization of organic matter by free-living prokaryotes in the water column. If that conclusion appears valid in the context of weak river discharges, further studies will be necessary to test its potential extension to other environmental conditions (rainy periods, river flood, tidal amplitude...).

4.2 Relationship between biogeochemical conditions and prokaryotic activities

Our analysis of biogeochemical trends in Terminos Lagoon has been combined with the study of the spatial distribution of prokaryotic extracellular activity. Bacterial aminopeptidase and lipase extracellular activities play a key function in the transformation of biopolymer into small monomers, since a large part of organic matter is in the form of large size molecules but only small molecules (<600 Da) are directly assimilable by bacteria (Weiss *et al.*, 1991). The expression of aminopeptidase activity indicates the absence of direct bacterial assimilation of dissolved organic matter and their ability to actively release enzymes outside the cells (Van Wambeke *et al.*, 2009). Moderate but significant negative correlations were found between aminopeptidase activity per cells and DOC concentration in Terminos Lagoon ($\rho=0.27$, $n=35$, $p<0.01$). Higher DOC concentrations associated with lower aminopeptidase activity suggest a higher amount of labile organic matter for bacteria. The high aminopeptidase activity in the Palizada River plume confirmed the presence of recalcitrant organic matter from terrestrial origin, as opposed to minimum activities in Puerto Real marine waters or in Candelaria mouths, where DOC concentrations were maximal. Lipase activities showed different trends, with higher activities found in the middle of the lagoon up North from Chumpan River to Carmen Island. We previously published that ambient quantity and quality of hydrolysable acyl-lipids clearly coupled with the measurement of their *in situ* hydrolysis rates (Bourguet *et al.*, 2009). The differences between spatial distributions of ectoenzymatic aminopeptidase and lipase activities suggest that organic matter from different composition resided in the central zone of Terminos Lagoon, a result in strong agreement with a recent study on hydrodynamics that identified a large circulation cell in the same central area (Contreras Ruiz *et al.*, 2014).

Unfortunately, the contribution of the protein or lipid pool to total organic matter was not measured at the time of the study, which may have strengthened our hypothesis on the role of the composition of organic matter in the spatial distribution of extracellular enzyme activities. This lack of information may explain the very low or
455 absence of correlation found between extracellular activities and measured biogeochemical parameters.

Phosphatase activity is well known to be controlled by the availability of soluble reactive phosphorus (Van Wambeke et al., 2009). This activity was essentially observed in the vicinity of Palizada River, and not in Puerto Real inlet, the two P-depleted zones which indirectly influence the stoichiometry of particulate organic matter, as discussed above (Fig. 8). Thus, a clear phosphatase activity but P-depleted zone means very low P-
460 availability for phytoplanktonic growth. This observation is consistent with the low phytoplankton productivity indicating a weak C-fixation rates in this zone, strengthens our bottom-up control hypothesis. Extracellular phosphatase activity was significantly ($p < 0.05$, $n = 35$) negatively correlated with PO_4 ($\rho = -0.46$) and positively with PP ($\rho = 0.60$). Our data therefore converge with the model previously proposed by Robadue *et al.* (2004) predicting a different behaviour between the Eastern and Western sides of the lagoon in terms of water budget as
465 well as ecosystem functioning, this distinction being mostly driven by the respective influences of the Palizada River discharge in the West and the Puerto Real marine water inputs in the North East.

4.3 Prokaryotic community structure and ectoenzyme activities

Molecular fingerprinting (such as CE-SSCP) and next-generation sequencing technologies generally
470 yielded converging results (Ghiglione et al., 2005; Ghiglione and Murray, 2012; Ortega-Retuerta et al., 2012; Sauret et al., 2015) evidencing clear shifts in bacterial community structure as a function of changes in biogeochemical characteristics (Ghiglione et al., 2005). Numerous factors can regulate microorganism population dynamics, often simultaneously, and several evidences found in the literature (Berdjeb et al., 2011; Fuhrman et al., 2013; Ghiglione et al., 2008) underlined the importance of relevant statistical analysis to
475 investigate the relative importance of environmental factors in predicting the bacterial community dynamics.

It is generally recognized that the expression of ectoenzyme activities could result from species selection and population dynamics (Martinez et al., 1996) and the zonation of prokaryotic community structure in the Eastern, middle and Western parts of the lagoon agreed with such a paradigm. The community composition in the Eastern part could be divided into two sub-clusters corresponding to the respective influences
480 of Palizada River mouth and El Carmen inlet. Both DNA- and RNA-based fingerprinting showed that Palizada River and El Carmen inlet hosted distinct prokaryotic communities, as previously observed in transition zones such as river (Ortega-Retuerta et al., 2012) or lagoon mouth (Rappé et al., 2000). The relation between

community composition and ectoenzyme activities was particularly evident when considering the lipase and aminopeptidase rates. Lipase activity was magnified in the middle of the lagoon with a South to North increasing
485 gradient from Chumpan River to Carmen Island that coincided with specific communities (cluster II in both DNA- and RNA-based fingerprinting). Other communities were found in the Western part under the influence of the Palizada River, where higher aminopeptidase activity was measured.

The combination of DNA and RNA strengthen our observations as DNA-based analysis alone would have failed to distinguish between active, dormant, senescent or dead cells hence preventing to assess the level of
490 activity of each detected bacterial population (Rodríguez-Blanco et al., 2010). Even though the abundance of bacteria in the sea is high, only a small fraction is considered to be metabolically active (Del Giorgio and Bouvier, 2002). Bacterial growth rate has been shown to correlate with cellular rRNA content (Kemp et al., 1993); therefore, information on cellular activity may be obtained by tracking reverse-transcribed 16S rRNA (Lami et al., 2009). In the present study, we focused on the free-living prokaryotes and disregarded the particle-
495 attached fraction by pre-filtrating the water by 3 µm, which allowed eliminating the problem of DNA eukaryotic chloroplasts that may have biased our results in the context of gradients of productive zones. The combination of DNA and RNA results in Terminos Lagoon showed similar trends with total and active communities presenting a strong zonation between the eastern, middle and western parts of the lagoon, to which could be added smaller transition zones located around major sources of coastal (El Carmen inlet) and river inputs (Palizada and
500 Candelaria). Here, the combination of DNA and RNA showed similar tendencies within the total and active communities presenting eastern, middle and western distribution among the lagoon. These results indicated that most of the free-living bacterial communities detected by molecular fingerprinting (DNA-based) were active (RNA-based) among the lagoon, with the exception of the local transition zones between the lagoon waters and the coastal (El Carmen inlet) or rivers (Palizada and Candelaria).

505

4.4 Biogeochemical parameters and PAHs are driving the prokaryotic community structure

Through the use of direct gradient multivariate ordination analyses, we demonstrated that a complex array of biogeochemical parameters was the driving force behind prokaryotic community structure shifts in Terminos Lagoon. Physico-chemical parameters such as nitrate, oxygen, dissolved organic matter (DOC, DON,
510 DOP) and chlorophyll *a* acted in synergy to explain bacterial assemblage changes on rDNA level. Some differences were observed to explain the geographical patterns of the metabolically active bacterial communities (rRNA level), which salinity, particulate organic matter (PON, PP) and phosphate were needed in addition to nitrate, oxygen and CHL parameters already outlined on rDNA level. The variance explained by the

environmental variables selected by the statistical model only represented 27 % and 40 % of the variability at the DNA and RNA level, respectively. So, further studies will be needed to elucidate the unexplained variance of the model, due to other parameters not taken into account in our study, such as ecological relationships between bacterial community themselves or top-down control by predation and viral lysis (Ghiglione et al., 2016).

The concentration of dissolved total PAHs was also a significant explanatory variable of the metabolically active bacterial community structure. PAHs are considered the most toxic component of crude oil to marine life and are ubiquitous pollutants in the coastal environment (Kennish, 1992). Our study was performed just before the 2010 Deepwater Horizon (DWH) blowout in the Gulf of Mexico, but several offshore oil platforms exist in the shallow waters of Campeche Bank in the southern part of the Gulf of Mexico, such as the one from the Campeche field (Cheek-1) which is only 60 km North from Terminos Lagoon (Warr et al., 2013). The coast of Campeche itself has been also impacted by the 1979 oil spill of the *Ixtoc I* platform, just about 100 km Northwest of Terminos Lagoon (Warr et al., 2013). PAHs concentrations in Terminos Lagoon indicated an input into the lagoon from El Carmen inlet (maximal concentration of 332 ng L⁻¹) that mostly impacted the Eastern part, with concentration <130 ng L⁻¹ in the rest of the lagoon. We observed a high enrichment of PAH-degraders in the South eastern part of the lagoon, with low but significant correlation with PAH concentrations ($\rho=0.37$, $p<0.05$, $n=35$). This enrichment was particularly high (estimated at $4.6 \cdot 10^4$ cell mL⁻¹, equivalent to 4.4 % of the free-living prokaryotes abundance) in the Palizada River mouth. Nitrogen fertilization from allochthonous inputs from the Palizada River may be crucial for PAHs degradation potential in Terminos Lagoon. Indeed, it is well accepted that bacterial degradation of hydrocarbon (carbon source for bacteria) is dependent on the nutrients to re-equilibrate the C:N:P ratio (Sauret et al., 2015; Sauret et al., 2016). Some halotolerant bacteria may have the capability to degrade PAHs and survive in river, lagoon and seawater, such as *Marinobacter hydrocarbonoclasticus* sp. 17 (Grimaud et al., 2012). Further studies using PAH-stable isotopes coupled with pyrosequencing (Dombrowski et al., 2016; Sauret et al., 2016) are necessary to identify the dynamic of these functional communities in Terminos Lagoon. Using similar approaches, previous reports showed that the pollutant content and PAHs, in particular, were responsible for the dynamic of bacterial community structure in the sediment of Bizerte lagoon, Tunisia (Ben Said et al., 2010). Such massive impact of pollutants was not observed here, possibly because of the difference in the degree of pollution between the two areas (moderately contaminated in Terminos Lagoon *versus* highly contaminated in Bizerte). Metabolically active bacterial community structure in the Terminos Lagoon was significantly impacted by PAH-pollution even though it did not exceed the effect of other environmental parameters and their specificity at each geographical location.

5. Conclusions

This study provides a new original set of biogeochemical characteristics for one of the largest shallow tropical coastal lagoon and due to the 2009-2010 El Niño Modoki episode, climatic conditions in Terminos Lagoon region were exceptionally dry at the time of our sampling, hence potentially indicative of future environmental conditions resulting from the predicted trends in climate change in the Centro American region. We evidenced a clear distinction in ecosystem functioning between the east and west parts of the lagoon. Most of the oceanic water entering through the inlets spread toward the south-east where dissolved organic matter accumulated. This area did not support significant phytoplankton development. In the West, we hypothesized a balance shift between a top-down and a bottom-up control to explain the different responses in terms of phytoplankton productivity. The decoupling between nitrogen inputs respectively brought by oceanic waters and the Palizada River, and phosphate inputs from the Chumpan River did not allow for phytoplankton C-fixation. Most of the phytoplankton biomass was aggregated around the Palizada River mouth (which brought most of the freshwater in the lagoon), in a P-depleted area (low phosphate concentration and high bacterial phosphatase activity). Bacterial ectoenzyme activities were mainly observed in the middle of the lagoon, along a south to north cross section stretching from the Chumpan River up to Carmen Island. Maximum mineralization activities were found in this area, which coincided with high extracellular lipase and aminopeptidase activities and low DOC and O₂ concentrations. The lagoon produced significant quantities of particulate and dissolved organic matter thanks to nutrients inputs from the rivers, to uncoupling between nitrogen and phosphate, and to prokaryotic activities, but in the end most of it was internally processed or stored and only a few of this autochthonous matter was exported to the Gulf of Mexico coastal waters. Hence during our study, the water column of Terminos Lagoon functioned globally as a sink, and especially as a "nitrogen assimilator". Highest PAH concentrations were measured in El Carmen inlet, suggesting an anthropogenic pollution of the zone probably related to the oil platform exploitation activities in the shallow waters of the South of the Gulf of Mexico and, more locally, to the efflux from El Carmen harbour that serves as a logistical support to the oil extraction industry. We also evidenced the importance of nitrogen fertilization from the Palizada River, which seems to support an abundant prokaryotic community of PAH-degraders.

Another significant outcome from our study was (i) to link the spatial distribution of ectoenzymatic activities with changes in prokaryotic community structure and (ii) to show that a combination of a complex set of physical and biogeochemical parameters was necessary to explain the changes in prokaryotic community

structure. This study also emphasizes the use of direct multivariate statistical analysis to keep the influence of pollutants in perspective, without denying the role of other physico-chemical variables to explain the dynamic of prokaryotic community structure in polluted areas.

580 Our study provided an extensive dataset efficiently mixing biogeochemical status with information on phytoplankton and prokaryotic structure and dynamics, which has never been measured before in Terminos Lagoon and its outcomes offers a strong base of information and reflexion for future studies on this essential coastal system and the potential environmental conditions that might prevail as a consequence of incoming climate change. Further studies will be needed to compare our dataset with high river input regime conditions and asses how it might affect the observed uncoupling between nitrogen and phosphate as well as the dominant
585 source of phosphorus and its consequence on the primary production and prokaryotic activities. Also, the role of the top-down control should also be investigated in order to better understand the variability of the observed responses.

590

Acknowledgments: The present work was conducted within the frame of the Joint Environmental Study of Terminos Lagoon (JEST) and jointly financed by the French National Program EC2CO-DRIL, the Institut de Recherche pour le Développement (IRD), the Centre National de la Recherche Scientifique (CNRS), the University of “Université de Lille-I”, and the Universidad Autonoma Metropolitana-Iztapalapa (UAM-I). The
595 authors are strongly gratefully to the Instituto de Ciencias del Mar y Limnologia, Universidad Nacional Autonoma de México (ICML-UNAM) for providing full access to their field station in Ciudad del Carmen. We also acknowledge P.A. and M.V. Ghighi for carefully proofreading. M.O-M was financially supported by Bonafont S.A De .C.V. and CONACyT during her PhD work.

600 **References**

- Abreu, P. C., Biddanda, B. B., and Odebrecht, C.: Bacterial dynamics of the Patos Lagoon estuary, southern Brazil (32°S, 52°W): Relationship with phytoplankton production and suspended material, *Estuarine, Coastal and Shelf Science*, 35, 621-635, 1992.
- 605 Aguayo, P., Gonzalez, C., Barra, R., Becerra, J., and Martinez, M.: Herbicides induce change in metabolic and genetic diversity of bacterial community from a cold oligotrophic lake, *World journal of microbiology & biotechnology*, 30, 1101-1110, 2014.
- Alexander, M.: Most probable number method for microbial populations. In: *Methods of Soil Analysis*, Page, A. L., Miller, R.H., Keeney, D.R. (Ed.), American Society of Agronomy, Madison, WI, 815-820 pp, 1982.
- 610 Amado, A. M., Meirelles-Pereira, F., Vidal, L. O., Sarmiento, H., Suhett, A. L., Farjalla, V. F., Cotner, J. B., and Roland, F.: Tropical freshwater ecosystems have lower bacterial growth efficiency than temperate ones, *Front Microbiol*, 4, 167, 2013.
- Aminot, A. and K erouel, R.: Dosage automatique des nutriments dans les eaux marines. *M ethodes en flux continu*, Ed Ifremer-Quae, 2007.
- 615 Arnosti, C. and Steen, A. D.: Patterns of extracellular enzyme activities and microbial metabolism in an Arctic fjord of Svalbard and in the northern Gulf of Mexico: contrasts in carbon processing by pelagic microbial communities, *Frontiers in Microbiology*, 4, 318, 2013.
- Arnosti, C., Steen, A. D., Ziervogel, K., Ghobrial, S., and Jeffrey, W. H.: Latitudinal Gradients in Degradation of Marine Dissolved Organic Carbon, *PLoS ONE*, 6, e28900, 2011.
- 620 Ben Said, O., Goni-Urriza, M., El Bour, M., Aissa, P., and Duran, R.: Bacterial community structure of sediments of the bizerte lagoon (Tunisia), a southern Mediterranean coastal anthropized lagoon, *Microbial ecology*, 59, 445-456, 2010.
- Berdjeb, L., Ghiglione, J. F., Domaizon, I., and Jacquet, S.: A 2-Year Assessment of the Main Environmental Factors Driving the Free-Living Bacterial Community Structure in Lake Bourget (France), *Microbial ecology*, 61, 941-954, 2011.
- 625 Bourguet, N., Goutx, M., Ghiglione, J.-F., Pujo-Pay, M., M evel, G., Momzikoff, A., Mousseau, L., Guigue, C., Garcia, N., Raimbault, P., Pete, R., Oriol, L., and Lef evre, D.: Lipid biomarkers and bacterial lipase activities as indicators of organic matter and bacterial dynamics in contrasted regimes at the DYFAMED site, NW Mediterranean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 1454-1469, 2009.
- 630 Brosius, J., Dull, T. J., Sleeter, D. D., and Noller, H. F.: Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*, *Journal of molecular biology*, 148, 107-127, 1981.
- Cauwet, G.: Determination of dissolved organic carbon (DOC) and nitrogen (DON) by high temperature combustion. In: *Methods of seawater analysis*, Grashoff, K., Kremling, K., and Ehrhard, M. (Eds.), 1999.
- Cincinelli, A., Stortini, A. M., Perugini, M., Checchini, L., and Lepri, L.: Organic pollutants in sea-surface microlayer and aerosol in the coastal environment of Leghorn—(Tyrrhenian Sea), *Marine Chemistry*, 76, 77-98, 2001.
- 635 Cole, J. J., Findlay, S., and Pace, M. L.: Bacterial production in fresh and saltwater ecosystems: a cross-system overview, *Marine ecology progress series*. Oldendorf, 43, 1-10, 1988.
- Conan, P., S ndergaard, M., Kragh, T., Thingstad, F., Pujo-Pay, M., Williams, P. J. I. B., Markager, S., Cauwet, G., Borch, N. H., Evans, D., and Riemann, B.: Partitioning of organic production in marine plankton communities: The effects of inorganic nutrient ratios and community composition on new dissolved organic matter, *Limnology and Oceanography*, 52, 753-765, 2007.
- 640 Conan, P., Turley, C. M., Stutt, E., Pujo-Pay, M., and Van Wambeke, F.: Relationship between Phytoplankton Efficiency and the Proportion of Bacterial Production to Primary Production in the Mediterranean Sea, *Aquatic Microbial Ecology*, 17, 131-144, 1999.
- 645 Contreras Ruiz, A., Douillet, P., and Zavala hidalgo, J.: Tidal dynamics of the Terminos Lagoon, Mexico: observations and 3D numerical modelling, *Ocean Dynamics*, 64, 1349-1371, 2014.
- Contreras Ruiz Esparza, A., Douillet, P., and Zavala-Hidalgo, J.: Tidal dynamics of the Terminos Lagoon, Mexico: observations and 3D numerical modelling, *Ocean Dynamics*, 64, 1349-1371, 2014.
- 650 David, L. T.: Laguna de T rminos, Campeche, Netherlands Institute for Sea Research, Texel, NL, 9-15 pp., 1999.
- David, L. T. and Kjerfve, B.: Tides and currents in a two-inlet coastal lagoon: Laguna de T rminos, M xico, *Continental Shelf Research*, 18, 1057-1079, 1998.
- Day, J. W. J., Day, R. H., Barreiro, M. T., Ley-Lou, F., and Madden, C. J.: Primary production in the Laguna de Terminos, a tropical estuary in the southern Gulf of Mexico, *Oceanologica Acta*, 5, 269-276, 1982.

- 655 Del Giorgio, P. A. and Bouvier, T. C.: Linking the physiologic and phylogenetic successions in free-living bacterial communities along an estuarine salinity gradient, *Limnology and Oceanography*, 47, 471-486, 2002.
- Dombrowski, N., Donaho, J. A., Gutierrez, T., Seitz, K. W., Teske, A. P., and Baker, B. J.: Reconstructing metabolic pathways of hydrocarbon-degrading bacteria from the Deepwater Horizon oil spill, *Nature Microbiology*, 1, 16057, 2016.
- 660 Ferguson, R. L., Buckley, E., and Palumbo, A.: Response of marine bacterioplankton to differential filtration and confinement, *Applied and Environmental Microbiology*, 47, 49-55, 1984.
- Fichez, R., Archundia, D., Grenz, C., Douillet, P., Gutiérrez Mendieta, F., Origel Moreno, M., Denis, L., Contreras Ruiz Esparza, A., and Zavala-Hidalgo, J.: Global climate change and local watershed management as potential drivers of salinity variation in a tropical coastal lagoon (Laguna de Terminos, Mexico), *Aquatic Sciences*, doi: 10.1007/s00027-016-0492-1, 2016. 2016.
- 665 Fitzwater, S. E., Knauer, G. A., and Martin, J.-M.: Metal contamination and its effect on primary production measurements, *Limnology and Oceanography*, 27, 544-551, 1982.
- Fuhrman, J., Follows, M., and Forde, S.: Applying “-omics” Data in Marine Microbial Oceanography, *Eos, Transactions American Geophysical Union*, 94, 241-241, 2013.
- 670 García-Ríos, V., Alpuche-Gual, L., Herrera-Silveira, J., Montero-Muñoz, J., Morales-Ojeda, S., Pech, D., Cepeda-González, M. F., Zapata-Pérez, O., and Gold-Bouchot, G.: Towards a coastal condition assessment and monitoring of the Gulf of Mexico Large Marine Ecosystem (GoM LME): Terminos Lagoon pilot site, *Environmental Development*, 7, 72-79, 2013.
- Ghiglione, J.-F., Larcher, M., and Lebaron, P.: Spatial and temporal scales of variation in bacterioplankton community structure in the NW Mediterranean Sea, *Aquatic Microbial Ecology*, 40, 229-240, 2005.
- 675 Ghiglione, J.-F., Martin-Laurent, F., and Pesce, S.: Microbial ecotoxicology: an emerging discipline facing contemporary environmental threats, *Environ Sci Pollut Res*, 23, 3981-3983, 2016.
- Ghiglione, J.-F., Philippot, L., Normand, P., Lensi, R., and Potier, P.: Disruption of narG, the gene encoding the catalytic subunit of respiratory nitrate reductase, also affects nitrite respiration in *Pseudomonas fluorescens* YT101, *Journal of bacteriology*, 181, 5099-5102, 1999.
- 680 Ghiglione, J. F. and Murray, A. E.: Pronounced summer to winter differences and higher wintertime richness in coastal Antarctic marine bacterioplankton, *Environmental Microbiology*, 14, 617-629, 2012.
- Ghiglione, J. F., Palacios, C., Marty, J. C., Mével, G., Labrune, C., Conan, P., Pujo-Pay, M., Garcia, N., and Goutx, M.: Role of environmental factors for the vertical distribution (0–1000 m) of marine bacterial communities in the NW Mediterranean Sea, *Biogeosciences Discussions*, 5, 2131-2164, 2008.
- 685 González-Gaya, B., Fernandez-Pinos, M.-C., Morales, L., Mejanelle, L., Abad, E., Pina, B., Duarte, C. M., Jimenez, B., and Dachs, J.: High atmosphere-ocean exchange of semivolatile aromatic hydrocarbons, *Nature Geosci*, 9, 438-442, 2016.
- Grenz, C., Denis, L., Pringault, O., and Fichez, R.: Spatial and seasonal variability of sediment oxygen consumption and nutrient fluxes at the sediment water interface in a sub-tropical lagoon (New Caledonia), *Marine Pollution Bulletin*, 61, 399-412, 2010.
- 690 Grenz, C., Origel Moreno, M., Denis, L., Gutiérrez Mendieta, F. J., Fichez, R., Douillet, P., Marquez Garcia, A. Z., Torres Alvarado, R., Calva Benítez, L. G., Álvarez Silva, C., Diaz Ruiz, S., and Gallegos Martinez, M. E.: A review of current knowledge on Terminos Lagoon (Mexico): a major site for Subtropical Marine Ecosystems Ecology studies, *Frontiers in Marine Science*, in rev. in rev.
- 695 Grimaud, R., Ghiglione, J.-F., Cagnon, C., Lauga, B., Vaysse, P.-J., Rodriguez-Blanco, A., Mangenot, S., Cruveiller, S., Barbe, V., Duran, R., Wu, L.-F., Talla, E., Bonin, P., and Michotey, V.: Genome Sequence of the Marine Bacterium *Marinobacter hydrocarbonoclasticus* SP17, Which Forms Biofilms on Hydrophobic Organic Compounds, *Journal of Bacteriology*, 194, 3539-3540, 2012.
- 700 Gullian-Klanian, M., Herrera-Silveira, J. A., Rodríguez-Canul, R., and Aguirre-Macedo, L.: Factors associated with the prevalence of *Perkinsus marinus* in *Crassostrea virginica* from the southern Gulf of Mexico, *Diseases of Aquatic Organisms*, 79, 237-247, 2008.
- Gutierrez, T., Rhodes, G., Mishamandani, S., Berry, D., Whitman, W. B., Nichols, P. D., Semple, K. T., and Aitken, M. D.: Polycyclic aromatic hydrocarbon degradation of phytoplankton-associated *Arenibacter* spp. and description of *Arenibacter algicola* sp. nov., an aromatic hydrocarbon-degrading bacterium, *Appl Environ Microbiol*, 80, 618-628, 2014.
- 705 Hauenstein, E. and Ramírez, C.: The influence of salinity on the distribution of *Egeria densa* in the Valdivian river basin Chile, *Arch. Hydrobiol.*, 1074, 511–519, 1986.
- 710 Head, I. M., Jones, D. M., and Roling, W. F.: Marine microorganisms make a meal of oil, *Nature reviews. Microbiology*, 4, 173-182, 2006.

- Hobbie, J. E.: A comparison of the ecology of planktonic bacteria in fresh and salt water, *Limnology and Oceanography*, 33, 750–764, 1988.
- 715 Holmes, R. M., Aminot, A., K erouel, R., Hooker, B. A., and Peterson, B. J.: A simple and precise method for measuring ammonium in marine and freshwater ecosystems, *Canadian Journal of Fisheries and Aquatic Sciences*, 56, 1801-1808, 1999.
- Hoppe, H. G., Arnosti, C., and Herndl, G. J.: Ecological significance of bacterial enzymes in the marine environment. In: *Enzymes in the Environment: Activity, Ecology, and Applications*, Burns, R. G. and Dick, R. P. (Eds.), Taylor & Francis, Marcel Dekker: New York, NY, USA., 2002.
- 720 Hsieh, W.-C., Chen, C.-C., Shiah, F.-K., Hung, J.-J., Chiang, K.-P., Meng, P.-J., and Fan, K.-S.: Community Metabolism in a Tropical Lagoon: Carbon Cycling and Autotrophic Ecosystem Induced by a Natural Nutrient Pulse, *Environmental Engineering Science*, 29, 776-782, 2012.
- Jassby, A. D. and Platt, T.: Mathematical formulation of the relationship between photosynthesis and light for phytoplankton, *Limnology and Oceanography*, 21, 540-547, 1976.
- 725 Jensen, J. R., Kjerfve, B., Ramsey Iii, E. W., Magill, K. E., Medeiros, C., and Sneed, J. E.: Remote sensing and numerical modeling of suspended sediment in Laguna de terminos, Campeche, Mexico, *Remote Sensing of Environment*, 28, 33-44, 1989.
- Jim enez, N., Vi nas, M., Guiu-Aragon es, C., Bayona, J. M., Albaig es, J., and Solanas, A. M.: Polyphasic approach for assessing changes in an autochthonous marine bacterial community in the presence of Prestige fuel oil and its biodegradation potential, *Applied Microbiology and Biotechnology*, 91, 823-834, 2011.
- 730 Kemp, P. F., Lee, S., and Laroche, J.: Estimating the growth rate of slowly growing marine bacteria from RNA content, *Applied and Environmental Microbiology*, 59, 2594-2601, 1993.
- Kennish, M. J.: Polynuclear aromatic hydrocarbons. Ecology of estuaries: anthropogenic effects, CRC Press, Boca Raton, 1992.
- 735 Lami, R., Ghiglione, J.-F., Desdevises, Y., West, N. J., and Lebaron, P.: Annual patterns of presence and activity of marine bacteria monitored by 16S rDNA-16S rRNA fingerprints in the coastal NW Mediterranean Sea, *Aquatic Microbial Ecology*, 54, 199, 2009.
- Liz rraga-Partida, M. L., Carballo Cruz, R., Izquierdo-Vicuna, F. B., Colwell, R. R., and Chang, I. W.: Bacteriologia de la Laguna de Terminos, Campeche, Mexico, *Anales del Instituto de Ciencias del Mar y Limnologi*, 14, 97-108, 1987.
- 740 Liz rraga-Partida, M. L., Mu oz-Rubio, J., Porras-Aguirre, J., Izquierdo-Vicuna, F. B., and Wong Chang, I.: Taxonomy and distribution of hydrocarbonoclastic bacteria from the Ixtoc-1 area, GERBAM — Deuxi me Colloque International de Bact riologie marine, 633-638, 1986.
- Lorenzen, C. J.: A method for the continuous measurement of in vivo chlorophyll concentration, *Deep Sea Research and Oceanographic Abstracts*, 13, 223-227, 1966.
- 745 MacCord, F., Azevedo, F. D. A., Esteves, F. A., and Farjalla, V. F.: Regulation of bacterioplankton density and biomass in tropical shallow coastal lagoons, *Acta Limnologica Brasiliensia*, 25, 224-234, 2013.
- Machado, E. C. and Knoppers, B. A.: Sediment oxygen consumption in an organic-rich, subtropical lagoon, Brazil, *Science of The Total Environment*, 75, 341-349, 1988.
- 750 Marker, A. F. H.: The use of acetone and methanol in the estimation of chlorophyll in the presence of phaeophytin, *Freshwater Biology*, 2, 361-385, 1972.
- Martinez, J., Smith, D. C., Steward, G. F., and Azam, F.: Variability in ectohydrolytic enzyme activities of pelagic marine bacteria and its significance for substrate processing in the sea, *Aquatic Microbial Ecology*, 10, 223-230, 1996.
- 755 M vel, G., Vernet, M., Goutx, M., and Ghiglione, J. F.: Seasonal to hour variation scales in abundance and production of total and particle-attached bacteria in the open NW Mediterranean Sea (0–1000 m), *Biogeosciences*, 5, 1573-1586, 2008.
- Milessi, A. C., Danilo, C., Laura, R.-G., Daniel, C., Javier, S., and Rodr guez-Gallego, L.: Trophic mass-balance model of a subtropical coastal lagoon, including a comparison with a stable isotope analysis of the food-web, *Ecological Modelling*, 221, 2859-2869, 2010.
- 760 Nore na-Barroso, E., Gold-Bouchot, G., and Sericano, J. L.: Polynuclear aromatic hydrocarbons in American oysters *Crassostrea virginica* from the Terminos Lagoon, Campeche, Mexico, *Marine Pollution Bulletin*, 38, 637-645, 1999.
- 765 Origel Moreno, M.: Variabilit  spatiale et temporelle des cycles biog ochimiques   l'interface eau-s diment dans la lagune de T rminos, Mexique, 2015. Th se de doctorat, Institut M diterran en d'Oc anologie, Universit  d'Aix-Marseille; Ecole Doctorale des Sciences de l'Environnement, 250 pp., 2015.

- Ortega-Retuerta, E., Jeffrey, W. H., Babin, M., Bélanger, S., Benner, R., Marie, D., Matsuoka, A., Raimbault, P., and Joux, F.: Carbon fluxes in the Canadian Arctic: patterns and drivers of bacterial abundance, production and respiration on the Beaufort Sea margin, *Biogeosciences*, 9, 3679-3692, 2012.
- 770 Osten-von Rendon, J., Memije, M., Ortiz, A., and Benitez, J.: Potential sources of PAHs in sediments from Terminos lagoon, Campeche, Mexico, *Toxicology Letters*, 172, Supplement, S162, 2007.
- Pedrós-Alió, C., Calderón-Paz, J. I., MacLean, M. H., Medina, G., Marrasé, C., Gasol, J. M., and Guixa-Boixereu, N.: The microbial food web along salinity gradients, *FEMS Microbiology Ecology*, 32, 143-155, 2000.
- 775 Pujo-Pay, M., Conan, P., and Raimbault, P.: Excretion of dissolved organic nitrogen by phytoplankton assessed by wet oxidation and N-15 tracer procedures, *Marine Ecology Progress Series*, 153, 99-111, 1997.
- Pujo-Pay, M. and Raimbault, P.: Improvement of the wet-oxidation procedure for simultaneous determination of particulate organic nitrogen and phosphorus collected on filters, *Marine Ecology Progress Series*, 105, 203-207, 1994.
- 780 Rappé, M. S., Vergin, K., and Giovannoni, S. J.: Phylogenetic comparisons of a coastal bacterioplankton community with its counterparts in open ocean and freshwater systems, *FEMS Microbiology Ecology*, 33, 219-232, 2000.
- Rivera-Monroy, V. H., Day, J. W., Twilley, R. R., Vera-Herrera, F., and Coronado-Molina, C.: Flux of nitrogen and sediment in a fringe mangrove forest in terminos lagoon, Mexico, *Estuarine, Coastal and Shelf Science*, 40, 139-160, 1995a.
- 785 Rivera-Monroy, V. H., de Mutsert, K., Twilley, R. R., Castañeda-Moya, E., Romigh, M. M., and Davis, I., Stephen E.: Patterns of nutrient exchange in a riverine mangrove forest in the Shark River Estuary, Florida, USA, *Hidrobiológica*, 17, 169-178, 2007.
- Rivera-Monroy, V. H., Madden, C. J., Day, J. W., Twilley, R. R., Vera-Herrera, F., and Alvarez-Guillén, H.: Seasonal coupling of a tropical mangrove forest and an estuarine water column: enhancement of aquatic primary productivity, *Hydrobiologia*, 379, 41-53, 1998.
- 790 Rivera-Monroy, V. H., Twilley, R. R., Boustany, R. G., Day, J. W., Vera-Herrera, F., and del Carmen Ramirez, M.: Direct denitrification in mangrove sediments in Terminos Lagoon, Mexico, *Marine Ecology Progress Series*, 126, 97-109, 1995b.
- 795 Robadue, D. J., Oczkowski, A., Calderon, R., Bach, L., and Cepeda, M. F.: Characterization of the Region of the Términos Lagoon: Campeche, Mexico, University of Rhode Island, 50 pp., 2004.
- Rodríguez-Blanco, A., Antoine, V., Pelletier, E., Delille, D., and Ghiglione, J.-F.: Effects of temperature and fertilization on total vs. active bacterial communities exposed to crude and diesel oil pollution in NW Mediterranean Sea, *Environmental Pollution*, 158, 663-673, 2010.
- 800 Roland, F., Lobão, L. M., Vidal, L. O., Jeppesen, E., Paranhos, R., and Huszar, V. L. M.: Relationships between pelagic bacteria and phytoplankton abundances in contrasting tropical freshwaters, *Aquatic Microbial Ecology*, 60, 261-272, 2010.
- Sauret, C., Böttjer, D., Talarmin, A., Guigue, C., Conan, P., Pujo-Pay, M., and Ghiglione, J.-F.: Top-Down Control of Diesel-Degrading Prokaryotic Communities, *Microbial ecology*, 70, 445-458, 2015.
- 805 Sauret, C., Tedetti, M., Guigue, C., Dumas, C., Lami, R., Pujo-Pay, M., Conan, P., Goutx, M., and Ghiglione, J.-F.: Influence of PAHs among other coastal environmental variables on total and PAH-degrading bacterial communities, *Environ Sci Pollut Res*, 23, 4242-4256, 2016.
- Severin, T., Conan, P., Durrieu de Madron, X., Houpert, L., Oliver, M. J., Oriol, L., Caparros, J., Ghiglione, J. F., and Pujo-Pay, M.: Impact of open-ocean convection on nutrients, phytoplankton biomass and activity, *Deep Sea Research Part I: Oceanographic Research Papers*, 94, 62-71, 2014.
- 810 Sundbäck, K., Miles, A., and Göransson, E.: Nitrogen fluxes, denitrification and the role of microphytobenthos in microtidal shallow-water sediments: an annual study, *Marine Ecology Progress Series*, 200, 59-76, 2000.
- Tedetti, M., Guigue, C., and Goutx, M.: Utilization of a submersible UV fluorometer for monitoring anthropogenic inputs in the Mediterranean coastal waters, *Marine Pollution Bulletin*, 60, 350-362, 2010.
- 815 They, N. H., Ferreira, L. M. H., Marins, L. F., and Abreu, P. C.: Stability of Bacterial Composition and Activity in Different Salinity Waters in the Dynamic Patos Lagoon Estuary: Evidence from a Lagrangian-Like Approach, *Microbial ecology*, 66, 551-562, 2013.
- Twilley, R. R.: Coupling of Mangroves to the Productivity of Estuarine and Coastal Waters. In: *Coastal-Offshore Ecosystem Interactions*, Springer-Verlag, 2013.
- 820 Tyler, A. C., McGlathery, K. J., and Anderson, I. C.: Benthic algae control sediment—water column fluxes of organic and inorganic nitrogen compounds in a temperate lagoon, *Limnology and Oceanography*, 48, 2125-2137, 2003.

- Van Wambeke, F., Ghiglione, J. F., Nedoma, J., Mével, G., and Raimbault, P.: Bottom up effects on bacterioplankton growth and composition during summer-autumn transition in the open NW Mediterranean Sea, *Biogeosciences*, 6, 705-720, 2009.
- 825 Warr, L. N., Friese, A., Schwarz, F., Schauer, F., Portier, R. J., Basirico, L. M., and Olson, G. M.: Bioremediating Oil Spills in Nutrient Poor Ocean Waters Using Fertilized Clay Mineral Flakes: Some Experimental Constraints, *Biotechnology Research International*, 2013, 9, 2013.
- Weiss, M., Abele, U., Weckesser, J., Welte, W., Schiltz, E., and Schulz, G.: Molecular architecture and electrostatic properties of a bacterial porin, *Science (New York, N.Y.)*, 254, 1627-1630, 1991.
- 830 Zimmerman, A. E., Martiny, A. C., and Allison, S. D.: Microdiversity of extracellular enzyme genes among sequenced prokaryotic genomes, *ISME J*, 7, 1187-1199, 2013.

Figure legends

Figure 1: Study site location and distribution of the 35 sampled stations in the lagoon.

835 **Figure 2:** Maps distribution of the physico-chemical parameters measured in the Terminos lagoon in October 2009 for **A.** Temperature ($^{\circ}\text{C}$); **B.** Salinity; **C.** nitrate concentrations (NO_3 in μM); **D.** ammonium concentrations (NH_4 in μM); **E.** phosphate concentrations (PO_4 in μM); **F.** dissolved organic carbon concentrations (DOC in μM); **G.** dissolved organic nitrogen concentrations (DON in μM); and **H.** dissolved organic phosphorus concentrations (DOP in μM).

840 **Figure 3:** as Figure 2 for **A.** particulate organic nitrogen concentrations (PON in μM); **B.** particulate organic phosphorus concentrations (PP in μM); **C.** Total chlorophyll concentrations (CHL in $\text{mg}\cdot\text{m}^{-3}$); **D.** phaeopigments (Phaeo in $\text{mg}\cdot\text{m}^{-3}$); **E.** maximum photosynthetic rate normalized to chlorophyll (P_m^b in $\text{mgC}\cdot\text{mgCHL}^{-1}\cdot\text{h}^{-1}$); **F.** Free-living prokaryotes abundance ($10^6 \text{ cell}\cdot\text{mL}^{-1}$)

Figure 4: as Figure 2 for **A.** aminopeptidase activities ($\text{fmol}\cdot\text{L}^{-1}\cdot\text{h}^{-1}\cdot\text{cell}^{-1}$); **B.** phosphatase activities ($\text{fmol}\cdot\text{L}^{-1}\cdot\text{h}^{-1}\cdot\text{cell}^{-1}$); and **C.** Lipase activities ($\text{fmol}\cdot\text{L}^{-1}\cdot\text{h}^{-1}\cdot\text{cell}^{-1}$)

845

Figure 5: as Figure 2 for **A.** Total dissolved PAHs ($\text{ng}\cdot\text{L}^{-1}$); and **B.** the most-probable-number (MPN in count)


Figure 6: Multidimensional scaling (MDS) plot of the total (**A**) and metabolically active (**B**) prokaryotic community structures as determined from CE-SSCP profiles based on Bray–Curtis similarity index. Clusters were determined according to similarity profile test SIMPROF ($p < 0.05$).

850 **Figure 7:** Canonical correspondence analysis of total (A) and active (B) bacterioplankton community structure from the 35 samples using physico-chemical parameters. Arrows point in the direction of increasing values of each variable. The length of the arrows indicates the degree of correlation with the represented axes. The position of samples relative to arrows is interpreted by projecting the points on the arrow and indicates the extent to which a sample prokaryotic community composition is influenced by the environmental parameter represented by that arrow. The variance explained by the environmental variables selected by the model represent 27 % and 40 % of the variability at the DNA and RNA level, respectively.

855

Figure 8: as Figure 2 for NOP:PP ratio

Figure 1



860

Figure 2




Figure 3






Figure 4




865

Figure 5



870

Figure 6



875

Figure 7




Figure 8

