

Associate Editor Comments

General Comments

Many thanks for your revised manuscript. You have managed to address the referees' comments well, and I'm happy in principle for this study to be published. I am also glad to see that you have made your source data available via figshare. Could I kindly ask you to address a number of remaining, mostly minor points that I list below? These are mainly clarifications in the text.

Specific Comments

Abstract: Please include a real reference when quoting annual Rs figures, i.e. "14.5 ± 1.2 t C ha⁻¹ broadleaved; 12.8 ± 1.0 t C ha⁻¹".

AR: Added.

Line 6: Either "... serve as a C sink or source...", or "... serve as C sinks or sources...".

AR: Changed to "serve as C sinks or sources".

Line 61: Should this be "0-10, 10-20 and 20-30 cm"?

AR: Sorry, yes, it was a typo. It should be "0-10, 10-20 and 20-30 cm"

Line 184: This use of input as a proxy for stock is unclear. Do you have data to back this up? (see also comment below).

AR: Due to a strong seasonal variation in litter input/decomposition (at both sites, but especially at the broadleaved site), litter layer depth and C stocks will vary with time. We think that thoroughly measured total annual litter input rates represent a reasonable estimation for this year's litter stocks and is more accurate than a single point in time measurement of the litter layer depth/stock. A thorough assessment of litter C stocks (multiple-time assessment throughout the seasons) was not carried out during this study. We therefore decided to stick to the litter input as proxy. Nevertheless, a rough estimate during spring 2015 showed that stocks roughly matched annual 2015 litter input rates. We added this to the method text and better explained the effects on uncertainty in the discussion (see below).

Methods (L183-187): We used the litter Q₁₀ together with continuous temperature at 5 cm soil depth to model daily Rh from the litter layer. In order to scale to field fluxes, we used the annual litter input (Table 1) as a proxy for field litter C stocks. A first rough litter assessment in March 2015 showed that litter stocks were in a similar range as the annual litter input at both sites.

Line 258: I'm not sure the word "disabled" works here. Stating that belowground allocation of C by trees is limited should be appropriate to make this point.

AR: We removed the word "disabled".

Line 260: Better: "However, there is..."

AR: Corrected.

Line 269: This is unclear to me. In steady state, litter C input is balanced by litter C losses, but the pool size of litter is not necessarily equal to one year's amount of litter. That depends on turnover rates. Please clarify.

AR: We agree, that annual litter input is not necessarily equal to the actual amount of litter on the forest floor. However, both, litter input rates and litter decomposition rates underlie a strong seasonal variation (especially in the broadleaved forest); depending on the time of the year, an exact quantification of annual litter stocks can thus be quite challenging (e.g. spring VS. fall measurements of litter layer thickness). We therefore think that total litter input rates as a rough proxy for the C stocks of this year's litter constitutes fair a compromise. We are aware that our upscaling approach of annual heterotrophic respiration rates to the field holds a lot of uncertainties; the estimation of litter C stocks is recognized as one of them (please see discussion section). However, the respiratory contribution from the litter layer to total heterotrophic respiration rates was comparably low at both sites. Although the magnitude of predictions seems to be overestimated, we nevertheless think that our method represents an applicable and easy tool to investigate the seasonal course of heterotrophic and autotrophic soil respiration rates. We changed the discussion to:

L 276-279: Using annual litter input as proxy for litter C stocks is a further source of uncertainty. Litter input has temporal patterns and thereby affects litter decomposition dynamics. Such temporal patterns in litter input/decomposition were not reflected in our model. The modeled contribution of the litter layer to total soil Rh was, however, small (Fig. 4), and therefore, the uncertainty related to temporal litter layer dynamics can also be considered as small.

Line 310: It was either similar or not... Do you mean "almost identical"?

AR: Changed to "almost identical".

Lines 315-322: I'm not sure why you give a very approximate soil C budget when stating that you don't actually have the data to do so confidently. This was not your aim, and I don't see that this is a meaningful contribution at this point.

AR: We deleted this chapter and moved the root and litter discussion (L317-323) upwards (now L249-254).

Soil CO₂ efflux from two mountain forests in the Eastern Himalayas Bhutan: components and controls

Norbu Wangdi^{*1,2,}, Mathias Mayer^{*1,}, Mani Prasad Nirola^{1,4,}, Norbu Zangmo^{2,}, Karma Orong^{2,}, Iftekhar Uddin Ahmed^{1,}, Andras Darabant^{1,}, Robert Jandl^{3,}, Georg Gratzner^{1,}, Andreas Schindlbacher³

¹Institute of Forest Ecology, University of Natural Resources and Life Sciences, 1180 Peter Jordan Strasse, Vienna, Austria

²Ugyen Wangchuck Institute for Conservation and Environmental Research, Department of Forests and Park Services Lamai Goempa, Bumthang, Bhutan

³Federal Research and Training Centre for Forests, Natural Hazards and Landscape – BFW, A-1131 Vienna, Austria

⁴National Biodiversity Center, Ministry of Agriculture and Forests, Thimphu, Bhutan

*These authors contributed equally to this work.

Correspondence to: Norbu Wangdi (norwangs@gmail.com)

Abstract

The biogeochemistry of mountain forests in the Hindu Kush-Himalaya range is poorly studied although climate change is expected to disproportionally affect the region. We measured the soil CO₂ efflux (Rs) at a high elevation (3260 m) [mixed](#), and a lower elevation (2460 m) broadleaved forest in Bhutan, eastern Himalayas, during 2015. Trenching was applied to estimate the contribution of autotrophic (Ra) and heterotrophic (Rh) soil respiration. The temperature (Q_{10}) and the moisture sensitivities of Rh were determined under controlled laboratory conditions and were used to model Rh in the field. The higher elevation mixed forest had a higher standing tree stock, reflected in higher soil C stocks and basal soil respiration. Annual Rs was similar between the two forest sites (14.5 ± 1.2 t C ha⁻¹ broadleaved; 12.8 ± 1.0 t C ha⁻¹ [mixed](#)). [Modelled annual contribution of Rh was ~ 65 % of Rs at both sites with a higher heterotrophic contribution during winter and lower contribution during the monsoon season. Rh, estimated from trenching, was in the range of modelled Rh but showed higher temporal variability. Measured temperature sensitivity of Rh was similar at the mixed and broadleaved forest site \(\$Q_{10}\$ 2.2- 2.3\) under intermediate soil moisture but decreased \(\$Q_{10}\$ 1.5 at both sites\) in dry soil. Rs closely followed the annual course of field soil temperature at both sites. Co-variation between soil temperature and moisture \(cold-dry winters, warm-wet summers\) likely was the main cause for this tight relationship. Under the \[prevailing\]\(#\) weather conditions, a simple temperature-driven model was able to explain more than 90 % of the temporal variation in Rs. Longer time series and/or experimental climate manipulations are required to understand the effects of eventually occurring climate extremes such as monsoon failures.](#)

Keywords: Himalaya, soil CO₂ efflux, autotrophic soil respiration, heterotrophic soil respiration, incubation, temperature sensitivity, moisture sensitivity

1 Introduction

Carbon dioxide (CO₂) efflux from soil (= soil respiration; Rs) is one of the major fluxes in the global C cycle, affects atmospheric CO₂ concentrations and [potentially](#) feeds back on global climate change (Reichstein et al., 2003; [Frey et al., 2013](#); [Wang et al., 2014](#); Hashimoto et al., 2015). Counteracting to C uptake via photosynthesis, Rs primarily determines whether forest ecosystems serve as C sinks or sources to the atmosphere (Bolstad et al., 2004; Dixon et al., 1994; Schlesinger and Andrews, 2000). The current function of forests as global C sink (Stocker, 2014; Janssens et al., 2003) could weaken or even turn into the opposite if climate change disproportionally accelerates respiratory processes such as Rs (Cox et al., 2000). Rs consists of an autotrophic component (Ra; root and rhizosphere respiration), which is closely linked to C gain by photosynthesis and a heterotrophic component (Rh), which is the respiratory product of soil organic matter (SOM) decomposition. While the source of Ra is recently assimilated CO₂, Rh can release stored soil C to the atmosphere. For better prediction of the response of forest C cycling to climate change, it is crucial to understand how Rs and its components are affected by changing environmental parameters such as temperature and moisture (Davidson and Janssens, 2006; [Sierra et al., 2015](#)). Rates and climate sensitivity of Rs, Ra and Rh can vary among forest ecosystem type and climatic region (Hashimoto et al., 2015). So far, research has focused on the temperate and boreal areas of the northern hemisphere whereas remote forested areas are still largely uninvestigated (Bond-Lamberty and Thomson, 2010).

The Hindu Kush-Himalaya range represents a region, where research on forest biogeochemistry is gaining momentum (Pandey et al., 2010; Sundarapandian and Dar, 2013; Sharma et al., 2010b; Dorji et al., 2014b; Ohsawa, 1991; Wangda and Ohsawa, 2006a; Tashi et al., 2016; Verma et al., 2012). It extends over 4.3 million km² across eight countries with an average forest cover of approximately 20 % (Schild, 2008), ranging from lowland tropical forest to high altitudinal forests up to ~ 4900 m (Liang et al., 2016; Schickhoff, 2005). Situated in the eastern Himalayas, Bhutan shows a forest cover of 70 % (DoFPS, 2011). Most forests in Bhutan are natural old growth (Ohsawa, 1987), store high amounts of C in biomass and soil (Dorji et al., 2014a; Sharma and Rai, 2007) and serve as an important regional C sink (FAO, 2010). As climate change is expected to intensify in the Himalaya region (Shrestha et al., 2012; Singh, 2011; Xu and Grumbine, 2014; Tsering et al., 2010; Xu et al., 2009), the effects on forest C cycling could have implications not only regionally, but also on a global scale.

[With the objective of a better understanding of soil C cycling of mountain forest ecosystems, we studied Rs, its components \(Ra, Rh\), as well as the effects of environmental drivers such as temperature and moisture at a moderately high altitude cool temperate mixed forest and a lower altitude cool temperate broadleaved forest in Bhutan. These forest types cover large areas of the eastern Himalayas.](#)

2 Materials and methods

2.1 Site description

Two representative forest ecosystems for the eastern Himalayas (Wikramanayake, 2002), a cool temperate conifer dominated **mixed** forest and a cool temperate broadleaved forest, were studied at Thimphu and Wangduephodrang districts, Bhutan. The cool temperate **mixed** forest (Grierson and Long, 1983) was situated on a south-east facing slope close to the top of a mountain ridge (elevation 3260 m a.s.l.). The cool temperate broadleaved forest was situated on an east facing gentle slope along the same mountain ridge ~ 11 km eastwards (elevation 2640 m a.s.l.). Sites will be referred to as “**mixed** forest” and “broadleaved forest” in the further text. The mixed forest was dominated by *Tsuga dumosa* along with *Picea spinulosa*, *Quercus semecarpifolia*, *Abies densa*, and *Taxus baccata*. The broadleaved forest was dominated by *Quercus lanata* and *Quercus griffithii*. Soils at the mixed forest were Cambisols. Soils at the broadleaved forest were Luvisols. A detailed site and soil description and the comparison are given in Table 1. The current study was aligned within a larger-scale throughfall manipulation experiment, which consisted of control and temporarily roofed areas within each forest type. For this study, we randomly distributed all our plots within the control areas (~ 1500 m² each) of the throughfall manipulation experiment.

2.2 Field measurements

Basic climate parameters were measured using automatic weather stations located at a distance of approx. one kilometer from the sites at the same elevation. Data was recorded at 15 min intervals on a Decagon-EM50 data logger (Decagon Devices Inc., Pullman, WA, USA). The automatic weather stations recorded precipitation with an ECRN-100 rain gauge (Decagon Devices Inc., Pullman, WA, USA), and air temperature and relative humidity with a VP-3 vapor pressure, temperature and relative humidity sensor (Decagon Devices Inc., Pullman, WA, USA).

Stand and soil inventories were carried out in March and April 2014 at both **sites** covering an area of ~ 1500 m² each. The location, height and the diameter at breast height of all trees having a dbh > 10 cm were assessed. The basal area was calculated for each tree species. Standing volume was estimated based on species-specific volume equations developed by Paul Lawmans (1994), Forest Survey of India (1996) and Department of Forests and Park Services, Bhutan (2005). Aboveground litter-fall was collected monthly using mesh-traps (n = 10) per site, with an area of 1.0 m² (100 × 100 cm). Litter was dried at 80 °C and the C content was assumed to be 50 % of the dry weight (de Wit et al., 2006). Soil samples were collected from the 0-10, 10-20 and 20-30 cm mineral soil layers of four locations at both sites in May 2014. Soil samples were sieved (2 mm) and dried (105 °C, 48 h). Soil organic C (SOC) of a ground (Pulverisette 5, Fritsch, Germany), 0.1 g subsample was measured by means of the dry combustion technique using a CN Analyser (TruSpec® CN, LECO Inc., Michigan, USA). Soil organic C stocks (t ha⁻¹) were calculated for each horizon by multiplying the SOC concentration (%) by the bulk density (g cm⁻³) and the depth of the horizon (cm). **Fine root (≤ 2 mm) biomass was estimated by the soil-core method (Makkonen and Helmisaari, 1999) in spring 2014 at both sites. We used a cylindrical soil corer (7.5 cm diameter) for sampling. The extracted samples were divided into three depth sections of 0-10, 10-20 and 20-30 cm. After washing and**

sorting (live roots and necromass), roots were dried at 70 °C to constant mass before weighing dry biomass. Contribution of fine root C was estimated as 50 % of the plant tissue.

Rs was measured at both sites once every three weeks from April 2015 to December 2015 at 10 randomly chosen plots (n = 10) at each. To cover the within-plot variability, Rs was measured at four positions within each plot (total 40 positions per site). We used a portable infrared gas analyzer (EGM-4, PP-Systems, Amesbury, USA) with an attached soil respiration chamber (SRC-1, PP-Systems, Amesbury, USA) for Rs measurements. Prior to measurements (March 2015), we installed permanent collars (total height 5 cm, 2-3 cm inserted into the soil, diameter 10 cm) at each plot which served as a base for Rs measurements. Rs was estimated by a linear fit to the increasing headspace CO₂ concentration over time (chamber closure time 90 seconds). A soil respiration measurement campaign lasted for ~ 5 h at each site. Measurement order among plots and collars was fully random to avoid bias from temporal variations in Rs.

We installed two trenching plots at each site in April 2014 (1 yr prior soil efflux CO₂ measurements) to estimate the relative contributions of Ra and Rh. Trenches (1.5 x 1.5 m) were dug to ~ 1 m depth, and all roots within the trenches were cut. The trenches were sealed with double layered plastic foil in order to restrict tree root ingrowth. Adjoining to each trenched plot, a corresponding control plot of the same size was established. Each trenched and control plot hosted three collars for Rs measurements. We measured soil CO₂ efflux at trenched and corresponding control plots after finishing regular Rs measurements (same day).

Volumetric soil water content (0-20 cm soil depth; (vol. %)) was measured in the center of each plot (Rs plots, trenched plots, control plots) using a portable Field Scout TDR meter (Spectrum Technologies, Inc. Aurora, USA) during Rs measurements. Soil temperature at 5 cm soil depth was measured with a handheld thermometer probe (Hanna Instruments, Germany) at each Rs measurement location. Soil temperature and soil moisture were measured continuously at soil profile pits (two pits per site) with five combined soil temperature-moisture sensors (TM-5; Decagon Devices, Inc., Pullman, WA, USA) at soil depths ranging from 5 to 120 cm. Data was recorded at 15 min intervals on Decagon-EM50 data loggers (Decagon Devices, Inc., Pullman, WA, USA).

2.3 Laboratory incubation

About 500 g of mineral soil (0-10 cm depth) and approximately 250 g of forest floor litter were sampled at six random locations (n = 6) at each site in mid-September 2015. The mineral soil was homogenized and sieved (2 mm mesh) and stored at 4 °C at field moisture for one week prior to transport from Bhutan to Austria for further processing. Forest floor litter was not sieved. Upon arrival in Austria, mineral soil samples were further divided into 3 sub-samples to account for potential soil heterogeneity at individual sampling locations. Samples were filled into 200 cm³ stainless steel cylinders at approximate field bulk density (~ 0.5 g dry weight cm⁻³ for mineral soil; ~ 0.1 g dry weight cm⁻³ for forest floor). In total, we incubated 36 sub-samples (cylinders) for mineral soil and 12 sub-samples for the forest floor litter. Filled cylinders were kept at 4 °C for 5 days for equilibration before incubation. Soil CO₂ efflux (= Rh) was measured using a fully automated incubation system. During incubation,

samples were put into 2 l containers and their CO₂ efflux was determined by a dynamic closed – chamber system (Pumpanen et al., 2009). For CO₂ measurements, containers were sequentially connected to an infrared gas analyzer (SBA-4, PP Systems International Inc., Amesbury, MA, USA) by means of a tubing system. In the meanwhile, disconnected containers were ventilated by means of an air pump in order to prevent internal CO₂ enrichment. Wet tissues were put into containers in order to prevent samples from drying out during incubations; moisture loss was thereby negligible (< 2 vol. %). CO₂ concentration within connected containers were measured for 6 minutes with a recording interval of 10 sec. Rates of CO₂ efflux were calculated from the headspace CO₂ increase during 2 – 6 minutes, after Pumpanen et al. (2009).

Incubations proceeded in two steps. We first incubated at different soil temperatures to assess the temperature sensitivity of Rh. In a second step, we incubated under different soil moisture contents to assess the sensitivity of Rh to changes in soil moisture. In addition, we repeated the temperature-runs with wet (140 % gravimetric water content (grav. %)) and dry (30 grav. %) soil in order to test for effects of soil moisture on the temperature sensitivity of Rh. In-between incubations, soil cores were stored in a cold room (+ 4 °C). During storage, soil moisture was kept constant by periodical water addition.

Temperature-incubation started with mineral soil. Soil temperature was increased from 5 °C until 25 °C in 5 °C steps, with each temperature step lasting for 6 h. At each temperature step, efflux measurements were repeated three times for each cylinder; to account for a warm up period between the individual temperature steps only a calculated mean value of the latter two measurements was used for further analysis. After finishing the temperature run, we re-measured Rh at 10 °C to assess and correct for potential effects of labile C loss during the ~ 30 h incubation. The forest floor litter was incubated under the same procedure as mineral soil.

After the temperature-incubation, we set soil moisture of all mineral soil sub-samples to 80 grav. %, incubated at constant 15 °C for 6 h and measured Rh as described above. Afterwards, the three sub-samples from each sampling location were split into (i) a sub-sample that was kept at constant soil moisture (80 grav. %), (ii) a sub-sample that was allowed to dry out (60 to 15 grav. %), and (iii) a sub-sample that was progressively watered (100 to 160 grav. %). In-between repeated incubations (all at 15 °C for 6 h) cylinders were taken out from incubation containers and were stored at 4 °C. The whole moisture-incubation procedure lasted for 10 weeks with ~ two-weekly intervals between incubations (time limiting step was soil drying). We used Rh from the sub-samples which had been kept at constant moisture to correct for potential decreases in Rh due to a loss in labile C throughout the experiment. After finishing all incubations, samples were dried and actual bulk density, as well as actual gravimetric (grav. %) and volumetric soil moisture (vol. %) of each sub-sample (cylinder), was calculated and their total C content was determined (TruSpec® CN, LECO Inc., Michigan, USA). Rh rates were expressed as $\mu\text{mol CO}_2 \text{ kg C}^{-1} \text{ s}^{-1}$.

2.4 Data analysis

Effects of site on field Rs, soil temperature and moisture were tested by means of repeated-measures ANOVA with a mixed-effects model structure (Pinheiro and Bates, 2000). The significance level for this and all other analyses

was set at $P < 0.05$. The relationship between soil temperature and Rs was fitted by an exponential function (Buchmann, 2000):

$$R = \beta_0 \cdot e^{(\beta_1 \cdot T)} \quad (1)$$

where R ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) is the measured Rs, T ($^{\circ}\text{C}$) is the soil temperature at 5 cm depth, and b_i are model parameters. Equation (1) was fitted to the daily averages of each site as well as to the individual plot data. Basal respiration rates at 10°C soil temperature (R_{s10}) were subsequently calculated (using Eq. (1)) for each site. One sampling date (2015 Jul 16) was excluded from this analysis because heavy rain occurred during measurements. The relationship between Rs and soil moisture was tested by fitting a polynomial function obtained from lab incubation (see further below). Cumulative annual Rs of both sites and both years were calculated by linear interpolation of field Rs between measurement dates of each individual plot (the area beneath the curves in Fig. 1 d). In addition, model parameters of Eq. (1), together with daily field soil temperatures at 5 cm depth were used to calculate a projected daily field Rs. To account for a spatial variation in soil temperature, continuously measured data were adjusted to discontinuously measured plot-data by linear modelling. Cumulative annual Rs rates were calculated by averaging the summed-up daily plot Rs values.

Average Rh rates from laboratory incubations were calculated for each site, soil horizon (mineral soil, forest floor litter) and temperature step ($5 - 25^{\circ}\text{C}$), respectively. Equation 1 was fitted to the temperature-incubation data separately for each site and soil horizon. Basal heterotrophic respiration rates at 10°C (R_{h10}) were calculated for each site. Temperature sensitivity (Q_{10}) of Rh was calculated as follows:

$$Q_{10} = e^{(10 \cdot \beta_1)} \quad (2)$$

where Q_{10} is the factor by which Rh changes at a temperature change of 10°C , and β_1 is the model parameter derived from Eq. (1). To determine the relationship between soil moisture and Rh, we fitted a polynomial function to the moisture-incubation data:

$$R = \beta_0 + \beta_1 \cdot VWC + \beta_2 \cdot VWC^2 \quad (3)$$

where R is the measured CO_2 efflux from soil samples (R_h), β_i are model parameters and VWC is the volumetric water content of the samples. Effects of soil moisture on Q_{10} values were tested by means of one-way ANOVA and Tukey's post-hoc tests.

We followed two approaches to estimate the contribution of R_a and R_h in the field. In a first approach, we used the trenching data, assuming that the CO_2 efflux from the trenched plots represented solely R_h , while the CO_2 efflux from adjacent control plots represented R_s , and accordingly, the difference between trenched and control plot CO_2 efflux represented R_a . As trenched plots lack water uptake by tree roots, they were regularly wetter than control plots. We accounted for that by correcting the soil CO_2 efflux for the difference in soil moisture by using Eq. (3) (see Schindlbacher et al. (2009) for details).

In a second approach, we applied the response functions of R_h derived during laboratory incubation together with field soil C stocks and field climate data. This allowed an alternative way to estimate the contribution of R_h in the field (Gough et al., 2007; Kutsch et al., 2010). Model parameters derived from Eq. (1) together with continuously measured temperature data from 5 cm soil depth were used to model daily R_h from the litter and from the mineral soil in 0 – 10 cm depth respectively. Model parameters for mineral soils together with continuous measurements of soil temperature in 20 cm depth were further used to model daily R_h from the mineral soil in 10 – 30 cm depth. Predicted R_h rates ($\mu\text{mol CO}_2 \text{ kg C}^{-1}$) were multiplied by the C stocks (kg C m^{-2}) of the respective soil layer. We used the litter Q_{10} together with continuous temperature at 5 cm soil depth to model daily R_h from the litter layer. In order to scale to field fluxes, we used the annual litter input (Table 1) as a proxy for field litter C stocks. A first rough litter assessment in March 2015 showed that litter stocks were in a similar range as the annual litter input at both sites. This procedure enabled us to upscale R_h to the whole soil profile in the field (Kutsch et al., 2010). To account for a moisture response as well, predicted R_h rates were also corrected for soil moisture conditions in the field. For that, model parameters derived from Eq. (3) were used to calculate R_h rates at actual moisture conditions in the field (from continuous moisture data) and at initial moisture conditions of the soil samples during incubation (mixed forest: 33 vol. %, broadleaved forest: 35 vol. %, litter: 46 vol. %); their relative difference was subsequently used to correct R_h rates predicted with Eq. (1). Since litter soil moisture was not regularly measured in the field, we applied the same moisture parameters and continuous soil moisture records as for mineral soil (0-10 cm). The R code of the empirical model is provided in the supplement information (S1).

3. Results

Air and soil temperatures were $\sim 4^{\circ}\text{C}$ higher at the lower elevation broadleaved forest (Table 1) with a stable trend throughout both study years (Fig. 1a). Air temperatures reached a maximum of 29.6°C and 22.6°C at the broadleaved and mixed forest, respectively. Winter air temperatures dropped slightly below freezing at the mixed forest which showed ephemeral snow cover. Soil temperatures remained above freezing at both sites during the full study period (Fig. 1b). Annual precipitation in 2015 was similar at both sites (mixed 1167 mm, broadleaved 1120 mm). Both sites received the maximum rainfall (60-75 % of annual precipitation) during the peak monsoon months (Jun, Jul and Aug). Soil moisture was significantly higher at the broadleaved forest during summer (Fig. 1c). During the dry season (Nov – Apr), manually measured soil moisture decreased to < 20 vol. % at both sites. Continuous soil moisture records indicated accelerated drying at the broadleaved forest (Fig. 1c).

Aboveground and below-ground C stocks were markedly higher in the mixed forest (Table 1). Standing volume was 1066 and $464\text{ m}^3\text{ ha}^{-1}$, at the mixed and broadleaved forest, respectively. Mineral soil organic C stocks down to 30 cm soil depth were 142 and 90 t ha^{-1} and leaf litter inputs (2015) were 3.5 and 3.4 t C ha^{-1} at the mixed and broadleaved forest, respectively. Fine root biomass (0-30 cm mineral soil) was lower at the mixed forest (2.3 t C ha^{-1}) when compared to the broadleaved forest (3.2 t C ha^{-1}).

Rs did not differ significantly between the two sites (mean Rs broadleaved: $4.2 \pm 0.7\text{ }\mu\text{mol CO}_2\text{-C m}^{-2}\text{ s}^{-1}$, mixed: $4.0 \pm 0.6\text{ }\mu\text{mol CO}_2\text{-C m}^{-2}\text{ s}^{-1}$) but basal respiration rates (Rs_{10}) were higher at the mixed forest (Fig. 2a). Cumulative annual Rs were $14.3 \pm 0.5\text{ t C ha}^{-1}$ for the broadleaved and $13.0 \pm 0.5\text{ t C ha}^{-1}$ for the mixed forest when calculated by linear interpolation between measurement dates. These values were very close to the ones obtained by the modelling approach (Eq. (1), $14.5 \pm 1.2\text{ t C ha}^{-1}$, broadleaved and $12.8 \pm 1.0\text{ t C ha}^{-1}$, mixed). Rs showed a higher spatial variability at the mixed forest (21 - 87 % coefficient of variation (CV)) than at the broadleaved forest (23 - 46 % CV). Between 89 and 96 % of the annual temporal variation in measured Rs was explained by field soil temperature (Eq. (1), Fig. 2a). Rs showed a weak relationship with soil moisture at the broadleaved forest site, whereas there was no significant correlation between Rs and soil moisture at the mixed forest site (Fig. 2b).

Laboratory incubations showed a strong positive, exponential, relationship between soil temperature and Rh (Fig. 2c). Temperature sensitivity of mineral soil Rh was similar between sites (mixed $Q_{10} = 2.2$, broadleaved $Q_{10} = 2.3$; Fig. 2c, Table 2) and slightly lower for forest floor litter (mixed $Q_{10} = 1.9$; broadleaved $Q_{10} = 2.2$; Table 2). Q_{10} values of dry soil (mixed $Q_{10} = 1.6$; broadleaved $Q_{10} = 1.5$) were significantly lower than Q_{10} from the soil which remained at intermediate moisture content ($P < 0.05$, Table 2). Q_{10} values obtained from dry and wet soil did not differ significantly (Table 2). Rh and soil moisture showed a unimodal relationship with highest rates of Rh at intermediate soil moisture (40 - 50 vol. %) and decreasing rates at lower and higher moisture levels (Fig. 2d). Soil from both sites responded overall similarly to changes in soil moisture. Mixed forest soil showed a slightly sharper decrease in Rh at lower and at higher soil moisture (Fig. 2d).

Trenching plots indicated an average autotrophic and heterotrophic contribution of 29 and 27 % and 71 and 73 % at the mixed and broadleaved forest sites during the whole 2015 season, respectively (Fig. 3). The contribution of R_a and R_h to R_s , obtained by trenching, showed high temporal variability and strong fluctuations between individual measurement dates at the mixed forest site (Fig. 3).

The modelling approach yielded annual heterotrophic contributions of 67 % in mixed forest and 63 % in broadleaved forest. Modelled cumulative annual R_h and R_a were 8.6 and 4.2 t C ha⁻¹ at the mixed and 9.5 and 5.0 t C ha⁻¹ at the broadleaved forest respectively. Modelled R_h was in the range of field R_s during the cold season (Fig. 3). The gap between R_h and R_s became larger during the growing season, implying highest contribution of R_a during the warm monsoon months at both sites (Fig. 3 and 4). The strong temporal fluctuation in sources (R_a , R_h) which was obtained from trenching was not confirmed by R_h model output (Fig. 3).

4. Discussion

Annual R_s of both forest sites ($12.8 - 14.5 \text{ t C ha}^{-1}$) was in the range of values reported for similar ecosystems ($10.1-13 \text{ t C ha}^{-1}$ (Dar et al., 2015); $10-12 \text{ t C ha}^{-1}$ (Li et al., 2008); 13.7 t C ha^{-1} (Yang et al., 2007) and 14.7 t C ha^{-1} (Wang et al., 2010)). The higher altitude mixed forest had double tree basal area and standing stock, indicating that this specific site is exceptionally productive (Singh et al., 1994; Sharma et al., 2010a; Tashi et al., 2016; Wangda and Ohsawa, 2006b). Soil C stocks of $\sim 140 \text{ t ha}^{-1}$ (0-30 cm depth mineral soil) indicate that these mixed forests are likely among those ecosystems with the highest C storage capacity in the eastern Himalayas (Wangda and Ohsawa, 2006a; Sheikh et al., 2009; Dorji et al., 2014a; Tashi et al., 2016). High soil C contents and stocks were reflected in generally higher basal respiration (R_{s10}) at the mixed forest explaining the comparatively high annual R_s rates at this cooler, higher altitude, site. Soil C input via aboveground litter-fall was almost identical between sites ($\sim 3.5 \text{ t C ha}^{-1}$) although tree basal area was substantially lower at the broadleaved forest. This can be attributed to a generally higher leaf litter production in broadleaved ecosystems (Tiwari and Joshi, 2015; Bisht et al., 2014). Fine root stocks at both sites fall within the upper range of estimates from other surveys in the Himalayan region (Adhikari et al., 1995; Usman et al., 1999; Garkoti, 2008; Rana et al., 2015), especially if it is considered that fine root contents in this study were estimated solely for 0-30 cm mineral soil depth.

At both forests, R_s tightly followed the seasonal course of soil temperature because soil temperature and soil moisture co-varied with dry and cold winters and optimal soil moisture during the warm summer months (Fig. 1b, c; Fig 2a, b). R_s can also be affected by labile C allocation to soil (Gu et al., 2004). During the growing season, trees tend to allocate higher amounts of labile C belowground, thereby potentially increasing the contribution of R_a and simultaneously accelerating SOM decomposition by increased availability of labile C and rhizosphere priming (Kuznyakov, 2010; Bader and Cheng, 2007; Bengtson et al., 2012; Dijkstra and Cheng, 2007; Schindlbacher et al., 2009). Such processes would further increase R_s and R_a during the warm summer months. Our modelled R_h and R_a data suggests that this was also likely the case in the studied forests (significant increase in R_a contribution during the summer months; Fig. 3).

Our model generated wintertime R_h fluxes which were in the range of, or slightly below, R_s fluxes (Fig. 4). During frost periods, downward C-flux from the tree canopy is limited and the contribution of R_a to R_s is considered low during winter (Rey et al., 2002; Hanson et al., 2000). Our modelled wintertime (and overall) R_h therefore lay in a realistic range. However, there is evidence that the contribution of R_a can be significant even during cold winters (Schindlbacher et al., 2007; Tucker et al., 2014). Roots in deeper and warmer soil layers can remain active and add to the soil CO_2 efflux. Accordingly, modelled R_h rather represents the upper edge of potential R_h at our site. Our modelling approach was based on a relatively simple set of soil C stocks combined with temperature and moisture sensitivities, and holds corresponding uncertainty with regard to quantity of R_h and its temporal dynamics. C stocks from deeper soil layers ($> 30 \text{ cm}$ depth) were not accounted for and a single Q_{10} (obtained from 0-10 cm depth) was used for the whole mineral soil layer. Stabilization of SOC is usually increasing with soil depth (Fontaine et al., 2007). Our R_h predictions for deeper layers (10-30cm) might therefore overestimate the real rate.

Using annual litter input as proxy for litter C stocks is a further source of uncertainty. Litter input has temporal patterns and thereby affects litter decomposition dynamics. Such temporal patterns in litter input/decomposition were not reflected in our model. The modeled contribution of the litter layer to total soil Rh was, however, small (Fig. 4), and therefore, the uncertainty related to temporal litter layer dynamics can also be considered as small. We further used a constant Q_{10} throughout the year, although the Q_{10} may vary with season due to changes in substrate supply and quality (Davidson and Janssens, 2006; Gu et al., 2004) and/or interactions with soil moisture (Sierra et al., 2015). We showed that soil moisture affected the temperature sensitivity of Rh by significantly lower Q_{10} under dry conditions (lab incubation, Table 2). Such dry conditions were, however, not observed in the field. We therefore assume that ignoring potential moisture effects on Q_{10} in our model had only minimal effects on the Rh estimate. Rhizosphere priming could have affected Rh dynamics as well, but we were not able to account for that in our model. Moreover, soil sieving could have positively affected Rh rates during incubation by releasing physically protected SOM and/or providing additional C sources via disrupted fungal hyphae and fine root fragments (Datta et al., 2014). Nevertheless, the modelled annual ~ 65 % contribution of Rh falls well within estimates from similar forests (Lee et al., 2010). Even if we overestimated the real contribution of Rh, we are confident that the model relatively robustly reflected the temporal dynamics of Rh/Ra throughout the year.

In contrast to the modelling approach, trenching was applied as an attempt to estimate Ra *in situ*. The trenching method, although highly invasive, was shown to provide reasonable estimates of Ra for several forest types (Hanson et al., 2000; Subke et al., 2006). Trenching suggested slightly higher contributions of Rh at both sites (average 72 % both sites) but showed much stronger temporal variations in Rh/Ra, especially at the mixed forest (Fig. 3). Trenching has several drawbacks. Soil moisture is usually higher in trenched plots because water uptake by roots is interrupted. This bias was accounted for as we used the moisture response function (Eq. (3)) for correction. However, trenched fine roots can maintain respiration for a comparatively long time after cutting (Lee et al., 2003) and when fine roots finally die, their decomposition can add to the soil CO₂ efflux from the trenched plots (Hanson et al., 2000). Assuming a dead fine root mass loss of roughly one-third during the second year after trenching (Díaz-Pinés et al., 2010) and accounting for the corresponding effects on soil CO₂ efflux (additional efflux ~ 1 t C ha⁻¹), the estimated annual contribution of Rh decreases to ~ 65 % of Rs, which is in the range of our modelling results. Potential effects of root decomposition, however, do not explain the atypically strong temporal variation in Ra at the mixed forest site. Soil CO₂ efflux from trenched plots was similar or even higher than from corresponding control plots, suggesting a steep decrease in Ra between July and September (Fig. 3). We do not have a straightforward explanation for this pattern. Probably we did not trench deep enough and missed a larger proportion of roots which added to the summertime CO₂ efflux from trenched plots. A further explanation could be altered nutrient availability to decomposers in the trenched plots. In trenched plot soil, roots do not compete any more for nutrients, potentially increasing nutrient availability to decomposers. This could accelerate SOM decomposition and soil CO₂ efflux. In summary, trenching showed a less clear outcome at the two study sites when compared to other forests. Therefore, other methods, such as girdling or isotope labeling might be alternatively applied in the forest types studied.

Our simple empirical temperature-driven Rs model explained most of the temporal variation in Rs under the typical monsoon weather patterns during 2015. However, monsoon failures and drought periods have occurred in the past and may even increase in frequency and/or severity of climate change (Schewe and Levermann, 2012; Menon et al., 2013; Cook et al., 2010; Sharmila et al., 2015). To model drought effects, it is necessary to further develop the model by integrating potential soil moisture response of Rs. To do so, longer Rs time series including dry years and/or data from artificial drought experiments are needed for model parameterization and testing.

5. Conclusion

The monsoon climate allows for highly productive mountain forests in the eastern Himalayas. Such forests can store high amounts of C in plant biomass and soil, which was particularly evident in the high altitude mixed forest in our study. At both forests studied, a simple temperature-driven model was sufficient to explain most of the temporal variation in Rs during the study year. The sites experienced typical monsoon climate with dry and cold winters and monsoon rain during the warm season. Further research and model development is, however, warranted to better understand how infrequent/extreme events such as monsoon failure and drought affect soil/ecosystem C cycling and Rs in these forest ecosystems.

6. Author contribution

N. Wangdi carried out the field research, analyzed data and drafted the manuscript. M. Mayer performed modelling and contributed to writing the manuscript. M. P. Nirola carried out the incubation experiment and analysed the data. N. Zangmo and K. Orong collected the data and continuously monitored the research sites. I.U Ahmed carried out the root and the soil analyses. G. Gratzner designed the larger-scale throughfall manipulation experiment. R. Jandl, G. Gratzner and A. Darabant designed this study and provided feedback on the manuscript. A. Schindlbacher supervised the overall work, designed the experiment and critically revised the manuscript.

7. Data Availability

All relevant soil respiration, soil moisture, and soil temperature data from the field and the laboratory incubations are freely available from open source figshare repository (<https://figshare.com>) via <https://dx.doi.org/10.6084/m9.figshare.4239122>

8. Acknowledgements

We are highly grateful to the management and staff of the Ugyen Wangchuck Institute for Conservation and Environment, Bumthang for supporting the study. This study was part of the work package I of the BC-CAP project (Climate Change Adaptation potentials of forests in Bhutan –Building human capacities and knowledge base) jointly implemented by the Department of Forest and Park Services, Bhutan and University of Natural Resources and Life Sciences (BOKU), Austria with funding by the Austrian Ministry of Agriculture, Forestry, Environment and Water Management.

9. Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the views of any institutions of the Royal Government of Bhutan or the Government of Austria.

References

- Adhikari, B., Rawat, Y., and Singh, S.: Structure and function of high altitude forests of central Himalaya I. Dry matter dynamics, *Ann. Bot.*, 75, 237-248, 1995.
- Bader, N. E., and Cheng, W.: Rhizosphere priming effect of *Populus fremontii* obscures the temperature sensitivity of soil organic carbon respiration, *Soil Biol. Biochem.*, 39, 600-606, 2007.
- Bengtson, P., Barker, J., and Grayston, S. J.: Evidence of a strong coupling between root exudation, C and N availability, and stimulated SOM decomposition caused by rhizosphere priming effects, *Ecology and Evolution*, 2, 1843-1852, 10.1002/ece3.311, 2012.
- Bisht, V. K., Nautiyal, B. P., Kuniyal, C. P., Prasad, P., and Sundriyal, R. C.: Litter Production, Decomposition, and Nutrient Release in Subalpine Forest Communities of the Northwest Himalaya, *J. Ecosyst.*, 2014, 2014.
- Bolstad, P., Davis, K., Martin, J., Cook, B., and Wang, W.: Component and whole-system respiration fluxes in northern deciduous forests, *Tree Physiol.*, 24, 493-504, 2004.
- Bond-Lamberty, B., and Thomson, A.: A global database of soil respiration data, *Biogeosciences*, 7, 1915-1926, 10.5194/bg-7-1915-2010, 2010.
- Buchmann, N.: Biotic and abiotic factors controlling soil respiration rates in *Picea abies* stands, *Soil Biology and Biochemistry*, 32, 1625-1635, 2000.
- Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D'Arrigo, R. D., Jacoby, G. C., and Wright, W. E.: Asian monsoon failure and megadrought during the last millennium, *Science*, 328, 486-489, 2010.
- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., and Totterdell, I. J.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 184-187, 2000.
- Dar, J. A., Ganie, K. A., and Sundarapandian, S.: Soil CO₂ efflux among four coniferous forest types of Kashmir Himalaya, India, *Environmental monitoring and assessment*, 187, 715, 10.1007/s10661-015-4927-2, 2015.
- Davidson, E. A., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165-173, 2006.
- de Wit, H. A., Palosuo, T., Hylen, G., and Liski, J.: A carbon budget of forest biomass and soils in southeast Norway calculated using a widely applicable method, *For. Ecol. Manage.*, 225, 15-26, 2006.
- Datta, R., Vranová, V., Pavelka, M., Rejšek, K., Formánek, P.: Effect of soil sieving on respiration induced by low-molecular-weight substrates, *International Agrophysics*, 28, 119-124, 2014.
- Díaz-Pinés, E., Schindlbacher, A., Pfeffer, M., Jandl, R., Zechmeister-Boltenstern, S., and Rubio, A.: Root trenching - A useful tool to estimate autotrophic soil respiration? Case study in an Austrian mountain forest, *European Journal of Forest Research*, 129, 101-109, 2010.
- Dijkstra, F. A., and Cheng, W.: Interactions between soil and tree roots accelerate long-term soil carbon decomposition, *Ecology Letters*, 10, 1046-1053, 10.1111/j.1461-0248.2007.01095.x, 2007.
- Dixon, R. K., Solomon, A., Brown, S., Houghton, R., Trexler, M., and Wisniewski, J.: Carbon pools and flux of global forest ecosystems, *Science*, 263, 185-190, 1994.

391 Dorji, T., Odeh, I., and Field, D.: Vertical Distribution of Soil Organic Carbon Density in Relation to Land
392 Use/Cover, Altitude and Slope Aspect in the Eastern Himalayas, *Land*, 3, 1232-1250, 10.3390/land3041232,
393 2014a.

394 Dorji, T., Odeh, I. O., Field, D. J., and Baillie, I. C.: Digital soil mapping of soil organic carbon stocks under
395 different land use and land cover types in montane ecosystems, Eastern Himalayas, *For. Ecol. Manage.*, 318, 91-
396 102, 2014b.

397 FAO: Global Forest Resources Assessment :Country Report -Bhutan, Food and Agriculture Organization of the
398 United Nations, Rome, 56, 2010.

399 Fontaine, S., Barot, S., Barre, P., Bdioui, N., Mary, B., and Rumpel, C.: Stability of organic carbon in deep soil
400 layers controlled by fresh carbon supply, *Nature*, 450, 277-280,
401 http://www.nature.com/nature/journal/v450/n7167/supinfo/nature06275_S1.html, 2007.

402 FRDD: Local Volume Table, Forest Resources Development Division, Department of Forests and Park Services,
403 Bhutan, 2005.

404 Frey, S. D., Lee, J., Melillo, J. M., and Six, J.: The temperature response of soil microbial efficiency and its
405 feedback to climate, *Nature Climate Change*, 3, 395-398, 2013.

406 FSI: Volume Equations for Forests of India, Nepal and Bhutan, Ministry of Environment and Forests, Govt. of
407 India, 249 pp., 1996.

408 Garkoti, S.: Estimates of biomass and primary productivity in a high-altitude maple forest of the west central
409 Himalayas, *Ecol. Res.*, 23, 41-49, 2008.

410 Gough, C. M., Vogel, C. S., Harrold, K. H., George, K., and Curtis, P. S.: The legacy of harvest and fire on
411 ecosystem carbon storage in a north temperate forest, *Global Change Biology*, 13, 1935-1949, 2007.

412 Grierson, A. J. C., and Long, D. G.: *Flora of Bhutan*, Royal Botanical Garden Edinburgh, Edinburgh, U.K., 1983.

413 Gu, L., Post, W. M., and King, A. W.: Fast labile carbon turnover obscures sensitivity of heterotrophic respiration
414 from soil to temperature: a model analysis, *Global Biogeochem. Cycles*, 18, 2004.

415 Hanson, P., Edwards, N., Garten, C., and Andrews, J.: Separating root and soil microbial contributions to soil
416 respiration: a review of methods and observations, *Biogeochemistry*, 48, 115-146, 2000.

417 Hashimoto, S., Carvalhais, N., Ito, A., Migliavacca, M., Nishina, K., and Reichstein, M.: Global spatiotemporal
418 distribution of soil respiration modeled using a global database, *Biogeosciences*, 12, 4121-4132, 10.5194/bg-12-
419 4121-2015, 2015.

420 Janssens, I. A., Freibauer, A., Ciais, P., Smith, P., Nabuurs, G.-J., Folberth, G., Schlamadinger, B., Hutjes, R. W.,
421 Ceulemans, R., and Schulze, E.-D.: Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic
422 CO₂ emissions, *Science*, 300, 1538-1542, 2003.

423 Kutsch, W. L., Persson, T., Schrumpf, M., Moyano, F. E., Mund, M., Andersson, S., and Schulze, E.-D.:
424 Heterotrophic soil respiration and soil carbon dynamics in the deciduous Hainich forest obtained by three
425 approaches, *Biogeochemistry*, 100, 167-183, 2010.

426 Kuzyakov, Y.: Priming effects: Interactions between living and dead organic matter, *Soil Biology and*
427 *Biochemistry*, 42, 1363-1371, <http://dx.doi.org/10.1016/j.soilbio.2010.04.003>, 2010.

428 Laumans, P.: Height -Diameter Functions form a Country Level Site classification and Local Volume Table
429 Selection., UNDP/FAO Forest Resources Management and Institutional

430 Development Project, Thimphu, Bhutan, 1994.

431

432 Lee, M.-s., Nakane, K., Nakatsubo, T., and Koizumi, H.: Seasonal changes in the contribution of root respiration to
433 total soil respiration in a cool-temperate deciduous forest, in: *Roots: The Dynamic Interface between Plants and the*
434 *Earth: The 6th Symposium of the International Society of Root Research*, 11–15 November 2001, Nagoya, Japan,
435 edited by: Abe, J., Springer Netherlands, Dordrecht, 311–318, 2003.

436 Lee, N.-y., Koo, J.-W., Noh, N. J., Kim, J., and Son, Y.: Autotrophic and heterotrophic respiration in needle fir and
437 Quercus-dominated stands in a cool-temperate forest, central Korea, *J. Plant Res.*, 123, 485–495, 10.1007/s10265-
438 010-0316-7, 2010.

439 Li, H.-j., Yan, J.-x., Yue, X.-f., and Wang, M.-b.: Significance of soil temperature and moisture for soil respiration
440 in a Chinese mountain area, *Agric. For. Meteorol.*, 148, 490–503, 2008.

441 Liang, E., Wang, Y., Piao, S., Lu, X., Camarero, J. J., Zhu, H., Zhu, L., Ellison, A. M., Ciais, P., and Peñuelas, J.:
442 Species interactions slow warming-induced upward shifts of treelines on the Tibetan Plateau, *Proceedings of the*
443 *National Academy of Sciences*, 113, 4380–4385, 10.1073/pnas.1520582113, 2016.

444 Makkonen, K., and Helmisaari, H.-S.: Assessing fine-root biomass and production in a Scots pine stand—
445 comparison of soil core and root ingrowth core methods, *Plant Soil*, 210, 43–50, 1999.

446 Menon, A., Levermann, A., Schewe, J., Lehmann, J., and Frieler, K.: Consistent increase in Indian monsoon
447 rainfall and its variability across CMIP-5 models, *Earth Syst. Dynam. Discuss.*, 4, 1–24, 10.5194/esdd-4-1-2013,
448 2013.

449 Pandey, R., Sharma, G., Singh, T., and Tripathi, S.: Factors influencing soil CO₂ efflux in a north eastern Indian
450 oak forest and plantation, *Afr. J. Plant Sci.*, 4, 280–289, 2010.

451 Pinheiro, J. C., and Bates, D. M.: *Mixed-Effects Models in S and S-Plus*, Springer - Verlag, New York Berlin
452 Heidelberg, 530 pp., 2000.

453 Pumpanen, J., Longdoz, B., and Kutsch, W.: Field measurements of soil respiration: principles and constraints,
454 potentials and limitations of different methods, in: *Soil Carbon Dynamics - An Integrated Methodology*, edited by:
455 Kutsch, W., Bahn, M., and Heinemeyer, A., Cambridge University Press, Cambridge, 16–33, 2009.

456 Rana, S., Bargali, K., and Bargali, S.: Assessment of plant diversity, regeneration status, biomass and carbon stock
457 in a Central Himalayan cypress forest, *Int. J. Biodivers. Conserv.*, 7, 321–329, 2015.

458 Reichstein, M., Rey, A., Freibauer, A., Tenhunen, J., Valentini, R., Banza, J., Casals, P., Cheng, Y., Grünzweig, J.
459 M., and Irvine, J.: Modeling temporal and large-scale spatial variability of soil respiration from soil water
460 availability, temperature and vegetation productivity indices, *Global Biogeochem. Cycles*, 17, 2003.

461 Rey, A., Pegoraro, E., Tedeschi, V., De Parri, I., Jarvis, P. G., and Valentini, R.: Annual variation in soil respiration
462 and its components in a coppice oak forest in Central Italy, *Global Change Biol.*, 8, 851–866, 2002.

463 Schewe, J., and Levermann, A.: A statistically predictive model for future monsoon failure in India, *Environ. Res.*
464 *Lett.*, 7, 044023, 2012.

465 Schickhoff, U.: The upper timberline in the Himalayas, Hindu Kush and Karakorum: a review of geographical and
466 ecological aspects, in: *Mountain Ecosystems*, Springer, 275–354, 2005.

467 Schild, A.: ICIMOD's Position on Climate Change and Mountain Systems, *Mountain Research and Development*,
468 28, 328–331, 10.1659/mrd.mp009, 2008.

469 Schindlbacher, A., Zechmeister-Boltenstern, S., Glatzel, G., and Jandl, R.: Winter soil respiration from an Austrian
470 mountain forest, *Agricultural and Forest Meteorology* 146, 205-215, 2007.

471

472 Schindlbacher, A., Zechmeister-Boltenstern, S., and Jandl, R.: Carbon losses due to soil warming: Do autotrophic
473 and heterotrophic soil respiration respond equally?, *Global Change Biology*, 15, 901-913, doi: 10.1111/j.1365-
474 2486.2008.01757.x, 2009.

475 Schlesinger, W. H., and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7-20,
476 2000.

477 Sharma, C. M., Baduni, N. P., Gairola, S., Ghildiyal, S. K., and Suyal, S.: Tree diversity and carbon stocks of some
478 major forest types of Garhwal Himalaya, India, *For. Ecol. Manage.*, 260, 2170-2179, 2010a.

479 Sharma, E., Chettri, N., and Oli, K. P.: Mountain biodiversity conservation and management: a paradigm shift in
480 policies and practices in the Hindu Kush-Himalayas, *Ecol. Res.*, 25, 909-923, 10.1007/s11284-010-0747-6, 2010b.

481 Sharma, P., and Rai, S. C.: Carbon sequestration with land-use cover change in a Himalayan watershed, *Geoderma*,
482 139, 371-378, 10.1016/j.geoderma.2007.02.016, 2007.

483 Sharmila, S., Joseph, S., Sahai, A. K., Abhilash, S., and Chattopadhyay, R.: Future projection of Indian summer
484 monsoon variability under climate change scenario: An assessment from CMIP5 climate models, *Global Planet.*
485 *Change*, 124, 62-78, 10.1016/j.gloplacha.2014.11.004, 2015.

486 Sheikh, M. A., Kumar, M., and Bussmann, R. W.: Altitudinal variation in soil organic carbon stock in coniferous
487 subtropical and broadleaf temperate forests in Garhwal Himalaya, *Carbon balance and management*, 4, 1-6, 2009.

488 Shrestha, U. B., Gautam, S., and Bawa, K. S.: Widespread climate change in the Himalayas and associated changes
489 in local ecosystems, *PLoS One*, 7, e36741, 10.1371/journal.pone.0036741, 2012.

490 Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of
491 soil organic matter with respect to simultaneous changes in temperature and moisture, *Journal of Advances in*
492 *Modeling Earth Systems*, 7, 335-356, 2015.

493 Singh, S. B. K., Isabella; Singh Karky, Bhaskar; Sharma, Eklabya: Climate change in the Hindu Kush-Himalayas:
494 the state of current knowledge, *ICIMOD*, 88 2011.

495 Singh, S. P., Adhikari, B. S., and Zobel, D. B.: Biomass, productivity, leaf longevity, and forest structure in the
496 central Himalaya, *Ecol. Monogr.*, 401-421, 1994.

497 Stocker, T. F.: Climate change 2013: the physical science basis: Working Group I contribution to the Fifth
498 assessment report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2014.

499 Subke, J.-A., Inglima, I., and Cotrufo, F. M.: Trends and methodological impacts in soil CO₂ efflux partitioning: A
500 metaanalytical review, *Global Change Biology*, 12, 921-943, doi: 10.1111/j.1365-2486.2006.01117.x, 2006.

501 Sundarapandian, S., and Dar, J.: Variation in Soil CO₂ Efflux in *Pinus Wallichiana* and *Abies Pindrow* Temperate
502 Forests of Western Himalayas, India. *Forest Res*, 3, 2, 2013.

503 Tashi, S., Singh, B., Keitel, C., and Adams, M.: Soil carbon and nitrogen stocks in forests along an altitudinal
504 gradient in the eastern Himalayas and a meta-analysis of global data, *Global Change Biol.*, 2016.

505 Tiwari, S., and Joshi, R.: Litter-fall production in cool-temperate forests of the Nanda Devi Biosphere Reserve,
506 Uttarakhand Himalaya, India, *Braz. J. Biolo. Sci.*, 2, 147-154, 2015.

507 Tsering, K., Sharma, E., Chettri, N., and Shrestha, A.: Climate Change Impact and Vulnerability in the Eastern
508 Himalayas – Synthesis Report. ICIMOD, ICIMOD, Kathmandu, Nepal, 2010.

509 Tucker, C.L., Young, J.M., Williams, D.G., and Ogle, K.: Process-based isotope partitioning of winter soil
510 respiration in a subalpine ecosystem reveals importance of rhizospheric respiration, *Biogeochemistry*, 121, 389-
511 408, 2014.

512

513 Usman, S., Singh, S., and Rawat, Y.: Fine root productivity and turnover in two evergreen central Himalayan
514 forests, *Ann. Bot.*, 84, 87-94, 1999.

515 Verma, A., Tewari, A., and Shah, S.: Carbon storage capacity of high altitude *Quercus semecarpifolia*, forests of
516 Central Himalayan region, *Scand. J. For. Res.*, 27, 609-618, 10.1080/02827581.2012.689003, 2012.

517 Wang, X., Jiang, Y., Jia, B., Wang, F., and Zhou, G.: Comparison of soil respiration among three temperate forests
518 in Changbai Mountains, China, *Can. J. For. Res.*, 40, 788-795, 2010.

519 Wang, X., Liu, L., Piao, S., Janssens, I. A., Tang, J., Liu, W., Chi, Y., Wang, J., and Xu, S.: Soil respiration under
520 climate warming: differential response of heterotrophic and autotrophic respiration, *Global Change Biol.*, 20, 3229-
521 3237, 2014.

522 Wangda, P., and Ohsawa, M.: Gradational forest change along the climatically dry valley slopes of Bhutan in the
523 midst of humid eastern Himalaya, *Plant Ecol.*, 186, 109-128, 2006a.

524 Wangda, P., and Ohsawa, M.: Structure and regeneration dynamics of dominant tree species along altitudinal
525 gradient in a dry valley slopes of the Bhutan Himalaya, *For. Ecol. Manage.*, 230, 136-150, 2006b.

526 Wikramanayake, E. D.: Terrestrial ecoregions of the Indo-Pacific: a conservation assessment, Island Press, 2002.

527 Xu, J., Grumbine, R. E., Shrestha, A., Eriksson, M., Yang, X., Wang, Y., and Wilkes, A.: The melting Himalayas:
528 cascading effects of climate change on water, biodiversity, and livelihoods, *Conserv. Biol.*, 23, 10.1111/j.1523-
529 1739.2009.01237.x, 2009.

530 Xu, J., and Grumbine, R. E.: Building ecosystem resilience for climate change adaptation in the Asian highlands,
531 *Wiley Interdisciplinary Reviews: Climate Change*, 5, 709-718, 10.1002/wcc.302, 2014.

532 Yang, Y.-S., Chen, G.-S., Guo, J.-F., Xie, J.-S., and Wang, X.-G.: Soil respiration and carbon balance in a
533 subtropical native forest and two managed plantations, *Plant Ecol.*, 193, 71-84, 2007.

Table 1 Site characteristics of the two studied forests

Parameter	Mixed forest	Broadleaved forest
Elevation (m)	3260	2460
Latitude	27°28' 00'' N	28°28'51.06'' N
Longitude	89°44'30.79''E	89°51'27.73'' E
Annual Precipitation 2015 (mm)	1167	1120
Mean Air Temperature 2015 (°C)	7.8	12.0
Dominant Overstorey species	<i>Tsuga dumosa</i> (59.5%) <i>Quercus semecarpifolia</i> (29.3%) <i>Picea spinulosa</i> (6.3%) <i>Abies densa</i> (4.1%) <i>Taxus baccata</i> (0.3%)	<i>Quercus lanata</i> (63.5%) <i>Quercus griffithii</i> (29.6%)
Dominant Understory species	<i>Ilex dipreyana</i> (0.2%) <i>Rhododendron arboreum</i> (0.1%)	<i>Symplocos sp.</i> (0.8%) <i>Lyonia ovalifolia</i> , (2.2%) <i>Rhododendron arboreum</i> (3.4%)
Tree density (No. ha ⁻¹)	364 ± 50	569 ± 19
Mean Tree height (m) Overstorey	24.4 ± 2.1	23.6 ± 1.4
Mean Tree Height (m) Understorey	7.8 ± 3.5	9.8 ± 0.4
Mean DBH (cm) Overstorey	50.7 ± 5.8	37.8 ± 2.3
Mean DBH (cm) Understorey	13.8 ± 1.4	16.1 ± 0.9
Tree basal area (m ² ha ⁻¹)	77.5 ± 4.6	39.9 ± 4.4
Standing volume (m ³ ha ⁻¹)	1066 ± 2.3	464 ± 25
Soil organic C (t ha ⁻¹) 0-30 cm	142.0 ± 25.4	90.1 ± 9.0
Soil organic C (t ha ⁻¹) 0-10 cm	61.9 ± 5.3	55.5 ± 6.9
Soil organic C (t ha ⁻¹) 10-30 cm	80.1 ± 8.0	34.6 ± 2.4
Soil N (t ha ⁻¹) 0-30 cm	7.4 ± 0.5	4.3 ± 0.4
Soil N (t ha ⁻¹) 0-10 cm	3.2 ± 0.2	2.4 ± 0.3
Soil N (t ha ⁻¹) 10-30 cm	4.2 ± 0.4	1.9 ± 0.1
pH (0-10 cm)	5.2 ± 0.1	5.0 ± 0.1
Bulk density (g cm ⁻³) 0-10 cm	0.61 ± 0.02	0.61 ± 0.01
Fine Root biomass (t C ha ⁻¹) 0-30 cm	2.3 ± 0.3	3.2 ± 0.5
Litter input (t C ha ⁻¹ yr ⁻¹)	3.5 ± 0.10	3.4 ± 0.03

*All stand and soil parameters are expressed as the mean ± standard error.

Layer	Incubation	Moisture (vol.%)	Rh ₁₀ ($\mu\text{mol CO}_2 \text{ kg C}^{-1} \text{ s}^{-1}$)	Q ₁₀
<u>Broadleaved</u>				
Litter	1	46 \pm 1	0.58 \pm 0.04	2.22 \pm 0.09
Mineral	1	35 \pm 2	0.22 \pm 0.03	2.31 \pm 0.06
Mineral	2	dry (10 \pm 1)	0.10 \pm 0.01	1.54 \pm 0.11a
Mineral	2	interm. (33 \pm 1)	0.14 \pm 0.02	2.39 \pm 0.22b
Mineral	2	wet (56 \pm 1)	0.18 \pm 0.03	2.12 \pm 0.25ab
<u>Mixed</u>				
Litter	1	46 \pm 2	1.05 \pm 0.24	1.93 \pm 0.06
Mineral	1	33 \pm 1	0.16 \pm 0.02	2.25 \pm 0.06
Mineral	2	dry (9 \pm 1)	0.08 \pm 0.01	1.55 \pm 0.18a
Mineral	2	interm. (29 \pm 1)	0.10 \pm 0.01	2.63 \pm 0.39b
Mineral	2	wet (51 \pm 1)	0.13 \pm 0.01	2.06 \pm 0.10ab

Table 2. Basal respiration rates (Rh₁₀) and temperature sensitivity (Q₁₀) of litter and mineral soil (0-10 cm depth) samples derived from laboratory incubations. Incubations took place initially after sampling (Incubation 1) using a set of 3 samples per plot (6 plots per site). Subsequently, sets were split and the moisture sensitivity of Rh was tested (Fig. 2d). Subsequent to moisture incubations, the different subsets (Dry, Interm., Wet) were re-incubated to test temperature sensitivities at different moisture contents (Incubation 2). The time-lag between Incubation 1 and Incubation 2 was approximately 10 weeks. Different letters indicate significant differences in Q₁₀ between soil moisture levels of the mineral soil samples.




Figure 1. Seasonal course of air temperature and precipitation (a), soil temperature (b), volumetric soil water content (c), and soil respiration (d) measured at a mixed and a broadleaved forest in Bhutan Himalayas in 2015. Circles represent daily mean values of manual measurements. Solid lines (a, b, c) represent daily mean values of continuous measurements. Error bars indicate standard error of the mean.




Figure 2. (a) Relationship between soil CO₂ efflux (R_s) and soil temperature, and (b) R_s and soil moisture (vol.%) at a broadleaved and a mixed forest in Bhutan Himalayas. (c) Relationship between heterotrophic soil respiration (R_h) and soil temperature, and (d) R_h and soil moisture (vol. %) as determined during a laboratory incubation. A temperature response was fitted with an exponential function (Eq. (1)) and a moisture response was fitted with a polynomial function (Eq. (3)). Error bars represent standard error of the mean (SE). Basal respiration rates at 10 $^{\circ}\text{C}$ (Rs_{10} , Rh_{10}) and temperature sensitivity of R_h (Q_{10}) are given (mean \pm SE).




Figure 3: Contribution of autotrophic soil respiration (R_a) to total soil CO_2 efflux (R_s) at a (a) broadleaved and (b) mixed forest in Bhutan Himalayas. Autotrophic contribution was derived from the differences between R_s measured at control and trenched plots (bars) and from the differences between modelled R_s and heterotrophic soil respiration rates (lines), respectively.




Figure 4. Seasonal course of modelled soil CO₂ efflux (Rs) and heterotrophic soil respiration rates (Rh) at a broadleaved and mixed forest in Bhutan Himalayas in 2015. Open circles are measured Rs rates. Error bars and shaded areas represent standard error of the daily mean. Dashed and dotted lines indicate the CO₂ contributions of litter and mineral soil layers to Rh. The area between the full line (total Rh) and the dashed line represents the contribution from litter, the area between dashed and dotted line represents the contribution of the topsoil (0-10 cm), and the area below the dotted line represents the contribution from the 10-30 cm mineral soil layer.