

## Supplementary Information

Table S1. Selection parameters for rainfall event samples from the Black Burn used in irradiation experiments. Top row denotes winter rainfall event values and bottom row denotes summer rainfall event values.

	<b>Rainfall event</b>	<b>a<sub>254</sub></b>	<b>E4:E6</b>	<b>POC mg L<sup>-1</sup></b>
Median	Winter	1.63	6.70	11.0
	Summer	1.36	6.73	5.83
Maximum	Winter	1.98	8.63	63.5
	Summer	1.47	11.0	10.6
Minimum	Winter	1.34	3.57	4.10
	Summer	1.30	4.62	3.88

Table S2. Carbon budget calculations for individual C species and total C balance in samples collected in May 2014 at the Black Burn and Loch Katrine. CH<sub>4</sub> concentrations are not included in budget calculations as they were < 0.05% of the total budget. Differences are expressed as % of non-irradiated samples. Values are means of 4 replicates. LOD = limit of detection

	DOC mg L <sup>-1</sup>	DIC mg L <sup>-1</sup>	CO <sub>2</sub> mg L <sup>-1</sup>	CO mg L <sup>-1</sup>	∑ C mg L <sup>-1</sup>
<b>Black Burn</b>					
Irradiated	48.9	1.18	2.82	0.13	<b>53.0</b>
Non irradiated	50.9	1.38	1.20	<LOD	<b>53.5</b>
Difference	-3.9 %	-14 %	135 %	---	<b>-0.8 %</b>
<b>Loch Katrine</b>					
Irradiated	5.30	1.42	0.82	0.02	<b>7.56</b>
Non irradiated	5.05	1.55	0.80	<LOD	<b>7.40</b>
Difference	5.0 %	-8.4 %	2.5 %	---	<b>2.2 %</b>

Figure S1. Spectral output (240-800 nm) of UV-B 313 lamps employed in irradiation experiments in this study.




Figure S2. Black Burn discharge during and 2 weeks before a) winter and b) summer rainfall events. The events are demarcated by two vertical lines on each of the plots.




Figure S3. Pearson correlation between total concentration of lignin phenols  $\Sigma_{11}$  ( $\mu\text{g L}^{-1}$ ) and DOC concentration in all Black Burn water samples (n=28), including monthly and rainfall event samples.

