

Rooting and plant ~~density coverage strongly~~ determine greenhouse gas budget of water hyacinth (*Eichhornia crassipes*) ~~mats~~

Ernandes Sobreira Oliveira Junior^{1,4*}; Yingying Tang^{1*}; Sanne J.P. van den Berg²; Leon P. M. Lamers^{1, 3};

5 Sarian Kosten¹

¹Department of Aquatic Ecology and Environmental Biology, Institute for Water and Wetland Research, Radboud University [Nijmegen](#), P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

²B-WARE Research Centre, P.O. Box 6558, 6503 GB, Nijmegen, The Netherlands.

³Department of Aquatic Ecology and Water Quality Management, Wageningen University and Research
10 Centre, P.O. Box 47, 6700 AA Wageningen, The Netherlands

⁴[Universidade Estadual de Mato Grosso, Laboratory of Ichthyology, Centro de Limnologia, Biodiversidade e Etnoecologia do Pantanal – previous institution.](#)

*Corresponding author:

15 Ernandes Sobreira Oliveira Junior

Dept. of Aquatic Ecology & Environmental Biology

Radboud University Nijmegen,

Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

Email: e.sobreira@science.ru.nl

*These authors equally contributed to the paper.

Running head: Rooting and plant [densitycoverage](#) strongly determine greenhouse gas budget of water hyacinth.

Keywords: Greenhouse gases, invasive species, experimental ecology, nutrient cycling, population

5 Abstract

Water hyacinth occurs in numerous tropical and subtropical countries, either as a native or as an invasive exotic species, where it can establish large and dense mats. The plant is also frequently used for water purification and bioremediation purposes. Although it is a free-floating species, the plant roots into the sediment of shallow waters, tapping into the sediment nutrient pool. Its long and extensive root system strongly increases nutrient
10 absorption, resulting in high growth rates and concurring high carbon sequestration rates. On the other hand, the plants may also fuel methane (CH₄) production as dense mats may deplete oxygen in the surface water and sediment below, which in combination with the high production of organic matter creates favorable conditions for methanogenesis. We hypothesize that water hyacinth vegetation acts as a strong greenhouse gas (GHG) sink due to its high growth rates, especially when (sediment) nutrient availability is high. Still, this sink may be
15 counterbalanced by CH₄ release, which will be most pronounced when the plants are rooting in the sediment due to potential CH₄ shuttling from the sediment through the roots and leaves into the atmosphere ([plant-mediated transport-chimney effect](#)). To mechanistically unravel the influence of water hyacinth on nutrient dynamics and greenhouse gas fluxes, we performed an aquarium experiment in which plant [densitycoverage](#) and root access to the sediment were manipulated. Although plant cover led to lower concentrations of dissolved total
20 phosphorus (DTP) and phosphate, there were no effects of [coveragedensity](#) or rooting. We found no vegetation

effect on the ebullition of CH₄, but its diffusion was 4.5 times higher at high plant coverage. Rooting increased CH₄ diffusion by 1.3 (high [coveredensity](#)) and 4 times (low [densitycoverage](#)), demonstrating the [plant-mediated transport chimney effect](#) that we hypothesized. Independent of rooting, however, water hyacinth at high [coveredensity](#) sequestered less carbon compared to low [coverage-density](#), possibly due to space limited growth and self-shading. Overall, water hyacinth enhanced CH₄ emissions, especially when rooted. Due to water hyacinth's high CO₂ sequestration rates, the overall GHG budget in terms of CO₂ equivalents still resulted in water hyacinth mats being near-neutral or even a GHG sink, depending on water hyacinth [densitycoverage](#). Our results show that the effect of water hyacinth mats on GHG fluxes strongly depends on both plant [densitycoverage](#) and contact with the sediment. This indicates that, when making regional GHG balances, not only plant presence but also its [densitycoverage](#) and water depth – regulating sediment-root contact – should be taken into account.

Key words: floating plant, nutrient dynamic, CH₄ emission, carbon dioxide sequestration, greenhouse warming potential.

1. Introduction

Water hyacinth (*Eichhornia crassipes*) is notorious [worldwide-at a global scale](#) because of the problems it poses to economy, society and ecology when occurring at high [densitiescoverages](#) (Villamagna and Murphy, 2010; Malik et al., 2007). Its high tolerance range for environmental conditions including pH, temperature and nutrients (Gutierrez et al. 2001; Wilson et al. 2005) provides an ample spectrum of colonization, and explains its widespread occurrence [and nuisance](#) around the world. Its fast growth rates and rapid dispersal through asexual reproduction explains its ability to form large floating mats comprising high biomass (Pinto-Coelho and Greco, 1999).

The search for useful end products for this large amount of biomass has resulted in its appliance as a source for bioenergy production (Chanakya et al. 1993), fodder (Jianqing et al. 2001), bio fertilizer (Zhao et al. 2012), and paper fiber (Reddy and Tucker 1983). Water hyacinth is also frequently used for water purification and bioremediation purposes because of its high nutrient uptake rates (Aoyama and Nishizaki, 1993; Mandi, 1994; Polprasert and Khatiwada, 1998). Harvested plants can be used for bioenergy production, fodder, bio fertilizer and paper fiber (Chanakya et al. 1993); (Jianqing et al. 2001); (Zhao et al. 2012); (Reddy and Tucker 1983))

Nutrient availability strongly determines *Eicchornia's* growth rate as well as nutrient allocation (Xie et al., 2004). Maximum nutrient uptake efficiency is typically reached in the early growth stage (Reddy, Agami & Tucker, 1989; Reddy, Agami & Tucker, 1990), explaining the formation of big mats in a few days (Tellez et al., 2008).

Water hyacinth's high growth rate results in high carbon dioxide (CO₂) uptake at rates of 3.4 - 5.4 g C-CO₂ m⁻² day⁻¹ as reported for tropical lakes (Peixoto et al., 2016). In these lakes, the vegetation even sets off open water CO₂ emissions, turning the system into CO₂ sink. While water hyacinth growth will decrease CO₂ emissions, its presence may simultaneously increase the emission of methane (CH₄) (Banik, Sen, and Sen, 1993), having a global warming potential (GWP) of 34 times CO₂ over a 100 year time scale (Myhre, Shindell and Bréon, 2013). Therefore, even relatively low rates of CH₄ emissions could offset the high CO₂ assimilation, turning water hyacinth mats into a greenhouse gas (GHG) source. The high density coverage of water hyacinth suppresses light penetration and therefore photosynthetic activity in the water below. In combination with reduced O₂ diffusion from the atmosphere into the water by its cover, this can result in anaerobic conditions below the plant water hyacinth mat (Reddy and DeBusk 1991) as has also been described for other floating plants (e.g. Caraco and Cole, 2002; Grasset et. al, 2016). Research performed in ditches and tanks showed that the combination of decreasing O₂ concentrations and high organic matter production by water hyacinth favors CH₄ emission. This effect was strongest after multiple years, probably due to organic matter accumulation (Banik et al., 1993).

Aquatic plants rooting in the sediment tend to enhance CH₄ emissions by transporting CH₄ directly from the sediment to the atmosphere. This plant-mediated CH₄ emission is an important pathway, which may even explain more than 50% of the total emission of inland waters (Dacey and Klug 1979; Grosse, Armstrong & Armstrong 1996). Plant-mediated CH₄ transport may take place through convective flow (pressurized flow), a common mechanism for aquatic plants, or by passive molecular diffusion (Cronk and Fennessy 2016; Grosse, Armstrong & Armstrong 1996; Konnerup et al. 2011)–chimney effect. Although water hyacinth is generally reported as a floating plant, the plant species can root in the sediment when the water level is sufficiently low (less than 50 cm; personal observation). Plant-mediated CH₄ emission is expected to increase when the plant is rooted in the sediment as transport rates are determined by concentration differences between compartments, and CH₄ concentrations tend to be much higher in sediment porewater than in surface water. Possibly, the increase in plant-mediated transport of pore water CH₄ to the atmosphere may inhibit the formation of bubbles in the sediment, thereby decreasing ebullition.

On the other hand, methanotrophic microbial communities associated with the plant roots (Yoshida et al. 2014) may oxidize a considerable portion of the CH₄ dissolved in the surface water or porewater (Kosten et al., 2016). The overall effect of floating plants on CH₄ emissions is therefore not straightforward (compare, for instance, findings of Bolpagni et al. (2007) and Ribaudo et al. (2012) who found that floating plants increase CH₄ emissions with Bharati et al. (2000) who found a reduction of CH₄ emissions; see also Kosten et al. (2016) for a review of field observations of the effect of floating plants on CH₄ emissions). Even when aquatic plants increase CH₄ emissions, their overall GHG budget may still be counterbalanced by their high growth rates - and therefore CO₂ uptake rates. CO₂ uptake rates can be expected to be highest when plants have access to nutrients in both sediment and water. enhanced growth rates and therefore CO₂ uptake rates, due to increased nutrient uptake from both sediment and water.

All in all, the effects of water hyacinth mats on GHG emissions are ~~therefore~~ not at all straightforward. Only few studies have investigated the effects of water hyacinth on total GHG (CH₄ and CO₂) emissions (~~CH₄ and CO₂~~) (Banik et al., 1993; Peixoto et al., 2016; Attermeyer et al., 2016), and none have included the effects of plant density/coverage or rooting. Moreover, the few studies that investigated the effect of water hyacinth on GHG balance showed contrasting results (Banik et al., 1993; enhanced CH₄ emissions; and Attermeyer, et al., 2016; decreased CH₄ emissions).

We therefore used an experimental approach under controlled conditions, to elucidate the roles of water hyacinth coverage and rooting on GHG dynamics. We hypothesize that ~~the~~ differences ~~found~~ may be 1) due to variation in density/coverage, and 2) related to whether or not the plants are rooted in the sediment. An increase in coverage may either increase or decrease CH₄ emissions, depending on the dominant process: enhanced methanogenesis due to lower oxygen (O₂) concentrations below the plant layers or enhanced CH₄ oxidation due to a higher root biomass and associated metanotrophic communities. With respect to hypothesis 2) we expect CO₂ uptake rates to be highest when plants have access to nutrients in the sediment, also leading to lower pore-water nutrient concentrations and C:N and C:P ratios. In addition, we expect diffusive CH₄ emissions – including plant-mediated emissions - to increase when plants are rooted. To elucidate the roles of coverage and rooting of water hyacinth vegetation on GHG fluxes and nutrient dynamics, and to study the directions of their effects, we used a full-factorial, controlled indoor aquarium experiment, ~~aiming to elucidate the effects of plant density/coverage and sediment rooting on the nutrient dynamics and GHG fluxes of water hyacinth vegetation.~~

2. Materials and methods

2.1. Experimental set-up

The experiment was conducted in 24 glass aquaria of 24 L (20 × 20 × 60 cm; length × width × height) filled with a layer of 7 cm of fresh sediment, and a layer of 38 cm of demineralized water. The sediment was collected from a eutrophic drainage ditch (Ede, The Netherlands; 51°59'43.58"N, 5°38'38.91"EL) in September 2014, and was sieved with a 5.0 mm sieve to remove stones and vegetation remnants. Sediment characteristics were determined at the beginning of the experiment (Table 1). The aquaria were placed in a water bath at 23°C in the greenhouse facilities of the Radboud University (Nijmegen, The Netherlands). A light frame of 220 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PAR (16h light/8h dark) was provided by Philips Green Power 400V/1000 WE lamps in a New E-Papillon 1000 W armature, to provide sufficient light in case of cloudy conditions.

Water hyacinth was collected from a commercial breeder (Nijmegen, the Netherlands) and cultivated in the greenhouse for approximately 10 months prior to the experiment, on organic sediment to which slow-release phosphorus granules were added. The experiment lasted for 59 days, from October to December 2014. The aquaria were randomly assigned to controls without plants, low densitycoverage (50% of water hyacinth coverage) or high coverage-density (100% coverage) ($n = 8$ for each treatment). In half of the treatments, a mesh (1.0 mm mesh size) was placed just above the sediment ($n=12$) to prevent the plants from rooting in the sediment, dividing the plant treatments into rooted and non-rooted treatments. There were 4 controls without, and 4 with a mesh (jointly referred to as 'controls'). We added individual water hyacinths to each aquarium: 1) 160 g (fresh weight - FW) to the low coveredensity treatment with mesh (non-rooted – 50%nR) or without mesh (rooted – 50%R); and 2) $413 \pm 2.63 \text{ g (FW} \pm \text{SD)}$ to the high densitycoverage treatment with mesh (non-rooted – 100%nR) or without mesh (rooted – 100%R) (Fig. 1). The fresh weight of the total biomass was measured using paper towel to carefully blot the plants dry and remove water attached, and then place the plants on a digital scale. To maintain ~~the~~ low coverage, water hyacinth was harvested partially at day 31 and 45.

2.2. Chemical analyses

Dissolved oxygen (DO), pH and temperature were measured weekly at both the surface and bottom of the water column, using a portable multi-meter (HQ40d multi, HACH, Loveland, Colorado, U.S.A.). Surface and porewater samples were collected anaerobically every week during the experiment using ceramic soil moisture
5 samplers (SMS rhizons, Eijkelkamp, Giesbeek, Netherlands). Total inorganic carbon (TIC) of water samples was measured with an Infra-red Gas Analyzer (IRGA; ABB Analytical, Frankfurt, Germany). Concentrations of PO_4^{3-} , NO_3^- and NH_4^+ in the water samples were measured colorimetrically on an Auto-Analyzer 3 system (Bran & Luebbe, Norderstedt, Germany) by using ammonium molybdate (Henriksen 1965), hydrazine sulphate (Kamphake et al. 1967) and salicylate (Grasshoff and Johannsen 1972), respectively. Concentrations of dissolved total P (DTP) were
10 measured by inductively coupled plasma-optical emission spectrometry (ICP-OES; IRIS Intrepid II, Thermo Fisher Scientific, Franklin, MA, U.S.A.). Dissolved organic carbon (DOC) in water samples was measured with a TOC-L CPH/CPN analyzer (Shimadzu, Kyoto, Japan) at the end of the experiment.

Sediment samples were collected at the start and end of the experiment, and subsequently dried for 48h at 60°C. Dry samples were heated for 4 hours at 550°C and re-weighed to determine organic matter content.
15 Dried sediment (200 mg) was digested in a microwave oven (MLS-1200 Mega, Milestone Inc., Sorisole, Italy) using 4 ml 65% HNO_3 and 1 ml 30% H_2O_2 to determine total sediment Fe, Al, Ca and P concentrations. Digested solutions were analyzed by ICP-OES (see above). Olsen P extracts (plant available P) was determined by extraction according to Olsen (1954), whereas a NaCl-extraction (exchangeable NH_4^+ and NO_3^-) was performed as described by Tomassen et al. (2004).

20

2.3. Greenhouse gas flux measurements

2.3.1. *Diffusive flux*

After 30, 38 and 45 days greenhouse gas (CO₂ and CH₄) diffusive fluxes were measured during the day and night using a lid on top of the aquarium to establish a closed system connected to a Picarro G2508 Greenhouse Gas Analyzer (Picarro Inc., Santa Clara, CA, USA). The lid was sealed air-tight with paste (Terostat IX, Teroson GmbH, Heidelberg, Germany). Measurements were conducted until a clear linear increase in CO₂ and CH₄ occurred (typically for 5 minutes). The slope of the relationship between gas concentration and time was used to calculate the gas flux as explained by Almeida et al. (2016). When the linear increase was interrupted by a sudden increase in gas concentration – due to bubbling – the lid was removed to restore ambient air concentrations, and the diffusive flux measurement was repeated.

2.3.2. *Ebullitive flux*

Total CH₄ fluxes (ebullitive + diffusive) were measured 3 times (on day 31, 39 and 46) during a period of 24 hours. During this time the glass lid (equipped with a rubber septum) was closed as described before. The increase in CH₄ concentration during 24 hours was determined by sampling the headspace (in duplicate) using a 1ml plastic syringe through the septum at the start and the end of the incubation and subsequent directly injected 0.5 ml into the gas chromatograph (HP 5890 equipped with a Porapak Q column (80/100 mesh), a flame ionization detector (GC-FID, Hewlett Packard, USA) and oven temperature 120°C). The total amount of CH₄ emitted was calculated by multiplying the change in CH₄ concentration in the headspace between t=0 and at the end with the volume of the headspace. The ebullitive fluxes were calculated by subtracting diffusive CH₄ fluxes determined the day before from the total amount of CH₄ emitted. Diffusion back into the water, occurring when the concentration in the headspace becomes higher than the concentration in the water, may lead to an underestimation of the ebullitive flux. We calculated the CH₄ concentration in the water based on the diffusive

[flux and a gas transfer velocity of 0.05 m/d. We therefore calculated the flux into the water for these cases, using the same gas transfer velocity, the headspace concentration at the end of the 24 hour flux incubation, and the calculated concentration in the water. Finally, we added this flux to the ebullitive flux.](#)

5

2.3.3. *Global Warming Potential*

To evaluate the net GHG effect we used a global warming potentials (GWPs) of 34 for CH₄ converting to CO₂-eq fluxes as described by Myhre et al. (2013).

10

2.4. *Plant measurements*

At the start of the experiment four extra plants were dried (148.37 ± 13.18 g FW [± SD](#) and 9.00 ± 0.94 g DW [± SD](#)) and used to analyze initial nutrient contents. At the end of the experiment all plants were collected. Water hyacinths were divided into leaves, petioles, and roots. The fresh plant samples were weighed and dried for 48h at 60°C, after which they were weighed again, grinded and homogenized. Subsequently, 200 mg of dry plant material was grinded and digested to determine total P concentrations in plants as described for the chemical analysis of sediment. An additional 3 mg of dry plant samples was combusted to determine C and N content with an elemental analyzer (Carlo Erba NA 1500, Thermo Fisher Scientific, Waltham, MA, USA).

15

20

2.5. [Plant measurements](#)[Statistical analyses](#)

Shapiro-Wilk's test and Bartlett's test were conducted to test normality of residuals and equality of error variances, respectively. Non-normal or heteroscedastic data were log transformed to meet these two requisites. ~~Linear mixed models were used to test the main effects and interactions of treatments on water characteristics, DO, GHG fluxes, GWPs and the ratios of C:N, C:P, and N:P in different plant tissues with the aquarium number as a random effect, by using R package nlme.~~ Linear mixed models were used to test the main effects and interactions of treatments over time on water characteristics, DO, GHG fluxes, and GWP. The effects on these variables, and ~~the on ratios of C:N, C:P, and N:P ratios and contents of N and P contents in different plant tissues~~ were tested with the aquarium number as a random effect, by using R package nlme. Tukey tests were performed to find differences between treatments by using R package multcomp. The effects of treatments and differences between treatments were considered significant if $P < 0.05$. All statistical analyses were carried out using the software program R (version 3.2.1; R development Core Team, 2015). All graphs were plotted by using SigmaPlot (v.11 Systat Software Inc, 2008).

3. Results

3.1. Biogeochemistry in water column and sediment

Dissolved oxygen (DO) concentrations did not vary in time ($P > 0.05$ for all treatments; data not shown), and were below saturation value (8.3 mg L^{-1} at 22°C). Average DO concentrations in the water layer were significantly lower in aquaria with plants ($3.5 \pm 0.2 \text{ mg L}^{-1}$) compared to the control aquaria without plants ($5.7 \pm 0.4 \text{ mg L}^{-1}$, $X^2 = 9.20$, $P < 0.05$). ~~The effect of coverage ($X^2 = 4.15$; $P < 0.05$) without a density effect was also seen on DO concentration where rooted treatments presented higher values compared to those than non-rooted~~ (Fig 2).

The treatments including water hyacinth had about 10-50 % lower concentrations of DTP and phosphate (PO_4^{3-}) in the surface water compared to the controls ($X^2=79.82$, $P < 0.001$ and $X^2=84.03$, $P < 0.001$ for DTP and PO_4^{3-} respectively; Table 2). In addition, they had lower NO_3^- concentrations in the surface water ($X^2=69.38$, $P < 0.001$)

5

3.2. Nutrient concentrations in different plant tissues

Plant coverage or rooting did not show effects on nutrient concentrations in different plant tissues ($P > 0.05$).

For low [coveredensity](#), P concentrations in petioles were higher in rooted plants than in not-rooted plants ($P < 0.01$; Fig S1A). In addition, only low-[densitycoverage](#) treatments showed higher P concentrations in petioles at the end of the experiment compared to the start ($P < 0.01$; Fig S1A). For all treatments P concentrations in roots were significantly higher at the end of the experiment than at the start of the experiment ($P < 0.001$; Fig S1B). Furthermore, only the high [densitycoverage](#) rooted plants had higher N concentrations in petioles at the end of the experiment compared to the start ($P < 0.001$; Fig S2). N concentrations in petioles were higher in the treatment with high [densitycoverage](#) rooted plants than all other treatments ($P < 0.001$; Fig S2).

15

In general, nutrient did not change in time or differ between treatments. There were some exceptions however (Fig 3). Rooted plants growing at high [densitycoverage](#) had a higher N: P ratio in the petioles (1.72, compared to 0.85) ($X^2=38.75$, $P < 0.001$), and the N: P ratio in the roots was lower at the end than at the start ($X^2=24.44$, $P < 0.001$).

20

3.3. Greenhouse gas fluxes

Total CH₄ fluxes were highest at high density (ANOVA for density effect; $X^2 = 9.27$, $P < 0.05$). This was mostly due to the high diffusion rates in the high coverage. Diffusion rates showed statistical difference between the treatments ($X^2 = 68.30$ $P < 0.05$). At low density coverage, diffusive CH₄ emissions were significantly higher in aquaria with rooted water hyacinth ($X^2 = 9.59$, $P < 0.01$) (Fig. 4A). At high coverage the rooted water hyacinth tended to have higher diffusive CH₄ fluxes as well, albeit non significant ($P = 0.90$). On average, CH₄ ebullition was 34.8 ± 238.27 mg CH₄ m⁻² d⁻¹ and differed due to the for high density coverage of plants compared to low density coverage and control treatments ($X^2 = 9.46$; $P < 0.05$) and did not significantly differ between the treatments (Fig 4A). As diffusion back into the water might have caused an underestimation of the ebullitive fluxes in the controls and the 50% nR treatments, this was corrected for (see Materials and Methods). Maximum loss of CH₄ from the headspace, i.e. assuming ebullition took place at the beginning of the 24 hour measurement, the loss was 14% on average. In terms of CO₂, the controls without water hyacinth functioned as a source, whereas the treatments with water hyacinth functioned as a CO₂ sink ($X^2 = 14.707$, $P < 0.001$) (Fig 4B). Day and night CO₂ fluxes for the treatments with plants were found to be opposite, showing CO₂ uptake during the day and emission during the night (Fig. S3). Differences in time were not found for GHG fluxes ($X^2 = 0.87$; $P > 0.05$ for CH₄; and $X^2 = 4.99$; $P > 0.05$ for GWP) except for CO₂ ($X^2 = 30.80$; $P > 0.05$)

4. Discussion

We found that water hyacinth presence significantly increased diffusive CH₄ fluxes. These fluxes increased with plant density coverage and, especially when plants were rooting in the sediment. The latter confirms our hypothesis regarding the role of plant-mediated CH₄ fluxes. It also, stresses the role importance of water depth for CH₄ emissions. Contrary to our expectations, however, we found no effect of rooting on CO₂ uptake

[rates](#). Due to water hyacinth's high CO₂ sequestration rates, the overall GHG budget in terms of CO₂ equivalents still resulted in water hyacinth mats being near-neutral or even a GHG sink, depending on water hyacinth [densitycoverage](#).

4.1. *The effect of water hyacinth on oxygen and CH₄ emissions*

Water hyacinth's cover lead to lower oxygen concentrations in the water column (Fig. 2). Low oxygen concentrations below other floating plant species have been reported in field and laboratory studies (Masifwa, Twongo and Denny, 2001; Nahlik and Mitsch, 2006) and have been attributed to the suppression of O₂ diffusion across the air-water interface, decrease of primary production in the water column due to lower light availability and the high oxygen demand of decomposing plant material (Reddy and DeBusk 1991).

Low oxygen concentrations [in the water](#) may, however, result in [a lower oxygenO₂ penetration depth in the sediment, in turn](#) increased [CH₄ emissions](#) (~~Bastviken et al., 2008~~) ([Huttunen et al. 2006](#)) which may, at least partially, explain why the diffusive CH₄ emission was up to 17 times higher in aquaria with water hyacinth compared to the controls. Water hyacinth has previously been reported as a CH₄ enhancer (2 to 5 times more CH₄ emissions from water hyacinth mats compared to open waters) (Banik et al., 2013). Other studies, in contrast, showed 2.6 times higher CH₄ fluxes from open waters compared to from water hyacinth mats (Attermeyer et al., 2016). We postulate that this discrepancy may well be driven by different underlying mechanisms. For one, the [densitiescoverages in these two studies](#) might have differed, with higher [densitiescoverages](#) leading to higher methanogenic rates. Additionally, along the roots of water hyacinth, CH₄ oxidation takes place due to the metanotrophic activity (Yoshida, et al., 2014), and due to a radial oxygen loss provided by this plant (Kosten, et al., [submitted2016](#)). Variation in root biomass and exudate loss, the composition and activity of microbial

communities, and water and sediment composition can be expected to affect CH₄ oxidation rates and hence CH₄ emission rates. In our study, we found that rooting led to 1.3 and 4 times higher diffusive-CH₄ emissions at high and low plant ~~density~~coverage, respectively, most possibly caused by the direct transportation of the CH₄ produced in the sediment to the atmosphere ~~through the aerenchyma of roots and leaves~~, thereby escaping CH₄ oxidation (Bastviken, 2009; Thomas, et al., 1995). Previous studies also found that shallow systems might show more effective plant-mediated CH₄ transport due to relative short distance between the sediment and the floating parts (Hamilton et al., 2014), resulting in higher emission rates.

Higher coverage led to higher diffusive CH₄ fluxes, presumably due to the production of (dissolved) organic matter substance and further lowering of O₂ concentrations. If the plant-mediated transport chimney effect indeed occurs, rooting in the sediment might avoid the formation of bubbles in the sediment, thereby decreasing ebullition and enhancing the proportion of CH₄ emitted by the plant tissue. -We did, however, not find a significant effect of rooting on ebullition in our study, only for the density of plants~~plant coverage~~.

4.2. The effect of water hyacinth on nutrient dynamics and carbon dioxide emissions

The percentage coverage by plants and their access to the sediment did not change nutrient uptake and allocation in a consistent way, although the rooting plants grown at high density~~coverage~~ ~~rooting plants~~ showed higher N: P ratios (Fig. 3), due to higher N concentration (Fig S2). Relatively high N concentrations under high-density~~coverage~~ conditions have been found in the field as well and have been related to N supply to the plant (Reddy et al., 1989). The general absence of a strong effect of root access to the sediment on plant nutrient contents suggests that the plants are capable of mobilizing nutrients from the sediment even without direct contact. This has also been demonstrated for the floating macrophyte *Stratiotes aloides*, for which the lowering

of O₂ levels due to high coverage can promote P release from the sediment by weakening the bonds of Fe-P complexes (Harpenslager, et al., 2016). The fact that water column nutrient concentrations tended to be higher in the treatments where the plants are rooted in the sediment (Table 2) suggests, however, that the plants preferably tap into the rich sediment nutrient pools directly. Pore-water concentrations of N and P were 220 and 30 times higher than in the surface water (results not shown). *Egeria densa*, *Hydrilla verticillata*, and *Myriophyllum spicatum* have been reported to only take up P from the sediment (Barko and Smart, 1980).

More efficient nutrient uptake when rooted in the sediment could lead to higher growth rates and concomitant CO₂ sequestration. We did not find, however, higher CO₂ sequestration in our rooted treatments. We only found a clear difference between the [densitiescoverages](#), with on average 1.6 times lower CO₂ sequestration rates at high [densitiescoverages](#), which we attribute to the limited space for growth.

On average our plant treatments sequestered -3.4 ± 2.2 g CO₂ m⁻² day⁻¹, regardless of [densitycoverage](#) and the position of the roots. This is notably higher than sequestration rates of other [rooted](#) aquatic plants, such as *Typha domingensis* and *Eichhornia azurea*, showing sequestration rates around -0.09 g CO₂ m⁻² day⁻¹ (Gripp et al. 2013), [for instance. This results shows that other rooted plant species presented less CO₂ uptake than water hyacinth, rooted or not. Our results match with](#) ~~The rates we measured are in~~ the range [found previously for water hyacinth](#) between -3.4 and -5.4 g CO₂ m⁻² day⁻¹, [found for E.c.](#) in field conditions (Peixoto et al., 2016; Attermeyer et al., 2016).

In aquaria without water hyacinth, CO₂ fluxes took place leading a 24h net emission of, on average, 0.3 g CO₂ m⁻² day⁻¹ (Fig. 4). The net emissions from the non-vegetated [aquaria controls](#) contrast the net CO₂ sequestration in the plant treatments indicating that the plants offset the CO₂ emissions from [the systems without](#)

plant/open waters. Clearly, the net emissions from both vegetated and non-vegetated systems strongly depends on sediment CO₂ production rates related to the input of decaying plant material.

4.3. *Effects of water hyacinth on the overall GHG balance*

Under the experimental conditions of our study in the absence of water hyacinth, CH₄ emissions were modest and net CO₂ emissions took place, leading to an overall emission of GHGs (Fig. 5). At low density/coverage, however, water hyacinth was a net sink of GHGs, regardless of the position of the roots. At high density/coverage CO₂ sequestration only partially counterbalanced CH₄ emissions, thereby making the system become a small GHG source.

Whether water hyacinth acts as a GHG sink or source depends on the balance of its effects on CH₄ emissions and CO₂ uptake rates. We here show that this balance strongly depends on plant coverage and rooting, with partly opposite effects with respect to CO₂ and CH₄. The plants tend to enhance CH₄ emissions especially at high coverage and when rooting in the sediment, whereas CO₂ uptake rates are highest at low coverage where growth was not space-limited and nutrient availability per plant is higher.

Ebullition played an important role in the overall GHG balance, since it accounted, on average, for 58% of the total CH₄ emissions for all treatments and even reached 86% at low density/coverage. This underlines once more that ebullition is one of the most important forms/pathways of CH₄ contribution/emission from wetlands to the atmosphere (Bastviken et al, 2008; Sawakuchi et al., 2014; Segarra et al., 2013; Coulthard, et al., 2009).

Our results highlight that the presence of water hyacinth mats can alter GHG emissions. CO₂ sequestration rates are enhanced and hence can trigger a regional effect offsetting the greenhouse gas emissions for open waters. Using water hyacinth for nutrient-rich wastewater purification under a relatively low [densitycoverage](#) (like 50% coverage) by regular harvest will likely reduce the emission of CH₄ and increase the sequestration of CO₂, especially when roots are prevented from reaching the sediment. As a main conclusion, we here showed that access to the sediment, as related to water depth, and plant [densitycoverage](#) are crucial factors influencing both nutrient dynamics and GHG emissions, which may explain the discrepancies reported in literature and should be taken into account when making regional GHG balances.

5. Author contribution

The experiment was designed by Ernandes S. Oliveira Junior, Yingying Tang, Sarian Kosten and Leon P. M. Lamers, and executed by Ernandes S. Oliveira Junior, Yingying Tang and Sanne van den Berg. The manuscript was written by Ernandes S. Oliveira Junior and Yingying Tang with contribution of all co-authors.

6. Acknowledgements

The authors would like to thank Walter Hendrickx, Yvette Evers for the Water hyacinth maintenance and Gerard van der Weerden for providing experimental space in the greenhouse facilities. We thank Paul van der Ven, Ankie De Vries-Brock, Germa Verheggen, Sebastian Krosse, and Roy Peters for their assistance with the chemical analyses, [and Peter Walraven for the construction of the aquaria set](#). Han Wang was essential for the experiment implementation and data acquisition. [We also thank Ralf Aben for developing a macro to calculate the diffusive fluxes-and for his fruitful contributions to our discussions about greenhouse gas fluxes. Simone Jaqueline Cardoso](#)

[played a key role for the statistical analyses of the data.](#) Ernandes Sobreira was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES by the Science without Borders scholarship program Process BEX 13607/13-8, and the project was funded by Koninklijke Nederlandse Akademie van Wetenschappen (KNAW) and Fundação de Amparo a Pesquisa de Mato Grosso (FAPEMAT). Yingying Tang was funded by the China
5 Scholarship Council (CSC; file number 201206140019). Sarian Kosten was supported by NWO-VENI grant 86312012.

7. References

10 [Almeida, R. et al. 2016. High Primary Production Contrasts with Intense Carbon Emission in a Eutrophic Tropical Reservoir. *Frontiers in Microbiology*. 7, \(2016\), 717.](#)

Aoyama I. & Nishizaki H. (1993) Uptake of nitrogen and phosphate, and water purification by water hyacinth *Eichhornia crassipes* (Mart.) Solms. *Water Science and Technology* 28(7), 47-53.

15 Attermeyer K., Flury S., Jayakumar R., Fiener P., Steger K., Arya V., Wilken F., van Geldern R. & Premke K. (2016) Invasive floating macrophytes reduce greenhouse gas emissions from a small tropical lake. *Scientific reports* 6.

Banik A., Sen M. & Sen, S. (1993) Methane emissions from waterhyacinth-infested freshwater ecosystems. *Chemosphere* 27(8), 1539-1552.

Barko J. & Smart R. (1980) Mobilization of sediment phosphorus by submersed freshwater macrophytes. *Freshwater Biology*. 10, 3 (1980), 229–238.

20 Bastviken D., Cole J. J., Pace M. L. & Van de Bogert M. C. (2008) Fates of methane from different lake habitats: Connecting whole-lake budgets and methane emissions. *Journal of Geophysical Research: Biogeosciences* 113(G2).

25 Bastviken D. (2009) Methane, p. 783–805. *In* G. E. Likens [ed.], *Encyclopedia of inland waters*. Elsevier.
Chanakya H., Borgaonkar S., Meena G. & Jagadish K. (1993) Solid-phase biogas production with garbage or water hyacinth. *Bioresource technology* 46(3), 227-231.

[Bharati, K., Mohanty, S.R., Singh, D.P., Rao, V.R., Adhya, T.K., 2000. Influence of incorporation or dual cropping of Azolla on methane emission from a flooded alluvial soil planted to rice in eastern India. *Agric. Ecosyst. Environ.* 79 \(1\), 73-83.](#)

[Caraco, N. F., and J. J. Cole. 2002. Contrasting impacts of a native and alien macrophyte on dissolved oxygen in a large river. *Ecol. Appl* 12: 1496–1509.](#)

[Chanakya, H., S. Borgaonkar, G. Meena, and K. Jagadish. 1993. Solid-phase biogas production with garbage or water hyacinth. *Bioresour. Technol.* **46**: 227-231.](#)

5 [Cronk, J. K., and M. S. Fennessy. 2016. *Wetland plants: biology and ecology*. CRC press.](#)

~~Coulthard T. J., Baird A. J., Ramirez J. & Waddington M. (2009). Methane dynamics in peat: importance of shallow peats and a novel reduced complexity approach for modeling ebullition. In: Baird AJ, Belyea LR, Comas X, Reeve AS, Slater LD (eds) Carbon cycling in northern peatlands, geophysical monograph series 184. AGU, Washington DC, pp 173–185.~~

10 [Dacey, J., and M. Klug. 1979. Methane efflux from lake sediments through water lilies. *Science* **203**: 1253-1255.](#)

[Grasshoff K. & Johannsen H. \(1972\) A new sensitive and direct method for the automatic determination of ammonia in sea water. *Journal du Conseil* 34\(3\), 516-521.](#)

15 [Grasset, C., Abril, G., Guillard, L., Delolme, C., & Bornette, G. \(2016\). Carbon emission along a eutrophication gradient in temperate riverine wetlands: effect of primary productivity and plant community composition. *Freshwater Biology*. 61 \(9\), 1405–1420.](#)

Gripp A. d. R., Marinho C. C., Sanches L. F., Petruzzella A. & Esteves, F. d. A. (2013) The role played by aquatic macrophytes regarding CO₂ balance in a tropical coastal lagoon (Cabiúnas Lagoon, Macaé, RJ). *Acta Limnologica Brasiliensia* 25(3), 291-301.

20 [Grosse, W., Armstrong, J., & Armstrong, W. \(1996\). A history of pressurised gas-flow studies in plants. *Aquatic Botany*, 54\(2\), 87-100.](#)

Gutierrez E., Ruiz E., Uribe E. & Martinez J. (2001) Biomass and productivity of water hyacinth and their application in control programs. *Biological and Integrated Control of Water Hyacinth, Eichhornia crassipes* 102, 109-199.

25 [Harpenslager S., Lamers L., van der Heide T., Roelofs J. & Smolders A. \(2016\) Harnessing facilitation: Why successful re-introduction of *Stratiotes aloides* requires high densities under high nitrogen loading. *Biological Conservation*. 195, 17–23.](#)

[Hamilton, S.K. S.J. Sippel, J.P. Chanton, and J.M. Melack. 2014. Plant-mediated transport and isotopic composition of methane from shallow tropical wetlands. *Inland Waters* 4: 369-376. doi 10.5268/IW-4.4.734](#)

30 [Henriksen A. \(1965\) An automatic method for determining low-level concentrations of phosphates in fresh and saline waters. *Analyst* 90\(1066\), 29-34.](#)

[Huttunen, J. T., T. S. Väisänen, S. K. Hellsten, and P. J. Martikainen. 2006. Methane fluxes at the sediment-water interface in some boreal lakes and reservoirs. *Boreal Environment Research* **11**: 27-34.](#)

[Jiangqing D., Ren W., Weidong F. & Guoliang Z. \(2001\) Water hyacinth in China: Its distribution, problems and control status. *Biological and integrated control of water hyacinth, Eichhornia crassipes* 102.](#)

5 Kamphake L., Hannah S. & Cohen J. (1967) Automated analysis for nitrate by hydrazine reduction. *Water research* 1(3), 205-216.

Kosten S., Piñeiro M., de Goede E., de Klein J. & Ettwig, K. [submitted 2016](#). Fate of methane in free floating plant dominated systems. *Water Research* [104\(1\), 200-207](#).

10 [Konnerup, D., B. K. Sorrell, and H. Brix. 2011. Do tropical wetland plants possess convective gas flow mechanisms? *New Phytologist* **190**: 379-386.](#)

Malik A. (2007) Environmental challenge vis a vis opportunity: the case of water hyacinth. *Environment international* 33(1), 122-138.

Mandi L. (1994) Marrakesh wastewater purification experiment using vascular aquatic plants *Eichhornia crassipes* and *Lemna gibba*. *Water Science and Technology* 29(4), 283-287.

15 Masifwa W., Twongo T. & Denny, P. (2001) The impact of water hyacinth, *Eichhornia crassipes* (Mart) Solms on the abundance and diversity of aquatic macroinvertebrates along the shores of northern Lake Victoria, Uganda. *Hydrobiologia*. 452, 1/3, 79–88.

20 Myhre, G., Shindell, D. and Bréon F. M. (2013) Anthropogenic and Natural Radiative Forcing. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.

Nahlik A. & Mitsch W. (2006) Tropical treatment wetlands dominated by free-floating macrophytes for water quality improvement in Costa Rica. *Ecological Engineering*. 28, 3, 246–257

Olsen S. R. (1954) Estimation of available phosphorus in soils by extraction with sodium bicarbonate.

25 Peixoto R. B., Marotta H., Bastviken D., & Enrich-Prast A. (2016) Floating Aquatic Macrophytes Can Substantially Offset Open Water CO₂ Emissions from Tropical Floodplain Lake Ecosystems. *Ecosystems*.

Pinto-Coelho R. & Greco M. (1999) The contribution of water hyacinth (*Eichhornia crassipes*) and zooplankton to the internal cycling of phosphorus in the eutrophic Pampulha Reservoir, Brazil. *Hydrobiologia*. 411, 115–127.

30 Polprasert C. & Khatiwada N. R. (1998) An integrated kinetic model for water hyacinth ponds used for wastewater treatment. *Water Research* 32(1), 179-185.

~~Reddy K. & Tucker J. (1983) Productivity and nutrient uptake of water hyacinth, *Eichhornia crassipes* L. Effect of nitrogen source. *Economic botany* 37(2), 237–247.~~

~~Reddy K. & Debusk W. (1984) Growth characteristics of aquatic macrophytes cultured in nutrient enriched water: I. Water hyacinth, water lettuce, and pennywort. *Economic Botany*. 38, 2, 229–239~~

5 Reddy K. & DeBusk W. (1991) Decomposition of water hyacinth detritus in eutrophic lake water. *Hydrobiologia* 211(2), 101-109.

Reddy K., Agami M. & Tucker J. (1989) Influence of nitrogen supply rates on growth and nutrient storage by water hyacinth (*Eichhornia crassipes*) plants. *Aquatic Botany* 36(1), 33-.

10 Reddy K., Agami M. & Tucker J. (1990) Influence of phosphorus on growth and nutrient storage by water hyacinth (*Eichhornia crassipes* (Mart.) Solms) plants. *Aquatic botany* 37(4), 355-365.

~~Ribaudo, C., Bartoli, M., Longhi, D., Castaldi, S., Neubauer, S. C., & Viaroli, P. (2012). CO₂ and CH₄ fluxes across a *Nuphar lutea* (L.) Sm. stand. *Journal of Limnology*, 71(1), 21.~~

15 ~~Sawakuchi, H., Bastviken, D., Sawakuchi, A., Krusche, A., Ballester, M. and Richey, J. 2014. Methane emissions from Amazonian Rivers and their contribution to the global methane budget. *Global Change Biology*. 20, 9 (2014), 2829–2840~~

~~Segarra, K. E. A., Samarkin, V., King, E., Meile, C., Joye, S. B. (2013) Seasonal variations of methane fluxes from an unvegetated tidal freshwater mudflat (Hammersmith Creek, GA). *Biogeochemistry* 115: 349-361.~~

Thomas K., Benstead J., Davies K. & Lloyd D. (1996) Role of wetland plants in the diurnal control of CH₄ and CO₂ fluxes in peat. *Soil Biology and Biochemistry*. 28, 1, 17–23.

20 Tomassen H., Smolders A. J., Limpens J., Lamers L. P. & Roelofs, J.G. (2004) Expansion of invasive species on ombrotrophic bogs: desiccation or high N deposition? *Journal of applied Ecology* 41(1), 139-150.

Villamagna A. & Murphy, B. (2010) Ecological and socio-economic impacts of invasive water hyacinth (*Eichhornia crassipes*): a review. *Freshwater biology* 55(2), 282-298.

25 Wilson J. R., Holst N. & Rees M. (2005) Determinants and patterns of population growth in water hyacinth. *Aquatic Botany* 81(1), 51-67.

Xie Y., Wen M., Yu D. & Li Y. (2004) Growth and resource allocation of water hyacinth as affected by gradually increasing nutrient concentrations. *Aquatic Botany*. 79, 3, 257–266.

Yoshida N., Iguchi H., Yurimoto H., Murakami A. & Sakai Y. (2014) Aquatic plant surface as a niche for methanotrophs. *Frontiers in Microbiology* 5: 30.

30 ~~Zhao F., Xi S., Yang X., Yang W., Li J., Gu B. & He Z. (2012) Purifying eutrophic river waters with integrated floating island systems. *Ecological Engineering* 40, 53–60.~~

Tables

Table 1 Sediment characteristics at the beginning of the experiment (mean ± SEM; n=3). All analyses were performed using fresh or dry sediment (see text 2.2).

<u>Characteristics</u>	<u>Unit</u>	<u>Sediment</u>
<u>Organic matter content</u>	<u>%</u>	<u>3.32±0.42</u>
<u>Total-P</u>	<u>μmol g⁻¹ DW</u>	<u>15.65±1.27</u>
<u>Olsen-P</u>	<u>μmol g⁻¹ DW</u>	<u>0.85±0.10</u>
<u>Salt-extractable NH₄⁺</u>	<u>μmol g⁻¹ DW</u>	<u>0.25±0.08</u>
<u>Salt-extractable NO₃⁻</u>	<u>μmol g⁻¹ DW</u>	<u>0.02±0.00</u>
<u>Total-Fe</u>	<u>μmol g⁻¹ DW</u>	<u>88.13±4.65</u>
<u>Total-Al</u>	<u>μmol g⁻¹ DW</u>	<u>81.49±3.80</u>
<u>Total-Ca</u>	<u>μmol g⁻¹ DW</u>	<u>100.43±6.63</u>

5

<u>Characteristics</u>	<u>Unit</u>	<u>Sediment</u>
<u>Organic matter content</u>	<u>%</u>	<u>3.32±0.24</u>
<u>Total-P</u>	<u>μmol g⁻¹ DW</u>	<u>15.65±0.73</u>
<u>Olsen-P</u>	<u>μmol g⁻¹ DW</u>	<u>0.85±0.06</u>

Salt extractable NH₄⁺	μmol g⁻¹ DW	0.25±0.05
Salt extractable NO₃⁻	μmol g⁻¹ DW	0.02±0.00
Total Fe	μmol g⁻¹ DW	88.13±2.69
Total Al	μmol g⁻¹ DW	81.49±2.20
Total Ca	μmol g⁻¹ DW	100.43±3.83

5

10

15

Table 2 Water characteristics in surface water during the experiment. All concentrations are given in $\mu\text{mol L}^{-1}$. DOC concentrations were determined at the end of the experiment (mean \pm SDEM; n=4), whereas other parameters were analyzed multiple times during the experiment (overall average are given, mean \pm SDEM; n=4).

Characteristics	C w/o mesh	C with mesh	50%nR	50%R	100%nR	100%R
DTP	17.4±10.3 ^a	11.1±2.7 ^a	2.2±0.3 ^b	7.8±9.2 ^b	6.4±8.3 ^b	3.7±2.6 ^b
PO ₄ ³⁻	12.8±5.8 ^a	9.5±2.7 ^a	1.6±0.3 ^b	7.8±10.8 ^b	1.4±0.2 ^b	2.8±2.0 ^b
NH ₄ ⁺	16.9±12.4	6.2±1.1	6.8±0.9	60.6±108.4	39.8±67.5	6.2±1.1
NO ₃ ⁻	9.5±1.6 ^a	1.2±1.1 ^b	1.8±2.1 ^b	0.4±0.2 ^b	0.6±0.4 ^b	0.4±0.3 ^b
TIC	734.3±198.4 ^b	1020.4±133.3 ^a	1052.0±201.7 ^a	1417.0±556.1 ^a	1319.0±259.4 ^a	1212.0±206.5 ^a
DOC	944.0±271.8	890.7±663.1	855.5±396.0	469.3±548.5	997.2±547.3	629.0±290.5

Significant differences among treatments are indicated by different lower case letters.

5

Characteristics	C w/o mesh	C with mesh	50%nR	50%R	100%nR	100%R
DTP	17.4±5.2 ^a	11.1±1.4 ^a	2.2±0.1 ^b	7.8±4.6 ^b	6.4±4.2 ^b	3.7±1.3 ^b
PO ₄ ³⁻	12.8±2.9 ^a	9.5±1.4 ^a	1.6±0.2 ^b	7.8±5.4 ^b	1.4±0.1 ^b	2.8±1.0 ^b
NH ₄ ⁺	16.9±6.2	6.2±0.6	6.8±0.5	60.6±54.2	39.8±33.8	6.2±0.6

NO ₃ ⁻	9.5±0.8 ^a	1.2±0.6 ^b	1.8±1.1 ^b	0.4±0.1 ^b	0.6±0.2 ^b	0.4±0.2 ^b
TIC	734.3±99.2 ^b	1020.4±66.6 ^a	1052.0±100.9 ^a	1417.0±278.0 ^a	1319.0±129.7 ^a	1212.0±103.2 ^a
DOC	944.0±135.9	890.7±331.6	855.5±198.0	469.3±274.3	997.2±273.6	629.0±145.3

Significant differences among treatments are indicated by different lower case letters.

Figure legends

Figure 1. Experimental design. C w/o mesh represents control without mesh; C w mesh represents control with mesh; 50% nR represents low [densitycoverage](#) with mesh; 50% R represents low [densitycoverage](#) without mesh; 100% nR represents high [densitycoverage](#) with mesh; 100% R represents high [densitycoverage](#) without mesh.

Figure 2. Mean dissolved oxygen concentrations (\pm [DEM](#)) of the water layer at 20 cm depth for controls (C), low [densitycoverage](#) (50%), and high [densitycoverage](#) (100%) of water hyacinth with (R) or without rooting (nR) in the sediment. Different lower case letters indicate significant differences between treatments by post hoc test ($P < 0.001$).

Figure 3. Ratios (\pm [SD](#)) between C and P ([whitelight grey](#)), between C and N ([dark shadedgrey](#)), and between N and P ([dottedblack](#)) in petioles (left panel) and roots (right panel) of water hyacinth for low [densitycoverage](#) (50%) and high [densitycoverage](#) (100%) with (R) or without (nR) roots in the sediment at the end and start of the experiment. All nutrient ratios are given in mol mol⁻¹. Different lower case letters indicate significant differences between treatments including the start of the experiment ($P < 0.05$). Note the log₁₀ scale on the y-axis.

Figure 4. CH₄ (left) and CO₂ (right) fluxes (\pm [DEM](#)) for controls (C), low [densitycoverage](#) (50%), and high [densitycoverage](#) (100%) of water hyacinth with (R) or without rooting (nR) in the sediment. ~~Different lower case letters and upper case letters indicate significant differences between treatments ($P < 0.05$) indicating, from top to bottom: total fluxes, ebullitive flux, and diffusive flux.~~ ~~for diffusion and the total flux, respectively. No statistic difference was found for ebullition.~~ Note different scales for the y-axis. Negative numbers refer to sequestration.

Figure 5. Global warming potential (GWP; mean \pm SEM) for controls (C), low densitycoverage (50%) and high densitycoverage (100%) water hyacinth coverage with or without rooting in the sediment. Different lower case letters indicate significant differences between treatments ($P < 0.001$).

5

10

15

5




Fig 1

10





Fig 2




Fig 3








Fig 4






Fig 5