1	Bacterial production in subarctic peatland lakes enriched by thawing permafrost
2	
3	Bethany N. Deshpande ¹ , Sophie Crevecoeur ¹ , Alex Matveev ¹ , and Warwick F. Vincent ¹
4	
5	¹ Centre for Northern Studies (CEN), Biology Department and Takuvik Joint International
6	Laboratory, Université Laval, Québec, QC, Canada
7	Running title: Bacterial production in thermokarst lakes
8	Correspondence to: B. N. Deshpande (bethany.deshpande.1@ulaval.ca)

10 **Abstract.** Peatlands extend over vast areas of the northern landscape. Within some of these areas, 11 lakes and ponds are changing in size as a result of permafrost thawing and erosion, resulting in 12 mobilisation of the carbon-rich peatland soils. Our aims in the present study were to characterize 13 the particle, carbon and nutrient regime of a set of thermokarst (thaw) lakes and their adjacent 14 peatland permafrost soils in a rapidly degrading landscape in subarctic Ouébec, Canada, and by 15 way of fluorescence microscopy, flow cytometry, production measurements and an in situ 16 enrichment experiment, determine the bacterial characteristics of these waters relative to other 17 thaw lakes in more mineral terrain and to non-permafrost-affected rock-basin lakes in the region. 18 The soil active layer in a degrading palsa (peatland permafrost mound) adjacent to one of the lakes contained an elevated carbon content (51% of dry weight), low C:N ratios (17:1 by mass), 19 20 and large stocks of other elements including N (3% of dry weight), Fe (0.6%), S (0.5%), Ca 21 (0.5%) and P (0.05%). Two permafrost cores were obtained to a depth of 2.77 m in the palsa, and 22 computerized tomography scans of the cores confirmed that they contained high concentrations 23 (>80%) of ice. Upon thawing, the cores released nitrate and dissolved organic carbon (from all 24 core depths sampled), and soluble reactive phosphorus (from bottom depths), at concentrations 25 well above those in the adjacent lake waters. The active layer soil showed a range of particle 26 sizes with a peak at 229 µm, and this was similar to the distribution of particles in the upper 27 permafrost cores. The particle spectrum for the lake water overlapped with those for the soil, but 28 extended to larger (surface water) or finer (bottom water) particles. On average, more than 50% 29 of the bacterial cells and bacterial production was associated with particles $>3 \mu m$. This relatively 30 low contribution of free-living cells (operationally defined as the $<1 \mu m$ fraction) to bacterial 31 production was a general feature of all of the northern lakes sampled, including other thaw lakes 32 and shallow rock-basin lakes (average \pm SE of 25 \pm 6%). However, a distinguishing feature of the 33 peatland thaw lakes was significantly higher bacterial specific growth rates, which averaged 4 to 34 7 times higher values than in the other lake types. An in situ enrichment experiment showed no 35 difference between organic carbon or phosphorus enrichment treatments at day 5 relative to a 36 control treatment, however there was a significant, >100% increase in bacterial growth rates 37 between days 1 and 5 in the soil and the carbon plus phosphorus enrichments. Collectively these 38 results indicate that particles, nutrients and carbon are released by degrading permafrost peatland 39 soils into their associated thermokarst lakes, creating favorable conditions for production by 40 particle-based as well as free-living aquatic bacterial communities. The reduced bacterial 41 concentrations despite high cellular growth rates imply that there is strong control of their 42 population size by loss-related factors such as grazing and viral lysis.

43

44 Keywords: bacteria, climate change, northern lakes, peatlands, permafrost, thermokarst

46 1 Introduction

47 Aquatic environments in permafrost landscapes are biogeochemical hotspots that provide 48 microbial communities with access to substrates that have been previously locked in frozen soils 49 for centuries to millennia (Kling et al., 1991; Vonk et al., 2015). The highest soil organic carbon contents of the permafrost environment are found in areas of sporadic or isolated permafrost 50 peatland, which span more than 5×10^5 km² of land in the circumpolar North (Tarnocai et al., 51 52 2009). The availability of organic substrates from these vast permafrost peatland environments to 53 bacterial communities in aquatic ecosystems is likely to be influenced by a variety of factors, 54 including the soil biogeochemical composition, particle and molecular size, and potentially other 55 environmental factors, such as nutrient concentrations and temperature. Lakes in the boreal zone 56 with flat catchments dominated by forest and peatland areas are strongly heterotrophic systems 57 that have high rates of bacterial production (e.g., Kankaala et al., 2006), but less is known about 58 the bacterial characteristics of peatland lakes further northwards, in permafrost regions.

Decomposition processes in aquatic ecosystems are mediated by bacterial communities that can be separated into two ecological groups: (1) free-living bacteria, in which solitary cells are suspended in the medium and break down dissolved organic matter, and (2) particle-attached bacteria, in which the communities are associated with various size-fractions of particles and break down this particulate organic material, as well as accessing dissolved materials (Kirchman and Mitchell, 1982). In both cases, these activities result in consumption of oxygen, production of carbon dioxide, and the mineralization of organic materials.

In most lake and ocean environments, free-living bacteria dominate the total bacterial
community in terms of numerical abundance (Unanue et al., 1992; Kirchman and Mitchell, 1982;
Simon et al., 1990; Lami et al., 2009). However, there are many environments in which particle-

69 based bacteria play a major role. Several studies from estuarine- and marine-waters show that 70 particle-attached populations can account for 10 to 100 times more bacterial cells and activity 71 than free-living bacteria, depending on particle concentrations (Crump et al., 1999; Ghiglione et 72 al., 2007; Acinas et al., 1999; Lami et al., 2009). A study of seven freshwater ecosystems 73 including two marsh-systems showed that a small proportion of particle-attached bacteria (<10%) 74 was associated with $\geq 40\%$ of the bacterial community activity, and it was hypothesized that lake 75 size may influence the presence of particulate matter for colonization via interactions between 76 sediments and the water column (Kirchman and Mitchell, 1982). In the turbid waters of the 77 Mackenzie River, which are strongly influenced by permafrost as well as fluvial erosion, more 78 than 90% of the total bacterial production was associated with particles $>3 \mu m$ (Vallières et al., 79 2008). Similarly, in the coastal waters of the Beaufort Sea, the >3 µm size-fraction often 80 accounted for >50% of total bacterial production (Garneau et al., 2006; Galand et al., 2008). 81 Other research has identified that a dominant portion (>50%) of bacterial cells were attached to 82 particles (Masin et al., 2012), especially in turbid systems with high concentrations of particulate 83 organic matter (Fletcher 1991; Lami et al., 2009).

84 Particle localisation of microbial processes has wide-ranging implications for trophic 85 processes in the water column and material fluxes to the sediments. In the oceanographic literature, large particles are referred to as "marine snow", and many studies that have shown the 86 87 importance of these aggregates for food-web dynamics (Michaels and Silver, 1988; Alldredge 88 and Silver, 1988; Kiørboe, 2001). Similarly, there has been attention given to "lake snow", 89 although lake particulate matter is likely to be more of terrestrial origin, and therefore of different 90 chemical composition than marine snow. Grossart and Simon (1993) found lake particle aggregates to be densely colonized by bacteria, with approximately 10⁸ bacterial cells per mL, 91

92 100 times higher than concentrations in the bulk water. The presence of particle-attached bacteria 93 may also result in greater community diversity, as has been demonstrated in studies of estuarine 94 mixing zones (Crump et al., 1998; Waidner and Kirchman, 2007). A study by Ploug and Grossart 95 (2000) identified a positive correlation between bacterial production, particulate organic carbon 96 and aggregate size. Size also affects sinking velocities through Stokes' Law, and therefore the 97 flux rates for transfer of materials from the pelagic zone to the sediments (Ploug and Grossart, 98 2000).

99 Permafrost thaw lakes are rich in suspended particulate material as a result of permafrost 100 erosion (Breton et al., 2009; Watanabe et al., 2011). They are also continuously supplied with 101 organic as well as inorganic inputs from the thermokarst processes in their surrounding 102 catchments (Vonk et al., 2015), and there is increasing attention to the bacterial responses to such 103 inputs. In the Mackenzie River Delta region, a mesocosm assay showed decreased planktonic 104 bacterial production rates in response to permafrost soil enrichment, but a strong increase in 105 benthic bacterial production rates (Moquin and Wrona, 2014). In a set of subarctic thermokarst 106 ponds, late winter bacterial production was low and dominated by free-living bacterioplankton, 107 while much higher production rates in summer were dominated by particle-attached bacteria, 108 correlated with terrestrial carbon concentrations (Roiha et al., 2015). Permafrost peatland lakes 109 are known to be strong emitters of greenhouse gases with diverse microbiota (Liebner et al., 110 2015), but little is known about the controls on their bacterial populations and productivity.

Our first aim in the present study was to investigate the physical and geochemical properties of permafrost peatland soils in order to better understand the allochthonous carbon, nutrient and particle inputs to peatland thermokarst lakes and their potential effects on bacterial activity. Secondly, we aimed to determine the distribution of bacterial production between the suspended 115 and free-living size fractions in these peatland lakes enriched with particles from eroding 116 permafrost. Our final aim was to experimentally determine the responses of peatland lake 117 bacteria to permafrost soil inputs. Two hypotheses were evaluated: firstly, that particle-attached 118 bacteria account for the dominant (>50%) fraction of the total microbial activity, and secondly 119 that organic carbon and nutrients originating from the erosion of permafrost soils promote 120 bacterial metabolic activity and cellular growth. We undertook this research in subarctic Québec 121 and first determined the chemical composition of the soil active layer and underlying permafrost 122 in a palsa (organic permafrost mound) that was adjacent to and eroding into a peatland thaw lake. 123 We examined the distribution of soil particle sizes and compared these with particle distributions 124 in permafrost peatland lakes, with comparisons of particle size spectra, nutrients and organic 125 carbon with other types of lakes in the region. We then measured bacterial abundance (BA) and 126 production rates (BP) associated with different size fractions in the peatland lakes and others, and 127 tested the effects of phosphorus, carbon and soil enrichment on bacterial concentrations and 128 activity by way of an in situ incubation experiment.

129 2 Materials and methods

130 **2.1** Study sites

The primary sampling sites were near Kuujjuarapik-Whapmagoostui (K-W, 55° 17' N, 77° 47' W) in thermokarst lakes and their surrounding soils in the peatland palsa valley of the Sasapimakwananistikw River (SAS; Fig. 1). Additional sampling for comparison took place in thermokarst lakes in the mineral lithalsa (mineral permafrost mounds) valley of the Kwakwatanikapistikw River (KWK) near K-W; glaciated rock-basin lakes, also near K-W, but not influenced by permafrost (RBL); and lithalsa lakes up to 200 km further to the North (SEC, BGR, NAS). This coastal region of northern Québec has experienced rapid climate change, including an increase in mean annual temperatures from -4.2 °C for 1932–1960 to -2.6 ± 1.2 °C for 2001-2010, accompanied by changes in the permafrost landscapes, including expansion of the tree line, changes to snow conditions, and various changes in thermokarst lakes frequency and extent (Bhiry et al., 2011). A total of 15 lakes were sampled at five permafrost-affected sites and one non-permafrost-affected area (Fig. 1; detailed location and depth information is given in Table S1 in the Supplement).

144 The SAS valley is an area of sporadic permafrost (<2% frozen ground) mainly colonized 145 by Sphagnum and Carex, with dark brown and black colored thermokarst lakes (Arlen-Pouliot 146 and Bhiry, 2005; Fillion et al., 2014). This peat bog contains more than 50 isolated permafrost 147 mounds (palsas; Fig. 2) covered by a 60 cm soil active layer (Arlen-Pouliot and Bhiry, 2005; 148 Fillion et al., 2014). Two sets of lakes were sampled in the SAS valley: the SAS1 set of lakes was 149 located to the south and the SAS2 lakes were located to the north of the river. The SAS lakes are 150 stratified, heterotrophic lakes with anoxic zones throughout the majority of the water column, and 151 surface waters that are under-saturated in oxygen (Deshpande et al., 2015). These lakes have abundant and active methanotrophic bacterial communities indicating their high methane supply 152 153 rates (Crevecoeur et al., 2015), and they emit greenhouse gases to the atmosphere at flux rates up to 13 mmol CH₄ m⁻² d⁻¹ and 240 mmol CO₂ m⁻² d⁻¹ (Matveev et al. 2016). 154

The valley of the Kwakwatanikapistikw River (KWK) is also in an area of sporadic permafrost, located only 8 km from SAS. It is colonized by lichen-moss, shrub-tundra vegetation, and forest patches in wind-sheltered areas (Arlen-Pouliot and Bhiry, 2005), and contains numerous thermokarst lakes that vary in color from blue-green to brown. The SEC site (55° 42.067' N, 76° 38.604' W) was located 86.0 km north-east of K-W and the BGR site was located 160 near the Hudson Bay and the village of Umijuag (56° 36.63' N, 76° 12.85' W), both in areas of 161 discontinuous permafrost. BGR1 appears to be a rapidly expanding lake, with a maximum depth 162 of 3.2 in 2006 (Breton et al., 2009) that had increased to 4.0 m in 2013 (Deshpande et al., 2015), 163 and 4.2 m in the present study. The Nastapoka River valley (NAS) contains thermokarst lakes 164 that were formed by thawing lithalsa mounds in this region of continuous permafrost. The non-165 permafrost-affected, rock-basin lakes (RBL) in the K-W region are of similar area and shallow 166 depth (0.8 - 2.1 m) to the thermokarst systems, but with rock substrate catchments rather than 167 permafrost soils.

168 2.2 Lake sampling and profiling

169 Sampling took place during three consecutive summers from 2012 to 2014. At each visit, surface 170 water was sampled using sterile plastic containers. A Van Dorn sampler was used to retrieve 171 samples from the oxycline or the bottom waters, 0.2 m above the bottom sediments. Water samples were analysed for soluble reactive phosphorus (SRP; detection limit of 0.5 μ g P L⁻¹), 172 total nitrogen (TN; detection limit of 0.008 mg N L⁻¹), dissolved organic carbon (DOC; detection 173 limit of 0.05 mg DOC L⁻¹), Chlorophyll a (Chl a), and total suspended solids (TSS) according to 174 175 methods in Laurion et al., (2010). Ammonium and nitrate (including nitrite) concentrations were 176 measured in water from the SAS valley only using a Lachat autoanalyzer (detection limit of 10 µg N L⁻¹). Soil was sampled from the active layer directly next to a lake at each site and stored in 177 178 sterile plastic bags. Frozen samples of both water and soils were returned to Université Laval 179 (Laboratoire de géomorphologie et de sédimentologie, FFGG) to determine particle size 180 distribution via Laser Particle Size Analyzer (LA-950V2, Horiba Ltd., Japan).

For CDOM (colored dissolved organic matter) analysis, water samples from the surface
waters of the SAS lakes were filtered through sample-rinsed 0.47 mm, 0.2 μm pore size cellulose

183 acetate membrane filters and the filtrate was analyzed with the Varian Cary 100 UV-VIS 184 spectrophotometer. The absorbance scans were obtained from the spectrophotometer over the 185 wavelength range of 200-850 nm at natural pH using a 1-cm acid-cleaned quartz cuvette on dual beam mode, at a speed of 240 nm min⁻¹ with a slit width of 1 nm. The absorbance was measured 186 187 against ultra-pure water and corrected against the blanks prepared on site. Absorption coefficients 188 were calculated as $a(\lambda) = 2.303 \times A(\lambda)/L$, where $a(\lambda)$ is the absorption coefficient at a wavelength 189 λ , $A(\lambda)$ is the optical density (absorbance) for wavelength λ , and L is the path length of the 190 cuvette (m). Spectral slope parameters were also determined from absorbance at 275nm (a_{275}) and 295 nm (a_{295}) using the formula $a_{295} = a_{275}e^{-S(295-275)}$, where S (nm⁻¹) is the spectral slope 191 192 parameter (Fichot and Benner, 2012).

193 2.3 Active layer and permafrost core sampling

Triplicate soil samples were retrieved from the surface active layer of a palsa adjacent to lake SAS1B on 3 August 2013; the soil was sampled from half way up the face of the palsa (approximately 1 m above the lake surface) that was eroding and collapsing into the lake by pushing tubes with an internal diameter of 6 cm and a length of 20 cm horizontally into the palsa (Fig. 2). Subsamples were frozen and transported to Québec City for CHN (LECO analyzer, CEN, Québec) and other geochemical analyses (ICP-AES; INRS-ETE, Québec). Enrichment Factors (EF) for the soil samples were determined as per Gorham and Janssens (2005):

$$EF_{soil} = \frac{[x]_{sample} / [Al]_{sample}}{[x]_{soil} / [Al]_{soil}}$$

where $[x]_{sample}$ is the concentration of a given element, x, in the study sample, $[Al]_{sample}$ is the concentration of the element Al in the study sample, $[x]_{soil}$ is the world median concentration of element *x*, and $[Al]_{soil}$ is the world median concentration of the element Al. Global average values were taken from Bowen (1979).

205 Permafrost cores were sampled from two sites 15 m apart on large an organic-rich palsa 206 mound at the SAS2 site (Fig. 1) on 5 August 2013. For permafrost core sampling, a soil pit was 207 dug into the palsa to the base of the active layer. The permafrost cores were then retrieved using a 208 portable drill system (Calmels et al., 2005). The recovered cores were 10-cm in diameter, with 209 lengths of 2.8 and 3.0 m. Samples were returned to the Centre for Northern Studies (Université 210 Laval, Québec) and maintained frozen. The frozen core was characterized by computerized 211 tomography (CT) scanning (Calmels and Allard, 2004) and later subsampled. Subsamples were 212 1.5-cm thick and centered at depths of 1.25, 1.75, and 2.77 m from each of the two permafrost 213 cores. Sample depths were chosen to span the full-length of the permafrost core, and based on 214 CT-scan results. Each sub-sample was placed in a sterilized container filled with ultrapure Milli-215 Q water and thawed for a period of 48-hours. Samples were then lightly shaken and then filtered 216 through 0.47-mm, 0.2-µm pore size cellulose acetate membrane filter. The filtrate and a blank 217 (Milli-Q water) were analysed for DOC, NO₃⁻ (including NO₂⁻) and SRP (INRS-ETE, Québec). DOC was analyzed using a Shimadzu 5000A TOC Analyzer (detection limits of 0.05 mg C L^{-1}). 218 219 SRP and NO₃⁻ were analyzed using a Lachat autoanalyzer (detection limit of 1 μ g N and P L⁻¹). 220 DOC, SRP, and NO_3^{-1} concentrations were normalized to per unit volume of ice in the sample, to 221 determine the potential contributions from melted ice. CDOM was analyzed as described above. 222 Unfiltered diluted samples were analyzed for particle-size distribution as described above. 223 Additional subsamples from throughout the core were analysed for total CHN content (LECO 224 analyzer, CEN).

225 2.4 Size-fractionation

Surface water was sampled from the pelagic zones of lakes BGR1, SEC, KWK12, SAS2A, NAS1A and RBL4K for size fractionation. Phytoplankton and particle-attached bacterial communities were filtered through a 35- μ m Nitex screen. The bacteria were then further separated into two size fractions: either by passing through a glass fibre filter (Pall Canada Ltd.) with a nominal pore size of 3.0 μ m, or by further filtration through a Whatman GF/B glass fibre filter with a nominal pore size of 1.0 μ m. Approximately 2.0 L was filtered for each size fraction, and an unfiltered total community sample was also preserved for concurrent analysis.

233 **2.5 Bacterial abundance**

234 The bacterial communities in the size-fractionated samples were counted after staining with 4',6-235 diamidino-2-phenylindole (DAPI). Samples were preserved with formaldehyde (3.7% final 236 concentration) in pre-washed plastic bottles, pre-rinsed with the sample and stored at 4°C for up to 6 months. Samples were later shaken to break up larger particles and then filtered onto 237 238 Nuclepore black polycarbonate membranes (0.22 µm, 25 mm) placed on cellulose acetate backing filters (0.7 µm, 25 mm) under low pressure. DAPI was added to 1.5 mL of sample (4 µg 239 mL⁻¹ final concentration) and left to stain for 10 min before filtering until dry. Filters were 240 241 mounted on slides with coverslips and non-fluorescent immersion oil, and stored at -20°C until 242 counting on a Zeiss Axioskip 2 epifluorescence microscope, under UV light and 1000× 243 magnification with immersion oil. A minimum of 15 fields or 400 cells were counted of each 244 sample triplicate.

All other bacterial samples were counted via flow cytometer (FACSCalibur, BD Biosciences Inc., USA) following the procedure described in Comte et al. (2016). Samples were

fixed with glutaraldehyde (1% final concentration) and stored at -80°C until analysis. Bacterial 247 248 samples were then shaken and a 500 µL aliquot was stained with the nucleic acid dye 50X SYBR Green I (20 uL mL⁻¹ final concentration: Life Technologies, Thermo Fisher Scientific, Waltham 249 250 MA, USA) for 10 min in the dark, and processed at the lowest flow rate (12 μ L min⁻¹) using 1 μ m 251 yellow-green microspheres (Polysciences Inc., Warrington PA, USA) solution as an internal 252 standard. Bacterial cells were enumerated based on fluorescence (FL1) and side-scatter 253 characteristics while excited at 488 nm, with data analyzed using the CellQuest Pro software with 254 manual gating.

255 2.6 Bacterial production

256 Bacterial production (BP) rates were measured via protein synthesis using radio-labelled leucine 257 incorporation (Kirchman, 2001). Triplicate microfuge tubes plus a killed control each containing 1.5 mL of sample and 40 nM of [4,5-³H]leucine (60 Ci mmol⁻¹, PerkinElmer Inc.) were incubated 258 259 for 1 h at 15°C. After one hour, biomass production was halted by the addition of 100 µL of 260 trichloroacetic acid (100%) and stored at 4°C. Later, the samples were returned to the Takuvik 261 radio-isotope laboratory at Université Laval. Samples were processed using the centrifugation 262 method of Smith and Azam (1992) and then radio-assayed in a scintillation counter to measure the ³H-leucine uptake. Net bacterial C production was estimated using a conversion factor of 3.1 263 264 kg C per mole of leucine (Kirchman et al., 1993; Iriberri et al., 1990). Bacterial community 265 specific growth rates (BG, d^{-1}) were determined (as described in Kirchman 2001) via BG = BP/B, where BP are bacterial production rates ($\mu g C L^{-1} h^{-1}$), and B is the bacterial biomass (fg C mL⁻¹) 266 that was converted from bacterial cellular counts (cells mL⁻¹) to units of carbon using an average 267 cellular carbon content of 18 fg C cell⁻¹, as measured in the nearby KWK lakes (Roiha et al., 268 269 2015).

270 2.7 Enrichment experiment

271 At lake SAS1B, the following treatments were added to triplicate 20 L containers of lake water: 43 µmol L⁻¹ of glucose (G; to achieve an estimated 20% increase of ambient DOC based on 272 previous measurements), 1.8 µg L⁻¹ of phosphorus (P; 10% enrichment of total P), glucose and 273 phosphorus together (G+P), and 6.2 mg L^{-1} of an aqueous suspension of the soil from the SAS1B 274 275 palsa face, sampled and analyzed as described above (S; equivalent to 20% increase in total 276 carbon relative to ambient DOC). A sample control was also prepared (C). Samples were taken 277 immediately after enrichment (t_0) , and from each container after 5 days (t_5) of incubation in situ 278 in the near-surface waters of SAS1B. Chl a, BA and BP rates were measured in each treatment.

279 2.8 Statistical analysis

280 Statistical differences were examined between the SAS peatland lakes and the other study lakes. 281 Environmental and bacterial variables for the comparison of lake waters were analysed by 1-way ANOVA for three lake groups: the SAS lakes, the other thermokarst lakes, and the non-282 283 thermokarst rock-basin lakes. A 2-way ANOVA was used to compare environmental and 284 biological differences between surface and bottom waters of the lake groups mentioned above. 285 For the size-fractionation data, a 2-way ANOVA was performed on pre-standardized variables, 286 with the factors of lake and size-fraction following the Shapiro normality test. The enrichment 287 experiment was analyzed via 2-factor repeated measures ANOVA, with each replicate 288 individually tracked. Variables before and after enrichment were also compared via Holm-Sidak 289 pairwise comparisons. BG were normalized by square route transformation, and BA was log 290 transformed. Analyses were done using GraphPad Prism (v. 6.0f, GraphPad Software Inc., USA) 291 and SigmaPlot (v. 12, Systat Software Inc., USA).

292 3 Results

293 **3.1** Active layer soil analyses

The SAS active layer extended to 60 cm depth and contained 509 g kg⁻¹ of carbon, or 50.9% of 294 the soil by mass, and 30 g kg⁻¹ of nitrogen (Table 1). Several elements were present in substantial 295 concentrations, including phosphorus at a mean value (\pm SD) of 545 (19) mg kg⁻¹, iron at 6.0 (1.6) 296 g kg⁻¹, and calcium at 5.2 (3.4) mg kg⁻¹. The soil enrichment factors (EF) were used to compare 297 298 concentrations of soil elements with world medians (Bowen, 1979) to determine whether the 299 elements were derived uniquely from wind-blown mineral soil (EF ≤ 1), or were more likely to 300 accumulate due to multiple sources (EF >>1) (Gorham and Janssens, 2005). Several elements had 301 EF values <1, including K, Na, and Mg. For Al, Mn, Pb, Cr, Zn, and Ba, the EF values were 302 between 1 and 2. Phosphorus had an EF of 16, and S had the highest EF value of 132.

303 **3.2** Permafrost core analysis and comparison with lake water

The frozen permafrost cores were composed mainly of organic matter intercalated with ice (>80%) in the two upper levels, with high concentrations of sediment only in the deepest level at ~ 2.77 m (Table 2). The soils throughout the core released water containing nitrate and high concentrations of DOC, and the sediment-containing matter near the bottom of the cores additionally released SRP at concentrations up to 30 times those measured in the SAS lakes.

The $a_{CDOM}(\lambda)$ spectra for the DOC from the upper two permafrost core samples were almost identical, but differed as much as 185 m⁻¹ from those for the bottom-core samples (Fig. S1). Mean values (±SD) of $a_{CDOM}(440)$ ranged from 9.96±0.28 to 16.2±0.18 m⁻¹ in all permafrost core samples except the bottom of core A, which was 51.5±0.4 m⁻¹. The spectral slope parameter $(S_{275-295})$ ranged from 0.0087 to 0.014 nm⁻¹ in the permafrost cores (mean for all values of 0.0012 ± 0.0018). For SAS lake water samples, $a_{CDOM}(\lambda)$ spectra were similar to those observed from the permafrost core samples (Fig. S1); $a_{CDOM}(440)$ ranged from 8.63 ± 0.051 to 18.1 ± 0.068 m^{-1} , and was consistently higher in bottom waters than in surface. Spectral slope parameters (S₂₇₅. 295) for the SAS water were within a notably smaller range than for the permafrost cores, and ranged from 0.011 to 0.012 nm⁻¹ (mean for all values of 0.0115\pm0.0004).

319 **3.3** Particle size distribution in soils and lakes

320 The particle-size distributions from the SAS permafrost cores followed a bimodal distribution as 321 observed in the SAS active layer soil and the SAS lake water (Fig. 4). The maximum frequency 322 of particle size (PSmax) ranged from 152 to 200 µm for the permafrost layers at 1.25 and 1.75 m 323 for both duplicates. The bottom layer of core A had a distribution maximum of 67 µm while the 324 maximum for core B was 51 µm. The active layer soil was similar to the distribution of particles 325 in the upper permafrost cores, but with a maximum frequency at a slightly higher size (229 μ m), 326 and an even distribution of particles from 80 to 300 µm. The particle spectrum for the SAS lake 327 water overlapped with those for the soil, but the surface water spectrum extended to higher sizes 328 and peaked at 394 µm, while the bottom water spectrum peaked at 200 µm and extended to finer 329 particles (Fig 4).

As a further analysis of particle size distribution, difference spectra were calculated relative to the SAS2A surface waters for the active layer soil, upper permafrost soil and bottom waters. These curves emphasize the strong overlap in size spectra, with differences ranging from only -3 to +6%, depending on size class (Fig. 5). These curves also illustrate how the surface water was slightly enriched in larger particles that ranged from 229 to 2010 μ m in diameter, while the bottom waters were enriched in finer particles that ranged from 39 to 200 μ m.

Within-lake particle sizes showed highly variable distributions across the other lakes studied, with modes that ranged from 4 to 1337 μ m (Table 3; Fig. 4). The smallest particles were identified in RBL4K, a rock-basin lake unaffected by permafrost, with a mode particle size of 4 μ m. Of the thaw lakes, the lakes from areas of sporadic, degrading permafrost had larger particles than the continuous- and discontinuous-permafrost lakes. For example, the mode particle sizes in NAS1A, BGR1, and SEC were 102, 59, and 153 μ m, respectively. In contrast, the particle-size modes in SAS2A and KWK12 were 300 μ m and 592 μ m, respectively.

The concordance of particle distributions between the lake waters and their adjacent soils varied among the other sample sites (Fig. 4). For example, the distribution of particles in KWK12 was strikingly similar to that of KWK active layer soil, while other lake samples tended to have modes than were higher than the mode observed in the soil, as in SAS. For example, the mode for BGR lake water was 59 μ m, but 10 μ m for the adjacent mineral soil. As at SAS, the distribution of soil and lake water particles at SEC was bimodal, and the particles encompassed a much wider range in soil than in the lakes.

350 **3.4 Lake properties**

351 The SAS lake surface waters differed significantly in their limnological properties from those of 352 other thaw lakes and also from the rock-basin lakes (Table 3; 1-way ANOVA). The SAS lakes had a mean (\pm SD) surface DOC value of 11.5 \pm 2.4 mg L⁻¹, that was significantly higher than the 353 two other lake categories (p < 0.001), with an average of 4.08 ± 1.4 mg L⁻¹ for the other 354 thermokarst lakes, and 10.6 \pm 3.1 mg L⁻¹ for the rock-basin lakes. The average surface Chl a 355 concentration was also significantly higher for the SAS lakes (mean \pm SD = 2.29 \pm 1.09 µg L⁻¹) 356 relative to the other categories (other thermokarst lakes mean = 0.897 ± 0.43 µg L⁻¹; rock-basin 357 lakes mean = $1.39\pm0.50 \ \mu g \ L^{-1}$; p = 0.03), as was surface TN, which averaged (±SD) 0.58 ± 0.10 358

mg L⁻¹ in the SAS lakes, and 0.30 ± 0.08 and 0.61 ± 0.20 mg L⁻¹ in each of the other two categories, respectively (p < 0.01). TSS and maximum particle size (PSmax) values were variable within each lake type (Table 3), with no significant differences; the surface PSmax values were strikingly variable in rock-basin lakes, ranging over three orders of magnitude, but much more homogeneous among the SAS waters (mean of 384 µm with a CV of 37%).

364 Despite their shallow depths, the SAS lakes showed substantial differences between the 365 top and bottom of the water column, which was also a feature of the other thaw lakes (Table 3). A 366 two-way ANOVA for depth and the two thaw lake types (SAS versus the other thaw lakes) showed that overall there were significant increases at depth in SRP (F = 23; p = 0.03); TN (F =367 18; p = 0.02) and Chl a (F = 28; p = 0.03), with no significant (p > 0.05) top-bottom differences 368 369 in DOC, TSS or PSmax. Consistent with the 1-way analysis for all three lake types, this 2-way 370 analysis (both depths) showed that SAS lakes had significantly higher concentrations of DOC (F = 65; p < 0.0001) and TN (F = 32; p = 0.004) relative to other thaw lakes. However, the SAS 371 lakes had significantly lower concentrations of TSS (F = 23; p = 0.05), which were especially 372 373 high in NAS1A and BGR2, and there were no significant differences (p > 0.05) between the two 374 groups of thaw lakes in terms of SRP, Chl a or PSmax.

Ammonium and nitrate were not analyzed routinely, but the surface waters of the SAS lakes sampled during late June 2014 contained undetectable concentrations of nitrate (<10 μ g N L⁻¹) and substantial concentrations of ammonium: 64 (SAS1A), 58 (SAS1B), 49 (SAS2A), 60 (SAS2B) and 74 (SAS2C) μ g NH₄⁺-N L⁻¹. Much higher values were recorded in the anoxic bottom waters of two of the lakes: 1021 (SAS1A) and 178 (SAS1B) μ g NH₄⁺-N L⁻¹, but values were similar to surface waters in SAS2A (48 μ g NH₄⁺-N L⁻¹).

381 3.5 Bacterial abundance and productivity

382 BA in the surface waters varied over an order of magnitude among SAS lakes, from 0.2 to $3.1 \times$ 10^6 cells mL⁻¹, and did not significantly differ from the other lake types (Table 4; p >0.05). BP 383 rates varied by a factor of six among SAS lakes, from 0.25 to 1.68 µg C L⁻¹ h⁻¹, and as for cell 384 385 concentration, there were no significant differences among the categories of lakes. BG were similarly variable, ranging from 1.9 to 113 fg cell⁻¹ d⁻¹. Four of the five values from SAS surface 386 387 waters were well above (factor of two or greater) any of the thermokarst or rock-basin lakes of 388 the two other categories. Overall, the mean value for SAS waters was significantly higher than 389 the others (F = 5.0; p = 0.026).

390 **3.6** Size-fractionated solids and associated bacteria

391 In SAS2A, the mass of suspended solids decreased sharply with filter size, and the $<1 \mu m$ 392 fraction contained only 5% of the total value (Fig. 6). Similar trends were observed in the other 393 lakes, with the exception of NAS1A where almost all of the suspended solids passed through the 394 35 and 3 µm filters, and 34% even passed through the 1 µm filter. In terms of bacterial cells, 395 highest counts were in the total or $<35 \,\mu m$ fraction, with the exception of NAS1A, which peaked 396 in the $<3 \mu m$ fraction. Only 28% of the maximum counts occurred in the $<1 \mu m$ fraction in 397 SAS2A samples, and this fraction was similarly low in SEC (23%). Higher percentages of 398 maximum occurred in the <1 µm fraction in RBL4K (43%), KWK12 (52%), BGR1 (52%) and 399 NAS1A (87%).

400 The BP data showed a more striking pattern (Fig. 6). Total production was maximal in the 401 unfiltered (total) or $<35 \ \mu m$ fraction, but the $<1 \ \mu m$ fraction accounted for only 26% of the 402 maximum in SAS2A, and was even lower in most of the other lakes: 11% (NAS1A), 11%

403 (RBL4K), 16% (KWK12), and 14% (BGR1). An average value of 39% was recorded in SEC. 404 BG (bacterial community specific growth rates) were highly variable among size fractions, ranging from a minimum of 0.021 ± 0.0035 d⁻¹ in the <1 μ m size fraction of BGR1 to a maximum 405 of 1.1 ± 0.032 d⁻¹ in the <3 µm fraction of SEC (Fig. 6). In the SAS2A analysis, rates were two 406 407 times higher in the $<3 \mu m$ than in the $<1 \mu m$ fraction. Similarly, in all of the other lake waters, 408 highest BG were in fractions above 1 µm: Total (NAS1A); Total and <3 µm (BGR1); <3 µm 409 (SEC); Total (KWK12) and Total and $<35 \mu m$ (RBL4K). The free-living cell fraction ($<1 \mu m$) 410 showed rates that were a small percentage of the production per cell in the unfiltered samples in 411 RBL4K (27%); KWK12 (16%); NAS1A (8%) and BGR1 (16%).

412 **3.7 Enrichment experiment**

413 Over the course of the experiment (Fig. 7), there was no significant change in Chl aconcentrations, either with time or among treatments (repeated measures ANOVA, p > 0.05). 414 415 There was a significant drop in bacterial cell abundance in all treatments by day 5 (BA; effect of time F = 39; p = 0.025) but no significant change in production rates (BP, p > 0.05). Although 416 417 there was no significant overall effect of time and treatment on bacterial growth rates (BG, 418 carbon production rates per cell; p > 0.05), there was a significant interaction effect between time and treatment (F = 4.9; p = 0.026). Further analysis (Holm-Sidak pairwise comparisons) showed 419 420 that while most of BG had not changed statistically, there was a large significant increase 421 between days 1 and 5 in the soil addition treatment (146% average increase; t = 2.9; p = 0.034) 422 and in the carbon+phosphorus (G+P) enrichment (187% increase; t = 3.34; p = 0.022).

423 4 Discussion

424 The permafrost peatland soils analyzed here contained abundant resources for bacterial 425 development, including biologically-accessible carbon, nitrogen, and phosphorus. Degradation of 426 the SAS permafrost peatland landscape was observed to be already underway, with the transfer of 427 material from permafrost palsa mounds and surface soils into thermokarst lakes. This transfer 428 was evident at the landscape level, from patterns of palsa size and degradation (Fig. 2), as well as 429 via within-lake evidence of similarities between particles, and at the microbial scale, via the 430 abundant particle-attached bacterial communities that likely benefit from access to particles and 431 flocculent matter. The favorable conditions available to bacterial communities in permafrost 432 peatland thaw lakes resulted in abundant bacterial communities with high BG. These bacterial 433 communities seemed less likely to be controlled by bottom-up processes, as is commonly 434 observed elsewhere (Pace and Cole 1994), than by top-down regulation.

435

4.1 Organic carbon enrichment

There is an estimated 1.048×10^6 km² of permafrost peatland in North America, half of which is 436 437 located in areas of sporadic or isolated permafrost like the SAS region of the present study 438 (Tarnocai et al., 2009). The surface soil and permafrost cores from the SAS region were abundant 439 in carbon, with ~50% C-content (by dry-weight) down to 2 m depth in the permafrost cores (Fig. 440 3), comparable to peatland soils elsewhere. A study of peat cores from five North-American sites 441 showed that the concentration was consistently around 50% C dry-weight (Gorham and Janssens, 442 2005), as was also observed in the surface SAS1 active layer and in the upper layers of the two 443 permafrost cores. Similarly, a synthesis of peat data from 9 studies in the northern hemisphere 444 (Gorham et al., 2012) yielded a mean value (±SD) of 50.1 (2.9)%. These carbon values are much higher than observed in other types of Arctic soils. For example, the carbon content of active 445

layer soil from a polygonal tundra of the Lena Delta ranged from 1.5 to 3% dry-weight (Höfle et
al., 2013), an order-of-magnitude less than values in the SAS soil. Lower carbon concentrations
were also measured in a study of permafrost soils of the Arctic tundra of Northwest Territories,
which were 6.5% C (Moquin and Wrona, 2015), and in the lithalsa mounds surrounding BGR1
(Fig. 1), where the carbon content of the active layer was less than 1% (ADAPT, 2014).

451 The degrading palsas provided a rich potential source of organic carbon to their associated 452 lakes (Fig. 2). Paleoecological studies (Arlen-Pouliot and Bhiry, 2005; Fillion et al., 2014) have 453 shown that the peat in the SAS valley accumulated in bogs over a 1400 year period from around 454 1800 to 400 years before the present, and was then uplifted in palsas during the period of 455 permafrost aggradation during the Little Ice Age, 400 to 200 years before present day. From the end of the 19th century onwards, the SAS palsas have been thawing and eroding, with the 456 457 production and eventual infilling of associated thermokarst lakes (palsa lakes), and this warming 458 phase appears to have accelerated over the last two decades (Bhiry et al., 2011). As shown in the 459 permafrost core analysis (Table 2), the potential organic input to thermokarst lakes includes not 460 only particulate organic carbon but also DOC. Two contrasting scenarios have been identified for 461 the effects of thawing permafrost on carbon transport in northern catchments (Vonk et al., 2015): 462 (1) decreased net export of carbon as a result of increased flow path lengths, residence times and 463 decomposition in a deeper active layer, or (2) increased net export as a result of thermal erosion 464 and mobilisation of particulate and dissolved organic carbon. The SAS valley is currently in the 465 'pulse phase' of carbon mobilisation, which as noted by Vonk et al. (2015), is typically 466 dominated by the export of particulate materials. However, the high DOC of the SAS lake waters, 467 significantly above that in the other lake types, also attests to a substantial export of DOC from 468 the SAS palsas to their associated lakes, and is consistent with the permafrost thaw analyses. The

low spectral slope values in the SAS lakes indicate a predominantly terrestrial origin of DOC in
these lakes (Fichot and Benner, 2012), and the similarity in slope values between the permafrost
cores and the lake waters also supports this close linkage between land and water (Fig. S1).

472 **4.2** Nutrient enrichment

473 In addition to carbon-enrichment, the SAS soil also likely represents a nutrient-rich substrate for 474 bacterial metabolic activity (Table 1). The soil N concentrations were low relative to C, reflecting 475 the stoichiometry of ancient moss tissue, however there were large standing stocks of N, and the 476 high ammonium levels in the lake water implies that nitrogen was in abundant supply for 477 planktonic microbial growth processes. The SAS surface soil had nearly three times as much 478 phosphorus and more than five times as much iron relative to concentrations found in other 479 Sphagnum peat bogs (Gorham and Janssens, 2005). It also had 1.6 times the concentration of 480 phosphorus and three times the amount of calcium when compared to permafrost soil from the 481 Northwest Territories (Moquin and Wrona, 2015), though only a fraction (5%) of the magnesium 482 observed in the latter study.

483 Part of the observed differences when compared to these other regions is related to the 484 long-term history of the Hudson Bay region. Enrichment Factors (EF) indicated that phosphorus 485 was above the global average for phosphorus concentrations in soil (EF=15.9), demonstrating 486 long-term transfer of this elements from other sources, such as via atmospheric deposition or 487 hydrological transfer (from surrounding rocks or terrestrial zones). The permafrost thawing 488 analyses showed a release of nitrate but no evidence of soluble phosphorus release, except from 489 the bottom-core samples that produced SRP concentrations that were 2- to 30-times above current 490 lake water values (Table 2). This coastal region of eastern Hudson Bay was submerged beneath 491 the Tyrell Sea from immediately after the retreat of the ice cap about 7900 years ago to emersion

from the sea about 6000 years before the present, and previous paleoecological studies on the deeper strata of the SAS valley sampled in the bottom core show that these are ancient marine sediments (Arlen-Pouliot and Bhiry, 2005; Fillion et al., 2014). This would account for their higher phosphorus release relative to that from the upper core. Deep thawing of the palsas during the final stages of their collapse could therefore have a major P-enrichment effect on the associated thermokarst lake ecosystems.

498 Inorganic nutrient release from thawing and eroding permafrost has been documented at 499 sites elsewhere, but with large variations among regions and landscape types (Vonk et al., 2015). 500 Slumping of permafrost soils into Alaskan streams increases their inorganic as well as organic 501 concentrations (Bowden et al., 2008), but phosphorus adsorption onto clays may decrease SRP 502 concentrations in some mineral permafrost environments (Breton et al., 2009). Nitrate export 503 from the Kuparuk River catchment on the Alaska North Slope increased 5-fold between the late 504 1970s and early 2000s and may have been related to permafrost soil and vegetation effects 505 (McClelland et al., 2007). These processes may also operate in the peatland permafrost sites 506 studied here.

507 **4.3 Particle enrichment**

The SAS permafrost peatland lakes were rich in particulate material, with substantially larger particles than non-permafrost-affected rock-basin lake RBL4K, and they contained some of the largest particles amongst thermokarst lakes studied (Fig. 4). Much finer particles over a narrower size range were recorded in the northern permafrost lakes NAS1A and BGR1, reflecting their mineral permafrost catchments. The close correspondence between the active layer, permafrost and lake particle size distribution at SAS2 (Fig. 5), is evidence that this richness of particles in the peatland lakes is derived from the thermo-erosion of their associated palsas. The small but 515 conspicuous shift to larger particles in the SAS surface waters may indicate the production of 516 aggregates by flocculation that are maintained in the upper water column by virtue of their low 517 density and therefore slow sinking speeds. The production of flocs was observed when SAS 518 water was brought into the laboratory and stored in clear glass bottles without mixing for several 519 days (B. N. Deshpande, unpublished observations), and may be favored by the high DOC 520 concentration in these waters. In contrast, the deeper waters showed an increase in fine particles, 521 contrary to what might be expected from Stokes' Law based upon size relationships. The origin 522 of this difference is unknown at this time, but it could be the result of fine, high-density mineral 523 particles settling out from the overlying water column. The extreme thermal stratification and 524 physical stability of these waters (Deshpande et al., 2015; Przytulska et al., 2016) indicates that 525 only particles of small size or low density would be maintained in the water column.

526 4.4

.4 Bacterial association with particles

527 There was clear evidence that the high particle-loading of peatland SAS waters affected its 528 bacterial ecology. Less than 30% of the total BA and BP rates were associated with the free-529 living category, operationally defined here as passing through a 1 μ m filter, and this effect was 530 even more pronounced for bacterial growth rates (BG), which were 50% lower in the $<1 \mu m$ than 531 in the $<3 \mu m$ fraction, implying much faster growth rates of bacteria on particles in the size range 532 1-3 µm. In SAS2A, as in many of the other northern lakes, bacterial populations and production 533 appeared to be distributed across the full range of particle sizes (Fig. 6). This is consistent with 534 studies across a range of salt- and fresh-water environments showing that particle-attached 535 bacteria are favored in environments with a high availability of particulate habitats (Lami et al., 536 2009; Kirchman and Ducklow, 1987; Kirchman and Mitchell, 1982).

537 There are several sources of experimental error that must be considered when interpreting 538 particle analyses of this sort. Firstly, it is difficult to obtain a reliable count of particle-based 539 bacteria. Our shaking of the samples would have broken up and released cells from larger 540 aggregates, but dislodging bacteria from particles is notoriously difficult. Even sonication is not 541 completely effective, and addition of a detergent along with sonication can result in a pronounced 542 increase in cell concentrations in sediment-laden waters (Yoon and Rosson, 1990). It is therefore 543 likely that the cell concentrations in the present study were underestimated, especially in the >1544 μm fractions. This would imply an even greater representation by particle-based cells to the total 545 community abundance, but it would also have led to an overestimation of the production rates per 546 cell biomass. Experiments based on size-fractionation by filtering are always challenging because 547 of the interactions between particle and the filters. Filtering water results in the breakup of 548 particles upon collision with the filter surface, causing two important effects: (1) an increased 549 bioavailability of organic particles (Amon and Brenner, 1996); and (2) potential detachment of 550 particle-attached bacterial from their organic-matter habitat. Filter blocking may have also 551 retained smaller fractions, including free-living cells. Although the exact rates per fraction in the 552 present study, as in all such studies, must be interpreted with caution, the distribution of cells and 553 activities across multiple fractions and the low values in the <1 µm fraction point to the 554 importance of particle-based bacteria in this system. This was not unique to the organic-rich SAS 555 waters, but was found in several of the other lakes. These results are consistent with a previous 556 study of five lakes in the mineral-rich KWK area, which showed a substantially higher proportion 557 of particle-attached bacteria ($62\pm30\%$ in the >3 µm size-fraction) and high BP ($59\pm30\%$) by 558 these groups (Roiha et al., 2015). The BG patterns are also consistent with data elsewhere; for 559 example, Kirchman and Mitchell (1982) found that particle-attached bacteria accounted for <10% 560 of the total bacterial populations but contributed substantially to total microbial activity.

561 **4.5** Controls on bacterial abundance and productivity

Bacterial abundance in the thermokarst lakes sampled here ranged from 0.20 to 3.5×10^6 cells mL⁻¹, which falls towards the higher end of the range for comparable aquatic ecosystems (Table 5). For example, in study of 16 lakes located in subarctic Sweden, bacterial concentrations ranged from 0.4 to 2.4×10^6 cells mL⁻¹ (Karlsson et al., 2001), and in a humic, boreal lake, bacterial concentrations ranged from 1.2 to 1.7×10^6 cells mL⁻¹ in winter (Tulonen et al., 1994).

The BP rates in the thermokarst lakes sampled here were also at the high end of the range in comparable systems (Table 5), from 0.25 to 2.3 μ g C L⁻¹ h⁻¹, with an average (±SD) of 0.92 (±0.46) μ g C L⁻¹ h⁻¹. These rates are more comparable with those observed in southern Québec aquatic ecosystems, and productive systems elsewhere, for example the high-nutrient Butron River, Vizcaya, Spain (Table 4). In this river, free-living bacteria had a production rate of 5.8 μ g C L⁻¹ h⁻¹, with particle-attached communities contributing 2.97 μ g C L⁻¹ h⁻¹ of bacterialproduction (Iriberri et al., 1990).

574 Such comparisons with other sites need to be made with caution because they rest on a 575 number of untested assumptions. Notably, we used a single factor for conversion between leucine 576 uptake to carbon production concentrations, and assumed a 2-fold isotopic dilution of leucine in the natural environment, observed in certain subarctic ecosystems (Simon and Kirchman, 1988; 577 578 Simon and Azam, 1989) and as has been employed for a wide range of bacterial production 579 studies elsewhere (Judd et al., 2006; del Giorgio et al., 1997; del Giorgio and Bouvier, 2002). 580 However, given the wide-range of DOC concentrations and unique characteristics of thermokarst 581 lakes, it is likely that the true conversion factors would lie on a wider range, as found in Pulido-582 Villena and Reche (2003). If ambient leucine concentrations follow the trend of measured total 583 DOC concentrations, then production rates may have been especially underestimated in the SAS incubations. This should not have influenced the pattern amongst size fractions, however, as noted above, errors associated with cell counts for the particle-associated fractions may have affected the estimates of rates per biomass.

587 The enrichment experiment in the present study showed that there was no significant 588 effect of either glucose or phosphorus when added separately to the SAS peatland lake waters 589 (Fig. 7). These results imply that bacterial stocks and production were not strongly limited by 590 carbon supply, nor were they subject to primary phosphorus limitation. This contrasts with results 591 from the permafrost-influenced, turbid waters of the Mackenzie River, Northwest Territories 592 Canada, where short-term glucose enrichment resulted in a 4-fold increase in leucine uptake rates 593 (Vallières et al., 2008). They also contrast with enzymatic analyses of Greenland lakes that 594 implied strong primary phosphorus limitation (Burpee et al. 2016). Chl a levels also showed no 595 significant effect of enrichment, in contrast to such experiments in High Arctic lakes (Bonilla et 596 al., 2005). However, for the addition of glucose plus phosphorus, and also for the soil addition 597 treatment, there was a significant increase in BG (Fig. 7). This implies a combined stimulatory 598 enrichment effect on bacterial production of the palsa permafrost soils eroding into the SAS 599 thermokarst lakes.

Other studies have shown that DOM-enrichment increased heterotrophic metabolism, sometimes at the expense of autotrophic metabolic activity (Forsström et al., 2013). In an Arctic tundra lake of the Northwest Territories, addition of permafrost soil resulted in decreased bacterial production in the pelagic zone, but increased rates in the benthic zone (Moquin and Wrona, 2015). A recent study of Arctic lakes in southwest Greenland found that microbial populations were primarily phosphorus-limited, and that this P-limitation increased with the availability of DOM (Burpee et al., 2016). In an area of discontinuous permafrost in northern

Sweden, addition of terrestrial carbon to aquatic bacterial samples resulted in a four- to sevenfold increase in respiration rates, with only a small proportion of the available carbon (1 - 12%)contributing to cellular biomass and the dominant portion respired to CO₂ (Roehm et al., 2009).

610 All of the limnological and microbial measurements point to the SAS peatland lakes as 611 favorable environments for BP. The BG values show that SAS lakes were outliers amongst 612 thermokarst lakes, with significantly elevated rates (Table 4). These rates are also on the higher 613 end of observations made across a wide trophic range of lakes from other regions (del Giorgio et 614 al., 1997; Smith and Prairie, 2004). The lack of response of cellular rates to either phosphorus or 615 carbon added singly also implies sufficiency, and even although there was a significant rise in 616 growth in response to soil and carbon+phosphorus (G+P), this was not translated into an increase 617 in BA (Fig. 7). In fact populations fell in all of the incubations, despite BG rates that remained 618 the same or higher. In combination, these observations suggest that bacterial biomass in the SAS 619 peatland lakes is controlled by top down processes rather than by bottom-up, substrate supply 620 rates. This might involve viral lysis, as in planktonic bacterial systems elsewhere (e.g., Lymer et 621 al., 2008) or it may be the result of intense grazing by bactivores. These waters contain high 622 concentrations of zooplankton, including Daphnia (Bégin 2015) that can filter picocyanobacteria 623 (Przytulska et al., 2015) and probably heterotrophic bacteria (Rautio and Vincent, 2006). Studies 624 on SAS2A have shown that they contain abundant ciliates, heterotrophic nanoflagellates and 625 mixotrophic phytoplankton (Przytulska et al., 2016), all of which may exert a control on bacterial 626 densities. In these environments, enriched by soil particles, carbon and nutrients released from 627 thawing and eroding permafrost, bacterial stocks may be ultimately capped by loss rather than 628 gain processes.

630 5 Acknowledgements

631 This study was made possible with financial support from the Natural Sciences and Engineering 632 Research Council of Canada (NSERC), the Canada Research Chair program, and the Québec 633 nature and technology research funds (FRQNT). Financial and logistical support are further 634 provided by the Centre for Northern Studies (Université Laval), the ADAPT (Arctic 635 Development and Adaptation to Permafrost in Transition) Discovery Frontiers grant, and the 636 Northern Science Training Program. We are grateful to P. Bégin, J. Comte, A. Przytulska-637 Bartosiewicz, V. Mohit, and C. Tremblay for their help in the field, and M.-J. Martineau, and J. 638 Comte for laboratory support. We thank I. Laurion (INRS-ETE) for flow cytometry and F. 639 Bruyant (Takuvik, Université Laval) for the radioactivity laboratory. We thank D. Sarrazin and S. 640 Aebischer for permafrost core sampling, and the CEN-ADAPT team for core and CHN analysis: 641 G. Labrecque, A.-S. Carbonneau, J. Roger and M. Lemay. We thank S. Prémont at the INRS-642 ETE laboratory and D. Cayer at Université Laval FFGG laboratory for support and guidance with 643 sample analysis.

645 6 References

- Acinas, S. G., Antón, J., and Rodríguez-Valera, F.: Diversity of free-living and attached bacteria
 in offshore western Mediterranean waters as depicted by analysis of genes encoding 16S
 rRNA, Appl. Environ. Microbiol., 65, 514-522, 1999.
- ADAPT: Carbon, nitrogen and water content of the active layer from sites across the Canadian
 Arctic, v. 1.0, Nordicana D21, doi:10.5885/45327AD-5245D08606AB4F52, 2014.
- Alldredge, A. L., and Silver, M. W.: Characteristics, dynamics and significance of marine snow,
 Progr. Oceanogr., 20, 41-82, 1988.
- Amon, R. M. W., and Benner, R.: Bacterial utilization of different size classes of dissolved
 organic matter, Limnol. Oceanogr., 41, 41-51, 1996.
- Arlen-Pouliot, Y., and Bhiry, N.: Palaeoecology of a palsa and a filled thermokarst pond in a
 permafrost peatland, subarctic Québec, Canada, Holocene, 15, 408-419,
 doi:10.1191/0959683605hl818rp, 2005.
- Bégin, P.-N.: Abondance et diversité des rotifères dans les mares de thermokarst subartiques,
 M.Sc., Départment de biologie, Université Laval, Canada, 2015.
- Bhiry, N., Delwaide, A., Allard, M., Bégin, Y., Filion, L., Lavoie, M., Nozais, C., Payette, S.,
 Pienitz, R., Saulnier-Talbot, É., and Vincent, W. F.: Environmental change in the Great
 Whale River region, Hudson Bay: Five decades of multidisciplinary research by Centre
 d'études nordiques (CEN), Ecoscience, 18, 182-203, doi:10.2980/18-3-3469, 2011.
- Bonilla, S., Villeneuve, V. and Vincent W. F.: Benthic and planktonic algal communities in a
 high arctic lake: Pigment structure and contrasting responses to nutrient enrichment, J.
 Phycol., 41, 1120-1130, doi:10.1111/j.1529-8817.2005.00154.x, 2005.
- Bowden, W. B., Gooseff, M. N., Balser, A., Green, A., Peterson, B. J., and Bradford, J.:
 Sediment and nutrient delivery from thermokarst features in the foothills of the North
 Slope, Alaska: Potential impacts on headwater stream ecosystems, J. Geophys. Res.,113,
 G02026, doi:10.1029/2007JG000470, 2008.
- Bowen, H. J. M.: Environmental Chemistry of the Elements, Academic Press, London, UK,
 1979.
- Breton, J., Prairie, Y., Vallières, C., and Laurion, I.: Limnological properties of permafrost thaw
 ponds in northeastern Canada, Can. J. Fish. Aquat. Sci., 66, 1635-1648, doi:10.1139/f09108, 2009.
- Burpee, B., Saros, J. E., Northington, R. M., and Simon, K. S.: Microbial nutrient limitation in
 Arctic lakes in a permafrost landscape of southwest Greenland, Biogeosciences, 13, 365374, doi:10.5194/bg-13-365-2016, 2016.
- Calmels, F., and Allard, M.: Ice segregation and gas distribution in permafrost using
 tomodensitometric analysis, Permafrost Periglac., 15, 367-378, doi:10.1002/ppp.508,
 2004.
- Calmels, F., Gagnon, O., and Allard, M.: A portable earth-drill system for permafrost studies,
 Permafrost Periglac., 16, 311-315, doi:10.1002/ppp.529, 2005.

- Comte, J., Lovejoy, C., Crevecoeur, S., and Vincent, W.F.: Co-occurrence patterns in aquatic
 bacterial communities across changing permafrost landscapes, Biogeosciences, 13, 175 doi:10.5194/bg-13-175-2016, 2016.
- 687 Crevecoeur, S., Vincent, W. F., Comte, J., and Lovejoy, C.: Bacterial community structure across
 688 environmental gradients in permafrost thaw ponds: methanotroph-rich ecosystems, Front.
 689 Microbiol., 6, 192, doi:10.3389/fmicb.2015.00192, 2015.
- 690 Crump, B. C., Armbrust, E. V., and Baross, J. A.: Phylogenetic analysis of particle-attached and
 691 free-living bacterial communities in the Columbia River, its estuary, and the adjacent
 692 coastal ocean, Appl. Environ. Microbiol., 65, 3192-3204, 1999.
- 693 Crump, B. C., Baross, J. A., and Simenstad, C. A.: Dominance of particle-attached bacteria in the
 694 Columbia River estuary, USA, Aquat. Microb. Ecol., 14, 7-18, 1998.
- del Giorgio, P. A., and Bouvier, T. C.: Linking the physiologic and phylogenetic successions in
 free-living bacterial communities along an estuarine salinity gradient, Limnol. Oceanogr.,
 47, 471-486, doi:10.4319/lo.2002.47.2.0471, 2002.
- del Giorgio, P. A., Prairie, Y., and Bird, D. F.: Coupling between rates of bacterial production
 and the abundance of metabolically active bacteria in lakes, enumerated using CTC
 reduction and flow cytometry, Microb. Ecol., 34, 144-154, doi:10.1007/s002489900044,
 1997.
- Deshpande, B. N., MacIntyre, S., Matveev, A., and Vincent, W. F.: Oxygen dynamics in permafrost thaw lakes: Anaerobic bioreactors in the Canadian subarctic, Limnol. Oceanogr., 60, 1656–1670, doi:10.1002/lno.10126, 2015.
- Fichot, C. G., and Benner, R.: The spectral slope coefficient of chromophoric dissolved organic
 matter (S275–295) as a tracer of terrigenous dissolved organic carbon in river-influenced
 ocean margins, Limnol. Oceanogr., 57, 1452-1466, doi:10.4319/lo.2012.57.5.1453, 2012.
- Fillion, M.-E., Bhiry, N., and Touazi, M. Differential development of two palsa fields in a peatland located near Whapmagoostui-Kuujjuarapik, Northern Quebec, Canada, Arct.
 Antarct. Alp. Res., 46, 40-54, doi:10.1657/1938-4246-46.1.40, 2014.
- Fletcher, M.: The physiological activity of bacteria attached to solid surfaces, Adv. Microb.
 Physiol., 32, 53-85, 1991.
- Forsström, L., Roiha, T., and Rautio, M.: Responses of microbial food web to increased
 allochthonous DOM in an oligotrophic subarctic lake, Aquat. Microb. Ecol., 68, 171-184,
 2013.
- Galand, P. E., Lovejoy, C., Pouliot, J., and Vincent, W. F.: Heterogeneous archaeal communities
 in the particle-rich environment of an arctic shelf ecosystem, J. Marine Syst., 74, 774-782,
 doi:10.1016/j.jmarsys.2007.12.001, 2008.
- Garneau, M.-È., Vincent, W. F., Alonso-Sáez, L., Gratton, Y., and Lovejoy, C.: Prokaryotic
 community structure and heterotrophic production in a river-influenced coastal arctic
 ecosystem, Aquat. Microb. Ecol., 42, 27-40, doi:10.3354/ame042027, 2006.
- Ghiglione, J. F., Mevel, G., Pujo-Pay, M., Mousseau, L., Lebaron, P., and Goutx, M.: Diel and
 seasonal variations in abundance, activity, and community structure of particle-attached
 and free-living bacteria in NW Mediterranean Sea, Microb. Ecol., 54, 217-231, 2007.

- Gorham, E., and Janssens, J. A.: The distribution and accumulation of chemical elements in five
 peat cores from the mid-continent to the eastern coast of North America, Wetlands, 25,
 259-278, 2005.
- Gorham, E., Lehman, C., Dyke, A., Clymo, D., and Janssens, J., Long-term carbon sequestration
 in North American peatlands, Quat. Sci. Rev., 58, 77-82,
 doi:10.1016/j.quascirev.2012.09.018, 2012.
- Grossart, H. P., and Simon, M.: Limnetic macroscopic organic aggregates (lake snow):
 Occurrence, characteristics, and microbial dynamics in Lake Constance, Limnol.
 Oceanogr., 38, 532-546, 1993.
- Höfle, S., Rethemeyer, J., Mueller, C. W., and John, S.: Organic matter composition and
 stabilization in a polygonal tundra soil of the Lena-Delta, Biogeosciences, 10, 3145-3158,
 doi:10.5194/bg-10-3145-2013, 2013.
- Iriberri, J., Unanue, M., Ayo, B., Barcina, I., and Egea, L.: Bacterial production and growth rate
 estimation from [3H] thymidine incorporation for attached and free-living bacteria in
 aquatic systems, Appl. Environ. Microbiol., 56, 483-487, 1990.
- Judd, K. E., Crump, B. C., and Kling, G. W.: Variation in dissolved organic matter controls
 bacterial production and community composition, Ecology, 87, 2068-2079, 2006.
- Kankaala, P., Huotari, J., Peltomaa, E., Saloranta, T., and Ojala, A. Methanotrophic activity in
 relation to methane efflux and total heterotrophic bacterial production in a stratified,
 humic, boreal lake, Limnol. Oceanogr., 51, 1195-1204, doi:10.4319/lo.2006.51.2.1195,
 2006.
- Karlsson, J., Jonsson, A., and Jansson, M.: Bacterioplankton production in lakes along an altitude
 gradient in the subarctic north of Sweden, Microb. Ecol., 42, 372-382,
 doi:10.1007/s00248-001-0009-9, 2001.
- Kirchman, D. L., and Ducklow, H. W.: Trophic dynamics of particle-bound bacteria in pelagic
 ecosystems: a review, in: Moriaty, D. J. W., and Pullin, R. S. V. (eds.) Detritus and
 Microbial Ecology in Aquaculture, ICLARM Conference Proceedings 14, International
 Center for Living Aquatic Resources Management, Manila, Philippines, 54-82, 1987.
- Kirchman, D. L., Keel, R. G., Simon, M., and Welschmeyer, N. A.: Biomass and production of
 heterotrophic bacterioplankton in the oceanic subarctic Pacific, Deep Sea Res. Part 1
 Oceanogr. Res. Pap., 40, 967-988, 1993.
- Kirchman, D., and Mitchell, R.: Contribution of particle-bound bacteria to total microheterotrophic activity in five ponds and two marshes, Appl. Environ. Microbiol., 43, 200-209, 1982.
- Kirchman, D.: Measuring bacterial biomass production and growth rates from leucine
 incorporation in natural aquatic environments, Methods Microbiol., 30, 227-237,
 doi:10.1016/S0580-9517(01)30047-8, 2001.
- Kling, G. W., Kipphut, G. W., and Miller, M. C.: Arctic lakes and streams as gas conduits to the atmosphere: Implications for tundra carbon budgets, Science, 251, 298-301, 1991.

- Lami, R., Cuperoa, Z., Ras, J., Lebaron, P., and Koblízek, M.: Distribution of free-living and
 particle-attached aerobic anoxygenic phototrophic bacteria in marine environments,
 Aquat. Microb. Ecol., 55, 31-38, 2009.
- Laurion, I., Vincent, W. F., Retamal, L., Dupont, C., Francus, P., MacIntyre, S., and Pienitz, R.:
 Variability in greenhouse gas emissions from permafrost thaw ponds, Limnol. Oceanogr.,
 55, 115-133, doi:10.4319/lo.2010.55.1.0115, 2010.
- Liebner, S., Ganzert, L., Kiss, A., Yang, S., Wagner, D., and Svenning, M. M.: Shifts in methanogenic community composition and methane fluxes along the degradation of discontinuous permafrost, Front. Microbiol., 6, 356, doi:10.3389/fmicb.2015.00356, 2015.
- Lymer, D., Lindström, E. S. and Vrede, K.: Variable importance of viral-induced bacterial
 mortality along gradients of trophic status and humic content in lakes, Freshwater Biol.,
 53, 1101–1113, doi:10.1111/j.1365-2427.2008.02015.x, 2008.
- Masin, M., Uperova, Z. C., Hojerova, E., Salka, I., Grossart, H.-P., and Koblizek, M.:
 Distribution of aerobic anoxygenic phototrophic bacteria in glacial lakes of northern
 Europe, Aquat. Microb. Ecol., 66, 77-86, 2012.
- Matveev, A., Laurion, I., Deshpande, B.N., Bhiry, N., and Vincent, W.F.: High methane
 emissions from thermokarst lakes in subarctic peatlands. Limnol. Oceanogr., 61, doi:
 0.1002/lno.10311, 2016.
- McClelland, J. W., Stieglitz, M., Pan, F., Holmes, R. M., and Peterson, B. J.: Recent changes in nitrate and dissolved organic carbon export from the upper Kuparuk River, North Slope, Alaska, J Geophys Res, 112, G04S60, doi:10.1029/2006JG000371, 2007.
- Michaels, A. F., and Silver, M. W.: Primary production, sinking fluxes and the microbial food
 web, Deep Sea Res. Part 1 Oceanogr. Res. Pap., 35, 473-490, 1988.
- Moquin, P. A., and Wrona, F. J.: Effects of permafrost degradation on water and sediment quality
 and heterotrophic bacterial production of Arctic tundra lakes: An experimental approach,
 Limnol. Oceanogr., 60, 1484–1497, doi:10.1002/lno.10110, 2015.
- Pace, M. L., and Cole, J. J.: Comparative and experimental approaches to top-down and bottom up regulation of bacteria. Microb. Ecol., 28, 181-193, 1994.
- Ploug, H., and Grossart, H. P.: Bacterial growth and grazing on diatom aggregates: Respiratory
 carbon turnover as a function of aggregate size and sinking velocity, Limnol. Oceanogr.,
 45, 1467-1475, 2000.
- Przytulska, A., Bartosiewicz, M., Rautio, M., Dufresne, F., and Vincent, W.F.: Climate effects on
 Arctic *Daphnia* via food quality and thresholds, PLoS ONE, 10, e0126231,
 doi:10.1371/journal.pone.0126231, 2015.
- Przytulska, A., Comte, J., Crevecoeur, S., Lovejoy, C., Laurion, I., and Vincent, W. F.:
 Phototrophic pigment diversity and picophytoplankton in permafrost thaw lakes,
 Biogeosciences, 13, 13-26, doi:10.5194/bg-13-13-2016, 2016.
- Pulido-Villena, E., and Reche, I.: Exploring bacterioplankton growth and protein synthesis to
 determine conversion factors across a gradient of dissolved organic matter, Microb Ecol,
 46, 33-42, doi:10.1007/s00248-002-0004-9, 2003.

- Rautio, M., and Vincent, W. F.: Benthic and pelagic food resources for zooplankton in shallow
 high-latitude lakes and ponds, Freshwater Biol., 51, 1038-1052, doi:10.1111/j.13652427.2006.01550.x, 2006.
- Roehm, C. L., Giesler, R., and Karlsson, J.: Bioavailability of terrestrial organic carbon to lake
 bacteria: The case of a degrading subarctic permafrost mire complex, J. Geophys. Res.,
 114, G03006, doi:10.1029/2008JG000863, 2009.
- Roiha, T., Laurion, I., and Rautio, M.: Carbon dynamics in highly heterotrophic subarctic thaw
 ponds, Biogeosciences, 12, 7223-7237, doi:10.5194/bg-12-7223-2015, 2015.
- Roiha, T., Tiirola, M., Cazzanelli, M., and Rautio, M.: Carbon quantity defines productivity
 while its quality defines community composition of bacterioplankton in subarctic ponds,
 Aquat. Sci., 74, 513-525, doi:10.1007/s00027-011-0244-1, 2012.
- Simon, M., Alldredge, A. L., and Azam, F.: Bacterial carbon dynamics on marine snow, Mar.
 Ecol. Prog. Ser., 65, 205-211, 1990.
- Simon, M., and Azam, F.: Protein content and protein synthesis rates of planktonic marine
 bacteria, Mar. Ecol. Prog. Ser, 51, 201-213, 1989.
- Simon, M., and Kirchman, D.: Isotope dilution during bacterial amino acid uptake in the
 subarctic Pacific: evidence for ammonium utilization by bacteria, EOS, 69, 1146, 1988.
- Smith, D. C., and Azam, F.: A simple, economical method for measuring bacterial protein
 synthesis rates in seawater using 3H-leucine, Mar. Microb. Food Webs, 6, 107-114, 1992.
- Smith, E. M., and Prairie, Y. T.: Bacterial metabolism and growth efficiency in lakes: The importance of phosphorus availability, Limnol. Oceanogr., 49, 137–147, 2004.
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil
 organic carbon pools in the northern circumpolar permafrost region, Global Biogeochem.
 Cy., 23, GB2023, doi:10.1029/2008GB003327, 2009.
- Tuolonen, T., Kankaala, P., Ojala, A., and Arvola, L.: Factors controlling production of
 phytoplankton and bacterial under ice in a humic, boreal lake, J. Plankton Res., 16, 1411 1432, 1994.
- Unanue, M., Ayo, B., Azúa, I., Barcina, I., and Iriberri, J.: Temporal variability of attached and
 free-living bacteria in coastal waters, Microb. Ecol., 23, 27-39, 1992.
- Vallières, C., Retamal, L., Ramlal, P., Osburn, C. L., and Vincent, W. F.: Bacterial production
 and microbial food web structure in a large arctic river and the coastal Arctic Ocean, J.
 Mar. Syst., 74, 756-773, doi:10.1016/j.jmarsys.2007.12.002, 2008.
- Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot,
 M., Billet, M. F., Canário, J., Cory, R. M., Deshpande, B. N., Helbig, M., Jammet, M.,
 Karlsson, J., Larouche, J., MacMillan, G., Rautio, M., Walter Anthony, K. M., and
 Wickland, K.P.: Effects of permafrost thaw on arctic aquatic ecosystems, Biogeosciences
 12, 7129-7167, doi:10.5194/bg-12-7129-2015, 2015.
- Waidner, L. A., and Kirchman, D. L.: Aerobic anoxygenic phototrophic bacteria attached to
 particles in turbid waters of the Delaware and Chesapeake estuaries, Appl. Environ.
 Microbiol., 73, 3936-3944, 2007.

- Watanabe, S., Laurion, I., Chokmani, K., Pienitz, R., and Vincent, W. F.: Optical diversity of
 thaw ponds in discontinuous permafrost: A model system for water color analysis, J.
 Geophys. Res., 116, G02003, doi:10.1029/2010jg001380, 2011.
- Yoon, W. B., and Rosson, R. A.: Improved method of enumeration of attached bacteria for study
 of fluctuation in the abundance of attached and free-living bacteria in response to diel
 variation in seawater turbidity, Appl. Environ. Microbiol., 56, 595-600, 1990.

7 Tables

853 Table 1. Chemical properties of permafrost soil samples from the palsa surface at SAS1. Each

854	value is the mean	(standard deviation)) of triplicates.
		`	

Major constituents	Mean elemental concentration (SD) (g kg ⁻¹ dry weight)	Minor constituents	Mean elemental concentration (SD) (mg kg ⁻¹ dry weight)
С	509 (10.9)	Ba	43.5 (10)
Ν	30.0 (2.6)	Mn	24.5 (12)
Fe	6.04 (1.6)	Cu	6.5 (1.1)
Ca	5.23 (3.4)	Zn	5.8 (0.5)
S	4.86 (0.52)	Cr	5.8 (0.5)
Al	3.67 (0.60)	Ni	5.3 (0.8)
Р	0.545 (0.019)	Мо	1.03 (0.18)
Κ	0.369 (0.178)	Pb	0.93 (0.46)
Na	0.344 (0.123)	Cd	0.12 (0.04)
Mg	0.339 (0.189)		

Table 2. Analysis from duplicate permafrost cores taken from palsas at the SAS2 study site. Gas, ice, and sediment (Sed) proportions (%) are by volume. a_{CDOM} (m⁻¹) is the CDOM absorption at 440 nm. Dissolved organic carbon (DOC, in mg L⁻¹), soluble reactive phosphorus (SRP, in μ g L⁻¹), and nitrate-nitrogen (NO₃, in μ g N L⁻¹) concentrations are potential contributions from melted core ice. N/A* = sample was below detection limit.

Core	Depth (m)	Gas (%)	Ice (%)	Sed (%)	a _{CDOM}	DOC	SRP	NO ₃
А	1.25	5	95	0	12.0	14.7	N/A*	42.1
	1.75	18	82	0	13.3	13.9	N/A*	36.5
	2.77	1	41	58	51.5	28.0	33.1	341
В	1.25	3	97	0	16.2	26.9	N/A*	62.2
	1.75	5	95	0	14.0	17.9	N/A*	63.1
	2.77	2	53	44	10.0	28.8	20.8	93.6

Table 3. Limnological properties of study sites. Zs, sample depth, soluble reactive phosphorus (SRP, in μ g L⁻¹), total nitrogen (TN, in mg L⁻¹), dissolved organic carbon (DOC, in mg L⁻¹), Chlorophyll *a* (Chl *a*, in μ g L⁻¹), total suspended solids (TSS, in mg L⁻¹); PSmax, maximum frequency of particle size distribution (μ m). The S samples correspond to surface at 0 m, and the B samples correspond to the maximum depth of each lake.

Lake	Zs	SRP	TN	DOC	Chl a	TSS	PSmax
SAS1A	S	1.01	0.58	10.4	3.31	5.31	452
	В	5.48	1.76	13.4	4.13	33.0	200
SAS1B	S	1.44	0.70	14.7	3.44	7.43	592
	В	2.18	0.89	14.6	6.22	11.3	262
SAS2A	S	1.42	0.42	8.50	0.94	17.6	300
	В	1.17	0.40	9.00	3.34	4.56	200
SAS2B	S	1.12	0.61	12.8	1.54	6.00	344
	В	8.34	1.88	22.0	7.78	16.5	1338
SAS2C	S	1.03	0.59	11.0	2.24	4.62	229
	В	5.76	1.71	21.0	1.84	15.8	229
KWK12	S	0.41	0.41	6.30	1.12	6.52	592
	В	1.78	0.63	7.60	6.24	27.1	200
SEC	S	0.75	0.26	2.90	1.02	6.97	153
	В	1.56	0.35	3.90	1.58	61.6	175
BGR1	S	0.42	0.24	2.40	0.49	4.57	59
	В	2.06	0.51	3.20	13.4	34.7	59
BGR2	S	1.85	0.38	5.10	0.92	30.8	229
NAS1A	S	4.18	0.25	3.80	0.32	112	102
	В	3.25	0.23	3.30	0.07	163	102
NAS1H	S	1.80	0.24	4.00	1.51	10.9	102
RBL4K	S	0.47	0.74	8.10	1.48	4.00	4
POND9K	S	4.84	0.81	15.1	1.16	3.22	13
RBL9K	S	3.66	0.54	10.2	0.88	1.28	200
RBLOlsha	S	1.09	0.36	9.10	2.05	1.46	175

Table 4. Bacterial properties of study sites. Zs, sample depth, bacterial abundance (BA, in 10^6 cells mL⁻¹, bacterial production rate (BP, in μ g C L⁻¹ h⁻¹), and bacterial community specific growth rates (BG, in d⁻¹). The S samples correspond to surface at 0 m, and the B samples correspond to the maximum depth of each lake.

Lake	Zs	BA	BP	BG
SAS1A	S	0.319	0.896	3.72
	В	2.59	1.074	0.550
SAS1B	S	0.203	0.855	5.58
	В	0.867	0.923	1.41
SAS2A	S	3.15	0.254	0.107
	В	1.57	1.676	1.42
SAS2B	S	0.321	0.804	3.32
	В	1.26	0.588	0.619
SAS2C	S	0.282	1.34	6.30
	В	1.99	0.547	0.364
KWK12	S	3.53	2.34	0.879
	В	1.55	1.164	0.996
SEC	S	1.98	0.885	0.593
	В	0.994	0.650	0.867
BGR1	S	2.79	0.264	0.125
	В	1.03	0.609	0.784
BGR2	S	1.89	1.312	0.920
NAS1A	S	1.71	1.188	0.921
	В	1.16	0.785	0.897
NAS1H	S	0.864	1.253	1.92
RBL4K	S	2.46	0.371	0.200
POND9K	S	2.64	0.791	0.397
RBL9K	S	2.78	0.597	0.285
RBLOlsha	S	0.819	0.849	1.37

Table 5. Comparison of bacterial abundances (BA, 10^6 cells mL⁻¹) and production rates (BP, μ g C L⁻¹ h⁻¹) from thermokarst lakes with other aquatic environments. * Trophic status based on Chl *a* concentrations only.

Name	Location	Trophic status	BA	BP	Reference
Pääjärvi	Finland, in winter	Dystrophic	1.2 - 1.7	0.0083 - 0.096	Tulonen et al., 1994
16 ponds	Subarctic Finland	Oligotrophic	1.75 - 2.75	0.063 - 0.15	Roiha et al., 2012
Mesocosm of small tundra lake	Mackenzie Delta, Northwest Territories	Mesotrophic		0.16	Moquin and Wrona, 2015
SAS2A	Subarctic Québec	Dystrophic, mesotrophic	3.1	0.25	This study
BGR1	Subarctic Québec	Oligotrophic	2.8	0.26	This study
Stukely	Southern Québec	Mesotrophic*	2.7	0.30	del Giorgio et al., 1997
16 lakes	Subarctic Sweden	Oligotrophic, dystrophic	2.4 - 0.4	0.012 - 0.32	Karlsson et al., 2001
RBL4K	Subarctic Québec	Oligotrophic	2.5	0.37	This study
SEC	Subarctic Québec	Oligotrophic, mesotrophic	2.0	0.88	This study
SASIA	Subarctic Québec	Dystrophic, mesotrophic	0.32	0.90	This study
NASIA	Subarctic Québec	Oligotrophic, dystrophic	1.7	1.2	This study
SAS2C	Subarctic Québec	Dystrophic, mesotrophic	0.28	1.3	This study
Lovering	Southern Québec	Oligotrophic*	3.4	1.35	del Giorgio et al., 1997
Massawippi	Southern Québec	Mesotrophic*	3.0	1.69	del Giorgio et al., 1997
Magog	Southern Québec	Eutrophic*	3.38	1.88	del Giorgio et al., 1997
Central	Southern Québec	Mesotrophic*	3.3	2.15	del Giorgio et al., 1997
Brome	Southern Québec	Eutrophic*	3.4	2.24	del Giorgio et al., 1997
KWK12	Subarctic Québec	Mesotrophic, dystrophic	3.5	2.34	This study

Waterloo	Southern Québec Hypereutrophic*	5.9	3.81	del Giorgio et al., 1997
----------	------------------------------------	-----	------	-----------------------------

877 8 Figure captions

Figure 1. Map of the study regions and photos of the 6 study areas.

Figure 2. Palsas and their associated thermokarst lakes in the SAS valley. Upper panel: aerial
view of the SAS1 area; lower panel ground-level view of SAS1B (photograph by Alexander
Culley).

Figure 3. Percent carbon and nitrogen by dry weight in two permafrost cores from SAS palsa
mounds, and proportions of gas, ice, and sediment by volume.

Figure 4. Distribution of particle sizes in lake water and permafrost soil samples. Insert in the top panel shows the soil sample particle-size distribution for both SAS valleys. The insert in the middle panel shows the particle-size distribution in three SAS lakes: SAS1A, SAS1B, and SAS2A. The bottom panel shows the particle-size distributions from the SAS2 permafrost core samples.

Figure 5. Difference spectra between the frequency distribution of particles in surface lake water
(SAS2A) relative to active layer soil, permafrost and bottom lake water.

Figure 6. Size-fractionated total suspended solids (TSS), bacterial abundance (BA), bacterial
production (BP), and bacterial cell-specific growth rates (BG) from five thermokarst lakes and
one rock-basin lake. Each is the mean ± SD of triplicate samples.

Figure 7. Planktonic variables in the 5-day experiment with amendments of glucose (G), glucose

plus phosphorus (G+P), phosphorus (P), and palsa soil (S) to SAS1B lake water. The control (C)

is shown on both day 1 and day 5. Each value is the mean \pm SD of triplicate samples.

Figure 1. Map of the study regions and photos of the 6 study areas. NAS is a mineral-rich site located in an area of continuous permafrost. BGR and SEC are also mineral-rich sites, but located in areas of discontinuous permafrost. The KWK and SAS lakes are in areas of sporadic permafrost, but KWK is mineral-rich while SAS is in a peatland landscape. The rock-basin lakes (RBL) are not of permafrost-origin.



903

Figure 2. Palsas and their associated thermokarst lakes in the SAS peatland valley. Upper panel:
aerial view of the SAS1 area; lower panel ground-level view of SAS1B (photograph by
Alexander Culley).





Figure 3. Percent carbon and nitrogen by dry weight in two permafrost cores from SAS palsamounds, and proportions of gas, ice, and sediment by volume.

Figure 4. Distribution of particle sizes in lake water and permafrost soil samples. Insert in the top panel shows the soil sample particle-size distribution for both SAS valleys. The insert in the middle panel shows the particle-size distribution in three SAS lakes: SAS1A, SAS1B, and SAS2A. The bottom panel shows the particle-size distributions from the SAS2 permafrost core samples.





921 Figure 5. Difference spectra between the frequency distribution of particles in surface lake water922 (SAS2A) relative to active layer soil, permafrost and bottom lake water.



Figure 6. Size-fractionated total suspended solids (TSS), bacterial abundance (BA), bacterial production (BP), and bacterial cell-specific growth rates (BG) from five thermokarst lakes and one rock-basin lake. Each is the mean ± SD of triplicate samples.



Figure 7. Planktonic variables in the 5-day experiment with amendments of glucose (G), glucose plus phosphorus (G+P), phosphorus (P), and palsa soil (S) to permafrost peatland SAS1B lake water. The control (C) is shown on both day 1 and day 5. Each value is the mean \pm SD of triplicate samples.



Lake	Zm	Latitude	Longitude
SAS1A	1.6	55.218800	77.707950
SAS1B	1.2	55.219013	77.707801
SAS2A	2.7	55.225018	77.696580
SAS2B	1.5	55.225212	77.696017
SAS2C	2.1	55.225083	77.694933
KWK12	2.7	55.330072	77.503849
SEC	3.4	55.701033	76.643850
BGR1	4.2	56.610833	76.215000
BGR2	1.1	56.610833	76.215617
NAS1A	3.2	56.923779	76.378293
NAS1H	3.6	56.924121	76.377038
RBL4K	0.7	55.331798	77.699393
POND9K	0.5	55.361083	77.650861
RBL9K	1.2	55.361083	77.650861
RBLOlsha	1.6	55.282452	77.739834

935 Table S1. Physical properties of study sites. Zm, maximum depth (m); location given by latitude

936 (°N), and longitude (°W).

937

Figure S1. CDOM absorbance curves for SAS permafrost soil (thawed in Milli-Q water) and lakewater.

