

# Analytical solution of nitracline with the evolution of subsurface chlorophyll maximum in stratified water columns

Xiang Gong<sup>1,2</sup>, Wensheng Jiang<sup>2,3</sup>, Linhui Wang<sup>2</sup>, Huiwang Gao<sup>2,4\*</sup>, Emmanuel Boss<sup>5</sup>, Xiaohong Yao<sup>2,4</sup>, Shuh-Ji Kao<sup>6</sup>, Jie Shi<sup>2</sup>

5 <sup>1</sup> School of Mathematics and Physics, Qingdao University of Science and Technology,  
Qingdao 266061, P. R. China

<sup>2</sup> Key Laboratory of Marine Environment and Ecology (Ministry of Education of China), Ocean University of China, Qingdao 266100, P. R. China

10 <sup>3</sup> Key Laboratory of Physical Oceanography (Ministry of Education of China), Ocean University of China, Qingdao 266100, P. R. China

<sup>4</sup> Qingdao Collaborative Center of Marine Science and Technology, Ocean University of China, Qingdao 266100, P. R. China

<sup>5</sup> School of Marine Sciences, University of Maine, Orono 04469-5706, USA

15 <sup>6</sup> State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 361005, P. R. China

\*Corresponding author: Huiwang Gao, hwgao@ouc.edu.cn

## Abstract:

In a stratified water column, the nitracline is a layer where the nitrate concentration increases below the nutrient-depleted upper layer, exhibiting a strong vertical gradient in the euphotic zone. The subsurface chlorophyll maximum layer (SCML) forms near the bottom of euphotic zone, acting as a trap to diminish the upward nutrient supply. Depth and steepness of the nitracline are important measurable parameters related to the vertical transport of nitrate into the euphotic zone. The correlation between the SCML and the nitracline has been widely reported in the literature, but the analytic solution for the relationship between them is not well established. By incorporating a piecewise function for the approximate Gaussian vertical profile of chlorophyll, we derive analytical solutions of a specified nutrient-phytoplankton model. The model is well suited to explain basic dependencies between a nitracline and a SCML. The

- 30 analytical solution shows that the nitracline depth is deeper than the depth of SCML, shoaling with an increase in light attenuation coefficient and with a decrease in surface light intensity. The inverse proportional relationship between the light level at the nitracline depth and the maximum rate of new primary production is derived. Analytic solutions also show that a thinner SCML corresponds to a steeper nitracline.
- 35 The nitracline steepness is positively related to light attenuation coefficient, but independent of surface light intensity. The derived equations of the nitracline in relation to the SCML provide further insight into the important role of the nitracline in marine pelagic ecosystems.

## 1 Introduction

40 Nitrogen availability, especially the nitrate upward supply to the euphotic zone where light intensity is sufficient to support net photosynthesis, limits the primary productivity in a stratified water column (Falkowski et al., 1998). Specifically, the nitrate supply from below and the light attenuated from above with the depth collaboratively affect the growth of phytoplankton and lead to the subsurface  
45 chlorophyll maximum (SCM) (Riley et al., 1949; Steele and Yentsch, 1960; Herbland and Voituriez, 1979; Cullen, 1982). The SCM layer (SCML) has attracted much attention since Riley (1949) because the layer contributes significantly to new primary production (NPP) in stratified waters (Probyn et al., 1995; Ross and Sharples, 2007; Fernand et al., 2013). The synergistic physical and biological interaction leads to a  
50 strong vertical nitrate gradient, conventionally referred to as the nitracline (Eppley et al., 1978; Herbland and Voituriez, 1979; Cullen and Eppley, 1981). Depth and steepness of the nitracline are important measurable parameters in regulating the supply of nitrate to the euphotic zone, and hence affecting NPP (Lewis et al., 1986; Bahamón et al., 2003; Aksnes et al., 2007; Cermeno et al., 2008; Omand and  
55 Mahadevan, 2015).

The nitracline depth physically depends on the degree of water column stratification and the magnitude of momentum transfer associated with wind stress (Denman and Gargett, 1983; Laanemets et al., 2004). It also depends on momentum transfer from below (Lipschultz et al., 2002) and in some cases, vertical advection such as  
60 upwelling (Laanemets et al., 2004). However, in a relatively stable environment, the SCML may restrict the diffusive flux of nitrates to the euphotic zone and continually erode the nitracline supposing that sufficient light is available (Probyn et al., 1995). The SCML thereby acts as an effective nutrient trap, regulating the nitracline depth (Banse, 1982; Beckmann and Hense, 2007; Klausmeier and Litchman, 2001; Probyn  
65 et al., 1995). On the other hand, variation of nitracline steepness, which is critical to determine the nitrate supply, was poorly understood due to lack of high vertical resolution data, e.g., both bottle data and Argo data tend to have low vertical

resolution sampling. Some studies showed that nitrogen flux is dependent more on the nitracline steepness than on the density gradients regulating turbulent diffusion (Bahamón and Cruzado, 2003; Bahamón et al., 2003; Lavigne et al., 2015). Thus, these measurable features of nitracline and their correlation with SCML may provide insightful information for mechanisms of the productivity in pelagic ecosystem, and the analytic solutions for these parameters may fill the knowledge gap.

Although a close relationship between the nitracline and SCML is always observed, the quantitative nature of nitracline in relation to the SCML formation has not been studied. The system of phytoplankton and the limiting nutrient on the vertical axis was often utilized to study the depth, intensity, and persistence of the SCML. Major theoretical results include photoacclimation (increase of chlorophyll per cell) (Steele, 1964; Fennel and Boss, 2003), bistability (Yoshiyama and Nakajima, 2002; Ryabov et al., 2010), oscillating SCM (Huisman et al., 2006), hysteresis conditions (Kiefer and Kremer, 1981; Navarro and Ruiz, 2013), and the ESS (evolutionary stable strategy) depth obtained by game-theory approach (Klausmeier and Litchman, 2001; Mellard et al., 2011). Recent mathematical studies solved the persistence and uniqueness of the steady state solution (Du and Hsu, 2010; Hsu and Yuan, 2010; Du and Mei, 2011), and gave rigorous proofs for the above-mentioned ESS depth and the game-theory approach (Du and Hsu, 2008a, b). Additionally, several modeling studies have been conducted to quantitatively assess the importance of different physic-biological processes leading to SCML (Jamart et al., 1977; Jamart et al., 1979; Varela et al., 1994; Klausmeier and Litchman, 2001; Hodges and Rudnick, 2004; Beckmann and Hense, 2007).

Among the studies using the nutrient-phytoplankton model, Klausmeier and Litchman (2001) first analytically derived the vertical nutrient distribution with the development of the SCML. In that model, the concentration of the limiting nutrient was found to be low and constant above the SCML and linearly increasing with depth below this layer in poorly mixed water column. Building on that model, Mellard et al. (2011) added stratification and surface nutrient input, which can make phytoplankton

grow in both the surface mixed layer and deep layer (SCML) simultaneously. Fennel and Boss (2003) derived that the sum of nutrients and phytoplankton at steady state will increase monotonically below the surface mixed layer until it equals the fixed  
100 nutrient concentration. By incorporating a generalized Gaussian function for vertical chlorophyll profile into the nutrient-phytoplankton dynamic equation, Gong et al. (2015) obtained that the steady-state nitrate concentration increased from the upper community compensation depth to the SCML depth. None of the studies, however, focused on the quantitative nature of nitracline in relation to the SCML in the  
105 stratified waters.

In this paper, we modified the nutrient-phytoplankton model by Gong et al. (2015) to study the roles of SCM in reshaping the nitracline. Two additional terms, atmospheric input, which promotes the growth of phytoplankton in the surface mixed layer, and the phytoplankton self-shading, which regulates the light penetration, were  
110 introduced into the previous model. Accordingly, a piecewise function comprising a constant value within the surface mixed layer and a Gaussian function below this layer was used as a fit to the steady state vertical chlorophyll profiles simulated by the nutrient-phytoplankton model. By incorporating the piecewise function into the nutrient-phytoplankton model, we derived the analytic solutions for the properties of  
115 the nitracline and the SCML in steady state, respectively, and the relationship between them was examined in response to light availability, surface nutrient input, and vertical diffusivity.

## **2 Definitions and Models**

### *2.1 Models*

#### *2.1.1 Dynamic equations*

  
120

We consider the following equations for phytoplankton and nutrient dynamics in stratified waters (Eqs. 1-2), where light and nitrogen are two limiting factors for phytoplankton growth (Fig. 1). The change in phytoplankton at depth  $z$  is the balance of the growth and death, and the passive moving (sinking and mixing) (Eq. 1). An

125 eddy diffusion coefficient  $K_v$  redistributes phytoplankton in the water column. Depth  $z$  is increasing toward the seabed.

$$\left\{ \begin{array}{l} \frac{\partial P}{\partial t} = \mu_m \min(f(I), g(N))P - \varepsilon P - w \frac{\partial P}{\partial z} + \frac{\partial}{\partial z} \left( K_v(z) \frac{\partial P}{\partial z} \right) \\ \frac{\partial N}{\partial t} = -\gamma \mu_m \min(f(I), g(N))P + \gamma \alpha \varepsilon P + N_{in}(z) + \frac{\partial}{\partial z} \left( K_v(z) \frac{\partial N}{\partial z} \right) \end{array} \right. \quad (1) \quad (2)$$

where  $P$  denotes the chlorophyll concentration ( $\text{mg m}^{-3}$ ). We assume the chlorophyll distribution to represent the distribution of phytoplankton biomass (that means that the photoacclimation of phytoplankton is ignored, and the SCM refers to the subsurface biomass maximum, SBM). This is a significant simplification. In fact, phytoplankton increases inter-cellular pigment concentration when light level decreases (Cullen, 1982; Fennel and Boss, 2003; Cullen, 2015). Usually, the depths of SCML and SBML are separate, and the latter is shallower than the former.

135 Nitrogen  $N$  (in unit:  $\text{mmol N m}^{-3}$ ) taken up by phytoplankton includes three sources, i.e., recycling from dead phytoplankton, atmospheric input to the surface mixed layer, and supply by mixing from deep water (Eq. 2).  $\gamma$  is the nitrogen content of the phytoplankton ( $\text{mmol N per mg Chl}$ ). Following Mellard et al. (2011), we consider the nitrogen input from atmosphere at the rate of  $N_{in}(z)$ , setting as a delta function at  $z=0$

140 with the total nutrient input in the surface mixed layer  $N_{inML}$ ,  $N_{inML} = \int_0^{z_s} N_{in}(z) dz$ , where  $z_s$  is the depth of surface mixed layer. Note that nitrogen input through the activity of nitrogen fixers is excluded. However, trichodesmium, if they are a mat at the surface, will be modeled similarly to the atmospheric input term.

$\mu_m$  is the maximum growth rate of phytoplankton,  $\varepsilon$  is the loss rate of phytoplankton (including respiration, mortality, zooplankton grazing), and  $\alpha$  is the recycling rate of dead phytoplankton ( $0 < \alpha < 1$ ). The specific rate of loss processes ( $\varepsilon$ ) is assumed to be vertically homogeneous due to lack of data (similar to Sverdrup, 1953). The treatment of grazing loss, is, in the least, an oversimplification, though many numerical models used a similar one (e.g., Klausmeier and Litchman, 2001; Fennel and Boss, 2003; Huisman et al., 2006). The growth-limited function for light  $I$  and nitrogen  $N$  is given

by  $\mu_m \min(f(I), g(N))$ . The Michaelis-Menten form for the light-limiting function  $f(I)$  and nitrogen-limiting function  $g(N)$ ,  $f(I)=I/(K_I+I)$  and  $g(N)=N/(K_N+N)$  is used, where  $K_I$  and  $K_N$  denote the half-saturation constants for light and nitrogen, respectively. The net growth rate,  $\mu_m \min(f(I), g(N)) - \varepsilon$ , is positive only if both light-limiting term  
155  $\mu_m f(I)$  and nitrogen-limiting term  $\mu_m g(N)$  are larger than the loss rate  $\varepsilon$ .

Light intensity  $I$  decreases exponentially with depth according to the Lambert-Beer law,

$$I(z) = I_0 \exp\left(-K_w z - K_c \int_0^z P dz\right) \quad (3)$$

where  $I_0$  is the surface light intensity,  $K_w$  and  $K_c$  are light attenuation coefficients of  
160 water and phytoplankton, respectively. For the sake of simplicity, we assume that both  $K_w$  and  $K_c$  are constant with depth.

The sinking velocity of phytoplankton  $w$  is non-negative in the chosen coordinate system. We assumed it to be constant with depths, neglecting the influencing of density gradients (pycnoclines), which may cause vertical variations in sinking.

165 To describe the water column stratification, we assume that the vertical eddy diffusivity  $K_v$  depends on depths,

$$K_v(z) = \begin{cases} K_{v1} & 0 \leq z \leq z_s \\ K_{v2} & z_s < z < z_b \end{cases} \quad (4)$$

where  $z_b$  is set to the bottom boundary of this model and is assumed to be sufficiently deep where the chlorophyll concentration approaches to zero. We assume that surface  
170 diffusivity ( $K_{v1}$ ) and subsurface diffusivity ( $K_{v2}$ ) (Lande and Wood, 1987; Hodges and Rudnick, 2004) are constant and  $K_{v1}$  is large enough to homogenize the chlorophyll and nitrogen concentration in the surface mixed layer. A gradual transition from the surface mixed layer to the deep one (in the vicinity of  $z_s$ ) can be written in terms of a generalized Fermi function (Ryabov et al., 2010),  $K_v(z) = K_{v2} + \frac{K_{v1} - K_{v2}}{1 + e^{z - z_s/l}}$ , where  
175 parameter  $l$  characterizes the width of the transition layer. In our numerical study, we assumed this transition layer is 2 m. For simplicity, parameter  $l$  is set to infinitely thin

in analytic solutions. A comprehensive list of symbols is given in Table 1.

The zero-flux boundary condition for the phytoplankton at the surface is used. At the bottom boundary of the model domain ( $z=z_b$ ) the Dirichlet boundary condition is used, i.e.,  $P \rightarrow 0$  for  $z \rightarrow z_b$ . Fennel and Boss (2003) used an infinite depth as  $z_b$  ( $z_b \rightarrow \infty$ ).  
 180 For the nitrogen distribution we set  $N=N_{inML}$  at the surface and diffusing into the water column a prescribed nitrate gradient,  $n$ , at the bottom. That is,

$$\begin{cases} \left( wP - K_{v1} \frac{\partial P}{\partial z} \right) \Big|_{z=0} = 0, & P(z_b) = 0. \\ N \Big|_{z=0} = N_{inML}, & \frac{\partial N}{\partial z} \Big|_{z=z_b} = n. \end{cases} \quad (5)$$

### 2.1.2 Fitting equation for vertical chlorophyll profile

In many stratified water columns, the vertical distribution of chlorophyll concentration is homogeneous within the surface mixed layer and appears a Gaussian below this layer (Fig. 2a), which is typical in open oceans (Uitz et al., 2006), shelf seas (Sharples et al., 2001), stratified estuary (Lund-Hansen, 2011), and arctic waters (Martin et al., 2012). The non-uniform vertical profile of chlorophyll with an SCML was first modeled by a generalized Gaussian function (Lewis et al., 1983), which has  
 190 subsequently been widely used with small modifications. For example, Platt et al. (1988) superimposed a constant background on the generalized Gaussian, and fitted it to field data on the vertical distribution of chlorophyll from coastal, upwelling, open oceans and Arctic waters. Afterward, some studies introduced a parameter to represent the slope of the Gaussian curve (Matsumura and Shiomoto, 1993; Mu Oz Anderson et al., 2015). In particular, to account for the observed characteristic that surface values always exceed the bottom ones (Fig. 2a), the generalized Gaussian functional form has been modified with a superimposition of a background linearly or exponentially decreasing with depth (Uitz et al., 2006; Mignot et al., 2011; Ardyna et al., 2013).  
 195

For simplicity, to analytically study the role of the SCML on shaping the nitracline, we therefore propose a piecewise function comprising a constant value in the surface mixed layer and below a general Gaussian function (Eq. 6) to approximate the vertical  
 200



profile of chlorophyll concentration in Fig. 2a.

$$P = \begin{cases} P_0 & 0 \leq z \leq z_s \\ P_{\max} \exp\left[-\frac{(z-z_m)^2}{2\sigma^2}\right] & z_s < z < z_b \end{cases} \quad (6)$$

205 where  $P$  is chlorophyll concentration as a function of depth  $z$ ,  $P_0$  is the chlorophyll concentration within the surface mixed layer, and  $P_{\max} = h/(\sigma\sqrt{2\pi})$  represents the maximum value of chlorophyll below the surface mixed layer. Considering the influence of the surface mixed layer on the chlorophyll vertical distribution,  $h$  is less than the total chlorophyll concentration integrated through the water column. Note

210 that the vertical distribution of chlorophyll is an incomplete general Gaussian function below the surface mixed layer (*see* Fig. 2a). The three Gaussian parameters ( $P_{\max}$ ,  $z_m$ ,  $\sigma$ ) can characterize the SCM phenomenon. Thus,  $z_m$  is the depth of the maximum chlorophyll (the peak of the bell shape), and  $\sigma$  is the standard deviation of Gaussian function, which controls the width of the SCML. The upper and lower boundary of

215 SCML can be defined as  $z_m - \sigma$  and  $z_m + \sigma$ , respectively, which are at the depths where there is the balance between phytoplankton growth and losses and thus reflect the physical-biological ecosystem dynamics associated with SCML (Beckmann and Hense, 2007; Gong et al., 2015). To explore the SCML in stratified waters, we assume the surface mixed layer is shallower than the upper boundary of the SCML, i.e.,

220  $z_s < z_m - \sigma$ . Examples of the piecewise function that reasonably well fitted to vertical chlorophyll profiles in the northern South China Sea (SCS) can be found in Gong et al. (2014).

The piecewise function approximation (Eq. 6) was evaluated and justified through numerical simulation of the nutrient-phytoplankton system (Eqs. 1-2), which is solved

225 with a semi-implicit time stepping scheme. The vertical resolution is uniform (2 m), extending down to 200 m. We assumed a small uniformly distributed concentration of phytoplankton ( $P(z,0)=0.1 \text{ mg m}^{-3}$ ) and nitrogen ( $N(z,0)=0.1 \text{ mmol N m}^{-3}$ ) as the initial conditions and run the model until in converge to a steady state (The modeling results show that the steady state has no relationship with the initial values of

230 phytoplankton and nitrogen). We use the biologically reasonable parameter values given in Table 1 to represent the system at Station SEATS (South East Asia Time Series) in the northern SCS. Thus, the specific (calibrated) model solution is considered as an example to obtain the analytic solutions of nitracline.

Fig. 3 shows the numerically simulated equilibrium distributions of nitrogen, light, and chlorophyll. In addition, the simulated vertical profile of chlorophyll is fitted well by the piecewise function of chlorophyll using the least square method (Fig. 3). Many numerical solutions of the nutrient-phytoplankton system have reproduced the vertical chlorophyll profile with the SCML (Fennel and Boss, 2003; Huisman et al., 2006; Ryabov et al., 2010). Thus, analogous to the study by Klausmeier and Litchman (2001), we incorporate the piecewise function (Eq. 6) to the nutrient-phytoplankton system (Eqs. 1-2) at steady state to examine the roles of the SCML in reshaping the nitracline. We note that the useful delta function approximation in Klausmeier and Litchman (2001) was verified by both simulation and rigorous mathematics (Du and Hsu, 2008a, b). As presented above, the assumption of the piecewise function approximation is physically practical.

## 2.2 Definition of the nitracline

The vertical distributions of nitrate often exhibit a strong gradient in depth (the nitracline), but the feature of nitracline (depth, steepness) is variable in euphotic zones due to the combined effect of physical and biological processes.

250 Many studies define the nitracline depth as the location where the maximum vertical gradient in nitrate concentrations occurs (Eppley et al., 1979; Bahamón et al., 2003; Wong et al., 2007; Beckmann and Hense, 2007; Martin et al., 2010). To measure the defined depth, a high vertical resolution of nitrate concentrations is needed and this is a big technique challenge existing for a long time. Thus, some definitions were also proposed to make the depth measurable. For example, one definition is the depth where the nitrate concentration reaches a prescribed concentration, e.g., 0.05, 0.1, 1.0, or 12 mmol N m<sup>-3</sup> (Cullen and Eppley, 1981; Koeve

et al., 1993; Martin and Pondaven, 2003). Some studies choose it to be the first depth where nitrogen is detectable (e.g., 0.05 or 0.1 mmol N m<sup>-3</sup>) (Cermeno et al., 2008; 260 Hickman et al., 2012) or where the nitrogen concentration exceeds the mixed layer value by a prescribed concentration difference (e.g., 0.05 mmol N m<sup>-3</sup>) (Laanemets et al., 2004). Significant differences exist between these defined depths, i.e., the depth of maximal nitrate gradient was found to be deeper by 10 m from the first depth where nitrate can be detected (Eppley et al., 1978), while the nitrate gradient at the first 265 detectable depth of nitrate is nearly zero (Cermeno et al., 2008).

With the development of nearly continuous nitrate profile measurement using the In Situ Ultraviolet Spectrophotometer (ISUS) optical nitrate sensor (Johnson and Coletti, 2002; Johnson et al., 2010), the detection of the maximum nitrate gradient could be more accurate than before. In this study, we adopt the location of the maximum nitrate 270 gradient ( $\max(dN/dz)$ ) in the euphotic zone as the nitracline depth ( $z_n$ ), which can be

$$\text{expressed by } \left. \frac{d^2N}{dz^2} \right|_{z_n} = 0 \text{ and } \left. \frac{d^3N}{dz^3} \right|_{z_n} < 0.$$

Below the surface mixed layer, the steady-state version of reduces to  $\gamma\mu_m \min(f(I), g(N))P - \gamma\alpha\varepsilon P = K_{v2} \frac{d^2N}{dz^2}$ . Thus, according to our model approach (Eq. 2) 275 the nitracline depth where  $\frac{d^2N}{dz^2} = 0$  represents a balance between the nutrient uptake and the recycling of phytoplankton loss, i.e.,  $\mu_m \min(f(I), g(N)) = \alpha\varepsilon$ .

The nitracline steepness is defined as the nitrate gradient at the nitracline depth ( $\left. \frac{dN}{dz} \right|_{z_n}$ ) in this study (Laanemets et al., 2004; Aksnes et al., 2007).

### 2.3 Data sources

The nitrate profiles were obtained from the ISUS measurement at the SEATS 280 station during the CHOICE-C 2012 summer cruise. 9 casts were conducted during Aug. 6-7, 2012. The raw ISUS nitrate data, which employed temperature-compensation, were first calibrated by the AutoAnalyzer 3 (AA3), and

then smoothed to remove noise. The sampling frequency was set at 5 Hz and the raw data were thus smoothed with a 25-point moving average in the surface mixed layer, a 5-point moving average in the SCML, and a 15-point moving average below the SCML. The data were then interpolated by a cubic spline function. The corresponding temperature was obtained from Conductivity-Temperature-Depth (CTD) measurements. Overall, nine sets of profiles are available to examine our analytical solutions.

## 3 Results

### 3.1 Relations between nitracline and SCML

#### 3.1.1 Nitracline depth and SCML

At steady state, multiplying Eq. (1) by  $\gamma$  then adding Eqs. (1) and (2) leads to:

$$(\alpha - 1)\varepsilon P - w \frac{dP}{dz} + \frac{d}{dz} \left( K_v(z) \frac{dP}{dz} \right) + \frac{1}{\gamma} \frac{d}{dz} \left( K_v(z) \frac{dN}{dz} \right) + \frac{N_m(z)}{\gamma} = 0 \quad (7)$$

By substituting the expression of eddy diffusivity (Eq. 4) and the fitted, depth dependent function of chlorophyll ( $P(z)$ , Eq. 6) into Eq. (7), we obtain the diffusive term of nitrate below the surface mixed layer, that is,

$$K_{v2} \frac{d^2 N}{dz^2} = \left[ -\frac{K_{v2}}{\sigma^2} \left( \frac{z - z_m}{\sigma} \right)^2 - \frac{w}{\sigma} \left( \frac{z - z_m}{\sigma} \right) + \frac{K_{v2}}{\sigma^2} + (1 - \alpha) \varepsilon \right] \gamma P \quad z_s < z < z_b \quad (8)$$

Letting  $d^2 N/dz^2 = 0$  in Eq. (8), for  $P > 0$  one gets

$$-\frac{K_{v2}}{\sigma^2} \left( \frac{z - z_m}{\sigma} \right)^2 - \frac{w}{\sigma} \left( \frac{z - z_m}{\sigma} \right) + \frac{K_{v2}}{\sigma^2} + (1 - \alpha) \varepsilon = 0 \quad z_s < z < z_b \quad (9)$$

Solving this quadratic equation of depth  $z$ , we obtain the depths  $z_{n1}$  and  $z_{n2}$ ,

$$\begin{cases} z_{n1} = z_m - \frac{w\sigma^2}{2K_{v2}} - \sqrt{\frac{w^2\sigma^4}{4K_{v2}^2} + \frac{(1-\alpha)\varepsilon\sigma^4}{K_{v2}} + \sigma^2} \\ z_{n2} = z_m - \frac{w\sigma^2}{2K_{v2}} + \sqrt{\frac{w^2\sigma^4}{4K_{v2}^2} + \frac{(1-\alpha)\varepsilon\sigma^4}{K_{v2}} + \sigma^2} \end{cases} \quad (10)$$

Taking the derivative of Eq. (8) with respect to depth  $z$ , we get  
 $K_{v2} \frac{d^3 N}{dz^3} = \left[ -\frac{2K_{v2}}{\sigma^4} (z - z_m) - \frac{w}{\sigma^2} \right] \gamma P - K_{v2} \frac{d^2 N}{dz^2} \frac{z - z_m}{\sigma^2}$ . Obviously, at depth  $z_{n1}$ ,  $d^3 N/dz^3 > 0$ ,

305 and at depth  $z_{n2}$ ,  $d^3 N/dz^3 < 0$ . That is,  $z_{n2}$  is the location of maximum nitrate gradients.

We obtain that the nitracline depth refers to the depth  $z_{n2}$ , i.e.,

$$z_n = z_{n2} = z_m - \frac{w\sigma^2}{2K_{v2}} + \sqrt{\frac{w^2\sigma^4}{4K_{v2}^2} + \frac{(1-\alpha)\varepsilon\sigma^4}{K_{v2}} + \sigma^2} \quad (11)$$

Particularly, Eq. (9) became a linear function of depth  $z$  when the second order item coefficient ( $K_{v2}/\sigma^4$ ) is zero, thus has only one solution. In fact, in typical stratified  
 310 waters the diffusivity below the surface mixed layer ( $K_{v2}$ ) is  $1.9 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ , and the thickness of SCML ( $2\sigma$ ) is from several meters to tens of meters (Cullen, 2015), thus,  $K_{v2}/\sigma^4$  (values from  $8.64 \times 10^{-9}$  to  $7.78 \text{ m}^2 \text{ s}^{-1}$ ) can be neglected in some cases. When  $K_{v2}/\sigma^4 \rightarrow 0$ , for non-zero sinking velocity we get one solution from Eq. (9),

$$z_n = z_m + \frac{(1-\alpha)\varepsilon\sigma^2}{w} \quad w \neq 0 \quad (12)$$

315 Both Eqs. (11) and (12) show that the nitracline depth is located below the SCML depth, i.e.,  $z_n > z_m$  (Fig. 1). A numerical study in weak vertical mixing environments showed a similar result (Beckmann and Hense, 2007). Note that the SCML represents the SBML in our model. In some oligotrophic oceans, the SCML will be deeper than the SBML due to the effect of photoacclimation on the vertical distribution of  
 320 chlorophyll (Fennel and Boss, 2003). For example, Li et al. (2015) showed that the modeled maximum nitrate gradient (nitracline) occurred below the depth of SCML in the northern SCS, and then we can deduce that the nitracline depth is also deeper than the depth of SBML. In the Mediterranean Sea, Bahamón et al. (2003) found that the nitracline occurred below the depth of SCML at 88% of the stations (50 out of 57  
 325 stations). As well known, the SCML in the Mediterranean is often due to photoacclimation. It is not surprised for the other 7 stations against  $z_n > z_m$  in the Mediterranean Sea if photoacclimation leads to a much deeper SCML than the SBML.

Thus, we conclude that the nitracline depth is located below the SBML depth, while the locations may not be necessarily true between the depths of nitracline and SCML.

### 330 3.2.2 Nitracline steepness and SCML

To illustrate the relationship between the nitracline steepness and the SCML, by integrating Eq. (8) from depth  $z_n$  to  $z_b$  and using the assumption for phytoplankton at the bottom boundary, i.e.,  $P \rightarrow 0$  for  $z \rightarrow z_b$  (Eq. 5), we get

$$\frac{dN}{dz}\Big|_{z_n} = \frac{dN}{dz}\Big|_{z_b} + \left( \frac{z_n - z_m}{\sigma^2} + \frac{w}{K_{v2}} \right) \gamma P\Big|_{z_n} - \frac{(1-\alpha)\varepsilon\gamma}{K_{v2}} \int_{z_n}^{z_b} P dz \quad (13)$$

335 This equality indicates that the nitracline gets steeper as the distance between the depths of nitracline layer and SCML is increased.

Incorporating Eq. (11) into Eq. (13) leads to

$$\frac{dN}{dz}\Big|_{z_n} = \frac{dN}{dz}\Big|_{z_b} + \left( \sqrt{\frac{w^2}{4K_{v2}^2} + \frac{(1-\alpha)\varepsilon}{K_{v2}} + \frac{1}{\sigma^2}} + \frac{w}{2K_{v2}} \right) \gamma P\Big|_{z_n} - \frac{(1-\alpha)\varepsilon\gamma}{K_{v2}} \int_{z_n}^{z_b} P dz \quad (14)$$

Equation (14) indicates that the nitracline steepness is negatively related to the  
340 thickness of SCML.

## 3.2 Analytical solutions for nitracline features

### 3.2.1 Depth of the nitracline

By substituting the general Gaussian function for chlorophyll below the surface mixed layer (Eq. 6) into Eq. (1), we obtain the steady-state net growth rate of  
345 phytoplankton below the surface mixed layer:

$$\mu_m \min(f(I), g(N)) - \varepsilon = -K_{v2} \left( z - \left( z_m - \frac{w\sigma^2}{2K_{v2}} \right) \right)^2 / \left( \sigma^4 + w^2/4K_{v2} + K_{v2}/\sigma^2 \right) \quad (15)$$

Let  $z_0 = z_m - w\sigma^2/(2K_{v2})$  in the first term of the right-hand of Eq. (15). From the result given by Gong et al. (2015), we know that  $z_0$  is the location of the maximum growth rate of phytoplankton (hereafter named as the depth of optimal growth), where

350 the transition from nutrients limitation to light limitation occurs (i.e.,  $f(I)=g(N)$  at depth  $z_0$ ).

Clearly,  $z_n > z_0$  (Eq. 11). Hence, the growth of phytoplankton at the nitracline depth  $z_n$  is limited by light, i.e.,  $\mu_m \min(f(I), g(N))|_{z_n} = \mu_m f(I)|_{z_n}$ . In other words, the growth rate of phytoplankton at the nitracline depth is a function of the light level at  
 355 the nitracline depth,  $I(z_n)$ . Thus, from Eqs. (11) and (15), we obtain the growth rate of phytoplankton at the nitracline depth, that is,

$$\mu_m f(I)|_{z_n} = \alpha \varepsilon \quad (16)$$

Note that the derivation of Eq. (16) only works when light and nutrient limitation (Blackman's law of limiting factors) is applied. Substituting the Michaelis-Menten  
 360 form for  $f(I)$  into Eq. (16), we have

$$I(z_n) = \frac{K_I}{\mu_m / \alpha \varepsilon - 1} \quad (17)$$

Rearranging Eq. (17), we find  $\mu_m - \alpha \varepsilon = \alpha \varepsilon K_I / I(z_n)$ . This equality indicates that the maximum rate of NPP,  $(\mu_m - \alpha \varepsilon)P$ , is inversely proportional to the light level at the nitracline depth,  $I(z_n)$ . Lande et al. (1989) found that higher maximum rates of  
 365 population growth corresponded to shallower nitracline depths in the central North Atlantic.

Furthermore, insertion of Eq. (3) into Eq. (17) yields another expression of the nitracline depth:

$$z_n = \frac{1}{K_w} \ln \frac{I_0 (\mu_m / \alpha \varepsilon - 1)}{K_I} - \frac{K_c}{K_w} \int_0^{z_n} P dz \quad (18)$$

370 Note that Eq. (18) is obtained on the premise that the nitracline depth exists. This equality shows that the nitracline depth is inversely proportional to the light attenuation coefficient of water ( $K_w$ ), and it deepens logarithmically with increasing surface light intensity ( $I_0$ ). It is noted that the nitracline depth has a negative relation

with the self-shading of phytoplankton ( $K_c \int_0^{z_n} P dz$ ).

375 Importantly, Eq. (18) predicts that the nitracline depth has no relation with subsurface diffusivity. Aksnes et al. (2007) also proposed a similar result that a shoaling nitracline per se cannot be taken as an unequivocal sign of increased upwelling, as well as eddy diffusion. However, this does not mean that fluid dynamics are unimportant in shaping vertical distribution of nitrate.

380 Equation (18) also indicates that both a higher recycling rate ( $\alpha$ ) and a larger loss rate ( $\varepsilon$ ) lead to a shallower nitracline, while the enhanced maximum growth rate of phytoplankton ( $\mu_m$ ) moves the nitracline depth downward. Modeling results showed that the nitracline was shoaled by 24% (from 84 m upwards to 64 m) when both the recycling rate ( $\alpha$ , from 0.6 to 0.8) and the loss rate ( $\varepsilon$ , from 0.3 d<sup>-1</sup> to 0.4 d<sup>-1</sup>) were increased by 33%. Accordingly, the predicted nitracline depth from Eq. (18) varied 385 from 86 m to 71 m. Increasing  $\mu_m$  by 33% (from 0.9 d<sup>-1</sup> to 1.2 d<sup>-1</sup>) makes the simulated nitracline deepening slightly (from 84 m to 88 m), leading to the predicted nitracline depth changing from 86 m to 92 m. The experiments with varying parameter values indicate that the updated  $z_n$  (based on the model runs) matches well 390 the predicted  $z_n$  of Eq. (18).

### 3.2.2 Steepness of the nitracline

In steady state, integrating Eq. (2) from the nitracline depth  $z_n$  to the bottom boundary  $z_b$ , and considering the light limitation of phytoplankton growth below depth  $z_n$  (Eq. 15), we obtain the nitrate gradient below the surface mixed layer,

395 
$$\frac{dN}{dz} \Big|_{z_n} = \frac{dN}{dz} \Big|_{z_b} + \frac{1}{K_{v2}} \int_{z_n}^{z_b} (\alpha\varepsilon - \mu_m f(I)) \gamma P dz \quad (19)$$

This equality shows that the nitracline steepness enhances with increasing nitrate gradient at the bottom boundary ( $\frac{dN}{dz} \Big|_{z_b}$ ) which depends on the intensity of nitrate intrusion from below. The vertical diffusion negatively influences the nitracline



steepness. The modeled time-series distributions of nitrate gradients and diffusive  
 400 nitrate fluxes in the northern SCS and the upstream Kuroshio Current showed similar  
 results (Li et al., 2015). Beckmann and Hense (2007) conducted sensitivity analysis of  
 both vertical diffusivity and nutrient concentration at the bottom boundary to examine  
 the vertical phytoplankton and nutrient profiles in oligotrophic oceans and their  
 numerical results support the relations presented in Eq. (19).

### 405 3.3 Analytical solutions for SCM characteristics

Similar to methods used by Gong et al. (2015), the piecewise function for vertical  
 chlorophyll profile (Eq. 6) was incorporated into the nutrient-phytoplankton model  
 (Eqs. 1-2) at steady state to derive the three SCM characteristics (SCML thickness, its  
 depth and intensity).

410 For  $z = z_m$  and  $z = z_m + \sigma$ , the net growth rate of phytoplankton (Eq. 15) can be  
 respectively expressed as following:

$$\mu_m f(I)|_{z=z_m} - \varepsilon = K_{v2}/\sigma^2 \quad (20)$$

$$\mu_m f(I)|_{z=z_m+\sigma} - \varepsilon = -w/\sigma \quad (21)$$

By substituting the growth limitation function for light to Eq. (20) or Eq. (21), we  
 415 obtain the expression of parameter  $z_m$ , i.e.,

$$z_m = \frac{1}{K_w} \ln \left[ \left( \frac{\mu_m}{\varepsilon + K_{v2}/\sigma^2} - 1 \right) \frac{I_0}{K_l} \right] - \frac{K_c}{K_w} \int_0^{z_m} P dz \quad (22)$$

or

$$z_m = \frac{1}{K_w} \ln \left[ \left( \frac{\mu_m}{\varepsilon - w/\sigma} - 1 \right) \frac{I_0}{K_l} \right] - \frac{K_c}{K_w} \int_0^{z_m+\sigma} P dz - \sigma \quad (23)$$

Subtracting Eqs. (22) and (23), and rearranging, we obtain the expression of  
 420 parameter  $\sigma$ :

$$\left( \frac{\mu_m}{\mu_m - \varepsilon + w/\sigma} - 1 \right) \exp^{K_w \sigma + K_c \int_{z_m}^{z_m + \sigma} P dz} = \frac{\mu_m}{\mu_m - \varepsilon - K_{v2}/\sigma^2} - 1 \quad (24)$$

Neglecting terms including self-shading of phytoplankton ( $K_c$ ) in Eqs. (22-24), both the analytical solutions of the depth and thickness of SCML are the same as the results presented in Gong et al. (2015). The self-shading effect of phytoplankton plays an important role in vertical pattern of chlorophyll (Shigesada and Okubo, 1981; Klausmeier and Litchman, 2001; Beckmann and Hense, 2007). In line with common sense, our analytic results indicate that a higher self-shading of phytoplankton negatively influences the depth and thickness of the SCML (Eq. 22 and Eq. 24), having similarly effect as an increasing light attenuation coefficient of water,  $K_w$ .

The expression of the SCML intensity is also different from the results presented in Gong et al. (2015). Integrating Eq. (7) from the surface of water to the bottom of surface mixed layer ( $z_s$ ), and from the bottom of surface mixed layer to the base of our model domain ( $z_b$ ) respectively, gives:

$$(1 - \alpha) \varepsilon \gamma P_0 z_s = K_{v2} \frac{dN}{dz} \Big|_{z_s+0} + N_{inML} \quad (25)$$

$$(1 - \alpha) \varepsilon \gamma \lambda h = K_{v2} \frac{dN}{dz} \Big|_{z_b} - K_{v2} \frac{dN}{dz} \Big|_{z_s+0} \quad (26)$$

Parameter  $\lambda h$  ( $\text{mg}/\text{m}^2$ ) is assumed as the integrated chlorophyll concentration below the surface mixed layer. Based on the properties of Gaussian function,  $\lambda$  can be expressed as  $\lambda = \Phi\left(\frac{z_b - z_m}{\sigma}\right) - \Phi\left(\frac{z_s - z_m}{\sigma}\right)$ , where  $\Phi\left(\frac{z_b - z_m}{\sigma}\right)$  and  $\Phi\left(\frac{z_s - z_m}{\sigma}\right)$  can be obtained from the standard normal table. According to the property of Gaussian function, we have  $0.68 < \lambda < 1.00$ ,  $z_s < z < z_b$ , under the assumption of  $z_s < z_m - \sigma$ .

Adding Eqs. (25) and (26) yields:

$$(1 - \alpha) \varepsilon \gamma (\lambda h + P_0 z_s) = K_{v2} \frac{dN}{dz} \Big|_{z_b} + N_{inML} \quad (27)$$

Nitrogen input to the surface mixed layer ( $N_{inML}$ ) causes an increase of surface

chlorophyll concentration (Eq. 25). Hence, the total chlorophyll in stratified water  
 445 columns ( $\lambda h + P_0 z_s$ ) increases with increasing  $N_{inML}$  (Eq. 27), which has also been  
 predicted by the numerical study (Mellard et al., 2011) and supported by the  
 experimental test (Mellard et al., 2012).

Because recycling processes are assumed to not immediately convert dead  
 phytoplankton back into dissolved nutrients below the surface mixed layer, i.e.,  $\alpha \neq 1$ ,  
 450 the total chlorophyll concentration below the surface mixed layer and the intensity of  
 SCML can be respectively expressed as:

$$\lambda h = \frac{K_{v2}}{(1-\alpha)\varepsilon\gamma} \frac{dN}{dz} \Big|_{z_b} + \frac{N_{inML} - (1-\alpha)\varepsilon P_0 z_s}{(1-\alpha)\varepsilon\gamma} \quad (28)$$

$$P_{\max} = \frac{1}{\lambda\sqrt{2\pi}\sigma} \left( \frac{K_{v2}}{(1-\alpha)\varepsilon\gamma} \frac{dN}{dz} \Big|_{z_b} + \frac{N_{inML} - (1-\alpha)\varepsilon P_0 z_s}{(1-\alpha)\varepsilon\gamma} \right) \quad (29)$$

The integrated chlorophyll concentration below the surface mixed layer ( $\lambda h$ ) and the  
 455 intensity of SCML ( $P_{\max}$ ) are influenced by  $N_{inML}$  positively and by  $P_0$  negatively (Eqs.  
 28-29). That is to say, the influence of nitrate input to the surface mixed layer on the  
 SCML intensity (also on the integrated chlorophyll concentration below the surface  
 mixed layer) is non-linear. Hence, their changes ( $\lambda h$  and  $P_{\max}$ ) with varying  $N_{inML}$   
 cannot be obtained from the steady-state solutions straight forwardly, depending on  
 460 the specific parameter combinations in the model. For example,  $\lambda h$  and  $P_{\max}$  decrease  
 when increasing nutrient enrichment directly to the surface mixed layer in the  
 ecosystem given by Mellard et al. (2011), while they are nearly unchanged in  
 oligotrophic oceans (Varela et al., 1994).

Our results (Eqs. 28-29) also show that enhanced subsurface diffusivity ( $K_{v2}$ )  
 465 increases the integrated chlorophyll concentration and the intensity of the SCML ( $\lambda h$   
 and  $P_{\max}$ ), as a result of a higher nitrate flux ( $K_{v2}n$ ). Physical upward transport of  
 nitrate across the bottom of nitracline is indeed the main nitrogen source for NPP in  
 the euphotic zone (Ward et al., 1989).

## 4 Discussion

### 470 4.1 Light effects on nitracline with SCML

We now examine how the steady state nitracline in relate to SCML depends on light availability, especially light level at the nitracline depth.

Substituting Eq. (17) to Eq. (28) and rearranging, we have

$$\frac{K_{v2}}{\lambda h + P_0 z_s} \frac{dN}{dz} \Big|_{z_b} + \frac{N_{inML}}{\lambda h + P_0 z_s} = \varepsilon \gamma - \frac{\mu_m \gamma}{K_I / I(z_n) + 1} \quad (30)$$

475 This equality indicates that the light level at the nitracline depth,  $I(z_n)$ , is positively related to the integrated chlorophyll concentration in the whole water column,  $\lambda h + P_0 z_s$ . we can derive from Eq. (3) that the nitracline depth ( $z_n$ ) is inversely related to integrated chlorophyll. This inverse relationship has been observed in many regions. In southern California coastal waters, the phytoplankton standing stock and its  
480 primary production rate were positively related to the reciprocal nitracline depth (Eppley et al., 1978; Eppley et al., 1979). Bahamón et al. (2003) found that larger depth-integrated chlorophyll with an average deeper SCML and nitracline (~129m, ~136m, respectively) occurred in the Western Sargasso, Central Sargasso and Eastern Atlantic, compared with that in the Canary Current zone.

485 The nitracline depth deepens with increasing surface light intensity but with decreasing light attenuation coefficients ( $K_w$  and  $K_c$ ). These results were consistent with observations, e.g., Letelier et al. (2004) found the depth of the nitracline to coincide with an isolume, a depth of constant light level in the North Pacific Subtropical Gyre.

490 The predicted effect of surface light intensity and light attenuation coefficient on the nitracline depth (Eq. 18) implies that the nitracline depth in stratified waters may have seasonal variations. In the North Pacific Subtropical Gyre, Litelier et al. (2004) found that the nitracline depth differences between winter and summer disappeared when nitrate concentrations were plotted against light level in the water column.

495 Aksnes et al. (2007) found that the seasonal pattern of nitracline depth was governed by seasonality in light attenuation coefficient, rather than in surface light intensity. Particularly, the inverse proportional relationship between the nitracline depth and light attenuation coefficient (Eq. 18) has also been derived from a steady-state model  
500 region off Southern California (Aksnes et al., 2007). Tiera et al. (2005) found a significant positive correlation between the nitracline depth and the depth of 1% surface light intensity (the proportion of reciprocal light attenuation coefficient) in the Eastern North Atlantic Subtropical Gyre. Bahamón et al. (2003) showed that the nitracline depth remained relatively constant around 1% surface light intensity depth  
505 in Western Sargasso.

The nitracline steepens with a higher light attenuation coefficient ( $K_w$  and  $K_c$ ) due to  $K_w$  and  $K_c$  negatively influencing SCML thickness (Eqs. 14 and 25). Numerical modeling showed that a higher  $K_w$  leads to a thinner SCML and thus a steeper nitracline layer (Beckmann and Hense, 2007). Aksnes et al. (2007) also found that the  
510 fluctuations in the nitracline steepness were positively correlated with the fluctuations in reciprocal Secchi depth (i.e., light attenuation coefficient) in the upwelling area off the coast of the Southern California. We further point out that the nitracline steepness almost stays constant when changing surface light intensity ( $I_0$ ), because surface light intensity has no relation to the SCML thickness (Eq. 25). The sensitivity analysis of a  
515 one-dimensional (vertical) model showed that the vertical nutrient profiles were almost paralleling with each other when increasing surface light intensity (Beckmann and Hense, 2007).

The inverse effects of light attenuation coefficient on the nitracline steepness and its depth imply that the nitracline becomes steeper as the nitracline shoals. Aksnes et al.  
520 (2007) found this consistent pattern in the upwelling area off the coast of the Southern California.

#### *4.2 In presence of surface nutrient input*

Current evidences and modeling analyses suggest that climate warming will increase ocean stratification, and hence reduce nutrient exchange between the ocean interior and the upper mixed layer (Cermeno et al., 2008; Chavez et al., 2011). Therefore, nutrient input directly to the euphotic layer due to atmospheric deposition may become a relatively more important nutrient supply mechanism to the euphotic layer (Mackey et al., 2010; Okin et al., 2011; Mellard et al., 2011). However, few model studies (e.g., Mellard et al. 2011) have explored the influences of external surface nutrient supply on vertical phytoplankton distribution.

Observations show that an inter-zone exists between the transition of the surface mixed layer and the deep layer, where the nutrient gradient equals nearly zero  $\frac{dN}{dz}\Big|_{z_s+0} = 0$  (Fig. 3), leading to the solution in Eq. (26). It follows that the total chlorophyll in the surface mixed layer depends on the surface nutrient supply ( $N_{inML}$ ) (Eq. 26). In this case, if  $N_{inML}$  is negligible, Eq. (26) degenerates to

$$(1-\alpha)\varepsilon\gamma P_0 z_s = 0 \quad (31)$$

In this case, the dead phytoplankton in surface mixed layer must be fully recycled, i.e.,  $\alpha=1$ , in order to sustain the positive chlorophyll concentration ( $P_0>0$ ). In other words, if the dead phytoplankton cannot be fully recycled in the surface mixed layer, external nutrient supply to the layer is needed to fuel the growth of phytoplankton. Thus, the term, external nutrient supply to the surface mixed layer, should be included in the system equations at steady state to make a positive surface chlorophyll concentration. Numerical results by Mallard et al. (2011) also showed that phytoplankton populations can grow in the mixed layer and in the deep layer together, when there is nutrient input directly to the mixed layer. However, a surface nutrient source is not a necessary term for a model approach where dissolved organic matter and detritus are explicitly resolved (Beckmann and Hense, 2007).

Accordingly, we treat the vertical phytoplankton distribution as a piecewise function, comprised by a linear function in the surface mixed layer and a Gaussian

550 function below, which is more realistic than the general Gaussian function. The  
 assumption of the piecewise function for phytoplankton is also consistent with the  
 assumption of piecewise vertical diffusivity. For simplicity, we assume that the  
 transition layer between the surface mixed layer and the deep one is infinitely thin,  
 and the chlorophyll is continuous within the transition layer. By assuming the SCML  
 555 depth is significantly deeper than the base of the surface mixed layer, we obtain the  
 steady state solutions for the SCML depth and thickness, similar to the solutions using  
 the general Gaussian function. However, the intensity of the SCML is affected by  
 surface nutrient supply with an associated positive increase in phytoplankton  
 concentration.

#### 560 4.3 SCML trapping Nutrient

Indeed, observations and numerical simulations showed that SCML played a role as  
 a nutrient trap in some regions, restricting the diffusive flux of nitrates to the surface  
 mixed layer (Anderson, 1969; Klausmeier and Litchman, 2001; Navarro and Ruiz,  
 2013).

565 From Eq. (10), we know  $z_s < z_{n1} < z_0 - \sigma < z_m - \sigma$  (Fig. 1). That is, the SCML occurred  
 below depth  $z_{n1}$ . For  $z_m < z_n$  (Eq. 10), we know that the upward diffusive nitrate  
 concentration is enrichment for phytoplankton growth in the lower part of the SCML  
 ( $z_m < z < z_m + \sigma$ ). To explore the SCML restricting nitrates into the surface mixed layer,  
 next, we examine if the nitrate concentration at depth  $z_{n1}$  above the upper boundary of  
 570 the SCML ( $z_{n1} < z_m - \sigma$ ) is depleted.

According to the definition of the depth  $z_0$  (where  $f(I) = g(N)$  holds) and  $z_0 > z_{n1}$  (Eq.  
 10, Fig. 1), we know that the growth of phytoplankton at depth  $z_{n1}$  is nitrate-limited,  
 i.e.,  $\mu_m \min(f(I), g(N))|_{z_{n1}} = \mu_m g(N)|_{z_{n1}}$ . From Eq. (14), we get that at depth  $z_{n1}$ ,  
 the growth rate of phytoplankton equals the recycling rate of dead phytoplankton, i.e.,  
 575  $\mu_m g(N)|_{z_{n1}} = \alpha \varepsilon$ . Inserting the Michaelis-Menten form for  $g(N)$  into this equality  
 yields:  $N(z_{n1}) = K_N / (\mu_m / \alpha \varepsilon - 1)$ . Phytoplankton maximum growth rates ( $\mu_m$ ) of 0.2 to

1 per day are typical in optical environmental conditions (Banse, 1982; Timmermans et al., 2005). We choose 0.5 per day to illustrate the result. Loss rate ( $\varepsilon$ ), although not well documented, is often quoted as 10% of the growth rate (Parsons et al., 1984). A reasonable choice for the remineralization efficiency seems to be  $\alpha=0.5$  (Huisman et al., 2006). The typical value of half-saturation constants for nitrate ( $K_N$ ) is between 0.1 and 0.7 mmol N m<sup>-3</sup> in oceans (Eppley et al., 1969). We adopt 0.4 mmol N m<sup>-3</sup>. Thus, we obtain that the nitrate concentration at depth  $z_{n1}$ ,  $N(z_{n1})$ , is equal to 0.03 mmol N m<sup>-3</sup>, a value lower than the detection limit, indicating the depletion of nitrate above depth  $z_{n1}$ .

Because the SCML acts as a nutrient barrier, it is easy to understand that the rate of NPP in the SCML ( $(\mu_m \min(f(I), g(N)) - \alpha\varepsilon)P, z_m - \sigma < z < z_m + \sigma$ ) is positively related to upward nitrate flux that is trapped. This condition can simply be derived by integrating Eq. (2) vertically at steady state, i.e.,

$\int_{z_m - \sigma}^{z_m + \sigma} (\mu_m \min(f(I), g(N)) - \alpha\varepsilon) \gamma P dz = K_{v2} dN/dz \Big|_{z_m - \sigma}^{z_m + \sigma}$ . This result suggests that, the production within the SCML is fuelled mainly, by nitrate and is thus NPP. Because at the nitracline depth the gross growth rate  $\mu_m \min(f(I), g(N))$  equals the recycling of dead phytoplankton  $\alpha\varepsilon$ , for the constant  $\alpha\varepsilon$  we assumed, within the nitracline layer ( $z_{n1} < z_m - \sigma$  and  $z_m < z_{n2}$ ) the nitrate uptake by phytoplankton has to be supplied by the vertical diffusion. Observations also showed that most of the primary production in SCML was supported by nitrate from vertical diffusion, with an average  $f$ -ratio (i.e., relative contribution of the nitrate uptake to the total nitrogen uptake) of  $0.74 \pm 0.26$  during early summer in Canadian Arctic waters (Martin et al., 2012).

#### 4.4 Vertical profile of nitrate gradients

From the monotonicity of the quadratic function of depth  $z$  in the left-hand of Eq. (9), we know that  $d^2N/dz^2 < 0$  when  $z_s < z < z_{n1}$  and  $z_{n2} < z < z_b$ , but  $d^2N/dz^2 > 0$  when  $z_{n1} < z < z_{n2}$ . In other words, the vertical gradient of the nitrate concentration ( $dN/dz$ ) decreases with depth on the interval  $(z_s, z_{n1})$ , while increases on the interval  $(z_{n1}, z_{n2})$ , and then decreases on the interval  $(z_{n2}, z_b)$ . If we consider the distribution of vertical



605 nitrate gradients as continuous across the base of the surface mixed layer, then we get  $dN/dz < 0$  for  $z_s < z < z_{n1}$  under the assumptions of the uniform nitrate distribution within the surface mixed layer (i.e.,  $dN/dz = 0$ ,  $0 < z < z_s$ ). The schematic of vertical profiles of nitrate gradients and chlorophyll concentrations in stratified waters is shown in Fig. 1.

The negative gradient of nitrate below the surface mixed layer ( $dN/dz < 0$  for  
610  $z_s < z < z_{n1}$ ) indicates that the nitrate concentration decreases with depth on the interval  $(z_s, z_{n1})$ . This decreasing nitrate feature has rarely been observed by traditional measurements probably due to the technique-limited low resolution. Some float data showed this feature in vertical nitrate profiles, for example, Sakamoto et al. (2009) found it at depths below the base of surface mixed layer (~45-50 m) by the ISUS  
615 temperature-compensated data at an eastern Pacific oligotrophic station. Our in situ time series measurements using the ISUS at SEATS station also showed this decreasing feature at depths ~25-30 m (Fig. 4). We note that this decreasing nitrate feature will disappear in our derivation if the subsurface vertical diffusion is too weak (Eq. 12) or the surface mixed layer is deeper than depth  $z_{n1}$ . Simulating results showed  
620 that the negative gradient of nitrate became smaller with increasing the sinking velocity ( $w$ ) and the recycling rate ( $\alpha$ ). The finding implied that the negative gradient is likely the result of the phytoplankton eating holes in the nitrate distributions.

#### 4.5 Limitation and application

The model in this study integrates a number of physical, chemical, and biological  
625 processes that act together to determine the vertical distribution of phytoplankton and nitrate, under the assumption that the system is strictly vertical and in steady state. A few processes such as oxygen status, photoacclimation, luxury uptake of nutrients, phytoplankton motility, concentration-dependent-herbivory, and depth-dependent herbivory are not included, although they can affect the vertical distribution of  
630 phytoplankton and nitrate. Detritus, dissolved organic matter, and zooplankton are not included explicitly, and all loss processes, except sinking, are set to be linearly proportional to phytoplankton. The sinking velocity of phytoplankton is assumed independent on density gradients. Further, the vertical transport of nutrients is only by

eddy diffusion in our model; in reality, nutrients can be supplied by many processes  
635 (turbulence, internal waves, storms, slant-wise and vertical convection), especially by  
upwelling (Katsumi and Hitomi, 2003; Aksnes et al., 2007).

In this study, the sinking velocity of phytoplankton is set independent on nitrate  
concentration. Vertically-varying sinking velocity have been observed as  
physiological response to variations in light or nutrient levels (Steele and Yentsch,  
640 1960; Bienfang and Harrison, 1984; Richardson and Cullen, 1995). The sinking  
velocity reduced with decreased light level and with increased nutrient concentration,  
and the resulting divergence in sinking velocity can be large enough to affect the  
location of the phytoplankton particle maximum. However, numerical results given by  
Fennel and Boss (2003) showed analytically that the divergence of the sinking rate  
645 contributes to the location of the SBM layer in a significant way only when the  
divergence in sinking rates occurred above the compensation depth in stable,  
oligotrophic environments. They also derived that in stable, oligotrophic  
environments with a predominance of small cells, the biomass maximum is located at  
the depth where growth and losses are equal, leaving few influence by sinking  
650 divergence.

It is worth pointing out that, in extreme oligotrophic regions, the SCML is very  
deep and attributable mostly to photoacclimation of chlorophyll content rather than to  
a peak of biomass (Steele 1964; Fennel and Boss, 2003; Cullen, 2015). The process of  
photoacclimation is also important for the nutrient-phytoplankton system (with  
655 stratified conditions) we focused on. To explore the influence of photoacclimation on  
the nitracline, we parameterized Chl:C using the mathematical description by Cloern  
et al. (1995), i.e.,  $\text{Chl:C} = 0.003 + 0.0154 \exp(0.050T) \exp(-0.059I) \mu'$ . That is, Chl:C  
is the ratio of Chl a to C in phytoplankton growth at steady state under defined  
temperature  $T$  ( $^{\circ}\text{C}$ ), daily irradiance  $I$  ( $\text{mol quanta m}^{-2} \text{d}^{-1}$ ), and at nutrient-limited  
660 growth rate  $\mu'$  ( $\mu' = N/(K_N + N)$ ). The ratio Chl:C increases when temperature  $T$  or  
nitrogen concentration  $N$  increases, while decreases with increasing daily irradiance  $I$ .

Let  $R = \text{Chl:C}$ , then the nitrogen content of phytoplankton can be written as  $\gamma = 1/(6.625 \cdot 12 \cdot R)$ , corresponding to a C:N ratio of 6.625 and a carbon atomic mass of 12. From the expression of nitracline depth (Eq. 18), we know that the ratio Chl:C has  
665 no influence on the nitracline depth. While the nitracline steepness increases with increasing parameter  $\gamma$  (Eq. 19). In other words, the nitracline gets steeper with a lower ratio Chl:C. Note that a certainly more realistic model would be one with equations that explicitly resolve variations of the Chl a-to-carbon and nitrogen-to-carbon ratio of phytoplankton.

670 The piecewise equation (Eq.6) can be used to mimic a large variety of vertical chlorophyll profiles from coastal, upwelling, open oceans and high latitude waters (Fig. 2). For example, for  $z_s > 0$ , when the depth of surface mixed layer equals or is deeper than the depth of SCML, the vertical profiles like Fig. 2b and 2c are often found in well-mixed waters (Uitz et al., 2006). For  $z_s = 0$ , the vertical distribution of  
675 chlorophyll concentration (Fig. 2d) can be expressed by a Gaussian function, which is usually found in coastal upwelling waters (Xiu et al., 2008). Particularly, when  $z_s = 0$  and  $z_m = 0$ , the surface bloom occurs (Fig. 2e). In general, the vertical profiles of chlorophyll can be classified into two types, i.e., one peak distribution or uniform distribution in large regions of lakes and oceans (Uitz et al., 2006; Lavigne et al.,  
680 2015). Note that the skewed profiles of chlorophyll with a sharp SCM was not considered in this study. The small-scale (1 m or less) vertical heterogeneity in chlorophyll distribution has been shown to be common features in coastal waters (Sullivan et al., 2010; Prairie et al., 2011; Durham and Stocker, 2012), named as thin layer.

685 Choosing the values of model parameters represented the system in the northern SCS (given in Table 1), we can retrieve the nitracline depth and steepness, the optimal depth and the three SCM characteristics. To make calculation easy, we neglect the term of self-shading by phytoplankton in the calculation, because a higher self-shading parameter has the same effect as an increasing light attenuation  
690 coefficient by water. The calculated and observed values of these parameters are listed

at Table 2. All these parameters calculated are in a reasonable range, although there are some discrepancies compared with observations. In fact, this is not surprising, considering that we assume a single phytoplankton group and neglect the microbial loop and the dynamics of the dissolved organic matter and detritus pools.

695 We stress that the analytical solutions of nitracline are valid only for estimates of  $z_m$ ,  
 $h$ , and  $\sigma$  that are consistent with the model's numerical steady-state solution. The  
numerical steady-state solution in turn depends on the combination of parameter  
values and on the forcing, boundary conditions. The approximations of  $z_m$ ,  $h$ , and  $\sigma$   
are entirely conditioned by the modeling results and thus also depend on the  
700 combination of model parameter values. To combine the analytical steady state  
solutions with observed  $z_m$ ,  $h$ , and  $\sigma$  (as derived from vertical profiles of chlorophyll *a*  
concentration) is only meaningful after model calibration (identifying a model  
solution that is in some agreement with the observed  $z_m$ ,  $h$ , and  $\sigma$ ).

## 5 Summary

705 We have presented a theoretical framework to investigate the interaction of  
phytoplankton and nutrient in stratified water column. A piecewise function for  
chlorophyll profiles comprising a linear function in the surface mixed layer and a  
Gaussian function below is assumed in the nutrient-phytoplankton model at steady  
state. A number of important findings are obtained under conditions of the model  
710 equations imposed.

In steady state, the nitracline is confined between two depths where the gross growth rate equals the recycling rate of dead phytoplankton, indicating that within the nitracline, nitrate consumption by phytoplankton has to be replenished by the upward flux of nitrate. This layer thereby is the major contributor to new primary production.

715 The nitracline depth always locates below the SCML depth, meanwhile, both depths deepened as the relaxation of the light limitation (decreasing light attenuation coefficient or increasing surface light intensity). The nitracline depth does not depend on the value of the subsurface diffusivity. The nitracline is steeper with a thinner

720 SCML. The nitracline steepness is positively influenced by the light attenuation coefficient, yet, responds insignificantly to surface light intensity.

Our analytical solutions show that phytoplankton in the SCML acts as an efficient nutrient trap, filtering out the upward nitrate supply. The light level at the nitracline depth has a positive relation with the depth-integrated chlorophyll concentration in the whole water column and with the maximum rate of NPP, acting as the indicator of integrated NPP. The NPP is constructed from the model equations that rely in Blackman's law of limiting factor for the growth rate. These findings are based on the assumption that a prominent instant recycling process exists.

725


*Acknowledgements.* The authors thank State Key Laboratory of Marine Environmental Science, Xiamen University for providing ISUS nitrate and CTD data, especially acknowledge C. J. Du. We are very grateful to A. W. Omta and another anonymous reviewer for their constructive and helpful suggestions. We also would like to thank Xiaohuan Liu, Yang Yu, and Xiaokun Ding for valuable advice and programming assistance. This work is funded in part by the National Key Basic Research Program of China 2014CB953700, and the National Nature Science Foundation of China (41406010, 41210008, and 91328202).

730

735


Figures

Figure 1



740 Fig. 1 Schematic picture of vertical profiles of nitrate gradient and chlorophyll (Chl a) in stratified  
745 water columns. (blue solid line is the vertical profile of nitrate gradient; green solid line is Chl a  
concentration as a function of depth; red solid line represents the growth limitation by light, red  
dotted line by nitrate; horizontal green solid lines indicate the locations of the upper and lower  
SCML,  $z_m - \sigma$ ,  $z_m + \sigma$ , respectively; horizontal black dotted line indicates the depth of the surface  
mixed layer,  $z_s$ ; vertical dotted black line represents zero nitrate gradient;  $z_{n1}$  and  $z_n$  are the  
locations of extreme nitrate gradients,  $z_n$  is the nitracline depth, and  $z_m$  is the depth of maximum  
chlorophyll concentration)

Figure 2



750

Fig. 2 Examples of the vertical profiles of chlorophyll (black solid line). (red dotted lines represent the parts of Gaussian fitting curves, not the actual chlorophyll)

Figure 3

755




Fig. 3 Steady-state vertical distributions of chlorophyll, nitrate, and light determined by numerical solutions of Eqs. (1) and (2). Horizontal red dash-dotted line indicates the depth of the surface mixed layer. Black dash line represents the fitting curve of vertical chl a profile. The fitting

760 equation is 
$$P = \begin{cases} 0.013 & 0 \leq z \leq 30 \\ 0.33 \exp \left[ -\frac{(z-63)^2}{2 \times 9^2} \right] & 30 < z < 200 \end{cases}$$



Figure 4




Fig. 4 Vertical nitrate gradient, ISUS nitrate and temperature at SEATS station (2012, cast 36) (horizontal line indicates the depth of the surface mixed layer, horizontal dotted line indicates the depth of nitracline, and the vertical dash-dotted line represents zero nitrate gradient).

765

Table

Table 1 List of symbols and their values used in models at SEATS station in northern SCS

Model parameters	Description (unit)	Values (range)
$I_0$	Surface light intensity ( $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ )	900 (200-1700) <sup>(1, 2)</sup>
$K_w$	Light attenuation coefficient of water ( $\text{m}^{-1}$ )	0.052 <sup>(1, 3)</sup>
$K_c$	Light attenuation coefficient of phytoplankton ( $\text{m}^2 (\text{mmol N})^{-1}$ )	0.05 <sup>(1, 3)</sup>
$K_I$	Half-saturation constant of light limited growth ( $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ )	40 <sup>(4)</sup>
$K_{v1}$	Surface diffusivity ( $(\times 10^{-4}) \text{m}^2 \text{s}^{-1}$ )	2 <sup>(5)</sup>
$K_{v2}$	Subsurface diffusivity ( $(\times 10^{-5}) \text{m}^2 \text{s}^{-1}$ )	5 <sup>(5)</sup>
$w$	Sinking velocity of phytoplankton ( $\text{m d}^{-1}$ )	1 <sup>(6)</sup>
$\varepsilon$	Loss rate of phytoplankton ( $\text{d}^{-1}$ )	0.3 <sup>(7)</sup>
$\alpha$	Nutrient recycling coefficient (dimensionless)	0.6 <sup>(7)</sup>
$K_N$	Half-saturation constant of nutrient uptake ( $\text{mmol N m}^{-3}$ )	0.4 <sup>(8)</sup>
$\mu_m$	Maximum growth rate of phytoplankton ( $\text{d}^{-1}$ )	0.9 <sup>(5, 7)</sup>
$N_{inML}$	Mixed layer nitrate input ( $(\times 10^{-7}) \text{mmol N m}^{-2} \text{s}^{-1}$ )	4 <sup>(9, 10)</sup>
$\gamma$	Nitrate content of phytoplankton ( $\text{mmol N per mg Chl}$ )	1/1.59 <sup>(11, 12)</sup>
$\lambda$	Proportion of integrated chlorophyll below surface mixed layer	0.9
$z_s$	Depth of surface mixed layer ( $\text{m}$ )	30 (10-80) <sup>(13, 14)</sup>
$z_b$	Bottom boundary of model domain ( $\text{m}$ )	200
$\frac{dN}{dz} \Big _{z=z_b}$	Nitrate gradient at the bottom boundary of model domain ( $\text{mmol N m}^{-4}$ )	0.2 (0-0.2) <sup>(15, 16, 17)</sup>

Superscripts refer to the references that provide the source for the parameter value and the citations are as follows: <sup>(1)</sup><http://oceandata.sci.gsfc.nasa.gov/SeaWiFS/Mapped/Annual/9km/>;  
770 <sup>(2)</sup>Wu and Gao, 2011; <sup>(3)</sup>Lee Chen et al., 2005; <sup>(4)</sup>Raven and Richardson, 1986; <sup>(5)</sup>Lu et al., 2010;  
<sup>(6)</sup>Bienfang and Harrison, 1984; <sup>(7)</sup>Liu et al., 2007; <sup>(8)</sup>Eppley et al., 1969; <sup>(9)</sup>Kim et al., 2014;  
<sup>(10)</sup>Duce et al., 2008; <sup>(11)</sup>Cloern et al., 1995; <sup>(12)</sup>Oschlies, 2001; <sup>(13)</sup>Wong et al., 2002; <sup>(14)</sup>Tseng et  
al., 2005; <sup>(15)</sup>Chen et al., 2006; <sup>(16)</sup>Our observations; <sup>(17)</sup>Li et al., 2015.

Table 2 Estimated results and observed values at SEATS station

Variables	Estimated results	Observations
Nitracline depth (m)	86	20-90 <sup>(1, 2, 3)</sup>
Nitracline steepness ( $mmol\ N\ m^{-4}$ )	0.21	0.30-0.50 <sup>(3)</sup>
Depth of SCML (m)	70	10-75 <sup>(4, 5, 6)</sup>
Intensity of SCML ( $mg\ m^{-3}$ )	0.47	0.40-0.90 <sup>(4, 5, 6)</sup>
Thickness of SCML (m)	24	10-55 <sup>(4, 5, 6)</sup>

775 Superscripts refer to the references that provide the source for the parameter value and the citations are as follows: <sup>(1)</sup>Tseng et al., 2005; <sup>(2)</sup>Wong et al., 2007; <sup>(3)</sup>Our observations; <sup>(4)</sup>Chen et al., 2006; <sup>(5)</sup>Liu et al., 2002; <sup>(6)</sup>Liu et al., 2007.

## References:

- 780 Aksnes, D. L., Ohman, M. D., Pascal, R.: Optical effect on the nitracline in a coastal upwelling area, *Limnol Oceanogr*, 3, 1179-1187, 2007.
- Anderson, G. C.: Subsurface chlorophyll maximum in the northeast Pacific Ocean, *Limnol Oceanogr*, 14, 386-391, 1969.
- Ardyna, M., Babin, M., Gosselin, M., Devred, E., Bélanger, S., Matsuoka, A., Tremblay, J. É.:  
785 Parameterization of vertical chlorophyll a in the Arctic Ocean: impact of the subsurface chlorophyll maximum on regional, seasonal, and annual primary production estimates, *Biogeosciences*, 10, 4383-4404, 2013.
- Bahamón, N., Velásquez, Z., Cruzado, A.: Chlorophyll a and nitrogen flux in the tropical North Atlantic Ocean, *Deep Sea Research Part I*, 50, 1189-1203, 2003.
- 790 Bahamón, N., Cruzado, A.: Modelling nitrogen fluxes in oligotrophic environments: NW Mediterranean and NE Atlantic, *Ecol Model*, 163, 223-244, 2003.
- Banse, K.: Cell volumes, maximal growth rates of unicellular algae and ciliates, and the role of ciliates in the marine pelagial1, 2, *Limnol Oceanogr*, 27, 1059-1071, 1982.
- Beckmann, A., Hense, I.: Beneath the surface: Characteristics of oceanic ecosystems under weak mixing conditions-A theoretical investigation, *Prog Oceanogr*, 75, 771-796, 2007.
- 795 Bienfang, P. K., Harrison, P. J.: Sinking-rate response of natural assemblages of temperate and subtropical phytoplankton to nutrient depletion, *Mar Biol*, 83, 293-300, 1984.
- Cermeno, P., Dutkiewicz, S., Harris, R. P., Follows, M., Schofield, O., Falkowski, P. G.: The role of nutricline depth in regulating the ocean carbon cycle., *P. Natl Acad Sci Usa*, 105, 20344-20349, 2008.
- 800 Chavez, F. P., Messié, M., Pennington, J. T.: Marine primary production in relation to climate variability and change, *Annu Rev Mar Sci*, 3, 227-260, 2011.
- Chen, C. C., Shiah, F. K., Chung, S. W., Liu, K. K.: Winter phytoplankton blooms in the shallow mixed layer of the South China Sea enhanced by upwelling, *J. Marine Syst*, 59, 97-110, 2006.
- Cloern, J. E., Grenz, C., Videgar-Lucas, L.: An empirical model of the phytoplankton chlorophyll:  
805 carbon ratio-the conservation factor between productivity and growth rate, *Limnol. Oceanogr.*, 40, 1313-1321, 1995.
- Cullen, J. J.: The deep chlorophyll maximum: comparing vertical profiles of chlorophyll a, *Can J. Fish Aquat Sci*, 39, 791-803, 1982.
- Cullen, J. J., Eppley, R. W.: Chlorophyll maximum layers of the Southern California Bight and  
810 possible mechanisms of their formation and maintenance, *Oceanologica Acta*, 1, 23-32, 1981.
- Cullen, J. J.: Subsurface Chlorophyll Maximum Layers: Enduring Enigma or Mystery Solved? *Annu Rev Mar Sci*, 7, 207-239, 2015.
- Denman, K. L., Gargett, A. E.: Time and space scales of vertical mixing and advection of' phytoplankton in the upper ocean, *Limnol Oceanogr*, 28, 801-815, 1983.
- 815 Du, Y., Mei, L.: On a nonlocal reaction - diffusion - advection equation modelling phytoplankton dynamics, *Nonlinearity*, 24, 319, 2011.
- Du, Y., Hsu, S. B.: On a Nonlocal Reaction-Diffusion Problem Arising from the Modeling of Phytoplankton Growth., *Siam J.math.anal*, 42, 1305-1333, 2010.

- 820 Du, Y., Hsu, S. B.: Concentration Phenomena in a Nonlocal Quasi-linear Problem Modelling  
Phytoplankton I: Existence, *Siam J. Math Anal*, 40, 1419-1448, 2008a.
- Du, Y., Hsu, S. B.: Concentration phenomena in a nonlocal quasilinear problem modelling  
phytoplankton II: Limiting profile, *SIAM J. Math. Anal*, 40, 1441-1470, 2008b.
- 825 Duce, R. A., Laroche, J., Altieri, K., Arrigo, K. R., Baker, A. R., Capone, D. G., Cornell, S., Dentener,  
F., Galloway, J., Ganeshram, R. S.: Impacts of atmospheric anthropogenic nitrogen on the open ocean.,  
*Science*, 320, 893-897, 2008.
- Durham, W. M., Stocker, R.: Thin phytoplankton layers: characteristics, mechanisms, and  
consequences, *Annu Rev Mar Sci*, 4, 177, 2012.**
- Eppley, R. W., Sapienza, C., Renger, E. H.: Gradients in phytoplankton stocks and nutrients off  
southern California in 1974-76, *Estuarine and coastal marine science*, 7, 291-301, 1978.
- 830 Eppley, R. W., Renger, E. H., Harrison, W. G.: Nitrate and phytoplankton production in southern  
California coastal waters, *Limnol Oceanogr*, 24, 483-494, 1979.
- Eppley, R. W., Rogers, J. N., McCarthy, J. J.: Half-saturation constant for uptake of nitrate and  
ammonium by marine phytoplankton, *Limnol Oceanogr*, 14, 912-920, 1969.
- 835 Falkowski, P. G., Barber, R. T., Smetacek, V.: Biogeochemical controls and feedbacks on ocean  
primary production, *Science*, 281, 200, 1998.
- Fennel, K., Boss, E.: Subsurface maxima of phytoplankton and chlorophyll: Steady-state solutions  
from a simple model, *Limnology & Oceanography*, 48, 1521-1534, 2003.
- Fernand, L., Weston, K., Morris, T., Greenwood, N., Brown, J., Jickells, T.: The contribution of the  
deep chlorophyll maximum to primary production in a seasonally stratified shelf sea, the North Sea,  
840 *Biogeochemistry*, 1-14, 2013.
- Gong, X., Shi, J., Gao, H. W., Yao, X. H.: Steady-state solutions for subsurface chlorophyll maximum  
in stratified water columns with a bell-shaped vertical profile of chlorophyll, *Biogeosciences*, 12,  
905-919, 2015.
- Gong, X., Shi, J., Gao, H.: Modeling seasonal variations of subsurface chlorophyll maximum in South  
845 China Sea, *J. Ocean U. China*, 13, 561-571, 2014.
- Herbland, A., Voituriez, B.: Hydrological structure analysis for estimating the primary production in  
the tropical Atlantic Ocean, *J. Mar Res*, 37, 87-101, 1979.
- Hickman, A. E., Moore, C., Sharples, J., Lucas, M. I., Tilstone, G. H., Krivtsov, V., Holligan, P. M.:  
Primary production and nitrate uptake within the seasonal thermocline of a stratified shelf sea, *Mar*  
850 *Ecol Prog Ser*, 463, 39-57, 2012.
- Hodges, B. A., Rudnick, D. L.: Simple models of steady deep maxima in chlorophyll and biomass,  
*Deep Sea Research Part I: Oceanographic Research Papers*, 51, 999-1015, 2004.
- Hsu, S. B., Yuan, L.: Single Phytoplankton Species Growth with Light and Advection in a Water  
Column, *Siam J. Appl Math*, 70, 2942-2974, 2010.
- 855 Huisman, J., Thi, N., Karl, D. M., Sommeijer, B.: Reduced mixing generates oscillations and chaos in  
the oceanic deep chlorophyll maximum, *Nature*, 439, 322-325, 2006.
- Jamart, B. M., Winter, D. F., Banse, K., Anderson, G. C., Lam, R. K.: A theoretical study of  
phytoplankton growth and nutrient distribution in the Pacific Ocean off the northwestern US coast,  
*Deep-Sea Research*, 24, 753-773, 1977.
- 860 Jamart, B. M., Winter, D. F., Banse, K.: Sensitivity analysis of a mathematical model of phytoplankton

- growth and nutrient distribution in the Pacific Ocean off the northwestern US coast, *J. Plankton Res.*, 1, 267-290, 1979.
- Johnson, K. S., Coletti, L. J.: In situ ultraviolet spectrophotometry for high resolution and long-term monitoring of nitrate, bromide and bisulfide in the ocean, *Deep Sea Research Part I*, 49, 1291-1305, 2002.
- 865 Johnson, K. S., Riser, S. C., Karl, D. M.: Nitrate supply from deep to near-surface waters of the North Pacific subtropical gyre, *Nature*, 465, 1062-1065, 2010.
- Katsumi, H., Hitomi, K.: Vertical Nutrient Distributions in the Western North Pacific Ocean: Simple Model for Estimating Nutrient Upwelling, Export Flux and Consumption Rates, *J. Oceanogr.*, 59, 870 149-161, 2003.
- Kiefer, D. A., Kremer, J. N.: Origins of vertical patterns of phytoplankton and nutrients in the temperate, open ocean: a stratigraphic hypothesis, *Deep Sea Research Part A. Oceanographic Research Papers*, 28, 1087-1105, 1981.
- Kim, T. W., Lee, K., Duce, R., Liss, P.: Impact of atmospheric nitrogen deposition on phytoplankton productivity in the South China Sea, *Geophys Res Lett*, 41, 3156-3162, 2014.
- 875 Klausmeier, C. A., Litchman, E.: Algal games: The vertical distribution of phytoplankton in poorly mixed water columns, *Limnol Oceanogr.*, 8, 1998-2007, 2001.
- Koeve, W., Eppley, R. W., Podewski, S., Zeitzschel, B.: An unexpected nitrate distribution in the tropical North Atlantic at 18° N, 30° W—implications for new production, *Deep Sea Research Part II*, 880 40, 521-536, 1993.
- Laanemets, J., Kononen, K., Pavelson, J., Poutanen, E. L.: Vertical location of seasonal nutriclines in the western Gulf of Finland, *J. Marine Syst.*, 52, 1-13, 2004.
- Lande, R., Wood, A. M.: Suspension times of particles in the upper ocean, *Deep Sea Research Part A. Oceanographic Research Papers*, 34, 61-72, 1987.
- 885 Lande, R., Li, W. K. W., Horne, E. P. W., Wood, A. M.: Phytoplankton growth rates estimated from depth profiles of cell concentration and turbulent diffusion, *Deep Sea Research Part A. Oceanographic Research Papers*, 36, 1141-1159, 1989.
- Lavigne, H., D'Ortenzio, F., Ribera D'Alcalà, M., Claustre, H., Sauzède, R., Gacic, M.: On the vertical distribution of the chlorophyll a concentration in the Mediterranean Sea: a basin scale and seasonal approach, *Biogeosciences Discussions*, 12, 4139-4181, 2015.
- 890 Lee Chen, Y.: Spatial and seasonal variations of nitrate-based new production and primary production in the South China Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 52, 319-340, 2005.
- Letelier, R. M., Karl, D. M., Abbott, M. R., Bidigare, R. R.: Light driven seasonal patterns of chlorophyll and nitrate in the lower euphotic zone of the North Pacific Subtropical Gyre, *Limnol Oceanogr.*, 49, 508-519, 2004.
- 895 Lewis, M. R., Harrison, W. G., Oakey, N. S., Hebert, D., Platt, T.: Vertical nitrate fluxes in the oligotrophic ocean, *Science*, 234, 870-873, 1986.
- Lewis, M. R., Cullen, J. J., Platt, T.: Phytoplankton and thermal structure in the upper ocean: consequences of nonuniformity in chlorophyll profile, *J. Geophys Res.*, 88, 2565-2570, 1983.
- 900 Li, Q. P., Wang, Y., Dong, Y., Gan, J.: Modeling long - term change of planktonic ecosystems in the northern South China Sea and the upstream Kuroshio Current, *Journal of Geophysical Research Oceans*, 2015.

- Lipschultz, F., Bates, N. R., Carlson, C. A., Hansell, D. A.: New production in the Sargasso Sea: History and current status, *Global Biogeochem Cy*, 16, 1, 2002.
- 905 Liu, K. K., Chao, S. Y., Shaw, P. T., Gong, G. C., Chen, C. C., Tang, T. Y.: Monsoon-forced chlorophyll distribution and primary production in the South China Sea: observations and a numerical study, *Deep-Sea Res. Pt. I*, 49, 1387-1412, 2002.
- Liu, K. K., Chen, Y. J., Tseng, C. M., Lin, I. I., Liu, H. B., Snidvongs, A.: The significance of phytoplankton photo-adaptation and benthic-pelagic coupling to primary production in the South China Sea: Observations and numerical investigations, *Deep Sea Research Part II: Topical Studies in Oceanography*, 2007, 1546-1574, 2007.
- 910 Lu, Z., Gan, J., Dai, M., Cheung, A.: The influence of coastal upwelling and a river plume on the subsurface chlorophyll maximum over the shelf of the northeastern South China Sea, *J. Marine Syst*, 82, 35-46, 2010.
- 915 Lund-Hansen, L. C.: Subsurface chlorophyll maximum (SCM) location and extension in the water column as governed by a density interface in the strongly stratified Kattegat estuary, *Hydrobiologia*, 1-14, 2011.
- Mackey, K. R. M., van Dijken, G. L., Mazloom, S., Erhardt, A. M., Ryan, J., Arrigo, K. R., Paytan, A.: Influence of atmospheric nutrients on primary productivity in a coastal upwelling region, *Global Biogeochem Cy*, 24, B4027, 2010.
- 920 Martin, A. P., Pondaven, P.: On estimates for the vertical nitrate flux due to eddy pumping, *Journal of Geophysical Research: Oceans* (1978 - 2012), 108, 2003.
- Martin, J., Tremblay, J. É., Price, N. M.: Nutritive and photosynthetic ecology of subsurface chlorophyll maxima in Canadian Arctic waters, *Biogeosciences*, 9, 5353-5371, 2012.
- 925 Martin, J., Tremblay, J., Gagnon, J., Tremblay, G., Lapoussière, A., Jose, C., Poulin, M., Gosselin, M., Gratton, Y., Michel, C.: Prevalence, structure and properties of subsurface chlorophyll maxima in Canadian Arctic waters, *Mar Ecol Prog Ser*, 412, 69-84, 2010.
- Matsumura, S., Shiimoto, A.: Vertical distribution of primary productivity function F (II) for the estimation of primary productivity using by satellite remote sensing, *Bull. Nat. Res. Inst. Far Seas Fish*, 30, 227-270, 1993.
- 930 Mellard, J. P., Yoshiyama, K., Litchman, E., Klausmeier, C. A.: The vertical distribution of phytoplankton in stratified water columns, *J. Theor Biol*, 269, 16-30, 2011.
- Mellard, J. P., Yoshiyama, K., Klausmeier, C. A., Litchman, E.: Experimental test of phytoplankton competition for nutrients and light in poorly mixed water columns, *Ecol Monogr*, 82, 239-256, 2012.
- 935 Mignot, A., Claustre, H., D'Ortenzio, F., Xing, X., Poteau, A., Ras, J.: From the shape of the vertical profile of in vivo fluorescence to Chlorophyll-a concentration, *Biogeosciences*, 8, 2391-2406, 2011.
- Mu Oz Anderson, M., Hernández Walls, R., Rojas Mayoral, E., Galindo Bect, S.: Fitting vertical chlorophyll profiles in the California Current using two Gaussian curves, *Limnology & Oceanography Methods*, 2015.
- 940 Navarro, G., Ruiz, J.: Hysteresis conditions the vertical position of deep chlorophyll maximum in the temperate ocean, *Global Biogeochem Cy*, 1-10, 2013.
- Okin, G. S., Baker, A. R., Tegen, I., Mahowald, N. M., Dentener, F. J., Duce, R. A., Galloway, J. N., Hunter, K., Kanakidou, M., Kubilay, N., Prospero, J. M., Sarin, M., Surapipith, V., Uematsu, M., Zhu, T.: Impacts of atmospheric nutrient deposition on marine productivity: Roles of nitrogen, phosphorus,



- 945 and iron, *Global Biogeochem Cy*, 25, B2022, 2011.
- Omand, M. M., Mahadevan, A.: The shape of the oceanic nitracline, *Biogeosciences*, 12, 3273-3287, 2015.
- Oschlies, A.: Model-derived estimates of new production: New results point towards lower values, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 2173-2197, 2001.
- 950 Parsons, T. R., Maita, Y., Lalli, C. M., 1984. A manual of chemical and biological methods for seawater analysis, xiv + 173 pp. Oxford: Pergamon Press.
- Platt, T., Sathyendranath, S., Caverhill, C. M., Lewis, M. R.: Ocean primary production and available light: further algorithms for remote sensing, *Deep Sea Research Part A. Oceanographic Research Papers*, 35, 855-879, 1988.
- 955 **Prairie, J. C., Franks, P. J. S., Jaffe, J. S., Doubell, M. J., Yamazaki, H.: Physical and biological controls of vertical gradients in phytoplankton, *Limnology & Oceanography: Fluids & Environments*, 1, 75-90, 2011.**
- Probyn, T. A., Mitchell-Innes, B. A., Searson, S.: Primary productivity and nitrogen uptake in the subsurface chlorophyll maximum on the Eastern Agulhas Bank, *Cont Shelf Res*, 15, 1903-1920, 1995.
- 960 Richardson, T. L., Cullen, J. J.: Changes in buoyancy and chemical composition during growth of a coastal marine diatom: Ecological and biogeochemical consequences, *Marine Ecology Progress*, 128, 77-90, 1995.
- Riley, G. A., Stommel, H., Bumpus, D. F.: Quantitative ecology of the plankton of the western North Atlantic, *Bulletin Bingham Oceanographical Collection*, 12, 1-69, 1949.
- 965 Ross, O. N., Sharples, J.: Phytoplankton motility and the competition for nutrients in the thermocline, *Mar Ecol Prog Ser*, 347, 21-38, 2007.
- Ryabov, A. B., Rudolf, L., Blasius, B.: Vertical distribution and composition of phytoplankton under the influence of an upper mixed layer, *J. Theor Biol*, 263, 120-133, 2010.
- Sakamoto, C. M., Johnson, K. S., Coletti, L. J.: Improved algorithm for the computation of nitrate concentrations in seawater using an in situ ultraviolet spectrophotometer, *Limnology and Oceanography: Methods*, 7, 132-143, 2009.
- 970 Sharples, J., Moore, C. M., Rippeth, T. P., Holligan, P. M., Hydes, D. J., Fisher, N. R., Simpson, J. H.: Phytoplankton distribution and survival in the thermocline, *Limnol Oceanogr*, 46, 486-496, 2001.
- Shigesada, N., Okubo, A.: Analysis of the self-shading effect on algal vertical distribution in natural waters, *J. Math Biol*, 12, 311-326, 1981.
- 975 Steele, J. H., Yentsch, C. S.: The vertical distribution of chlorophyll, *Journal of the Marine Biological Association of the United Kingdom*, 39, 217-226, 1960.
- Steele, J. H.: A study of production in the Gulf of Mexico, *J. mar. Res*, 22, 211-222, 1964.
- Sullivan, J. M., Donaghay, P. L., Rines, J. E.: Coastal thin layer dynamics: consequences to biology and optics, *Cont Shelf Res*, 30, 50-65, 2010.**
- 980 Sverdrup, H. U.: On conditions for the vernal blooming of phytoplankton, *J. Cons. int. Explor. Mer*, 18, 287-295, 1953.
- Teira, E., Mourino, B., Marañón, E., Pérez, V., Pazo, M. J., Serret, P., de Armas, D., Escanez, J., Woodward, E., Fernández, E.: Variability of chlorophyll and primary production in the Eastern North Atlantic Subtropical Gyre: potential factors affecting phytoplankton activity, *Deep-Sea Research Part I*, 52, 569-588, 2005.
- 985

- Timmermans, K. R., Van der Wagt, B., Veldhuis, M., Maatman, A., De Baar, H.: Physiological responses of three species of marine pico-phytoplankton to ammonium, phosphate, iron and light limitation, *J. Sea Res*, 53, 109-120, 2005.
- 990 Uitz, J., Claustre, H., Morel, A., Hooker, S. B.: Vertical distribution of phytoplankton communities in open ocean: An assessment based on surface chlorophyll, *J. Geophys. Res*, 111, C8005, 2006.
- Varela, R. A., Cruzado, A., Tintoré, J.: A simulation analysis of various biological and physical factors influencing the deep-chlorophyll maximum structure in oligotrophic areas, *J. Marine Syst*, 5, 143-157, 1994.
- 995 Ward, B. B., Kilpatrick, K. A., Renger, E. H., Eppley, R. W.: Biological nitrogen cycling in the nitracline., *Limnol Oceanogr*, 34, 493-513, 1989.
- Wong, G. T. F., Tseng, C. M., Wen, L. S., Chung, S. W.: Nutrient dynamics and N-anomaly at the SEATS station, *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 1528-1545, 2007.
- Xiu, P., Liu, Y., Tang, J.: Variations of ocean colour parameters with nonuniform vertical profiles of chlorophyll concentration, *Int J. Remote Sens*, 29, 831-850, 2008.
- 1000 Yoshiyama, K., Nakajima, H.: Catastrophic transition in vertical distributions of phytoplankton: Alternative equilibria in a water column, *J. Theor Biol*, 216, 397-408, 2002.

**List of what we changed in the revised version of manuscript:**

1005

Original	Revised
Page 1, line 27-28	<b>Line 27-29:</b> Change “we derive analytical solutions for the system of phytoplankton and nutrient.” to “we derive analytical solutions of a specified nutrient-phytoplankton model. The model is well suited to explain basic dependencies between a nitracline and a SCML.”
Page 2, line 33-34	Delete the sentence “suggesting that the light level at the nitracline can be used as an indicator for integrated new primary production.”
Page 5, line 110-112	<b>Line 110-113:</b> Change “Accordingly, a piecewise function comprising a constant value within the surface mixed layer and a Gaussian function below this layer was used to fit vertical chlorophyll profiles.” To “Accordingly, a piecewise function comprising a constant value within the surface mixed layer and a Gaussian function below this layer was used as a fit to the steady state vertical chlorophyll profiles simulated by the nutrient-phytoplankton model.”
Page 10, after line 237	<b>Line 232-233:</b> Add a sentence “Thus, the specific (calibrated) model solution is considered as an example to obtain the analytic solutions of nitracline.”
Page 11, line 273-275	<b>Line 269-271:</b> Rewrite the definition of nitracline depth “In this study, we adopt the location of the maximum nitrate gradient ( $\max(dN/dz)$ ) in the euphotic zone as the nitracline depth ( $z_n$ ), which can be expressed by $\frac{d^2N}{dz^2}\Big _{z_n} = 0$ and $\frac{d^3N}{dz^3}\Big _{z_n} < 0$ .”

<p>Page 11, line 277-278</p>	<p><b>Line 273-275:</b> Change the sentence “Thus, at the nitracline depth the balance between uptake and recycling terms can be derived: <math>\mu_{\min}(f(I), g(N))=\alpha\varepsilon</math>.” To “Thus, according to our model approach (Eq. 2) the nitracline depth where <math>\frac{d^2N}{dz^2} = 0</math> represents a balance between the nutrient uptake and the recycling of phytoplankton loss, i.e., <math>\mu_{\min}(f(I), g(N))=\alpha\varepsilon</math>.”</p>
<p>Page 12, line 286-288</p>	<p><b>Line 283-286:</b> Add the explanation of ISUS data processing method “The sampling frequency was set at 5 Hz and the raw data were thus smoothed with a 25-point moving average in the surface mixed layer, a 5-point moving average in the SCML, and a 15-point moving average below the SCML. The data were then interpolated by a cubic spline function.”</p>
<p>Page 12, line 296-297</p>	<p>Spell out chlorophyll concentration <math>P</math> is a fitted and depth dependent variable, <b>Line 295-296:</b> “the fitted, depth dependent function of chlorophyll (<math>P(z)</math>, Eq. 6)”.</p>
<p>Page 12, line 299 and 301</p>	<p><b>Line 298 and 300:</b> Add the domain of Eqs. (8) and (9), i.e., “<math>z_s &lt; z &lt; z_b</math>”.</p>
<p>Page 13, line 310</p>	<p><b>Line 309:</b> Delete the minus sign in <math>-K_{v2}/\sigma^4</math>.</p>
<p>Page 14, line 335 and 339</p>	<p><b>Line 200:</b> Correct Eqs. (13) and (14) as <math>\frac{dN}{dz}\Big _{z_n} = \frac{dN}{dz}\Big _{z_b} + \left(\frac{z_n - z_m}{\sigma^2} + \frac{w}{K_{v2}}\right)\gamma P\Big _{z_n} - \frac{(1-\alpha)\varepsilon\gamma}{K_{v2}} \int_{z_n}^{z_b} P dz</math> and <math>\frac{dN}{dz}\Big _{z_n} = \frac{dN}{dz}\Big _{z_b} + \left(\sqrt{\frac{w^2}{4K_{v2}^2} + \frac{(1-\alpha)\varepsilon}{K_{v2}} + \frac{1}{\sigma^2}} + \frac{w}{2K_{v2}}\right)\gamma P\Big _{z_n} - \frac{(1-\alpha)\varepsilon\gamma}{K_{v2}} \int_{z_n}^{z_b} P dz</math></p>
<p>Page 15, line 359</p>	<p><b>Line 358-359:</b> Spell out the condition by Eq. (16) “Note that the derivation of Eq. (16) only works when light and nutrient limitation (Blackman’s law of limiting factors) is applied.”</p>

<p>Page 15, Line 367</p>	<p><b>Line 210-213 :</b> Correct the wrong typo of Eq. (18)</p> $z_n = \frac{1}{K_w} \ln \frac{I_0 (\mu_m / \alpha \varepsilon - 1)}{K_I} - \frac{K_c}{K_w} \int_0^{z_n} P dz$
<p>Page 16, After line 380</p>	<p><b>Line 382-390 :</b> Add numerical experiments with varying parameter values “Modeling results showed that the nitracline was shoaled by 24% (from 84 m upwards to 64 m) when both the recycling rate (<math>\alpha</math>, from 0.6 to 0.8) and the loss rate (<math>\varepsilon</math>, from 0.3 d<sup>-1</sup> to 0.4 d<sup>-1</sup>) were increased by 33%. Accordingly, the predicted nitracline depth from Eq. (18) varied from 86 m to 71 m. Increasing <math>\mu_m</math> by 33% (from 0.9 d<sup>-1</sup> to 1.2 d<sup>-1</sup>) makes the simulated nitracline deepening slightly (from 84 m to 88 m), leading to the predicted nitracline depth changing from 86 m to 92 m. The experiments with varying parameter values indicate that the updated <math>z_n</math> (based on the model runs) matches well the predicted <math>z_n</math> of Eq. (18).”</p>
<p>Page 17, line 416-419</p>	<p><b>Page 18, Line 426-429:</b> Rewrite the sentence “In line with common sense, our analytic results indicate that a higher self-shading of phytoplankton negatively influences the depth and thickness of the SCML (Eq. 22 and Eq. 24), having similarly effect as an increasing light attenuation coefficient of water, <math>K_w</math>.”</p>
<p>Page 18, after line 425</p>	<p>Move the introduction of <math>\lambda</math> from <b>Line 223-227</b> to <b>Line 436-440</b>.</p>
<p>Page 22, after line 535</p>	<p><b>Line 545-547:</b> Add the sentence “However, a surface nutrient source is not a necessary term for a model approach where dissolved organic matter and detritus are explicitly resolved (Beckmann and Hense, 2007).”</p>
<p>Page 25, line 607-608</p>	<p><b>Line 619-622:</b> Add the possible mechanism of negative vertical gradient in nitrate “Simulating results showed that the negative</p>

	<p>gradient of nitrate became smaller with increasing the sinking velocity (<math>w</math>) and the recycling rate (<math>\alpha</math>). The finding implied that the negative gradient is likely the result of the phytoplankton eating holes in the nitrate distributions.”</p>
<p>Page 25, after line 636</p>	<p><b>Line 651-669:</b> Add the influence of photoacclimation on nitracline “It is worth pointing out that, in extreme oligotrophic regions, the SCML is very deep and attributable mostly to photoacclimation of chlorophyll content rather than to a peak of biomass (Steele 1964; Fennel and Boss, 2003; Cullen, 2015). The process of photoacclimation is also important for the nutrient-phytoplankton system (with stratified conditions) we focused on. To explore the influence of photoacclimation on the nitracline, we parameterized Chl:C using the mathematical description by Cloern et al. (1995), i.e., <math>\text{Chl:C} = 0.003 + 0.0154 \exp(0.050T) \exp(-0.059I) \mu'</math>. That is, Chl:C is the ratio of Chl a to C in phytoplankton growth at steady state under defined temperature <math>T</math> (<math>^{\circ}\text{C}</math>), daily irradiance <math>I</math> (<math>\text{mol quanta m}^{-2} \text{d}^{-1}</math>), and at nutrient-limited growth rate <math>\mu'</math> (<math>\mu' = N/(K_N + N)</math>). The ratio Chl:C increases when temperature <math>T</math> or nitrogen concentration <math>N</math> increases, while decreases with increasing daily irradiance <math>I</math>. Let <math>R = \text{Chl:C}</math>, then the nitrogen content of phytoplankton can be written as <math>\gamma = 1/(6.625 * 12 * R)</math>, corresponding to a C:N ratio of 6.625 and a carbon atomic mass of 12. From the expression of nitracline depth (Eq. 18), we know that the ratio Chl:C has no influence on the nitracline depth. While the nitracline steepness increases with increasing parameter <math>\gamma</math> (Eq. 19). In other words, the nitracline gets steeper with a lower ratio Chl:C. Note that a certainly more realistic</p>

	<p>model would be one with equations that explicitly resolve variations of the Chl a-to-carbon and nitrogen-to-carbon ratio of phytoplankton.”</p>
<p>Page 26, after line 647</p>	<p><b>Line 680-683:</b> Spell out the neglect of the skewed profiles of chlorophyll with a sharp SCM in this study, “Note that the skewed profiles of chlorophyll with a sharp SCM was not considered in this study. The small-scale (1 m or less) vertical heterogeneity in chlorophyll distribution has been shown to be common features in coastal waters (Sullivan et al., 2010; Prairie et al., 2011), named as thin layer.”</p>
<p>Page 26, after line 657</p>	<p><b>Line 694-702:</b> Stress validity of the analytic solutions for nitracline, i.e., “We stress that the analytical solutions of nitracline are valid only for estimates of <math>z_m</math>, <math>h</math>, and <math>\sigma</math> that are consistent with the model’s numerical steady-state solution. The numerical steady-state solution in turn depends on the combination of parameter values and on the forcing, boundary conditions. The approximations of <math>z_m</math>, <math>h</math>, and <math>\sigma</math> are entirely conditioned by the modeling results and thus also depend on the combination of model parameter values. To combine the analytical steady state solutions with observed <math>z_m</math>, <math>h</math>, and <math>\sigma</math> (as derived from vertical profiles of chlorophyll <math>a</math> concentration) is only meaningful after model calibration (identifying a model solution that is in some agreement with the observed <math>z_m</math>, <math>h</math>, and <math>\sigma</math>).”</p>
<p>Page 26, line 663</p>	<p><b>Line 709-710:</b> Change the sentence “A number of important findings are obtained” to “A number of important findings are obtained under conditions of the model equations imposed.”</p>
<p>Page 27, line 665-678</p>	<p><b>Line 721-726:</b> Rewrite this sentence “The light level at the nitracline depth has a positive relation with the depth-integrated</p>

	<p>chlorophyll concentration in the whole water column and with the maximum rate of NPP, acting as the indicator of integrated NPP. The NPP is constructed from the model equations that rely in Blackman’s law of limiting factor for the growth rate. These findings are based on the assumption that a prominent instant recycling process exists.”</p>
Acknowledgements	<p><b>Line 729-732:</b> Add many thanks to the two referees and other three friends “We are very grateful to A. W. Omta and another anonymous reviewer for their constructive and helpful suggestions. We also would like to thank Xiaohuan Liu, Yang Yu, and Xiaokun Ding for valuable advice and programming assistance.”</p>
References	<p>Add 3 references in <b>line 826-827, 954-956, 978-979.</b></p>
Figures	<p>Redraw the profiles of ISUS Nitrate and Nitrate gradient by using new data processing method in Fig. 4.</p>
Tables	<p>Correct the wrong typo of <math>\gamma</math> in Table 1. Correct the estimated results in Table 2.</p>



***Interactive comment on “Analytical solution of nitracline with the evolution of subsurface chlorophyll maximum in stratified water columns” by Xiang Gong et al.***

A. W. Omta (Referee #1) [omta@mit.edu](mailto:omta@mit.edu)

*The manuscript is an analytical study of the relationship between the vertical distributions of phytoplankton and nutrients. An earlier paper by Gong et al. (2015) investigated the impact of light intensity, vertical diffusion, and the phytoplankton sinking velocity on the depth and width of the subsurface biomass maximum. Now, Gong et al. expand upon this earlier work with a careful study of what may determine the nutricline depth. The overall setup is good and there is a logical progression in the development of the text. Although analytical studies such as this one tend to be somewhat difficult to read, in my opinion they ought to have a much more prominent place in the field than they currently have, because they can provide much deeper insights than either (forward) numerical simulations or (inverse) parameter/state estimations. Having said all this, I think that at two points in the study, some further analysis is warranted before publication:*

**Response:** We are very grateful for the helpful comments and we have revised our manuscript accordingly.

*1) The authors admit that the assumption that the chlorophyll distribution represents the phytoplankton biomass distribution "is a significant simplification. In fact, phytoplankton increases inter-cellular pigment concentration when light level decreases (Cullen, 1982; Fennel and Boss, 2003; Cullen, 2015)." (p. 6, l. 129-131) Now, there happen to be fairly precise mathematical descriptions of this effect, e.g., Cloern et al. (1995). Thus, the authors ought to be able to investigate how and to which extent photoacclimation would impact their predictions regarding the relationship between the subsurface chlorophyll maximum and the nutricline depth.*

**Response:** Agree. We parameterize Chl: C using Eq. 15 of Cloern et al. Then let  $R = \text{Chl: C}$ , the nitrogen content of phytoplankton  $\gamma$  will be written as  $\gamma = 1/(6.625 \cdot 12 \cdot R)$ , corresponding to a C:N ratio of 6.625 and a carbon atomic mass of 12. The detailed results have been added in Section 4.5 to illustrate how and to which extent photoacclimation influence the relationships between a nitracline and a SCM.

*2) An unexpected prediction is the possible existence of nitrate minima below the surface mixed layer. According to the authors, these features disappear "if the subsurface vertical diffusion is too weak or the surface mixed layer is deeper than depth  $z_{n1}$ . The possible mechanism deserves to be explored." (p. 24/25, l. 606-608). I*

*think I may understand the origin of these remarkable features. Consider a situation without phytoplankton sinking and with full recycling of dead phytoplankton ( $w=0$ ,  $\alpha=1$ ). In that case, the nitrate distribution is simply the inverse of the phytoplankton distribution: if  $P$  has a maximum, then  $N$  has a minimum. When the sinking speed  $w$  increases and/or the recycling  $\alpha$  decreases, a background vertical  $N$  gradient develops which makes the  $N$  minimum shallower, until it has disappeared. Essentially, the  $N$  minima are then the result of the phytoplankton eating holes in the  $N$  distributions. All this is illustrated in the attached figure. In my view, it would be very interesting, if the authors would investigate this hypothesis by varying the sinking velocity and the recycling coefficient, starting from  $w=0$ ,  $\alpha=1$ .*

**Response:** Many thanks for this suggestion. Following this idea, we adopted numerical simulation to examine this hypothesis by varying the sinking velocity and the recycling coefficient. The results have been added in the revision.

**Interactive comment on “Analytical solution of nitracline with the evolution of subsurface chlorophyll maximum in stratified water columns” by Xiang Gong et al.**

*Anonymous Referee #2*

**5 1 General comments**

*The study of Xiang Gong and coauthors is concerned with the existence and characteristics of a nitracline in the presence of subsurface chlorophyll maxima (SCM). The authors derive analytical solutions that describe possible steady state results of a one-dimensional vertical model of nutrients (dissolved inorganic nitrogen) and phytoplankton biomass. Analytical steady state solutions are nicely derived for stratified conditions, with some weak mixing below a shallow upper mixed layer. A piecewise function is introduced as an approximation of the vertical distribution of phytoplankton biomass. This elegant approach was described and applied in an earlier study by X. Gong, J. Shi, H. W. Gao, and X.H. Yao, published 2015 in Biogeosciences, 12, 905-919. The authors take various different perspectives on the steady state solution. One of their main conclusions is that nitrate consumption by the phytoplankton has to be replenished by an upward flux of nitrate, which is interpreted as the major contribution to new primary production.*

*It is still fascinating to realise how much can be learned from analytical solutions of a model. X. Gong and his coauthors derive and explore steady state model solutions, elucidating interrelations between the characteristics of the nitracline and SCM. The stepwise derivation of particular solutions is generally good, but some readers may eventually lose track of all initial/original model assumptions. While reading about half of their study it became increasingly difficult to understand the actual meaning of the derived solutions, albeit mathematical steps were reproducible in most cases. For example, after the introduction of the depth of maximum growth ( $z_0$ ), many statements are made and conclusions are drawn that may lead readers astray. The authors tend to interpret their analytical solutions to be indicative for true conditions. But the solutions only reflect steady state conditions of model results. Furthermore, the authors give the impression that their analytical solutions are straightforward and can be used to make inference about nitracline features, once  $z_m$ ,  $h$ , and  $\sigma$  have been derived from observed profiles of chlorophyll a (Chl a) concentration. To do so would be inappropriate, which should be explicitly stated in the study. It is a conceptual problem that has to be reasonably addressed by the authors. Some major revision of the manuscript is therefore needed before the study can be recommended for publication in Biogeosciences.*

**Response:** Many thanks for the helpful suggestions and comments. We try best to revise and make physical meanings more obvious with those derived solutions. We also tone down the statements and conclusions to avoid any misleading. For example,  
40 we moved the statements drawn from Equation (17) (line 361-364) to the Discussion. The challenge and uncertainty have been included when we present those implications. Please see the revision.

*The analytical solutions presented are, apart from Equation (18) (see specific comments), correct. However, the author's should stress that the analytical solutions  
45 are valid only for estimates of  $z_m$ ,  $h$ , and  $\sigma$  that are consistent with the model's numerical steady state solution. The numerical steady state solution in turn depends on the forcing, boundary conditions and on the combination of parameter values. The approximations of  $z_m$ ,  $h$ , and  $\sigma$  are entirely conditioned by the model results and thus also depend on the combination of model parameter values. To combine the analytical  
50 steady state solutions with observed  $z_m$ ,  $h$ , and  $\sigma$  (as derived from vertical profiles of chlorophyll *a* concentration) is only meaningful after model calibration (identifying a model solution that is in some agreement with the observed  $z_m$ ,  $h$ , and  $\sigma$ ). A calibration requires the numerical model to be run in the first place. In other words, the equations, e.g. for the depth of the nitracline ( $z_n$ ), are valid only for  $z_m$ ,  $h$ , and  $\sigma$   
55 that remain dynamically consistent with the imposed model. Otherwise, the derived equations are not applicable.*

**Response:** Agree. We state that the analytical steady state solutions of nitracline are applicable only for estimates of  $z_m$ ,  $h$ , and  $\sigma$  that are consistent with the model's numerical steady state solution. The challenge and uncertainty have been included  
60 when we present those implications.

*Another concern is, although already addressed/discussed by the authors, the neglect of photoacclimation dynamics. The process of photoacclimation is essential for those systems (with stratified conditions) the authors focus on, and such a model approach would be better suited to make inference about the basic interrelations between a  
65 nitracline and a SCM. A possibility would be to include some additional parameterization that could yield variable  $\gamma$ , which can be derived from e.g. Cloern et al. (1995, L&O, 40(7), 1313-1321). When resorting to a parameterisation of Cloern et al. (e.g. their Eq. 15), some care has to be taken only with respect to the temporal integral of daily irradiance that is averaged over the upper mixed layer in their study.  
70 A certainly more realistic model would be one with equations that explicitly resolve variations of the Chl *a*-to-carbon and nitrogen-to-carbon ratio of the algae. An interesting aspect would be to see whether the "symmetric", piecewise Gaussian function would still be useful to approximate profiles of simulated Chl *a*, even if still*

75 *applicable to fit phytoplankton nitrogen biomass. The authors only discuss possible shifts in depth (location) of the SCM. They do not consider skewed profiles of Chl a, with a sharp SCM, as can be seen in many Chl a observational profiles.*

**Response:** In the revision, we parameterize Chl: C using Eq. 15 of Cloern *et al.* (1995). Then let  $R = \text{Chl: C}$ , the nitrogen content of phytoplankton  $\gamma$  will be written as  $\gamma = 1/(6.625 \cdot 12 \cdot R)$ , corresponding to a C:N ratio of 6.625 and a carbon atomic mass  
80 of 12. The detailed results will be added in Section 4.5 to illustrate how and to which extent photoacclimation influence the relationships between a nitracline and a SCM. The limitation, i.e., the skewed profiles of Chl a with a sharp SCM was not considered, have been added in Section 4.5.

## 2 Specific comments

### 85 Abstract

*lines 26-27: "..., we derive analytical solutions for the system of phytoplankton and nutrient."*

*The authors derive analytical solutions of a specified model. The model is well suited to explain basic dependencies between a nitracline and a deep chlorophyll a maximum.*  
90

**Response:** Agree. This sentence have been rewritten, i.e., we derive analytical solutions of a specified nutrient-phytoplankton model. The model is well suited to explain basic dependencies between a nitracline and a SCML.

*lines 31-34: "The inverse proportional relationship..., suggesting that the light level at the nitracline can be used as an indicator for integrated new primary production."*  
95

*It is not clear whether the model approach is appropriate to clearly distinguish between regenerated and new production. The dynamical model equations only resolve some instant remineralisation, with a direct mass flux from the phytoplankton back to the nutrient pool.*

100 **Response:** Agree. We delete this sentence and modify the related results in the text accordingly.

### 1 Introduction

*The introduction is nice. It is well written and informative.*

*line 112: "... was used to fit vertical chlorophyll profiles."*

105 *Here the authors should clarify that the Gaussian function is used as a fit to the steady state solution of the model.*

**Response:** Agree. We spell out that the Gaussian function is used as a fit to the steady state solution of the model. In reality, the Gaussian function is also applicable for

many profiles of Chl a in stratified waters, especially open ocean.

## 110 **2 Definition and models**

*pages 5 - 9: The model is nicely described and sufficient details are provided. I would suggest to introduce  $\lambda$  not here but where it is needed (on page 18).*

**Response:** Agree. We move the introduction of  $\lambda$  to page 18.

*page 9, lines 235 - 237: “We use the biologically reasonable parameter values given in Table 1 to represent the system at station SEATS...”*

*Thus, a specific (calibrated) model solution is considered as an example.*

**Response:** Agree. We add this sentence in the revision.

## pages 10 - 11: **Definition of the nitracline**

*The text is well written. The concept described in the final paragraph (lines 270 - 280) is clear. However, it is still confusing because simulated as well as observed profiles of  $N$  yield  $\frac{d^2N}{dz^2} \approx 0$  (or  $\frac{dN}{dz} \approx \text{constant}$ ) over some distinct depth range, e.g. as depicted in Fig. (2).*

**Response:** Many thanks for noticing this issue. The depth of nitracline in our study was defined as the location of maximum nitrate gradient in the euphotic zone, which can be expressed by  $\frac{d^2N}{dz^2} = 0$  and  $\frac{d^3N}{dz^3} < 0$ , not implying  $\frac{dN}{dz} = \text{constant}$ . The equality  $\frac{d^2N}{dz^2} = 0$  means  $\frac{dN}{dz} \approx \text{constant}$  only when  $\frac{d^2N}{dz^2} \equiv 0$  in the domain. Thus, to determine the depth of nitracline from simulated as well as observed profiles, we have to plot the profile of nitrate gradient and find the maximum nitrate gradient ( $\left. \frac{dN}{dz} \right|_{\max}$ ). We have rewritten this sentence in the revision.

*The described balance between uptake and recycling only works for this particular kind of model approach. The authors may add “According to our model approach (Eq. 2) the depth where  $\frac{d^2N}{dz^2}=0$  represents a balance between the growth rate and the phytoplankton loss rate.”*

**Response:** Agree. This has been added in the revision.

## 135 **3 Results**

*page 12: You may add here the depth range that is considered ( $z_s < z < z_b$ ).*

**Response:** Agree. The depth range  $z_s < z < z_b$  has been added in Eq. (8).

line 112: "... the fitted function of chlorophyll..."

140 *Suggestion: "... the fitted, depth depend function of chlorophyll ( $\gamma P(z)$ ). This reminds the reader that  $P$  actually includes an exponential in Eq. (8).*

**Response:** Agree. This sentence has been revised as "... the fitted, depth dependent function of chlorophyll ( $P(z)$ )..."

page 13: The minus sign ( $-K_{v2}/\sigma^4$ ) is confusing.

**Response:** We have delete the minus sigh in the revision.

145 *line 313: do the authors mean... "... (values from 8.64 to  $7.78 \cdot 10^{-9} \text{ m}^{-2} \text{ s}^{-1}$ )..."?*

**Response:** Because  $K_{v2}$  is  $1.9 \cdot 10^{-9} \text{ m}^2 \text{ s}^{-1}$ ,  $\sigma$  is from several meters to tens of meters, thus the ratio of  $K_{v2}$  to  $\sigma^4$  is from  $8.64 \cdot 10^{-9}$  to  $7.78 \cdot \text{m}^{-2} \text{ s}^{-1}$ . This will be clarified in the revision.

Equation (12): for non-zero  $w$ .

150 **Response:** We add the condition of non-zero  $w$  for Eq. (12) in the revision.

### **Depth of the nitracline**

page 15, lines 353 - 364: *This derivation only works when Blackman's law of limiting factors (light and nutrient limitation) is applied. Hence, it is a particular model assumption. The maximum rate discussed here first of all represents a net primary production term. Only in the context of this particular model version it is also interpreted as new primary production. The sentence "It follows that the light level at the nitracline is an indicator of integrated NPP in the water column." is a strong statement. This finding strongly depends on the underlying model equations. It would be good to see different steady state solutions of the model while varying values of  $\epsilon$  and  $\alpha$  (e.g. increasing  $\epsilon$  while decreasing  $\alpha$  and vice versa). This way the authors may substantiate their conclusion.*

155  
160

**Response:** We spell out that the derivation only works when Blackman's law of limiting factors (light and nutrient limitation) is applied in the revision. We will also examine the simulated results by varying values of  $\epsilon$  and  $\alpha$ , please see the revision.

165 *Equation (18): The inclusion of  $\gamma$  in the last term is incorrect. The parameter  $\gamma$  can be removed. This is because  $K_c$  is normalised to nitrogen biomass and not to Chl  $a$ .*

**Response:** Sorry for the typo, we have removed " $\gamma$ ".

page 16, lines 377 - 380: *"Equation (18) also indicates that both a higher recycling rate ( $\alpha$ ) of dead phytoplankton and a larger loss rate ( $\epsilon$ ) lead to a shallower nitracline, while the enhanced maximum growth rate of the phytoplankton ( $\mu_m$ ) moves*

170

*the nitracline depth down.”*

*It would be good to see this conclusion consolidated by some model results. This way the authors can also demonstrate the predictive power of applying Eq. (18). The parameters could be varied just as discussed by the authors and it would be interesting to see how well an updated  $z_n$  (based on the model runs with the parameter values varied) matches the predicted  $z_n$  of Eq. (18) (based on the previous model results, e.g. of P).*

**Response:** Thank you for this suggestion. We run the N-P model to examine how well the modelled  $z_n$  matches the predicted value and the results have been added in the revision.

*page 17, lines 416 - 419: “Our results indicate... self-shading negatively influences depth and thickness of the SCML,...”*

*This is comprehensible.*

**Response:** We rewrite this sentence.

## 185 **4 Discussion**

### **In presence of surface nutrient input**

*page 22, lines 527 - 535: This is certainly the case for the model assumption of an instant remineralisation of organic matter that originates directly from the phytoplankton.*

190 *Must this (the need to include a surface nutrient source) also be expected for a model approach where dissolved organic matter (DOM) and detritus are explicitly resolved?*

**Response:** After examining the N-P-D model given by Beckmann and Hense (2007), we found that the surface nutrient input is not necessary for a model approach where dissolved organic matter (DOM) and detritus are explicitly resolved. In the revision, we will spell out this assumption.

### **Vertical profiles of nitrate gradients**

*page 24, line 605: In Fig. (4) the profile of  $\frac{dN}{dz} \cdot 20$  does not correspond with the shown profile of N. The N profile clearly indicates a constant  $\frac{dN}{dz}$  (of approximately  $0.38 \text{ mmol N m}^{-4} \rightarrow 7.6 \text{ mmol N m}^{-4} = \frac{dN}{dz} \cdot 20$ ) in the depth range of 50 -70 m. The shown  $\frac{dN}{dz} \cdot 20$  does not reveal this feature. The authors need to clarify this.*

**Response:** Thank your comments. We redraw this figure by using new data



processing method in the revision.

### **Limitation and application**

205 *page 24, lines 610- 636: As important as the model assumptions for the sinking and remineralisation of particulate organic matter is photoacclimation. The authors should consider to include one or two figures with profiles of Chl a concentrations with typical but different shapes of the SCM.*

**Response:** In the revision, we spell out the limitation of photoacclimation in this Section and revise the manuscript accordingly.

210 *page 26, lines 648 - 657: I used the parameter values of Table (1) and the values for  $z_m$ ,  $h$ , and  $\sigma$  from Table (2) to calculate the corresponding  $z_n$  and  $\frac{dN}{dz}$  (Eqs. 11 and 14).*

*I obtain  $z_n=70$  m and  $\frac{dN}{dz} = 0.025$  mmol N m<sup>-4</sup>. In Table the solutions are  $z_n=79$  m and  $\frac{dN}{dz} = 0.24$  mmol N m<sup>-4</sup>. I cross-checked my equations and all values and have not found any explanation for this discrepancy. I thought that all values presented are*  
215 *consistent with the imposed model dynamics and thus valid for any of the analytical steady state solutions presented.*

**Response:** Sorry for the typo. We found that the value of  $\gamma$  should be 1/1.59, not 1.59. We will recalculate the values. Please see the revision.

### **Summary**

220 *pages 26 - 27: The authors may here stress that the important findings are conditioned by the model equations imposed. The interpretation of NPP is not straightforward and becomes particularly difficult to specify under steady state conditions of a weakly mixed water column. The authors construct NPP from the model equations that rely in Blackman's law of limiting factor for the growth rate. I*  
225 *suggest to the authors to refine their statements, clarifying their findings are based on the assumption that a prominent instant recycling process exists.*

**Response:** Thank you for the helpful suggestions. We rewrite the summary by stressing the assumptions of our model results, especially the statements about NPP.

230