



1 **Physiological response of a golden tide alga (*Sargassum muticum*) to the**
2 **interaction of ocean acidification and phosphorus enrichment**

3

4 Zhiguang Xu^a, Guang Gao^{b,c*}, Juntian Xu^b, Hongyan Wu^d

5

6

7 ^aMarine Biology Institute of Shandong Province, Qingdao 266104, China

8 ^bMarine Resources Development Institute of Jiangsu, Huaihai Institute of Technology,

9 Lianyungang, 222005, China

10 ^cSchool of Marine Science and Technology, Ridley Building, Newcastle University,

11 Newcastle upon Tyne, NE1 7RU, England, UK

12 ^dHubei University of Technology, Wuhan, 430068, China

13

14 *Corresponding author, g.gao@ncl.ac.uk, Phone/Fax: +44(0)1912085048



15 Abstract

16 The evolvement of golden tides would be influenced by global change factors,
17 such as ocean acidification and eutrophication, but the related studies are very scarce.
18 In this study, we cultured a golden tide alga, *Sargassum muticum*, at two levels of
19 $p\text{CO}_2$ (400, 1000 μatm) and phosphate (0.5 μM , 40 μM) conditions to investigate the
20 interactive effects of elevated $p\text{CO}_2$ and phosphate on physiological properties of the
21 thalli. The higher $p\text{CO}_2$ level and phosphate (P) level alone increased the relative
22 growth rate by 40.82% and 47.78%, net photosynthetic rate by 46.34% and 55.16%,
23 soluble carbohydrates by 32.78% and 61.83% respectively whilst the combination of
24 these two levels did not promote growth or soluble carbohydrates further. The higher
25 levels of $p\text{CO}_2$ and P alone also enhanced the nitrate uptake rate by 68.27% and
26 35.89%, nitrate reductase activity by 89.08% and 39.31%, and soluble protein by
27 19.05% and 15.13% respectively. The nitrate uptake rate and soluble protein was
28 further enhanced although the nitrate reductase activity was reduced when the higher
29 levels of $p\text{CO}_2$ and P worked together. The higher $p\text{CO}_2$ level and higher P level alone
30 did not affect the dark respiration rate of thalli but they together increased it by 32.30%
31 compared to the condition of the lower $p\text{CO}_2$ and lower P. The mute effect of the
32 higher level of $p\text{CO}_2$ and higher P on growth, soluble carbohydrates, combined with
33 the promoting effect of it on soluble protein and dark respiration, suggests more
34 energy was drawn from carbon assimilation to nitrogen assimilation at the condition
35 of higher $p\text{CO}_2$ and higher P, probably to act against the higher $p\text{CO}_2$ caused
36 acid-base perturbation via synthesizing H^+ transport-related protein. Our results
37 indicate ocean acidification and eutrophication may not boost the gold tides events
38 synergistically although each of them alone has a promoting effect.

39 Key words: carbohydrates, growth, photosynthesis, protein, respiration, *Sargassum*
40 *muticum*

41 1. Introduction

42 *Sargassum* C. Agardh (1820) is the most species-rich genus in the Phaeophyta
43 and has a global distribution (Mattio and Payri, 2011). The species of this genus
44 constitutes an important part of the marine flora and is considered as a valuable and



45 unique habitat for a number of highly adapted marine animal species (Laffoley et al.,
46 2011). Some species of *Sargassum* are economically important, being used as animal
47 fodder, manure in agriculture, as well as alginates production (Ashok-Kumar et al.,
48 2012; Fenoradosa et al., 2010; González-López et al., 2012). On the other hand,
49 *Sargassum* is an aggressive genus and it can rapidly spread and invade new areas
50 (Sfriso and Facca, 2013). The invasion of *Sargassum* would accordingly compete
51 with indigenous species for nutrients and light and lead to the alteration of macroalgal
52 community structure (Rueness, 1989; Stæhr et al., 2000). For instance, the increased
53 abundance of *S. muticum* in Limfjorden (Denmark) between 1990 and 1997 led to
54 decreased cover of several indigenous species belonging to the genera of *Codium*,
55 *Fucus*, and *Laminaria*, and thus reduced species richness and diversity of the
56 macroalgal community (Stæhr et al., 2000). Recently, the species of *Sargassum*
57 inundate the coasts along Gulf of Mexico, West African, Caribbean, and Brazil in
58 unprecedented biomass, termed as golden tides (Schell et al., 2015; Smetacek and
59 Zingone, 2013). Apart from the negative effect on aesthetics and tourism, the
60 occurrence of golden tides could kill the fish within the algal mass, mainly due to
61 hypoxia or anoxia in the waters caused by decomposition of *Sargassum* thalli
62 (Cruzrivera et al., 2015). In addition, the dense *Sargassum* accumulation could clog
63 fishing nets and impede the passage of boats, leading to food shortages for local
64 people who live on artisanal fisheries (Smetacek and Zingone, 2013). The occurrence
65 of golden tides has been linked to higher nutrient levels in the seawaters (Lapointe,
66 1995; Smetacek and Zingone, 2013). The distribution pattern and biomass of
67 *Sargassum* spp. are environment (temperature, light, nutrients, etc.)-dependent (Ang,
68 2006; Sfriso and Facca, 2013).

69 Due to burning fossil fuels and changes to land use, the atmospheric
70 concentrations of carbon dioxide have increased to the level of 401.72 ppm in July
71 2016 (<http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html>), which is
72 unprecedentedly high in at least the last 800,000 years (IPCC, 2013). When CO₂
73 dissolves in seawater it forms carbonic acid and as more CO₂ is taken up by the
74 ocean's surface, the pH decreases, moving towards a less alkaline and therefore more



75 acidic state, termed ocean acidification. The mean surface ocean pH has already
76 decreased by 0.1 units since the beginning of the industrial era, corresponding to a 26%
77 increase in hydrogen ion concentration (IPCC, 2013). By 2100, concentrations of CO₂
78 (aq) and HCO₃⁻ are predicted to increase by 192% and 14%, respectively, and CO₃²⁻
79 to decrease by 56%, with a concomitant decline in pH to 7.65 (Raven et al., 2005).
80 Increased CO₂ could exert positive, neutral, or negative on physiological properties of
81 macroalgae (Ji et al., 2016; Wu et al., 2008). In terms of *Sargassum* species, increased
82 CO₂ (800 ppm) enhanced photosynthetic rate (based on CO₂ uptake) in *S. muticum*
83 (Longphuir et al., 2014). On the other side, the same level of increased CO₂ (750
84 ppm) did not affect growth, Rubisco's maximal activity, affinity for CO₂ or quantity
85 in *S. vulgare* (Alvaro and Mazal, 2002). Furthermore, increased CO₂ (750 ppm)
86 significantly decreased net photosynthetic rate and light saturation point of *S.*
87 *henslowianum* (Chen and Zou, 2014).

88 Apart from ocean acidification, eutrophication is another environmental challenge.
89 Eutrophication can occur naturally in lakes via transferring nutrients from the
90 sediment to water by living or decomposing macrophytes, resuspension, diffusion,
91 and bioturbation (Carpenter, 1981). However, anthropogenic activities have
92 accelerated the rate and extent of eutrophication (Carpenter et al., 1998). Inevitable
93 urbanization of a growing human population, increased use of coastal areas, and rising
94 fertilizer use for agricultural intensification has led to accelerated nutrient inputs from
95 land-water to coastal waters (Smith et al., 1999). These changes in nutrient
96 availability result in eutrophication, an increasing threat for coastal ecosystems
97 (Bricker et al., 2008). One consequence of eutrophication is that it can lead to algal
98 bloom, such as green tides and golden tides (Smetacek and Zingone, 2013). There are
99 relatively intensive studies regarding the effect of nutrients on physiological
100 properties of *Sargassum* species (Hwang et al., 2004; Incera et al., 2009; Lapointe,
101 1995; Liu and Tan, 2014; Nakahara, 1990). Enrichment of nutrients usually can
102 enhance the growth and photosynthetic parameters of *Sargassum*. For instance, the
103 growth rate of *S. baccularia* almost doubled when nutrients increased from 3 μM
104 ammonium plus 0.3 μM phosphate to 5 μM ammonium plus 0.5 μM phosphate



105 (Schaffelke and Klumpp, 1998) and the photosynthetic rates of *S. fluitans* and *S.*
106 *natans* were also two-fold higher with 0.2 mM PO₃⁻ enrichment compared to the
107 control (Lapointe, 1986). Furthermore, some studies have demonstrated that
108 macroalgae experience more phosphorus limit instead of nitrogen limit (Lapointe,
109 1986; Lapointe et al., 1987, 1992; Littler et al., 1991). For instance, nitrogen
110 enrichment did not affect growth rates of *S. fluitans* or *S. natans* whilst phosphorus
111 enrichment increased them from 0.03–0.04 (control) to 0.05–0.08 doublings d⁻¹
112 (Lapointe, 1986).

113 Neither ocean acidification nor eutrophication is proceeding in isolation; rather
114 they occur simultaneously, particularly in coastal areas. The interactive effects of two
115 factors may be completely different, or be of greater magnitude, compared to effects
116 of any single stressor. To the best of our knowledge, no studies have been reported in
117 regard to the interactive effects of ocean acidification and eutrophication on
118 *Sargassum*. In this study, we chose the species *S. muticum* to investigate its responses
119 to interaction of ocean acidification and eutrophication. *S. muticum* is an invasive
120 macroalga and commonly habitats on rocky shores (Karlsson and Loo, 1999). It
121 originates from Japan and was introduced to the northern Pacific coast of the United
122 States in the early 20th century (Scagel, 1956), and was also introduced to Europe
123 along with the imported Japanese oyster in the late 1960s (Jones and Farnham., 1973).
124 Nowadays, its distribution is worldwide due to the introduction and the subsequent
125 rapid expansion (Cheang et al., 2010). Our study would supply insight into how ocean
126 acidification and eutrophication affect the physiological properties of *S. muticum* and
127 thus the evolution of golden tides.

128 **2. Materials and methods**

129 *2.1. Sample collection and experiment design*

130 *S. muticum* was collected from lower intertidal rocks on the coast of Lidao,
131 Rongcheng, China (37 °15'N, 122 °35'E). The samples were transported to the
132 laboratory in an insulated polystyrene cooler (4–6 °C) within 3 hours. Healthy thalli
133 were selected and rinsed with sterile seawater to remove sediments, epiphytes and
134 small grazers. The thalli were maintained in an intelligent illumination incubator



135 (MGC-250P, Yiheng Technical Co. Ltd., Shanghai, China) for 24 hours before the
136 experiment. The temperature in the incubator was set as 20°C with a 12h: 12h
137 (light/dark) photoperiod of 150 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ photosynthetically active
138 radiation (PAR). After the maintenance, a two-way factorial experiment was set up to
139 investigate the interactive effects of $p\text{CO}_2$ and phosphate on *S. muticum*. The thalli
140 were placed in 3 L flasks with 2 L sterile seawater (one thallus per flask) and cultured
141 at fully crossed two $p\text{CO}_2$ (400 μatm , LC; 1000 μatm , HC) and two phosphate (0.5
142 μM , LP; 40 μM , HP) levels with continuous aeration for 13 days. Phosphorus was
143 selected as a nutrient variable since some findings have displayed that phosphorus,
144 rather than nitrogen, is the primary limiting nutrient for macroalgae (Lapointe, 1986;
145 Lapointe et al., 1987, 1992; Littler et al., 1991). The 400 $\mu\text{atm } p\text{CO}_2$ and 0.5 μM
146 phosphate are the conditions of natural seawater. The 400 $\mu\text{atm } p\text{CO}_2$ was achieved
147 by bubbling ambient air and 1000 $\mu\text{atm } p\text{CO}_2$ was obtained through a CO_2 plant
148 chamber (HP1000 G-D, Wuhan Ruihua Instrument & Equipment Ltd, China) with the
149 variation of CO_2 less than 5%. The higher P level (40 μM) was achieved by adding
150 NaH_2PO_4 to natural seawater and the nitrate concentration was set as 200 μM for all
151 treatments to avoid N limit. The media were refreshed every day.

152 2.2. Carbonate chemistry parameters

153 The seawater pH was recorded with a pH meter (pH 700, Eutech Instruments,
154 Singapore) and total alkalinity (TA) was measured by titrations. The salinity of
155 seawater was 29. Other carbonate system parameters, which were not directly
156 measured, were calculated via CO2SYS (Pierrot et al., 2006), using the equilibrium
157 constants of K_1 and K_2 for carbonic acid dissociation (Roy et al., 1993).

158 2.3. Measurement of growth

159 The growth of *S. muticum* was determined by weighing fresh thalli. The thalli of *S.*
160 *muticum* were blotted gently with tissue paper to remove water on the surface of the
161 thalli before weighing. The relative growth rate (RGR) was estimated as follows:
162 $\text{RGR} = (\ln W_t - \ln W_0) / t \times 100$, where W_0 is the initial fresh weight (FW) and W_t is
163 the weight after t days culture.

164 2.4. Determination of photosynthesis and respiration



165 The net photosynthetic rate of thalli was measured by a Clark-type oxygen
166 electrode (Chlorolab-3, Hansatech, Norfolk, UK) at the end of the experiment.
167 Approximately 0.1 g of fresh weight algae harvested from the culture flask was
168 transferred to the oxygen electrode cuvette with 8 ml sterilized media, and the media
169 were stirred during measurement. The irradiance and temperature conditions were set
170 as the same as that in the growth incubators. The increase of oxygen content in
171 seawater within five minutes was defined as net photosynthetic rate and the decrease
172 of oxygen content in seawater in darkness within ten minutes was defined as
173 respiration rate. Net photosynthetic rate (NPR) and respiration rate were presented as
174 $\mu\text{mol O}_2 \text{ g}^{-1} \text{ FW h}^{-1}$.

175 Photosynthetic rates at different dissolved inorganic carbon (DIC) levels were
176 measured under saturating irradiance of $600 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ at the end of the
177 experiment. The various DIC concentrations (0–13.2 mM) were obtained by adding
178 different amounts of NaHCO_3 to the Tris buffered DIC-free seawater. DIC was
179 removed from the natural seawater by reducing pH to approximately 4.0 with the
180 addition of 1.0 M HCl, and then sparging for 2 h with pure N_2 gas (99.999%). Finally,
181 Tris buffer (25mM) was added and the pH was adjusted to 8.1 with freshly prepared 1
182 M NaOH and 1 M HCl. The parameters, maximum photosynthetic rate (V_{max}) and the
183 half saturation constant ($K_{0.5}$, i.e., the DIC concentration required to give half of
184 Ci-saturated maximum rate of photosynthetic O_2 evolution), were calculated from the
185 Michaelis-Menten kinetics equation (Caemmerer and Farquhar, 1981):

186 $V = V_{max} \times [S] / (K_{0.5} + [S])$, where [S] is the DIC concentration.

187 2.5. Assessment of photosynthetic pigments

188 Approximately 100 mg of fresh weight thalli from each culture condition at the end
189 of the experiment was ground thoroughly in 2 ml 80% acetone and placed in darkness
190 for 12 hours. Then the homogenate was centrifuged for 10 minutes at 5,000 g and the
191 supernatant was used to determine Chl *a* content spectrophotometrically according to
192 the equation of Lichtenthaler (1987).

193 2.6. Measurement of nitrate uptake rate

194 The nitrate uptake rate (NUR) of thalli was estimated from the decrease of NO_3^-



195 concentration in the culture medium over a given time interval (12 hours) during light
196 period using the following equation: $NUR = (N_0 - N_t) \times V / W / 12$, where N_0 is the
197 initial concentration of NO_3^- , N_t is the concentration after 12 hours, V is the volume of
198 the culture medium, and W is the fresh weight of the thalli in culture. NO_3^-
199 concentration in the seawater was measured according to Strickland and Parsons
200 (1972).

201 2.7. Estimate of nitrate reductase activity

202 Nitrate reductase activity of thalli was assayed according to modified method of
203 Corzo and Niell (1991). The measurement was conducted during the local noon
204 period (13:00) since the activity of nitrate reductase usually displays circadian
205 periodicity a maximum during the light period and a minimum in darkness (Deng et
206 al., 1991; Velasco and Whitaker, 1989). Approximately 0.3 g (FW) of thalli from each
207 culture condition was incubated for 1 h at 20°C in darkness in the reaction solution
208 (10 mL), which contained 0.1 M phosphate buffer, 0.1% propanol (v/v), 50 mM
209 KNO_3 , 0.01 mM glucose, and 0.5 mM EDTA, with a pH of 8.0. The mixture was
210 flushed with pure N_2 gas (99.999%) for 2 minutes to obtain an anaerobic state before
211 the incubation. The concentration of nitrite produced was determined colorimetrically
212 at 540 nm (Zou, 2005). The NR activity was expressed as $\mu\text{mol } NO_2^- \text{ g}^{-1} \text{ FW h}^{-1}$.

213 2.8. Analysis of biochemical composition

214 About 0.2 g of FW thalli from each culture condition at the end of the experiment
215 were ground in a mortar with distilled water and soluble carbohydrates were extracted
216 in a water bath of 80°C for 30 min. After being centrifuged for 10 minutes at 5,000 g,
217 supernatant was volumed to 25 ml with distilled water, and soluble carbohydrates
218 content was determined by phenol-sulfuric acid method (Kochert, 1978).

219 Approximately 0.2 g of FW thalli from each culture condition at the end of the
220 experiment were ground in a mortar with extraction buffer (0.1 mol L^{-1} phosphate
221 buffer, pH 6.8) and then centrifuged for 10 minutes at 5,000 g. Soluble protein was
222 estimated from the supernatant using the Bradford (1976) assay with bovine serum
223 albumin as a standard.

224 2.9. Data Analysis



225 Results were expressed as means of replicates \pm standard deviation. Data were
226 analyzed using the software SPSS v.21. The data under every treatment conformed to
227 a normal distribution (Shapiro-Wilk, $P > 0.05$) and the variances could be considered
228 equal (Levene's test, $P > 0.05$). Two-way ANOVA was conducted to assess the effects
229 of $p\text{CO}_2$ and P on carbonate parameters, relative growth rate, net photosynthesis rate,
230 V_{max} , $K_{0.5}$, Chl *a*, nitrate uptake rate, nitrate reductase activity, soluble carbohydrates,
231 soluble protein, and dark respiration rate. Tukey HSD was conducted for *post hoc*
232 investigation. A confidence interval of 95% was set for all tests.

233 3. Results

234 The effects of ocean acidification and P enrichment on seawater carbonate
235 parameters were detected first (Table 1). Two-way ANOVA analysis ($P = 0.05$)
236 showed that $p\text{CO}_2$ had a main effect on all parameters except TA whilst P did not
237 affect any parameter. *Post hoc* Tukey HSD comparison ($P = 0.05$) showed projected
238 ocean acidification decreased pH by 0.31 unit at both LP and HP, CO_3^{2-} by 45.24%
239 (LP) and 44.70% (HP), but increased $p\text{CO}_2$ by 138.29% (LP) and 134.08% (HP), DIC
240 by 9.53% (LP) and 9.26% (HP), HCO_3^- by 14.11% (LP) and 13.79% (HP), and CO_2
241 by 138.88% (LP) and 134.20% (HP).

242 The growth of *S. muticum* cultured at different $p\text{CO}_2$ and P conditions was
243 recorded (Fig. 1). $p\text{CO}_2$ and P had an interactive effect on the relative growth rate of *S.*
244 *muticum* (ANOVA, $F = 5.776$, $df = 1, 8$, $P = 0.043$) and each factor had a main effect
245 (ANOVA, $F = 19.145$, $df = 1, 8$, $P = 0.002$ for $p\text{CO}_2$; ANOVA, $F = 30.592$, $df = 1, 8$,
246 $P = 0.001$ for P). *Post hoc* Tukey HSD comparison ($P = 0.05$) showed that the higher
247 levels of $p\text{CO}_2$ and higher P alone increased the relative growth rate by 40.82% and
248 47.78% respectively, compared to the relative growth rate ($3.05 \pm 0.36\%$) at the
249 condition of lower $p\text{CO}_2$ and lower P. The combination of the higher $p\text{CO}_2$ and higher
250 P levels did not enhance the relative growth rate as much as the sum of the higher
251 $p\text{CO}_2$ alone plus the higher P alone, with an increase of 59.66%. Although the higher
252 P level increased the relative growth rate at the condition of lower $p\text{CO}_2$, it did not
253 affect the relative growth rate at the condition of higher $p\text{CO}_2$.



254 In terms of the net photosynthetic rate (Fig. 2), both $p\text{CO}_2$ (ANOVA, $F = 26.556$,
255 $df = 1, 8, P = 0.001$) and P had main effects (ANOVA, $F = 38.963, df = 1, 8, P <$
256 0.001) on it. *Post hoc* Tukey HSD comparison ($P = 0.05$) showed the higher $p\text{CO}_2$
257 level increased the net photosynthetic rate by 46.34% and 23.96% at the conditions of
258 lower P and higher P respectively. The higher P level increased the net photosynthetic
259 rate by 55.16% and 31.43% at the conditions of lower $p\text{CO}_2$ and higher $p\text{CO}_2$
260 respectively. The difference in the net photosynthetic rate between LCHP and HCLP
261 was statistically insignificant.

262 The carbon-saturating maximum photosynthetic rate (V_{\max}) and the half saturation
263 constant ($K_{0.5}$), obtained from the photosynthesis versus DIC curves (Fig. 3), are
264 shown in Table 2. The $p\text{CO}_2$ and P had an interactive effect on V_{\max} of *S. muticum*
265 (ANOVA, $F = 10.095, df = 1, 8, P = 0.013$) and each factor had a main effect
266 (ANOVA, $F = 31.402, df = 1, 8, P = 0.001$ for $p\text{CO}_2$; ANOVA, $F = 105.116, df = 1, 8,$
267 $P < 0.001$ for P). *Post hoc* Tukey HSD comparison ($P = 0.05$) showed the higher
268 $p\text{CO}_2$ level increased the V_{\max} by 42.44% at the condition of lower P while the
269 increase at the condition of higher P was statistically insignificant. The higher P level
270 increased the V_{\max} at the conditions of both lower $p\text{CO}_2$ (64.90%) and higher $p\text{CO}_2$
271 (24.01%), with the larger promoting effect at the condition of lower $p\text{CO}_2$.

272 $p\text{CO}_2$ and P interacted on the $K_{0.5}$ of *S. muticum* (ANOVA, $F = 5.928, df = 1, 8, P$
273 $= 0.041$) and each factor had a main effect (ANOVA, $F = 14.713, df = 1, 8, P = 0.005$
274 for $p\text{CO}_2$; ANOVA, $F = 20.857, df = 1, 8, P = 0.002$ for P). *Post hoc* Tukey HSD
275 comparison ($P = 0.05$) showed the higher $p\text{CO}_2$ level increased the $K_{0.5}$ by 97.85% at
276 the condition of lower P but did not affect it at the condition of higher P. In contrast,
277 the higher P level decreased the $K_{0.5}$ by 55.22% at the condition of higher $p\text{CO}_2$ and
278 the negative effect of the higher P level at the condition of lower $p\text{CO}_2$ was
279 insignificant.

280 The contents of photosynthetic pigment Chl *a* under various treatments were also
281 estimated (Fig. 4). $p\text{CO}_2$ and P had an interactive effect on the Chl *a* content
282 (ANOVA, $F = 8.184, df = 1, 8, P = 0.021$), P had a main effect (ANOVA, $F = 22.828,$
283 $df = 1, 8, P = 0.001$), while $p\text{CO}_2$ did not affect it (ANOVA, $F = 0.676, df = 1, 8, P =$



284 0.435). *Post hoc* Tukey HSD comparison ($P = 0.05$) showed the higher P level
285 increased the Chl *a* content from 0.17 ± 0.00 to 0.25 ± 0.02 mg g⁻¹ FW at the
286 condition of lower $p\text{CO}_2$ whereas the difference in the Chl *a* content between HCLP
287 (0.21 ± 0.02 mg g⁻¹ FW) and HCHP (0.23 ± 0.02 mg g⁻¹ FW) was not statistically
288 significant.

289 To assess the effects of ocean acidification and P enrichment on the nitrogen
290 assimilation in *S. muticum*, nitrate uptake rate under various $p\text{CO}_2$ and P treatments
291 was investigated first (Fig. 5). Both $p\text{CO}_2$ (ANOVA, $F = 139.916$, $df = 1, 8$, $P < 0.001$)
292 and P (ANOVA, $F = 43.923$, $df = 1, 8$, $P < 0.001$) had main effects on the nitrate
293 uptake rate of *S. muticum*. The nitrate uptake rates at the conditions of lower $p\text{CO}_2$
294 were 0.18 ± 0.01 (LP) and 0.25 ± 0.03 $\mu\text{mol NO}_3^- \text{g}^{-1} \text{FW h}^{-1}$ (HP) respectively. *Post*
295 *hoc* Tukey HSD comparison ($P = 0.05$) showed the higher $p\text{CO}_2$ level increased the
296 nitrate uptake rate to 0.31 ± 0.02 $\mu\text{mol NO}_3^- \text{g}^{-1} \text{FW h}^{-1}$ at the condition of lower P and
297 to 0.39 ± 0.01 $\mu\text{mol NO}_3^- \text{g}^{-1} \text{FW h}^{-1}$ at the condition of higher P, compared to those at
298 the conditions of lower $p\text{CO}_2$. The higher P level also increased the nitrate uptake rate
299 by 35.89% at the condition of lower $p\text{CO}_2$ and by 27.71% at the condition of higher
300 $p\text{CO}_2$, compared to those at the conditions of lower P.

301 Apart from nitrate uptake, the nitrate reductase activity (NRA) of *S. muticum*
302 under various $p\text{CO}_2$ and P treatments was also detected (Fig. 6). $p\text{CO}_2$ and P
303 interacted on NRA of *S. muticum* (ANOVA, $F = 28.435$, $df = 1, 8$, $P = 0.001$) and
304 $p\text{CO}_2$ had a main effect (ANOVA, $F = 59.038$, $df = 1, 8$, $P < 0.001$). The NRA at the
305 conditions of lower $p\text{CO}_2$ were 0.10 ± 0.01 (LP) and 0.14 ± 0.02 $\mu\text{mol NO}_2^- \text{g}^{-1} \text{FW}$
306 h^{-1} (HP) respectively, and the higher $p\text{CO}_2$ level increased it to 0.19 ± 0.00 $\mu\text{mol NO}_2^-$
307 $\text{g}^{-1} \text{FW h}^{-1}$ at the condition of lower P and to 0.15 ± 0.02 $\mu\text{mol NO}_2^- \text{g}^{-1} \text{FW h}^{-1}$ at the
308 condition of higher P. The higher P level increased the NRA by 39.31% at the
309 condition of lower $p\text{CO}_2$, however, it decreased NRA by 17.81% at the condition of
310 higher $p\text{CO}_2$.

311 The soluble carbohydrates (Fig. 7a) and protein (Fig. 7b) were estimated to
312 understand the effects of ocean acidification and P enrichment on the products of
313 carbon and nitrogen assimilation in *S. muticum*. $p\text{CO}_2$ and P had an interactive effect



314 on the soluble carbohydrates (ANOVA, $F = 18.294$, $df = 1, 8$, $P = 0.003$) and P had a
315 main effect (ANOVA, $F = 23.129$, $df = 1, 8$, $P = 0.001$). The higher P level increased
316 the soluble carbohydrates from 25.40 ± 1.66 to 41.10 ± 1.74 mg g⁻¹ FW at the
317 condition of lower $p\text{CO}_2$ but did not alter it at the condition of higher $p\text{CO}_2$. The
318 higher $p\text{CO}_2$ level increased the soluble carbohydrates to 33.72 ± 3.31 mg g⁻¹ FW at
319 the condition of lower P while the decrease of soluble carbohydrates caused by the
320 higher $p\text{CO}_2$ level was not statistically significant at the condition of higher P.

321 Both $p\text{CO}_2$ (ANOVA, $F = 106.663$, $df = 1, 8$, $P < 0.001$) and P (ANOVA, $F =$
322 75.003 , $df = 1, 8$, $P < 0.001$) had main effects on the soluble protein of *S. muticum*
323 and the interactive effect of the two factors was not detected (ANOVA, $F = 4.961$, df
324 $= 1, 8$, $P = 0.057$). The soluble protein contents at the conditions of lower $p\text{CO}_2$ were
325 8.49 ± 0.49 (LP) and 9.77 ± 0.14 mg g⁻¹ FW (HP) respectively. The higher $p\text{CO}_2$ level
326 increased it to 10.11 ± 0.16 mg g⁻¹ FW at the condition of lower P and to 12.28 ± 0.44
327 mg g⁻¹ FW at the condition of higher P. The higher P level also increased the soluble
328 protein contents by 15.13% at the condition of lower $p\text{CO}_2$ and by 21.51% at
329 condition of higher $p\text{CO}_2$.

330 Finally, the effects of ocean acidification and P enrichment on the dark respiration
331 rate of *S. muticum* were investigated (Fig. 8). $p\text{CO}_2$ and P had an interactive effect on
332 the dark respiration rate (ANOVA, $F = 19.584$, $df = 1, 8$, $P = 0.002$) and each factor
333 had a main effect (ANOVA, $F = 6.428$, $df = 1, 8$, $P = 0.035$ for $p\text{CO}_2$; ANOVA, $F =$
334 6.754 , $df = 1, 8$, $P = 0.032$ for P). The higher $p\text{CO}_2$ level increased the dark
335 respiration rate from 14.21 ± 1.94 to 21.24 ± 1.28 $\mu\text{mol O}_2$ g⁻¹ FW h⁻¹ at the condition
336 of higher P but did not affect it at the condition of lower P. Likewise, The higher P
337 level increased the respiration rate from 14.15 ± 0.65 to 21.24 ± 1.28 $\mu\text{mol O}_2$ g⁻¹ FW
338 h⁻¹ at the condition of higher $p\text{CO}_2$ but did not change it at the condition of lower
339 $p\text{CO}_2$.

340 4. Discussion

341 4.1. Effects of $p\text{CO}_2$ and P on carbon assimilation

342 The higher $p\text{CO}_2$ level increased the net photosynthetic rate in *S. muticum* at the
343 condition of lower P in the present study. Although the dissolved inorganic carbon in



344 seawater is around 2 mM, the dominant form is HCO_3^- , with CO_2 typically accounting
345 for less than 1% (Dickson, 2010). In addition, CO_2 in seawater diffuses ~8,000 times
346 slower than in air (Gao and Campbell, 2014). Furthermore, the marine macroalgae
347 have high $K_{0.5}$ values (40–70 $\mu\text{M CO}_2$) for Rubisco, the carbon assimilating enzyme
348 (Ji et al., 2016). The evidence above indicates that the CO_2 in seawater should be
349 carbon limited for marine macroalgae. The promoting effect of elevated CO_2 on
350 photosynthesis was also reported in other macroalgae species, such as green algae
351 *Ulva linza* (Gao et al., 1999), red algae *Pyropia haitanensis* (Zou and Gao, 2002), and
352 brown algae *Petalonia binghamiae* (Gao and Kunshan, 2010). Meanwhile, the higher
353 $p\text{CO}_2$ level increased $K_{0.5}$ of *S. muticum* at the condition of lower P in the present
354 study, which indicates the plant grown at the condition of higher $p\text{CO}_2$ reduced its
355 photosynthetic affinity for DIC. This phenomenon is commonly found in both
356 microalgae and macroalgae (Gao and Campbell, 2014; Ji et al., 2016; Wu et al., 2008)
357 and is considered as a sign of down-regulated CCMs at high CO_2 conditions (Gao and
358 Campbell, 2014). But this decrease of photosynthetic affinity for DIC did not lead to
359 reduced photosynthesis in *S. muticum* in the present study, mainly because of
360 increased CO_2 availability for Rubisco and depressed photorespiration at the elevated
361 ratio of CO_2 to O_2 , which has been confirmed in red seaweed *Lomentaria articulata*
362 (Kübler et al., 1999).

363 The higher P level also increased the net photosynthetic rate of *S. muticum* in the
364 present study, which can be partially explained by the decreased $K_{0.5}$ at the condition
365 of higher P. The decreased $K_{0.5}$ is an indication of increased photosynthetic carbon-use
366 capability. Phosphorus is a key macronutrient component for organisms and high
367 levels of P availability is not only essential for chloroplast DNA and RNA synthesis
368 (Vered and Shlomit, 2008), but is required for various chloroplast functions, referring
369 to phosphorylation of photosynthetic proteins, synthesis of phospholipids and
370 generation of ATP (Zer and Ohad, 2003). Therefore, High P levels could speed up the
371 transport of C_i from media to the site of Rubisco by supplying necessary energy. In
372 addition, P enrichment can increase both activity and amount of Rubisco (Lauer et al.,
373 1989). Meanwhile, phosphorus, with low concentration in seawater, is generally



374 considered to be limiting for marine primary producer (Elser et al., 2007; Howarth,
375 1988; Müller and Mitrovic, 2015). Therefore, adding extra phosphorus to natural
376 seawater can stimulate photosynthesis of algae. For instance, the midday (12:00)
377 photosynthetic rates increased from 1.3 to 2.3 mg C g⁻¹ DW h⁻¹ for *S. natans*, from 0.9
378 to 2.1 mg C g⁻¹ DW h⁻¹ for *S. fluitans* when 0.2 mM P was added (Lapointe, 1986). In
379 the present study, the addition of 40 µmol P also resulted in nearly two-fold increase
380 of the net photosynthetic rate and V_{max} , which suggests the significant importance of P
381 in photosynthesis of this alga. In addition, the higher P level promoted the synthesis of
382 Chl *a* at the condition of lower pCO_2 , which may also contribute to the increased net
383 photosynthetic rate in *S. muticum* at the condition of higher P. Although P is not the
384 component constituting Chl *a*, higher P supply may stimulate the content of Chl *a*
385 synthesis-related enzymes and thus the production of Chl *a*. The positive effect of P
386 on Chl *a* was also reported in *S. thunbergii* (Nakahara, 1990). On the other hand, the
387 higher P level did not increase the Chl *a* content at the condition of higher pCO_2 in the
388 present study. The possible reason is that there is more ATP available at the condition
389 of higher pCO_2 due to the down-regulation of CCMs and thus there is no need to
390 synthesize more Chl *a* to capture more light for cells as excessive energy can lead to
391 the harm to photosynthesis and growth of algae (Gao et al., 2012; Xu and Gao, 2012).

392 4.2. Effect of pCO_2 and P on nitrogen assimilation

393 The higher pCO_2 level noticeably enhanced the nitrate uptake rate in *S. muticum*
394 regardless of P concentration in the present study. This could be attributed to the
395 increased nitrate reductase activity (NRA) at the condition of higher pCO_2 . The
396 enhanced NRA at the conditions of high CO_2 was also reported in *U. rigida* (Gordillo
397 et al., 2001), *Hizikia fusiforme* (Zou, 2005), *P. haitanensis* (Liu and Zou, 2015),
398 *Corallina officinalis* (Hofmann et al., 2013), as well as the higher plants *Plantago*
399 *major* (Fonseca et al., 1997), tomato (Yelle et al., 1987), etc. Taken together, these
400 findings indicate that the response of NRA in plants to elevated CO_2 may be
401 homogeneous.

402 The higher P level also enhanced the nitrate uptake in *S. muticum* regardless of
403 pCO_2 level, which can be partially due to the increased NRA at the condition of



404 higher P. This is very evident at the condition of lower $p\text{CO}_2$. However, the higher P
405 level decreased the NRA at the condition of higher $p\text{CO}_2$, which did not lead to
406 reduced nitrate uptake. This indicates there should be other mechanisms to account
407 for the promoting effect of the higher P level on the nitrate uptake. One possible
408 mechanism is that the higher P level can increase the availability of ATP that is
409 required for the active uptake of nitrate across the plasma membrane. The
410 phenomenon that ATP concentration increases with P level has been found in higher
411 plants (Olivera et al., 2004; Rychter et al., 2006). Apart from *S. muticum*, the positive
412 effect of higher P level on nitrate uptake was also reported in red macroalgae
413 *Gracilaria lemaneiformis* (Xu et al., 2010) and higher plant *Phaseolus vulgaris*
414 (Gniazdowska and Rychter, 2000). The increased nitrate uptake, NRA and soluble
415 protein at the condition of higher P in the present study suggest that high P
416 availability promoted nitrogen assimilation in *S. muticum*.

417 The higher P level increased the nitrate uptake rate and soluble protein at the
418 conditions of both lower $p\text{CO}_2$ and higher $p\text{CO}_2$ but it only increased the NRA in *S.*
419 *muticum* at the condition of lower $p\text{CO}_2$ in the present study. Surprisingly, it
420 decreased the NRA at the condition of higher $p\text{CO}_2$. The reason for that may be not
421 onefold but must be related to interaction of $p\text{CO}_2$ and P. High $p\text{CO}_2$, on one hand,
422 could enhance photosynthetic carbon fixation and thus growth by supplying sufficient
423 CO_2 . On the other hand, it also results in the decrease of pH and increase of seawater
424 acidity, which can disturb the acid-base balance on cell surface of algae (Flynn et al.,
425 2012). Algae may accordingly allocate additional energy to act against the acid-base
426 perturbation in some way. This hypothesis is supported by increased respiration at the
427 condition of higher $p\text{CO}_2$ and higher P in the present study. The increased soluble
428 protein and decreased NRA at the condition of higher $p\text{CO}_2$ and higher P suggest
429 some H^+ transport-related protein, such as plasma membrane H^+ -ATPase, might be
430 synthesized to counteract the acid-base perturbation caused by increased $p\text{CO}_2$ and
431 H^+ . The plasma membrane H^+ -ATPase plays an essential role in maintaining an
432 electrochemical proton gradient across the plasma membrane (Morth et al., 2011;
433 Sondergaard et al., 2004). The additional production of H^+ transport-related protein



434 like plasma membrane H⁺-ATPase could competitively decrease the synthesis of
435 nitrate reductase. This hypothesis needs further experimental evidence to stand even
436 though it could explain the results in the present study. Meanwhile, the higher *p*CO₂
437 can also deliver the signal to induce the synthesis of H⁺ transport-related protein, but
438 low P supply may limit the synthesis. Accordingly, the nitrate reductase activity did
439 not decrease at the condition of higher *p*CO₂ and lower P.

440 4.3. Connection between carbon and nitrogen assimilation

441 The increased net photosynthetic rate at the condition of higher *p*CO₂ and higher
442 P did not result in higher soluble carbohydrates compared to the condition of higher
443 *p*CO₂ and lower P. The additional ATP produced by photosynthetic electron transport
444 at the condition of higher *p*CO₂ and higher P may be drawn to nitrogen assimilation as
445 more soluble protein was synthesized at the condition of higher *p*CO₂ and higher P.
446 The additional energy allocation to protein synthesis, possibly H⁺ transport-related
447 protein to maintain the balance of acid-base, hindered the increase of growth, which
448 may be the reason that the higher P increased the net photosynthetic rate but not the
449 growth rate at the condition of higher *p*CO₂. Although synthesized protein can also
450 contribute to the increase of thalli weight, it is not as energy-effective as
451 carbohydrates (Norici et al., 2011; Raven, 1982). It seems that *S. muticum* tends to
452 maintain a steady state in vivo even if it can sacrifice growth to some extent,
453 considering that regulation of intracellular acid-base balance is crucial for organismal
454 homeostasis (Flynn et al., 2012; Smith and Raven, 1979). The increased respiration
455 at HC was also demonstrated in *G. lemaneiformis* (Xu et al., 2010) and *U. prolifera*
456 (Xu and Gao, 2012). The respiration at the condition of higher *p*CO₂ and lower P did
457 not increase compared to at the condition of lower *p*CO₂ and lower P in the present
458 study, suggesting the action against acid-base perturbation did not commence. The
459 acid-base perturbation at the condition of higher *p*CO₂ and lower P may lead to the
460 decreased photosynthetic rate compared to that at the condition of lower *p*CO₂ and
461 lower P.

462 5. Conclusion

463 Our study, for the first time, demonstrates the combined effects of elevated *p*CO₂



464 and P enrichment on the physiological traits of a golden alga, *S. muticum*. It suggests
465 current ocean environment is both CO₂ and P limited for the photosynthesis and grow
466 of *S. muticum*. Therefore, future ocean acidification and eutrophication may promote
467 the growth of *S. muticum* and thus occurrence of gold tide events. Meanwhile, *S.*
468 *muticum* tends to maintain homeostasis taking advantage of phosphate enrichment,
469 at the cost of growth. Accordingly, the combination of ocean acidification and
470 eutrophication may not boost gold tides further compared to ocean acidification or
471 eutrophication alone.

472 **Acknowledgements**

473 This study was supported by the National Natural Science Foundation of China
474 (Nos. 41376129, 41476097 and 31270452), the Public Science and Technology
475 Research Funds Projects of Ocean (Nos. 201505022, 201405040 and 201305021), the
476 Earmarked Fund for Modern Agro-industry Technology Research System in
477 Shandong Province (SDAIT-26), and the Experimental Study Project on Ecological
478 Simulation in Coastal Waters of Shandong Peninsula.

479 **References**

- 480 Alvaro, I. and Mazal, H.: Growth, photosynthetic properties and Rubisco activities
481 and amounts of marine macroalgae grown under current and elevated seawater
482 CO₂ concentrations, *Glob. Chang. Biol.*, 30, 831-840, 2002.
- 483 Ang, P. O.: Phenology of *Sargassum* spp. in Tung Ping Chau Marine Park, Hong
484 Kong SAR, China, *J. Appl. Phycol.*, 18, 403-410, 2006.
- 485 Ashok-Kumar, N., Vanlalzarzova, B., Sridhar, S., and Baluswami, M.: Effect of liquid
486 seaweed fertilizer of *Sargassum wightii* Grev. on the growth and biochemical
487 content of green gram (*Vigna radiata* (L.) R. Wilczek), *Recent Res. Sci. Technol.*,
488 4, 40-45, 2012.
- 489 Bradford, M. M.: A rapid and sensitive method for the quantitation of microgram
490 quantities of protein utilizing the principle of protein-dye binding, *Anal.*
491 *Biochem.*, 72, 248-254, 1976.
- 492 Bricker, S. B., Longstaff, B., Dennison, W., Jones, A., Boicourt, K., Wicks, C., and
493 Woerner, J.: Effects of nutrient enrichment in the nation's estuaries: a decade of



- 494 change, *Harmful Algae*, 8, 21-32, 2008.
- 495 Caemmerer, S. V. and Farquhar, G. D.: Some relationships between the biochemistry
496 of photosynthesis and the gas exchange of leaves, *Planta*, 153, 376-387, 1981.
- 497 Carpenter, S. R.: Submersed vegetation: an internal factor in lake ecosystem
498 succession, *Am. Nat.*, 1981. 372-383, 1981.
- 499 Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and
500 Smith, V. H.: Nonpoint pollution of surface waters with phosphorus and nitrogen,
501 *Ecol. Appl.*, 8, 559-568, 1998.
- 502 Cheang, C. C., Chu, K. H., Fujita, D., Yoshida, G., Hiraoka, M., Critchley, A., Choi, H.
503 G., Duan, D., Serisawa, Y., and Ang, P. O.: Low genetic variability of *Sargassum*
504 *muticum* (Phaeophyceae) revealed by a global analysis of native and introduced
505 populations, *J. Phycol.*, 46, 1063-1074, 2010.
- 506 Chen, B. and Zou, D.: Growth and photosynthetic activity of *Sargassum*
507 *henslowianum* (Fucales, Phaeophyta) seedlings in responses to different light
508 intensities, temperatures and CO₂ levels under laboratory conditions, *Mar. Biol.*
509 *Res.*, 10, 1019-1026, 2014.
- 510 Corzo, A. and Niell, F. X.: Determination of nitrate reductase activity in *Ulva rigida* C.
511 Agardh by the in situ method, *J. Exp. Mar. Bio. Ecol.*, 146, 181-191, 1991.
- 512 Cruzrivera, E., Floresd áz, M., and Hawkins, A.: A fish kill coincident with dense
513 *Sargassum accumulation* in a tropical bay, *Bull. Mar. Sci.*, 91, 455-456, 2015.
- 514 Deng, M. D., Moureaux, T., Cherel, I., Boutin, J. P., and Caboche, M.: Effects of
515 nitrogen metabolites on the regulation and circadian expression of tobacco nitrate
516 reductase, *Plant Physiol. Biochem.*, 29, 239-247, 1991.
- 517 Dickson, A. G.: The carbon dioxide system in seawater: Equilibrium chemistry and
518 measurements. In: Guide to best practices for ocean acidification research and
519 data reporting, Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J. P. (Eds.),
520 Publications Office of the European Union, Luxembourg, 2010.
- 521 Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S.,
522 Hillebrand, H., Ngai, J. T., Seabloom, E. W., Shurin, J. B., and Smith, J. E.:
523 Global analysis of nitrogen and phosphorus limitation of primary producers in



- 524 freshwater, marine and terrestrial ecosystems, *Ecol. Lett.*, 10, 1135-1142, 2007.
- 525 Fenoradosoa, T. A., Ali, G., Delattre, C., Laroche, C., Petit, E., Wadouachi, A., and
526 Michaud, P.: Extraction and characterization of an alginate from the brown
527 seaweed *Sargassum turbinarioides* Grunow, *J. Appl. Phycol.*, 22, 131-137, 2010.
- 528 Flynn, K. J., Blackford, J. C., Baird, M. E., Raven, J. A., Clark, D. R., Beardall, J.,
529 Brownlee, C., Fabian, H., and Wheeler, G. L.: Changes in pH at the exterior
530 surface of plankton with ocean acidification, *Nat. Clim. Chang.*, 2, 510-513,
531 2012.
- 532 Fonseca, F., Bowsher, C. G., and Stulen, I.: Impact of elevated atmospheric CO₂ on
533 nitrate reductase transcription and activity in leaves and roots of *Plantago major*,
534 *Physiol. Plant.*, 100, 940-948, 1997.
- 535 Gao, D. Z. and Kunshan: Acquisition of inorganic carbon by *Endarachne binghamiae*
536 (Scytosiphonales, Phaeophyceae), *Eur. J. Phycol.*, 45, 117-126, 2010.
- 537 Gao, K. and Campbell, D. A.: Photophysiological responses of marine diatoms to
538 elevated CO₂ and decreased pH: a review, *Funct. Plant Biol.*, 41, 449-459, 2014.
- 539 Gao, K., Xu, J., Gao, G., Li, Y., Hutchins, D. A., Huang, B., Wang, L., Zheng, Y., Jin,
540 P., and Cai, X.: Rising CO₂ and increased light exposure synergistically reduce
541 marine primary productivity, *Nat. Clim. Change*, 2, 519-523, 2012.
- 542 Gao, K., Yan, J., and Aruga, Y.: Relationship of CO₂ concentrations to photosynthesis
543 of intertidal macroalgae during emersion, *Hydrobiologia*, 398/399, 355-359,
544 1999.
- 545 Gniazdowska, A. and Rychter, A. M.: Nitrate uptake by bean (*Phaseolus vulgaris* L.)
546 roots under phosphate deficiency, *Plant & Soil*, 226, 79-85, 2000.
- 547 González-López, N., Moure, A., and Domínguez, H.: Hydrothermal fractionation of
548 *Sargassum muticum* biomass, *J. Appl. Phycol.*, 24, 1569-1578, 2012.
- 549 Gordillo, F. J. L., Niell, F. X., and Figueroa, F. L.: Non-photosynthetic enhancement
550 of growth by high CO₂ level in the nitrophilic seaweed *Ulva rigida* C. Agardh
551 (Chlorophyta), *Planta*, 213, 64-70, 2001.
- 552 Hofmann, L., Straub, S., and Bischof, K.: Elevated CO₂ levels affect the activity of
553 nitrate reductase and carbonic anhydrase in the calcifying rhodophyte *Corallina*



- 554 *officinalis*, J. Exp. Bot., 64, 899–908, 2013.
- 555 Howarth, R. W.: Nutrient limitation of net primary production in marine ecosystems,
556 Annu. Rev. Ecol. Syst., 1988. 89-110, 1988.
- 557 Hwang, R. L., Tsai, C. C., and Lee, T. M.: Assessment of temperature and nutrient
558 limitation on seasonal dynamics among species of sargassum from a coral reef in
559 southern taiwan, J. Phycol., 40, 463-473, 2004.
- 560 Incera, M., Olabarria, C., Troncoso, J. S., and López, J.: Response of the invader
561 *Sargassum muticum* to variability in nutrient supply, Mar. Ecol. Prog. Ser., 377,
562 91-101, 2009.
- 563 IPCC: Climate change 2013: The physical science basis. In: Working Group I
564 Contribution to the Fifth Assessment Report of the Intergovernmental Panel on
565 Climate Change, Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K.,
566 Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M. (Eds.), Cambridge
567 Univ Press, New York, 2013.
- 568 Ji, Y., Xu, Z., Zou, D., and Gao, K.: Ecophysiological responses of marine macroalgae
569 to climate change factors, J. Appl. Phycol., 2016. 1-15, 2016.
- 570 Jones, G. and Farnham, W.: Japweed: new threat to British coasts, New Sci., 60,
571 394-395, 1973.
- 572 Kübler, J. E., Johnston, A. M., and Raven, J. A.: The effects of reduced and elevated
573 CO₂ and O₂ on the seaweed *Lomentaria articulata*, Plant Cell & Environment,
574 22, 1303-1310, 1999.
- 575 Karlsson, J. and Loo, L. O.: On the distribution and continuous expansion of the
576 Japanese seaweed - *Sargassum muticum* - in Sweden, Bot. Mar., 42, 285-294,
577 1999.
- 578 Kochert, G.: Carbohydrate determination by the phenol-sulfuric acid method. In:
579 Handbook of Phycological Methods: Physiological and Biochemical Methods,
580 Hellebust, J. A. and Graigie, J. S. (Eds.), Cambridge University Press,
581 Cambridge, 1978.
- 582 Laffoley, D. A., Roe, H. S. J., Angel, M. V., Ardron, J., Bates, N. R., Boyd, I. L.,
583 Brooke, S., Buck, K. N., Carlson, C. A., and Causey, B.: The protection and



- 584 management of the Sargasso Sea: The golden floating rainforest of the Atlantic
585 Ocean, 2011.
- 586 Lapointe, B. E.: A comparison of nutrient - limited productivity in *Sargassum natans*
587 from neritic vs. oceanic waters of the western North Atlantic Ocean, *Limnol. &*
588 *Oceanogr.*, 40, 625-633, 1995.
- 589 Lapointe, B. E.: Phosphorus-limited photosynthesis and growth of *Sargassum natans*
590 and *Sargassum fluitans* (Phaeophyceae) in the western North Atlantic, *Deep Sea*
591 *Res. Part A Oceanogr. Res. Pap.*, 33, 391-399, 1986.
- 592 Lapointe, B. E., Littler, M. M., and Littler, D. S.: A comparison of nutrient-limited
593 productivity in macroalgae from a Caribbean barrier reef and from a mangrove
594 ecosystem, *Aquat. Bot.*, 28, 243-255, 1987.
- 595 Lapointe, B. E., Littler, M. M., and Littler, D. S.: Nutrient availability to marine
596 macroalgae in siliciclastic versus carbonate-rich coastal waters, *Estuaries &*
597 *Coasts*, 15, 75-82, 1992.
- 598 Lauer, M. J., Pallardy, S. G., Blevins, D. G., and Randall, D. D.: Whole leaf carbon
599 exchange characteristics of phosphate deficient soybeans (*Glycine max* L.), *Plant*
600 *Physiol.*, 91, 848-854, 1989.
- 601 Lichtenthaler, H. K.: Chlorophylls and carotenoids: Pigments of photosynthetic
602 biomembranes, *Methods Enzymol.*, 148, 350-382, 1987.
- 603 Littler, M. M., Littler, D. S., and Titlyanov, E. A.: Comparisons of N- and P-limited
604 productivity between high granitic islands versus low carbonate atolls in the
605 Seychelles Archipelago: a test of the relative-dominance paradigm, *Coral Reefs*,
606 10, 199-209, 1991.
- 607 Liu, C. and Zou, D.: Effects of elevated CO₂ on the photosynthesis and nitrate
608 reductase activity of *Pyropia haitanensis* (Bangiales, Rhodophyta) grown at
609 different nutrient levels, *Chin. J. Oceanol. Limnol.*, 33, 419-429, 2015.
- 610 Liu, Y. and Tan, H.: Changes of growth and nutrient-relating enzymatic activities of
611 *Sargassum thunbergii* when exposed to different nutrient conditions, *Aquat. Sci.*
612 *Technol.*, 2, 1-13, 2014.
- 613 Longphuir, S. N., Eschmann, C., Russell, C., and Stengel, D. B.: Seasonal and



- 614 species specific response of five brown macroalgae to high atmospheric CO₂,
615 Mar. Ecol. Prog. Ser., 493, 91-102, 2014.
- 616 Müller, S. and Mitrovic, S. M.: Phytoplankton co-limitation by nitrogen and
617 phosphorus in a shallow reservoir: progressing from the phosphorus limitation
618 paradigm, Hydrobiologia, 744, 255-269, 2015.
- 619 Mattio, L. and Payri, C. E.: 190 years of *Sargassum* taxonomy, facing the advent of
620 DNA phylogenies, Bot. Rev., 77, 31-70, 2011.
- 621 Morth, J. P., Pedersen, B. P., Buch-Pedersen M. J., Andersen, J. P., Vilsen, B.,
622 Palmgren, M.G. and Nissen, P. A: Structural overview of the plasma membrane
623 Na⁺, K⁺-ATPase and H⁺-ATPase ion pumps, Nat. Rev. Mol.Cell Biol., 12, 60-70,
624 2011.
- 625 Nakahara, K. G. H.: Effects of nutrients on the photosynthesis of *Sargassum*
626 *thunbergii*, Bot. Mar., 33, 375-384, 1990.
- 627 Norici, A., Bazzoni, A. M., Pugnetti, A., Raven, J. A., and Giordano, M.: Impact of
628 irradiance on the C allocation in the coastal marine diatom *Skeletonema marinoi*
629 Sarno and Zingone, Plant Cell Environ., 34, 1666–1677, 2011.
- 630 Olivera, M., Tejera, N., Iribarne, C., Ocaña, A., and Lluch, C.: Growth, nitrogen
631 fixation and ammonium assimilation in common bean (*Phaseolus vulgaris*):
632 effect of phosphorus, Physiol. Plant., 121, 498–505, 2004.
- 633 Pierrot, D., Lewis, E., and Wallace, D. W. R.: MS Excel program developed for CO₂
634 system calculations, ORNL/CDIAC-105a. Carbon Dioxide Information Analysis
635 Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge,
636 Tennessee, 2006.
- 637 Raven, J., Caldeira, K., Elderfield, H., Hoegh-Guldberg, O., Liss, P., Riebesell, U.,
638 Shepherd, J., Turley, C., and Watson, A.: Ocean acidification due to increasing
639 atmospheric carbon dioxide, The Royal Society, London, 2005.
- 640 Raven, J. A.: The energetics of freshwater algae; energy requirements for biosynthesis
641 and volume regulation, New Phytol., 92, 1–20, 1982.
- 642 Roy, R. N., Roy, L. N., Vogel, K. M., Porter-Moore, C., Pearson, T., Good, C. E.,
643 Millero, F. J., and Campbell, D. M.: The dissociation constants of carbonic acid



- 644 in seawater at salinities 5 to 45 and temperatures 0 to 45°C, Mar. Chem., 44,
645 249-267, 1993.
- 646 Rueness, J.: *Sargassum muticum* and other introduced Japanese macroalgae:
647 Biological pollution of European coasts, Mar. Pollut. Bull., 20, 173-176, 1989.
- 648 Rychter, A. M., Chauveau, M., Bomsel, J. L., and Lance, C.: The effect of phosphate
649 deficiency on mitochondrial activity and adenylate levels in bean roots, Physiol.
650 Plant., 84, 80-86, 2006.
- 651 Scagel, R. F.: Introduction of a Japanese alga, *Sargassum muticum*, into the northeast
652 Pacific, Fisheries Research Papers, 1, 49-58, 1956.
- 653 Schaffelke, B. and Klumpp, D. W.: Nutrient-limited growth of the coral reef
654 macroalga *Sargassum baccularia* and experimental growth enhancement by
655 nutrient addition in continuous flow culture, Mar. Ecol. Prog. Ser., 164, 199-211,
656 1998.
- 657 Schell, J. M., Goodwin, D. S., and Siuda, A. N. S.: Recent sargassum inundation
658 events in the Caribbean, Oceanography, 28, 8-10, 2015.
- 659 Sfriso, A. and Facca, C.: Annual growth and environmental relationships of the
660 invasive species *Sargassum muticum* and *Undaria pinnatifida* in the lagoon of
661 Venice, Estuar. Coast. Shelf Sci., 129, 162-172, 2013.
- 662 Smetacek, V. and Zingone, A.: Green and golden seaweed tides on the rise, Nature,
663 504, 84-88, 2013.
- 664 Smith, F. A. and Raven, J. A.: Intracellular pH and its regulation, Annu. Rev. Plant
665 Physiol., 30, 289-311, 1979.
- 666 Smith, V. H., Tilman, G. D., and Nekola, J. C.: Eutrophication: impacts of excess
667 nutrient inputs on freshwater, marine, and terrestrial ecosystems, Environ. Pollut.,
668 100, 179-196, 1999.
- 669 Sondergaard, T. E., Schulz, A. and Palmgren, M. G.: Energization of transport
670 processes in plants. Roles of the plasma membrane H⁺-ATPase, Plant Physiol.,
671 136, 2475-2482, 2004.
- 672 Stæhr, P. A., Pedersen, M. F., Thomsen, M. S., Wernberg, T., and KrauseJensen, D.:
673 Invasion of *Sargassum muticum* in Limfjorden (Denmark) and its possible



674 impact on the indigenous macroalgal community, Mar. Ecol. Prog. Ser., 207,
675 79-88, 2000.

676 Strickland, J. D. H. and Parsons, T. R.: A practical handbook of seawater analysis, 2nd
677 ed., Fisheries Research Board of Canada, Ottawa, 1972.

678 Velasco, P. J. and Whitaker, J. R.: Synthesis and degradation of nitrate reductase
679 during the cell cycle of *Chlorella sorokiniana*, Plant Physiol., 89, 220-224, 1989.

680 Vered, I. and Shlomit, Y. R.: Phosphate and sulfur limitation responses in the
681 chloroplast of *Chlamydomonas reinhardtii*, FEMS Microbiol. Lett., 283, 1-8,
682 2008.

683 Wu, H. Y., Zou, D. H., and Gao, K. S.: Impacts of increased atmospheric CO₂
684 concentration on photosynthesis and growth of micro- and macro-algae, Sci.
685 China Ser. C Life Sci., 51, 1144-1150, 2008.

686 Xu, J. and Gao, K.: Future CO₂-induced ocean acidification mediates the
687 physiological performance of a green tide alga, Plant Physiol., 160, 1762-1769,
688 2012.

689 Xu, Z., Zou, D. H., and Gao, K.: Effects of elevated CO₂ and phosphorus supply on
690 growth, photosynthesis and nutrient uptake in the marine macroalga *Gracilaria*
691 *lemaneiformis* (Rhodophyta), Bot. Mar., 53, 123-129, 2010.

692 Yelle, Gosselin, and Trudel: Effect of atmospheric CO₂ concentration and root-zone
693 temperature on growth, mineral nutrition, and nitrate reductase activity of
694 greenhouse tomato, J. Am. Soc. Hort. Sci., 112, 1036-1040, 1987.

695 Zer, H. and Ohad, I.: Light, redox state, thylakoid-protein phosphorylation and
696 signaling gene expression, Trends Biochem. Sci., 28, 467-470, 2003.

697 Zou, D.: Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen
698 metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae,
699 Phaeophyta), Aquaculture, 250, 726-735, 2005.

700 Zou, D. and Gao, K.: Effects of desiccation and CO₂ concentrations on emersed
701 photosynthesis in *Porphyra haitanensis* (Bangiales, Rhodophyta), a species
702 farmed in China, Eur. J. Phycol., 37, 587-592, 2002.

703



707 **Table 1.** Parameters of the seawater carbonate system at different CO₂ and phosphate conditions. Measurements and estimation of the
 708 parameters are described in Materials and Methods. Data are the means ±SD (n = 3). LCLP, the low pCO₂ and low P condition, LCHP, the low
 709 pCO₂ and high P condition, HCLP, the high pCO₂ and low P condition, HCHP, the high pCO₂ and P condition, DIC = dissolved inorganic carbon,
 710 TA = total alkalinity. Different superscript letters indicate significant differences in one parameter between treatments (*P* < 0.05).

Treatment	pH	pCO ₂ (µatm)	HCO ₃ ⁻ (µmol kg ⁻¹)	CO ₃ ²⁻ (µmol kg ⁻¹)	CO ₂ (µmol kg ⁻¹)	DIC (µmol kg ⁻¹)	TA (µmol kg ⁻¹)
LCLP	8.07±0.02 ^b	426.9±31.1 ^a	2000.2±51.7 ^a	200.9±5.8 ^b	14.2±1.0 ^a	2215.3±49.7 ^a	2475.2±44.2
LCHP	8.07±0.02 ^b	423.9±21.1 ^a	1987.6±10.9 ^a	199.8±11.4 ^b	14.1±0.7 ^a	2201.5±19.3 ^a	2504.7±33.8
HCLP	7.76±0.02 ^a	1017.2±83.2 ^b	2282.5±27.6 ^b	110.0±10.0 ^a	34.0±2.9 ^b	2426.5±32.5 ^b	2541.5±44.2
HCHP	7.76±0.02 ^a	992.2±44.9 ^b	2261.8±35.9 ^b	110.5±5.9 ^a	33.1±1.5 ^b	2405.4±39.4 ^b	2563.6±44.2



712 **Table 2.** The carbon-saturating maximum photosynthetic rate (V_{max} , $\mu\text{mol O}_2 \text{ g}^{-1} \text{ FW}$
713 h^{-1}) and half saturation constant ($K_{0.5}$, mM) for *S. muticum* cultured under different
714 $p\text{CO}_2$ and P conditions for 13 days. Different superscript letters indicate significant
715 differences in one parameter between treatments ($P < 0.05$).

	LCLP	LCHP	HCLP	HCHP
V_{max}	57.00 ± 2.88^a	93.99 ± 0.98^c	81.18 ± 5.94^b	100.67 ± 6.81^c
$K_{0.5}$	0.21 ± 0.02^a	0.14 ± 0.05^a	0.42 ± 0.08^b	0.19 ± 0.05^a



716 **Fig. 1.** Relative growth rate (RGR) of *S. muticum* grown at different $p\text{CO}_2$ and P
717 conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low $p\text{CO}_2$ and low
718 P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high $p\text{CO}_2$ and low
719 P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different letters above error
720 bars indicate significant differences between treatments ($P < 0.05$).

721 **Fig. 2.** Net photosynthetic rate (RGR) of *S. muticum* after being grown at different
722 $p\text{CO}_2$ and P conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low
723 $p\text{CO}_2$ and low P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high
724 $p\text{CO}_2$ and low P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different
725 letters above error bars indicate significant differences between treatments ($P < 0.05$).

726 **Fig. 3.** The photosynthesis versus DIC curves of *S. muticum* after being cultured
727 under $p\text{CO}_2$ and P conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the
728 low $p\text{CO}_2$ and low P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the
729 high $p\text{CO}_2$ and low P condition, HCHP, the high $p\text{CO}_2$ and high P condition. DIC =
730 dissolved inorganic carbon.

731 **Fig. 4.** Chl *a* content of *S. muticum* after being grown at different $p\text{CO}_2$ and P
732 conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low $p\text{CO}_2$ and low
733 P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high $p\text{CO}_2$ and low
734 P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different letters above error
735 bars indicate significant differences between treatments ($P < 0.05$).

736 **Fig. 5.** Nitrate uptake rate of *S. muticum* after being grown at different $p\text{CO}_2$ and P
737 conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low $p\text{CO}_2$ and low
738 P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high $p\text{CO}_2$ and low
739 P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different letters above error
740 bars indicate significant differences between treatments ($P < 0.05$).

741 **Fig. 6.** Nitrate reductase activity (NRA) of *S. muticum* after being grown at different
742 $p\text{CO}_2$ and P conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low
743 $p\text{CO}_2$ and low P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high
744 $p\text{CO}_2$ and low P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different
745 letters above error bars indicate significant differences between treatments ($P < 0.05$).

746 **Fig. 7.** The contents of soluble carbohydrates (a) and protein (b) of *S. muticum* after
747 being grown at different $p\text{CO}_2$ and P conditions for 13 days. Data are the means \pm SD
748 ($n = 3$). LCLP, the low $p\text{CO}_2$ and low P condition, LCHP, the low $p\text{CO}_2$ and high P
749 condition, HCLP, the high $p\text{CO}_2$ and low P condition, HCHP, the high $p\text{CO}_2$ and high



750 P condition. Different letters above error bars indicate significant differences between
751 treatments ($P < 0.05$).

752 **Fig. 8.** Dark respiration rate of *S. muticum* after being grown at different $p\text{CO}_2$ and P
753 conditions for 13 days. Data are the means \pm SD ($n = 3$). LCLP, the low $p\text{CO}_2$ and low
754 P condition, LCHP, the low $p\text{CO}_2$ and high P condition, HCLP, the high $p\text{CO}_2$ and low
755 P condition, HCHP, the high $p\text{CO}_2$ and high P condition. Different letters above error
756 bars indicate significant differences between treatments ($P < 0.05$).




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8