

High resolution regional modeling of natural and anthropogenic radiocarbon in the Mediterranean Sea

Mohamed Ayache¹, Jean-Claude Dutay¹, Anne Mouchet¹, Nadine Tisnérat-Laborde¹, Paolo Montagna², Toste Tanhua³, Giuseppe Siani⁴, and Philippe Jean-Baptiste¹

¹Laboratoire des Sciences du Climat et de l'Environnement LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

²Istituto di Scienze Marine, CNR, Bologna, Italy

³GEOMAR Helmholtz Centre for Ocean Research Kiel, Dusternbrooker Weg 20, 24105 Kiel, Germany

⁴Laboratoire Interactions et Dynamique des Environnements de Surface IDES, UMR 8148 CNRS - Université Paris-Sud 11, Orsay, France

Correspondence to: M Ayache (mohamed.ayache@lsce.ipsl.fr)

We thank Prof. Christoph Heinze (Editor), and anonymous Referees for their constructive comments and suggestions, which have helped to improve the manuscript. We have carefully considered all questions and concerns raised.

We provide a marked-up manuscript version; all change is recalled in red (see the revised manuscript with point-by-point responses below, after the responses to the reviewers' comments).

The structure of our reply is as follows; each comment from the anonymous reviewer is recalled in blue, and our reply in black.

Reply to Referree#1

The authors use a high-resolution dynamical model to simulate the distribution of radiocarbon in the Mediterranean Sea. While I feel the topic is relevant and the treatment new, I would like the author would better specify different aspects treated in the paper:

a) the role of the Atlantic Water: sensitivity experiments at Gibraltar should be discussed;

We have performed two simulations with different boundary conditions at Gibraltar (red and blue boxes and lines, see Fig.1, below); the first time series (red box) gives very low level of radiocarbon in the Mediterranean Sea (as represented in Fig.2, below). In the second simulation we used a larger box (blue in Fig.1), where results are more realistic compared to some data from the North Atlantic (Tisnérat-Laborde et al., 2013; Tisnérat-Laborde, personal communication). The radiocarbon simulation greatly improves when using the larger box as boundary conditions, hence this was used to simulate ^{14}C in the Mediterranean Sea.

This part has been clarified in the revised version of the manuscript.

[See changes p 7, line 18-25 in the revised manuscript.]




Fig. 1: The concentration of radiocarbon in the Atlantic inflow (NEMO global model, Mouchet et al. 2016).




Fig.2: $\Delta^{14}\text{C}$ values (in ‰) in the Ligurian sub-basin from 1765 to 2008 for the surface water (0-10 m depth), together with available in-situ observations (Tisnérat-Laborde et al., 2013) from coral (black dashed line). Simulated data obtained using the smaller box (blue line, simu 1), and the larger box (blue dashed line, simu 2).

b) the artificial modifications performed in order to simulate the EMT should be deeply discussed;

To improve dense-water fluxes through the Cretan Arc during the EMT (1992-1993) the ARPERA forcings were modified over the Aegean sub-basin (Beuvier et al., 2012a), by increasing mean values as done by Herrmann and Somot (2008) for the Gulf of Lions. More specifically, from November to March for the winters 1991-1992 and 1992-1993, daily surface heat loss was increased by 40 W m^{-2} , daily water loss by 1.5 mm and the daily wind stress modulus by 0.02 N m^{-2} . These changes accelerate the transfer of surface temperature and salinity perturbations into intermediate and deep layers of the Aegean subbasin, and improve the dense-water formation in the Aegean sub-basin during the EMT, with more intense mixing from winter convection.

The artificial modifications performed in order to simulate the EMT were fully discussed by Beuvier et al (2010), and in our previous work on anthropogenic tritium modelling (Ayache et al., 2015a). In this study we have used the same parametrization and method as in those previous work, therefore we do not think it is necessary to deeply discuss the details in the present manuscript.

[See changes, p 07 line 30-35 in the revised manuscript.]

c) how can convective penetration be increased in the simulations?

The convective penetration is more important in the classical area of deep convection in the Mediterranean Sea (i.e. Gulf of Lion, Adriatic and the Aegean sub-basins...) where the surface heat loss, water loss and the wind stress are more important on those areas.

In this work we have used a high resolution dynamical model (NEMO-MED12, Beuvier et al. 2012) based on the tagged version nemo v3.2 of the NEMO ocean general circulation model (Madec et al., 2008). This model was only forced by the atmospheric model AREPERA and we prescribe the initial and boundary conditions (as detailed in the manuscript section 2.1). Increasing the convective penetration could be obtained by changing air sea fluxes or in the Adriatic changing river runoff, but it is not the goal of this paper.

d) overall, a more critical discussion about limitations of the model simulations should be addressed.

We agree with the reviewer that limitations of the model simulations should be more critically discussed.

Previous passive tracer evaluations of NEMO-MED12 (e.g., Ayache et al., 2016; Ayache et al., 2015a; Palmiéri et al., 2015) have shown that the model satisfactorily simulates the main structures of the thermohaline circulation of the Mediterranean Sea, with mechanisms having a realistic timescale compared to observations.

However tritium/helium-3 simulations from Ayache et al. (2015) have highlighted the too-weak formation of Adriatic Deep Water (AdDW), followed by a weak contribution to the Eastern Mediterranean Deep Water (EMDW) in the Ionian sub-basin. In the western basin, the

production of WMDW is correctly simulated, but the spreading of the recently ventilated deep water to the south of the basin is too weak. The consequences of these weaknesses in the model's skill in simulating some important aspects of the dynamics of the deep ventilation of the Mediterranean will have to be kept in mind when analyzing the model output.

We thank the reviewer for this suggestion. This part has been extended in the revised version.

[See changes p 12 line 34, in the revised manuscript.]

Reply to Referree#2

p.2, 1.1-3 'The Mediterranean Sea can be considered as a "miniature ocean", where global change can be studied at smaller/shorter spatial and temporal scales (100 yr compared to more than 1000 yr for the global ocean ...).' The mentioned time scales of 100 vs. 1000 years refer to the overturning time of the Mediterranean/world ocean. Is that really identical with the time scale on which global change is going on, as it is implied by this sentence?

In many study the Mediterranean Sea is described as a miniature ocean (e.g. Lascaratos et al 1999) based on the difference of the overturning time of the Mediterranean/world ocean. Most of the physical processes that characterize the global general ocean circulation (e.g. intermediate and deep water formation) also occur in the Mediterranean Sea but at shorter time scale. This allows investigating human-induced climate modifications that are rapidly transferred to sub-surface waters in the entire Mediterranean Sea. For example, the increase of seawater temperature at intermediate and deep level due to the effect of the present global warming is stronger in the Mediterranean Sea compared to that observed at similar depths in the global ocean. Similarly, acidification due to uptake of anthropogenic carbon is already affecting all deep water masses of the Mediterranean Sea (Palmieri et al, 2015) Owing to its small size and limited exchange with the Atlantic Ocean, the Mediterranean Sea amplifies the effects of global changes, which can be then studied at shorter temporal scales.

The shorter Mediterranean turnover timescales permits to perform longer and more computational efficient simulations.

For the sake of clarity, we have modified this sentence in the revised version of the manuscript.

[See changes p 2, line 2-4 in the revised manuscript.]

p.2, 1.19-22 In this paragraph, ^{14}C is characterized as conservative tracer such as CFCs and tritium. This is not exactly true, as ^{14}C is changed by biology, especially the remineralisation of organic matter. This effect is small and often neglected, but it still is a conceptual difference.

We thank the referee for this suggestion; this conceptual difference between ^{14}C and the other tracers has been clarified in the revised manuscript.

[See changes p 2, line 25-26 in the revised manuscript.]

p.3, l.6-8 and p.33, l.31 Here and at some other passages in the paper the role of ^{14}C for the determination of water mass ages and constraining the deep water circulation is mentioned. This is not wrong, but regarding ages, ^{14}C is normally used in older waters with ages of order 1000 yr (comparable to the half-life time). For the Mediterranean, tracers with shorter input histories such as CFCs and tritium are more useful. They are also more useful in constraining the deep water pathways in circulation models because the number of observations is much larger than for ^{14}C . This should be made clear somewhere in the text.

We agree with the referee, that tracers with shorter input histories are more adapted to investigate water mass ages and circulation in the Mediterranean Sea (see for example Ayache et al., 2015a for anthropogenic tritium and Palmieri et al., 2015 for CFCs).

The present radiocarbon simulation aims at implementing a geochemical tracer with a longer time scale allowing more paleo-oriented applications. This ^{14}C modelling would help improving the knowledge of the natural distribution of ^{14}C in the Mediterranean, providing a unique opportunity to explore the impact of the interannual/decadal variability on radiocarbon distribution in the Med Sea.

Clarified in revised version

[See changes p 4, line 19-21 in the revised manuscript.]

p.7, first paragraph on the choice of kw: It seems to me that the choice of kw is the main work regarding the tuning of the circulation model on the base of ^{14}C data. So this topic might be given more room in the description and discussion.

For the Mediterranean Sea we have studied the impact of Kw on the radiocarbon distribution in this semi-enclosed basin, and we have chosen a value that gives the best agreement with available in-situ data. On the other hand the present simulation was done in a computationally efficient off-line mode (as mentioned in section 2.2), i.e. the dynamic was run independently from the ^{14}C module and the Kw parametrization was adapted for the Mediterranean Sea.

A sentence was added to clarify this point.

[See changes p 7, line 4-7 in the revised manuscript.]

p.8, l.17-18 '... leading to a relatively higher ^{14}C level in the EMed surface water closer to -46 ‰' Has the value of -46 ‰ a special meaning? Then this should be mentioned in the text. According to Fig. 2a, the values are close to -44 ‰

The referee is correct, the value is closer to -44 and this has been corrected in the revised version. The -46 ‰ has no special meaning.

[See changes p 8, line 20 in the revised manuscript.]

p.8, l.18-20 'For both western and eastern surface water, the model simulates ^{14}C concentrations slightly higher than the in-situ observations...' I don't see this from the data. In Fig. 2d, 2e and 3, the data are sometimes smaller and sometimes higher than the model results.

The values given in table 1 for model and observations are almost identical for the WMed and EMed, only smaller subregions show significant differences.

The referee is correct, there are no significant differences between WMed and the EMed average values. However, if we look at Fig. 3 (data from Siani et al. 2000) there is an important spatial gradient across the different sub-basins in the Mediterranean Sea as a consequence of old carbon impact near the coastal areas. This effect is not represented in the present simulation. On the other hand, our model results are in good agreement with average values provided by Reimer and McCormac (2002) for each sub basin in the Mediterranean. Other in-situ data would help to improve the model parametrization.

p.8, l.20-21 'A careful comparison between model outputs and seawater observations (1959) reveals a more pronounced dis-agreement, especially in the EMed surface water where the model overestimates the ^{14}C values by more than 10‰ (Fig.4a).' Where is the profile shown in Fig. 4a located? Or is it a composite from different locations? If it is one complete profile, the location should be indicated in the inlet map of Fig. 2e or given in coordinates. Second, the measured EMed surface value shown in Fig. 2e is much larger than the value from Fig. 4a, around -45 ‰ So how representative is the profile shown in Fig. 4a for the whole EMed?

The vertical profile shown in Fig. 4a is a composite of seawater observations from different locations (Brocker et al., 1969) as represented in Fig. 2 for the pre-bomb situation. The measured EMed surface value shown in Fig. 2e is much larger than the value from Fig. 4a because the latter presents the average value of all in-situ data and model output on the same position. However the representation of the pre-bomb distribution is more contrasted in the simulation, where several issues complicate the simulation of the natural steady state distribution of ^{14}C using ocean-model circulation (e.g. the uncertainty associated with the boundary conditions).

For the sake of clarity, we have modified Fig.4 caption in the revised manuscript.

p.10, first paragraph Only the higher ^{14}C values in the deep water in the Levantine basin are mentioned here, although in the western Med. the values are comparably high between 4 °E and 10 °E.

Added in the revised manuscript.

[See changes p 10, line 11 in the revised manuscript.]

p.10, l.13-14 'However the model simulates well the ^{14}C values in the surface and deep water of Adriatic sub-bassin (Figure 7a and 7c) compared to Meteor M84/3cruise data (Tanhua et al., 2013).' According to Fig. 7a and 7c the model values are too high, which is even more pronounced in Fig. 7b for the intermediate layers?

We agree with the referee that the model values in the Adriatic deep water are higher compared to those obtained for the Meteor M84/3 cruise. However the high ^{14}C level in the deep water proved that the model simulates deep convection in the Adriatic sub-basin. Nevertheless, the outflow of this deep water through the Strait of Otranto is weaker in the model and the simulated signal of deep-water ventilation from the Adriatic sub-basin is propagating at too shallow depth compared to the observations. This shortcoming was also noticed for the other tracer simulations with the same model NEMO-MED12 (e.g. Ayache et al. 2015a; Palmiéri et al.,

2015; Ayache et al. 2015b; Ayache et al. 2016). In the Adriatic sub-basin, the contribution of rivers is very important; however, the atmospheric forcing ARPERA combined with the river runoff data set overestimates the freshwater flux, and provides too much freshwater on this domain. This results in unusually low-salinity water compared to observations, preventing winter convection and the propagation of AddW to the bottom of the Ionian sub-basin.

Figures:

Fig.2: In subfigures b and c, the y-labels have a larger fontsize than the x-labels. The fontsize of the colour bar is too small, and the space between the colour bar and the upper maps should be enhanced.

Adjusted

Fig.3: The font size of the axis labels is too large and of the labels of the color bar too small.

Corrected

Fig.5: Exactly the same as for Fig.2.

Adjusted

Fig.7: Exactly the same as for Fig.2.

Adjusted

Fig.11: The ylabel 'Time (yr)' should be centered.

Corrected

Minor comments/corrections:

p.6, Eq.1 the vector 'u' should be notated in bold math

Corrected

p.9, 1.26 '... when we compare...' (not compared)

Corrected

p.10. 1.13 '...values in the surface and deep water of the Adriatic sub-basin' ('the' is missing and 'basin' is misspelled)

Corrected

p.12, l.3 'However the representation of the pre-bomb distribution is more contrasted in the simulation' I don't understand the meaning of 'contrasted'.

Replaced by 'more difficult'

p.13 l.7 '... to prolonged exposure of the surface water to the atmosphere.' (add 'the' before 'atmosphere').

Done

p.13 l.7-8 'where it depends on convection processes with higher convection occurring especially during the bomb peak' I don't see why higher convection has occurred during the bomb peak. Maybe it is meant that the amount of ^{14}C entering the deep water was higher during that time.

The transfer of radiocarbon was higher during the bomb peak as a consequence of large amount of ^{14}C in the atmosphere. We agree with the referee that this sentence is not clear and it can be easily misinterpreted. It has been modified in the revised version.

[See changes p 13, line 16-17 in the revised manuscript.]

p.13, l.18 '... at the bottom of the Levantine sub-basin' ('the' is missing)

Corrected.

Reply to Referee#3

Abstract

The simulation was run until 2010 to give the post-bomb distribution. I believe the simulation is run until 2008 although model outputs are compared with in situ measurements taken in 2011.

The ^{14}C simulation was done in a computationally efficient off-line mode, which allows us to run simulation of different tracers (e.g. Palmieri et al 2015; Ayache et al. 2015), in pre-computed transport fields instead of re-computing them, which is very costly.

The dynamical fields are available until 2011 from Beuvier et al. (2012). Starting from the end of the pre-industrial equilibrium run, the model was integrated from 1765 to 2011 (as mentioned page 7 line 13) covering the Suess effect (SUESS, 1955), the entire radiocarbon transient generated by the atmospheric nuclear weapon tests performed in the 1950s and early 1960s as well as the anthropogenic CO_2 increase.

Corrected in the revised manuscript

[See changes p 1, line 4 in the revised manuscript.]

Introduction Page 1, Lines 15-21: The whole paragraph seems to be out of scope. I do not see the relationship between the stresses suffered by the MedSea and the distribution of radiocarbon

In addition to providing constraints on radiocarbon distribution in the Mediterranean, our simulations provide information on the ventilation of the deep Mediterranean waters which is useful for assessing NEMO-MED12 performance.

This study is part of the work carried out to assess the robustness of the NEMO-MED12 model, which will be used to study the evolution of the climate and its effect on the biogeochemical cycles in the Mediterranean Sea, and to improve our ability to predict the future evolution of the Mediterranean Sea under the increasing anthropogenic pressure (e.g. Drobinski et al., 2012; Beuvier et al., 2010; Herrmann et al., 2010; Somot et al., 2006).

Page 2, Lines 6-10. The excess of evaporation versus precipitation does not transform Atlantic waters into Mediterranean waters and leads AW to sink offshore. It is actually the process that drives the entry of Atlantic waters through the Strait of Gibraltar to compensate water loss and keep the mass balance. Water masses formation in the basin are subsequently the result of other phenomena more related with atmospheric forcing and density gradients. It might be a small detail but conceptually it is important, especially for readers not familiar with the MedSea. I would suggest to re-write the paragraph.

We thank the referee for this suggestion, this paragraph is rephrased in the revised version according to reviewer's comments.

[See changes p 2, line 7-10 in the revised manuscript.]

Page 2, Line 21. I would not compare CFC and tritium with ^{14}C , as this one is not entirely passive. I understand what the authors mean by the sentence but radiocarbon is indeed used by the biological community so I would recommend to state that circumstance or simply not to equate the tracers.

This point has been also raised by reviewer #2 and it has been addressed in the revised version of the manuscript.

Results

Page 8, lines 17-18: According to Figs 2 and 3, the model does not overestimate the radiocarbon concentration in surface everywhere in the basin but it depends on the particular region. Also, plots in Fig. 2 could be manifestly enhanced as it is hard to distinguish in situ observations over the contour in the graphs.

The number of in-situ data for the pre-industrial period is very limited in the Mediterranean Sea. Figure 2 shows that the east-west gradient was satisfactorily captured by the model, with a slight overestimation of ^{14}C concentrations in the surface water of western basin. However results on figure 3 are reservoir ages and as discussed in the manuscript they are influenced by other sources (e.g. coastal input of "old carbon") so that only the spatial structure is discussed (east-west gradient).

More paleo-data from the pre-industrial period would help improving the knowledge of the natural distribution of ^{14}C in the Mediterranean.

Fig.2 was improved as suggested by the reviewer.

Page 8, Line 20. The careful comparison between vertical profiles of model outputs and seawater observations in Fig. 4 is restricted to the Eastern basin. Why the Western basin is not considered if according to Fig 3 there are also some disagreements? Not enough in situ data to compare? Please clarify

The careful comparison between vertical profiles of model outputs and in-situ data in Fig.4 is restricted to the EMed because there is no data available for the WMed for the bomb situation from Stuiver et al. (1983).

Fig. 3 presents only the surface values from paleo-reconstruction, and there is no data for the deep waters.

Page 8, Line 27. Any idea why the pre-bomb radiocarbon levels differ so much between the in situ data and the model outputs in the Aegean sub-basin (Table 1)? I guess there must be some circulation patterns not resolved in the model. Plus, I do not understand the sentence the range in the observations is also high.

The simulation of natural radiocarbon is particularly difficult because the average dynamical circulation used in the present study does not produce enough convection to pull-up the old carbon accumulated in the deep water. In addition, our simulation does not take into account the potential impact of old carbon in the coastal area, which could be the case in the Aegean sub-basin.

The sentence ‘the range in the observations is also high’ means that the range of uncertainty is higher in the observations.

Clarified in revised version.

[See changes p 8, line 30 in the revised manuscript.]

Page 9, Line 10. At depth, the model tends to underestimate the ^{14}C penetration in the deep Ionian sub-basin, where it fails to reproduce the high ^{14}C levels associated with EMDW formation (Fig. 4b). Where is this disagreement shown in Fig. 4b? Does the plot correspond to that particular sub-basin or to the entire Eastern basin?

The vertical profile plotted in Fig.4b represents the model result in the Ionian sub-basin together with in-situ data measured by Stuiver et al. 1983 at 18 °E.

As mentioned in the Introduction, in this study we used the NEMO-Med12 dynamical model that was already tested and evaluated with other tracers, such as tritium (Ayache et al., 2015a), helium (Ayache et al., 2015b) and CFC (Palmieri et al., 2015). Those tracers have highlighted that the model simulates a too-weak formation of Adriatic Deep Water (AdDW), followed by a weak contribution to the Eastern Mediterranean Deep Water (EMDW) in the Ionian sub-basin. The EMDW formed in the Adriatic basin is propagating to the entire deep eastern basin so that the consequences of weak formation of this water mass in the model are observed in the whole sub-basin.

Page 9, Line 24. The greater is the mixing layer depth, the weaker is the amplitude and the peak is delayed. Is this sentence grammatically correct?

We thank the referee for this suggestion, this sentence has been changed for the sake of clarity. **[See changes p 9, line 28-29 in the revised manuscript.]**

Page 9 and 10: To me, Fig. 7 depicts too much disagreement between simulated distributions and in situ data in many regions, not only in deep convection areas. For instance, even though it is hard to see the symbols in Fig. 7a, it seems that in surface waters of the Strait of Gibraltar the model overestimates the radiocarbon concentration by more than 20 ‰. Plus, in the discussion section it is stated that there is no time series data of ^{14}C concentration in that area, while in the graph there are at least 4 measurements in the gulf of Cadiz and within the channel of the Strait. Could have they been used to fuel the model? In addition, explanation of data indicated in Figs 7 and 8 is confused, as description of patterns jumps from one to another without a logic sequence.

Although we partly agree with the reviewer, we think that the disagreement between simulated distribution and in situ data is particularly evident in deep convection areas as represented in Fig. 7.

In this simulation we used the same parametrization for the whole basin with same boundary conditions at the surface (first level), with ^{14}C and the atmospheric CO_2 values extracted from Orr et al. (2001). The radiocarbon values in the buffer zone are prescribed from a global simulation of radiocarbon by Mouchet et al (2016), and the ocean ^{14}C is initially set to a constant value of 0.85 ($\Delta^{14}\text{C} = -150$ ‰, appropriate for the deep ocean; (Key et al., 2004)).

Hence we did not use any in-situ measurements to fuel the model for a specific region, because here we aimed to develop and optimize a ^{14}C modelling method for the whole Mediterranean Sea basin. Moreover no time series exists close to Gibraltar exist to force the model in the duration of the simulation. Data in the gulf of Cadiz represent a single date. As we used in-situ data to evaluate our results they cannot be employed to force the simulation.

Figures 7 and 8 present the same in-situ data of METEOR M84/3. Fig.7 provides a descriptive overview of the global horizontal distribution of ^{14}C , where the vertical profile in Fig.8 permits to quantify the difference between the model and in-situ data at different levels. Hence the description of these figures was mad at the same time in the text.

Page 10: The radiocarbon time evolution spans from 1925 to 2008, why this particular year? In fact, Fig. 7 shows comparison between the model outputs and data of the 2011 Meteor cruise, which included measurements throughout the whole basin. Why the simulated evolution does not run until then? It would be interesting to confirm that evolution follows the pattern indicated in Fig. 7. Also, I would keep the same vertical scale in all plots to facilitate comparisons. The response found in intermediate-deep waters of the gulf of Lions is somehow unexpected, as deep convection events during winter should favor the sink of radiocarbon, particularly in extreme winters, such as that occurring in the area in 2004/2005. In Fig 9d, the intermediate layer of the gulf of Lions exhibit the lowest radiocarbon levels after the bomb episodes and deeper waters are characterized by values even lower than those found in the Tyrrhenian sub-basin. Is there any explanation for that? Moreover, are data in plot 9d integrated values through

the whole water column? These results are not explained in the text. Plus, the title is wrong, it should say whole water columns.

The model was integrated from 1765 to 2011 as mentioned in section 2.3 page 7 line 14. The ^{14}C evolution was plotted from 1925 to 2008 in Fig.9 just to zoom on the period affected by the Suess effect (SUESS, 1955), and the entire radiocarbon transient generated by the atmospheric nuclear weapon tests performed in the 1950s and early 1960s as well as the anthropogenic CO_2 increase.

The intermediate-deep waters of the Gulf of Lions, characterized by ^{14}C values that are lower compared to the other sub-basins, are the result of a mixture of local water masses with Levantine Intermediate Water that is formed in the Levantine sub-basin, i.e. this water mass is isolated from the bomb signal in the atmosphere until arrived to the Gulf of Lion (deep convection area), hence the peak-bomb appears delayed in this area.

The data in plot 9d integrate values through the whole water column and present the same pattern as in Fig.9c for the deep water where the content of radiocarbon in the deep water control the distribution of $\Delta^{14}\text{C}$ in the whole water column.

This has been clarified in the revised text, and the title has been corrected as suggested by the reviewer. However we didn't use the same scale, because in the vertical section $\Delta^{14}\text{C}$ value up to -70 (due to the AdDW waters shortcomings, as mentioned provisory), and if we use the same scale for the horizontal maps, many information will be not clear as well (i.e. the gradient between the different basins).

[See changes p 11, line 3-4 in the revised manuscript.]

Discussion

Page 11, Line 10. The radiocarbon simulations provide independent and additional constraints on the thermohaline circulation and deep-water ventilation in the Mediterranean Sea. I do not see this in the manuscript. It would be the other way around. Data are interpreted according to the general circulation mechanisms known to proceed in the Med Sea.

Unlike the other tracers (e.g. CFC and Tritium), radiocarbon simulation provide additional constraints on the thermohaline circulation from the seasonal cycle to decadal and centennial timescales (e.g. Naegler, 2009; Muller et al., 2006; Rodgers et al., 1997; Guilderson et al., 1998).

In this study we have implemented the ^{14}C module in high resolution regional model, and we work mainly on the validation on this ^{14}C modelling method in the Mediterranean Sea basin, this work will allow many other applications especially in paleo-context e.g. Sapropel events. However direct comparison with in-situ ^{14}C data is a new constraint for the model and it has revealed or confirmed some shortcomings such the weak EMDW formation. This will be clarified in the conclusion of the paper.

Clarified in the revised manuscript

[See changes p 3, line 2-7 in the revised manuscript.]

Page 12. The comparison between the model outputs and the ^{14}C values from insitu data reported by Broecker and Gerard (1969), Stuiver et al. (1983) and Tanhua et al. (2013) reveals a good model performance in simulating the bomb/post-bomb radiocarbon distribution (Fig. 4b, Fig. 8). However the representation of the pre-bomb distribution is more contrasted in the simulation (Fig. 4a). I do not understand this paragraph. In fact, those two figures in particular show the largest disagreements, particularly in intermediate-deep waters and for the bomb-produced radiocarbon.

The reviewer is correct, there is a quite large disagreement between the model and in-situ data in some regions at intermediate-deep water depths for the bomb produced radiocarbon. However the mentioned sentence refers to Fig.6 where an important disagreement ($\sim 15\%$) is observed between the model outputs and the sea-surface ^{14}C record obtained from a 50-year-old shallow-water coral from Tisnérat-Laborde et al., (2013) for the natural pre-bomb signal. On the other hand the model nicely represents the bomb signal with the good timing and the amplitude of the peak in the near-surface water compared to in-situ data (Fig.6).

For the sake of clarity, this sentence has been modified in the revised version.

[See changes p 12, line 6-9 in the revised manuscript.]

Page 13, Line 7: with higher convection occurring especially during the bomb peak.

Where is this shown in the paper?

We agree with the reviewer that the sentence is not clear. There is no higher convection during the period of bomb peak, but the surface water masses undergo transfer or convection with different intensity in the different sectors of the Mediterranean Basin. This argument has been clarified in the manuscript

[See changes p 13, line 16-17 in the revised manuscript.]

Conclusions

The natural distribution of ^{14}C in the Mediterranean Sea is mainly affected by the inflow of Atlantic water through the Strait of Gibraltar.

As far as I understood, the concentration of radiocarbon in the Atlantic inflow did not come from in situ data or available measurements since it was taken from previous modeling approaches (as indicated in different sections of the paper). Therefore, this study does not show per se, the influence of the Atlantic radiocarbon on the distribution of this tracer in the Med Sea, as it is a fixed value used to fuel the model. The paper actually demonstrates that the entry of Atlantic waters is essential for water masses formation and circulation in the Mediterranean, which is a very well-known topic and which, in turn, regulates the distribution of radiocarbon. In fact, it would have been interesting to perform the same simulations by changing for instance the values of the water masses transport through the Strait or the radiocarbon concentration associated to the Atlantic jet. To me, such conclusion cannot be drawn from the data. I would omit it here and in the abstract or at least, the sentence should be re-written.

Unfortunately, there is no time series data of ^{14}C concentration close to the Strait of Gibraltar. Hence simulated ^{14}C levels in the model's Atlantic water (AW) are determined from global

model estimates. As mentioned in the paper, we have performed sensitivity tests on the imposed value (as presented in Fig.1, below).

We have performed two simulations with different boundary conditions at Gibraltar (red and blue boxes and lines); the first time series (red box) gives very low level of radiocarbon in the Mediterranean Sea (as represented in Fig.2, below). In the second simulation we used a larger box (blue in Fig.1), where results are more realistic compared to some data from the North Atlantic (Tisnérat-Laborde et al., 2013; Tisnerat-Laborde, personal communication). The radiocarbon simulation greatly improves when using the larger box as boundary conditions, hence this was used to simulate ^{14}C in the Mediterranean Sea.

This point has been also raised by reviewer #1 and it has been addressed in the revised version of the manuscript.

[See changes p 7, line 18-25 in the revised manuscript.]




Fig. 1: The concentration of radiocarbon in the Atlantic inflow (NEMO global model, Mouchet et al. 2016).




Fig.2: $\Delta^{14}\text{C}$ values (in ‰) in the Ligurian sub-basin from 1765 to 2008 for the surface water (0-10 m depth), together with available in-situ observations (Tisnérat-Laborde et al., 2013) from coral (black dashed line). Simulated data obtained using the smaller box (blue line, simu 1), and the larger box (blue dashed line, simu 2).

High resolution regional modeling of natural and anthropogenic radiocarbon in the Mediterranean Sea

Mohamed Ayache¹, Jean-Claude Dutay¹, Anne Mouchet¹, Nadine Tisnérat-Laborde¹, Paolo Montagna², Toste Tanhua³, Giuseppe Siani⁴, and Philippe Jean-Baptiste¹

¹Laboratoire des Sciences du Climat et de l'Environnement LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France.

²Istituto di Scienze Marine, CNR, Bologna, Italy

³GEOMAR Helmholtz Centre for Ocean Research Kiel, Dusternbrooker Weg 20, 24105 Kiel, Germany.

⁴Laboratoire Interactions et Dynamique des Environnements de Surface IDES, UMR 8148 CNRS- Université Paris-Sud 11, Orsay, France.

Correspondence to: M Ayache (mohamed.ayache@lsce.ipsl.fr)

Abstract. A high-resolution dynamical model (NEMO-MED12) was used to give the first simulation of the distribution of radiocarbon (^{14}C) across the whole Mediterranean Sea. The simulation provides a descriptive overview of both the natural pre-bomb ^{14}C and the entire anthropogenic radiocarbon transient generated by the atmospheric bomb tests performed in the 1950s and early 1960s. The simulation **was run until 2011** to give the post-bomb distribution. The results are compared to available in-situ measurements and proxy-based reconstructions. The radiocarbon simulation allows an additional and independent test of the dynamical model, NEMO-MED12, and its performance to produce the thermohaline circulation and deep-water ventilation. The model produces a generally realistic distribution of radiocarbon when compared with available in-situ data. The results demonstrate the major influence of the flux of Atlantic water through the strait of Gibraltar on the inter-basin natural radiocarbon distribution, and characterize the ventilation of intermediate and deep water ventilation especially through the propagation of the anthropogenic radiocarbon signal. We explored the impact of the interannual variability on the radiocarbon distribution during the Eastern Mediterranean transient event (EMT). It reveals a significant increase in ^{14}C concentration (by more than 60 ‰) in the Aegean deep water, and at intermediate level (value up to 10 ‰) in the western basin. The model shows that the EMT makes a major contribution to the accumulation of radiocarbon in the eastern Mediterranean deep waters.

1 Introduction

The Mediterranean region has been identified as a hot-spot for future climatic changes (Giorgi, 2006; Giorgi and Lionello, 2008; MerMex-Group, 2011; Diffenbaugh and Giorgi, 2012). Because this mid-latitude almost-enclosed sea is surrounded by countries with high population growth to the south and highly industrialized countries to the north, it is under strong anthropogenic pressures. This stress is expected to intensify due to factors such as warming and substantial precipitation decrease (Attané and Courbage, 2004). In the context of global change (IPCC, 2013) we need to improve our understanding of how changes in the climate and circulation of the Mediterranean Sea interact with the biogeochemical processes that define its functioning.

The Mediterranean Sea can be considered as a "miniature ocean", where global change can be studied at smaller/shorter spatial and temporal scales, e.g. **warming at intermediate water depths in the Mediterranean Sea is about 10 times larger than trends reported in literature (~~~100-yr compared to more than 1000-yr for the global ocean~~)** (e.g., Millot and Taupier-Letage, 2005; Schroeder et al., 2016). The Mediterranean Sea has a well-defined overturning circulation with distinct surface, intermediate and deep water masses circulating in the western and the eastern basins, and varying at interannual time scales. This makes it an excellent test bed for studying basic processes that will also affect the global thermohaline circulation.

The Mediterranean is a concentration basin in which evaporation exceeds precipitation and river runoff. Warmer, fresher water enters at the surface from the Atlantic (AW) through Gibraltar, and colder saline water leaves below. Relatively fresh waters of Atlantic origin circulating in the Mediterranean increase in density, and then form new water masses via convection events driven by intense local cooling from winter storms. The Levantine Intermediate Water (LIW) represents one of the main water masses of the Mediterranean Sea. It spreads throughout the entire Mediterranean basin at intermediate depths (between ~150 and 700 m) (Pinardi and Masetti, 2000), and is the major contributor to the Mediterranean outflow into the North Atlantic (Bryden and Stommel, 1984). Furthermore, the LIW participates in the deep convection processes of the Western Mediterranean deep water (WMDW) occurring in the Gulf of Lions, and the Adriatic sub-basin for the Eastern Mediterranean deep water (EMDW) (Millot and Taupier-Letage, 2005). The formation of deep water in the Mediterranean Sea is also characterized by interannual/decadal variability such as the Eastern Mediterranean Transient (EMT) event, known to create a major shift in deep water formation in the East Mediterranean Sea (EMed) at the beginning of the 1990s (Roether et al., 1996, 2007; Malanotte-Rizzoli et al., 1999; Lascaratos et al., 1999; Theocharis et al., 1992; Beuvier et al., 2012a). The EMT describes a change in the formation site for EMDW, when it temporarily switched from the Adriatic to the Aegean sub-basin.

In many respects, the most useful diagnostics of the ventilation of the ocean's interior come from geochemical tracers characterized by simple boundary conditions at the ocean's surface, and conservation in deep water (Key et al., 2004; Sarmiento and Gruber, 2006; Broecker and Peng, 1982). In particular, the passive transient tracers (CFC, ^{14}C , and tritium) do not affect the water mass densities (as opposed to active tracers such as temperature and salinity). Radiocarbon (^{14}C) is an ideal tracer for studying air-sea gas exchange, and for assessing the ventilation rate of the deep water masses at very long timescales (Toggweiler et al., 1989a, b). **Although ^{14}C is affected by biological processes, especially remineralization of organic matter, this effect can be considered minimal for the present simulation.**

Radiocarbon (^{14}C) is naturally formed by the reaction between nitrogen atoms in the atmosphere and slow-moving neutrons produced by a whole cascade of nuclear reactions between cosmic radiation and molecules in the upper atmosphere. Radiocarbon is not produced in the ocean's interior. All ^{14}C enters the ocean from the atmosphere through gas exchange with surface water with an equilibration time of 7-10 years (Broecker and Peng, 1982; Mahadevan, 2001). Radioactive decay of ^{14}C (the half-life being 5730 yr) reduces its concentration over time in the water column. Over the last 150 years, the natural distribution of radiocarbon has been disturbed by i) the dilution of atmospheric ^{14}C by the release of fossil fuel CO_2 , depleted in ^{14}C (the Suess Effect; Suess, 1955), and ii) the production of bomb ^{14}C by thermonuclear weapon testing in the late 1950s and early 1960s. The latter strongly increased the ^{14}C levels in the atmosphere (Rafter and Fergusson, 1957), and consequently the gradient between surface and subsurface waters (e.g., Broecker et al., 1985).

Knowledge of the timescale of the thermohaline circulation is of central importance in the debate on the sequestration of anthropogenic carbon in the deep ocean. Unlike the other tracers (e.g. CFC and Tritium), the radiocarbon concentration in the oceanic water masses is an invaluable tool allowing us to study the thermohaline circulation from the seasonal cycle, i.e., the near-surface circulation, vertical transport, and mixing (Naegler, 2009; Muller et al., 2006; Rodgers et al., 1997; Guilderson et al., 1998) at decadal and centennial timescales (e.g., Levin and Heshaimer, 2000; Stuiver et al., 1983). Radiocarbon plays a crucial role in carbon cycle investigations allowing us to assess the carbon fluxes between reservoirs (e.g., Levin et al., 2010), and the description of the air-sea gas exchange process (e.g., Wanninkhof, 1992; Sweeney et al., 2007).

Understanding the spatiotemporal variation of radiocarbon (Broecker and Peng, 1982) allows us to determine the ages of different water masses and to establish the overturning timescale and water-mass renewal time for individual basins and the global ocean (e.g., Matsumoto et al., 2004). Unlike other tracers, such as tritium, the ^{14}C in ocean surface water is not in equilibrium with atmosphere; this means that the surface ocean does not have the same ^{14}C age as the atmosphere (i.e., not zero age). This difference, also known as the "radiocarbon reservoir age" is caused both by the delay in exchange rates between atmospheric CO_2 and the carbonate system (Broecker and Peng, 1982), and the dilution effect due to the mixing of surface waters with intermediate or deep waters depleted in ^{14}C during seasonal vertical convection or upwelling, respectively. Indeed, when surface waters are isolated from the atmosphere, the radiocarbon clock begins to tick and ^{14}C content of water gradually decays.

Radiocarbon observations have played a crucial role as an experimental tool revealing the spatial and temporal variability of carbon sources and sinks (Roether and Weiss, 1980). Observational programmes (e.g., GEOSECS, WOCE and TTO) have provided snapshots of the large-scale distribution of radiocarbon in the world's oceans. However, few ^{14}C measurements have been made in the Mediterranean. Broecker and Gerard (1969) provided the first characterization of the natural radiocarbon in the surface and intermediate waters of the whole Mediterranean Sea from in-situ observations. More studies tried to determine the sea-surface radiocarbon reservoir ages of the Mediterranean, which are mainly affected by the Atlantic surface waters entering at Gibraltar, and/or by local factors related to freshwater input from rivers (Siani et al., 2000). The first ^{14}C reservoir age (360 ± 80 yr) was calculated by Broecker and Olson (1961) using the pre-bomb shells collected along the Algerian continental shelf. Later, Delibrias (1985) obtained an average ^{14}C reservoir age of 350 ± 35 yr through the analysis of pre-bomb mollusc shells from the French and Algerian shelves. The average marine reservoir age for the whole Mediterranean Sea was estimated by Siani et al. (2000) to be some 390 ± 85 yr.

Finally, mollusc shells were also used to yield a more significant dataset for the Mediterranean Sea with a mean sea-surface reservoir ^{14}C age of 400 ± 16 yr (Reimer and McCormac, 2002). More recently, the first annually-resolved sea-surface ^{14}C record was obtained from a 50-year-old shallow-water coral (*Cladocora caespitosa*) from the western Mediterranean Sea, covering the pre- and post-bomb period (Tisnérat-Laborde et al., 2013). However, all these observations are discrete both in time and/or space, they cannot give a clear description of radiocarbon evolution between the past and the actual situation now at either the regional or the global scale.

Although box-models have been extensively used to quantify the radiocarbon inventory (e.g., Broecker and Gerard, 1969; Craig, 1969), their application in deriving the oceanic distribution of radiocarbon is limited due to their very simple param-

terization. On the other hand, numerical modelling gives us a clear 4-D description of the water column, which provides an additional opportunity to better understand the ^{14}C distribution in seawater.

Several different ocean models have previously been used to study the global radiocarbon distribution (e.g., Toggweiler et al., 1989a, b; Duffy et al., 1995; Mouchet, 2013). However, these studies used coarse-resolution models which could not
5 represent satisfactorily the critical spatial and temporal scales of circulation in the Mediterranean Sea.

Here, we used a high-resolution regional model (NEMO-MED12, horizontal resolution $1/12^\circ$, ~ 7 km) of the entire Mediterranean Sea (Beuvier et al., 2010, 2012a). This model has been used previously for biogeochemical studies (Ayache et al., 2015b; Guyennon et al., 2015; Ayache et al., 2015a; Palmiéri et al., 2015; Ayache et al., 2016) and dynamical application (Soto-Navarro et al., 2014; Beuvier et al., 2012a; Lebeaupin Brossier et al., 2011). Here, we use the model to provide the first
10 simulation of radiocarbon distribution and the related reservoir age. The simulation covers the different states of ^{14}C from the steady natural distribution to the Suess effect, the ^{14}C bomb peak in the 1960s, and the post-bomb distribution until 2011. Our model results are compared to available ^{14}C measurements of seawater and marine carbonates reported by Broecker and Gerard (1969), Stuiver et al. (1983), Siani et al. (2000) and Tanhua et al. (2013), and to a 50-year high resolution ^{14}C record obtained from a shallow-water coral specimen (Tisnérat-Laborde et al., 2013).

Our work highlights the impact of anthropogenic perturbation (^{14}C bomb peak and the Suess effect) on the radiocarbon
15 distribution across the whole Mediterranean Sea, as well as the regional response across the different sub-basins. In addition, the simulation provides: i) constraints on the ^{14}C air-sea transfer; ii) a descriptive overview of the Mediterranean ^{14}C distribution, which gives an additional improvement of in-situ data interpretation, and iii) more perspectives on the impact of the interannual variability of the Mediterranean thermohaline circulation (e.g., EMT event) on the modelled ^{14}C distribution. **The present
20 radiocarbon simulation aims at improving the knowledge of the natural distribution of ^{14}C in the Mediterranean Sea and implementing a geochemical tracer with a longer time scale allowing more paleo-oriented applications.**

Furthermore, this study is part of the work under way to assess the robustness of the NEMO-MED12 model, and its use in studying the thermohaline circulation and the biogeochemical cycles in the Mediterranean Sea. The overarching objective of this work is to predict the future evolution of this basin under the increasing anthropogenic pressure.

25 **2 Method**

2.1 Circulation model

The Nucleus for European Modelling of the Ocean (NEMO) is a free surface-ocean circulation model (Madec and NEMO-
Team., 2008). Here, it is used in its Mediterranean configuration called NEMO-MED12 (Beuvier et al., 2012a) with a horizontal
30 resolution $1/12^\circ$ (~ 7 km), and 50 vertical z-coordinates ranging from 1 m at the surface to 450 m at depth with partial-step formulation.

NEMO-MED12 covers the whole Mediterranean Sea and includes part of the near Atlantic Ocean (buffer zone) from 11°W to 36°E and from 30°N to 47°N . The exchange with the Atlantic Ocean occurs through this buffer zone, where 3-D salinity and temperature fields are relaxed to the observed climatology (Beuvier et al., 2012a). The sea surface height (SSH) is restored

in the buffer zone from the GLORYS1 reanalysis (Ferry et al., 2010) in order to conserve the Mediterranean Sea water volume. The Black Sea is not explicitly represented in NEMO-MED12 configuration; exchanges with the Black Sea consist of a two-layer flow corresponding to the Dardanelles' net budget estimates of Stanev and Peneva (2002).

The atmospheric forcing of NEMO-MED12 is provided by daily mean fields of momentum, freshwater and heat fluxes from the high resolution atmospheric model (ARPERA) over the period 1958-2013 (Herrmann and Somot, 2008; Herrmann et al., 2010). The sea-surface temperature (SST) and water-flux correction term are applied using ERA-40 (Beuquier et al., 2012a). River runoff is derived from the interannual dataset of Ludwig et al. (2009) and Vörösmarty et al. (1996).

The initial conditions (temperature, salinity) are prescribed from the MedAtlas-II (Rixen et al., 2005; MEDAR-MedAtlas-group, 2002) climatology weighted by a low-pass filter with a time window of 10 years between 1955 and 1965 (Beuquier et al., 2012b). For the buffer zone (west of the Strait of Gibraltar) the initial state is based on the World Ocean Atlas 2005 (Antonov et al., 2006; Locarnini et al., 2006).

This model correctly simulates the main structures of the thermohaline circulation of the Mediterranean Sea, with mechanisms having a realistic timescale compared to observations (Ayache et al., 2015a). In particular, tritium (Ayache et al., 2015a) and helium isotope simulations (Ayache et al., 2015b) have shown that the EMT signal from the Aegean sub-basin is realistically simulated during early 1995. However, some aspects of the model still need to be improved: for instance the too-weak formation of Adriatic Deep Water (AdDW), followed by a low contribution to the EMDW in the Ionian sub-basin. In the western basin, the production of WMDW is reliable, but the spreading of the recently ventilated deep water to the south of the basin is too weak.

Full details of the model and its parameterizations are reported by Beuquier et al. (2012a, b), Palmiéri et al. (2015) and Ayache et al. (2015a).

2.2 The tracer model

The ^{14}C distribution in the ocean is often expressed as a delta notation relative to the $^{14}\text{C}/\text{C}$ ratio of the atmosphere ($\Delta^{14}\text{C} = (^{14}\text{R}/\text{Rref} - 1) \times 1000$; ^{14}R is the $^{14}\text{C}/\text{C}$ ratio of the ocean, and for the purpose of ocean ventilation studies Rref is set to one (Toggweiler et al., 1989a).

Here we use the approach of Toggweiler et al. (1989a, b) in which the ratio ^{14}R is transported by the model rather than the individual concentrations of C and ^{14}C . Several model studies adopted the simplified formulation of Toggweiler et al. (1989a) to describe the transport of ^{14}C in the ocean (Mouchet, 2013; Muller et al., 2006; Butzin et al., 2005; Orr et al., 2001; England and Rahmstorf, 1999; Maier-Reimer et al., 1993).

This approach is based on two main assumptions: i) the Dissolved Inorganic Carbon (DIC) field is constant and homogeneous, and ii) the air-sea fractionation processes and biological activity could be ignored (Mouchet, 2013; Toggweiler et al., 1989a). The first assumption reduces the capacity of the model to estimate the ^{14}C inventory and the ocean bomb- ^{14}C uptake (Mouchet, 2013) but does not much affect the equilibrium ^{14}C distribution in the ocean (Maier-Reimer et al., 1993; Orr et al., 2001; Mouchet, 2013). Modelled and observed ^{14}C may be directly compared since the observed ^{14}R ratios are corrected for the isotopic fractionation once converted to the standard $\Delta^{14}\text{C}$ notation (Stuiver and Polach, 1977).

This simplified approach is commonly used in model evaluation to critically examine the dynamics of the model (i.e., circulation and ventilation) against in-situ observation; because i) many oceanic ^{14}C data were obtained either by measuring ^{14}C in dissolved inorganic carbon in seawater or in corals and mollusc shells, and ii) it can be implemented in the ocean circulation models at relatively low computational cost allowing many sensitivity tests (e.g., Matsumoto et al., 2004).

- 5 Radiocarbon is implemented in the model as a passive conservative tracer, which does not affect ocean circulation. Hence its movement can be tracked in an off line mode using the pre-computed transport daily fields (U, V, W) of NEMO-MED12 dynamical model (Beuvier et al., 2012b). A time-step of 20 minutes is applied. The same approach was used to simulate the ϵNd (neodymium) distribution in the Mediterranean Sea (Ayache et al., 2016), and the mantle and crustal helium isotope signature (Ayache et al., 2015b), as well as to model the anthropogenic tritium invasion (Ayache et al., 2015a), and CFC and
10 anthropogenic carbon storage (Palmiéri et al., 2015).

Passive tracers are transported in the Mediterranean using a classical advection-diffusion equation, including the sources and sinks. The equation governing the transport of the dissolved inorganic carbon ^{14}R in the ocean is:

$$\frac{\delta}{\delta t} {}^{14}R = -\nabla \cdot (\mu {}^{14}R - K \cdot \nabla {}^{14}R) - \lambda {}^{14}R, \quad (1)$$

- Where λ is the radiocarbon decay rate, u the 3-D velocity field, and K the diffusivity tensor. Since radiocarbon is not
15 produced in the ocean, all ^{14}C enters the surface water through gas exchange. The radiocarbon flux through the sea-air boundary conditions is proportional to the difference in the ratios between the ocean and the atmosphere (Toggweiler et al., 1989a) and given as:

$$F = \kappa R ({}^{14}R - {}^{14}Ra), \quad (2)$$

- Where \mathcal{F} is the flux out of the ocean, and ${}^{14}Ra$ is the atmospheric $^{14}C/C$ ratio. The transfer velocity κR for the radiocarbon
20 ratio in Eq. (2) is computed as:

$$\kappa R = \frac{\kappa CO_2 K_0}{\overline{C_T}} P^a CO_2 \quad (3)$$

with κCO_2 being the carbon dioxide transfer velocity, K_0 the solubility of CO_2 in seawater taken from Weiss (1974), $P^a CO_2$ the atmospheric CO_2 pressure, and $\overline{C_T}$ the average sea-surface dissolved inorganic carbon concentration, classically set to $2 \text{ mol } m^{-3}$ (Toggweiler et al., 1989a; Orr et al., 2001; Butzin et al., 2005).

- 25 The CO_2 transfer velocity is computed with the help of surface-level wind speeds, w ($m \text{ s}^{-1}$), using the ARPERA forcing (Herrmann and Somot, 2008; Herrmann et al., 2010) following the Wanninkhof (1992) formulation:

$$\kappa CO_2 = k_w \times w^2 \sqrt{660/Sc}, \quad (4)$$

Where Sc is the Schmidt number computed with the model S and T fields.

The value of the empirical coefficient k_w depends on the wind field (Toggweiler et al., 1989a; Wanninkhof, 1992; Naegler, 2009). Sensitivity tests were performed to determine the value of the empirical coefficient k_w among the available values in the literature. Sensitivity tests were performed to determine the value of the empirical coefficient k_w among the available values in the literature. Accordingly, we have chosen a $k_w = 0.25 \times (0.01/3600) \text{ s m}^{-1}$ for our radiocarbon simulation, which produces the best agreement between model outputs and in-situ data for the pre-bomb period.

2.3 Model initialization and forcing

The natural radiocarbon distribution was first simulated using the atmospheric $^{14}R_a = 1$; the ocean ^{14}R is initially set to a constant value of 0.85 ($\Delta^{14}C = -150 \text{ ‰}$, appropriate for the deep ocean; (Key et al., 2004)). An atmospheric CO_2 of 280 ppm is prescribed for this steady state simulation. These simulations were integrated for 700 years using a 10-year interval of NEMO-MED12 circulation fields between 1965 and 1974 continuously repeated until they reached a quasi-steady state (i.e., the globally averaged drift was less than 0.001 ‰ per year). This forcing period was selected because it does not include any intense interannual variability, such as the event of the Eastern Mediterranean Transient (EMT, Roether et al. (2007); Schroeder et al. (2008)).

Starting from the end of the pre-industrial equilibrium run, the model was integrated from 1765 to 2011 covering the Suess effect (Suess, 1955), the entire radiocarbon (^{14}R) transient generated by the atmospheric nuclear weapon tests performed in the 1950s and early 1960s as well as the anthropogenic CO_2 increase. The ^{14}R level in the atmosphere (Fig.1) is taken from Orr et al. (2016) and references cited therein, and the atmospheric CO_2 from Orr et al. (2001). Unfortunately, there is no time series data of ^{14}C concentration around the Strait of Gibraltar. Hence simulated ^{14}C levels in the model's Atlantic water (AW) are determined by damping to global model estimates. The radiocarbon values in the buffer zone are prescribed from a global simulation of radiocarbon by Mouchet et al (2016). We have made two simulations with different boundary conditions at Gibraltar (see Supplement 1), the time series calculated from the larger box between 35°N and 55°N and from 0° to 46°W , improve a lot the radiocarbon simulation and the results are more realistic compared to some in-situ data (Tisnérat-Laborde et al., (2013); Tisnérat-Laborde, personal communication). So we have used this time series as boundary condition at Gibraltar to simulate ^{14}C in the Mediterranean Sea (see Supplement 1).

We also performed a sensitivity test on the impact of the EMT events on the radiocarbon distribution in the Mediterranean Sea. Two separate simulations were run for the period between 1990 and 2010 (i.e., covering the EMT event that occurred at the beginning of the 1990s). The NoEMT run was performed using the classical atmospheric forcing from ARPERA, as described in Sect. 2.1.

To improve dense-water fluxes through the Cretan Arc during the EMT (1992-1993) the ARPERA forcings were modified over the Aegean sub-basin (Beuvier et al., 2012a), by increasing mean values as done by Herrmann and Somot (2008) for the Gulf of Lions. More specifically, from November to March for the winters 1991-1992 and 1992-1993, daily surface heat loss was increased by 40 W m^{-2} , daily water loss by 1.5 mm and the daily wind stress modulus by 0.02 N m^{-2} . These changes accelerate the transfer of surface temperature and salinity perturbations into intermediate and deep layers of the Aegean sub-

basin, and improve the dense-water formation in the Aegean sub-basin during the EMT, with more intense mixing from winter convection.

3 Results

3.1 Steady state pre-bomb distribution

5 The ^{14}C model results of the radiocarbon natural distribution for March 1956 are expressed in $\Delta^{14}\text{C}$ (Fig.2) and in surface radiocarbon reservoir age (Fig.3). They provide a descriptive overview of the basin-wide distribution of radiocarbon before the anthropogenic perturbation. Figures 2a, 2b and 2c present the horizontal ^{14}C distribution of surface waters (between the surface and 200 m depth), intermediate (between 200 and 600 m) and deep waters (between 600 and 3500 m), respectively. Figures 2d and 2e show the radiocarbon distribution over the whole water column in the Mediterranean along a longitudinal
10 transect for both the eastern and western basins together with in-situ observations from Broecker and Gerard (1969). Figure 3 compares model results of reservoir ages and several marine reservoir ^{14}C age data available for the surface water of the Mediterranean; these data were obtained from pre-bomb calcareous marine shells between 1867 and 1948 and coral *Cladocora caespitosa* (Siani et al., 2000; Reimer and McCormac, 2002; Tisnérat-Laborde et al., 2013).

As illustrated in Figs. 2a and 3, there is a significant geographic heterogeneity in surface water for each sub-basin for "nat-
15 ural" (or pre-bomb) ^{14}C obtained both from model results and data. Table 1 shows that overall, the average $\Delta^{14}\text{C}$ values are generally lower in the WMed corresponding to older reservoir ^{14}C age (402 ± 27) compared to the EMed (349 ± 14), the Adriatic (373 ± 29) and the Aegean (349 ± 32) sub-basins that show younger reservoir ^{14}C ages than the data of Reimer and McCormac (2002). These figures clearly show the surface inflow of Atlantic waters through the Strait of Gibraltar were progressively enriched during their spreading into the EMed, leading to a relatively higher $\Delta^{14}\text{C}$ level in the EMed surface
20 water closer to -44‰ . For both western and eastern surface water, the model simulates ^{14}C concentrations slightly higher than the in-situ observations (Broecker and Gerard, 1969; Siani et al., 2000; Reimer and McCormac, 2002; Tisnérat-Laborde et al., 2013). A careful comparison between model outputs and seawater observations (1959) reveals a more pronounced disagreement, especially in the EMed surface water where the model overestimates the $\Delta^{14}\text{C}$ values by more than 10 ‰ (Fig.4a). However, the lack of more in-situ pre-bomb values greatly limits the comparison between model results and observations.

25 The model also simulates the rapid decrease of $\Delta^{14}\text{C}$ values with depth in the eastern basin, marking a significant vertical gradient and the most negative values of deep-water $\Delta^{14}\text{C}$ over the entire Mediterranean Sea ($-68 \pm 7 \text{‰}$). At depth, the model simulates low levels of $\Delta^{14}\text{C}$ in the eastern basin deep water (average value: $-64 \text{‰} \pm 7.4$), significantly lower than those simulated in the WMed deep waters (average value: $-48 \text{‰} \pm 6.9$) (Fig. 2c). To conclude, the model reproduced reasonably well the E-W gradient and the mean regional values of radiocarbon age, except for the Aegean sub-basin where the model
30 underestimates the regional mean value (Table 1), but the uncertainty in the data is also high.

3.2 Distribution of post-Bomb ^{14}C

The simulated bomb ^{14}C ocean distribution in the whole Mediterranean Sea in March 1977 is illustrated in Fig. 5. The large atmospheric $\Delta^{14}\text{C}$ increase is reasonably well captured by the model in the surface layer (values up to 120 ‰) over the whole basin. The lowest values are encountered in the known region of convection and formation of deep and intermediate waters (i.e., Gulf of Lions and Cyprus-Rhodes area; Fig.5a). Figure 5b shows a high concentration of radiocarbon at intermediate depths mainly in areas with recent water-mass ventilation. The radiocarbon distribution is more uniform in the deep water, except one location where relatively high radiocarbon levels are simulated in the deep layer as a result of mixing with the radiocarbon enriched surface water, particularly in the Cretan Sea (values up to ± 70 ‰) (Fig. 5c).

Figures 5d and 5e show the modelled $\Delta^{14}\text{C}$ results along vertical sections in the western and eastern basins compared with in-situ data obtained from seawater samples in the Ionian Sea during the GEOSECS expedition in 1977 (Station 404, 35.24°N, 17.12°E, Stuiver and ostlund, 1983). Similarly to the pre-bomb situation, the $\Delta^{14}\text{C}$ values decrease rapidly with depth, exhibiting a significant vertical gradient between the maximum in the surface water of around 120 ‰, and the minimum in the deep water values of around -50 ‰ in the western basin and around -60 ‰ in the eastern basin. The model correctly simulates the $\Delta^{14}\text{C}$ vertical distribution in the first 1500 m of the water column, in agreement with observations (Fig. 5e). At depth, the model tends to underestimate the ^{14}C penetration in the deep Ionian sub-basin, where it fails to reproduce the high $\Delta^{14}\text{C}$ levels associated with EMDW formation (Fig. 4b).

Figure 6 displays the modelled $\Delta^{14}\text{C}$ evolution between 1765 and 2008 for surface waters (average depth between 0-10 m in dashed line and between 0-100 m depths in solid line) in the Liguro-Provençal sub-basin, plotted against the in-situ values as reconstructed by Tisnerat-Laborde et al. (2013) from a 50-year old zooxanthellate coral *C. caespitosa* collected alive in 1998 along the coast of Bonassola (44°10'N, 9°36'E, NW Mediterranean, 28 m water depth), and from mollusc shells (Siani et al., 2000), Tisnerat-Laborde, personal communication).

Between 1900 and 1952, the modelled $\Delta^{14}\text{C}$ values show a slight decrease of ~ 12 ‰ resulting from the Suess effect (Druffel and Suess, 1983). The model slightly overestimates the observed pre-bomb mean value (-56 ± 3 ‰, in 1949-1955 Tisnérat-Laborde et al. (2013)) as noted previously. Between 1952 and 1980, the $\Delta^{14}\text{C}$ proxy values increase rapidly from -56 ‰ to almost + 85 ‰ in the Ligurian sub-basin due to a net uptake of atmospheric bomb ^{14}C .

The model represents well the uptake of bomb ^{14}C for the top layer (0-10m) and the sub-surface layer (0-100m) until 1965. Then, a slight difference of $\Delta^{14}\text{C}$ is simulated between these two layers, with a higher value in the top layer that is consistent with the observations. These differences are the result of vertical convective mixing (Mahadevan, 2001), **i.e. the mixing layer depth could impact the amplitude of $\Delta^{14}\text{C}$ peak in the surface layer**. Afterwards, the $\Delta^{14}\text{C}$ values decreased slowly with fluctuations, but reaching a value around +50 ‰ in 2008. This gradual decline of $\Delta^{14}\text{C}$ (values up to + 60 ‰) is well simulated in the surface water **when we compare** the modelled present day (March 2011) distribution of radiocarbon in the surface water (Fig. 7). These results demonstrate that the model simulates the bomb ^{14}C uptake in surface and sub-surface water with a realistic timescale comparable to in-situ data, and shows a good consistency between the observed and simulated bomb $\Delta^{14}\text{C}$ annual average value (Fig. 6).

Figure 7a shows the modelled present day (March 2011) distribution of radiocarbon in the surface water, against Meteor M84/3 cruise data (Tanhua et al., 2013). The $\Delta^{14}C$ distribution pattern for the surface water is similar to the model outputs obtained for years 1956 and 1977, with the eastern basin generally showing higher $\Delta^{14}C$ values compared to the western basin, except in areas of formation of deep and intermediate waters in the Mediterranean Sea (the Cyprus-Rhodes area, and in the Gulf of Lions), where the $\Delta^{14}C$ concentration decreases rapidly due to higher vertical convection (Fig. 7).

Figures 7d and 7e, present the simulated radiocarbon content for March 2011, at intermediate and deep depth along a W-E transect together with the available Meteor M84/3 cruise data (Tanhua et al., 2013). The two vertical sections show a $\Delta^{14}C$ maximum in the first 500 m depth ($\Delta^{14}C > 40 \text{ ‰}$). At deeper depths, $\Delta^{14}C$ values exhibit a significant vertical gradient up to 1500 m (Fig. 8), with low $\Delta^{14}C$ values simulated for the deep waters (values lower than -40 ‰), except for the central Levantine (i.e the area south of Crete sub-basin) and deep water, where high values $\Delta^{14}C$ are simulated (around -20 ‰) due to the intense deep convection in this areas. **Relatively high values are simulated in the Algerian basin (around -30 ‰ , Fig. 7d).**

The model correctly reproduces the $\Delta^{14}C$ content of the surface waters as noted previously, with values similar to observations (values about $+50 \text{ ‰}$, Fig. 7 and Fig. 8). For the deeper depths, the simulated $\Delta^{14}C$ levels tend to be underestimated by more than 20 ‰ in the WMed and by about 50 ‰ in the EMed compared to the observations. This is the result of the too-weak deep water overflow through the Otranto Strait from the Adriatic into the Ionian sub-basin, and the weak southern penetration of the new WMDW in the simulation compared to the values deduced from in-situ observations (Beuvier et al., 2012a, b; Ayache et al., 2015a). This underestimation leads to excessively low $\Delta^{14}C$ average values at depth of the eastern basin. However the model simulates well the $\Delta^{14}C$ values in the surface and deep water of **the Adriatic sub-basin** (Figure 7a and 7c) compared to Meteor M84/3 cruise data (Tanhua et al., 2013).

20 3.3 The spatial and temporal variability

The temporal variability of radiocarbon distribution was explored as a function of sub-basin location (Fig. 9). Specifically, we compared the annual average $\Delta^{14}C$ time series in different "boxes" following the LIW trajectory from the Levantine sub-basin to the Gulf of Lions (including the Tyrrhenian, Ligurian, Algerian and Cretan sub-basins) for the surface (Fig. 9a), intermediate (Fig. 9b), deep (Fig. 9c) and whole water column (Fig. 9d).

25 The $\Delta^{14}C$ evolution of the surface water is very similar within the different sub-basins until 1965. Afterwards, Tyrrhenian, Algerian and the Ligurian sub-basins have similar bomb $\Delta^{14}C$ peak record, while the Gulf of Lions, Levantine basin and Cretan Sea respond differently to the bomb-signal compared to the other sub-basins. The Levantine/Cretan Sea and the Gulf of Lions show surface values as high as 100 ‰ and as low as 60 ‰ , respectively (Fig. 9a). The differences between the western and eastern basins are more pronounced at intermediate depths (Fig. 9b), especially between the Cretan Sea and the Gulf of Lions, which shows an almost 40 ‰ difference in $\Delta^{14}C$. The Algerian and Ligurian sub-basins are characterized by a very similar $\Delta^{14}C$ evolution through time, showing intermediate values between the Cretan Sea and the Gulf of Lions. The results for the Tyrrhenian sub-basin and Cretan Sea indicate higher transfer in intermediate water compared to other sub-basins. Model outputs for the deep layers (600-3500 m) reveal much higher $\Delta^{14}C$ levels in the Cretan Sea compared to the other locations (Fig. 9c) because it has shallower bottom depth. The $\Delta^{14}C$ difference across the six sub-basins is more pronounced at deeper

depths than at the surface (Fig. 9a) and intermediate layers (Fig. 9b), especially after the ^{14}C bomb peak. This difference decreases gradually after 1995, particularly in the surface water where the $\Delta^{14}\text{C}$ values are almost the same among the different sub-basins. **The integrated values for the whole water column (Fig. 9d) show the same pattern as seen in the deep waters (Fig. 9c), suggesting a strong role of deep layers in controlling the distribution of $\Delta^{14}\text{C}$ in the water column.**

5 The impact of the EMT event on the radiocarbon distribution in the Mediterranean was analyzed by comparing the outputs of two model simulations (shown in Fig. 10 and Fig. 11): "EMT" and "NoEMT" for the years of 1995 and 1999, respectively (see Sect. 2.3). A substantial penetration of radiocarbon is observed in the deep water south of Crete in 1995, as a consequence of the EMT event that increased bottom $\Delta^{14}\text{C}$ values by more than 60 ‰, close to ^{14}C bomb peak values. On the other hand, the EMT reduces the $\Delta^{14}\text{C}$ value in the intermediate waters in the EMed (Fig. 10b). The EMT-related $\Delta^{14}\text{C}$ signal in the deep
10 waters decreases gradually after the event, with values around 30 ‰ in 1999 (Fig. 10d). For the WMed (Fig. 10a, 10c), the contrast is particularly pronounced at intermediate levels, with regional values shifted by almost 10 ‰ between 200 and 800 m depth in the Algerian basin (Fig. 10a), as a consequence of the abrupt change in the eastern basin during the EMT event. As shown in Fig. 11, the shift begins in 1992 in the Levantine sub-basin and reached a 60 ‰ difference in 1995 between these two simulations (Figs. 11a, 11b).

15 4 Discussion

The radiocarbon simulations provide independent and additional constraints on the thermohaline circulation and deep-water ventilation in the Mediterranean Sea. The relatively simple approach of radiocarbon modelling adopted here from Toggweiler et al. (1989a) and Mouchet et al (2016) using a high resolution regional model, led to a realistic simulation of the radiocarbon distribution relative to available in-situ data. It also enables the evaluation of the NEMO-MED12 model performance in the
20 Mediterranean Sea from the seasonal to decadal and centennial timescales. Furthermore, it provides a unique opportunity to better constrain the variability of the uptake of bomb ^{14}C in the whole Mediterranean Sea and to study the impact of important hydrological events such as the Eastern Mediterranean Transient (EMT).

The modelled radiocarbon distribution is very sensitive to the value of the empirical coefficient (K_W) (i.e. is the constant regulating air-sea flux). In this study we have used $K_W = 0.25 \times (0.01/3600) \text{ s } m^{-1}$, this value led to a better simulation
25 of $\Delta^{14}\text{C}$ in the Mediterranean compared to the other estimates available in the literature, i.e., $0.426 \times (0.01/3600) \text{ s } m^{-1}$ used in global scale simulation (Mouchet et al, 2016; Naegler, 2009). The K_W value depends on the wind field and the upper ocean mixing rate field (Wanninkhof, 1992; Toggweiler et al., 1989a). For the present work we used the wind fields from the ARPERA forcing (Herrmann and Somot, 2008; Herrmann et al., 2010) and the atmospheric CO_2 values from Orr et al. (2016). These boundary conditions enabled the model to produce satisfactory simulations of the bomb ^{14}C chronology. In particular,
30 the timing of the $\Delta^{14}\text{C}$ peak in the surface is consistent with the estimated $^{14}\text{CO}_2$ time transfer from the atmosphere to the ocean in the surface waters (~ 10 yr, (Broecker and Peng, 1982) as shown in Fig. 6.

Unlike the global ocean, where input/output of radiocarbon comes only from the exchange with the atmosphere, in the Mediterranean Sea there is also lateral exchange of ^{14}C through the Strait of Gibraltar. Unfortunately, there is no time series

data of ^{14}C concentration in that area. Hence simulated ^{14}C levels in the model's Atlantic water (AW) are determined by damping to global model estimates from Mouchet et al (2016) at the western boundary of the model domain, using the 3-D profile calculated between 35°N and 55°N and from 0° to 46°W (sensitivity tests were performed to determine this box). This large box in the North Atlantic gave the most representative signature of radiocarbon during the bomb peak (value up to 140
5 ‰ in 1980) from the global simulation of Mouchet et al (2016).

The comparison between the model outputs and the sea-surface $\Delta^{14}\text{C}$ record (Fig. 6) obtained from a 50-year-old shallow-water coral in the western Mediterranean Sea from (Tisnérat-Laborde et al., 2013) reveals a good model performance in simulating the bomb/post-bomb radiocarbon distribution (Fig. 4b, Fig. 8). However the representation of the pre-bomb distribution is more difficult in the simulation (Fig. 4a). Several issues complicate the simulation of the natural steady state distribution of
10 ^{14}C using ocean-model circulation: i) the uncertainty associated with the radiocarbon surface boundary conditions applied in ocean model experiments, ii) the climatological field to represent the wind forcing, often based on atmospheric model outputs and/or historical data, and iii) the significant changes due to the human activity which affects the radiocarbon distribution in the atmosphere and the ocean (e.g., Suess effect). In addition, the limited spatial and temporal resolution of seawater and carbonate organism measurements during the pre-bomb period limits our understanding of the natural radiocarbon distribution in
15 the Mediterranean Sea.

On the other hand, the ^{14}C reservoir ages for this period are exclusively localized over the continental shelf (mainly reconstructed from shallow-water corals and molluscs). These proxy data reveal a high regional variability as reconstructed by Siani et al. (2000) between 1837 and 1951, and Reimer and McCormac (2002), that can be attributed to both: (i) the interactions between the ocean and land by the transport of depleted freshwater, and (ii) the potential changes in the vertical mixing of the
20 water column, with an increase of air-sea CO_2 exchanges. These processes could be favoured by the atmospheric conditions, such as the North Atlantic Oscillation (NAO), East Atlantic Pattern (EA), East Atlantic/West Russian pattern (EA/WR) within stronger and frequent wind storms and stronger precipitation over northern Europe (Josey et al., 2011).

After the ^{14}C bomb peak, a large gradient of $\Delta^{14}\text{C}$ existed between the surface waters already enriched and saturated in bomb ^{14}C (values up to 120 ‰) and intermediate/deep waters with relatively low $\Delta^{14}\text{C}$ level (Fig. 5), associated with the long
25 equilibration time with the radiocarbon depleted deep waters, and to vertical mixing. Nevertheless the model simulation shows that the bomb-produced radiocarbon signal has reached the deep layers of the Mediterranean Sea due to the rapid transfer of surface waters to intermediate and deep depths, especially in the Cretan Sea, where a high $\Delta^{14}\text{C}$ is simulated in the deep waters (Fig. 7).

The new $\Delta^{14}\text{C}$ data obtained from the analysis of the seawater samples collected during the Meteor M84/3 cruise represent a
30 unique opportunity to critically assess the dynamics of the NEMO-MED12 ocean model and to evaluate its ability to reproduce the main features of the present day radiocarbon distribution in the Mediterranean. The model produces realistic simulated $\Delta^{14}\text{C}$ values in the surface layer that are in agreement with in-situ measurements, thus supporting our modelling approach. However, some important aspects of the model still need to be improved, particularly for deep water, where it underestimates $\Delta^{14}\text{C}$ (Fig. 7, Fig. 8). Previous passive tracer evaluations of NEMO-MED12 have shown that the ventilation rates of deep
35 waters are underestimated by the model for the whole Mediterranean (e.g., Ayache et al., 2015a; Palmiéri et al., 2015).

This is particularly evident in the Ionian sub-basin where the Eastern Mediterranean Deep Water is not properly simulated due to the too-weak formation of Adriatic Deep Water that flows at shallower depths compared to the observations. Similarly, the southwards propagation of the newly-formed WMDW in the model is slower than the observations as a consequence of a reduced salinity content (and hence density) in the formation area. Finally, tritium and helium (Ayache et al., 2015a) and CFC (Palmiéri et al., 2015) simulations have shown that the model overestimates the mixing near the Cretan Arc .

Several factors could control the radiocarbon distribution across the Mediterranean Sea. During the pre-industrial period, the AW inflow at Gibraltar, together with freshwater input from rivers could have played an important role on the radiocarbon distribution in the Mediterranean. The large amount of radiocarbon injected in the atmosphere during the thermonuclear weapon testing is now the dominant control on the ^{14}C distribution in the surface water, completely masking the natural radiocarbon background. This creates the opportunity to study the constraints on the $^{14}\text{CO}_2$ air-sea exchange. On the other hand, the ventilation rate is the key mechanism and the most important factor controlling the ^{14}C distribution in the deep layer.

The model has provided, for the first time, the evolution of $\Delta^{14}\text{C}$ in different parts of the basin and at different depths (Fig. 9). The difference in $\Delta^{14}\text{C}$ in surface water between the western and eastern basins reveals enrichment of $\Delta^{14}\text{C}$ along surface water-mass pathway due to prolonged exposure of the surface water to the atmosphere. It also shows the different mechanism of ^{14}C transfer at depth, where it depends on convection processes with higher convection occurring especially during the bomb peak with a large amount of radiocarbon in the atmosphere and the surface water masses undergo transfer with different intensity in the different sectors of the Mediterranean Basin.

The sequence of EMT events that occurred at the beginning of the 1990s in the eastern Mediterranean has substantially changed the deep water-mass structure in the whole basin. Different hypotheses concerning the preconditioning of the EMT and its timing have been proposed in the literature (Roether et al., 2007; Beuvier et al., 2012a; Lascaratos et al., 1999; Theocharis et al., 1999; Klein et al., 1999; Stanev and Peneva, 2002; Josey, 2003). The renewal of the deep water masses after the EMT is satisfyingly simulated by our regional model as illustrated by tritium-helium3 (Ayache et al., 2015a) and by neodymium simulations (Ayache et al., 2016). These findings allow us to study the impact of interannual variability on a very long timescale, including the exceptional events observed in the ventilation of the deep waters. The radiocarbon simulation documents a severe impact of the EMT on the water mass distribution, through the transfer of a large volume of ^{14}C -enriched near-surface water into the deep layers, with the highest contribution being observed in the area south of the Cretan Arc.

The EMT event generates an important accumulation of ^{14}C -enriched water at the bottom of the Levantine sub-basin with more than 60 ‰ of difference in 1995 compared to the pre-EMT situation. In our simulation the LIW layer is also affected by low values in the eastern Mediterranean, where the renewal of the bottom water masses (low concentration of radiocarbon) during the EMT could lead to a decrease in the ^{14}C content in the LIW layer (200-600 m, Fig. 10). On the other hand, higher values of radiocarbon are simulated at intermediate levels in the western Mediterranean during the EMT, with shifts up to 10 ‰ compared to the No-EMT values. During the EMT, part of the Levantine basin is filled by water masses originating in the Aegean Sea, with different characteristics compared to the Adriatic. Hence the EMT could modify water mass characteristics and potentially affect the formation of deep water masses in this basin.

35 5 Conclusions

The radiocarbon distribution of the whole Mediterranean Sea was simulated for the first time using a high-resolution model (NEMO-MED12) at $1/12^\circ$ horizontal resolution and compared to available in-situ measurements and proxy-based reconstructions. The present study provides a unique opportunity to improve the interpretation/understanding of the available in-situ data, and could help in the design of new observational programmes for the Mediterranean Sea. It also provides a new approach to understanding and better constraining air-sea gas exchange and dynamics of Mediterranean water masses over the last decade. The air-sea exchange parameterization led to a realistic simulation of bomb ^{14}C in the surface water, compared to in-situ data. The model correctly simulates the main features of radiocarbon distribution during and after the ^{14}C bomb perturbation, especially in the surface/intermediate layers. On the other hand, severe mismatches between model and observations in the deep layer are clearly associated with shortcomings in the model parameterization.

10 The natural distribution of ^{14}C in the Mediterranean Sea is mainly affected by the inflow of Atlantic water through the Strait of Gibraltar. Further, the large amount of radiocarbon injected into the atmosphere during the nuclear bomb-testing period has been the dominant factor defining the ^{14}C distribution in the surface water, largely masking the natural radiocarbon background. More paleo-data from the pre-industrial period would help improve the knowledge of the natural distribution of ^{14}C in the Mediterranean and better constrain the fluxes and exchange of radiocarbon between the different reservoirs.

15 This ^{14}C modelling provides a unique opportunity to explore the impact of the interannual variability on the radiocarbon distribution in the whole Mediterranean Sea and the interaction between its western and eastern basins. The outputs of the model simulation of the EMT event reveal a significant increase in $\Delta^{14}\text{C}$ (by more than 60 ‰) in the Aegean deep water, and at intermediate level (value up to 10 ‰) in the western basin. The model results with/without EMT show that the vertical transport of surface signals in the Mediterranean is strong, suggesting a major contribution of the EMT in the accumulation of radiocarbon in the eastern Mediterranean deep waters. Although the approach we adopted does not attempt to quantify the anthropogenic carbon, the model results and observations on the ^{14}C distribution support the contention that a large amount of anthropogenic carbon is being stored in the deep Mediterranean waters, in agreement with previous findings (e.g., Palmiéri et al., 2015).

6 Code availability

25 The model used in this work is a free surface ocean general circulation model NEMO (Madec and NEMO-Team., 2008) in a regional configuration called NEMO-MED12 (Beuvier et al., 2012a). (<http://www.nemo-ocean.eu/>)

References

- Antonov, J. I., Locarnini, R. A., Boyer, T. P., Mishonov, A. V., and Garcia, H. E.: World Ocean Atlas 2005, Volume 2: Salinity, S. Levitus, Ed, NOAA Atlas NESDIS 62, U.S. Government Printing Office, Washington, D.C., p. 182 pp, 2006.
- Attané, I. and Courbage, Y.: Demography in the Mediterranean region: situation and projections, Plan Bleu. Paris, Economica, 2004.
- Ayache, M., Dutay, J.-C., Jean-Baptiste, P., Beranger, K., Arsouze, T., Beuvier, J., Palmieri, J., Le-vu, B., and Roether, W.: Modelling of the anthropogenic tritium transient and its decay product helium-3 in the Mediterranean Sea using a high-resolution regional model, *Ocean Science*, 11, 323–342, doi:10.5194/os-11-323-2015, 2015a.
- Ayache, M., Dutay, J.-C., Jean-Baptiste, P., and Fourré, E.: Simulation of the mantle and crustal helium isotope signature in the Mediterranean Sea using a high-resolution regional circulation model, *Ocean Science*, 11, 965–978, doi:10.5194/os-11-965-2015, 2015b.
- Ayache, M., Dutay, J.-C., Arsouze, T., Révillon, S., Beuvier, J., and Jeandel, C.: High-resolution neodymium characterization along the Mediterranean margins and modelling of ϵ Nd distribution in the Mediterranean basins, *Biogeosciences*, 13, 5259–5276, doi:10.5194/bg-13-5259-2016, <http://www.biogeosciences.net/13/5259/2016/>, 2016.
- Beuvier, J., Sevault, F., Herrmann, M., Kontoyiannis, H., Ludwig, W., Rixen, M., Stanev, E., Béranger, K., and Somot, S.: Modeling the Mediterranean Sea interannual variability during 1961–2000: Focus on the Eastern Mediterranean Transient, *Journal of Geophysical Research*, 115, C08 017, doi:10.1029/2009JC005950, 2010.
- Beuvier, J., Béranger, K., Lebeaupin Brossier, C., Somot, S., Sevault, F., Drillet, Y., Bourdallé-Badie, R., Ferry, N., and Lyard, F.: Spreading of the Western Mediterranean Deep Water after winter 2005: Time scales and deep cyclone transport, *Journal of Geophysical Research*, 117, C07 022, doi:10.1029/2011JC007679, 2012a.
- Beuvier, J., Lebeaupin Brossier, C., Béranger, K., Arsouze, T., Bourdallé-Badie, R., Deltel, C., Drillet, Y., Drobinski, P., Lyard, F., Ferry, N., Sevault, F., , and Somot, S.: MED12, Oceanic component for the modelling of the regional Mediterranean Earth System, *Mercator Ocean Quarterly Newsletter*, 46, 60–66, 2012b.
- Broecker, S. and Gerard, R.: Natural radiocarbon in the Mediterranean Sea, Lamont-Doherty Geological Observatory of Columbia University, Palisades, New York 10964, 1969.
- Broecker, W. and Peng, T.-H.: Tracers in the Sea: Palisades, NY (Lamont-Doherty Geol. Observ.), 1982.
- Broecker, W. S. and Olson, E. A.: Lamont Radiocarbon Measurements VIII, *Radiocarbon*, 3, 176–204, doi:10.2458/AZU_JS_RC.3.1080, <https://journals.uair.arizona.edu/index.php/radiocarbon/article/view/1080>, 1961.
- Broecker, W. S., Peteet, D. M., and Rind, D.: Does the ocean–atmosphere system have more than one stable mode of operation?, *Nature*, 315, 21–26, doi:10.1038/315021a0, <http://www.nature.com/doi/10.1038/315021a0>, 1985.
- Butzin, M., Prange, M., and Lohmann, G.: Radiocarbon simulations for the glacial ocean: The effects of wind stress, Southern Ocean sea ice and Heinrich events, doi:10.1016/j.epsl.2005.03.003, 2005.
- Craig, H.: Abyssal carbon and radiocarbon in the Pacific, *Journal of Geophysical Research*, 74, 5491–5506, doi:10.1029/JC074i023p05491, <http://doi.wiley.com/10.1029/JC074i023p05491>, 1969.
- Diffenbaugh, N. S. and Giorgi, F.: Climate change hotspots in the CMIP5 global climate model ensemble., *Climatic change*, 114, 813–822, doi:10.1007/s10584-012-0570-x, 2012.
- Druffel, E. M. and Suess, H. E.: On the radiocarbon record in banded corals: Exchange parameters and net transport of $^{14}\text{CO}_2$ between atmosphere and surface ocean, *Journal of Geophysical Research*, 88, 1271, doi:10.1029/JC088iC02p01271, <http://doi.wiley.com/10.1029/JC088iC02p01271>, 1983.

- Duffy, P. B., Eliason, D. E., Bourgeois, A. J., and Covey, C. C.: Simulation of bomb radiocarbon in two global ocean general circulation models, *Journal of Geophysical Research*, 100, 22 545, doi:10.1029/95JC01937, <http://doi.wiley.com/10.1029/95JC01937>, 1995.
- England, M. H. and Rahmstorf, S.: Sensitivity of Ventilation Rates and Radiocarbon Uptake to Subgrid-Scale Mixing in Ocean Models, *Journal of Physical Oceanography*, 29, 2802–2828, doi:10.1175/1520-0485(1999)029<2802:SOVRAR>2.0.CO;2, 1999.
- Ferry, N., Parent, L., Garric, G., Barnier, B., and Jourdain, N. C.: Mercator Global Eddy Permitting Ocean Reanalysis GLORYS1V1: Description and Results, *Mercator Ocean Quarterly Newsletter*, 36, 15–28, 2010.
- Giorgi, F.: Climate change hot-spots, *Geophysical Research Letters*, 33, doi:10.1029/2006GL025734, 2006.
- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Global and Planetary Change*, 63, 90–104, doi:10.1016/j.gloplacha.2007.09.005, 2008.
- Guilderson, T. P., Schrag, D. P., Kashgarian, M., and Southon, J.: Radiocarbon variability in the western equatorial Pacific inferred from a high-resolution coral record from Nauru Island, *Journal of Geophysical Research: Oceans*, 103, 24 641–24 650, doi:10.1029/98JC02271, <http://doi.wiley.com/10.1029/98JC02271>, 1998.
- Guyennon, A., Baklouti, M., Diaz, F., Palmieri, J., Beuvier, J., Lebaupin-Brossier, C., Arsouze, T., Béranger, K., Dutay, J.-C., and Moutin, T.: New insights into the organic carbon export in the Mediterranean Sea from 3-D modeling, *Biogeosciences*, 12, 7025–7046, doi:10.5194/bg-12-7025-2015, 2015.
- Herrmann, M., Sevault, F., Beuvier, J., and Somot, S.: What induced the exceptional 2005 convection event in the northwestern Mediterranean basin? Answers from a modeling study, *Journal of Geophysical Research*, 115, C12 051, doi:10.1029/2010JC006162, 2010.
- Herrmann, M. J. and Somot, S.: Relevance of ERA40 dynamical downscaling for modeling deep convection in the Mediterranean Sea, *Geophysical Research Letters*, 35, L04 607, doi:10.1029/2007GL032442, 2008.
- IPCC: Climate Change 2013 .: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xi, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, p. 1535, doi:10.1017/CBO9781107415324., <http://www.climatechange2013.org/report/full-report/>, 2013.
- Josey, S. A.: Changes in the heat and freshwater forcing of the eastern Mediterranean and their influence on deep water formation, *Journal of Geophysical Research*, 108, 3237, doi:10.1029/2003JC001778, 2003.
- Josey, S. A., Somot, S., and Tsimplis, M.: Impacts of atmospheric modes of variability on Mediterranean Sea surface heat exchange, *Journal of Geophysical Research*, 116, C02 032, doi:10.1029/2010JC006685, <http://doi.wiley.com/10.1029/2010JC006685>, 2011.
- Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cycles*, 18, doi:10.1029/2004GB002247, 2004.
- Klein, B., Roether, W., Manca, B., Bregant, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: The large deep water transient in the Eastern Mediterranean, *Deep-Sea Research*, 46, 371 – 414, 1999.
- Lascaratos, A., Roether, W., Nittis, K., and Klein, B.: Recent changes in deep water formation and spreading in the eastern Mediterranean Sea: a review, *Progress in Oceanography*, 44, 5–36, doi:10.1016/S0079-6611(99)00019-1, 1999.
- Lebaupin Brossier, C., Béranger, K., Deltel, C., and Drobinski, P.: The Mediterranean response to different space–time resolution atmospheric forcings using perpetual mode sensitivity simulations, *Ocean Modelling*, 36, 1–25, doi:10.1016/j.ocemod.2010.10.008, 2011.
- Levin, I. and Heshaimer, V.: Radiocarbon – a unique tracer of global carbon cycle dynamics, *RADIOCARBON*, 42, 69–80, 2000.

- Levin, I., Naegler, T., Kromer, B., Diehl, M., Francey, R. J., Gomez-Pelaez, A. J., Steele, L. P., Wagenbach, D., Weller, R., and Worthy, D. E.: Observations and modelling of the global distribution and long-term trend of atmospheric $^{14}\text{CO}_2$, *Tellus B*, 62, doi:10.3402/tellusb.v62i1.16511, <http://www.tellusb.net/index.php/tellusb/article/view/16511>, 2010.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., and Garcia, H. E.: *World Ocean Atlas 2005, Volume 1: Temperature*, S. Levitus, Ed, NOAA Atlas NESDIS 61, U.S. Government Printing Office, Washington, D.C. 182 pp, 2006.
- 5 Ludwig, W., Dumont, E., Meybeck, M., and Heussner, S.: River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades?, *Progress in Oceanography*, 80, 199–217, doi:10.1016/j.pocean.2009.02.001, 2009.
- Madec, G. and NEMO-Team.: Note du Pole de modélisation, Institut Pierre-Simon Laplace (IPSL), France, NEMO ocean engine, 27, doi:ISSN N1288-1619, 2008.
- 10 Mahadevan, A.: An analysis of bomb radiocarbon trends in the Pacific, *Marine Chemistry*, 73, 273–290, doi:10.1016/S0304-4203(00)00113-4, 2001.
- Maier-Reimer, E., Mikolajewicz, U., and Hasselmann, K.: Mean Circulation of the Hamburg LSG OGCM and Its Sensitivity to the Thermal Surface Forcing, *Journal of Physical Oceanography*, 23, 731–757, doi:10.1175/1520-0485(1993)023<0731:MCOTHL>2.0.CO;2, 1993.
- 15 Malanotte-Rizzoli, P., Manca, B. B., D'Alcala, M. R., Theocharis, A., Brenner, S., Budillon, G., and Ozsoy, E.: The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations, *Dynamics of Atmospheres and Oceans*, 29, 365–395, doi:10.1016/S0377-0265(99)00011-1, 1999.
- Matsumoto, K., Nishihara, S., Kamimura, M., Shiraiishi, T., Otaguro, T., Uehara, M., Maeda, Y., Ogura, K., Lumsden, A., and Ogura, T.: The prepattern transcription factor *Irx2*, a target of the FGF8/MAP kinase cascade, is involved in cerebellum formation, *Nature Neuroscience*, 7, 605–612, doi:10.1038/nn1249, <http://www.nature.com/doi/finder/10.1038/nn1249>, 2004.
- 20 MEDAR-MedAtlas-group: Medar-Medatlas Protocol (Version 3) Part I: Exchange Format and Quality Checks for Observed Profiles, P. Rap. Int. IFREMER/TMSI/IDM/SIS002-006, 50, 2002.
- MerMex-Group: Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, *Progress in Oceanography*, 91, 97–166, doi:10.1016/j.pocean.2011.02.003, 2011.
- 25 Millot, C. and Taupier-Letage, I.: *The Mediterranean Sea*, vol. 5K, doi:10.1007/b107143., 2005.
- Mouchet, A.: The Ocean Bomb Radiocarbon Inventory Revisited, *Radiocarbon*, 55, doi:10.2458/azu_js_rc.55.16402, <https://journals.uair.arizona.edu/index.php/radiocarbon/article/view/16402>, 2013.
- Mouchet et al, A.: Assessing ventilation in NEMO v3.6 : radiocarbon and age., To be submitted to *Geophys. Mod. Dev.* NEMO special issue, 2016.
- 30 Muller, S. A., Joos, F., Edwards, N. R., and Stocker, T. F.: Water Mass Distribution and Ventilation Time Scales in a Cost-Efficient, Three-Dimensional Ocean Model, *J. Climate*, 19, 5479–5499, doi:10.1175/JCLI3911.1., 2006.
- Naegler, T.: Reconciliation of excess ^{14}C -constrained global CO_2 piston velocity estimates, *Tellus B*, 61, 372–384, doi:10.1111/j.1600-0889.2008.00408.x, <http://www.tellusb.net/index.php/tellusb/article/view/16836>, 2009.
- Orr, J. C., Maier-Reimer, E., Mikolajewicz, U., Monfray, P., Sarmiento, J. L., Toggweiler, J. R., Taylor, N. K., Palmer, J., Gruber, N., Sabine, C. L., Le Quéré, C., Key, R. M., and Boutin, J.: Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models, *Global Biogeochemical Cycles*, 15, 43–60, doi:10.1029/2000GB001273, <http://doi.wiley.com/10.1029/2000GB001273>, 2001.
- 35

- Orr, J. C., Najjar, R. G., Aumont, O., Bopp, L., Bullister, J. L., Danabasoglu, G., Doney, S. C., Dunne, J. P., Dutay, J.-C., Graven, H., Griffies, S. M., John, J. G., Joos, F., Levin, I., Lindsay, K., Matear, R. J., McKinley, G. A., Mouchet, A., Oschlies, A., Romanou, A., Schlitzer, R., Tagliabue, A., Tanhua, T., and Yool, A.: Biogeochemical protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project (OMIP), *Geoscientific Model Development Discussions*, pp. 1–45, doi:10.5194/gmd-2016-155, <http://www.geosci-model-dev-discuss.net/gmd-2016-155/>, 2016.
- 5 Palmiéri, J., Orr, J. C., Dutay, J.-C., Béranger, K., Schneider, A., Beuvier, J., and Somot, S.: Simulated anthropogenic CO₂ storage and acidification of the Mediterranean Sea, *Biogeosciences*, 12, 781–802, doi:10.5194/bg-12-781-2015, 2015.
- Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: a review, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 158, 153–173, doi:10.1016/S0031-0182(00)00048-1, 2000.
- Reimer, P. and McCormac, F.: Marine Radiocarbon Reservoir Corrections for the Mediterranean and Aegean Seas, *Radiocarbon*, 44, 159–
- 10 166, doi:10.2458/AZU_JS_RC.44.4088, 2002.
- Rixen, M., Beckers, J. M., Levitus, S., Antonov, J., Boyer, T., Maillard, C., Fichaut, M., Balopoulos, E., Iona, S., Dooley, H., Garcia, M. J., Manca, B., Giorgetti, A., Manzella, G., Mikhailov, N., Pinardi, N., and Zavatarelli, M.: The Western Mediterranean Deep Water: A proxy for climate change, *Geophysical Research Letters*, 32, 1–4, doi:10.1029/2005GL022702, 2005.
- Rodgers, K. B., Cane, M. A., and Schrag, D. P.: Seasonal variability of sea surface Delta C-14 in the equatorial Pacific in an ocean circulation
- 15 mode, *J. Geophys. Res.*, 102, 18 627– 18 639., 1997.
- Roether, W. and Weiss, W.: Die groß-skalige thermohaline Zirkulation des Mittelmeers, *Annal. Meteorol. (Neue Folge)*, 15, 1980.
- Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: Recent Changes in Eastern Mediterranean Deep Waters, *Science*, 271, 333–335, doi:10.1126/science.271.5247.333, 1996.
- Roether, W., Klein, B., Manca, B. B., Theocharis, A., and Kioroglou, S.: Transient Eastern Mediterranean deep waters in response to the
- 20 massive dense-water output of the Aegean Sea in the 1990s, *Progress in Oceanography*, 74, 540–571, doi:10.1016/j.pocean.2007.03.001, 2007.
- Sarmiento, J. L. and Gruber, N.: *Ocean Biogeochemical Dynamics*, 2006.
- Schroeder, K., Ribotti, a., Borghini, M., Sorgente, R., Perilli, a., and Gasparini, G. P.: An extensive western Mediterranean deep water renewal between 2004 and 2006, *Geophysical Research Letters*, 35, 1–7, doi:10.1029/2008GL035146, 2008.
- 25 Schroeder, K., Chiggiato, J., Bryden, H. L., Borghini, M., and Ben Ismail, S.: Abrupt climate shift in the Western Mediterranean Sea, *Scientific Reports*, 6, 23 009, doi:10.1038/srep23009, <http://www.nature.com/articles/srep23009>, 2016.
- Siani, G., Paterne, M., Arnold, M., Bard, E., Mativier, B., Tisnerat, N., and Bassinot, F.: Radiocarbon Reservoir Ages In The Mediterranean Sea And Black Sea, *Radiocarbon*, 42, 271–280, doi:http://dx.doi.org/10.2458/azu_js_rc.42.3844, 2000.
- Soto-Navarro, J., Somot, S., Sevault, F., Beuvier, J., Béranger, K., Criado-Aldeanueva, F., and García-Lafuente, J.: Evaluation of regional
- 30 ocean circulation models for the Mediterranean Sea at the Strait of Gibraltar : volume transport and thermohaline properties of the outflow, *Climate Dynamics*, doi:10.1007/s00382-014-2179-4, 2014.
- Stanev, E. V. and Peneva, E. L.: Regional sea level response to global climatic change : Black Sea examples, *Europe*, 32, 33 – 47, 2002.
- Stuiver, M. and Polach, H.: Discussion: Reporting of 14C data, *Radiocarbon*, 22, 964–966, 1977.
- Stuiver, M., Quay, P. D., and Ostlund, H. G.: Abyssal water carbon-14 distribution and the age of the world oceans., *Science (New York, N.Y.)*, 219, 849–51, doi:10.1126/science.219.4586.849, <http://www.ncbi.nlm.nih.gov/pubmed/17780221>, 1983.
- 35 SUESS, H. E.: Radiocarbon Concentration in Modern Wood, *Science*, 122, 1955.

- Sweeney, M. T., Thomson, M. J., Cho, Y. G., Park, Y. J., Williamson, S. H., Bustamante, C. D., and McCouch, S. R.: Global Dissemination of a Single Mutation Conferring White Pericarp in Rice, *PLoS Genetics*, 3, e133, doi:10.1371/journal.pgen.0030133, <http://dx.plos.org/10.1371/journal.pgen.0030133>, 2007.
- Tanhua, T., Hainbucher, D., Cardin, V., Álvarez, M., Civitarese, G., McNichol, A. P., and Key, R. M.: Repeat hydrography in the Mediterranean Sea, data from the Meteor cruise 84/3 in 2011, *Earth System Science Data*, 5, 289–294, doi:10.5194/essd-5-289-2013, <http://www.earth-syst-sci-data.net/5/289/2013/>, 2013.
- Theocharis, A., Georgopoulos, D., Karagevrekis, P., Iona, A., Perivoliotis, L., and Charalambidis, N.: Aegean influence in the deep layers of the Eastern Ionian Sea, *Rapport de la Commission internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 33, 235, 1992.
- Theocharis, A., Nittis, K., Kontoyiannis, H., Papageorgiou, E., and Balopoulos, E.: Climatic changes in the Aegean Sea influence the Eastern Mediterranean thermohaline circulation (1986–1997), *Geophysical Research Letters*, 26 (11), 1617–1620, 1999.
- 10 Tisnérat-Laborde, N., Montagna, P., McCulloch, M., Siani, G., Silenzi, S., and Frank, N.: A High-Resolution Coral-Based $\Delta 14C$ Record of Surface Water Processes in the Western Mediterranean Sea, *Radiocarbon*, 55, 1617–1630, doi:10.2458/azu_js_rc.55.16397, <https://journals.uair.arizona.edu/index.php/radiocarbon/article/view/16397>, 2013.
- Toggweiler, J. R., Dixon, K., and Bryan, K.: Simulations of radiocarbon in a coarse-resolution world ocean model: 1. Steady state prebomb distributions, *Journal of Geophysical Research*, 94, 8217, doi:10.1029/JC094iC06p08217, <http://doi.wiley.com/10.1029/JC094iC06p08217>, 1989a.
- 15 Toggweiler, J. R., Dixon, K., and Bryan, K.: Simulations of radiocarbon in a coarse-resolution world ocean model: 2. Distributions of bomb-produced carbon 14, *Journal of Geophysical Research*, 94, 8243, doi:10.1029/JC094iC06p08243, <http://doi.wiley.com/10.1029/JC094iC06p08243>, 1989b.
- Vörösmarty, C. J., Fekete, B. M., and Tucker, B. A.: Global River Discharge Database (RivDIS V1.0), International Hydrological Program, Global Hydrological Archive and Analysis Systems, UNESCO, Paris, 1996.
- 20 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *Journal of Geophysical Research*, 97, 7373, doi:10.1029/92JC00188, 1992.

Table 1. Regional means of radiocarbon reservoir age before 1950 AD. Column 2 gives the observations from Reimer and McCormac (2002); Column 3 the model values in 1940 AD. The uncertainty in the mean is the larger of the standard deviation based on counting statistics and the “standard deviation,” which is the square root of the variance.

Region	Age (Yr)	
	Reimer and McCormac (2002)	Model
Western Mediterranean	400 ± 22	402 ± 27
Eastern Mediterranean	353 ± 47	349 ± 14
Algerian sub-basin	413 ± 51	410 ± 27
Tyrrhenian sub-basin	390 ± 21	373 ± 29
Adriatic sub-basin	396 ± 61	349 ± 32
Aegean sub-basin	480 ± 72	336 ± 14
Whole Mediterranean	400 ± 16	379 ± 19




Figure 1. Atmospheric $\Delta^{14}\text{C}$ in ‰ (orange), atmospheric CO_2 in ppm (blue), from Orr et al. (2016) and references cited therein.




Figure 2. Model output for March 1956 showing the pre-bomb situation. Upper panel: mean $\Delta^{14}C$ (in ‰) in surface waters (0 to 200 m), intermediate waters (200 to 600 m), and deep waters (600 to 3500 m). Lower panel: $\Delta^{14}C$ along E-W section in (d) WMed, and (e) EMed, where colour-filled dots represent in-situ observations (Broecker and Gerard, 1969). Both model and data are reported with the same colour scale.




Figure 3. Average radiocarbon age (years) in the upper 50 m as computed with the model for 1940. Circles represent reservoir ages derived from measurements of the composition of shells (Siani et al., 2000; Reimer and McCormac, 2002), and from corals (Tisnérat-Laborde et al., 2013).




Figure 4. Model/data comparison of $\Delta^{14}C$ vertical profiles for (a) the pre-bomb distribution as a composite of seawater observations from different locations measured by Broecker and Gerard (1969), and (b) total radiocarbon distributions (natural + bomb) in the Eastern basin measured by Stuiver et al. (1983) at 18 °E. Model results are in blue, while black indicates the in-situ data.




Figure 5. Model output for March 1977 for the post-bomb situation. Upper panel: mean $\Delta^{14}C$ (in ‰) in surface waters (0 to 200 m), intermediate waters (200 to 600 m), and deep waters (600 to 3500 m). Lower panel: $\Delta^{14}C$ along E-W section in (d) WMed, and (e) EMed, where colour-filled dots represent in-situ observations (Stuiver et al., 1983). Both model and data are reported with the same colour scale.




Figure 6. $\Delta^{14}C$ values (in ‰) in the Ligurian sub-basin from 1765 to 2008 for the surface water (0-10 m depth; blue dashed line), and sub-surface water (0-100 m depth; blue solid line) together with available in-situ observations (Tisnérat-Laborde et al., 2013) from coral (black dashed line) and molluscs (cyan squares).




Figure 7. Model output in March 2011. Upper panel: mean $\Delta^{14}C$ (in ‰) surface water (0 to 200 m), intermediate water (200 to 600 m), and deep water (600 to 3500 m). Lower panel: $\Delta^{14}C$ along E-W section in (d) WMed, and (e) EMed, where colour-filled dots represent in-situ observations from Meteor M84 Tanhua et al. (2013). Both model and data are reported with the same colour scale.




Figure 8. Comparison of average vertical profiles of $\Delta^{14}C$ in the WMed (left) and in the EMed (right). Model results are in blue; red indicates the in-situ data




Figure 9. $\Delta^{14}\text{C}$ evolution from 1925 to 2008, in the Gulf of Lions (green), the Algerian sub-basin (red), the Levantine sub-basin (magenta), the Tyrrhenian sub-basin (black), the Cretan Sea (cyan) and the Ligurian (blue).




Figure 10. $\Delta^{14}C$ difference between EMT and NoEMT experiments along sections in the WMed (left column) and in the EMed (right column) for 1995 (top) and 1999 (bottom).




Figure 11. Mean $\Delta^{14}\text{C}$ obtained for experiment with EMT (red) and NoEMT (green), (a) in the Levantine sub-basin deep water (2000-3500 m depth) and (b) in the Algerian sub-basin at intermediate level (200-600 m). The right panels illustrate the difference between EMT and NoEMT for the corresponding left panels.