



The effects of environment on *Arctica islandica* shell formation and architecture

Stefania Milano^{1*}, Gernot Nehrke², Alan D. Wanamaker Jr.³, Irene Ballesta-Artero^{4,5}, Thomas Brey², Bernd R. Schöne¹

¹ Institute of Geosciences, University of Mainz, Joh.-J.-Becherweg 21, 55128 Mainz, Germany

² Alfred Wegener Institute for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

³ Department of Geological and Atmospheric Sciences, Iowa State University, Ames, Iowa, 50011-3212, USA

⁴ Royal Netherlands Institute for Sea Research and Utrecht University, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

⁵ Department of Animal Ecology, VU University Amsterdam, The Netherlands

* Corresponding author. Email: smilano@uni-mainz.de

Keywords: Confocal Raman microscopy; Scanning electron microscopy; Shell microstructure; Water temperature; Diet; Bivalve shell



25 Abstract

26 Mollusks record valuable information in their hard parts that reflect ambient environmental
27 conditions. For this reason, shells can serve as excellent archives to reconstruct past climate and
28 environmental variability. However, animal physiology and biomineralization, which are often
29 poorly understood, can make the decoding of environmental signals a challenging task. Many of
30 the routinely used shell-based proxies are sensitive to multiple different environmental and
31 physiological variables. Therefore, the identification and interpretation of individual environmental
32 signals (e.g. water temperature) often is particularly difficult. Additional proxies not influenced by
33 multiple environmental variables or animal physiology would be a great asset in the field of
34 paleoclimatology. The aim of this study is to investigate the potential use of structural properties
35 of *Arctica islandica* shells as an environmental proxy. A total of eleven specimens were analyzed
36 to study if changes of the microstructural organization of this marine bivalve are related to
37 environmental conditions. In order to limit the interference of multiple parameters, the samples
38 were cultured under controlled conditions. Three shells presented here were grown at two different
39 water temperatures (10 °C and 15 °C) for multiple weeks and exposed only to ambient food
40 conditions. An additional eight specimens were reared under three different dietary regimes. Shell
41 material was analyzed with two techniques: (1) Confocal Raman microscopy (CRM) was used to
42 quantify changes of the orientation of microstructural units and pigment distribution and (2)
43 Scanning electron microscopy (SEM) was used to detect changes in microstructural organization.
44 Our results indicate that *A. islandica* microstructure is not sensitive to changes in the food source,
45 and likely, shell pigment are not altered by diet. However, seawater temperature had a statistically
46 significant effect on the orientation of the biomineral. Although additional work is required, the



47 results presented here suggest that the crystallographic orientation of biomineral units of *A.*
48 *islandica* may serve as an alternative and independent proxy for seawater temperature.

49

50

51 1. Introduction

52 Biomineralization is a process through which living organisms produce a protective, mineralized
53 hard tissue. The considerable diversity of biomineralizing species contributes to high variability in
54 terms of shape, organization and mineralogy of the structures produced (Lowenstam and Weiner,
55 1989; Carter et al., 2012). Different architectures at the micrometer and nanometer scale and
56 different biochemical compositions determine material properties that serve specific functions
57 (Weiner and Addadi, 1997; Currey, 1999; Merkel et al., 2007). Besides these differences, all
58 mineralized tissues are hybrid materials consisting in hierarchical arrangements of biomineral units
59 surrounded by organic matrix, also known as “microstructures” (Bøggild, 1930; Carter and Clark,
60 1985; Rodriguez-Navarro et al., 2006) or “ultrastructures” (Blackwell et al., 1977; Olson et al.,
61 2012) or overall “fabrics” (Schöne, 2013; Schöne et al., 2013). The carbonate and organic phases
62 represent the fundamental level of the organization of biomaterials (Aizenberg et al., 2005; Meyers
63 et al., 2006). The mechanisms of microstructure formation and shaping, especially in mollusks, has
64 attracted increasing attention during recent decades. At present, it is commonly accepted that the
65 organic compounds play an important role in the formation of the inorganic phases of biominerals
66 (Weiner and Addadi, 1991; Berman et al., 1993; Dauphin et al., 2003; Nudelman et al., 2006).
67 However, the identification of the exact mechanisms driving biomineralization is still an open
68 research question. Previous studies conducted on mollusks show that environmental parameters



69 can influence microstructure formation (Lutz, 1984; Tan Tiu and Prezant, 1987; Tan Tiu, 1988;
70 Nishida et al., 2012). These results set the stage for a research interest toward the use of shell
71 microstructures as proxies for reconstructing environmental conditions (Tan Tiu, 1988; Tan Tiu
72 and Prezant, 1989; Olson et al., 2012; Milano et al., 2015).

73 Mollusks are routinely used as climate and environmental proxy archives because they can
74 record a large amount of environmental information in their shells (Richardson, 2001; Wanamaker
75 et al., 2011a; Schöne and Gillikin, 2013). Whereas structures at nanometric level are still
76 underexplored as potential environmental recorders, shell patterns at lower magnification, such as
77 individual growth increments, are commonly used for this purpose (Jones, 1983; Schöne et al.,
78 2005; Marali and Schöne, 2015; Mette et al., 2016). Mollusks deposit skeletal material on a
79 periodic basis and at different rates (Thompson et al., 1980; Deith, 1985). During periods of fast
80 growth, growth increments are formed whereas during periods of slower growth, growth lines are
81 formed (Schöne, 2008; Schöne and Gillikin, 2013). The periodicity of such structures ranges from
82 tidal to annual (Gordon and Carriker, 1978; Schöne and Surge, 2012). By crossdating time-series
83 with similar growth patterns it is possible to construct century and millennia-long master
84 chronologies (Marchitto et al., 2000; Black et al. 2008; Black et al., 2016; Butler et al., 2013). This
85 basic approach, in combination with geochemical methods, has great potential in reconstructing
86 past climatic conditions (Wanamaker et al., 2011b). At present, the most frequently used and well-
87 accepted geochemical proxy is oxygen isotopic composition of shell material ($\delta^{18}\text{O}_{\text{shell}}$) (Epstein,
88 1953; Grossman and Ku, 1986; Schöne et al., 2004; Wanamaker et al., 2007) which may serve as
89 a paleothermometer and/or paleosalinometer (Mook, 1971; Andrus, 2011); however, the $\delta^{18}\text{O}_{\text{shell}}$
90 is influenced by both seawater temperature and the isotopic composition of seawater ($\delta^{18}\text{O}_{\text{water}}$;
91 related to salinity). Thus, $\delta^{18}\text{O}_{\text{shell}}$ -based temperature reconstructions are particularly challenging
92 in habitats with fluctuating $\delta^{18}\text{O}_{\text{water}}$ conditions such as estuaries or restricted basins (Gillikin et al.,



2005). Because of the multiple impacts on $\delta^{18}\text{O}_{\text{shell}}$ values, there have been substantial efforts (including elemental ratios; cite Schone et al 2010 papers here) to develop alternative techniques to reconstruct environmental variables from mollusk shells.

This study investigates the possibility using shell microstructure properties to serve as a new environmental proxy. For this purpose, the effects of seawater temperature (grown at 10 °C and 15 °C) and dietary regime on the microstructural units of *Arctica islandica* cultured under controlled conditions were analyzed and quantified. *A. islandica* was chosen as model species because of its great potential in paleoclimatology and paleoceanography (see Schöne, 2013; Wanamaker et al., 2016). Its extreme longevity of up to more than 500 years makes this species a highly suitable archive for long-term paleoclimate and environmental reconstructions (Schöne et al., 2005; Wanamaker et al., 2008; Wanamaker et al., 2012; Butler et al., 2013).

2. Materials and Methods

The analyses were conducted on eleven *A. islandica* shells. Three juvenile *A. islandica* shells, sampled for the seawater temperature experiment, were collected alive on November 21, 2009 aboard the *F.V. Three of a Kind* off Jonesport, Maine USA (44° 26' 9.829"N, 67° 26' 18.045"W) in 82 m water depth. From 2009 to 2011, all animals were kept in a flowing seawater laboratory at the Darling Marine Center, Walpole, Maine, USA (see Beirne et al., 2012 for additional details). In 2011, clams were grown at two different temperature regimes for 16 weeks (Table 1). At the completion of the experiment, shells were estimated to be between 4 to 5 years old. Eight one-year



old juveniles were collected in July 2014 from Kiel Bay, Baltic Sea (54° 32' N, 10° 42' E; Fig. 1) and kept alive in tanks at 7 °C for six months at the Alfred Wegener Institute for Polar and Marine Research (AWI), Bremerhaven, Germany. During this time interval, the animals were fed with an algal mix of *Nannochloropsis* sp., *Isochrysis galbana* and *Pavlova lutheri*. Then, they were transferred to the Royal Netherlands Institute for Sea Research (NIOZ), Texel, The Netherlands, and cultured in tanks at three different dietary conditions for 11 weeks (Table 1).

121

122

2.1 Seawater temperature experiment

The seawater temperature experiment started on 27 March 2011 and ended on 21 July 2011. Prior to the start of the experiment the animals were marked with calcein. The staining leaves a clear fluorescent marker in the shells that can be used to identify which shell material has formed prior to and during the experiment. Initially, the animals were kept at 10.3 ± 0.2 °C for 48 days. Then, they were briefly removed from the tanks and marked again. Subsequently, the clams were cultured for 69 more days at 15.0 ± 0.3 °C. Ambient seawater was pumped from the adjacent Damariscotta River estuary and adjusted to desired temperature. The salinity was measured with a Hydrolab[®] MiniSonde. It ranged between 30.2 ± 0.7 and 30.7 ± 0.7 , in the two experimental phases, respectively. During the entire culture period, all clams were exposed to ambient food conditions. At the end of the experiment the soft tissues were removed.

134

135

2.2 Food experiment



137 The food experiment was carried out from 9 February 2015 to 29 April 2015. The animals were
138 placed in aquaria inside a climate room at 9 °C. Water temperature in the tanks ranged between 8
139 and 10 °C. Water salinity was measured by using an ENDECO 102 refractometer and ranged
140 between 29.6 and 29.9 ± 0.1 in each aquarium. The 15-liter tanks were constantly supplied with
141 aerated water from the Wadden Sea. The clams were acclimated for three weeks before the start of
142 the experiment. Three dietary regimes were chosen. One treatment consisted of feeding the animals
143 with Microalgae Mix (food type 1), a ready-made solution of four marine microalgae (25 %
144 *Isochrysis*, 25 % *Tetraselmis*, 25 % *Thalassiosira*, 25 % *Nannochloropsis*) with a particle size
145 range of 2 - 30 µm. A second treatment was based on PhytoMaxx (food type 2), a solution of living
146 *Nannochloropsis* algae with 2 - 5 µm size range. A third treatment served as control, i.e., the
147 animals were not fed with any additional food. In treatments with food type 1 and 2, microalgae
148 were provided at the constant optimum concentration of 20×10^6 cells/liter (Winter, 1969). A
149 dispenser equipped with a timer was used to distribute the food five times per day. At the end of
150 the experiment the soft tissues were removed. A distinct dark line in the shells indicated the
151 transposition to the NIOZ aquaria and the associated stress. This line marks the beginning of the
152 tank experiment.

153

154

155 2.3 Sample preparation

156 The right valve of each specimen was cut into two 1.5 millimeter-thick sections along the axis of
157 maximum growth. For this purpose, a low speed precision saw (Buehler Isomet 1000) was used.
158 Given the small size and fragility of the juvenile shells used in the food experiment, the valves
159 were fully embedded in a block of Struers EpoFix (epoxy) and air-dried overnight prior the



sectioning. Sections of the clams used in the temperature experiment were embedded in epoxy after the cutting. All samples were ground using a Buehler Metaserv 2000 machine equipped with Buehler silicon carbide papers of different grit sizes (P320, P600, P1200, P2500). In addition, the samples were manually ground with Buehler P4000 grit paper and polished with a Buehler diamond polycrystalline suspension (3 μ m). Sample surfaces were rinsed in demineralized water and air-dried. In the samples of the temperature experiment, the calcein marks were located under a fluorescence light microscope (Zeiss Axio Imager.A1m microscope equipped with a Zeiss HBO100 mercury lamp and filter set 38: excitation wavelength, ca. 450 - 500 nm; emission wavelength, ca. 500 - 550 nm).

169

170

2.4 *A. islandica* shell organization

The shell of *A. islandica* consists of pure aragonite and it is divided in two major layers, an outer (OSL) and the inner shell layer (ISL). The OSL is further subdivided in outer (oOSL) and inner portion (iOSL) (Schöne, 2013). These layers are characterized by specific microstructures (Ropes et al., 1984). The oOSL largely consists of homogenous microstructure with a bulky and granular aspect (Schöne et al., 2013). The iOSL and ISL are largely composed of crossed-lamellar to cross-acicular microstructures (Dunca et al., 2009). The present study focuses on ventral margin of the shells. Analyses were carried out exclusively in the OSL.

Similar to other mollusks, the shell of *A. islandica* contains pigment polyenes which are obviously visible when using CRM (Hedegaard et al., 2006). Polyenes are organic compounds containing single (C-C) and double (C=C) carbon-carbon bonds forming a polyenic chain. Their



182 distribution across the shell is not homogenous. The pigments are abundant in the oOSL whereas
183 they become scarce in the iOSL (Stemmer and Nehrke, 2014).

184

185

186 2.5 Confocal Raman microscopy and image processing

187 Shells were mapped with a WITec alpha 300 R (WITec GmbH, Germany) confocal Raman
188 microscope. Scans of $50 \times 50 \mu\text{m}$, $100 \times 50 \mu\text{m}$ and $150 \times 50 \mu\text{m}$ were performed using a
189 piezoelectric scanner table. All Raman measurements were carried out using a 488 nm diode laser.
190 A spectrometer (UHTS 300, WITec, Germany) was used with a 600 mm^{-1} grating, a 500 nm blaze
191 and an integration time of 0.03 s. On each sample two to six scans were made, depending of the
192 thickness of the shell. For instance, in juvenile shells (food experiment), two scans of each sample
193 were made. On larger shells used in the temperature experiment, six maps were completed, i.e.,
194 two maps in the oOSL, two in the middle of the iOSL and two in the inner portion of the iOSL.
195 Each scan contained between 40,000 and 120,000 data points, depending on the map size. The
196 spatial resolution equaled 250 nm. Half of the maps were performed on the shell portion formed
197 before the experiments. The other half were made on the shell portion formed under experimental
198 conditions. In order to avoid areas affected by transplantation or marking stress, the scans were
199 located far off the calcein and stress lines. Raman maps on food experiment shells were performed
200 $300 \mu\text{m}$ away from the stress line. In the shells from the temperature experiment, the scans were
201 made 1 mm away from the calcein mark.

202 Polarized Raman microscopy is known to provide comprehensive information about the
203 crystallographic properties of the materials (Hopkins and Farrow, 1985). The aragonite spectrum



is characterized by two lattice modes (translation mode T_a , 152cm^{-1} and librational mode L_a , 206cm^{-1}) and the two internal modes (in-plane band ν_4 , 705cm^{-1} and symmetric stretch ν_1 , 1085cm^{-1}). The ratio (R_{ν_1/T_a}) between peak intensities belonging to ν_1 and T_a is caused by different crystallographic orientations of the aragonitic units (Hopkins and Farrow, 1985; Nehrke and Nouet, 2011). Within each scan, R_{ν_1/T_a} was calculated for each data point. New spectral images were generated using WITecProject software (version 4.1, WITec GmbH, Germany). These images were then binarized by replacing all values above 2.5 with 1 and the others with 0. The orientation was quantified by calculating the area formed by pixels of value 1 over the total scan area. The imaging software Gwyddion (<http://gwyddion.net/> last checked: June 2016) was used for this purpose. The results were expressed in percentage.

The Raman scans of the food experiment shells were analyzed to investigate the pigment composition. Polyene peaks have definite positions in the spectrum according to the number of the C-C and C=C bonds of the chain, which are specific for certain types of pigments. The two major polyene peaks at ~ 1130 (R_1) and 1520 cm^{-1} (R_4) were identified by using the “multipeak fitting 2” routine of IGOR Pro (version 7.00, WaveMetrics, USA). Their exact position was determined adopting a Gaussian fitting function (Fig. 2). The number of single (N_1) and double carbon bonds (N_4) was calculated by applying the equations by Schaffer et al. (1991):

$$(1) \quad N_1 = 476 (R_1 - 1,082)$$

$$(2) \quad N_4 = 830 (R_4 - 1,438)$$

Spectral images of the R_4 band were used to locate the polyenes in the shell and measure the thickness of the pigmented layer. This analysis was conducted only on the shells of the food experiment. Given the larger size of the shells used in the temperature experiment, the spectral



226 maps were not sufficient for a correct localization of the pigmented layer boundaries and estimation
227 of its thickness.

228 To quantify changes of the orientation of individual biomineral units of the juvenile shells
229 (food experiment), the spectral maps were subdivided into two portions. The outermost shell
230 portion (oOSL) was enriched in pigments whereas the iOSL showed a decrease in polyene content.

231

232

233 2.6 Scanning electron microscopy

234 After performing Raman measurements, the samples were prepared for SEM analysis. Each shell
235 slab was ground with a Buehler Metaserv 2000 machine and Buehler silicon F2500 grit carbide
236 paper. To reduce the impact of grinding on the sample surface of juvenile shells, extra grinding
237 was done by hand. Then, the slabs were polished with a Buehler diamond polycrystalline
238 suspension (3 μm). Afterward, shell surfaces were etched in 0.12 N HCl solution for 10 (food
239 experiment samples) to 90 s (temperature experiment samples) and subsequently placed in 6 vol %
240 NaClO solution for 30 min. After being rinsed in demineralized water, air-dried samples were
241 sputter-coated with a 2 nm-thick platinum film by using a Low Vacuum Coater Leica EM ACE200.

242 A scanning electron microscope (LOT Quantum Design 2nd generation Phenom Pro desktop
243 SEM) with backscattered electron detector and 10 kV accelerating voltage was used to analyze the
244 shells. Images were taken at the same distances from the calcein and stress lines as in the case of
245 the Raman measurements to assure comparability of the data.

246 In addition, stitched SEM images of the ventral margins were used to accurately determine
247 the shell growth advance during the culturing experiments. Growth increment widths were



measured with the software Panopea (© Schöne and Peinl). Given the difference in duration of the two phases of the temperature experiment, the measurements were expressed as total growth and instantaneous growth rate (Fig. 3a + b). The latter was calculated using the following equation (Brey et al., 1990; Witbaard et al., 1997):

$$(3) \text{ Instantaneous growth rate} = (\ln(y_t / y_0) / a)$$

where y_0 represents the initial shell height, y_t is the final shell height and a is the duration of the experiment. In the case of the food experiment, only the total growth was calculated (Fig. 3c).

3. Results

3.1 Effect of seawater temperature and diet on *A. islandica* shell growth

When exposed to a water temperature of 10 °C, the shells grew between 11.67 and 14.17 mm during a period of 48 days. During a period of 69 days at 15 °C, the growth ranged between 2.32 and 5.77 mm (Fig. 3a). Instantaneous growth rate showed a decrease between the two experimental phases. At 10 and 15 °C, the average instantaneous growth per day was 0.0091 and 0.0013, respectively (Fig. 3b). The decrease in total growth and growth rate between the two temperatures was statistically significant (t -test, $p < 0.01$).

During the food experiment, shells grew between 0.37 and 3.71 mm with large differences due to the different food types. Growth of specimens exposed to food type 1 ranged between 1.87 to 3.71 mm, whereas those cultured with food type 2 grew between 0.55 to 0.96 mm. Both control



specimens added 0.37 mm of shell during the experimental phase (Fig. 3c). ANOVA and Tukey's HSD post hoc tests showed significant differences between specimens cultured with food type 1 and 2 ($p < 0.05$) and between food type 1 and control shells ($p < 0.05$).

3.2 Effect of seawater temperature on *A. islandica* microstructure

At a water temperature of 10 °C, the area occupied by microstructural units oriented with $R_{v1/Ta}$ higher than 2.5 a.u. (= arbitrary units) ranged between 31.3 and 50.6 % in the oOSL and between 21.3 and 33.5 % in the iOSL. When exposed to 15 °C, values ranged between 25.6 and 48.7 % and between 45.7 and 55.9 % in the oOSL and iOSL, respectively (Fig. 4). Whereas the slight difference of area with $R_{v1/Ta} > 2.5$ in the oOSL was not significant between the two water temperatures (t -test, $p = 0.62$), the area with $R_{v1/Ta} > 2.5$ in the iOSL significantly increased at 15 °C (t -test, $p = 0.02$). Under the SEM, no difference was visible between units formed at 10 °C and 15 °C (Fig. 5).

3.3 Effect of food on *A. islandica* microstructure and pigments

In the shells cultured with food type 1, the area occupied by biomineral units oriented with $R_{v1/Ta}$ higher than 2.5 a.u. during the experiment ranged between 24.8 % (oOSL) and 43.0 % (iOSL). In the shell portion deposited before the experiment, the ratio varied between 19.4 % (oOSL) and 36.2 % (iOSL). Although a trend was recognized, these variations were not statistically different (t -tests. OSL: $p = 0.43$; ISL: $p = 0.57$; Fig. 6a). On the contrary, in the clams exposed to food type 2, the



290 area occupied by units oriented with $R_{v1/Ta} > 2.5$ ranged between 11.7 % (oOSL) and 20.4 %
 291 (iOSL). Before the experiment, the proportions were higher, i.e., 18.1% (oOSL) and 26.3% (iOSL)
 292 (Fig. 6b). As for the other treatment, the difference was not significant (*t*-tests. oOSL: $p = 0.34$;
 293 iOSL: $p = 0.28$). In the control shells grown with no extra food supply, the area with $R_{v1/Ta} > 2.5$
 294 ranged between 24.6 % (oOSL) and 44.8 % (iOSL) during the experiment and 21.2 % (oOSL) and
 295 44.5 % (iOSL) before the experiment (Fig. 6c). Hence, no trend was visible and the two portions
 296 did not show significant differences (*t*-tests. oOSL: $p = 0.59$; iOSL: $p = 0.99$). As for the
 297 temperature experiment, under the SEM, the microstructure of the shells from the food experiment
 298 did not show any change (Fig. 7).

299 All treatments showed a slightly thicker pigmented layer formed during the experiment than
 300 during the acclimation phase (Fig. 8a). During the experiment, clams cultured with food type 1
 301 showed, on average, a thickening by 6.4 %. In the food type 2 specimens, the layer thickness
 302 increased by 9.9 %. Control shells showed an increase of 10.4 % (Fig. 8b). However, none of these
 303 differences was statistically significant (*t*-test. Food type 1: $p = 0.43$; Food type 2: $p = 0.39$; Control:
 304 $p = 0.10$). According to the position of the polyene peaks, the number of single carbon bonds in the
 305 pigment chain did not change between the acclimation and experimental phase ($N_1 = 10.1 \pm 1.3$
 306 and $N_1 = 10.0 \pm 0.9$, respectively). Likely, no significant variation was observed in the number of
 307 double carbon bonds ($N_4 = 10.5 \pm 0.2$ and $N_4 = 10.4 \pm 0.3$, respectively; Table 2).

308

309

310

311



312 4. Discussion

313 According to the results, variations of both food type and water temperature can influence the shell
314 production rate of *A. islandica*. However, the shell microstructure and pigmentation react
315 differently to these two environmental variables. Whereas changes of the dietary conditions do not
316 affect the shell architecture and pigment composition, the crystallographic orientation of the
317 biomineral units responds to seawater temperature fluctuations.

318

319

320 4.1 Environmental influence on shell microstructure

321 The environmental conditions experienced by mollusks during the process of biomineralization
322 appear to influence shell organization (Carter, 1980). Among the different environmental variables,
323 water temperature is the most studied driving force of structural changes of the shell. For instance,
324 shell mineralogy can vary depending on water temperature (Carter, 1980). According to the thermal
325 potentiation hypothesis, nucleation and growth of calcitic structural units is favored at low
326 temperatures by kinetic factors (Carter et al., 1998). As a consequence, bivalve species living in
327 cold water environments exhibit additional or thicker calcitic layers compared to the corresponding
328 species from warm waters (Lowenstam, 1954; Taylor and Kennedy, 1969). Changes in the calcium
329 carbonate polymorph also affect the type of microstructures (Milano et al., 2016). However,
330 architectural variations often occur without mineralogical impact (Carter, 1980).

331 The present results indicate that temperature induces a change in the crystallographic
332 orientation of the biomineral units of *A. islandica*. Although water temperature was previously
333 shown to have an impact on microstructure formation, the attention has been mainly addressed to



the effects on the morphometric characteristics (e.g. size and shape) or on the type of microstructure. Milano et al. (2015) demonstrated that size and elongation of prismatic structural units of *Cerastoderma edule* were positively correlated to seawater temperature variation throughout the growing season. Likely, low temperatures induced the formation of small nacre tablets in *Geukensia demissa* (Lutz, 1984). Seasonal changes of the microstructural type were reported in the freshwater bivalve *Corbicula fluminea* (Prezant and Tan Tiu, 1986; Tan Tiu and Prezant, 1989). During the warm months, crossed acicular structure was produced, whereas simple crossed-lamellae were formed during the winter period. So far, variations of the crystallographic properties of bivalve biominerals have been exclusively investigated as a response to hypercapnic (acidified) conditions. *Mytilus galloprovincialis* and *Mytilus edulis* showed a significant change in the orientation of the prisms forming shell calcitic layer when subjected to hypercapnia (Hahn et al., 2012; Fitzner et al., 2014). Altered crystallographic organization may derive from the animal exposure to suboptimal conditions. These findings together with the present results suggest that thermal- and hypercapnic-induced stress are likely to affect the ability of the bivalves to preserve the orientation of their microstructural units (Fitzner et al., 2015).

Different food sources do not significantly influence the orientation of the biomineral units or the composition and distribution of pigments in shells of *A. islandica*. In previous studies, the relationship between microstructure and diet was virtually overlooked resulting in a lack of data in the literature. As suggested by Hedegaard et al., (2006), however, the type of polyenes is influenced by food. The ingestion of pigment-enriched microalgae potentially leads to an accumulation of pigments in mollusk tissues and the shell (Soldatov et al., 2013). On the other hand, it has been argued that polyenes do not generate from food sources like other pigments (i.e., carotenoids), but they are locally synthesized (Karampelas et al., 2009). In accordance to Stemmer and Nehrke (2014), the results presented here support the view that the specific diets on which the animals rely



on do not influence shell pigment composition. The chemical characteristics of the polyenes are likely to be specie-specific and independent from the habitats.

4.2 Confocal Raman microscopy as tool for microstructural analysis

From a methodological perspective, the present study represents an innovative approach in the investigation of shell microstructural organization. Electron backscatter diffraction (EBSD) has been previously used to determine the crystallographic orientation of gastropod (Fryda et al., 2009; Pérez-Huerta et al., 2011) and bivalve microstructural units (Checa et al., 2006; Frenzel et al., 2012; Karney et al., 2012). Whereas, CRM on mollusk shells is generally applied within studies on taphonomic mineralogical alteration and pigment identification (Stemmer and Nehrke, 2014; Beierlein et al., 2015). Both techniques provide considerably high spatially resolved analysis up to 250 nm, allowing the identification of individual structural units at μm - and nm-scale (Cusack et al., 2008; Karney et al., 2012). CRM offers important advantages supporting a broader application of this methodology in the biomineralization research field. For instance, samples do not require any pre-treatment. Unlike EBSD, there is no need of preparing thin-sections ($\sim 150 \mu\text{m}$ thick) or etching the shell surface (Griesshaber et al., 2010; Hahn et al., 2012). Therefore, further structural and geochemical analyses can be easily performed on the same sections (Nehrke et al., 2012). In addition, the size of CRM scans can be remarkably large ($\sim 7\text{-}8 \text{ mm}^2$) without compromising the achievable resolution. By overlapping adjacent scans, it is possible to produce stitched scans allowing to further increase the region of interest on the shell surface.



SEM has previously been demonstrated to provide a convenient approach for the identification of individual structural units and the quantification of potential changes occurring within them (Milano et al., 2015, 2016b). However, SEM exclusively provides information about the morphometric characteristics of the microstructural units. As highlighted by the present study, to achieve an exhaustive examination, it is suggested to combine SEM with techniques assessing crystallographic properties of the biomaterials. For instance, our results show that the effect of water temperature is detectable in crystallographic orientation but not in morphometric features of the biomineral units.

4.3 Environmental influence on shell growth

Numerous previous studies demonstrated that growth rate of *A. islandica* - and many other bivalves - is linked to environmental variables (e.g., Witbaard et al., 1997, 1999; Schöne et al., 2004; Butler et al., 2010; Mette et al., 2016). However, the relative importance of the main factors, temperature and food supply/quality driving shell formation are still not well understood. Positive correlations between shell growth and water temperature have been identified (i.e., Schöne et al., 2005; Wanamaker et al., 2009; Marali et al., 2015), but the relationship between shell growth and environment is more complex (Marchitto et al., 2010; Stott et al., 2010; Schöne et al., 2013) and likely dependent on the synergic effect of food availability and water temperature (Butler et al., 2013; Lohmann and Schöne, 2013; Mette et al., 2016). Tank experiments were run in order to precisely identify the role of these two parameters of shell growth of *A. islandica* (Witbaard et al., 1997; Hiebenthal et al., 2012). A tenfold increase in instantaneous growth rate was observed between 1 and 12 °C, with the greatest variation occurring below 6 °C (Witbaard et al., 1997). On the contrary, a temperature increase between 4 and 16 °C was shown to produce a slowdown of



402 shell production (Hiebenthal et al., 2012). Our results are in agreement with the latter study and
403 show a decrease in the instantaneous growth rate between 10 and 15 °C. High temperatures are
404 often associated with an increase of free radical production (Abele et al., 2002). A large amount of
405 energy then has to be allocated to limit oxidative cellular damage (Abele and Puntarulo, 2004).
406 This translates into a higher accumulation of lipofuscin and slower shell production rate
407 (Hiebenthal et al., 2013). The contrasting results of previous studies may be explained by individual
408 differences in the tolerance toward temperature change (Marchitto et al., 2000).

409 Along with water temperature, food availability was also shown to influence *A. islandica*
410 shell growth (Witbaard et al., 1997). At high algal cell densities, the siphon activity increased. This,
411 in turn, was positively correlated to shell growth. Previous experiments used different combinations
412 of algae such as *Isochrysis galbana* and *Dunaliella marina* (Witbaard et al., 1997), or
413 *Nannochloropsis oculata*, *Phaeodactylum tricornutum* and *Chlorella* sp. (Hiebenthal et al., 2012)
414 to grow the clams. However, there are still uncertainties about the composition of the primary food
415 source for this species (Butler et al., 2010). Even though it is challenging to determine the preferred
416 algal species, our results show that the use of a mixture of different algal species results in
417 significantly faster shell growth than the used of just one algal species. In the natural environment,
418 suspension feeders such as *A. islandica* preferentially ingest certain particle sizes (Rubenstein and
419 Koehl, 1977; Jorgensen, 1996; Baker et al., 1998). The exposure to a limited algal size range, as in
420 the case of food type 2, may affect shell growth. Furthermore, multispecific solutions contain a
421 higher variability of biochemical components that better meet the nutritional requirements of the
422 animal (Widdows, 1991). Our results are in good agreement with previous findings. For instance,
423 it has been shown by Strömberg and Cary (1984) that *Mytilus edulis* shell growth increased as a
424 result of a diet based on three different algal species. Furthermore, Epifanio (1979) tested the
425 differences on the growth of *Crassostrea virginica* and *Mercenaria mercenaria* of a mixed diet



426 composed by *Isochrysis galbana* and *Thalassiosira pseudonana* and diets consisting of the single
427 species. Faster growth was measured in the mixed diet treatment, indicating a synergic effect of
428 the relative food composition (Epifanio, 1979). Likely, *Mytilus edulis* grew faster when reared with
429 different types of mixed diets as opposed to monospecific diets (Galley et al., 2010).

430

431

432

433 5. Conclusions

434 *Arctica islandica* shell growth and biomineral orientation vary with changes in sewerage
435 temperature. However, exposure to different food sources affect shell deposition rate but do not
436 influence the organization of the biomineral units. Given the exclusive sensitivity to one
437 environmental variable, the orientation of biomineral units may represent a promising new
438 temperature proxy for paleoenvironmental reconstructions. However, additional studies are needed
439 to further explore the subject. In particular, intra-individual variability influence on the results
440 needs to be assessed. In the present study, a variation in the orientation between individuals was
441 well visible and the risks associated have to be taken in account when considering further
442 application of the possible proxy. Furthermore, the effect of other environmental variables such as
443 salinity needs to be tested.

444 The innovative application of CRM for microstructural orientation and proxy development
445 proved that the technique has large potential in this research direction. More studies are needed to
446 validate its suitability in paleoclimatology experimental works.



447

448

449 Acknowledgements

450 The authors acknowledge the crew of the *F.V. Three of a Kind* for helping with the collection of
451 the animals. Design and execution of the seawater temperature experiment were successfully
452 realized thanks to the support of B. Beal, D. Gillikin, A. Lorrain and the Darling Marine Center
453 scientific team. Funding for this study was kindly provided by the EU within the framework of the
454 Marie Curie International Training Network ARAMACC (604802).

455

456

457 References

458 Abele, D., Heise, K., Pörtner, H. O. and Puntarulo, S.: Temperature-dependence of mitochondrial
459 function and production of reactive oxygen species in the intertidal mud clam *Mya arenaria*,
460 J. Exp. Biol., 205, 1831-1841, doi:10.1016/j.jeb.2002.04.007, 2002.

461 Abele, D. and Puntarulo, S.: Formation of reactive species and induction of antioxidant defence
462 systems in polar and temperate marine invertebrates and fish, Comp. Biochem. Physiol. Part
463 A, 138(4), 405-415, doi:10.1016/j.cbpa.2004.05.013, 2004.

464 Aizenberg, J., Weaver, J. C., Thanawala, M. S., Sundar, V. C., Morse, D. E. and Fratzl, P.: Skeleton
465 of *Euplectella* sp.: structural hierarchy from the nanoscale to the macroscale, Science,
466 309(5732), 275-278, doi:10.1126/science.1112255, 2005.



- 467 Andrus, C. F. T.: Shell midden sclerochronology, *Quat. Sci. Rev.*, 30(21-22), 2892-2905,
 468 doi:10.1016/j.quascirev.2011.07.016, 2011.
- 469 Baker, S. M., Levinton, J. S., Kurdziel, J. P. and Shumway, S. E.: Selective feeding and
 470 biodeposition by zebra mussels and their relation to changes in phytoplankton composition
 471 and seston load, *J. Shellfish Res.*, 17, 1207-1213, 1998.
- 472 Beierlein, L., Nehkre, G. and Brey, T.: Confocal Raman microscopy in sclerochronology: A
 473 powerful tool to visualize environmental information in recent and fossil biogenic archives,
 474 *Geochemistry Geophys. Geosystems*, 16, 325-335, doi:10.1002/2014GC005684.Key, 2015.
- 475 Beirne, E. C., Wanamaker, A. D. and Feindel, S. C.: Experimental validation of environmental
 476 controls on the $\delta^{13}\text{C}$ of *Arctica islandica* (ocean quahog) shell carbonate, *Geochim.*
 477 *Cosmochim. Acta*, 84, 395-409, doi:10.1016/j.gca.2012.01.021, 2012.
- 478 Berman, A., Hanson, J., Leiserowitz, L., Koetzle, T. F., Weiner, S. and Addadi, L.: Biological
 479 control of crystal texture: a widespread strategy for adapting crystal properties to function,
 480 *Science*, 259(5096), 776-779, doi:10.1126/science.259.5096.776, 1993.
- 481 Black, B. A., Gillespie, D. C., MacLellan, S. E. and Hand, C. M.: Establishing highly accurate
 482 production-age data using the tree-ring technique of crossdating: a case study for Pacific
 483 geoduck (*Panopea abrupta*), *Can. J. Fish. Aquat. Sci.*, 65, 2572-2578, doi:10.1139/F08-158,
 484 2008.
- 485 Black, B. A., Griffin, D., van der Sleen, P., Wanamaker, A. D., Speer, J. H., Frank, D. C., Stahle,
 486 D. W., Pederson, N., Copenheaver, C. A., Trouet, V., Griffin, S. and Gillanders, B. M.: The
 487 value of crossdating to retain high-frequency variability, climate signals, and extreme events



- 488 in environmental proxies, Glob. Chang. Biol., 22(7), 2582-2595, doi:10.1111/gcb.13256,
 489 2016.
- 490
- 491 Bøggild, O. B.: The shell structure of the mollusks, in Det Kongelige Danske Videnskabernes
 492 Selskabs Skrifter, Natruvidenskabelig og Mathematisk, pp. 231-326, Afdeling., 1930.
- 493 Brey, T., Arntz, W. E., Pauly, D. and Rumohr, H.: *Arctica (Cyprina) islandica* in Kiel Bay (Western
 494 Baltic): growth, production and ecological significance, J. Exp. Mar. Bio. Ecol., 136(3), 217-
 495 235, doi:10.1016/0022-0981(90)90162-6, 1990.
- 496 Butler, P. G., Richardson, C. A., Scourse, J. D., Wanamaker, A. D. Jr, Shammon, T. M. and
 497 Bennell, J. D.: Marine climate in the Irish Sea: analysis of a 489-year marine master
 498 chronology derived from growth increments in the shell of the clam *Arctica islandica*, Quat.
 499 Sci. Rev., 29(13-14), 1614-1632, doi:10.1016/j.quascirev.2009.07.010, 2010.
- 500 Butler, P. G., Wanamaker, A. D. Jr, Scourse, J. D., Richardson, C. A. and Reynolds, D. J.:
 501 Variability of marine climate on the North Icelandic Shelf in a 1357-year proxy archive based
 502 on growth increments in the bivalve *Arctica islandica*, Palaeogeogr. Palaeoclimatol.
 503 Palaeoecol., 373, 141-151, doi:10.1016/j.palaeo.2012.01.016, 2013.
- 504 Carter, J. G.: Environmental and biological controls of bivalve shell mineralogy and
 505 microstructure, in Skeletal Growth of Aquatic Organisms: Biological Records of
 506 Environmental Change (Topics in Geobiology), edited by D. C. Rhoads and R. A. Lutz, pp.
 507 69-113, Plenum, N. Y, 1980.



- 508 Carter, J. G. and Clark, G. R. I.: Classification and phylogenetic significance of molluscan shell
 509 microstructure, in Mollusks, Notes for a Short Course, edited by T. W. Broadhead, pp. 50-
 510 71., 1985.
- 511 Carter, J. G., Barrera, E. and Tevesz, M. T. S.: Thermal potentiation and mineralogical evolution
 512 in the Bivalvia (Mollusca), J. Paleontol., 72(6), 991-1010, 1998.
- 513 Carter, J. G., Harries, P. J., Malchus, N., Sartori, A. F., Anderson, L. C., Bieler, R., Bogan, A. E.,
 514 Coan, E. V., Cope, J. C. W., Cragg, S. M., Garcia-March, J. R., Hylleberg, J., Kelley, P.,
 515 Kleemann, K., Kriz, J., McRoberts, C., Mikkelsen, P. M., Pojeta, J. J., Temkin, I., Yancey,
 516 T. and Zieritz, A.: Illustrated glossary of the bivalvia, Treatise Online, 1(48), 2012.
- 517 Checa, A. G., Okamoto, T. and Ramírez, J.: Organization pattern of nacre in Pteriidae (Bivalvia:
 518 Mollusca) explained by crystal competition., Proc. Biol. Sci., 273(1592), 1329-37,
 519 doi:10.1098/rspb.2005.3460, 2006.
- 520 Currey, J. D.: The design of mineralised hard tissues for their mechanical functions, J. Exp. Biol.,
 521 202, 3285-3294, 1999.
- 522 Cusack, M., Parkinson, D., Freer, A., Pérez-Huerta, A., Fallick, A. E. and Curry, G. B.: Oxygen
 523 isotope composition in *Modiolus modiolus* aragonite in the context of biological and
 524 crystallographic control, Mineral. Mag., 72(2), 569-577,
 525 doi:10.1180/minmag.2008.072.2.569, 2008.
- 526 Dauphin, Y., Cuif, J. P., Doucet, J., Salomé, M., Susini, J. and Williams, C. T.: In situ chemical
 527 speciation of sulfur in calcitic biominerals and the simple prism concept, J. Struct. Biol.,
 528 142(2), 272-280, doi:10.1016/S1047-8477(03)00054-6, 2003.



- 529 Deith, M. R.: The composition of tidally deposited growth lines in the shell of the edible cockle,
 530 *Cerastoderma edule*, J. Mar. Biol. Assoc. United Kingdom, 65(3), 573-581, 1985.
- 531 Dunca, E., Mutvei, H., Göransson, P., Mörtz, C.-M., Schöne, B. R., Whitehouse, M. J., Elfman,
 532 M. and Baden, S. P.: Using ocean quahog (*Arctica islandica*) shells to reconstruct
 533 palaeoenvironment in Öresund, Kattegat and Skagerrak, Sweden, Int. J. Earth Sci., 98(1), 3-
 534 17, doi:10.1007/s00531-008-0348-6, 2009.
- 535 Epifanio, C. E.: Growth in bivalve molluscs: nutritional effects of two or more species of algae in
 536 diets fed to the American oyster *Crassostrea virginica* (Gmelin) and the hard clam
 537 *Mercenaria mercenaria* (L.), Aquaculture, 18, 1-12, 1979.
- 538 Epstein, S., Buchbaum, R., Lowenstam, H. A. and Urey, H. C.: Revised carbonate-water isotopic
 539 temperature scale. Bull. Geol. Soc. Am., 64, 1315-1326, 1953.
- 540 Fitzner, S. C., Cusack, M., Phoenix, V. R. and Kamenos, N. A.: Ocean acidification reduces the
 541 crystallographic control in juvenile mussel shells, J. Struct. Biol., 188, 39-45,
 542 doi:10.1016/j.jsb.2014.08.007, 2014.
- 543 Fitzner, S. C., Zhu, W., Tanner, K. E., Phoenix, V. R., Nicholas, A. K. and Cusack, M.: Ocean
 544 acidification alters the material properties of *Mytilus edulis* shells, J. R. Soc. Interface,
 545 12(103), doi:10.1098/rsif.2014.1227Published, 2015.
- 546 Frenzel, M., Harrison, R. J. and Harper, E. M.: Nanostructure and crystallography of aberrant
 547 columnar vaterite in *Corbicula fluminea* (Mollusca), J. Struct. Biol., 178(1), 8-18,
 548 doi:10.1016/j.jsb.2012.02.005, 2012.



- 549 Fryda, J., Bandel, K. and Frydova, B.: Crystallographic texture of Late Triassic gastropod nacre:
550 Evidence of long-term stability of the mechanism controlling its formation, Bull. Geosci.,
551 84(4), 745-754, doi:10.3140/bull.geosci.1169, 2009.
- 552 Galley, T. H., Batista, F. M., Braithwaite, R., King, J. and Beaumont, A. R.: Optimisation of larval
553 culture of the mussel *Mytilus edulis* (L.), Aquac. Int., 18(3), 315-325, doi:10.1007/s10499-
554 009-9245-7, 2010.
- 555 Gillikin, D. P., De Ridder, F., Ulens, H., Elskens, M., Keppens, E., Baeyens, W. and Dehairs, F.:
556 Assessing the reproducibility and reliability of estuarine bivalve shells (*Saxidomus*
557 *giganteus*) for sea surface temperature reconstruction: Implications for paleoclimate studies,
558 Palaeogeogr. Palaeoclimatol. Palaeoecol., 228(1-2), 70-85,
559 doi:10.1016/j.palaeo.2005.03.047, 2005.
- 560 Gordon, J. and Carriker, M. R.: Growth lines in a bivalve mollusk: subdaily patterns and dissolution
561 of the shell, Science, 202, 519-521, doi:10.1126/science.202.4367.519, 1978.
- 562 Griesshaber, E., Neuser, R. D. and Schmahl, W. W.: The application of EBSD analysis to
563 biomaterials: microstructural and crystallographic texture variations in marine carbonate
564 shells, Semin Soc Esp Miner., 7, 22-34, 2010.
- 565 Grossman, E. L. and Ku, T.: Oxygen and carbon isotope fractionation in biogenic aragonite:
566 Temperature effects, Chem. Geol., 59, 59-74, 1986.
- 567 Hahn, S., Rodolfo-Metalpa, R., Griesshaber, E., Schmahl, W. W., Buhl, D., Hall-Spencer, J. M.,
568 Baggini, C., Fehr, K. T. and Immenhauser, A.: Marine bivalve shell geochemistry and
569 ultrastructure from modern low pH environments: environmental effect versus experimental
570 bias, Biogeosciences, 9, 1897-1914, doi:10.5194/bg-9-1897-2012, 2012.



- 571 Hedegaard, C., Bardeau, J. F. and Chateigner, D.: Molluscan shell pigments: An in situ resonance
572 Raman study, *J. Molluscan Stud.*, 72(2), 157-162, doi:10.1093/mollus/eyi062, 2006.
- 573 Hiebenthal, C., Philipp, E., Eisenhauer, A. and Wahl, M.: Interactive effects of temperature and
574 salinity on shell formation and general condition in Baltic Sea *Mytilus edulis* and *Arctica*
575 *islandica*, *Aquat. Biol.*, 14(3), 289-298, doi:10.3354/ab00405, 2012.
- 576 Hiebenthal, C., Philipp, E. E. R., Eisenhauer, A. and Wahl, M.: Effects of seawater $p\text{CO}_2$ and
577 temperature on shell growth, shell stability, condition and cellular stress of Western Baltic
578 Sea *Mytilus edulis* (L.) and *Arctica islandica* (L.), *Mar. Biol.*, 160, 2073-2087,
579 doi:10.1007/s00227-012-2080-9, 2013.
- 580 Hopkins, J. B. and Farrow, L. A.: Raman microprobe determination of local crystal orientation, *J.*
581 *Appl. Phys.*, 59(4), 1103-1110, doi:10.1063/1.336547, 1985.
- 582 Jones, D. S.: Sclerochronology: Shell record of the molluscan shell, *Am. Sci.*, 71(4), 384-391,
583 1983.
- 584 Jorgensen, C. B.: Bivalve filter feeding revisited, *Mar. Ecol. Prog. Ser.*, 142(1-3), 287-302,
585 doi:10.3354/meps142287, 1996.
- 586 Karampelas, S., Fritsch, E., Mevellec, J. Y., Sklavounos, S. and Soldatos, T.: Role of polyenes in
587 the coloration of cultured freshwater pearls, *Eur. J. Mineral.*, 21(1), 85-97, doi:10.1127/0935-
588 1221/2009/0021-1897, 2009.
- 589 Karney, G. B., Butler, P. G., Speller, S., Scourse, J. D., Richardson, C. A., Schröder, M., Hughes,
590 G. M., Czernuszka, J. T. and Grovenor, C. R. M.: Characterizing the microstructure of
591 *Arctica islandica* shells using NanoSIMS and EBSD, *Geochemistry, Geophys. Geosystems*,
592 13(4), doi:10.1029/2011GC003961, 2012.



- 593 Lohmann, G. and Schöne, B. R.: Climate signatures on decadal to interdecadal time scales as
594 obtained from mollusk shells (*Arctica islandica*) from Iceland, *Palaeogeogr. Palaeoclimatol.*
595 *Palaeoecol.*, 373, 152-162, doi:10.1016/j.palaeo.2012.08.006, 2013.
- 596 Lowenstam, H. A.: Factors affecting the aragonite: calcite ratios in carbonate-secreting marine
597 organisms, *J. Geol.*, 62(3), 284-322, 1954.
- 598 Lowenstam, H. A. and Weiner, S.: *On biomineralization*, Oxford University Press, New York.,
599 1989, pp.324.
- 600 Lutz, R. A.: Paleoeological implications of environmentally-controlled variation in molluscan
601 shell microstructure, *Geobios*, 17, 93-99, doi:10.1016/S0016-6995(84)80161-8, 1984.
- 602 Marali, S. and Schöne, B. R.: Oceanographic control on shell growth of *Arctica islandica*
603 (*Bivalvia*) in surface waters of Northeast Iceland - Implications for paleoclimate
604 reconstructions, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 420, 138-149,
605 doi:10.1016/j.palaeo.2014.12.016, 2015.
- 606 Marchitto, T. M., Jones, G. A., Goodfriend, G. A. and Weidman, C. R.: Precise temporal
607 correlation of Holocene mollusk shells using sclerochronology, *Quat. Res.*, 53, 236-246,
608 doi:10.1006/qres.1999.2107, 2000.
- 609 Merkel, C., Griesshaber, E., Kelm, K., Neuser, R., Jordan, G., Logan, A., Mader, W. and Schmahl,
610 W. W.: Micromechanical properties and structural characterization of modern inarticulated
611 brachiopod shells, *J. Geophys. Res.*, 112(G02008), doi:10.1029/2006JG000253, 2007.
- 612 Mette, M. J., Wanamaker, A. D., Carroll, M. L., Ambrose, W. G. and Retelle, M. J.: Linking large-
613 scale climate variability with *Arctica islandica* shell growth and geochemistry in northern
614 Norway, *Limnol. Oceanogr.*, 61(2), 748-764, doi:10.1002/lno.10252, 2016.



- 615 Milano, S., Schöne, B. R. and Witbaard, R.: Changes of shell microstructural characteristics of
 616 *Cerastoderma edule* (Bivalvia) - A novel proxy for water temperature, *Palaeogeogr.*
 617 *Palaeoclimatol. Palaeoecol.*, doi:10.1016/j.palaeo.2015.09.051, 2015.
- 618 Milano, S., Prendergast, A. L. and Schöne, B. R.: Effects of cooking on mollusk shell structure and
 619 chemistry: Implications for archeology and paleoenvironmental reconstruction, *J. Archaeol.*
 620 *Sci. Reports*, 7, 14-26, doi: 10.1016/j.jasrep.2016.03.045, 2016.
- 621 Milano, S., Schöne, B. R., Wang, S. and Müller, W. E.: Impact of high $p\text{CO}_2$ on shell structure of
 622 the bivalve *Cerastoderma edule*, *Mar. Environ. Res.*, 119, 144-155,
 623 doi:10.1016/j.marenvres.2016.06.002, 2016b.
- 624 Mook, W.: Paleotemperatures and chlorinities from stable carbon and oxygen isotopes in shell
 625 carbonate, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 9(4), 245-263, doi:10.1016/0031-
 626 0182(71)90002-2, 1971.
- 627 Nehrke, G. and Nouet, J.: Confocal Raman microscope mapping as a tool to describe different
 628 mineral and organic phases at high spatial resolution within marine biogenic carbonates: case
 629 study on *Nerita undata* (Gastropoda, Neritopsina), *Biogeosciences*, 8, 3761-3769,
 630 doi:10.5194/bg-8-3761-2011, 2011.
- 631 Nishida, K., Ishimura, T., Suzuki, A. and Sasaki, T.: Seasonal changes in the shell microstructure
 632 of the bloody clam, *Scapharca broughtonii* (Mollusca: Bivalvia: Arcidae), *Palaeogeogr.*
 633 *Palaeoclimatol. Palaeoecol.*, 363-364, 99-108, doi:10.1016/j.palaeo.2012.08.017, 2012.
- 634 Nudelman, F., Gotliv, B. A., Addadi, L. and Weiner, S.: Mollusk shell formation: mapping the
 635 distribution of organic matrix components underlying a single aragonitic tablet in nacre, *J.*
 636 *Struct. Biol.*, 153, 176-187, doi:10.1016/j.jsb.2005.09.009, 2006.



- 637 Pérez-Huerta, A., Dauphin, Y., Cuif, J. P. and Cusack, M.: High resolution electron backscatter
638 diffraction (EBSD) data from calcite biominerals in recent gastropod shells, *Micron*, 42(3),
639 246-51, doi:10.1016/j.micron.2010.11.003, 2011.
- 640 Pérez-Huerta, A., Etayo-Cadavid, M. F., Andrus, C. F. T., Jeffries, T. E., Watkins, C., Street, S. C.
641 and Sandweiss, D. H.: El Niño impact on mollusk biomineralization-implications for trace
642 element proxy reconstructions and the paleo-archeological record. *PLoS One*, 8(2), e54274,
643 doi:10.1371/journal.pone.0054274, 2013.
- 644 Prezant, R. S. and Tan Tiu, A.: Spiral crossed-lamellar shell growth in *Corbicula* (Mollusca:
645 Bivalvia), *Trans. Am. Microsc. Soc.*, 105(4), 338-347, 1986.
- 646 Richardson, C. A.: Molluscs as archives of environmental change, *Oceanogr. Mar. Biol. an Annu.*
647 *Rev.*, 39, 103-164, 2001.
- 648 Rodríguez-Navarro, A. B., CabraldeMelo, C., Batista, N., Morimoto, N., Alvarez-Lloret, P.,
649 Ortega-Huertas, M., Fuenzalida, V. M., Arias, J. I., Wiff, J. P. and Arias, J. L.: Microstructure
650 and crystallographic-texture of giant barnacle (*Austromegabalanus psittacus*) shell, *J. Struct.*
651 *Biol.*, 156, 355-362, doi:10.1016/j.jsb.2006.04.009, 2006.
- 652 Ropes, J. W., Jones, D. S., Murawski, S. A., Serchuk, F. M. and Jearld, A.: Documentation of
653 annual growth lines in ocean quahogs, *Artica islandica* Linne, *Fish. Bull.*, 82(1), 1-19, 1984.
- 654 Rubenstein, D. I. and Koehl, M. A. R.: The mechanisms of filter feeding: some theoretical
655 considerations, *Am. Nat.*, 111, 981-994, 1977.
- 656 Schaffer, H. E., Chance, R. R., Silbey, R. J., Knoll, K. and Schrock, R. R.: Conjugation length
657 dependence of Raman scattering in a series of linear polyenes: Implications for
658 polyacetylene, *J. Chem. Phys.*, 94(6), 4161, doi:10.1063/1.460649, 1991.



- 659 Schöne, B. R.: The curse of physiology—challenges and opportunities in the interpretation of
660 geochemical data from mollusk shells, *Geo-Marine Lett.*, 28, 269-285, doi:10.1007/s00367-
661 008-0114-6, 2008.
- 662 Schöne, B. R.: *Arctica islandica* (Bivalvia): a unique paleoenvironmental archive of the northern
663 North Atlantic Ocean, *Glob. Planet. Change*, 111, 199-225,
664 doi:10.1016/j.gloplacha.2013.09.013, 2013.
- 665 Schöne, B. R. and Gillikin, D. P.: Unraveling environmental histories from skeletal diaries —
666 Advances in sclerochronology, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 373, 1-5,
667 doi:10.1016/j.palaeo.2012.11.026, 2013.
- 668 Schöne, B. R. and Surge, D. M.: Part N , revised , volume 1 , chapter 14 : bivalve sclerochronology
669 and geochemistry, *Treatise Online*, 1(46), 1-24, 2012.
- 670 Schöne, B. R., Freyre Castro, A. D., Fiebig, J., Houk, S. D., Oschmann, W. and Kröncke, I.: Sea
671 surface water temperatures over the period 1884-1983 reconstructed from oxygen isotope
672 ratios of a bivalve mollusk shell (*Arctica islandica*, southern North Sea), *Palaeogeogr.*
673 *Palaeoclimatol. Palaeoecol.*, 212, 215-232, doi:10.1016/j.palaeo.2004.05.024, 2004.
- 674 Schöne, B. R., Fiebig, J., Pfeiffer, M., Gleß, R., Hickson, J., Johnson, A. L. A., Dreyer, W. and
675 Oschmann, W.: Climate records from a bivalved Methuselah (*Arctica islandica*, Mollusca;
676 Iceland), *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 228(1-2), 130-148,
677 doi:10.1016/j.palaeo.2005.03.049, 2005.
- 678 Schöne, B. R., Radermacher, P., Zhang, Z. and Jacob, D. E.: Crystal fabrics and element impurities
679 (Sr/Ca, Mg/Ca, and Ba/Ca) in shells of *Arctica islandica*—Implications for paleoclimate



- 680 reconstructions, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 373, 50-59,
681 doi:10.1016/j.palaeo.2011.05.013, 2013.
- 682 Soldatov, A. A., Gostyukhina, O. L., Borodina, A. V. and Golovina, I. V.: Qualitative composition
683 of carotenoids, catalase and superoxide dismutase activities in tissues of bivalve mollusc
684 *Anadara inaequalis* (Bruguiere, 1789), *J. Evol. Biochem. Physiol.*, 49(4), 3889-398,
685 doi:10.1134/S0022093013040026, 2013.
- 686 Stemmer, K. and Nehrke, G.: The distribution of polyenes in the shell of *Arctica islandica* from
687 North Atlantic localities: a confocal Raman microscopy study, *J. Molluscan Stud.*, 80, 365-
688 370, doi:10.1093/mollus/eyu033, 2014.
- 689 Stott, K. J., Austin, W. E. N., Sayer, M. D. J., Weidman, C. R., Cage, A. G. and Wilson, R. J. S.:
690 The potential of *Arctica islandica* growth records to reconstruct coastal climate in north west
691 Scotland, UK, *Quat. Sci. Rev.*, 29(13-14), 1602-1613, doi:10.1016/j.quascirev.2009.06.016,
692 2010.
- 693 Strömberg, T. and Cary, C.: Growth in length of *Mytilus edulis* L. fed on different algal diets, *J.*
694 *Exp. Mar. Bio. Ecol.*, 76(1), 23-34, doi:10.1016/0022-0981(84)90014-5, 1984.
- 695 Tan Tiu, A.: Temporal and spatial variation of shell microstructure of *Polymesoda caroliniana*
696 (Bivalvia: Heterodonta), *Am. Malacol. Bull.*, 6(2), 199-206, 1988.
- 697 Tan Tiu, A. and Prezant, R. S.: Temporal variation in microstructure of the inner shell surface of
698 *Corbicula fluminea* (Bivalvia: Heterodonta), *Am. Malacol. Bull.*, 7(1), 65-71, 1989.
- 699 Tan Tiu, A. and Prezant, R. S.: Shell microstructural responses of *Geukensia demissa granosissima*
700 (Mollusca: Bivalvia) to continual submergence, *Am. Malacol. Bull.*, 5(2), 173-176, 1987.



- 701 Taylor, J.D. and Kennedy, W.J.: The shell structure and mineralogy of *Chama pellucida* inst
702 electron microscope, Veliger 11: 391-398, 1969.
- 703 Thompson, I., Jones, D. S. and Dreibelbis, D.: Annual internal growth banding and life history of
704 the ocean quahog *Arctica islandica* (Mollusca: Bivalvia), Mar. Biol., 57, 25-34, 1980.
- 705 Wanamaker, A. D., Kreutz, K. J., Borns, H. W., Introne, D. S., Feindel, S., Funder, S., Rawson, P.
706 D. and Barber, B. J.: Experimental determination of salinity, temperature, growth, and
707 metabolic effects on shell isotope chemistry of *Mytilus edulis* collected from Maine and
708 Greenland, Paleoceanography, 22(2), doi:10.1029/2006PA001352, 2007.
- 709 Wanamaker, A. D., Heinemeier, J., Scourse, J. D., Richardson, C. A., Butler, P. G., Eiríksson, J.
710 and Knudsen, K. L.: Very long-lived mollusks confirm 17th century ad tephra-based
711 radiocarbon reservoir ages for North Icelandic shelf waters, Radiocarbon, 50(3), 399-412,
712 2008.
- 713 Wanamaker, A. D., Kreutz, K. J., Schöne, B. R., Maasch, K. A., Pershing, A. J., Borns, H. W.,
714 Introne, D. S. and Feindel, S.: A late Holocene paleo-productivity record in the western Gulf
715 of Maine, USA, inferred from growth histories of the long-lived ocean quahog (*Arctica*
716 *islandica*), Int. J. Earth Sci., 98(1), 19-29, doi:10.1007/s00531-008-0318-z, 2009.
- 717 Wanamaker, A. D., Hetzinger, S. and Halfar, J.: Reconstructing mid- to high-latitude marine
718 climate and ocean variability using bivalves, coralline algae, and marine sediment cores from
719 the Northern Hemisphere, Palaeogeogr. Palaeoclimatol. Palaeoecol., 302(1), 1-9,
720 doi:10.1016/j.palaeo.2010.12.024, 2011a.



- 721 Wanamaker, A. D., Butler, P. G., Scourse, J. D., Heinemeier, J., Eiríksson, J., Knudsen, K. L. and
722 Richardson, C. A.: Surface changes in the North Atlantic meridional overturning circulation
723 during the last millennium, *Paleoceanography*, 3, 237-252, doi:10.1038/ncomms1901, 2012.
- 724 Wanamaker, A. D., Mette, M. J. and Whitney, N.: The potential for the long- lived bivalve *Arctica*
725 *islandica* to contribute to our understanding of past AMOC dynamics, *US CLIVAR Var.*, 14,
726 13-19, 2016.
- 727 Wanamaker, A. D., Kreutz, K. J., Schöne, B. R. and Introne, D. S.: Gulf of Maine shells reveal
728 changes in seawater temperature seasonality during the Medieval Climate Anomaly and the
729 Little Ice Age, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 302(1), 43-51,
730 doi:10.1016/j.palaeo.2010.06.005, 2011b.
- 731 Weiner, S. and Addadi, L.: Acidic macromolecules of mineralized tissues: the controllers of crystal
732 formation, *Trends Biochem. Sci.*, 16, 252-256, 1991.
- 733 Weiner, S. and Addadi, L.: Design strategies in mineralized biological materials, *J. Mater. Chem.*,
734 7, 689-702, doi:10.1039/a604512j, 1997.
- 735 Widdows, J.: Physiological ecology of mussel larvae, *Aquaculture*, 94(2-3), 147-163,
736 doi:10.1016/0044-8486(91)90115-N, 1991.
- 737 Winter, J. E.: Über den Einfluß der Nahrungskonzentration und anderer Faktoren auf
738 Filtrierleistung und Nahrungsausnutzung der Muscheln *Arctica islandica* und *Modiolus*
739 *modiolus*, *Mar. Biol.*, 4(2), 87-135, doi:10.1007/BF00347037, 1969.
- 740 Witbaard, R., Franken, R. and Visser, B.: Growth of juvenile *Arctica islandica* under experimental
741 conditions, *Helgolaender Meeresuntersuchungen*, 51, 417-431, doi:10.1007/BF02908724,
742 1997.




743 Witbaard, R., Duineveld, G. C. A. and de Wilde, P. A. W. J.: Geographical differences in growth
 744 rates of *Arctica islandica* (Mollusc: Bivalvia) from the North Sea and adjacent waters, J. Mar.
 745 Biol. Assoc. United Kingdom, 79, 907-915, 1999.

746

747

748 Figures and tables



749

750 **Fig. 1.** Shell of adult *Arctica islandica* used in the temperature experiment (left) and juvenile from the Baltic
 751 Sea used in the food experiment (right). The map indicates the localities where the two sets of shells were
 752 collected: Jonesport, Maine (circle) and Kiel Bay (square).




Fig. 2. Raman spectrum of *Arctica islandica* showing the typical aragonite peaks (grey line). The exact position of the polyene peaks R_1 and R_4 was determined by using a peak fitting routine based on a Gaussian function (black line).





Fig. 3. *Arctica islandica* shell growth under controlled conditions. (a) Total growth and (b) instantaneous growth rate during the temperature experiment. (c) Total growth during the food experiment.



765


766 **Fig. 4.** Effect of temperature increase on biomineral orientation. (a) Position of the Raman maps of the three
 767 specimens reared at 10 °C and 15 °C. Dotted lines indicate the location of the calcein marks. dog = direction
 768 of growth. (b) Raman spectral maps of R_{v1}/T_a . Left images of each column represents shell portion formed



769 at 10 °C, right images represent shell portions formed at 15 °C. First row of pairs refers to oOSL, the other
 770 two represent the iOSL. Scale bars = 10 µm. (c) Proportions of biominerals with $R_{v1/Ta} > 2.5$ a.u. with respect
 771 to the total map area. Asterisks indicate significant difference between the orientation of iOSL
 772 microstructures formed at 10 and 15°C ($p < 0.05$).

773

774



775

776



Fig. 5. SEM images of *Arctica islandica* shell microstructures formed at 10 °C (left column) and at 15 °C (right column). The sketch indicates the position of the images 1 mm away from the calcein mark (grey line). The first row of images refers to the oOSL, the other two row refers to the iOSL. Scale bars if not otherwise indicated = 5 µm.





Fig. 6. Effect of different diets based on (a) food type 1, (b) food type 2 and (c) ambient food on biomineral orientation. The optical microscope images indicate the position of the Raman scans. Dotted line marks the start of the experiment. The portion of shell prior the line was formed during the acclimation phase. dog = direction of growth. The Raman spectral maps indicate the ratio $R_{v1/Ta}$ for each data point of the scan. For



788 each shell, maps on the left represent shell portions during the experiment, maps on the right represent shell
789 portions formed during the acclimation phase. In the acclimation portion of the sample reared with ambient
790 food, a significant change in the microstructure orientation is visible. The respective area of the Raman map
791 was not considered in further calculations because it was influenced by the emersion and transportation
792 stress at the start of the experiment. Scale bars = 10 μm . The graphs show the proportions of biominerals of
793 oOSL and iOSL with $R_{\text{vI}/\text{Ta}} > 2.5$ a.u. with respect to the total map area.


794

795



796

797 **Fig. 7.** SEM images of *Arctica islandica* shell microstructures formed during the acclimation phase at AWI
 798 (left column) and during the food experiment (right column). Scale bars = 4 μ m.



799

800 **Fig. 8.** Effects of diet on shell pigment distribution. (a) Raman spectral maps of the 1524 cm⁻¹ band
 801 representing the distribution of the polyenes in the shell cultured with food type 2. Dotted line marks the
 802 start of the experiment. dog = direction of growth. (b) The graph shows the thickness of the pigmented layer
 803 over the whole shell thickness before and during the food experiments.

804

805



Table 1. List of the studied specimens of *Arctica islandica* and experimental conditions.

Sample ID	Locality	Age	Experiment	Treatment
A2	Maine	5	Temperature	10 °C + 15 °C
A4	Maine	4	Temperature	10 °C + 15 °C
A5	Maine	4	Temperature	10 °C + 15 °C
S12	Kiel Bay	1	Diet	Food 1
S14	Kiel Bay	1	Diet	Food 1
S15	Kiel Bay	1	Diet	Food 1
G11	Kiel Bay	1	Diet	Food 2
G12	Kiel Bay	1	Diet	Food 2
G15	Kiel Bay	1	Diet	Food 2
N13	Kiel Bay	1	Diet	No additional food
N15	Kiel Bay	1	Diet	No additional food

Table 2. Details of the pigment composition of the *Arctica islandica* shells used in the food experiment.

The position of the major polyene peaks R_1 and R_4 in the Raman spectrum is indicated together with the number of single and double carbon bonds of the pigment molecular chain (N_1 and N_4). Each shell was analyzed in the portions formed before and during the experimental phase.

Sample ID	Shell portion	R_1 (cm ⁻¹)	R_4 (cm ⁻¹)	N_1	N_4
S12	Acclimation	1130.9	1515.2	9.7	10.8
	Food 1	1121.4	1515.3	12.1	10.7
S14	Acclimation	1133.2	1519.4	9.3	10.2
	Food 1	1132.2	1518.6	9.5	10.3
S15	Acclimation	1129.5	1516.5	10.0	10.6
	Food 1	1132.1	1519.8	9.5	10.1
G11	Acclimation	1132.6	1518.4	9.4	10.3
	Food 2	1129.5	1517.0	10.0	10.5
G12	Acclimation	1131.7	1518.7	9.6	10.3
	Food 2	1132.1	1518.2	9.5	10.4
G15	Acclimation	1132.4	1519.5	9.4	10.2
	Food 2	1128.0	1520.9	10.3	10.0
N13	Acclimation	1130.2	1515.6	9.9	10.7
	Ambient food	1131.4	1514.1	9.6	10.9
N15	Acclimation	1117.9	1516.0	13.3	10.6
	Ambient food	1130.7	1517.0	9.8	10.5
Average		1129.7 ± 4.2	1517.5 ± 2.0	10.1 ± 1.1	10.4 ± 0.3