

1 **Biochar reduces yield-scaled emissions of reactive nitrogen gases from**  
2 **vegetable soils across China**

3 Changhua Fan, Hao Chen, Bo Li, Zhengqin Xiong\*

4 Jiangsu Key Laboratory of Low Carbon Agriculture and GHGs Mitigation, College of Resources and Environmental  
5 Sciences, Nanjing Agricultural University, Nanjing 210095, China




6 \*Corresponding author (Z. Xiong): E-mail: zqxiong@njau.edu.cn;

7 Tel: +86-25-84395148; Fax: +86-25-84395210

1 **Highlights**

- 2 1. Two contrasting biochars affected GNrI across 4 major vegetable soils in China.
- 3 2. Biochar affects gaseous Nr or yield largely depending on soil types.
- 4 3. Both biochars decreased GNrI with Bw mitigating gaseous Nr while Bm improving yield.


1 **Abstract**

2 Biochar amendment to soil has been proposed as a strategy for sequestering carbon, mitigating climate change and  
3 enhancing crop productivity.  few studies have demonstrated the general effects of different feedstock-derived  
4 biochars on the various gaseous reactive nitrogen emissions (GNrEs, N<sub>2</sub>O, NO and NH<sub>3</sub>) simultaneously across the  
5 typical vegetable soils in China. A greenhouse pot experiment with five consecutive vegetable crops was conducted to  
6 investigate the effects of two contrasting biochar, namely, wheat straw biochar (Bw) and swine manure biochar (Bm) on  
7 GNrEs, vegetable yield and gaseous reactive nitrogen intensity (GNrI) in four typical vegetable s from Acrisol  
8 (Hunan province), Anthrosol (Shanxi province), Cambisol (Shandong province) and Phaeozem (Heilongjiang province)  
9 which are representative of the intensive vegetable ecosystems across mainland China. Results showed that remarkable  
10 GNrE mitigation induced by biochar occurred in Anthrosol and Phaeozem, whereas enhancement of yield occurred  
11 in Cambisol and Phaeozem. Additionally, both biochars decreased GNrI through reducing N<sub>2</sub>O and NO emissions by  
12 36.4–59.1 % and 37.0–49.5 % for Bw (except for Cambisol), respectively,  through improving yield by 13.5–30.5 %  
13 for Bm (except for Acrisol and Anthrosol). Biochar amendments generally stimulated the NH<sub>3</sub> emissions with greater  
14 enhancement from Bm than Bw. We can infer that the biochar's effects on the GNrEs and vegetable yield strongly  
15 depend on the attributes of the soil and biochar. Therefore, in order to achieve the maximum benefits under intensive  
16 greenhouse vegetable agriculture, both soil type and biochar characteristics should be seriously considered before  
17 conducting large-scale biochar applications.

18 **Keyword:** Biochar, Intensive vegetable soil, Gaseous reactive nitrogen emissions (GNrEs), Gaseous reactive  
19 nitrogen intensity (GNrI)

# 1 Introduction

2 Agriculture accounts for an estimated emission of 4.1 (1.7–4.8) Tg N yr<sup>-1</sup> for nitrous oxide (N<sub>2</sub>O) and 3.7 Tg N yr<sup>-1</sup>  
3 for nitric oxide (NO), contributing 60 % and 10 %, respectively, to the total global anthropogenic emissions, largely due  
4 to increases of nitrogen (N) fertilizer application in cropland (Ciais, 2013). The concentration of atmospheric N<sub>2</sub>O, a  
5 powerful, long-lived, greenhouse gas, has increased from 270 parts per billion by volume (ppbv) in the pre-industrial era  
6 to ~ 324 ppbv (Ussiri and Lal, 2013); it has 265 times the global warming potential of carbon dioxide (CO<sub>2</sub>) on a  
7 100-year horizon (IPCC, 2013) and also causes depletion of the ozone layer in the atmosphere (Ravishankara et al.,  
8 2009). In contrast, NO<sub>x</sub>, which is mainly emitted as NO, does not directly affect the earth's radiative balance but  
9 catalyzes the production of tropospheric ozone (O<sub>3</sub>), which is a greenhouse gas associated with detrimental effects on  
10 human health (Anenberg et al., 2012) and crop production (Avnery et al., 2011). Finally, ammonia (NH<sub>3</sub>) volatilization is  
11 one of the major N loss pathways (Harrison and Webb, 2001) as well, with up to 90% coming from agricultural activities  
12 (Misselbrook et al., 2000; Boyer et al., 2002). As a natural component and a dominant atmospheric alkaline gas, NH<sub>3</sub>  
13 plays an important role in atmospheric chemistry and ambient aerosol formation (Langridge et al., 2012; Wang et al.,  
14 2015b). In addition to nutrient enrichment (eutrophication) of terrestrial and aquatic systems and global acidification of  
15 precipitation, NH<sub>3</sub> has also been shown to be a major factor in the formation of atmospheric particulate matter and  
16 secondary aerosols (Kim et al., 2006; Pinder et al., 2007), leading to potentially adverse effects on human and ecosystem  
17 health such as visibility degradation and threats to biodiversity (Powlson et al., 2008; Behera et al., 2013).

18 In China, vegetable production devotes an area of approximately  $24.7 \times 10^6$  ha, equivalent to 12.4% of the total  
19 available cropping area, and the production represented 52 % of the world vegetable production in 2012 (FAO, 2015).  
20 Intensified vegetable cultivation in China is characterized by high N application rates, high cropping index and frequent  
21 farm practices. Annual N fertilizer inputs for intensively managed vegetable cultivation in rapidly developing areas are  
22 3–6 times higher than in cereal grain cultivation in China (Ju et al., 2006; Diao et al., 2013; Wang et al., 2015a). As a  
23 result, great concern exists about excess N fertilizer application, leading to low use efficiency in intensive vegetable  
24 fields in China (Deng et al., 2013; Diao et al., 2013; Li et al., 2016). Meanwhile, intensive vegetable agriculture is  
25 considered to be an important source of N<sub>2</sub>O (Xiong et al., 2006; Jia et al., 2012; Li et al., 2015b; Zhang et al., 2015) and  
26 NO production (Mei et al., 2009). Moreover, NH<sub>3</sub> volatilization is another important N pathway in fertilized soil,  
27 resulting in large losses of soil-plant N (Pacholski et al., 2008; Zhang et al., 2011). Therefore, the reduction of reactive N  
28 loss is key to meet the joint challenges of high production and acceptable environmental consequences  intensive  
29 vegetable production (Zhang et al., 2013).

30 Biochar is the dark-colored, carbon (C)-rich residue of pyrolysis or gasification of plant biomass under oxygen

1 (O<sub>2</sub>)-limited conditions, specifically produced for use as a soil amendment (Sohi, 2012). The amendment of agricultural  
2 ecosystems with biochar has been proposed as an effective countermeasure for climate change (Smith, 2016). These  
3 additions have been suggested to increase soil carbon storage (Mukherjee and Zimmerman, 2013; Stavi and Lal, 2013),  
4 decrease greenhouse gas emissions (Li et al., 2016), and improve soil fertility and crop production (Major et al., 2010;  
5 Liu et al., 2013). However, some recent studies have reported no difference or even an increase in soil N<sub>2</sub>O emissions  
6 induced by biochar application for various soils (Saarnio et al., 2013; Wang et al., 2015a). Besides, NH<sub>3</sub> volatilization  
7 was enhanced by biochar application in pasture soil (Clough et al., 2010), vegetable soil (Sun et al., 2014) and paddy soil  
8 in the wheat-growing season (Zhao et al., 2014). Additionally, crop productivity responses to biochar amendments  
9 differed among various biochars (Cayuela et al., 2014). These inconsistent results suggest that current biochar application  
10 to soil is not a “one-size fit-all paradigm” because of the variation in the physical and chemical characteristics of the  
11 different biochars, soil types and crop species (Field et al., 2013; Cayuela et al., 2014). Moreover, limited types of  
12 biochar (Spokas and Reicosky, 2009) and soil (Sun et al., 2014) were involved in the experiments in previous studies.  
13 Thus, the evaluation of the different types of biochar under the typical soils is imperative to gain a comprehensive  
14 understanding of potential interactions before the large-scale application of biochars.

15 Therefore, a greenhouse pot experiment was conducted in an effort to investigate the effects of different types of  
16 biochar on gaseous reactive nitrogen emissions (GNrEs), namely, N<sub>2</sub>O, NO and NH<sub>3</sub>, simultaneously in four intensively  
17 cropped vegetable soils across main vegetable production areas of mainland China. We hypothesized that: 1) biochar  
18 amendment could affect GNrEs, vegetable yield and yield-scaled gaseous reactive nitrogen emissions, namely, gaseous  
19 reactive nitrogen intensity (GNrI) in vegetable soils across mainland China, 2) those influences would vary among  
20 biochar and soil types. Overall, the objectives of this research were to gain a comprehensive insight into the effects of  
21 two contrasting biochars on the GNrEs, vegetable yield and GNrI in intensively managed vegetable production in China.

## 1 **2 Materials and methods**

### 2 *2.1. Experimental soil and biochar*

3 Four typical greenhouse vegetable cultivation sites with a long history (more than 10 years) of conventional  
4 cultivation were selected from Northeast, Northwest, Central and Eastern China (Fig. S1): 1. a Phaeozem from Jiamusi  
5 (46°48' N, 130°12' E) in the Heilongjiang province, 2. an Anthrosol from Yangling (34°18' N, 108°2' E) in the Shanxi  
6 province, 3. an Acrisol from Changsha (28°32' N, 113°23' E) in the Hunan province, 4. a Cambisol from Shouguang  
7 (36°56' N, 118°38' E) in the Shandong province (FAO and ISRIC, 2012). Those four types of vegetable soil represented  
8 a range of differences in physicochemical properties and regions (Table S1). Soil samples were manually collected from  
9 the cultivated layer (0–20 cm) after the local vegetable harvest in April, 2015. The samples were air-dried and passed  
10 through a 5 mm stainless steel mesh sieve and homogenized thoroughly. Any visible roots and organic residues were  
11 removed manually before being packed with the necessary amount of soil to achieve the initial field bulk density. Each  
12 pot received 15 kg of 105 °C dry-weight-equivalent fresh soil. For each of the biochar amendment pot, 282.6 g pot<sup>-1</sup>  
13 sieved biochar (2 mm) was mixed with the soil thoroughly before the experiment, which was equivalent to a 40 t ha<sup>-1</sup>  
14 biochar dose (dry weight). No more biochar was added later in the experimental period.

15 The two types of biochar that were used in this experiment are derived from two common agricultural wastes in  
16 China: wheat straw and swine manure, hereafter referred to as Bw and Bm, respectively (Table S1). The Bw was  
17 produced at the Sanli New Energy Company in Henan, China, by pyrolysis and thermal decomposition at 400–500 °C.  
18 The Bm was produced through thermal decomposition at 400 °C by the State Key Laboratory of Soil Science and  
19 Sustainable Agricultural, Institute of Soil Science, Chinese Academy of Sciences. In accordance with Lu (2000), soil  
20 organic carbon (SOC) was measured by wet digestion with H<sub>2</sub>SO<sub>4</sub>–K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, total nitrogen (TN) was determined by  
21 semi-micro Kjeldahl digestion, and soil texture was determined with the pipette method. The soil pH and biochar pH  
22 were measured in deionized water at a volume ratio of 1:2.5 (soil to water) with a PHS-3C mv/pH detector (Shanghai  
23 Kangyi Inc. China). Biochar content of hydrogen (H) was measured by elemental analysis after dry combustion (Euro  
24 EA, Hekatech GmbH, Wegberg, Germany). The oxygen content of biochar was measured with the same device after  
25 pyrolysis of the sample at 1000 °C followed by reduction of the evolved O<sub>2</sub> to CO and quantified by GC-TCD. The soil  
26 nitrate (NO<sub>3</sub><sup>-</sup>-N) and ammonium (NH<sub>4</sub><sup>+</sup>-N) were measured following the two-wavelength ultraviolet spectrometry and  
27 indophenol blue method, respectively, using an ultraviolet spectrophotometer (HITACHI, UV-2900, Tokyo, Japan).  
28 Electric conductivity (EC) was measured by using a Mettler-Toledo instrument (FE30-K, Shanghai, China) at a 1:5 (w:v)  
29 soil to water ratio. Cation exchange capacity (CEC) was determined using the CH<sub>3</sub>COONH<sub>4</sub> method. Dissolved organic  
30 carbon (DOC) was extracted from 5 g of the biochar/soil with an addition of 50 ml deionized water and measured by a

1 TOC analyzer (TOC-2000/3000, Metash Instruments Co., LTD, Shanghai, China). Ash content was measured by heating  
2 the biochars at 750 °C for 4 h. The specific surface area of the biochar material was tested using the Brunauer–Emmett–  
3 Teller (BET) method, from which the N adsorption–desorption isotherms at 77 K were measured by an automated gas  
4 adsorption analyzer ASAP2000 (Micromeritics, Norcross, GA) with + 5% accuracy. Scanning electron microscopy (SEM)  
5 imaging analysis was conducted using a HITACHI S-3000N scanning electron microscope.

## 6 *2.2. Experimental set-up and management*

7 The pot experiments were performed at the greenhouse experimental station of Nanjing Agricultural University,  
8 China. Five vegetable crops were grown successively in the four vegetable soils during the experimental period. For each  
9 type of soil, three treatments with three replicates were arranged in a random design: urea without biochar (N), urea with  
10 wheat straw biochar (N+Bw), urea with swine manure biochar (N+Bm). In addition, phosphate and potassium fertilizers  
11 in the form of calcium magnesium phosphate and potassium chloride, together with urea, were broadcasted and mixed  
12 with soil thoroughly prior to sowing the vegetables. No topdressing events occurred because of the frequent cultivation  
13 and short growth period for the leafy vegetables. Based on the vegetable growth, all pots received equal amounts of water  
14 and no precipitation. Detailed information on the pot management practices is provided in Table S2.


15 Each pot consists of a 30 cm × 30 cm (height × diameter) cylinder made of polyvinyl chloride (PVC). The top of  
16 each pot was surrounded by a special water-filled trough collar, which allowed a chamber to sit on the pot and prevent  
17 gas exchange during the gas-sampling period. Small holes (diameter of 1 cm) at the bottom of the pots were designed for  
18 drainage. To prevent soil loss, a fine nylon mesh (< 0.5 mm) was attached to the base of the soil cores before packing.

## 19 *2.3. Measurement of N<sub>2</sub>O, NO and NH<sub>3</sub>*

20 The NO and N<sub>2</sub>O fluxes were measured simultaneously from each vegetable cultivation using a static opaque  
21 chamber method (Zheng et al., 2008; Yao et al., 2009). A square PVC chamber of 35 cm × 35 cm × 40 cm (length ×  
22 width × height) was temporarily mounted on the pot for gas flux measurement. The chamber was coated with sponge and  
23 aluminum foil outside to prevent solar radiation heating the chamber. Gas samples for flux measurements were collected  
24 between 8 and 10 a.m. on each measuring day to minimize the influence of diurnal temperature variation. Gas fluxes  
25 were usually measured once a week and every other day for one week following fertilizer application. To measure the  
26 N<sub>2</sub>O flux, four samples were collected from the headspace chamber using 20 ml polypropylene syringes at 0, 10, 20, and  
27 30 min after chamber closure. The gas concentrations in the samples were analyzed within 12 h after sampling using an  
28 Agilent 7890A gas chromatograph equipped with an electron capture detector (ECD) for N<sub>2</sub>O detection. Argon-methane  
29 (5 %) was used as the carrier gas at a flow rate of 40 ml min<sup>-1</sup>. The column and ECD temperatures were maintained at 40  
30 and 300 °C, respectively. The gas chromatography configurations described by Wang et al. (2013) were adopted for the

1 gas concentration analysis. N<sub>2</sub>O flux was calculated using the linear increases in gas concentration with time. Sample sets  
2 were rejected unless they yielded a linear regression value of R<sup>2</sup> > 0.90.

3 For each NO flux measurement, gas samples were collected from the same chamber that was used for the N<sub>2</sub>O flux  
4 measurements (Yao et al., 2009). Before closing the chamber, an approximately 1.0 L gas sample from the headspace of  
5 each chamber was extracted into an evacuated sampling bag (Delin Gas Packing Co., LTD, Dalian, China), and this  
6 measurement was regarded as time 0 min for NO analysis. After 30 min under chamber enclosure conditions (i.e., after  
7 the N<sub>2</sub>O sample collections were completed), another headspace gas sample with the same volume was extracted from  
8 each chamber into another evacuated bag. Within 1 h after sampling, NO concentrations were analyzed by a model 42i  
9 chemiluminescence NO–NO–NO<sub>x</sub> analyzer (Thermo Environmental Instruments Inc., Franklin, MA, USA). The NO  
10 fluxes were derived from the concentration differences between the two collected samples. The NO<sub>x</sub> analyzer was  
11 calibrated by a model 146i dynamic dilution calibrator system at the end of each crop-growing season.

12 The mean flux of N<sub>2</sub>O or NO during the experiment period is the average of all measured fluxes weighted by the  
13 interval between two neighboring measurements (Xiong et al., 2006). The cumulative N<sub>2</sub>O  calculated as the product  
14 of the mean flux and the entire duration.

15 The NH<sub>3</sub> volatilization was determined using the ventilation method (Zhao et al., 2010). The  
16 phosphoglycerol-soaked sponge was replaced every day after each fertilization event for approximately one week. The  
17 phosphoglycerol-soaked sponges used to collect the NH<sub>3</sub> samples were immediately extracted with 300 mL potassium  
18 chloride (KCl) solution (1 mol L<sup>-1</sup>) for 1 h. The concentration of NH<sub>4</sub><sup>+</sup>-N was measured using the indophenol blue  
19 method at 625 nm (Sororzano, 1969) by ultraviolet spectrophotometry (HITACHI, UV-2900, Tokyo, Japan, with 0.005  
20 absorbance of photometric accuracy). The cumulative seasonal NH<sub>3</sub> volatilization was the sum of the daily emissions  
21 during the measurement period.

#### 22 2.4. Auxiliary measurements

23 Simultaneously with the determination of trace gas fluxes, the air temperature and the soil temperature at a depth of  
24 5 cm were measured using thermally sensitive probes at each sampling date. Soil water content was also measured using  
25 a portable water detector (Mode TZS-1K, Zhejiang Top Instrument Corporation Ltd., China) by the frequency domain  
26 reflectometer method at a depth of 5 cm. Measured soil water contents (v/v) were converted to water filled pore space  
27 (WFPS) with the following equation:

$$28 \text{WFPS} = \text{volumetric water content (cm}^3 \text{ cm}^{-3}) / \text{total soil porosity (cm}^3 \text{ cm}^{-3}) \quad (1)$$

29 Here, total soil porosity = [1 - (soil bulk density (g cm<sup>-3</sup>) / 2.65)] with an assumed soil particle density of 2.65 (g cm<sup>-3</sup>).

30 The total soil bulk density was determined with the cutting ring method according to Lu (2000).



1 After each vegetable crop reached physiological maturity, the fresh vegetable yield was measured by weighing the  
2 whole aboveground and belowground biomass in each pot.

$$3 \text{GNrE} = \text{cumulative N}_2\text{O} + \text{cumulative NO} + \text{cumulative NH}_3 \text{ emissions (kg N t}^{-1}\text{)} \quad (2)$$

$$4 \text{GNrI} = \text{GNrE} / \text{vegetable fresh yield (kg N t}^{-1}\text{ yield)} \quad (3)$$

5 After the one-year pot experiment, a soil sample from each pot was blended carefully. One subsample was stored at  
6 4 °C for determination of microbial biomass carbon (MBC), potential nitrification rate (PNR) and denitrification enzyme  
7 activity (DEA) within 3 days. Another subsample was air-dried for analysis of SOC, TN, pH and EC. MBC was  
8 determined by substrate-induced respiration using a gas chromatography (Anderson and Domsch 1978). PNR was  
9 measured using the chlorate inhibition soil-slurry method as previously described (Kurola et al., 2005) with  
10 modifications (Hu et al., 2016). DEA was quantified as described by Smith and Tiedje (1979).

### 11 *2.5. Data processing and statistics*

12 Two-way ANOVA was used to analyze the effects of the biochar type, soil type, and their interactions on soil  
13 properties, N<sub>2</sub>O, NO and NH<sub>3</sub> emissions, vegetable yield, GNrE and GNrI throughout the experimental period. Multiple  
14 comparisons among the treatments were assessed using Tukey's HSD test. Significant differences were considered at  $P <$   
15 0.05. All statistical analyses were performed using JMP ver. 7.0 (SAS Institute, Cary, NC, USA, 2007). Pearson's  
16 correlation analysis was used to determine whether there were significant interrelationships between N<sub>2</sub>O/NO and PNR  
17 or DEA in each soil, using SPSS window version 18.0 (SPSS Inc., Chicago, USA).

### 1 3. Results

#### 2 3.1. Soil responses to biochar amendment

3 Appreciable differences in all observed soil properties existed among soil types (Table 1), suggesting the wide  
4 variations of soil characteristics across mainland China. Additionally, biochar amendments had significant influences on all  
5 the soil properties (Table 1,  $p < 0.05$ ). Compared with N treatments, biochar amendments increased the SOC, TN and EC  
6 by 20.4–135.0 %, 0.5–21.2 % and 2.4–38.1 %, respectively, across all the soils. Compared with Bw, Bm amendment  
7 increased SOC and TN by 5.8–20.5 % and 9.5–14.2 % ( $p < 0.05$ ), respectively, whereas EC values were higher by 3.3–  
8 21.5 % induced by Bw than Bm amendment over all soils. Additionally, biochar amendments significantly increased soil  
9 pH by 0.27–0.64 and 0.08–0.10 units compared with N treatment in Acrisol and Anthrosol soils ( $p < 0.05$ ), respectively,  
10 and Bm performed better than Bw on increasing soil pH in Acrisol. Furthermore, biochar amendments tended to increase  
11 MBC in Cambisol and Phaeozem, and Bm increased MBC relative to Bw in all soils.

12 As shown in Fig. 1, no consensus effects on PNR and DEA were observed with biochar amendments across all soils.  
13 Compared with N treatment, biochar amendments significantly increased PNR in Phaeozem while exerted no influences  
14 on Cambisol (Fig. 1a). Compared with Bw, Bm amendment significantly increased PNR in Acrisol and Anthrosol.  
15 Moreover, compared with N, biochar amendments reduced DEA in most soils, significantly in Anthrosol and Phaeozem  
16 by an average of 40.1 and 37.8 % (Fig. 1b,  $p < 0.05$ ), respectively. In comparison with Bw, enhancements in DEA were  
17 observed by 42.5 and 74.4 % with Bm amendment in Acrisol and Anthrosol, respectively ( $p < 0.05$ ).

#### 18 3.2. Seasonal variations of $N_2O$ and NO emissions


19 The dynamics of  $N_2O$  fluxes from all N-applied treatments in the four vegetable soils were relatively consistent and  
20 followed a sporadic and pulse-like pattern that was accompanied with fertilization, tillage and irrigation (Fig. 2). In  
21 addition, peak  $N_2O$  fluxes varied greatly. Most of the  $N_2O$  emissions occurred during the Amaranth and Tung choy  
22 growing periods, and there were several small emissions peaks during the Spinach and Coriander herb growing periods  
23 due to lower N application rate (Table S2), soil temperature and water content (Fig. S2). The highest peaks of  $N_2O$   
24 emissions from Acrisol, Anthrosol, Cambisol and Phaeozem were 4133.7, 1784.0, 432.4 and 1777.2  $\mu\text{g N m}^{-2} \text{h}^{-1}$ ,  
25 respectively. Although biochar (Bw and Bm) application did not significantly alter the seasonal pattern of the  $N_2O$  fluxes,  
26 they greatly lowered some peaks of  $N_2O$  emissions in the Anthrosol and Phaeozem by 8.7–74.4% and 23.6–73.6%,  
27 respectively (Fig. 2b and d).

28 Clearly, the NO fluxes demonstrated similar seasonal dynamics to the  $N_2O$  fluxes (Fig. 3). Some relatively high  
29 peak NO fluxes were still observed in the Spinach and Coriander herb planting seasons even though relatively low  
30 temperatures occurred during these periods, primarily due to lower soil moisture which was suitable for NO production.

1 The NO fluxes ranged from -44.6 to 377.6  $\mu\text{g N m}^{-2} \text{h}^{-1}$  across all soil types. Furthermore, some NO peaks were  
2 significantly weakened with the Bw and Bm in the Acrisol (Fig. 3a).

### 3 3.3. Cumulative $\text{N}_2\text{O}$ , NO and $\text{NH}_3$ emissions

4 Cumulative  $\text{N}_2\text{O}$  emissions varied greatly among soil types (Table 3a,  $p < 0.05$ ), from 1.97 to 31.56  $\text{kg N ha}^{-1}$  across  
5 all the soils during the vegetable cultivation period. Biochar amendments had significant influences on the cumulative  
6  $\text{N}_2\text{O}$  emissions (Table 2,  $p < 0.001$ ). In comparison with the N treatment, biochar amendment resulted in no consistent  
7 effects on  $\text{N}_2\text{O}$  emissions over all soils (Table 3a), indicating significant interactions between biochar and soil types  
8 (Table 2,  $p < 0.001$ ). Additionally, Bw amendment decreased  $\text{N}_2\text{O}$  emissions by 11.8–38.4 % across all the soils in  
9 relation to Bm, indicating that Bw performed better mitigation effects than Bm across all the soils, significantly in  
10 Acrisol (Table 3a,  $p < 0.05$ ). The values of cumulative NO emissions were much smaller than those of  $\text{N}_2\text{O}$  emissions,  
11 with a remarkable variation of 0.20–8.99  $\text{kg N ha}^{-1}$  across all soils (Table 3b). Biochar amendments had pronounced  
12 effects on NO emissions (Table 2,  $p < 0.001$ ), but their effects differed between vegetable soils (Table 3b), which  
13 suggested significant interactions between biochar and soil types (Table 2,  $p < 0.001$ ). Compared with Bm, Bw  
14 amendment significantly reduced NO emissions in Anthrosol and Phaeozem (Table 3b,  $p < 0.05$ ). Moreover,  $\text{N}_2\text{O}$   
15 emissions had positive relationships with DEA both in Anthrosol and Phaeozem, and were affected positively with PNR  
16 in Acrisol (Table 4). Additionally, NO emissions had positive correlations with both PNR and DEA in Anthrosol.  
17 However, neither  $\text{N}_2\text{O}$  nor NO emissions were influenced significantly by PNR and DEA in Cambisol.

18 As is shown in Table 3c, the cumulative  $\text{NH}_3$  emissions fluctuated greatly from 4.72–7.57  $\text{kg N ha}^{-1}$  across all the  
19 soils. Biochar amendments produced no significant influences on the  $\text{NH}_3$  emissions relative to N treatment in most soils  
20 (Table 3c). A tendency was found for the cumulative  $\text{NH}_3$  emissions in N  to be higher than those in the N+Bw  
21 treatment, although this difference was not remarkable within each soil. Additionally, stimulation effects were  
22 consistently present after the first fertilization event in each type of soil (Fig. 4).

### 23 3.4. Vegetable yield and gaseous reactive N intensity during the five-vegetable crop rotation

24 The vegetable yields for the five consecutive vegetable crops are presented in Table 3e. Pronounced differences  
25 existed among all soils (Table 3e,  $p < 0.05$ ). Additionally, biochar amendments exerted no significant effects on vegetable  
26 yield (Table 2). Compared with the N treatment, biochar amendments were prone to increase vegetable yield in Cambisol  
27 and Phaeozem against Acrisol and Anthrosol (Tables 3e), denoting pronounced interactions between soil and biochar  
28 (Table 2,  $p < 0.05$ ). Compared with Bm, Bw amendment lowered total yield over all the soils (Table 3e), significantly in  
29 Acrisol and Cambisol ( $p < 0.05$ ).

30 Table 3f presents the GNrI during the whole experiment period, with a pronounced variation among soils ( $p < 0.05$ ).


1 The GNrI was greatly affected by biochar amendment during the whole experiment period (Table 2,  $p < 0.01$ ). Compared  
2 to N treatment, biochar amendments reduced the GNrI by 4.3–27.8 % across all soils, significantly in Anthrosol and  
3 Phaeozem (Table 3f,  $p < 0.05$ ). Moreover, there were no remarkable differences between Bw and Bm throughout all soils.



## 1 **4. Discussion**

### 2 *4.1. Biochar effects on GNRs across different soil types*


3 The effects of biochar amendment on the N<sub>2</sub>O and NO emissions may be positive, negative or neutral, largely  
4 depending on the soil condition and the inherent characteristics of the biochar (Spokas and Reicosky, 2009; Nelissen et  
5 al., 2014). In our study, effects of two biochars on the N<sub>2</sub>O and NO emissions did not follow a consensus trend across the  
6 four typical vegetable soils (Table 3a, b). In agreement with Cayuela et al. (2014), who reported that the role of biochar in  
7 mitigating N<sub>2</sub>O emission was maximal in soils close to pH neutral, remarkable mitigation effects were observed in  
8 Anthrosol and Phaeozem with the biochar amendments (Table 3a). These findings potentially resulted from the effects of  
9 the biochars on soil aeration, C/N ratio and pH, which affected the N dynamics and N cycling processes (Zhang et al.,  
10 2010; Ameloot et al., 2015). In line with Obia et al. (2015), biochar decreased NO emissions in low-pH Acrisol (Table  
11 3b), probably by stimulating denitrification enzyme activity, which then resulted in less NO accumulation relative to N<sub>2</sub>  
12 production. Moreover, the liming effects of biochar may have prevented the chemical decomposition of NO<sub>2</sub><sup>-</sup> to NO  
13 (Islam et al., 2008), leaving only enzymatically produced NO to accumulate. However, different from the Acrisol soils,  
14 neither N<sub>2</sub>O nor NO emission was significantly influenced by PNR or DEA, suggesting other processes might play vital  
15 roles in Cambisol. Besides nitrification and denitrification, nitrifier denitrification (Wrage et al., 2001) and heterotrophic  
16 nitrification (Zhu et al., 2011) can be important processes for producing N<sub>2</sub>O/NO as well, especially in vegetable soils  
17 with low pH, low carbon content and high N content (Wrage et al., 2001). Ma et al. (2015) speculated that nitrifier  
18 denitrification was the main process producing N<sub>2</sub>O in the North China Plain (Cambisol within this region). In addition,  
19 surplus N input in vegetable systems probably masked the beneficial effects of the biochar addition on the N  
20 transformation (Wang et al., 2015a). Therefore, the underlying mechanism of how biochar affect those processes needs to  
21 be illustrated in their further research.

22 On the other hand, different biochars may not produce universal influences on N<sub>2</sub>O emissions for the same soil due  
23 to the distinct properties of the biochar (Spokas and Reicosky, 2009). In the current study, overall, in comparison with  
24 Bm, the Bw amendment had more effective mitigation effects on N<sub>2</sub>O and NO emissions (Table 3a, b), largely due to the  
25 following reasons. First, compared with Bw, Bm had lower the contents of the TN and DOC by 80% and 40% (Table S1),  
26 respectively, which might supply extra N or C source for heterotrophic nitrification in the acidic Acrisol, leading Bm to  
27 being ineffective for reducing the N<sub>2</sub>O emissions (Table 3a). This result was in accordance with Li et al. (2015a), who  
28 observed that biochar amendment had no significant influence on the cumulative N<sub>2</sub>O emissions, and even higher N<sub>2</sub>O  
29 emissions occurred when biochar was input. Additionally, as shown in Fig.1, Bm was more prone to stimulate PNR and  
30 DEA, thus displaying lower mitigation ability than Bw. Second, compared with Bm, the C/N ratio was approximately

1 twofold  Bw (Table S1), presumably leading to more inorganic nitrogen being immobilized in biochar with a higher  
2 C/N ratio (Ameloot et al., 2015), decreasing the available N for microorganisms. Last, as presented in Fig. S3 and Table  
3 S1, Bw had more pores and surface area, having a better advantage over Bm in absorbing NO accordingly. Others have  
4 found that the lower mitigation capacity of high-N biochars (e.g., manures or biosolids) is probably due to the increased  
5 N release in the soil from the biochar (Schouten et al., 2012). To our knowledge, very few studies have investigated  
6 biochar effects on NO emissions (Nelissen et al., 2014; Obia et al., 2015), and the mechanisms through which biochar  
7 influence NO emissions are not elucidated yet. Therefore, more research is needed to clarify the underlying mechanisms  
8 of biochar on NO emission.


9 Intensively managed soils receiving fertilizer such as urea or anhydrous NH<sub>3</sub> and ruminant urine patches are  
10 potential hot spots for NH<sub>3</sub> formation, where the use of biochar is expected to retain NH<sub>3</sub>-N in the soil system (Clough  
11 and Condon, 2010). Actual  the effects of biochar amendments on NH<sub>3</sub> volatilization largely depend on soil  
12 characteristics, biochar types. Soil texture is an important factor impacting NH<sub>3</sub> transfer and release. Moisture  lay contents  
13 were present in the Anthrosol (Table S1), which was limited in large soil pores, thus, the addition of porous biochar could  
14 enhance the soil aeration, promoting NH<sub>3</sub> volatilization (Sun et al., 2014). Additionally, it was worthy to note that  
15 cumulative NH<sub>3</sub> emissions were slightly higher in soils with the Bm than those with the Bw amendment (Fig. 4 and Table  
16 3c) and that difference could presumably be attributed to less surface area and the much higher pH of Bm (Fig. S3 and  
17 Table S1), resulting in weak adsorption and great liming effects.

#### 18 4.2. Biochar effects on vegetable yield and GNrI across different soil types

19 The application of biochar is usually intended to increase crop yields, and evidence suggests this may be successful  
20 (Schulz et al., 2013; Li et al., 2016). Due to its liming effect, biochar helps to improve the supply of essential macro- and  
21 micronutrients for plant growth (Chan and Xu, 2009; Major et al., 2010). Enhancement of vegetable yield with biochar  
22 amendment occurred in Cambisol and Phaeozem (Table 3e). Additionally, the effects of Bm and Bw on vegetable yield  
23 were inconsistent, which  ably due to the wide diversity of physicochemical characteristics of biochar that translates  
24 into variable reactions in soil (Novak et al., 2014). First, compared to Bw, Bm has a higher DOC content (Table S1),  
25 through which more nutrients may be directly introduced to the soil (Rajkovich et al., 2012). Secondly, besides their large  
26 amount of plant-available nutrients (Hass et al., 2012), biochars produced with manure have been generally considered  
27 significant for improving soil fertility by promoting soil structure development (Joseph et al., 2010), with the result that  
28 Bm was found superior to Bw in vegetable production enhancement in our case (Table 3e). As biochar effects on  
29 vegetable yield were variable, both biochar properties and soil conditions and crop species ought to be taken into account  
30 comprehensively before applying biochar to a certain soil condition.

1           However, no promotion of yield was observed with biochar amendments in Acrisol and Anthrosol. This could be  
2 attributed to exacerbated soil salinity which inhibited the uptake of nutrients and water (Ju et al., 2006; Zhou et al., 2010)  
3 and the growth of the soil microorganisms (Setia et al., 2011), leading to unsustainable greenhouse vegetable production.  
4 Compared with other biochar (Jia et al., 2012), the higher amounts of ash in Bw and Bm may contain high salts, which  
5 would result in soil salinity (Hussain et al., 2016). After the addition of the two salt-rich biochars, the EC values of  
6 Acrisol and Anthrosol vegetable soils increased, which might reach the limits to tolerance for the leafy vegetables  
7 (Shannon and Grieve, 1998). Here, we assessed two feedstock-derived biochar effects on GNrI in typical cultivated  
8 vegetable soils across mainland China. Overall, biochar amendments reduced GNrI over all the soils, with the magnitude  
9 largely depending on soil type. Remarkable reduction in GNrI had been detected due to the efficient mitigation induced  
10 by biochar in Anthrosol and Phaeozem (Table 3f). However, despite enhanced vegetable yield, no significant decreases in  
11 GNrI were observed in Cambisol, mainly because of the absence of mitigation effects on N<sub>2</sub>O, NO and NH<sub>3</sub> emissions of  
12 biochars (Table 3a, b and c). Overall, Bw was superior to Bm in mitigating the GNrE while Bm performed better in  
13 vegetable yield enhancement (Table 3d and e). Therefore, the mitigation efficacies on GNrI were not notably different  
14 between Bw and Bm amendments across the four soils.

1 **5. Conclusion**

2 The study demonstrated that biochar amendments mostly reduced N<sub>2</sub>O and NO emissions and slightly increased the  
3 NH<sub>3</sub> emiss  ile produced no consensus influences on yield though those effects were largely both biochar- and  
4 soil-specific. Additionally, biochar amendments did decrease GNrI in intensive vegetable soils across mainland China.  
5 Furthermore, Bw was superior to Bm in mitigating the GNrE whereas the Bm performed better in crop yield throughout  
6 all soils. Consequently, both soil type and biochar characteristics need to be seriously considered before large-scale  
7 biochar application under certain regions of intensive vegetable production.



1     **Acknowledge**

2     We would like to express our heartfelt gratitude to the editor and anonymous reviewers for their constructive comments  
3     that greatly improved the manuscript. This work was jointly supported by the National Natural Science Foundation of  
4     China (41471192), Special Fund for Agro-Scientific Research in the Public Interest (201503106) and the Ministry of  
5     Science and Technology (2013BAD11B01).

## 1 **References**

- 2 Ameloot, N., Sleutel, S., Das, K. C., Kanagaratnam, J., and Neve, S. D.: Biochar amendment to soils with contrasting  
3 organic matter level: effects on N mineralization and biological soil properties, *Global Change Biology Bioenergy*, 7,  
4 135–144, 2015.
- 5 Anderson, J. and Domsch, K.: A physiological method for the quantitative measurement of microbial biomass in soils,  
6 *Soil biology and biochemistry*, 10, 215–221, 1978.
- 7 Anenberg, S. C., Schwartz, J., Shindell, D., Amann, M., Faluvegi, G., Klimont, Z., Janssensmaenhout, G., Pozzoli, L.,  
8 Van, D. R., and Vignati, E.: Global Air Quality and Health Co-benefits of Mitigating Near-Term Climate Change  
9 through Methane and Black Carbon Emission Controls, *Environmental Health Perspectives*, 120, 831–839, 2012.
- 10 Avnery, S., Mauzerall, D. L., Liu, J., and Horowitz, L. W.: Global crop yield reductions due to surface ozone exposure: 1.  
11 Year 2000 crop production losses and economic damage, *Atmospheric Environment*, 45, 2284–2296, 2011.
- 12 Behera, S. N., Sharma, M., Aneja, V. P., and Balasubramanian, R.: Ammonia in the atmosphere: a review on emission  
13 sources, atmospheric chemistry and deposition on terrestrial bodies, *Environmental Science and Pollution Research*,  
14 20, 8092–8131, 2013.
- 15 Boyer, E. W., Goodale, C. L., Jaworski, N. A., and Howarth, R. W.: Anthropogenic nitrogen sources and relationships to  
16 riverine nitrogen export in the northeastern USA. In: *The Nitrogen Cycle at Regional to Global Scales*, Springer,  
17 2002.
- 18 Cayuela, M., Van Zwieten, L., Singh, B., Jeffery, S., Roig, A., and Sánchez-Monedero, M.: Biochar's role in mitigating  
19 soil nitrous oxide emissions: A review and meta-analysis, *Agriculture, Ecosystems & Environment*, 191, 5–16, 2014.
- 20 Chan, K. Y. and Xu, Z.: Biochar: nutrient properties and their enhancement, *Biochar for environmental management:*  
21 *science and technology*, 2009. 67–84, 2009.
- 22 Ciais, P.: Carbon and other biogeochemical cycles: Final draft underlying scientific technical assessment, IPCC  
23 Secretariat, Geneva, 2013. 2013.
- 24 Clough, T. J. and Condon, L. M.: Biochar and the nitrogen cycle: introduction, *Journal of Environmental Quality*,  
25 39, 1218-1223, 2010.
- 26 Clough, T. J., Bertram, J. E., Ray, J. L., Condon, L. M., O'Callaghan, M., Sherlock, R. R., and Wells, N. S.: Unweathered  
27 Wood Biochar Impact on Nitrous Oxide Emissions from a Bovine-Urine-Amended Pasture Soil, *Soil Science*  
28 *Society of America Journal*, 74, 852–860, 2010.
- 29 Deng, J., Zhou, Z., Zheng, X., and Li, C.: Modeling impacts of fertilization alternatives on nitrous oxide and nitric oxide  
30 emissions from conventional vegetable fields in southeastern China, *Atmospheric Environment*, 81, 642–650, 2013.

1 Diao, T., Xie, L., Guo, L., Yan, H., Lin, M., Zhang, H., Lin, J., and Lin, E.: Measurements of N<sub>2</sub>O emissions from  
2 different vegetable fields on the North China Plain, *Atmospheric Environment*, 72, 70–76, 2013.

3 FAO, IIASA, ISRIC, and ISSCAS: Harmonized World Soil Database Version 1.2, 2012. 2012.

4 Field, J. L., Keske, C. M. H., Birch, G. L., Defoort, M. W., and Cotrufo, M. F.: Distributed biochar and bioenergy  
5 coproduction: a regionally specific case study of environmental benefits and economic impacts, *Global Change  
6 Biology Bioenergy*, 5, 177–191, 2013.

7 Food and Agriculture Organization (FAO) (2015) FAOSTAT (Food and Agriculture Organization Statistical Data)  
8 Statistical Yearbook Vol. 4. Available at: <http://faostat.fao.org> (accessed 12 August 2015)

9 Harrison, R. and Webb, J.: A review of the effect of N fertilizer type on gaseous emissions, *Advances in Agronomy*, 73,  
10 65–108, 2001.

11 Hass, A., Gonzalez, J. M., Lima, I. M., Godwin, H. W., Halvorson, J. J., and Boyer, D. G.: Chicken manure biochar as  
12 liming and nutrient source for acid Appalachian soil, *Journal of Environmental Quality*, 41, 1096–1106, 2012.

13 Hu, H. W., Macdonald, C. A., Trivedi, P., Anderson, I. C., Zheng, Y., Holmes, B., Bodrossy, L., Wang, J. T., He, J. Z., and  
14 Singh, B. K.: Effects of climate warming and elevated CO<sub>2</sub> on autotrophic nitrification and nitrifiers in dryland  
15 ecosystems, *Soil Biology & Biochemistry*, 92, 1–15, 2016.

16 Hussain, M., Farooq, M., Nawaz, A., Al-Sadi, A. M., Solaiman, Z. M., Alghamdi, S. S., Ammara, U., Yong, S. O., and  
17 Siddique, K. H. M.: Biochar for crop production: potential benefits and risks, *Journal of Soils & Sediments*, 2016.  
18 1–32, 2016.

19 IPCC: Climate Change 2013: The Physical Science Basis: working group I contribution to the Fifth Assessment Report  
20 of the Intergovernmental Panel on Climate Change, Cambridge University Press, Stockholm, 2013.

21 Islam, A., Chen, D., White, R. E., and Weatherley, A. J.: Chemical decomposition and fixation of nitrite in acidic pasture  
22 soils and implications for measurement of nitrification, *Soil Biology & Biochemistry*, 40, 262–265, 2008.

23 Jia, J., Li, B., Chen, Z., Xie, Z., and Xiong, Z.: Effects of biochar application on vegetable production and emissions of  
24 N<sub>2</sub>O and CH<sub>4</sub>, *Soil Science and Plant Nutrition*, 58, 503–509, 2012.

25 Joseph, S. D., Campsarbertain, M., Lin, Y., Munroe, P., Chia, C. H., Hook, J., Van, Z. L., Kimber, S., Cowie, A., and  
26 Singh, B. P.: An investigation into the reactions of biochar in soil, *Australian Journal of Soil Research*, 48, 501–515,  
27 2010.

28 Ju, X. T., Kou, C. L., Zhang, F. S., and Christie, P.: Nitrogen balance and groundwater nitrate contamination: comparison  
29 among three intensive cropping systems on the North China Plain, *Environmental Pollution*, 143, 117–125, 2006.

30 Kim, J. Y., Song, C. H., Ghim, Y. S., Won, J. G., Yoon, S. C., Carmichael, G. R., and Woo, J. H.: An investigation on NH<sub>3</sub>

1 emissions and particulate  $\text{NH}_4^+$ - $\text{NO}_3^-$  formation in East Asia, *Atmospheric Environment*, 40, 2139–2150, 2006.

2 Kurola, J., Salkinoja-Salonen, M., Aarnio, T., Hultman, J., and Romantschuk, M.: Activity, diversity and population size  
3 of ammonia-oxidising bacteria in oil-contaminated landfarming soil, *FEMS Microbiology Letters*, 250, 33–38,  
4 2005.

5 Langridge, J. M., Lack, D., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, J. A., Nowak, J. B., Perring, A. E.,  
6 Schwarz, J. P., and Spackman, J. R.: Evolution of aerosol properties impacting visibility and direct climate forcing  
7 in an ammonia-rich urban environment, *Journal of Geophysical Research Atmospheres*, 117, 2240–2260, 2012.

8 Li, B., Bi, Z., and Xiong, Z.: Dynamic responses of nitrous oxide emission and nitrogen use efficiency to nitrogen and  
9 biochar amendment in an intensified vegetable field in southeastern China, *Global Change Biology Bioenergy*, DOI:  
10 10.1111/gcbb.12356,. 2016.

11 Li, B., Fan, C. H., Xiong, Z. Q., Li, Q. L., and Zhang, M.: The combined effects of nitrification inhibitor and biochar  
12 incorporation on yield-scaled  $\text{N}_2\text{O}$  emissions from an intensively managed vegetable field in southeastern China,  
13 *Biogeosciences*, 12, 15185–15214, 2015a.

14 Li, B., Fan, C. H., Zhang, H., Chen, Z. Z., Sun, L. Y., and Xiong, Z. Q.: Combined effects of nitrogen fertilization and  
15 biochar on the net global warming potential, greenhouse gas intensity and net ecosystem economic budget in  
16 intensive vegetable agriculture in southeastern China, *Atmospheric Environment*, 100, 10–19, 2015b.

17 Liu, X., Zhang, A., Ji, C., Joseph, S., Bian, R., Li, L., Pan, G., and Paz-Ferreiro, J.: Biochar's effect on crop productivity  
18 and the dependence on experimental conditions—a meta-analysis of literature data, *Plant and Soil*, 373, 583–594,  
19 2013.

20 Lu, R.: *Methods of soil and agro-chemical analysis*, China Agricultural Science and Technology Press, Beijing, 2000.  
21 127–332, 2000. (in Chinese)

22 Ma, L., Shan, J., Yan, X., 2015. Nitrite behavior accounts for the nitrous oxide peaks following fertilization in a  
23 fluvo-aquic soil. *Biology and Fertility of Soils* 51, 563-572.

24 Major, J., Lehmann, J., Rondon, M., and Goodale, C.: Fate of soil-applied black carbon: downward migration, leaching  
25 and soil respiration, *Global Change Biology*, 16, 1366–1379, 2010.

26 Mei, B. L., Zheng, X. H., Xie, B. H., Dong, H. B., Zhou, Z. X., Rui, W., Jia, D., Feng, C., Tong, H. J., and Zhu, J. G.:  
27 Nitric oxide emissions from conventional vegetable fields in southeastern China, *Atmospheric Environment*, 43,  
28 2762–2769, 2009.

29 Misselbrook, T. H., Weerden, T. J. V. D., Pain, B. F., Jarvis, S. C., Chambers, B. J., Smith, K. A., Phillips, V. R., and  
30 Demmers, T. G. M.: Ammonia emission factors for UK agriculture, *Atmospheric Environment*, 34, 871–880(810),

1 2000.

2 Mukherjee, A. and Zimmerman, A. R.: Organic carbon and nutrient release from a range of laboratory-produced biochars  
3 and biochar–soil mixtures, *Geoderma*, s 193–194, 122–130, 2013.

4 Nelissen, V.: Effect of different biochar and fertilizer types on N<sub>2</sub>O and NO emissions, *Soil Biology & Biochemistry*, 70,  
5 244–255, 2014.

6 Novak, J. M., Spokas, K. A., Cantrell, K. B., Ro, K. S., Watts, D. W., Glaz, B., Busscher, W. J., and Hunt, P. G.: Effects  
7 of biochars and hydrochars produced from lignocellulosic and animal manure on fertility of a Mollisol and Entisol,  
8 *Soil Use and Management*, 30, 175–181, 2014.

9 Obia, A., Cornelissen, G., Mulder, J., and Dörsch, P.: Effect of Soil pH Increase by Biochar on NO, N<sub>2</sub>O and N<sub>2</sub>  
10 Production during Denitrification in Acid Soils, *Plos One*, 10, 359–367, 2015.

11 Pacholski, A., Cai, G. X., Fan, X. H., Ding, H., Chen, D., Nieder, R., and Roelcke, M.: Comparison of different methods  
12 for the measurement of ammonia volatilization after urea application in Henan Province, China, *Journal of Plant  
13 Nutrition and Soil Science*, 171, 361–369, 2008.

14 Pinder, R. W., Adams, P. J., and Pandis, S. N.: Ammonia emission controls as a cost-effective strategy for reducing  
15 atmospheric particulate matter in the Eastern United States, *Environmental Science & Technology*, 41, 380–386,  
16 2007.

17 Powlson, D. S., Addiscott, T. M., Benjamin, N., Cassman, K. G., de Kok, T. M., Van, G. H., L'Hirondel, J. L., Avery, A.  
18 A., and Van, K. C.: When does nitrate become a risk for humans?, *Journal of Environmental Quality*, 37, 291–295,  
19 2008.

20 Rajkovich, S., Enders, A., Hanley, K., Hyland, C., Zimmerman, A. R., and Lehmann, J.: Corn growth and nitrogen  
21 nutrition after additions of biochars with varying properties to a temperate soil, *Biology & Fertility of Soils*, 48,  
22 271–284, 2012.

23 Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N<sub>2</sub>O): the dominant ozone-depleting substance  
24 emitted in the 21st century, *Science*, 326, 123–125, 2009.

25 Saarnio, S., Heimonen, K., and Kettunen, R.: Biochar addition indirectly affects N<sub>2</sub>O emissions via soil moisture and  
26 plant N uptake, *Soil Biology & Biochemistry*, 58, 99–106, 2013.

27 Schouten, S., Groenigen, J. W. V., Oenema, O., and Cayuela, M. L.: Bioenergy from cattle manure? Implications of  
28 anaerobic digestion and subsequent pyrolysis for carbon and nitrogen dynamics in soil, *Global Change Biology  
29 Bioenergy*, 4, 751–760, 2012.

30 Schulz, H., Dunst, G., and Glaser, B.: Positive effects of composted biochar on plant growth and soil fertility, *Agronomy*

1 for Sustainable Development, 33, 817–827, 2013.

2 Setia, R., Marschner, P., Baldock, J., Chittleborough, D., and Verma, V.: Relationships between carbon dioxide emission  
3 and soil properties in salt-affected landscapes, *Soil Biology & Biochemistry*, 43, 667–674, 2011.

4 Shannon, M. C. and Grieve, C. M.: Tolerance of vegetable crops to salinity, *Scientia Horticulturae*, 78, 5–38, 1998.

5 Smith, M. S. and Tiedje, J. M.: Phases of denitrification following oxygen depletion in soil, *Soil Biology & Biochemistry*,  
6 11, 261–267, 1979.

7 Smith, P.: Soil carbon sequestration and biochar as negative emission technologies, *Global Change Biology*, 51, 574–575,  
8 2016.

9 Sohi, S. P.: Agriculture. Carbon storage with benefits, *Science*, 338, 1034–1035, 2012.

10 Sororzano, L.: Determination of ammonia in natural waters by the phenolhypochlorite method, *Limnol. Oceanogr*, 14,  
11 799–801, 1969.

12 Spokas, K. A. and Reicosky, D. C.: Impacts of sixteen different biochars on soil greenhouse gas production, *Ann.*  
13 *Environ. Sci*, 3, 4, 2009.

14 Stavi, I. and Lal, R.: Agroforestry and biochar to offset climate change: a review, *Agronomy for Sustainable*  
15 *Development*, 33, 81–96, 2013.

16 Sun, L., Li, L., Chen, Z., Wang, J., and Xiong, Z.: Combined effects of nitrogen deposition and biochar application on  
17 emissions of N<sub>2</sub>O, CO<sub>2</sub> and NH<sub>3</sub> from agricultural and forest soils, *Soil science and plant nutrition*, 60, 254–265,  
18 2014.

19 Ussiri, D. and Lal, R.: *The Role of Nitrous Oxide on Climate Change*, Springer Netherlands, 2013.

20 Wang, J., Chen, Z., Ma, Y., Sun, L., Xiong, Z., Huang, Q., and Sheng, Q.: Methane and nitrous oxide emissions as  
21 affected by organic–inorganic mixed fertilizer from a rice paddy in southeast China, *Journal of Soils and Sediments*,  
22 13, 1408–1417, 2013.

23 Wang, J., Chen, Z., Xiong, Z., Chen, C., Xu, X., Zhou, Q., and Kuzyakov, Y.: Effects of biochar amendment on  
24 greenhouse gas emissions, net ecosystem carbon budget and properties of an acidic soil under intensive vegetable  
25 production, *Soil Use and Management*, 31, 375–383, 2015a.

26 Wang, S., Nan, J., Shi, C., Fu, Q., Gao, S., Wang, D., Cui, H., Saizlopez, A., and Zhou, B.: Atmospheric ammonia and its  
27 impacts on regional air quality over the megacity of Shanghai, China, *Scientific Reports*, 5, 2015b.

28 Wrage, N., Velthof, G., Van Beusichem, M., Oenema, O., 2001. Role of nitrifier denitrification in the production of  
29 nitrous oxide. *Soil Biology and Biochemistry* 33, 1723-1732.

30 Xiong, Z., Xie, Y., Xing, G., Zhu, Z., and Butenhoff, C.: Measurements of nitrous oxide emissions from vegetable

1 production in China, *Atmospheric Environment*, 40, 2225–2234, 2006.

2 Yao, Z., Zheng, X., Xie, B., Mei, B., Wang, R., Klaus, B. B., Zhu, J., and Yin, R.: Tillage and crop residue management  
3 significantly affects N-trace gas emissions during the non-rice season of a subtropical rice-wheat rotation, *Soil  
4 Biology & Biochemistry*, 41, 2131–2140, 2009.

5 Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J., and Crowley, D.: Effect of biochar amendment on  
6 yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China, *Agriculture  
7 Ecosystems & Environment*, 139, 469–475, 2010.

8 Zhang, F., Chen, X., and Vitousek, P.: Chinese agriculture: An experiment for the world, *Nature*, 497, 33-35, 2013.

9 Zhang, M., Fan, C. H., Li, Q. L., Li, B., Zhu, Y. Y., and Xiong, Z. Q.: A 2-yr field assessment of the effects of chemical  
10 and biological nitrification inhibitors on nitrous oxide emissions and nitrogen use efficiency in an intensively  
11 managed vegetable cropping system, *Agriculture Ecosystems & Environment*, 201, 43–50, 2015.

12 Zhang, Y., Luan, S., Chen, L., and Shao, M.: Estimating the volatilization of ammonia from synthetic nitrogenous  
13 fertilizers used in China, *Journal of Environmental Management*, 92, 480–493, 2011.

14 Zhao, L. M., Wu, L. H., Dong, C. J., and Li, Y. S.: Rice yield, nitrogen utilization and ammonia volatilization as  
15 influenced by modified rice cultivation at varying nitrogen rates, *Agricultural Sciences*, 01, 10–16, 2010.

16 Zhao, X., Wang, J., Wang, S., and Xing, G.: Successive straw biochar application as a strategy to sequester carbon and  
17 improve fertility: A pot experiment with two rice/wheat rotations in paddy soil, *Plant and Soil*, 378, 279–294, 2014.

18 Zheng, X., Mei, B., Wang, Y., Xie, B., Wang, Y., Dong, H., Xu, H., Chen, G., Cai, Z., and Yue, J.: Quantification of N<sub>2</sub>O  
19 fluxes from soil–plant systems may be biased by the applied gas chromatograph methodology, *Plant and Soil*, 311,  
20 211–234, 2008.

21 Zhou, J. B., Chen, Z. J., Liu, X. J., Zhai, B. N., and Powlson, D. S.: Nitrate accumulation in soil profiles under  
22 seasonally open ‘sunlight greenhouses’ in northwest China and potential for leaching loss during summer  
23 fallow, *Soil Use and Management*, 26, 332–339, 2010.

24 Zhu, T., Zhang, J., Cai, Z., 2011. The contribution of nitrogen transformation processes to total N<sub>2</sub>O emissions from  
25 soils used for intensive vegetable cultivation. *Plant and Soil* 343, 313-327.

1 **Table legends**

2 **Table 1**

3 Soil organic carbon (SOC), soil total nitrogen (TN), soil pH, electric conductivity (EC) and microbial biomass carbon  
 4 (MBC) as affected by different treatments across the four vegetable soils.

Soil	Treatment	SOC (g kg <sup>-1</sup> )	TN (g kg <sup>-1</sup> )	pH	EC (ds m <sup>-1</sup> )	MBC (mg kg <sup>-1</sup> )
Acrisol	N	8.0±0.8c	1.37±0.12b	4.37±0.04c	1.76±0.21b	1353±119a
	N+Bw	15.6±0.5b	1.47±0.07b	4.64±0.04b	2.43±0.31a	1173±49b
	N+Bm	18.8±0.6a	1.64±0.04a	5.01±0.03a	2.00±0.32ab	1234±50ab
Anthrosol	N	9.7±0.7c	1.55±0.04b	7.53±0.02b	1.74±0.27b	490±9a
	N+Bw	15.6±0.8b	1.62±0.06b	7.61±0.05a	2.25±0.22a	495±16a
	N+Bm	17.5±1.1a	1.79±0.03a	7.63±0.01a	1.96±0.06ab	504±18a
Cambisol	N	7.9±0.1b	1.13±0.04b	7.70±0.08a	0.85±0.03b	535±13b
	N+Bw	14.2±0.6a	1.20±0.04b	7.66±0.03a	0.92±0.04a	554±10ab
	N+Bm	15.5±1.4a	1.37±0.06a	7.71±0.03a	0.87±0.02ab	573±12a
Phaeozem	N	29.9±0.5b	2.19±0.04b	6.91±0.05a	0.83±0.03b	921±44b
	N+Bw	36.0±1.5a	2.20±0.03b	6.92±0.06a	0.95±0.03a	988±56b
	N+Bm	38.1±1.8a	2.41±0.01a	6.94±0.04a	0.92±0.06a	1242±196a
ANOVA results						
Biochar		***	***	***	***	*
Soil		***	***	***	***	***
Biochar×Soil		*	n.s.	***	n.s.	**

5 Data shown are means ± standard deviations of three replicates. See Fig. 1 for treatments codes. Different letters within  
 6 the same column indicate significant differences among treatments within the same soil at  $p < 0.05$  level.

7 \*\*\*Significant at  $p < 0.001$ ; \*\*significant at  $p < 0.01$ ; \*significant at  $p < 0.05$ ; n.s. not significant.



1 **Table 2**

2 Two-way ANOVA for the effects of biochar (Bc) and soil (S) types on cumulative N<sub>2</sub>O, NO and NH<sub>3</sub> emissions, gaseous reactive nitrogen emission (GNrE), vegetable yield  
3 and gaseous reactive nitrogen intensity (GNrI) during the entire sampling period.

Factors	DF	N <sub>2</sub> O emission			NO emission			NH <sub>3</sub> emission			GNrE			Vegetable yield			GNrI		
		SS	F	P	SS	F	P	SS	F	P	SS	F	P	SS	F	P	SS	F	P
Bc	2	271.9	65.1	***	46.4	174.7	***	0.5	0.8	n.s.	380.5	86.4	***	76.2	3.2	n.s.	0.1	7.9	**
S	3	1429.9	228.1	***	152.2	382.1	***	4.1	3.8	*	2322.6	351.5	***	4316.9	123.3	***	2.3	110.3	***
Bc×S	6	179.3	14.3	***	33.4	41.9	***	1.4	0.7	n.s.	234.5	17.7	***	230.4	3.3	*	0.1	1.6	n.s.
Model	11	4009.7	174.5	***	225.3	154.3	***	29.1	7.5	***	5290	218.3	***	15962.0	124.4	***	5.8	77.0	***
Error	24	50.1			3.2			8.5			52.9			280.0			0.2		

4 SS: the sum of squares.

5 F value: the ratio of mean squares of two independents samples.

6 P value: the index of differences between the control group and the experimental group. \*, \*\* and \*\*\* indicate significance at  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively.

7 n.s.: not significant.

1 **Table 3**

2 Cumulative gaseous nitrogen (N<sub>2</sub>O, NO and NH<sub>3</sub>) emissions, gaseous reactive nitrogen emission (GNrE), vegetable yield  
 3 and gaseous reactive nitrogen intensity (GNrI) under the different treatments across the four soils.

Treatments	Acrisol	Anthrosol	Cambisol	Phaeozem
(a) Cumulative N <sub>2</sub> O emissions (kg N ha <sup>-1</sup> )				
N	30.59±3.15aA	7.83±0.60aB	2.52±0.37aC	7.10±1.91aB
N+Bw	19.45±2.43bA	3.20±0.28bB	1.97±0.21aB	3.45±0.86bB
N+Bm	31.56±1.35aA	3.63±0.62bB	2.26±0.58aB	4.01±0.68bB
(b) Cumulative NO emissions (kg N ha <sup>-1</sup> )				
N	8.99±1.01aA	1.27±0.15aB	0.20±0.08aC	0.97±0.11aBC
N+Bw	4.54±0.60bA	0.80±0.13bB	0.33±0.19aB	0.52±0.03bB
N+Bm	3.87±0.30bA	1.16±0.17aB	0.21±0.10aC	0.94±0.03aB
(c) Cumulative NH <sub>3</sub> emissions (kg N ha <sup>-1</sup> )				
N	4.72±0.27aB	5.79±0.54bA	6.34±0.51aA	5.67±0.42aA
N+Bw	5.09±0.38aB	6.83±0.74abA	7.35±0.75aA	6.24±0.49aAB
N+Bm	5.32±0.42aB	7.57±0.57aA	7.37±1.11aA	6.48±0.43aAB
(d) GNrE (kg N ha <sup>-1</sup> )				
N	44.30±3.13aA	14.89±1.33aB	9.06±0.80aC	13.74±1.67aB
N+Bw	29.08±2.21bA	10.82±1.14bB	9.64±0.88aB	10.21±0.92bB
N+Bm	40.76±1.66aA	12.36±0.74bB	9.84±0.49aC	11.42±0.27bBC
(e) Vegetable yield (t ha <sup>-1</sup> )				
N	35.20±2.52aB	25.29±3.90aC	39.09±2.03bB	75.65±5.84bA
N+Bw	29.05±2.35bC	23.57±1.74aC	44.53±3.74bB	76.95±4.04abA
N+Bm	34.93±2.87aC	26.30±2.63aD	51.00±3.18aB	85.89±3.29aA
(f) GNrI (kg N t <sup>-1</sup> yield)				
N	1.27±0.18aA	0.59±0.08aB	0.23±0.02aC	0.18±0.04aC
N+Bw	1.01±0.12aA	0.46±0.05bB	0.22±0.04aC	0.13±0.02bC
N+Bm	1.17±0.15aA	0.47±0.04bB	0.19±0.01aC	0.13±0.01bC

4 Data shown are means ± standard deviations of the three replicates. See Fig. 1 for treatments codes. Different lowercase  
 5 letters within the same column indicate significant differences among treatments within the same soil at  $p < 0.05$  level.  
 6 Different capital letters within the same row indicate significant differences among soil types within the same treatment  
 7 at  $p < 0.05$  level.

1 **Table 4**

2 The correlations between N<sub>2</sub>O or NO emission and PNR or DEA in each soil.

Item	Acrisol		Anthrosol		Cambisol		Phaeozem	
	PNR	DEA	PNR	DEA	PNR	DEA	PNR	DEA
N <sub>2</sub> O	0.75*	0.66	0.49	0.76*	-0.10	0.16	-0.82**	0.70*
NO	0.62	-0.29	0.79*	0.69*	-0.54	0.01	-0.63	0.22

3 Asterisks indicated 0.05 level significances ( $*p < 0.05$ ) and 0.01 level significances ( $**p < 0.01$ ), n = 9.


1 **Figure legends**

2 **Fig. 1** Potential nitrification rate (PNR) and Denitrification enzyme activity (DEA) under different treatments in Acrisol,  
3 Anthrosol, Cambisol and Phaeozem. The three treatments with each soil were urea without biochar (N), urea with wheat  
4 straw biochar (N+Bw) and urea with swine manure biochar (N+Bm). Bars indicate standard deviation (mean + SD, n =  
5 3). Different letters above the bars indicate significant differences among the different treatments within the same soil, at  
6  $p < 0.05$ .


7 **Fig. 2** Temporal dynamics of soil  $\text{N}_2\text{O}$  ( $\mu\text{g N m}^{-2} \text{h}^{-1} \pm \text{SD}$ , n = 3) fluxes under different treatments in Acrisol (a),  
8 Anthrosol (b), Cambisol (c) and Phaeozem (d) with five consecutive vegetable crops. The inserted panels describe the  
9  $\text{N}_2\text{O}$  fluxes during the last two cropping seasons. The solid arrows indicate fertilization. See Fig. 1 for treatments codes.

10 **Fig. 3** Temporal dynamics of soil NO ( $\mu\text{g N m}^{-2} \text{h}^{-1} \pm \text{SD}$ , n = 3) fluxes under different treatments in Acrisol (a),  
11 Anthrosol (b), Cambisol (c) and Phaeozem (d) with five consecutive vegetable crops. The solid arrows indicate  
12 fertilization. See Fig. 1 for treatments codes.


13 **Fig. 4** Cumulative ammonia ( $\text{NH}_3$ ) emissions from the Acrisol (a), Anthrosol (b), Cambisol (c) and Phaeozem (d) during  
14 the four nitrogen fertilization events F: every N fertilization event. The bars indicate the standard deviation of the mean  
15 ( $\text{kg N ha}^{-1} \pm \text{SD}$ , n = 3) of each treatment for the sum of the four N fertilization events. See Fig. 1 for treatments codes.  
16 Different letters above the bars indicate significant differences among the different treatments for each soil, at  $p <$   
17 0.05.




1



1



1



1