

This file contains:

- 1. A point-by-point response to the editor and the referee, including a list of relevant changes**
- 2. A marked up manuscript version, thus, the manuscript including “track changes” from the present revision.**

Response to editor and referee

2016-09-10

First we would like to thank the editor Prof. Fennel and the anonymous referee for their helpful and thoughtful comments. In our response we repeat the comments/questions followed by our response/changes. We refer to page + line numbers in the now revised manuscript, if suited, in which “track changes” are included.

AR-Author reply

Editors comments:

Associate Editor Decision: Reconsider after major revisions (30 Jul 2016) by Prof. Katja Fennel

Comments to the Author:

Dear Authors,

Reviewer 1 provided a further detailed review of your revised manuscript. I encourage you to carefully consider these comments.

In addition I'm providing my own comments below. I encourage you to submit a revised manuscript for further consideration as well as detailed responses to each comment.

Page and line numbers refer to the revised manuscript with changes tracked.

Define what you mean by “coastal filter” (P1, L18; P1, L25 and L26). The definition provided on page 4, lines 6-8 is insufficient. Where do these nutrients ultimately go? Which processes are responsible?

AR: This is now shortly better defined already in the abstract (P1, L19-21; P2, L2-5), with a more thorough definition in the introduction (P3, L17-21). It is clarified that retention in this manuscript also includes removal of nutrients (P3-4, L27-5), since the point of view is how much of the nutrient load from land does not reach the open sea.

P1, L24: “good quality” Do you mean “the simulated values agree well with observations”? Unclear otherwise. Please rephrase.

AR: Yes, this is what we mean and it is now rephrased (P1,L25-27).

P1, L26: “take care of” is not a good word choice. Replace by “remove” or “retain” or similar.

AR: We agree and have changed it (P2, L1).

Also, please explain where all the nutrients ultimately go. The Archipelago cannot infinitely continue to retain nutrients.

AR: This is now more clear in the abstract (P2, L2-5) and better described in the introduction (mentioned above (P3-4, L27-5)). Fig. 4 in the previous manuscript schematically shows the coastal filter and the retention and is therefore now moved to the introduction section and re-numbered to Fig. 1. Consequently the following figures up to fig. 3 changes number as well.

P2, L1-7: Denitrification does not “retain” nutrients. Denitrification removes nutrients from the system! This is different from retention.

AR: We agree with the Prof. Fennel that it is a simplification to include the denitrification process in the retention definition. But, since one of the “simpler” methods to calculate retention is to calculate the difference between the load of nutrients and the outflow (described in section 2.4), which is used in a number different studies, the comparable numbers of the retention includes denitrification. From the point of view that the filter capacity is defined as the difference from what comes in to the area and what comes out from the area, then the denitrification process is included in the retention. We also consider e.g. burial to be a removal of nutrients since it is a sink for the model. This is now better explained in the manuscript where the definition and explanation is described in the introduction and further described also in the method section (section 2.4).

P4, L8: “There are not enough estimates of nutrient retention in different coastal systems around the world...”

I note that you are not providing references of those that exist – something that was also criticized by Reviewer 1. Here are just three examples that I’m familiar with (I’m sure there are many more):

Xue, Z., He, R., Fennel, K., Cai, W.-J., Lohrenz, S., and Hopkinson, C., Modeling ocean circulation and biogeochemical variability in the Gulf of Mexico, *Biogeosciences*, 10, 7219-7234, doi:10.5194/bg-10-7219-2013 (2013)

Fennel, K., The role of continental shelves in nitrogen and carbon cycling: Northwestern North Atlantic case study. *Ocean Science* 6, 539-548, doi:10.5194/os-6-539-2010 (2010)

Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., Haidvogel, D., Nitrogen cycling in the Mid Atlantic Bight and implications for the North Atlantic nitrogen budget: Results from a three-dimensional model. *Global Biogeochemical Cycles* 20, GB3007, doi:10.1029/2005GB002456. (2006)

Two other studies of some potential relevance as well as the references given within are:

Laurent, A., Fennel, K., Hu, J., and Hetland, R.: Simulating the effects of phosphorus limitation in the Mississippi and Atchafalaya River plumes, *Biogeosciences*, 9, 4707-4723, doi:10.5194/bg-9-4707-2012 (2012)

Laurent, A., Fennel, K., Simulated reduction of hypoxia in the northern Gulf of Mexico due to phosphorus limitation, *Elementa* 2:000022, doi:10.12952/journal.elementa.000022 (2014)

Please make an effort to properly place your findings in the context of the published literature.

AR: First we would like to thank Prof. Fennel for the literature suggestions. We have read them all and considered their interesting results. We have included results from some of the studies where both the river input and the removal of nutrients (retention) are given, section 3.2 (P18, L25-31) as well as given examples of model studies in the introduction (P4, L24-25). We have also changed the formulation of the statement (P4, L17-18). New references included in the revised manuscript:

Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., Haidvogel, D., Nitrogen cycling in the Mid Atlantic Bight and implications for the North Atlantic nitrogen budget: Results from a three-dimensional model. *Global Biogeochemical Cycles* 20, GB3007, doi:10.1029/2005GB002456. (2006)

Seitzinger, S. P. and Giblin, A. E.: Estimating denitrification in North Atlantic continental shelf sediments, *Biogeochemistry*, 35, 235-260, 1996.

Xue, Z., He, R., Fennel, K., Cai, W.-J., Lohrenz, S., and Hopkinson, C., Modeling ocean circulation and biogeochemical variability in the Gulf of Mexico, *Biogeosciences*, 10, 7219-7234, doi:10.5194/bg-10-7219-2013 (2013)

P11, L13: Please see my earlier comment about the difference between removal and retention. Please be very clear what you mean. At present these two terms are lumped together which is not appropriate. Always be clear about the processes that are underlying your results. Just quantifying

the difference between incoming and outgoing matter fluxes without explaining where the missing matter ended up is not sufficient.

AR: We do not totally agree. The retention has in many other studies been calculated as the difference between the nutrient load and the outflow e.g. Karlson (2010), Johnston (1991), Billen et al. (2011), and some other studies that we refer to in the manuscript. In a lot of areas the different processes are not known and a simpler method is the only choice. In our study we use both the simpler method, but we also calculate separately the burial and the denitrification processes. In this way we can compare the total “retention” with other studies. Also in Nixon et al. (1996), see e.g. fig. 3, the difference between the input and the outflow is defined as “retention”, and a part of that retained nitrogen is denitrified. From the point of view that fractions of the nutrients supply from land are prevented from being further transported to the open sea, then it is “retained” in the area and for nitrogen that includes both burial and denitrification in our model study. As mentioned above, this is now more described in the introduction (P3-4, L25-13) as well as in the section 2.4 (P11, 11-27).

P11, L16: Do not relegate important definitions to the supplement. Define retention in the main text.

AR: Yes, we agree. Because the used retention definitions are already described in the main text the supplementary 2 is not necessary to include. We remove it from the submission.

P12: I question your calculation of “freshwater” residence time. As per the Mosen reference you provide, residence time is defined as the ratio of an inventory and its inflow or outflow rate. You talk about freshwater residence time and divide the volume by the freshwater inflow rate, but the volume is not the freshwater volume but the total volume. Perhaps you do not mean freshwater residence time? Please clarify.

AR: We agree. We do mean the fresh water residence time and have now calculated it as the fresh water volume (given by a freshwater tracer in the model) divided by the freshwater inflow (Q_f). The description is revised in section 2.4 (P12, L14-23). This change also led to an updated Fig. 13 since the lower residence time affected the positions of our data points in the graph as well as the text in section 3.2.2 (P19-20, L25-1).

P12, L9-22: This text and Table 2 are not very informative. It is not clear what the baseline load is. Without this it is not particularly useful to state a percentage reduction. Consider giving less detail about treatment plants and instead describe better what nutrient inputs were used in the different experiments.

AR: We agree. The baseline load (forcing from 2010, the annual loads of N and P) are now given in the revised text in section 2.4.2 (P13). The level of details has been reduced. Also table 2 has been reduced since some of the numbers are now given in the text.

P13, L4-6: “A clear relationship between river outflow and nutrient load is observed ...”

This is not a result that can be observed, but something that is imposed!!! Does not belong in the results but in the Methods.

AR: This sentence is now re-formulated (P14, L4-6). We hope the re-formulation makes this clearer.

Fig. 9: Please separate input from land and atmosphere instead of providing one combined number.

AR: Done.

The concept of temporary retention does not make much sense to me. Do you consider your system to be in a dynamic steady state? How do you distinguish between processes leading to temporary versus permanent retention? Do the N and P inventories in the water column differ between the beginning and end of the simulation period? Please clarify this in section 3.2.

AR: Permanent retention for P means burial. For N benthic denitrification represented as much as almost 92 % of the permanent retention, burial for less than 8 % and pelagic denitrification was below 1 %. This is written on P17, L7-10.

The temporary retention on the other hand is considered to be changes in the pelagic and benthic pools of nutrients. The nutrient pools include both the inorganic and organic nutrients. Factors that affect the benthic N and P pool are supply of organic material from the water column settling on the sediment, the decomposition of organic material and the release of inorganic nutrients as well as by burial

(which in the model is a permanent sink of nutrients). The pelagic N and P pools are affected by the supply from land, the sinking of organic material, the release of nutrients from the sediment to the water column and to the net export of nutrients to areas down-stream. These processes are handled by the model and what we present in the paper is the total temporary retention. This is now written in section 2.4 (P11, L20-27).

The benthic and the pelagic pools can thus change during the simulation period, which is described in section 3.2 (P16,L12-14; P18, L14-18) and can be observed in fig. 12.

Katja Fennel
Editor

Comments from reviewer 1:

P6L5: "transparency"

AR: Done (P6, L18).

P12L9: section title suggestion: Nutrient load reduction scenario

AR: Done.

P12L21 (and everywhere else in the manuscript, there are many occurrences): change "with" to "by"

AR: Done.

P12L22: replace "to" by "in"

AR: This paragraph is re-written and the comment is therefore not valid any more.

P16L1-3: change to something like "Permanent retention was relatively stable during the simulated period, while fluctuations in the temporal retention reflects the effect of varying riverine nutrient input (Fig. 10c,d)."

AR: Done (P17, L4-5).

Also, temporal retention of P has increased, Can you comment on that?

AR: Yes, this is now mentioned in the section 3.2 (P18, 17-20).

P17L11-16: This new part could be polish for clarity.

AR: This section is now revised.

P17L14: change "has" to "have"

AR: Done.

P18L23: change "parameters" to "processes"

AR: The whole paragraph is revised, section 3.2.2 (P19).

P18L26: change "relation" to "relationship"

AR: Done (P19, L27).

P18L27-28: replace to "... the filter efficiency. Nixon et al (1996) show that including the depth in the analysis of retention"

AR: Done, P1928-30.

P19L1-31: This added text could be improved.

P19L1-4: suggested replacement for the first 2 sentences: "In the model, denitrification is an O2 dependent process that has a maximum rate at O2 concentration...".

AR: Done, but parts of this section is also moved to the new method section 2.4.1 (P12-13).

P19L5-9: Are you talking about the model here or in general? This is not described in the Methods so we don't know if this relates to your results or not.

AR: It is mentioned in the method section 2.2.2 that the release of phosphate is redox dependent (P8, L18-20) and it is now also explained in the section 2.4.1 (P12-13).

P19L9-12: What do you mean by manually? This is vague, you need to provide details on what you did. It could be mentioned in the Methods.

AR: This is now rewritten and described in the new section 2.4.1 in the revised manuscript (P12-13).

P19L17: Is this due only to denitrification? what about higher primary production due to more P available? Did you compare the different effects and found that denitrification is the dominant process? You mention this later in the load reduction experiment.?

AR: The permanent retention of N increased and the proportion of the N that was denitrified also increased. The permanent retention, burial, of P decreased. Thus there was a change in the N:P ratio. However, the temporary retention of N did not change significantly, but there was a decreased export of N to the open sea, while the P export increased compared to the original run. This paragraph is reformulated (P20, L14-30)

P19L17-25: Suggested changes for clarity: "Denitrification increased in magnitude and as a proportion of the permanent retention from 92% to 94%, while the proportion of burial decreased. The decrease in P retention was due to higher release of P from the sediment. Denitrification has been shown to increase in areas with longer residence times (Nixon et al., 1996, Nielsen et al., 2001, Finlay et al., 2013). In the simulation the residence time was only six days on average and the filter efficiencies of N and P were estimated to be 10% and 9%, respectively, which were lower than in the Stockholm Archipelago..."

AR: Thank you for the suggested changes, however, large parts of this paragraph is re-formulated (P20, L14-30)

Also, I don't see why the results L23-25 are included here. Can you be more explicit about the comparison with your system?

AR: Denitrification is one of the removal processes, permanent retention, which is discussed here. The magnitude of the denitrification has been shown to increase with areas having longer residence time compared to shorter. The result in the study in the Randers fjord is here compared to the Stockholm archipelago where we have both longer residence time and more retention due to denitrification. This is reformulated in the revised manuscript (P20, L14-19).

P19L29-31: This is true everywhere. Please clarify your statement.

AR: This sentence is now rewritten (P20, L20-22).

P19-20L33-4: Algae uptake changes the pool of nutrients, please rephrase to something like this: Assimilation of nutrients by benthic algae does not directly change the inventory of N and P but transfers inorganic N and P into organic material..."

AR: This sentence is now rewritten (P20-21, L32-1).

What is the expected effect on retention? You need to relate this to your results, this is part of the uncertainty of your results.

AR: A discussion about this is now added (P21, L3-12).

P20L13: suggested change to title: "Response to nutrient load reduction."

AR: Done (P21, L21).

P21L11: suggested change: "is" instead of "becomes"

AR: Done (P22, L19).

P21L14: Change "From these experiments it seems that" to "These results indicate that".

AR: Done (P22, L22).

P21L17-18: suggested change: "... concentrations might be expected locally. However, this effect largely depends on the water residence time and on which nutrient limits the seasonal phytoplankton production initially..."

AR: Done (P22, L24-26).

P21L20-29: I don't see the use of this paragraph.

AR: We agree, and large part of the paragraph is now removed.

P21L22-25: not sure what that means, please clarify.

AR: These sentences are now rewritten (P22, L28-30).

P23L6: replace "to" with "in".

AR: Done (P24, L6).

P23L8-10: is that permanent or temporary? Can you indicate their ratio?

AR: This retention is the total, thus a sum of permanent and temporary retention (defined in section 2.4), which now is clarified in the conclusions (P24, L9). The temporary retention in this case is negative, thus we have a decrease in the inventory of N and P. The permanent retention is larger about 98 %. In the conclusions we would prefer to give the total number of retention, however, the different values are given in fig. 9 from which the different ratios can be calculated.

P25L1: journal names should be abbreviated

AR: Done (P26-32).

P25L16-18: add database title: "Baltic Environmental Database" and use the short url provided: <http://www.balticnest.org/bed>

AR: Done (P26, L16).

P44: is there a reason to not start the x-axis at 0? It seems odd.

AR: No, there is no reason and we also took the opportunity to remake the figure. In the new figure (fig. 14) we think it is easier to compare the filter efficiencies of N and P.

Additional changes by the authors:

“- a model study” has been removed from the title, since the word “Modelling” should be enough.

The keyword “nutrient retention “ is already in the title and is therefore changed to “denitrification” .

Modelling nutrient retention in the coastal zone of an eutrophic sea

Borttaget: - a model study

Elin Almroth-Rosell^{*1}, Moa Edman¹, Kari Eilola¹, H.E. Markus Meier^{2,1}, Jörgen Sahlberg¹

¹Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

²Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany

*Corresponding author:

Elin Almroth-Rosell

Swedish Meteorological and Hydrological Institute

Sven Källfelts gata 15, SE-426 71 Västra Frölunda, Sweden.

Tel: +46(0)31 7518969. E-mail: elin.almroth.rosell@smhi.se

Keywords: Denitrification, phosphate, nitrogen, eutrophication, biogeochemistry, coastal zone, Baltic Sea

Borttaget: Nutrient retention

Abstract

The Swedish Coastal zone Model (SCM) was used at a test site, the Stockholm Archipelago located in the northern part of the central Baltic Sea, to study the retention capacity of the coastal filter on nitrogen (N) and phosphorus (P) loads from land and atmosphere. The efficiency of the coastal filter to permanently retain nutrients determines how much of the local nutrient loads actually reach the open sea. The SCM system is a NPDZ-type model coupled to a horizontally integrated, physical model in particular suitable for estuaries. In this study the Stockholm Archipelago consisting of 86 sub-basins was divided into three sub-areas: the inner, the intermediate and the outer archipelago. An evaluation of model results showed that the modelled freshwater supply agrees well with observations. The nutrient, salinity and temperature dynamics simulated by the SCM model are also found to be in good or acceptable agreement with observations. The analysis showed that the Stockholm Archipelago works as a filter for nutrients that enter the coastal zone from land, but the filter

Borttaget: with observations

Borttaget: .and tT

Borttaget: in

Borttaget: are of good quality

Borttaget: with observations

Borttaget: Further, tT

1 efficiency is not effective enough to retain all the supplied nutrients. However, at least 65 %
2 and 72 % of the P and N, respectively, are retained during the studied period (1990-2012). A
3 major part of the retention is permanent, which for P means burial. For N almost 92 % of the
4 permanent retention is represented by benthic denitrification, less than 8 % by burial, while
5 pelagic denitrification is below 1 %. Highest total amounts of P and N are retained in the
6 outer archipelago where the surface area is largest. The area specific retention of P and N,
7 however, is highest in the smaller inner archipelago and decreases towards the open sea. A
8 reduction scenario of the land loads of N and P showed that the filter efficiencies of N and P
9 increase and the export of N from the archipelago decreases. About 15 years after the
10 reduction the export of P changes into an import of P from the open sea to the archipelago.

Borttaget: take care of

1 Introduction

2 The worldwide increase of coastal eutrophication and anoxia has spread exponentially since
3 the 1960s. Coastal oxygen depletion is associated with dense population areas and large river
4 loads of nutrients (Diaz and Rosenberg, 2008). The use of industrially produced fertilizer
5 started in the late 1940s and has since then been contributing to the anthropogenic fertilization
6 of the global marine system (Galloway et al., 2008). The river load of nutrients originating
7 from agriculture activities has been shown to be controlled by the size of the river flow, e.g.
8 the flow from the Mississippi River has a large impact on the oxygen conditions in the
9 northern Gulf of Mexico, which suffers from severe hypoxia with “dead zones” as a result
10 (Rabalais et al., 2002).

11 With ambition to diminish eutrophication there has been a lot of efforts around the world to
12 reduce the land load of nutrients to sea, but the expected results of a healthier environment
13 have not been accomplished in all places (Kemp et al., 2009). The responses of eutrophication
14 and the extent of hypoxic area for changes in nutrient loads are different in different types of
15 systems. Also changes in climatic and hydrodynamic conditions might lead to a non-linear
16 recovery (Kemp et al., 2009). Nutrients transported from land to sea first enter the coastal
17 zones and are then further transported towards the open sea. However, not all of the supplied
18 nutrients reach the open sea as they are retained in the coastal zone (Fig. 1), which acts as a
19 filter (McGlathery et al., 2007). The retention capacity depends on different chemical,
20 physical and/or biological processes that involve nutrients e.g. denitrification, permanent
21 burial, algae and plant assimilation (Duarte and Cebrián, 1996; Voss et al., 2005). The filter
22 efficiency of the coastal zone might be of large importance for the water quality in open
23 waters.

24 Fig. 1.

25 Retention capacity is, however, not well defined. Johnston (1991) discussed that retention
26 processes are of different magnitudes and irreversibility, e.g. plant uptake and litter
27 decomposition provide short- to long-term retention of nutrients. Billen et al. (2011) and
28 Nixon et al. (1996) defined retention as the net effect of temporary and permanent removal
29 from the water phase through different biogeochemical processes. Burial and denitrification
30 lead to a permanent removal of nutrients from the ecological system (Voss et al., 2005). Plant
31 assimilation of nutrients and sedimentation of organic material might influence the temporary
32 retention, a build-up of active nutrient pools in the water and in the sediment. Some of the

Borttaget: Nutrients are involved in coastal

Borttaget: (

Borttaget:)

Borttaget: and are, thus, retained in coastal areas

Borttaget: The retention The and

1 organic material is more refractory than others e.g. parts of root systems, which also can
2 influence biogeochemical processes by enhanced sediment oxygen, nutrient and dissolved
3 organic material concentrations (McGlathery et al., 2007). Thus, temporary retention depends
4 on the release rates, translocations, and the longevity of plants, which causes variations in
5 retention capacity depending on the time scale of the study. The net effect of nutrient
6 retention in an area can be studied by the simple method of subtracting the output of nutrients
7 from the input (Johnston, 1991). This simple method of calculating the retention capacity of
8 nitrogen (N) and phosphorus (P) has been used in a number of studies (e.g. Eilola et al., 2014;
9 Hayn et al., 2014; Karlsson et al., 2010; Nixon et al., 1996; Sanders et al., 1997) for different
10 areas of the world. The retention capacity has been discussed to be related to the residence
11 time and depth in different water systems (Balls, 1994; Hayn et al., 2014; Nixon et al., 1996).
12 Hence, the longer a water parcel and its nutrient content stays within a system, the more the
13 containing nutrients are affected by the internal transformation and retention processes.
14 Filter efficiency is in the present study referred to as the capacity of the studied area to retain
15 the local nutrient loads from land and atmosphere (see Section 2.4). It is distinguished
16 between the permanent removal and temporal retention of which the latter is caused by
17 changes in the N and P inventory (Fig. 1). There are studies of nutrient retention in different
18 coastal zones around the world, but there are not enough estimates to evaluate and understand
19 its effect on the environmental status of coastal seas. Quantification of the filter efficiencies in
20 different coastal ecosystems as estuaries, archipelagos, lagoons and embayments would
21 increase the understanding and the knowledge necessary for managing the coastal zone.
22 Numerical models have been used to a larger extent for studies in lakes and freshwater
23 catchment areas (e.g. Ahlgren et al., 1988; Hejzlar et al., 2009) than for retention and filter
24 efficiency studies in coastal areas where only a few studies seem to exist in the literature (e.g.
25 Fennel et al. 2006; Seitzinger and Giblin, 1996; Xue et al. 2013),
26 The Baltic Sea (Fig. 2), located in northern Europe, is an example where the enhanced land
27 load of nutrients to the sea (Gustafsson et al., 2012) has led to eutrophication and
28 consequently increased frequency and intensity of cyanobacterial blooms, expanding bottom
29 hypoxia and dead bottom zones (e.g. Bergström et al., 2001; Conley et al., 2009; Diaz and
30 Rosenberg, 2008; Vahtera et al., 2007). Actually, the largest anthropogenically induced
31 hypoxic area in the world is found in the Baltic Sea (Carstensen et al., 2014), where it varied
32 between 70000 and 80000 km² during year 2010-2014 (Hansson and Andersson, 2014). In the

Borttaget: The importance of

Borttaget: affects the permanent

Borttaget: Billen et al. (2011) and Nixon et al. (1996) defined retention as the net effect of temporary and permanent removal from the water phase through different biogeochemical processes (e.g. biological uptake, burial and denitrification). Burial and denitrification lead to a removal of nutrients from the ecological system (Voss et al., 2005).

Borttaget: retentions on the water quality

Borttaget: of

Borttaget: larger

Borttaget: of nutrient retention in different coastal zones around the world

Borttaget: (see discussions below)

Borttaget: 1

1 Baltic Sea, most of the coastal zones and the open sea still suffer from eutrophication in spite
2 of reduced nutrient loads since the 1990s (HELCOM, 2010).

3 The aim with this study is to quantify the filter efficiency in the eutrophic Stockholm
4 Archipelago (see section 2.1) of N and P and to discuss the relative importance of different
5 physical and/or biological processes using the Swedish Coastal zone Model (SCM). In
6 addition, changes in the filter efficiency along the land-sea continuum, from the inner
7 archipelago, through the intermediate and outer archipelago to the open Baltic Sea, will be
8 studied in order to evaluate the effect of the size of the archipelago on the filter efficiency.
9 After a description of the model system (Section 2) and an evaluation of the results of SCM
10 (Section 3.1), the filter efficiency of the coastal zone is calculated and the effects of a reduced
11 land load of N and P are analyzed (Section 3.2). Conclusions finalize the study (Section 4).

12 2 Methods

13 2.1 Study site

14 The brackish archipelago of Stockholm (Fig. 2), located at the east coast of Sweden, is the
15 largest archipelago in Sweden and the second largest in the Baltic Sea. The archipelago is a
16 continuation of the river Norrström with an average discharge of about $160 \text{ m}^3 \text{ s}^{-1}$ from Lake
17 Mälaren (Lindh, 2013). The river outflow carries about 2600 metric tons (t) of N and 120 t of
18 P annually to the coastal basin “Strömmen” in the inner archipelago (Lännergren, 2010). The
19 rocky islands in the archipelago are surrounded by basins of different sizes and depths which
20 are connected by straits. In this study the archipelago has been divided into three areas: the
21 inner, intermediate and outer archipelagos. Several large islands form a natural border
22 between the inner and the intermediate archipelagos and the limited water exchange occurs
23 through five narrow sounds with shallow sills. The outflow from the inner to the intermediate
24 archipelago passes through the sounds in the surface layer, while inflows of more saline water
25 mainly occur at larger depths. The border between the intermediate and the outer archipelagos
26 follows the chain of islands in north-south direction with several connections between the
27 areas (Fig. 2).

Borttaget: 1

28 Fig. 2

Borttaget: 1

Borttaget: 1

29 The largest point sources of nutrients to the inner archipelago originate from waste water
30 treatment facilities of Stockholm, which is situated at the outlet of the Lake Mälaren. Signs of

1 eutrophication in the Stockholm Archipelago have been observed as increased ratio of
2 laminated sediments from the 1930s (Jonsson et al., 2003) and the eutrophication status in the
3 inner Stockholm Archipelago was in the early 1970s classified as highly eutrophic
4 (Lännergren et al., 2009). In the 1970s the sewage treatment facilities in Stockholm started to
5 chemically precipitate P, which reduced their P load from about 600 t yr⁻¹ to about 100 t yr⁻¹
6 (Fig. 3 in Lücke, 2014). The reduction led to some improvements of the marine environment
7 (Brattberg, 1986), but in the 1990s the areas were still eutrophic with poor bottom water
8 oxygen conditions (Jonsson et al., 2003; Rosenberg and Diaz, 1993). In the mid-1990s there
9 was a further reduction of the P to about 25 t yr⁻¹ and the sewage treatment facilities started to
10 reduce the N as well, from about 3000 t yr⁻¹ to 1250 t yr⁻¹ (Fig. 3 in Lücke, 2014), which led
11 to further improvement of the eutrophication status. In 2008 the bottom oxygen conditions
12 had clearly improved in the deeper parts and only enclosed bays, such as e.g. Stora Värtan,
13 suffered still from anoxia (Karlsson et al., 2010 and references therein). However, the annual
14 monitoring status report of the environmental status of the inner Stockholm Archipelago in
15 2014 still classified the area as unsatisfactory eutrophic (Lücke, 2015) according to the
16 national directives by the Swedish Environmental Protection Agency and the Swedish Agency
17 for Marine and Water Management (Naturvårdsverket, 2007; HaV, 2013) based on the EU
18 Water Framework Directive. The area still suffered from reduced water transparency, high
19 concentrations of phytoplankton chlorophyll and areas without any bottom fauna due to low
20 oxygen concentrations.

Borttaget: a

21 **2.2 Model description**

22 The Swedish Coastal zone Model (SCM) is a multi-basin 1D-model based on the equation
23 solver PROgram for Boundary layers in the Environment (PROBE; Svensson, 1998), coupled
24 to the Swedish Coastal and Ocean Biogeochemical model (SCOBI; Eilola et al., 2009;
25 Marmefelt et al., 1999). The model system was developed to calculate physical and
26 biogeochemical states in Swedish coastal waters. The inner, intermediate and outer Stockholm
27 archipelagoes (Fig. 2) are represented by 16, 44 and 26 sub-basins, respectively (see figure in
28 Supplement 1).

Borttaget: 1

29 **2.2.1 PROBE**

30 The physical model PROBE calculates horizontal velocities, temperature and salinity profiles
31 (Svensson, 1998; Omstedt, 2015). The surface mixing is calculated by a k - ϵ turbulence model

1 and the bottom mixing is a parameterization based on the stability in the bottom water. Light
2 transmission, as well as ice formation growth and decay, are also included in the model. The
3 vertical grid resolution is half a meter in the uppermost layers, one metre from 4-70 m, and
4 two metres between 70-100 m. The general differential equation of the PROBE solver is
5 formally written as

$$6 \quad \frac{\partial \phi}{\partial t} + \frac{\partial}{\partial x_i} u_i \phi = \frac{\partial}{\partial z} \left(\Gamma_\phi \frac{\partial \phi}{\partial z} \right) + S_\phi \quad (1)$$

7 Here ϕ is the dependent variable, t time, z vertical coordinate, x_i horizontal coordinates, u_i
8 horizontal velocities, Γ_ϕ vertical exchange coefficient, and S_ϕ source and sink terms. Vertical
9 advection (and moving surface) is included accounting for vertical transport in sub-basins due
10 to in and outflows. The sources and sinks determined by the ecosystem model are added to S_ϕ .

11 The water exchange between the sub-basins is controlled by the baroclinic pressure gradients.
12 The net flow through the sounds will be the same as the river discharge from land in order to
13 preserve volume. Inflowing water to a sub-basin is interleaved into its density level without
14 any entrainment, and heavy surface water in one sub-basin may thus reach the bottom level in
15 an adjacent basin. The sea level variations outside the boundary are of minor importance for
16 the SCM results and are therefore not included in the forcing. The water exchange across the
17 boundary between the coastal zone and the open sea is assumed to be in geostrophic balance,
18 since this boundary is open with a width greater than the internal Rossby radius. A time step
19 of 600 seconds was used in the present simulations.

20 **2.2.2 Biogeochemical model (SCOBI)**

21 The SCOBI model describes the biogeochemistry of marine waters in the Baltic Sea and
22 Kattegat (Eilola et al., 2009). Nine pelagic and two benthic variables (Fig. 3) are described in
23 the SCM-SCOBI model. In the pelagic zone three different phytoplankton groups (diatoms,
24 flagellates and others, and cyanobacteria), one zooplankton group, one pool for detritus and
25 three inorganic nutrients pools (nitrate, ammonium and phosphate) are represented. The
26 model also calculates oxygen and hydrogen sulfide concentrations, of which the latter are
27 represented by “negative oxygen” equivalents ($1 \text{ ml H}_2\text{S l}^{-1} = -2 \text{ ml O}_2 \text{ l}^{-1}$) and includes the
28 conversion of sulfate into hydrogen sulfide (Fonselius, 1969). Thus, the negative oxygen
29 corresponds to the amount of oxygen needed to oxidise the hydrogen sulfide. The sediment in
30 the present model is parameterized by one vertically integrated bulk sediment layer (level 3 in

Borttaget: 2

1 Soetaert et al, 2000). Organic material that sinks to the sediment is divided into one benthic
2 nitrogen pool (NBT) and one benthic phosphorus pool (PBT). SCOBI has been used and
3 validated in several studies, both coupled to the basin scale Baltic Sea model PROBE-Baltic
4 (e.g. Marmefelt et al., 1999) and to the three dimensional Rossby Center Ocean model (RCO;
5 e.g. Meier et al., 2011).

6 In the model the processes of phytoplankton assimilation, mortality and nitrogen fixation,
7 zooplankton grazing, excretion of detritus and dissolved inorganic nitrogen (DIN) and
8 phosphorus (DIP), the oxygen and temperature dependent mineralization of detritus, benthic
9 N and benthic P, nitrification and denitrification are described. Phytoplankton assimilates
10 carbon (C), N and P according to the Redfield molar ratio (C:N:P=106:16:1) and the biomass
11 is represented by chlorophyll (Chl) according to a constant carbon to chlorophyll mass (mg)
12 ratio (C:Chl=50:1). Light attenuation depends on background attenuation due to water and
13 humic substances and a variable attenuation caused by particulate organic matter
14 (phytoplankton, zooplankton and detritus). All particulate variables sink downward through
15 the water column. Predation is used as a closing term to parameterize interactions with higher
16 trophic levels in the ecosystem and move matter from zooplankton to the detrital and inorganic
17 pools. Resuspension of sediment that is important in the open Baltic Sea (Almroth-Rosell et
18 al., 2011) has not yet been implemented in this SCOBI version, but the sediment releases
19 dissolved inorganic nutrients back to the water mass, with the release of phosphate being
20 redox dependent. Some fractions of benthic N and P are assumed to be buried in the sediment
21 as a permanent sink, and are hence removed from the system. For further details of the
22 SCOBI model the reader is referred to Eilola et al. (2009; 2011).

23 | *Fig. 3.*

Borttaget: 2

24 **2.2.3 Forcing**

25 The SCM-SCOBI model system is forced by weather, the conditions in the sea outside the
26 archipelago, point sources, discharge of freshwater and nutrients from land and atmospheric
27 deposition of nutrients. The initial values for both the pelagic zone and the sediment are
28 derived from spin-up simulations.

29 There are two types of land derived forcing; discharge of water and nutrients from both rivers
30 and surface run-off from the drainage area given by the S-HYPE model (Lindström et al.,
31 2010) and point sources representing sewage plants and industries. The run-off is added to the
32 surface water of each basin and no reduction of river nutrients due to precipitation at river-

1 mouths is assumed in this model setup. The point sources of nutrient loads are assigned to the
2 depth levels mostly resembling the actual depth of the discharge. The inorganic riverine
3 nutrient loads are added as DIN and DIP to the SCM. The organic nutrients in the land loads
4 are calculated from the difference between total nitrogen (TN) and DIN, and total phosphorus
5 (TP) and DIP, respectively. The bioavailability and the composition (dissolved or particulate)
6 of the organic nitrogen and phosphorus loading from land are generally not known. In the
7 present model configuration the fraction of organic nutrient loads that follows the Redfield
8 ratio are assumed to be bioavailable and will be added to the detritus pool in the model, while
9 the remaining fractions of nutrient loads are treated as conservative tracers in the model..

10 The weather forcing consists of solar insolation, air temperature, wind, relative humidity and
11 cloudiness. The insolation and all the radiation and heat fluxes across the water-air interface
12 are calculated by the PROBE model. The weather variables are taken from a gridded database
13 developed at the Swedish Meteorological and Hydrological Institute (SMHI), using 3-hourly
14 meteorological synoptic monitoring station data, and the depositions of nitrogen species
15 (NHX and NOX) are calculated by the MATCH model (Robertson et al., 1999). For the
16 deposition of phosphate, a literature value of $0.5 \text{ kg m}^{-2} \text{ month}^{-1}$ is used (Areskoug, 1993).

17 The boundary conditions to the open Baltic Sea is set by vertical mean profiles calculated by a
18 one dimensional PROBE setup for each Baltic open water area and assimilation of monitoring
19 data. The monitoring data used in the assimilation are extracted from the stations MS4, US5B,
20 SR5, BY31 and BY29 (Fig. 2) depending on depth and time, to get the best representation of
21 the open sea's influence on the SCM model domain.

Borttaget: 1

22 2.3 Evaluation strategy

23 To quantify the fit between modelled values and observations a correlation coefficient, r , was
24 calculated (Eq. 2).

$$25 \quad r = \frac{\sum_{i=1}^n (P_i - \bar{P})(O_i - \bar{O})}{\sqrt{\sum_{i=1}^n (P_i - \bar{P})^2 \sum_{i=1}^n (O_i - \bar{O})^2}} \quad 2$$

26 where P is model value, O is observation of the analyzed parameter, i is the data number and n
27 is the total number of data points. Two series of observations and model values that are
28 identical will lead to an r value equal to one, while uncorrelated data result in a r value close

1 to zero. In addition to the r value, the average cost function (C) values (Eq. 3) for the different
2 parameters were used in the evaluation of the SCM results.

3

$$4 \quad C = \frac{\sum_{i=1}^n \frac{|P_i - O_i|}{sd(O_i)}}{n} \quad 3$$

5 A cost function describes the proximity of model results and observations by normalizing the
6 difference between them with the standard deviation (sd) of the observations. If average
7 model results fall within the standard deviation of observations, C is below one which is
8 regarded as good. Results that are within two standard deviations will be regarded as to be on
9 an acceptable level. The corresponding simulation levels, good and acceptable, for the
10 correlation coefficient are achieved when r is higher than two thirds (0.66) and one third
11 (0.33), respectively. This approach using r and C has been used in earlier studies (Edman and
12 Omstedt, 2013; Edman and Anderson, 2014) and is based on methods by Oschlies (2010).

13 The outflow from Lake Mälaren is three orders of magnitude larger than the sum of all other
14 S-HYPE fresh water components to the inner Stockholm Archipelago. The output from S-
15 HYPE of fresh water and nutrient loads from Mälaren to the Stockholm Archipelago was
16 therefore used in the evaluation of the fresh water forcing to SCM. Observations of freshwater
17 discharge were retrieved from the Baltic Environmental Database (BED, 2015) at the Baltic
18 Nest Institute, Stockholm University. The correlation between monthly mean of observed and
19 simulated discharge for the period (1990-2012) was then calculated.

20 In the evaluation of the results of the SCM in different basins, the long term averages (1990-
21 2012) of the vertical distribution of salinity, DIN, DIP and oxygen during winter (November-
22 February) and summer months (May-August) were compared to corresponding observations
23 for the whole modelled period. Further, the correlation r and the mean cost function C of the
24 vertical distribution of observations and model output were calculated. Also the long term
25 averages of the seasonal variations in surface temperature, DIN, DIP and bottom water
26 oxygen concentrations were used in the evaluation by calculating the corresponding r and C
27 values.

28 Observations from the Stockholm Archipelago (Fig. 4) were provided by Stockholm City and
29 Stockholm University. For the quantitative validation described above the quality of
30 observations from each site (Table 1) had to fulfil three requirements to be used in the

Borttaget: 3

1 validation process; 1) period coverage: 80% of the years sampled; 2) annual coverage: at least
2 7 of the 12 months sampled; and 3) vertical data coverage: at least 5 depth levels frequently
3 measured over the full depth of the basin. In addition at least 3 months with observations were
4 required for the evaluation of winter and summer conditions. Average values were then
5 calculated for periods and depth levels with dense data distribution. The model output was
6 used in the same way as observations, and the modelled averages were calculated for the same
7 time intervals and depth ranges.

8 *Fig. 4.*

Borttaget: 3

9 *Table 1.*

10 2.4 Calculation of retention

11 The retention of P and N in a region can be calculated as the difference between the load and
12 the outflow (Almroth-Rosell et al., 2015; Hayn et al., 2014; Johnston, 1991; Meier et al.,
13 2012). The input of nutrients is the sum of inflows from outer areas, rivers, land runoff, point
14 sources and atmospheric load, while the outflow of nutrients is the export from the area to
15 outer seas (Fig. 1). N₂-fixation is another process that needs to be taken into account as it is a
16 source of bioavailable N to the system. Retention in the present study can be temporal or

Borttaget: 4

17 permanent. Permanent retention removes the nutrients permanently from the pool of nutrients
18 in the modelled system. Burial is the only retention process that permanently removes P. For
19 N, in addition to burial, also benthic and pelagic denitrification is considered as permanent
20 removal. The temporal retention during a studied period can be negative or positive
21 depending on changes in the pelagic and benthic inventory of nutrients. The nutrient pools
22 include both the inorganic and organic nutrients. Factors that affect the benthic N and P pools
23 are the sedimentation of organic material from the water column, the decomposition of
24 organic material and the release of inorganic nutrients back to the water column, as well as
25 burial of nutrients. The pelagic N and P pools are affected by the supply from land, the export
26 of organic material to the sediment, the release of nutrients from the sediment to the water
27 column and to the net export of nutrients to downstream areas.

Borttaget: . A build-up of the pelagic and benthic active pools of P and N are referred to as temporal retention processes. P

Borttaget: For modelled P bB

Borttaget: ., while f

Borttaget: are

28 The different processes that affect retention have been calculated separately, as they are
29 included in the biogeochemical model SCOBI. Total retention (R_{tot}) is the sum of both
30 permanently and temporally retained P and N. Area specific retention, R_{AS} , is the retention
31 normalized to the area and is calculated from eq. 4:

Borttaget: ing

Borttaget: (burial, denitrification, benthic and pelagic nutrient pool changes)

Borttaget: (see supplement 2 for retention definitions)

Borttaget: The a

1 $R_{AS} = \frac{R_{tot}}{A}$ 4

2 where A is the size of the area. R_{AS} can be used to compare the retention in basins of different
3 sizes. The filter efficiency, F_{eff} , is calculated from eq. 5:

4 $F_{eff} = \frac{R_{tot}}{Nu_{land}} \times 100$ 5

5 where Nu_{land} is the sum of the nutrient load from land and the deposition from air. The F_{eff} is
6 an estimate of the proportion (%) of the nutrients from land and atmosphere that is retained
7 within the area. Similarly the retention efficiency (R_{eff}) can also be calculated, defined as the
8 proportion of the total nutrient load (sum of all sources, including import from surrounding
9 waters) that is retained within the area. In the present study, however, the focus is on the filter
10 efficiency.

Borttaget: (supplement 2)

Borttaget: effect and filter

11 The total retention efficiency was calculated for the entire Stockholm Archipelago, and also
12 separately for the inner, intermediate and outer archipelagos in order to investigate the spatial
13 gradient of retention capacity from the inner coastal zone towards the open Baltic Sea.

Borttaget: and the filter

14 The residence time is defined as the average time water, or a dissolved substance, spends
15 within a particular basin (Bolin and Rodhe, 1973). In the present study the residence time of
16 the freshwater is calculated to relate the filter efficiency to physical characteristics of the
17 archipelago as described by Nixon et al. (1996). A fresh water tracer in the model is used to
18 determine the freshwater volume (V_{gr}) in the different parts of the archipelago. The fresh
19 water residence time is estimated by the flushing time calculated from the freshwater volume
20 divided by the fresh water discharge received from land (Q_r) as in the freshwater fraction
21 method discussed by Sheldon and Alder (2006). The filter efficiency was calculated for: the
22 inner, the sum of the intermediate and inner archipelago, and the entire Stockholm
23 Archipelago.

24 **2.4.1 Oxygen reduction scenario**

Borttaget: Fig. 4. ¶

25 In the model, denitrification is an O_2 dependent process that has a maximum rate at O_2
26 concentration of about $45 \mu\text{mol l}^{-1}$ ($\sim 1 \text{ ml l}^{-1}$) while denitrification halts under anoxic
27 conditions. Also P is affected by oxygen since P has an oxygen dependent adsorption
28 behaviour on particulate iron(III)oxyhydroxides (Mortimer, 1941). The adsorption of P on
29 particles can lead to higher burial rates during oxic conditions compared to anoxic conditions
30 when the release rate of P from the sediment is higher (Viktorsson et al., 2012; Almroth-

Rosell et al., 2015). This O₂ dependent adsorption behaviour is also simulated by the model (Eilola et al., 2009) using a reduced release of P from the sediment when O₂ is present in the bottom water. The effect of the O₂ concentration on the filter efficiency is studied in an experiment where the O₂ concentration was reduced in the SCOBI model with a fixed amount, 134 μmol l⁻¹ (3 ml l⁻¹), during the simulation period (1990-2012).

2.4.2 Nutrient load reduction scenario

The SCM is also used to investigate the effect of a reduction of the nutrient load from land to the Stockholm Archipelago. The reductions are applied to the forcing from 2010 with a river load of 4027 t N yr⁻¹ and 163 t P yr⁻¹, and a load from point sources of 1805 t N yr⁻¹ and 30 t P yr⁻¹ to the entire Stockholm archipelago. Reductions of point sources were estimated from realistic minimum discharge concentrations of N and P from sewage treatment facilities based on technical feasibility, but not on economic or resource sustainability (Table 2, Kerstin Rosén Nilsson, County Administrative Board of Stockholm, personal communication). Point sources from different industries are assumed to decrease their discharge of N and P by 10 %. The minimum discharge concentrations, and the 10 % reduction from industries, resulted in reductions by approximately 51 % of N and 34 % of P from point sources. The reductions of N and P from land runoff, e.g. due to decreased nutrient load from agriculture and increased use of small sized sewage treatment plants by individual households, are set to 15 % for N and 10 % for P. The combined reductions in rivers and point sources results in a total decrease of N and P load by 20 % and 12 % relative to 2010. A SCM model spin-up run period of 45 years, with forcing from year 2010 provides the steady state initial conditions used for the reduction experiment. After the spin-up period the reductions of the nutrient loads are implemented.

Table 2.

Borttaget: R

Borttaget: of the nutrient land load

Borttaget: The SCM is also used to investigate the effect of a reduction of the nutrient load from land to the Stockholm Archipelago.

Borttaget: The loads of P and N from sewage treatment facilities depend on their size, i.e. the number of person equivalents (pe) for which they were built.

Borttaget: Estimates of realistic minimum discharge concentrations of P

Borttaget: from sewage treatment facilities used in the SCM reduction scenario, which are based on technical feasibility but not on economic or resource sustainability, are given in Table 2 (Kerstin Rosén Nilsson, County Administrative Board of Stockholm, personal communication). The reductions of P and N from land runoff, e.g. due to decreased nutrient load from agriculture and increased use of small sized sewage treatment plants by individual households where these are not connected to a municipal sewage treatment facility are set to 10 % and 15 % for P and N, respectively. Also point sources from different industries are assumed to decrease their discharge of N and P with 10 % as well. The reductions result in a total decrease of P and N load by 12 and 20 %, respectively, to the entire Stockholm Archipelago. ¶

Borttaget: river and weather

Borttaget: , according to Table 2

1 3 Results and discussion

2 3.1 Validation

3 The variability of the modelled discharge of water and nutrients by the S-HYPE model agrees
4 well with observations (Fig. 5 and Table 3) for the simulated period (1990-2012). A good
5 description of river runoff is needed because the nutrient loads are strongly related to the
6 magnitude of river outflow (Q_F), as seen in Fig. 5. The model seems to slightly underestimate
7 the spring discharge and overestimate low flow regimes relative to observations. However,
8 overall it captures a realistic annual variation of the discharge, which is reflected in high
9 correlation coefficients (Eq. 2) for all evaluated parameters (Table 3). Highest correlation
10 coefficients are found for Q_F and TN, compared to the slightly lower values for TP, DIP and
11 DIN, which is in accordance with previous studies (Grimvall et al., 2014; Strömquist et al.,
12 2012). An extensive validation is also available in Sahlberg et al. (2008).

13 |Fig. 5.

14 |Table 3.

15 Datasets from eight stations (Table 1) fulfilled the requirements of good data availability and
16 were used in the evaluation of the SCM model results. There are aspects that are important to
17 have in mind when comparing model results and observations. In the model the state variables
18 are horizontally averaged in each basin, while observations are measured at one station at a
19 certain location. The Stockholm Archipelago has relatively large spatial salinity gradients and
20 the representativeness of a station when compared to model results can be somewhat limited
21 if e.g. the position of the station is close to an out- or inlet of the basin. Observations may in
22 general also be influenced by local conditions, e.g. sewage effluents, high sediment fluxes or
23 stagnant conditions, which are smeared out in the average results of the model. Still we
24 assume for the present study that the station data are good enough for the quantitative model
25 validation and give a background for discussions about model strengths and weaknesses. As
26 an example, validation results are shown for one of the basins where the number of
27 observations is large enough both during summer and winter periods to be included in the
28 validation process. The example is from station Blockhusudden (Position G in Fig. 3), where
29 the largest data set of observations was found. The station is situated at the boundary between
30 the innermost basin Strömmen and the next adjacent sub-basin.

Borttaget: A clear relationship between the magnitude of

Borttaget: and the nutrient loads is observed both for monthly observations and S-HYPE output (Fig. 5).

1 The objective correlation coefficients (Eq. 2) and the cost function value (Eq. 3) for the
2 different state variables implied correspondingly that the model manage to simulate the
3 average vertical winter and summer profiles with good or acceptable skills in the basin
4 Strömmen (Fig. 6g), except for the average seasonal value of DIN that was described as not
5 good. The differences between model results and observations of DIN may be a result of the
6 location of the monitoring station.

7 The long term average summer depth profiles of modelled salinity and oxygen in the basin
8 Strömmen correlate well with observations, while the winter values of salinity were too low,
9 especially in the surface layers (Fig. 7a,b). This difference is partly due to the fact that the
10 salinity of a station at the entrance to the basin is more reflecting the boundary conditions of
11 the downstream basin than the mean conditions in Strömmen. The surface winter
12 concentrations of oxygen were too high, but decreased with depth and became too low in the
13 lower layers (Fig. 7b). It might be expected that winter surface oxygen concentrations in
14 observations should be higher than in summer because of the temperature effect on oxygen
15 saturation concentrations as seen from the model results. However, the number of
16 observations during winter are limited and occurred mostly in November and February, which
17 may influence the average values of the observations.

18 The results indicate that there is an impact from local conditions at the monitoring station that
19 are not captured by the model setup. The modelled DIN depth profiles show higher values at
20 about 15 m depth during both winter and summer (Fig. 7c), while the DIP profiles values
21 seems to be satisfactory at all depth and periods (Fig. 7d). Also the individual observations
22 show higher concentrations of both DIN and DIP around 15 m depth which is where the
23 halocline has its largest vertical gradient. This depth level corresponds to the depth where two
24 sewage water treatment plants relieve their sewage water in the model. The winter
25 stratification was stronger in the model because of the lower surface salinity. This hampers
26 the vertical transports of oxygen and has an influence on the winter oxygen conditions in the
27 deep water that were lower in the model compared to the observations from the more well
28 ventilated entrance area.

29 The average seasonal variation of the surface temperature and the bottom water oxygen
30 concentrations was captured by the model, but not the increase of surface nutrients, especially
31 DIN, during autumn (Fig. 8). The surface salinity was overall somewhat low, which probably
32 is a result of the location of the monitoring station, as described above.

1 In the other basins used in the evaluation (vertical and seasonal profiles are not shown) of the
2 SCM state variables during winter, summer and season were simulated with good or
3 acceptable skills, except for the average vertical summer profiles of DIN in the basin
4 Solöfjärden (Fig. 6c) and oxygen concentration in the basin Sandöfjärden (Fig. 6a). The
5 combined model skills, which were calculated as the average of the individual r and C values,
6 were good in six of the eight evaluated basins (purple cross in Fig. 6). In the remaining two
7 basins the skills were considered as acceptable.

8 Fig. 1

Borttaget: 6

9 Fig. 2

Borttaget: 7

10 Fig. 3

Borttaget: 8

11 3.2 Retention of nutrients in the Stockholm Archipelago

12 The load and the inventories of N and P may change and vary between the beginning and the
13 end of a studied period, thus the determination of total nutrient retention depends on the time
14 scales of consideration as discussed in section 3.2.3. During the period 1990-2012 on average
15 174 t P yr^{-1} and 5846 t N yr^{-1} entered the inner archipelago, mainly from the Lake Mälaren.
16 That is a major part of the 217 t P yr^{-1} and 8288 t N yr^{-1} which entered the entire Stockholm
17 Archipelago (Fig. 9). The P load from point sources was clearly lower than the river load (Fig.
18 10). However, the N load from point sources was higher than the river load in the beginning
19 of the studied period (Fig. 10b), but decreased in the middle of the 1990s due to the
20 implementation of a more effective method to remove N in the waste water treatment
21 facilities. The P supply to the intermediate archipelago mainly originated from runoff from
22 land, while for N there were also some point sources that contributed to the land load on the
23 same level. In the outer archipelago the nutrient load from land was almost negligible and
24 most of the nutrients were deposited from the atmosphere.

25 Fig. 9.

26 Fig. 10.

27 Largest amounts of P and N in the model were retained in the outer archipelago compared to
28 the intermediate and inner archipelagos (Fig. 9). The retentions of all supplied P and N,
29 including the net import from upstream areas, within the inner, intermediate and outer
30 Stockholm archipelagos amounts to 18 %, 23 % and 48 % for P, respectively, and 14 %, 26 %
31 and 60 % for N, respectively. The area of the three zones increases from inner (109 km^2), to

1 the intermediate (759 km²) and to the outer archipelago (2360 km²) and thus, the retention of
2 nutrients seems to increase with increased area. On the other hand, the average of the area
3 specific retention of P and N was for the simulation period highest in the inner archipelago,
4 and decreased towards the open sea (Fig. 11). The permanent retention was relatively stable
5 during the simulated period, while fluctuations in the temporal retention reflect the effect of
6 varying riverine nutrient input (Fig. 10c, d). The water depth and the residence time affect the
7 retention of nutrients, which will be further discussed in Section 3.2.2. The largest part of the
8 total retention in the entire Stockholm Archipelago was permanent, which for P means burial.
9 For N benthic denitrification represented as much as almost 92 % of the permanent retention,
10 burial for less than 8 % and pelagic denitrification was below 1 %.

11 *Fig. 11.*

12 Karlsson et al. (2010) found in their empirical study for 1982-2007 that about 15 % of the
13 total input of N and 10 to 13% of the total input of P were retained in the inner Stockholm
14 Archipelago. However, their numbers are based on the total input, thus both the land load and
15 an estimated input from outer areas, i.e. the intermediate Stockholm Archipelago. A
16 recalculation from the given numbers in their study resulted in a filter efficiency of about 25
17 and 24 % for N and about 21 and 30 % for P of the nutrient load from land and atmosphere
18 for the periods 1982-1995 and 1996-2007, respectively. These numbers of the filter efficiency
19 are higher than the numbers in the present model study. To be able to compare the numbers a
20 recalculation of the filter efficiency in the SCM for the latter period (1996-2007) in the inner
21 archipelago was performed, but did not change the SCM results considerably. The largest
22 difference between the two studies is caused by the calculation of net exchange of nutrients
23 through the sounds. The transport through the sounds was in Karlsson et al. (2010) calculated
24 from average volume flows estimated from mass balance calculations for salt together with
25 budget calculations using observations of average nutrient concentrations. In the present study
26 the exchange of nutrients between the inner and the intermediate archipelago was part of the
27 dynamic model calculations in the SCM. The SCM net outflow from the inner archipelago for
28 N and P was about 11 % and 8 %, respectively, larger compared to the net outflow of the
29 nutrients in the Karlsson et al. (2010) study. Another difference between the two studies was
30 the land load of P, which was about 8 % lower in the SCM. The difference in land load of N
31 was only about 1 %. Thus, calculations from an empirical model based on Knudsen's
32 relations (Knudsen, 1900) and calculations using long term average values resulted in about
33 10 % higher retention efficiency values compared to the calculations from SCM, a coupled

Borttaget: during

Borttaget: on a stable level

Borttaget: riverine

Borttaget: are

Borttaget: ing

1 numeric physical-biogeochemical model with high vertical resolution and small time step. In
2 spite of the difference in models the result are surprisingly close.

3 The average temporary retention in SCM for the entire simulated period is negative in all
4 three parts of the archipelago for both P and N (Fig. 9 and Fig. 10). The reason for negative
5 temporary retention is mainly a decrease in the benthic nutrient pools during the period (Fig.
6 12). The largest decrease (29 %) is found in the pelagic pool of N in the inner archipelago,
7 which coincides with the decrease in N load from point sources (Fig. 10). In the intermediate
8 and outer Stockholm archipelagos the pelagic pool of N remains on about the same level
9 through the whole simulation period. The large decreases in the benthic pools of N and P (14-
10 18 %) occur in the intermediate and outer archipelagos, while there are only small changes in
11 the pelagic and benthic pools of P in the inner archipelago. Because of the nutrient retention
12 there is a reduced net transport of N and P from the inner archipelago towards the
13 intermediate and outer archipelagos and further to the open sea during the simulated period
14 (Fig. 9). The annual temporary retention of P in the entire Stockholm Archipelago increases
15 with time during the simulated period (Fig. 10). There is a change to positive values at the end
16 of the period, when there again is a build-up of the benthic pools of P (fig. 12). The build-up
17 is most likely a result of better oxygen conditions in the modelled deep water (not shown)
18 during the end of the simulation period, which lead to a lower release of P from the sediment
19 to the water column (Eilola et al., 2009). For the temporary retention of N there is no visible
20 trend in the variation with time. In addition to the nutrient load from land, and the net export
21 of nutrients to outer areas, there is also an extensive circulation of nutrients between the coast
22 and the open sea. The importance of imported nutrients into the coastal zones from sea have
23 been discussed in earlier studies (e.g. Humborg et al., 2003) in which it was concluded that
24 many estuaries has a net import of DIN and DIP from sea, e.g. Chesapeake Bay (Boynton et
25 al. 1995). This is shown also for e.g. the Mid Atlantic Bight where almost three times the
26 riverine input of N is denitrified (Fennel et al., 2006). In different parts of the shelf in the Gulf
27 of Mexico the denitrified proportion of the land input of N is in total 86 %, where locally on
28 the different part of the shelves the denitrification fraction of the supply from land varied
29 between 68 % and 341 % (Xue et al., 2013). Thus, in many cases the import is larger than the
30 export and the coastal zones works as a filter not only for the nutrients from land, but also for
31 the nutrients from the open sea as also discussed in section 3.2.3.

32 Fig. 12.

Borttaget: s

Borttaget: these

1 3.2.1 The coastal filter

2 From the present results it can be concluded that the Stockholm Archipelago works like a
3 filter for nutrients that enter the coastal zone from land and atmosphere. However, a rather
4 large area of the archipelago is needed to effectively retain the nutrients. About 82 and 86 %
5 of P and N supplies, respectively, pass the small inner archipelago and are exported to the
6 intermediate archipelago. In the intermediate and the outer archipelago all local supplies of
7 nutrients from land and atmosphere are retained together with a fraction of the nutrients
8 imported from the inner archipelago. The filter efficiencies increase with increased coastal
9 area from land to the sea continuum (Fig. 13). However, the filter efficiency of the entire
10 Stockholm Archipelago is not effective enough to retain of all the nutrients that enter the
11 system from land and the atmosphere, but still, at least 65 % and 72 % of the supplied P and
12 N, respectively, are retained. The total retention numbers (permanent and temporary)
13 correspond to 141 t P yr⁻¹ and 5954 t N yr⁻¹ (Fig. 9). Since Stockholm Archipelago is the
14 largest archipelago in Sweden it might be that most of the other Swedish coastal areas with a
15 large run-off from land would be less effective as coastal filters and, thus, contribute to a
16 larger extent to the eutrophication in the open sea. This is one question in focus of an on-
17 going study where the entire Swedish coastal area will be evaluated similarly to the present
18 study.

Borttaget: take care

19 3.2.2 Processes affecting retention

20 The present study was performed in an area characterised as an eutrophic archipelago in an
21 inland sea with basins having oxic, hypoxic and anoxic bottom waters. Nixon et al. (1996)
22 showed that the retention of P and N correlated to the log scale of the ratio between the
23 average depth and the residence time of the study areas, which is confirmed by the results
24 from the studies by Billen et al. (2011), Hayn et al. (2014) and Nielsen et al. (2001) as well as
25 by the present study (Fig. 13). The freshwater residence time in the Stockholm Archipelago is
26 48 days in the inner, 108 days in the middle and inner, and 185 days in the entire area. No
27 clear relationship was found between the filter efficiency and the average depth, which vary
28 between 17 m and 20 m for the three areas. These results are in agreement with Nixon et al.
29 (1996) who showed that including the depth in the analysis of retention vs residence time did
30 not much improve their regression. In the present study the change of the filter efficiency with
31 residence time is about 0.5-0.6 % per day. The results of the present retention estimates are in
32 agreement with results from previous studies (Billen et al., 2011; Hayn et al., 2014; Nielsen et

Flyttad (infogning) [2]

Borttaget: 4-5 % yr⁻¹ with increasing fresh water residence time.

1 | al., 2001; Nixon et al., 1996), but with somewhat higher values in the entire archipelago (Fig.
 2 | 13). Their studies were performed in various types of systems: coastal lagoons, drowned river
 3 | estuaries, coastal embayments, and inland seas in North America and in Europe. Those
 4 | systems varied from being relatively pristine to systems with large point sources (eutrophic),
 5 | and they also varied between oxic to hypoxic and/or anoxic conditions. In shallow areas larger
 6 | parts of the sinking particulate organic material may reach all the way down to the sea floor
 7 | where it can be exposed to retention processes such as burial and denitrification. On the other
 8 | hand, in a much deeper area a larger part of the organic material may become re-mineralised
 9 | within the water column on its way down to the sea floor. The nutrients can then be re-used
 10 | by phytoplankton and/or be further transported out from the system. Long residence times in a
 11 | system increase the time of exposure for biogeochemical transformation processes and
 12 | sedimentation within the system and larger parts of the nutrients may be retained.

13 | Fig. 13.

14 | Denitrification increase the retention in areas with longer residence times (Nixon et al., 1996,
 15 | Finlay et al., 2013) as also seen from Fig. 13. In the Randers Fjord, the residence time was
 16 | short (six days) and the filter efficiencies of N and P were lower, 10 % and 9 %, respectively
 17 | (Nielsen et al., 2001), compared to the Stockholm Archipelago where the freshwater residence
 18 | time is longer. The denitrified proportion of the permanently retained N was also lower, about
 19 | 60 % compared to in the Stockholm Archipelago (92 %). Oxygen is an important factor
 20 | regulating the magnitude of denitrification. In waters with longer residence time the bottom
 21 | water might be less ventilated, and, thus, the bottom water oxygen concentrations lower with
 22 | higher denitrification as a result. As a result of the forced reduction of the oxygen
 23 | concentrations with 134 μ M the hypoxic areas increased by 49 km² (300 %) and the anoxic
 24 | area increased by 13 km² (360 %) in the entire Stockholm Archipelago. The reduced oxygen
 25 | concentration led to increased N retention (780 t yr⁻¹ or 14 %) due to increased denitrification
 26 | and to decreased P retention (49 t yr⁻¹ or 28 %) due to higher release of P from the sediment.
 27 | Denitrification increased the fraction of permanent retention from 92 % to 94 %, while the
 28 | buried fraction decreased. The inner archipelago had the largest increase of hypoxic and
 29 | anoxic areas and also the largest changes in retention of N and P. The N retention increased
 30 | there by 243 t yr⁻¹ (29 %) and the P retention decreased by 9 t yr⁻¹ (38 %).

31 | Benthic primary producers and benthic fauna are also important for the retention of nutrients
 32 | in shallow coastal ecosystems (McGlathery et al., 2007; Norkko et al., 2012). Assimilation of
 33 | nutrients during primary production does not directly change the inventory of N and P, but

Borttaget: lower

Borttaget: and inner plus intermediate

Borttaget: s

Flyttad uppåt [2]: Nixon et al. (1996) showed that the retention of P and N correlated to the log scale of the ratio between the average depth and the residence time of the study areas, which is confirmed by the results from the studies by Billen et al. (2011), Hayn et al. (2014) and Nielsen et al. (2001) as well as by the present study (Fig. 13).

Borttaget: These two parameters therefore affect the retention of nutrients (Fig. 13) and the correlation implies that the retention occurs mostly in the sediment due to processes such as burial and denitrification, which is the case in the present model study. However, in the present study no clear relation was found between the average depth (10-20 m for all three areas) and the filter efficiency. Actually, also Nixon et al. (1996) discussed this and concluded that including the depth in the analyze of the retention vs residence time did not much improve the regression. The filter efficiency in the present study did, however, increase with 4-5 % yr⁻¹ as the fresh water residence time increased in the land to sea continuum.

Flyttad uppåt [3]: Thus, a reduced oxygen concentration led to increased N retention due to increased denitrification.

Borttaget: The denitrification process has an oxygen dependency and the largest sensitivity of the denitrification rates to changes in oxygen concentrations in the model occur at concentrations below 180-220 μ mol l⁻¹ (~4-5 ml/l). The maximum ...

Flyttad (infogning) [3]

Borttaget: has in earlier studies been shown to

Borttaget: This is confirmed also in a study with a two-layer hydrodynamic ...

Borttaget: (Nielsen et al., 2001)

Borttaget: . T

Borttaget: only on average six

Borttaget: estimated to 10

Borttaget: 9

Borttaget: which were lower

Borttaget: in

Borttaget: with a

Borttaget: in the inner archipelago of about 110 days

Borttaget: Waters with longer residence times are less ventilated at the sediment ...

Borttaget: factors

Borttaget: by algae

Borttaget: pools

Borttaget: nutrients

1 | transfer the nutrients into organic material. Plant uptake at the bottoms can e.g. lead to
2 | increased burial and also influence on the oxygen dependent biogeochemical processes in the
3 | sediment due the plant metabolism (McGlathery et al., 2007). These processes are not yet
4 | implemented in the SCM that only include pelagic primary production, and are therefore not
5 | included in the present study. Including these processes may have some impact on the model
6 | dynamics e.g. on bottoms where seagrasses and burrowing macrofauna might influence the
7 | decomposition of organic material and the permanent burial of nutrients and organic matter.
8 | The evaluation of forcing and model results indicate, however, that the model system is able
9 | to reproduce much of the observed physics and nutrient dynamics in the archipelago which
10 | give confidence to the budget estimations of nutrient retention in the area. A quantitative
11 | evaluation of the effect and the implementation of benthic flora and fauna to the model are
12 | therefore left for future work.

13 | It is also important to know whether a system is in balance with the nutrient loads or not since
14 | it would affect the retention capacity. In this study the temporary retention is negative for both
15 | N and P in all three areas of the Stockholm Archipelago which implies that the system is not
16 | in a steady state. This imbalance is however expected since there are reductions of the nutrient
17 | loads in the first part of the simulation period (Fig. 10a, b). However, the possibility that the
18 | results may be influenced by unknown initial conditions of sediment concentrations should
19 | not be excluded. There are only few observations available and the knowledge about the
20 | amount of sediment nutrients involved in biogeochemical cycles is poor.

21 | 3.2.3 Response to nutrient load reduction

22 | The fastest response in the nutrient load reduction experiment is seen in the pelagic pool of N
23 | which rapidly decreases, but reaches a steady state after about three years with reduced loads
24 | (Fig. 14). The pelagic pool of P decreases in the inner archipelago but increases slightly in the
25 | outer areas. The changes in P pools are slower compared to those in N pools. The large and
26 | fast decrease of pelagic N in the inner archipelago, results in a decreased N:P ratio (Table 4),
27 | as well as (not shown) lower chlorophyll concentrations, reduced sedimentation, and
28 | increased export of P from the inner archipelago to the outer areas and the Baltic proper. Also
29 | the anoxic areas decrease by about 30% as a result of the lower deposition of organic material
30 | on the sea floor (not shown). The changes in the benthic pools of N and P occur over a longer
31 | time period, and the benthic P pool does not reach a steady state until about 40 years after the
32 | reduction.

Borttaget: scenario of the nutrient
land load

Borttaget: with

1 In the reduction scenario the transport of N to the open sea from the Stockholm Archipelago
2 decreases by 62% within four years (Table 4). The filter efficiency of N in the entire
3 archipelago increases at the same time from 79 % to 90 % as a result of the load reduction.
4 The longer response time of P compared to N is observed also in the filter efficiency (Fig. 14).
5 The filter efficiency of P at the end of the spin-up run is about 100 %. This implies that under
6 the 2010 conditions, all the P land load is retained in the Stockholm Archipelago when the
7 system is in steady state. This is not the case when the original model forcing is used, which
8 implies that the Stockholm archipelago is still adjusting to the load reductions already
9 implemented. Thus, the coastal region might under present conditions continue to improve
10 without further actions.

11 The filter efficiency of P decreases to 74 % during the first years after the reduction,
12 coinciding with the large decrease in the N pelagic pool and the decrease in N/P ratio. After
13 the initial decrease, the filter efficiency slowly increases to 106 % at the end of the simulation
14 period, i.e. retention is larger than the land and atmospheric load of P. As a consequence the
15 export of P from the archipelago to the Baltic proper decreases with time, and about 18 years
16 after the load reduction the direction of the transport changes. This coincides with the time
17 when the filter efficiency again reached 100 %. Thereafter the archipelago begins to import P
18 from the open sea. Thus, with the contemporary boundary conditions used at the open sea, P
19 from the Baltic proper is retained within the archipelago. For coastal management this
20 indicates the importance of the open sea nutrient conditions when effects of load reductions
21 are evaluated.

Borttaget: becomes

22 These results indicate that local nutrient load abatements can improve the environmental state
23 of a semi-enclosed coastal site (the inner archipelago) that is locally impacted by humans. The
24 results also imply that for the first 5-15 years, increased nutrient concentrations might be
25 expected locally. However, this effect largely depends on the water residence time and on
26 which nutrient limits the seasonal phytoplankton production initially. However, for the more
27 open coastal zone, represented in the present study by the intermediate and outer archipelago,
28 the response to further nutrient load reductions was minor. This exemplifies that for open
29 coastal areas the interactions between the open sea and the coastal zone is probably more
30 important than the land-sea connection.

Borttaget: From t

Borttaget: experiments

Borttaget: it seems

Borttaget: be expected, but the effect of this will largely depend on which nutrient that limit the seasonal phytoplankton production initially, and also the water residence time.¶

Borttaget: many

Borttaget: nutrient cycling in the

Borttaget: For the Baltic Sea this could mean that continued cooperation between the Baltic Sea countries will be an important factor for the environmental development of many coastal sites. Some of the coastal areas that are more open and influenced by the open water properties may have a less problematic environmental state to begin with, mainly due to better ventilation of deeper waters, and also less anthropogenic interference. Some other locations may be affected e.g. by large blooms of harmful algae that thrive in nutrient rich waters.¶

31 The present study can conclude that even the eutrophicated Stockholm Archipelago can, after
32 further nutrient load abatements, act as a sink for open water phosphorous. Similar behaviour

1 was found in the Chesapeake Bay (Boynton et al. 1995), which acts as a sink for the total load
2 of P, thus, P from land, atmosphere and from the open sea.

3 *Table 4.*

4 *Fig. 14.*

1 4 Conclusion

2 Archipelagos are complex areas with many basins and several shallow sounds, which affect
3 the transport of water and the dissolved and particulate nutrients. For the first time the SCM
4 model was used to study the capacity of the coastal filter of nutrients. An evaluation showed
5 that overall, model results agree with observations.

6 We focused our study in the northern Baltic proper and investigated retention of N and P in
7 the Stockholm Archipelago. The main findings are described below.

Borttaget: to

- 8 • The coastal zone works as an efficient filter for the land loads of nutrients. Under
9 prevailing conditions the total retention are 65 % and 72 % of P and N, respectively,
10 supplied from land,
- 11 • A sensitivity experiment reducing the land load of nutrients showed that the retention
12 capacity of N and P increased. In this case the export of N from the archipelago
13 decreased and P was imported from the open sea.
- 14 • The average filter efficiency is dependent on the spatial dimensions of the coastal area.
15 Thus, nutrient retention per area is largest in the inner archipelago and decreases
16 towards the open sea.
- 17 • Average water depth and water residence time regulate the retention of nutrients that
18 occurs mostly in the sediment due to processes such as burial and denitrification.
- 19 • The pools of nutrients in the water and in the sediment changes with nutrient loads on
20 different time scales and affects the temporal nutrient retention in the area. N has a
21 rather short response time of about three years while it takes about 40 years for P to
22 reach balance in a system with constant forcing. Changing N:P ratios in the
23 archipelago due to the different response time scales also have an impact on the
24 nutrient retention capacity on decadal time scales.
- 25 • Coastal management needs to take the aspects of time and balance between nutrient
26 loads and pools into account in the assessment of impacts from nutrient load
27 abatements. On shorter timescales the retention capacity of P might be less effective
28 when the nutrient load from land decreases.

Borttaget: , respectively, are retained

1 5 Acknowledgement

2 The research presented in this study is part of the Baltic Earth programme (Earth System
3 Science for the Baltic Sea region, see <http://www.baltex-research.eu/balticearth>) and is part of
4 the BONUS COCOA (Nutrient COcktails in COAstal zones of the Baltic Sea) project which
5 has received funding from BONUS, the joint Baltic Sea research and development
6 programme (Art 185), funded jointly from the European Union's Seventh Programme for
7 research, technological development and demonstration and from the Swedish Research
8 Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS), grant no.
9 2013-2056. Additional funding came from the EU Water Framework Directive programme at
10 the Swedish Meteorological and Hydrological Institute. We would like to thank Kerstin Rosén
11 Nilsson at the County Administrative Board of Stockholm for interesting discussions and
12 good advice. We are also grateful to the reviewers and the editor for their good comments and
13 suggestions to improve earlier versions of the manuscript.

Borttaget: also

14

6 References

- Ahlgren, I., Frisk, T., and Kamp-Nielsen, L.: Empirical and theoretical models of phosphorus loading, retention and concentration vs. lake trophic state. In: Phosphorus in Freshwater Ecosystems, Springer, 1988.
- Almroth-Rosell, E., Eilola, K., Hordoir, R., Meier, H. E. M., and Hall, P. O. J.: Transport of fresh and resuspended particulate organic material in the Baltic Sea — a model study, *J. Mar. Sys.*, 87, 1-12, 2011.
- Almroth-Rosell, E., Eilola, K., Kuznetsov, I., Hall, P. O. J., and Meier, H. E. M.: A new approach to model oxygen dependent benthic phosphate fluxes in the Baltic Sea, *J. Mar. Sys.*, 144, 127-141, 2015.
- Areskoug, H.: Nedfall av kväve och fosfor till Sverige, Östersjön och Västerhavet., Rapport 4148, 1993.
- Balls, P. W.: Nutrient Inputs to Estuaries from Nine Scottish East Coast Rivers; Influence of Estuarine Processes on Inputs to the North Sea, *Estuar. Coast Shelf S.*, 39, 329-352, 1994.
- [Baltic Environmental Database](http://www.balticnest.org/bed): <http://www.balticnest.org/bed>, 2015.
- Bergström, S., Alexandersson, H., Carlsson, B., Josefsson, W., Karlsson, K.-G., and Westring, G.: Climate and Hydrology of the Baltic Sea. In: A System Analysis of the Baltic Sea, Wulff, F., Rahm, L., and Larsson, P. (Eds.), Ecological Studies, Springer-Verlag, Berlin-Heidelberg, 2001.
- Billen, G., Silvestre, M., Grizzetti, B., Leip, A., Garnier, J., Voss, M., Howarth, R., Bouraoui, F., Lepistö, A., and Kortelainen, P.: Nitrogen flows from European regional watersheds to coastal marine waters, European nitrogen assessment, 271-297, 2011.
- Bolin, B. and Rodhe, H.: A note on the concepts of age distribution and transit time in natural reservoirs, *Tellus*, 25, 58-62, 1973.
- Boynton, W., Garber, J., Summers, R., and Kemp, W.: Inputs, transformations, and transport of nitrogen and phosphorus in Chesapeake Bay and selected tributaries, *Estuaries*, 18, 285-314, 1995.
- Brattberg, G.: Decreased phosphorus loading changes phytoplankton composition and biomass in the Stockholm archipelago, *Vatten*, 42, 1-153, 1986.

Borttaget: urnal

Borttaget: of Marine Systems

Borttaget: Journal of Marine Systems

Borttaget: inc,

Borttaget: al and

Borttaget:

Borttaget: cience

Borttaget: alticnest/thenestsystem/bed
balticenvironmentaldatabase.4.74cf9d041
3b817d9359e0.html

1 Carstensen, J., Conley, D., Bonsdorff, E., Gustafsson, B., Hietanen, S., Janas, U., Jilbert, T.,
2 Maximov, A., Norkko, A., Norkko, J., Reed, D., Slomp, C., Timmermann, K., and
3 Voss, M.: Hypoxia in the Baltic Sea: Biogeochemical Cycles, Benthic Fauna, and
4 Management, *AMBIO*, 43, 26-36, 2014.

5 Conley, D., Bjorck, S., Bonsdorff, E., Carstensen, J., Destouni, G., and Gustafsson, B.:
6 Hypoxia-Related Processes in the Baltic Sea, *Environ. Sci. Technol.*, 43, 3412-3420,
7 2009.

8 Diaz, R. J. and Rosenberg, R.: Spreading dead zones and consequences for marine
9 ecosystems, *Science*, 321, 926-929, 2008.

10 Duarte, C. M. and Cebrián, J.: The fate of marine autotrophic production, *Limnol. Oceanogr.*,
11 41, 1758-1766, 1996.

12 Edman, M. and Omstedt, A.: Modeling the dissolved CO₂ system in the redox environment of
13 the Baltic Sea, *Limnol. Oceanogr.*, 58, 74-92, 2013.

14 Edman, M. K. and Anderson, L. G.: Effect on pCO₂ by phytoplankton uptake of dissolved
15 organic nutrients in the Central and Northern Baltic Sea, a model study, *J. Mar. Sys.*,
16 139, 166-182, 2014.

17 Eilola, K., Almroth-Rosell, E., and Meier, H. E. M.: Impact of saltwater inflows on
18 phosphorus cycling and eutrophication in the Baltic Sea. A 3D model study., *Tellus A*,
19 66, 23985, 2014.

20 Eilola, K., Gustafsson, B. G., Kuznetsov, I., Meier, H. E. M., Neumann, T., and Savchuk, O.
21 P.: Evaluation of biogeochemical cycles in an ensemble of three state-of-the-art
22 numerical models of the Baltic Sea, *J. Mar. Sys.*, 88, 267-284, 2011.

23 Eilola, K., Meier, M. H. E., and Almroth, E.: On the dynamics of oxygen, phosphorus and
24 cyanobacteria in the Baltic Sea; A model study, *J. Mar. Sys.*, 75, 163-184, 2009.

25 Finlay, J. C., Small, G. E., and Sterner, R. W.: Human influences on nitrogen removal in
26 lakes, *Science*, 342, 247-250, 2013.

27 Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., and Haidvogel, D.: Nitrogen cycling
28 in the Mid Atlantic Bight and implications for the North Atlantic nitrogen budget:
29 Results from a three-dimensional model, *Global Biogeochem. Cycles*, 20, 2006.

30 Fonselius, S. H.: Hydrography of the Baltic deep basins III, 1969.

Borttaget: mental s

Borttaget: ence

Borttaget: &

Borttaget: t

Borttaget: ogy

Borttaget: s

Borttaget: ogy and

Borttaget: aphy

Borttaget: Limnology and Oceanography

Borttaget: Journal of Marine Systems

Borttaget: Journal of Marine Systems

Borttaget: Journal of Marine Systems

- 1 Galloway, J. N., Townsend, A. R., Erismann, J. W., Bekunda, M., Cai, Z., Freney, J. R.,
2 Martinelli, L. A., Seitzinger, S. P., and Sutton, M. A.: Transformation of the nitrogen
3 cycle: recent trends, questions, and potential solutions, *Science*, 320, 889-892, 2008.
- 4 Grimvall, A., von Brömssen, C., and Lindström, G.: Using process-based models to filter out
5 natural variability in observed concentrations of nitrogen and phosphorus in river water,
6 *Environ. Monit. Assess.*, 186, 5135-5152, 2014.
- 7 Gustafsson, B., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B.,
8 Neumann, T., Ruoho-Airola, T., Savchuk, O., and Zorita, E.: Reconstructing the
9 Development of Baltic Sea Eutrophication 1850–2006, *AMBIO*, 41, 534-548, 2012.
- 10 Hansson, M. and Andersson, L.: Oxygen Survey in the Baltic Sea 2013 - Extent of Anoxia
11 and Hypoxia, 1960-2013, Swedish Meteorological and Hydrological Institute,
12 Norrköping, Sweden, REPORT OCEANOGRAPHY 50, 22 pp., 2014.
- 13 HaV: Havs och Vattenmyndighetens föreskrifter om klassificering och miljö kvalitetsnormer
14 avseende ytvatten, HVMFS 2013:19, 2013.
- 15 Hayn, M., Howarth, R., Marino, R., Ganju, N., Berg, P., Foreman, K. H., Giblin, A. E., and
16 McGlathery, K.: Exchange of nitrogen and phosphorus between a shallow lagoon and
17 coastal waters, *Estuar. Coast.*, 37, 63-73, 2014.
- 18 Hejzlar, J., Anthony, S., Arheimer, B., Behrendt, H., Bouraoui, F., Grizzetti, B., Groenendijk,
19 P., Jeuken, M., Johnsson, H., and Porto, A. L.: Nitrogen and phosphorus retention in
20 surface waters: an inter-comparison of predictions by catchment models of different
21 complexity, *J. Environ. Monitor.*, 11, 584-593, 2009.
- 22 HELCOM: Ecosystem Health of the Baltic Sea 2003–2007: HELCOM Initial Holistic
23 Assessment., 2010.
- 24 Humborg, C., Danielsson, Å., Sjöberg, B., and Green, M.: Nutrient land–sea fluxes in
25 oligotrophic and pristine estuaries of the Gulf of Bothnia, Baltic Sea, *Estuar. Coast.*
26 *Shelf. S.*, 56, 781-793, 2003.
- 27 Johnston, C. A.: Sediment and nutrient retention by freshwater wetlands: effects on surface
28 water quality, *Crit. Rev. Env. Sci. Tec.*, 21, 491-565, 1991.
- 29 Jonsson, P., Persson, J., and Holmberg, P.: Skärgårdens bottnar, Naturvårdsverket rapport,
30 5212, 2003.

Borttaget: Environmental

Borttaget: oring and

Borttaget: ment

Borttaget: Estuaries and Coasts

Borttaget: Journal of Environmental
Monitoring

Borttaget: Estuarine, Coastal and Shelf
Science

Borttaget: Critical Reviews in
Environmental Science and Technology

- 1 Karlsson, O. M., Jonsson, P. O., Lindgren, D., Malmaeus, J. M., and Stehn, A.: Indications of
2 recovery from hypoxia in the inner Stockholm archipelago, *Ambio*, 39, 486-495, 2010.
- 3 Kemp, W., Testa, J., Conley, D., Gilbert, D., and Hagy, J.: Temporal responses of coastal
4 hypoxia to nutrient loading and physical controls, *Biogeosciences*, 6, 2985-3008, 2009.
- 5 Knudsen, M.: Ein hydrographischer lehrsatz, *Annalen der Hydrographie und Maritimen*
6 *Meteorologie*, 28, 316-320, 1900.
- 7 Lindh, G.: Miljörapport 2013, Stockholm Vatten VA AB, Stockholm Vatten, 2013.
- 8 Lindström, G., Pers, C., Rosberg, J., Strömqvist, J., and Arheimer, B.: Development and
9 testing of the HYPE (Hydrological Predictions for the Environment) water quality
10 model for different spatial scales, *Hydrol. Res.*, 41, 295-319, 2010.
- 11 Lücke, J. : Stockholms recipienter. Påverkan av Stockholms framtida avloppsrening.
12 Stockholm Vatten AB, 2014.
- 13 Lücke, J.: Undersökningar i Stockholms skärgård 2014. Vattenkemi och växtplankton, och
14 bottenfauna. Stockholm Vatten AB, 2015.
- 15 Lännergren, C.: Undersökningar i Östra Mälaren till och med 2009, Stockholm Vatten, 2010.
- 16 Lännergren, C., Eriksson, B., and Stehn, A.: Surveys in the Stockholm archipelago 2008,
17 Stockholm Vatten Report, Diary, No: 09SV139, 243, 2009.
- 18 Marmefelt, E., Arheimer, B., and Langner, J.: An integrated biochemical model system for the
19 Baltic Sea, *Hydrobiologia*, 393, 45-56, 1999.
- 20 McGlathery, K. J., Sundbäck, K., and Anderson, I. C.: Eutrophication in shallow coastal bays
21 and lagoons: the role of plants in the coastal filter, *Mar. Ecol-Prog. Ser.*, 348, 1-18,
22 2007.
- 23 Meier, H. E. M., Müller-Karulis, B., Andersson, H., Dieterich, C., Eilola, K., Gustafsson, B.,
24 Höglund, A., Hordoir, R., Kuznetsov, I., Neumann, T., Ranjbar, Z., Savchuk, O., and
25 Schimanke, S.: Impact of Climate Change on Ecological Quality Indicators and
26 Biogeochemical Fluxes in the Baltic Sea: A Multi-Model Ensemble Study, *AMBIO*, 41,
27 558-573, 2012.
- 28 Meier, M., Eilola, K., and Almroth, E.: Climate-related changes in marine ecosystems
29 simulated with a 3-dimensional coupled physical-biogeochemical model of the Baltic
30 Sea, *Clim. Res.*, 48, 31-55, 2011.

Borttaget: Hydrology research

Borttaget: Marine Ecology Progress Series

Borttaget: ate

Borttaget: earch (CR)

1 Monsen, N. E., Cloern, J. E., Lucas, L. V., and Monismith, S. G.: A comment on the use of
2 flushing time, residence time, and age as transport time scales, *Limnol. Oceanogr.*, 47,
3 1545-1553, 2002.

Borttaget: o

4 Mortimer, C. H.: The Exchange of Dissolved Substances Between Mud and Water in Lakes,
5 *J. Ecol.*, 29, 280-329, 1941.

Borttaget: Journal of Ecology

6 Naturvårdsverket: Status, potential och kvalitetskrav för sjöar, vattendrag, kustvatten och
7 vatten i övergångszon. En handbok om hur kvalitetskrav i ytvattenförekomster kan
8 bestämmas och följas upp, 2007.

9 Nielsen, K., Risgaard-Petersen, N., Sømod, B., Rysgaard, S., and Bergø, T.: Nitrogen and
10 phosphorus retention estimated independently by flux measurements and dynamic
11 modelling in the estuary, Randers Fjord, Denmark, *Mar. Ecol-Prog. Ser.*, 219, 25-40,
12 2001.

Borttaget: Marine Ecology Progress Series

13 Nixon, S. W., Ammerman, J. W., Atkinson, L. P., Berounsky, V. M., Billen, G., Boicourt, W.
14 C., Boynton, W. R., Church, T. M., Ditoro, D. M., Elmgren, R., Garber, J. H., Giblin, A.
15 E., Jahnke, R. A., Owens, N. J. P., Pilson, M. E. Q., and Seitzinger, S. P.: The fate of
16 nitrogen and phosphorus at the land-sea margin of the North Atlantic Ocean. In:
17 Nitrogen Cycling in the North Atlantic Ocean and its Watersheds, Howarth, R. (Ed.),
18 Springer Netherlands, 1996.

19 Norkko, J., Reed, D. C., Timmermann, K., Norkko, A., Gustafsson, B. G., Bonsdorff, E.,
20 Slomp, C. P., Carstensen, J., and Conley, D. J.: A welcome can of worms? Hypoxia
21 mitigation by an invasive species, *Global Change Biol.*, 18, 422-434, 2012.

Borttaget: ogy

22 Omstedt, A.: Guide to process based modeling of lakes and coastal seas, Springer Praxis-
23 Books in Geophysical Sciences, 2015.

24 Oeschle, A., J. Blackford, S. C. Doney, and M. Gehlen.: Modelling considerations, In U.
25 Riebesell, V. J. Fabry, L. Hansson, and J.-P. Gattuso [eds.], Guide to best practices for
26 ocean acidification research and data reporting., Publications Office of the European
27 Union., 233-242, 2010.

28 Rabalais, N. N., Turner, R. E., and Wiseman Jr, W. J.: Gulf of Mexico hypoxia, AKA "The
29 dead zone", *Annu. Rev. Ecol. Syst.*, 2002. 235-263, 2002.

Borttaget: Annual Review of ecology and Systematics

30 Robertson, L., Langner, J., and Engardt, M.: An Eulerian limited-area atmospheric transport
31 model, *J. App. Meteorol.*, 38, 190-210, 1999.

Borttaget: Journal of Applied Meteorology

- 1 Rosenberg, R. and Diaz, R. J.: Sulfur bacteria (*Beggiatoa* spp.) mats indicate hypoxic
2 conditions in the inner Stockholm Archipelago, *Ambio*, 22, 32-36, 1993.
- 3 Sahlberg, J., Marmefelt, E., Brandt, M., Hjerdt, N., and Lundholm, K.: HOME - Vatten i
4 Norra Östersjöns vattendistrikt, Integrerat modellsystem för vattenkvalitetsberäkningar,
5 Swedish Meteorological and Hydrological Institute, Oceanografi, SMHI reports 93,
6 2008.
- 7 Sanders, R., Jickells, T., Malcolm, S., Brown, J., Kirkwood, D., Reeve, A., Taylor, J.,
8 Horrobin, T., and Ashcroft, C.: Nutrient fluxes through the Humber estuary, *J. Sea*
9 *Res.*, 37, 3-23, 1997.
- 10 Sheldon, J. E. and Alber, M.: The calculation of estuarine turnover times using freshwater
11 fraction and tidal prism models: a critical evaluation, *Estuar. Coast.*, 29, 133-146, 2006.
- 12 Seitzinger, S. P. and Giblin, A. E.: Estimating denitrification in North Atlantic continental
13 shelf sediments, *Biogeochemistry*, 35, 235-260, 1996.
- 14 Soetaert, K., Middelburg, J. J., Herman, P. M., and Buis, K.: On the coupling of benthic and
15 pelagic biogeochemical models, *Earth-Sci. Rev.*, 51, 173-201, 2000.
- 16 Strömqvist, J., Arheimer, B., Dahné, J., Donnelly, C., and Lindström, G.: Water and nutrient
17 predictions in ungauged basins: set-up and evaluation of a model at the national scale,
18 *Hydrolog. Sci. J.*, 57, 229-247, 2012.
- 19 Svensson, U.: PROBE An instruction manual, SMHI, 1998.
- 20 Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P.,
21 Tamminen, T., Viitasalo, M., Voss, M., and Wasmund, N.: Internal ecosystem
22 feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management
23 in the Baltic Sea, *AMBIO*, 36, 186-194, 2007.
- 24 Viktorsson, L., Almroth-Rosell, E., Tengberg, A., Vankevich, R., Neelov, I., Isaev, A.,
25 Kravtsov, V., and Hall, P. O. J.: Benthic Phosphorus Dynamics in the Gulf of Finland,
26 Baltic Sea, *Aquat. Geochem.*, 18, 543-564, 2012.
- 27 Voss, M., Emeis, K., Hille, S., Neumann, T., and Dippner, J.: Nitrogen cycle of the Baltic Sea
28 from an isotopic perspective, *Global Biogeochem. Cy.*, 19, GB3001, 2005.

Borttaget: Journal of Sea Research

Borttaget: s

Borttaget: r

Borttaget: Hydrological Sciences
Journal

Borttaget: : A journal of the Human
Environment

Borttaget: Global biogeochemical
cycles

1 | [Xue, Z., He, R., Fennel, K., Cai, W., Lohrenz, S., and Hopkinson, C.: Modeling ocean](#)
2 | [circulation and biogeochemical variability in the Gulf of Mexico, Biogeosciences, 10,](#)
3 | [7219-7234, 2013.](#)
4 |
5 |

7 Figures

Flyttad (infogning) [1]




Fig. 1.

Borttaget: 4




Fig. 2

Borttaget: 1




Fig. 3

Borttaget: 2




Fig. 4

Borttaget: 3




Fig. 5.

Flyttad uppåt [1]: ¶

¶

Point sources

River load

Perr

Ret

Fig. 4.¶

Borttaget:Sidbrytning.....

¶




Fig. 6.




Fig. 7.




Fig. 8.





Fig. 9.






Fig. 10.




Fig. 11.




Fig. 12.





Fig. 13.







Fig. 14



Borttaqet:

8 Figure captions

Fig. 1. Simplified scheme of the retention calculations in the study area. Permanent retention is considered as a permanent removal of nutrients from the ecological system and includes burial and for nitrogen also denitrification. Temporary retention is defined as the changes in nutrient inventory in the active sediment layer and water column. The temporary retention may change sign depending on whether the nutrient inventory increases or declines.

Borttaget: 4

Borttaget: calculation scheme of retention in the study area.

Fig. 2. The Swedish Coastal zone Model can be used in different areas along the Swedish coast stretching from the Norwegian border in the West to the Finnish border in the North (different colours, left). In the present study the SCM model covers the northern Baltic proper (marked with a red square) and has been used to estimate the coastal filter efficiency of nutrients in the Stockholm inner (red), intermediate (orange) and outer (blue) archipelagos (right). The outlet of river Norrström is marked with a black arrow and the different basins are shown by the black contours.

Borttaget: 1

Fig. 3. Schematic figure of the Swedish Coastal and Biogeochemical model, SCOB. Oxygen and hydrogen sulphide are simplified for clarity.

Borttaget: 2

Fig. 4. Available locations with observations (circles and dots) in the Stockholm Archipelago. Model evaluation of temperature, salinity, DIN, DIP and bottom water oxygen concentration was performed at selected stations (circles marked with letters), which are described in Table 1.

Borttaget: 3

Fig. 5. Observed (stars) and modelled (line) monthly outflow (Q_F) and nutrient loads from Lake Mälaren, through Norrström to basin Strömmen for the modelled period (1990-2012). DIN is the sum of nitrate and ammonium.

Fig. 6. Average cost function (C) and correlation coefficients, adjusted ($1-r$) to the range 0-1, for an overview of the model skill at the eight different validation sites (A-G). The individual skills of the different parameters, average seasonal variation (black) and/or the vertical summer (red) and winter (turquoise) profiles of temperature (T), DIN, DIP and oxygen concentrations (O_2) are shown, as well as the combined model skills for all variables (purple cross). Variables within the inner quarter circle and between the two quarter circles are considered to be good and acceptable, respectively, while variables that are outside the quarter circles are not well simulated.

Fig. 7. The SCM modelled (lines) and observed (circle and diamond) vertical average profiles (1990-2012) of salinity (a) and concentrations of oxygen (O_2 ; b), DIN (c) and DIP (d) in the basin Strömmen during winter (turquoise) and summer (red) months. Depth layers with dense number of observations (grey stars) determined the vertical depth intervals (grey shaded area) used in the profile calculations. The standard deviations (horizontal lines) were calculated for the summer and winter values of the observations.

Fig. 8. Simulated (lines) and observed averages (squares) of the seasonal variation and the standard deviation (vertical lines) of the observations in the basin Strömmen (1990-2012) of surface temperature (Temp), salinity, DIN and DIP and of the bottom water oxygen concentrations. Time periods with dense number of observations (grey stars) determined the time intervals (grey shaded area) used in the calculations.

Fig. 9. Transport scheme of N and P ($t\ yr^{-1}$) from land (leaning top boxes) and atmosphere (top boxes), and the net exchange from the inner, intermediate and outer archipelago (ellipse) towards the open sea. Total retention is the sum of temporary retention (square) and permanent retention (square with round corners). For P burial is the only process that leads to permanent retention, while for N also denitrification removes N. Negative values for the temporary retention means a decrease in the benthic and/or pelagic pools of nutrients.

Fig.10. The external annual load and retention ($t\ yr^{-1}$) of P (a, c) and N (b, d) in the entire Stockholm Archipelago for the period 1990-2012. Total load (shaded area) and the contributions from the different sources; rivers and land run off (diamonds), point sources (circles) and atmosphere (solid line) is shown on the top row. The total retention (shaded area) as a sum of permanent retention (solid line) and temporary retention (diamonds) (c, d) is shown on the bottom row.

Fig.11. The retention per area unit ($t\ km^{-2}\ yr^{-1}$) of P (left) and N (right) in each basin of the Stockholm Archipelago.

Fig.12. The total content ($g\ m^{-2}$) of the pelagic (top) and benthic (bottom) P (left) and N (right) in the inner (diamonds), intermediate (circles), outer (triangles) and entire (black line) Stockholm archipelagos.

Fig.13. The filter efficiency of P (left) and N (right) versus the logarithmic ratio between the average depth and the freshwater residence time of the study areas ($month\ yr^{-1}$). Data from other studies are from Billen et al. (2011), Hayn et al. (2014), Nielsen et al. (2001) and Nixon et al. (1996). The straight line shows the logarithmic regression for the data from Nixon et al. (1996).

Formaterat: Svenska (Sverige)

Fig.14. Pelagic (upper) and benthic (middle) pools of P (left) and N (right) in the inner (red), intermediate (orange), outer (turquoise) and entire (black) Stockholm Archipelago. The filter efficiencies (%) of N (red) and P (blue) load from land and atmosphere are shown for the entire Stockholm Archipelago (lower), where the small peaks derive from leap years.

Borttaget: the nutrient

9 Tables

Table 1. Number of sampling occasions (Occ) during the number of years, number of months during each year, and number of depths levels that was frequently sampled at the different stations used for validation of model results. The position of the stations can be seen in [Fig. 3](#).

Borttaget: Fig. 3

| ID | Station name | Basin name | Occ | Years* | Months | Depths** |
|----|--------------------|--------------------|-----|--------|--------|----------|
| A | Nyvarp | Sandöfjärden | 209 | 23 | 8 | 14 |
| B | Kanholmsfjärden | Kanholmsfjärden | 206 | 23 | 9 | 13 |
| C | Solöfjärden | Solöfjärden | 213 | 23 | 8 | 14 |
| D | TrälhavetII | Trälhavet | 215 | 23 | 9 | 13 |
| E | S. Vaxholmsfjärden | S. Vaxholmsfjärden | 131 | 23 | 7 | 8 |
| F | Blomskär | Stora Värtan | 141 | 23 | 8 | 9 |
| G | Blockhusudden | Strömmen | 249 | 23 | 11 | 16 |
| H | Baggensfjärden | Baggensfjärden | 173 | 20 | 9 | 10 |

* Entire period is 23 years; **Sampled at least half of the sample occasions

Table 2. The maximum concentrations of P and N (mg l^{-1}) in the discharge from sewage treatment plants of different size (person equivalents, pe)

| Sewage treatment facilities (pe) | P (mg l^{-1}) | N (mg l^{-1}) |
|----------------------------------|--------------------------|--------------------------|
| >50 000 | 0.1 | 4 |
| 10 000-50 000 | 0.1 | 6 |
| <10 000 | 0.15 | 10 |

Borttaget: and the reduction of P and N loads from rivers and industries (%) in the scenario run with the SCM

Borttaget: Decrease in land load

Table 3. The correlation coefficients (r) between observations (obs) and model results (S-HYPE), and the long term (1990-2012) averages of river outflow (Q_F) and nutrient loads from Lake Mälaren.

| Variable | Units | Average obs | Average S-HYPE | r |
|----------|---------------------------------------|-------------|----------------|------|
| Q_F | $10^6 \text{ m}^3 \text{ month}^{-1}$ | 421 | 422 | 0.94 |
| TN | t month^{-1} | 270 | 271 | 0.93 |
| DIN | t month^{-1} | 83 | 76 | 0.86 |
| TP | t month^{-1} | 13 | 11 | 0.87 |
| DIP | t month^{-1} | 5.7 | 5.7 | 0.79 |

Table 4. The total land load (rivers, land run-off and atmosphere) of P and N ($t\ yr^{-1}$) to the Stockholm Archipelago, the size of the benthic and pelagic N and P pools (t), the export from the area ($t\ yr^{-1}$) and the filter efficiency (F_{eff}) before and after the nutrient reductions, as well as their percentage changes. The system is in both cases in steady state, thus the benthic and pelagic pools are in balance with the nutrient load.

| | Unit | Initial values | End of period | Change (%) | |
|--------------|-------------|----------------|---------------|------------|------|
| P | Total load | $t\ yr^{-1}$ | 213 | 186 | -13 |
| | Pool* | T | 10661 | 9952 | -7 |
| | Export | $t\ yr^{-1}$ | -3.5 | -11 | -207 |
| | F_{eff} | % | 101 | 106 | |
| N | Total load | $t\ yr^{-1}$ | 7690 | 6164 | -20 |
| | Pool* | T | 35196 | 33216 | -6 |
| | Export | $t\ yr^{-1}$ | 1585 | 609 | -62 |
| | F_{eff} | % | 79 | 90 | |
| N:P** | Molar ratio | | 42 | 35 | -16 |

*The sum of benthic and pelagic pools. ** In the inner archipelago.

