

Increased temperature causes different carbon and nitrogen processing patterns in two common intertidal foraminifera (*Ammonia tepida* and *Haynesina germanica*)

Julia Wukovits¹, Annekatrin Julie Enge¹, Wolfgang Wanek², Margarete Watzka², and Petra Heinz¹

¹University of Vienna, Department of Palaeontology, Vienna, Austria

²University of Vienna, Department of Microbiology and Ecosystem Science, Terrestrial Ecosystem Research, Vienna, Austria

Correspondence to: Julia Wukovits (julia.wukovits@univie.ac.at)

Abstract. Benthic foraminifera are highly abundant heterotrophic protists in marine sediments, but future environmental changes will challenge the tolerance limits of intertidal species. Metabolic rates and physiological processes in foraminifera are strongly depending on environmental temperatures. Temperature related stress could therefore impact foraminiferal food source processing efficiency and might result in altered nutrient fluxes through the intertidal food web. In this study, we performed a laboratory feeding experiment on *Ammonia tepida* and *Haynesina germanica*, two dominant foraminiferal species of the German Wadden Sea/Friedrichskoog, to test the effect of temperature on phytodetritus retention. The specimens were fed with ¹³C and ¹⁵N labelled freeze dried *Dunaliella tertiolecta* (green algae) at the start of the experiment and were incubated at 20°C, 25°C, and 30°C respectively. Dual labelling was applied to observe potential temperature effects on the relation of phytodetrital carbon and nitrogen retention. Samples were taken over a period of two weeks. Foraminiferal cytoplasm was isotopically analysed to investigate differences in carbon and nitrogen uptake derived from the food source. Both species showed a positive response to the provided food source, but carbon uptake rates of *A. tepida* were 10-fold higher compared to those of *H. germanica*. Increased temperatures had a far stronger impact on carbon uptake of *H. germanica* than on *A. tepida*. A distinct increase in levels of phytodetrital derived nitrogen (compared to more steady carbon levels) could be observed over the course of the experiment in both species. The results suggest that higher temperatures have a significant negative effect on the carbon exploitation of *H. germanica*. For *A. tepida*, higher carbon uptake rates and the enhanced tolerance range for higher temperatures could outline an advantage in warmer periods, if the main food source consists of chlorophyte phytodetritus. These conditions are likely to impact nutrient fluxes in *A. tepida*/*H. germanica* associations.

1 Introduction

The intertidal is an extreme environment, exposed to intense seasonal and diurnal fluctuations in temperature, challenging the physiological limits of benthic organisms, for instance foraminifera. Foraminifera are marine heterotrophic protists with a common worldwide occurrence in extant and fossil communities. Future environmental changes are expected to effect coastal foraminiferal communities and assemblage structures (Schafer et al., 1996; Culver and Buzas, 1995) since some intertidal species exhibit a fast response to rapid changes in their environment (e.g. warming). Temperature affects physiological per-

formances, resulting in a lack of fitness altering community structures and leading to shifts in nutrient fluxes and ecosystem balance (Brown et al., 2004; Allen et al., 2005; Petchey et al., 2010; O'Connor et al., 2009; Yvon-Durocher et al., 2010). In intertidal mudflats, smaller benthic foraminifera contribute up to 80% of the protist biomass (Lei et al., 2014) and are an important component of the food web (Lipps and Valentine, 1970; Buzas, 1978; Buzas and Carle, 1979; Nomaki et al., 2008).

5 But the allocation of their trophic role still lacks defined considerations. Due to their high abundances and substantial carbon incorporation, it is assumed that they play a major role in the carbon cycle of these environments (Moodley et al., 2000). *Haynesina germanica* and *Ammonia tepida* often co-occur with high abundances in intertidal sediments of the temperate zone. The dominance of these two species shows seasonal fluctuations and can be location-specific (Murray, 2014; Alve and Murray, 1994; Debenay et al., 2006; du Chatelet et al., 2009). Factors controlling these community shifts still need to be specified and 10 might include differential sensitivity to general disturbances, differences in food preferences, or variations in physiological limits to physical stress.

Temperature has been proven to play a major role in reproduction, growth and respiration rates of intertidal foraminifera (Bradshaw, 1955, 1957, 1961; Lee and Muller, 1973; Haynert and Schönfeld, 2014; Cesbron et al., 2016). Elevated temperatures around 35°C were reported to increase the expression of stress proteins in *A. tepida* (Heinz et al., 2012) and represent the range 15 of minimum respiratory activity, which subsequently drops until the lethal point of 45°C (Bradshaw, 1961). There is still a lack of data concerning physiological effects of temperature on *H. germanica*, though there is evidence about lower environmental temperatures (12°C) being most supportive for its reproduction. (Goldstein and Alve, 2011). However, possible restraints or advances of temperature effects on phytodetritus uptake have not been evaluated yet. Foraminiferal food sources include microalgae, phytodetritus or bacteria (Muller and Lee, 1969; Lee et al., 1966; Goldstein and Corliss, 1994), which are prevalent 20 elements of the intertidal particulate organic matter (POM) pool. Distributions of POM sources or microalgae groups are often used to correlate foraminiferal abundances and to relate the availability of different POM sources with population dynamics or food preferences (Alve and Murray, 1994, 2001; du Chatelet et al., 2009; Hayward et al., 1996; Murray and Alve, 2000; Ward et al., 2003; Topping et al., 2006; de Nooijer, 2007; Diz and Francés, 2008; Goineau et al., 2012; Papaspyrou et al., 2013; Melis and Covelli, 2013). Microalgae are proposed to be the preferred food source of *A. tepida*, in particular over bacteria (Pascal 25 et al., 2008b), while *H. germanica* possesses a mechanism to efficiently feed on diatoms (Austin et al., 2005). Foraminifera accumulate their food with a pseudopodial network to ingest and transport food particles to the endoplasm (Goldstein and Corliss, 1994; Bowser et al., 1992), which is often protected by an inorganic (e.g. calcareous) shell. Their ingestion rates of algae or phytodetritus are comparable to bacterial assimilation rates of detrital carbon (Moodley et al., 2000, 2002).

Although abundant data sources exist about food derived carbon in foraminiferal cytoplasm, information about nitrogen re- 30 mains scarce. Research about coupling of food derived carbon and nitrogen in foraminifera is limited to a few in situ study in the bathyal of the Arabian Sea, where foraminifera play an important role in benthic carbon fluxes (Enge et al., 2014, 2016). Tidal flats function as both an important source and sink for nutrients (Joye et al., 2009) and represent essential ecological components of earth's marine systems. The abundant intertidal primary producers like microalgae or diatoms control C and N flows of the sediments (Cook et al., 2004), with foraminifera being an important consumer of their biomass. Regarding the high 35 abundances of foraminifera in benthic communities, significant effects of temperature on their food processing could poten-

tially influence intertidal nutrient fluxes, in particular if they relate to foraminiferal carbon and nitrogen coupling. This study aims to investigate the effects of temperature on the food exploitation and on C and N cycling of *H. germanica* and *A. tepida*. The response to an artificially produced phytodetrital food source (*Dunaliella tertiolecta*, Chlorophyta) was tested under three temperature regimes. *Dunaliella tertiolecta* has been identified as a valuable food source for *A. tepida* (Pascal et al., 2008a), 5 but was not tested on *H. germanica* so far. This study also tests which species shows a more efficient response to chlorophyte detritus in a mesocosm setting. The laboratory feeding experiment was performed in incubation chambers, adjusted to 20°C, 25°C and 30°C. To obtain direct estimates of the uptake of carbon and nitrogen, phytodetritus was labelled with stable isotopes (^{13}C , ^{15}N). This method has provided important data on the in situ feeding behaviour of foraminifera in various environments (Moodley et al., 2002; Enge et al., 2016; Moodley et al., 2000; Middelburg et al., 2000; Witte et al., 2003; Nomaki et al., 10 2005, 2009, 2011; Sweetman et al., 2009; Enge et al., 2011; Jeffreys et al., 2013). The temperatures chosen for this experiment correspond to experimentally determined values that cover optimum or tolerance ranges of physiological processes in intertidal foraminifera (Bradshaw, 1957, 1961; Lee and Muller, 1973). Further, they lie in the range of seasonal and diurnal temperature amplitudes measured on intertidal surface sediments close to the sampling area (Al-Raei et al., 2009). Simulated variations 15 in temperature were assumed to influence the food uptake efficiency of the species, due to potential temperature stress. The amount of phytodetrital carbon (pC) and nitrogen (pN) uptake should reveal information about the nutrient processing potential of the two species. Simultaneous detection of both stable isotopes allows to determine the ratio in which pC relative to pN is retained in bulk foraminiferal cytoplasm over time. This helps to interpret phytodetrital uptake in relation with foraminiferal carbon and nitrogen coupling (Enge et al., 2016; Evrard et al., 2010; Hunter et al., 2012) and temperature influences might 20 show in imbalances of these ratios between treatments. Benthic foraminifera are used in the assessment of (paleo-) environmental data. Studies aiming to develop new foraminiferal proxies (e.g. for organic matter accumulation or physical parameters) generally apply statistical analysis to field surveys. There is however a comparably low amount of biological or ecological data available. This study offers an unique dataset on the effect of an altered environmental condition (temperature) on food resource 25 exploitation on the level of the cytoplasmatic balance of food derived carbon and nitrogen. The aim is to support definitions of the role of benthic foraminifera in intertidal carbon and nitrogen fluxes with respect to warming events and changes in POM availability.

2 Material and Methods

2.1 Sampling site & Material collection

Surface sediment was taken on 24th April 2014 during low tide in the intertidal mudflat of the German Wadden Sea near Friedrichskoog (Germany). Water temperature and salinity at the sampling site were 19.9°C and 31 (practical salinity units). 30 Sediment was collected and sieved at the sampling site through 500 μm and 63 μm meshes to remove larger meiofauna and organic particles. In the laboratory, samples were sieved again to obtain foraminiferal specimen in the size fraction of 125 - 355 μm . The sediment contained high abundances of *Ammonia tepida* and *Haynesina germanica* individuals, which were picked and collected for further processing. Living individuals were identified under the microscope regarding intact protoplasma and

particle accumulation around the aperture (Moodley et al., 2002; Nomaki et al., 2005; Moodley et al., 1997; Nomaki et al., 2006). Care was taken to achieve a homogenous size distribution within specimens in each replicate. Each species was kept separately in crystallising dishes and fed regularly with living *Dunaliella tertiolecta* until the start of the experiment.

2.1.1 Production of ^{13}C and ^{15}N enriched phytodetritus

5 At the start of the experiment, lyophilized powder of *Dunaliella tertiolecta* was used to simulate a phytodetritus pulse. The chlorophyte was grown in f/2 medium (Guillard and Ryther, 1962; Guillard, 1975) enriched with 98 atom % ^{13}C ($\text{NaH}^{13}\text{CO}_3$, Sigma-Aldrich) and 98 atom % ^{15}N ($\text{Na}^{15}\text{NO}_3$, Sigma-Aldrich) to final concentrations of 1.5 mmol L⁻¹ $\text{NaH}^{13}\text{CO}_3$ and 0.44 mmol L⁻¹ $\text{Na}^{15}\text{NO}_3$. Algal cultures were kept within incubators ($T = 20^\circ\text{C}$; dark: light = 16:8). After 20 days, they were harvested by centrifugation (800 G; 10 min) and rinsed three times in simplified artificial seawater (ASW, compare preparation 10 in Enge et al. (2011)) to remove unassimilated isotope tracer. The hereby obtained algal slurry was shock frozen with liquid nitrogen and lyophilized at -55°C and 0.180 mbar for 6 days.

2.1.2 Experimental setup

Separate time series for *A. tepida* and *H. germanica* were performed at three temperatures (20°C , 25°C , 30°C) in triplicate. Additionally, background samples of untreated specimens were taken to obtain natural abundance values for $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ 15 and initial cytoplasmatic total organic carbon (C) and nitrogen (N) content. Specimens were transferred into 72 experimental dishes (150 individuals of *A. tepida*/170 individuals of *H. germanica* per dish), each containing 280 mL modified synthetic seawater without sediment (SSW; (Kester et al., 1967; Dickson and Goeyt, 1994), adjusted to a pH of 8.10 and a salinity of 32. The dishes were incubated at a light: dark = 12:12 hrs cycle (type ST 2 POL-ECO Aparatura incubation chambers). After three days of acclimation, the labelled algal diet was added ($396.36 \pm 45.57 \text{ mg C m}^{-2}$; $40.26 \pm 9.43 \text{ mg N m}^{-2}$). No further 20 feed amendment was provided throughout the experiment. Algal particles were visible as a green layer on the bottom of the experimental vessels until the end of the experiment. Oxygen, salinity and pH were kept constant at optimum levels during incubation. Experimental dishes were sealed airtight and opened after two days to avoid hypoxia (measured O_2 at sampling days was $5 - 8 \text{ mg L}^{-1}$). Both species were subsampled to obtain data of phytodetritus processing on the 2nd, 4th, 7th and 14th day of the experiment.

25 2.1.3 Sample preparation

Foraminifera were removed from the experimental dishes and frozen at -20°C to stop metabolic activities. The foraminifera were cleaned from adhering particles with a hair brush. Further on, the organisms were carefully washed in simplified ASW. An amount of 50 (*A. tepida*) or 60 (*H. germanica*) individuals met the optimum range of 0.7 - 1.0 mg cytoplasmatic dry weight, necessary for isotope and elemental analysis. Only foraminifera meeting the criteria for live specimens as described 30 above were prepared for analysis. Figure 1 shows relative amounts of cytoplasm conditions and analysed individuals on the respective sampling days. Specimens were transferred to tin capsules, dried at 50°C , decalcified and dried in a final drying step




Figure 1. Relative amounts of individuals per sampling day and temperature treatment. The terms unusual and intact refer to the cytoplasmatic appearance of the specimen, additionally the fraction of analysed specimen is shown. Unusual cytoplasm includes individuals with patchy distribution of cytoplasm within the test or an unusual coloration. Individuals with intact cytoplasm includes former live specimens as described in the text.

for three days. All glassware used for preparation was combusted at 500°C for 5h, picking tools and tin capsules were cleaned in a solution of dichloromethane (CH_2Cl_2) and methanol (CH_4O) (1:1, v:v).

After opening the experimental dishes on the 2nd day, water samples (aliquots of 50 mL) were taken from the *H. germanica* series. They were transferred to 50 mL headspace vials. Some drops of HgCl_2 were added to stop respiratory activities and 5 biological production of CO_2 (Kroopnick et al., 1972). The vials were sealed airtight and stored at 4°C. To determine ^{13}C of the dissolved inorganic carbon (DIC) in the samples, 12 mL vials were flushed with He, filled with 0.5 mL 85% H_3PO_4 and 2 mL of these water samples, sealed airtight and stored to equilibrate for 48 h (Li et al., 2007; Taipale and Sonninen, 2009).

2.1.4 Sample analyses

Foraminiferal samples and water samples were analysed at the Stable Isotope Laboratory at the University of Vienna for Environmental Research (SILVER). Phytodetrital (dry weight) and foraminiferal content of organic carbon or nitrogen and ratios 10 of $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ were determined with an Isotope Ratio Mass Spectrometer (IRMS; DeltaPLUS, Thermo Finnigan) coupled with an interface (ConFlo III, Thermo Finnigan) to an elemental analyzer (EA 1110, CE Instruments). $\delta^{13}\text{C}$ of dissolved inorganic carbon (DIC- $\delta^{13}\text{C}$) was measured after release as CO_2 by H_3PO_4 addition in the headspace (headspace gas sampler: GasBench II, Thermo Fisher) of the prepared samples using an IRMS (Delta Advantage V, Thermo Fisher). Atom% 15 of the samples were derived from isotope ratio data and were calculated using the Vienna PeeDee Belemnite standard for C (RVPDB = 0.0112372) and atmospheric nitrogen for N (RatmN = 0.0036765), where X is ^{13}C or ^{15}N :

$$atom\% = \frac{100 * R_{\text{standard}} * \left(\frac{\delta X_{\text{sample}}}{1000} + 1 \right)}{1 + R_{\text{standard}} * \left(\frac{\delta X_{\text{sample}}}{1000} + 1 \right)} \quad (1)$$

Net uptake (uptake into the foraminiferal cell, excluding released amounts, hereafter referred to as 'uptake') of phytodetrital carbon and nitrogen in foraminiferal cytoplasm was calculated by determining the excess (E) of isotope content within the samples against the natural abundance of the isotopes in the foraminiferal cytoplasm (Middelburg et al., 2000):

$$E = \frac{\text{atom}X_{\text{sample}} - \text{atom}X_{\text{background}}}{100} \quad (2)$$

5 where X is ^{13}C or ^{15}N . Excess and total organic carbon and nitrogen (C, N, biomass normalised, per mg sample weight, or per individual) were used to calculate the amount of incorporated isotope $I_{\text{iso}} \mu\text{g mg}^{-1}$ or $\mu\text{g ind}^{-1} = E \times C (N) \mu\text{g mg}^{-1}$ or $\mu\text{g ind}^{-1}$, to obtain the amount of phytodetrital carbon (pC $\mu\text{g mg}^{-1}$ or $\mu\text{g ind}^{-1}$) and nitrogen pN ($\mu\text{g mg}^{-1}$ or $\mu\text{g ind}^{-1}$) within the foraminiferal cytoplasm (Hunter et al., 2012):

$$pX = \frac{\frac{I_{\text{iso}}}{\text{atom}\%X_{\text{phyto}}}}{100} \quad (3)$$

10 For a better comparison of the variation in uptake dynamics between the two species, the uptake of labelled particles was time normalized to obtain uptake rates of phytodetrital carbon and nitrogen ($\text{ng mg}^{-1} \text{h}^{-1}$). Exponential decay functions ($y = a (-b^*x)$) were applied with least squares global curve fitting for carbon uptake rates. The steepness of the decrease b (high b value = fast decrease) indicates whether there is a fast or slow drop in the uptake rates within the related samples over time. An F-test was carried out with mean values of treatment curves to compare the nonlinear regression models (exponential decrease) 15 for uptake rates of phytodetrital carbon and evaluate if a single model can be fitted for the two species at 20°C. The same procedure was used to prove if uptake rates of phytodetrital carbon vary within species between the three temperature treatments. Homogeneity of variances was tested using Fligner-Killeen's test which can be applied when population means are not known (Conover et al., 1981) and can overcome some problems with small sample sizes (Wasserstein and Boyer Jr, 1991). Additionally, graphical exploration of variance homogeneity was carried out by plotting the residuals for pC, pN, TOC and TN (Zuur 20 et al., 2010), for details see Suppl. Fig 1. Welch's t-test was used to detect differences of C:N ratios, pC:pN ratios and cytoplasmatic C and N content between species. This test is recommended for sample sizes < 10 , robust against heteroscedacity within the data (Moser and Stevens, 1992; Ruxton, 2006; McDonald, 2014) and can still be applied in case of unequal variances. To compare within species differences, a two-way ANOVA was carried out with time and temperature as independent factors 25 and pC, pN, C and N as dependent variables. The statistical analysis applied here were meant to emphasise the visualizations (graphical depictions) of the findings of this study. The applied tests were chosen to suit and represent the data in the most appropriate way (to retain robustness with respect to the design of the study). Graphs and data analysis were done using R (R Development Core Team, 2008) via RStudio (RStudio Team, 2015) and the packages ggplot 2 (Wickham, 2009) nlstools, (Florent Baty et al., 2015) and plyr (Wickham, 2011).

Table 1 Delta values for carbon and nitrogen isotopes. Natural $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of unlabelled background samples (BG), of isotopically enriched samples of food (*D. tertiolecta*) and foraminifera, and DIC- $\delta^{13}\text{C}$ within water samples at the 2nd day of the incubation period (standard deviation in parenthesis; letters denote Tukey Grouping of significant differences in DIC- $\delta^{13}\text{C}$ between temperatures; n.d. = no data).

		$\delta^{13}\text{C} \text{‰}$	$\delta^{15}\text{N} \text{‰}$	DIC $\delta^{13}\text{C}$
<i>D. tertiolecta</i>	BG	-18.3	16.2	n.d.
	labeled	30267	213298	n.d.
<i>A. tepida</i>	BG	-13.9 (± 0.2)	13.4 (± 0.6)	n.d.
	20°C	1742 (± 216)	7277 (± 1203)	n.d.
	25°C	1269 (± 318)	5758 (± 1426)	n.d.
	30°C	707 (± 122)	4109 (± 320)	n.d.
<i>H. germanica</i>	BG	-13.7 (± 0.7)	11.7 (± 0.4)	n.d.
	20°C	248 (± 26)	1087 (± 71)	1845 (± 76) a
	25°C	118 (± 20)	784 (± 86)	2186 (± 60) b
	30°C	116 (± 5)	879 (± 29)	2214 (± 66) b

3 Results

3.1 Carbon processing

Elevated ^{13}C values of the foraminiferal samples demonstrated the strong response to the labelled food source (Table 1). Temperature had a significant impact on pC levels in *H. germanica* (Table 2 and Fig 2). An interactive effect of time and 5 temperature caused a different time related processing of pC at 20°C compared to 25° and 30°C (Table 2 and Fig 2). The content of pC in *H. germanica* at 20°C was significantly higher than at 25°C and 30°C on each day of data collection ($p < 0.05$; no significant difference in 25°C and 30°C samples). Additionally, values for DIC- $\delta^{13}\text{C}$ measured in water samples (day 2) at elevated temperatures were significantly higher than those from the 20°C approach (Table 1). Time related variations 10 in cytoplasmatic C (biomass normalised) in *H. germanica* specimens were also temperature dependent (Table 2). In contrast, there was no relation between time and temperature concerning the pC content of *A. tepida*. Levels of pC in *A. tepida* were significantly increased at 25°C (25°C - 20°C: $p = 0.015$, 25°C - 30°C: $p < 0.001$, pairwise t-testing with pooled SDs, since no 15 significance of time effects), revealing a higher optimum grazing temperature for *A. tepida* compared to *H. germanica* (pC, Fig 2). Temperature had no effect on biomass C in *A. tepida*, while there were time dependent variations in biomass C (Table 2). These appeared at the start and end of the experiment but no significant influence by temperature treatments on the other days. Individual cytoplasmatic C content $\mu\text{g ind}^{-1}$ was different between the two species and proportions of individual uptake of pC to 20 individual C content showed similar proportions like biomass related carbon values (Fig 3). For detailed information about 25 individual pC, pN, TOC and TN content see Supplementary Table.

Table 2 Effects of time and temperature. Results of a two-way ANOVA displaying the main effects time and temperature on the dependent variables pC, pN, C, N in *A. tepida* and *H. germanica* and their interactions. Significant p values are printed in bold letters.

			Df	SM	F value	p value
<i>A. tepida</i>	pC	temperature	2	17.341	7.898	0.002
		day	3	5.298	2.413	0.092
		temperature x day	6	1.299	0.592	0.734
		Error	24	2.196		
	pN	temperature	2	0.728	9.284	0.001
		day	3	1.939	24.716	< 0.001
		temperature x day	6	0.084	1.073	0.406
		Error	24	0.079		
	C	temperature	2	81.230	2.373	0.110
		day	4	445.060	12.998	< 0.001
		temperature x day	8	42.650	1.246	0.308
		Error	30	34.240		
	N	temperature	2	2.112	2.848	0.074
		day	4	4.221	5.690	0.001
		temperature x day	8	0.601	0.810	0.600
		Error	30	0.742		
<i>H. germanica</i>	pC	temperature	2	0.296	81.300	< 0.001
		day	3	0.010	2.654	0.071
		temperature x day	6	0.010	2.826	0.032
		Error	24	0.004		
	pN	temperature	2	0.002	8.524	0.002
		day	3	0.027	133.994	< 0.001
		temperature x day	6	0.001	5.260	0.001
		Error	24	< 0.000		
	C	temperature	2	14.973	3.630	0.039
		day	4	40.819	9.898	< 0.001
		temperature x day	8	17.057	9.898	0.002
		Error	30	4.124		
	N	temperature	2	2.240	13.722	< 0.001
		day	4	2.978	18.248	< 0.001
		temperature x day	8	0.275	1.685	0.143
		Error	30	0.163		




Figure 2. Phytodetrital derived carbon and nitrogen in foraminiferal cytoplasm. Biomass normalized pC and pN of *A. tepida* and *H. germanica* at 20°C, 25°C, and 30°C. Error bars denote standard deviation.

Table 3 Carbon uptake rates. Regression variables, correlation coefficient R^2 , standard error of estimate S and decay b the exponential decrease of carbon uptake rates of *H. germanica* and *A. tepida* versus time.

		R^2	S	b
<i>A. tepida</i>	20°C	0.873	11.382	-0.262 (± 0.048)
	25°C	0.595	21.247	-0.130 (± 0.047)
	30°C	0.918	4.797	-0.130 (± 0.017)
	all T	0.628	17.048	-0.162 (± 0.030)
<i>H. germanica</i>	20°C	0.945	0.957	-0.247 (± 0.030)
	25°C	0.904	0.797	-0.343 (± 0.058)
	30°C	0.912	0.789	-0.523 (± 0.090)
	all T	0.732	1.680	-0.316 (± 0.053)




Figure 3. Cytoplasmatic carbon (C) and nitrogen (N) content and pC or pN uptake per foraminiferal individual. Data represents values for day 2 of the 20°C approach. Error bars denote standard deviation.




Figure 4. Carbon uptake rates. Uptake rates of phytodetrital carbon of *A. tepida* and *H. germanica* at three temperatures. Dots represent the calculated uptake rates derived from isotope data per time, curves show functions of exponential decrease (curves are combined for *A. tepida*, due to statistical similarity).

Carbon uptake rates showed an exponential decrease with time for both species (Fig 4), with a very good fit of curves except for *A. tepida* at 25°C where a relatively low determination coefficient R^2 was observed (Table 3). Mean uptake rates display a highly significant difference between species (F-test: $F = 33.74$, $p = 0.029$). *H. germanica* showed a much lower efficiency in the uptake of phytodetritus derived from *D. tertiolecta* compared to *A. tepida*. Uptake rates in *A. tepida* can be pooled within 5 a single function (F-test; $F = 7.664$, $p = 0.062$). Temperature causes a strong variation in rates for *H. germanica* (F-test; $F = 24.97$, $p = 0.012$). The rate of *A. tepida* showed a slightly faster decrease at 20°C relative to the higher temperatures and the highest deviation (S) of $21.247 \text{ ng mg}^{-1} \text{ h}^{-1}$ from average uptake rates at 25°C. In general, *H. germanica* exhibited a faster drop in the uptake rates at the higher temperatures (c.f. b, Table 3).

3.2 Nitrogen processing

10 The $\delta^{15}\text{N}$ signatures of all foraminiferal cytoplasm samples showed a strong increase after addition of labelled algae (Table 1). Like in pC, time and temperature showed interactive effects on the pN content in *H. germanica* (Table 2). *Haynesina germanica* showed a temperature related variation on day 4 and 14. Cytoplasmatic N showed less variation with temperature in *A. tepida*




Figure 5. Nitrogen uptake rates. Uptake rates of phytodetrital nitrogen by *A. tepida* and *H. germanica* at 20°C, 25°C, and 30°C. Error bars denote standard deviation

than in *H. germanica*. In contrast to the pC values, progressing time caused significant increases in pN content in both species (Fig 2, pN). N changes in *A. tepida* were time dependent and increased at the start of the 25°C and 30°C approach. Analogous to cytoplasmatic C, individual N and pN are proportional to total biomass related values (Fig 3).

In general, nitrogen uptake rates of *H. germanica* showed a steeper decline over time and at all temperatures in contrast to *A. tepida* (Fig 5). While *H. germanica* showed the highest uptake rates at 20°C, uptake rates for *A. tepida* were highest at 25°C. At 5 30°C, uptake rates of phytodetrital nitrogen showed fluctuating patterns for both species. In *Ammonia* samples, rates at 25°C and 30°C drop after the 4th day. The steepest decrease at 20°C can be found between day 2 and day 4. Uptake rates of nitrogen were about tenfold higher in *A. tepida* compared to *H. germanica*.

There was a noticeable peak in N uptake on the 4th day at 30°C in both species. In *A. tepida*, this peak describes an increase 10 of the uptake rate followed by a linear decrease. In contrast, all uptake rates showed a rapid drop in *H. germanica* especially at the higher temperatures with a recognisable negative peak at 30°C on day 4, and a flattening of nitrogen uptake rates at 25°C and 30°C on day 4.




Figure 6. Ratios of cytoplasmatic pC:pN ratios. Biomass normalized pC:pN within foraminiferal cytoplasm. Error bars denote standard deviation, stars indicate significant differences of pC:pN ratios between the two species on the respective day (Welch's t-test; *p <0.050, **p <0.010; ***p <0.001).

3.3 Relationship of cytoplasmatic and phytodetrital derived carbon and nitrogen content

Haynesina germanica showed a pN enrichment in contrast to pC loss or relatively stable pC over time (Fig 2). This enrichment of pN was detectable in both species, but was stronger in *H. germanica* with an almost equal increase at all temperatures until the end of the experiment. In contrast, the pN rise in *A. tepida* was restricted to the first week. Despite the constant enrichment of pN in *H. germanica*, pC levels were lower at higher temperatures. This faster loss of pC at 25°C and 30°C at similar levels of phytodetritus uptake results in lower pC:pN ratios at higher temperatures (Fig 6). The interactive effect of temperature and time on the food uptake of *H. germanica* caused a decoupling of pC and pN (Table 2 and Fig 7). This is visualized by a lack of correlation between phytodetrital carbon and nitrogen content in the cytoplasm of *H. germanica* over the course of the experiment (Fig 7). In contrast, pC and pN were strongly coupled in *A. tepida* across the entire time series (Fig 7). Cytoplasmatic C:N ratios were generally similar in both species, with no significant impact of temperature (Table 4).

4 Discussion

4.1 Effect of temperature on carbon and nitrogen uptake

The metabolism of *H. germanica* was significantly affected by elevated temperatures. Higher temperatures reduced the amount of pC in the foraminiferal cytoplasm, while a slighter effect on pN indicates a lower impact of temperature on the general




Figure 6. Carbon and nitrogen coupling. Phytodetrital derived nitrogen (pN) vs. carbon (pC) of *A. tepida* and *H. germanica* at 20°C (circles), 25°C (squares), and 30°C with linear regression for *A. tepida*.

Table 4 C:N ratios of algae and foraminiferal cytoplasm. Brackets contain standard deviation. Ratios in bold letters denote differences of

C:N between *A. tepida* and *H. germanica*. Welch's t-test, $p < 0.050$.

		20 °C	25 °C	30 °C
<i>D. tertiolecta</i>		6.08	-	-
<i>A. tepida</i>	0 days	4.20(± 0.42)	4.20 (± 0.42)	4.20 (± 0.42)
	2 days	5.32 (± 0.59)	5.32 (± 0.20)	5.79 (± 0.90)
	4 days	5.02 (± 0.21)	4.97 (± 0.07)	5.70 (± 0.36)
	7 days	5.30 (± 0.49)	5.60 (± 0.30)	4.99 (± 0.05)
	14 days	4.29 (± 0.87)	4.04 (± 0.44)	4.41 (± 0.35)
<i>H. germanica</i>	0 days	5.07 (± 0.31)	5.07 (± 0.31)	5.07 (± 0.31)
	2 days	4.77 (± 0.14)	4.85 (± 0.21)	4.68 (± 0.15)
	4 days	4.63 (± 0.43)	4.39 (± 0.32)	3.65 (± 0.38)
	7 days	4.57 (± 0.42)	4.34 (± 0.17)	4.67 (± 0.36)
	14 days	5.04 (± 0.09)	4.32 (± 0.32)	4.51 (± 0.18)

uptake of algal phytodetritus. In contrast, pC processing in *A. tepida* was favoured at the intermediate experimental temperature level of 25°C, demonstrating different optimum phytodetritus processing temperatures in the two species. The loss of pC (Fig 2) in *H. germanica* in the warmer environment or the faster decline of uptake rates at 25°C and 30°C (Fig 4 and Table 3) was most likely caused by elevated respiration rates. Accordingly, increased DIC-¹³C content within the SSW at elevated 5 temperatures suggests higher respiratory activities (Table 1). Elevated temperatures have been reported to increase respiration and food consumption in aquatic herbivores, detritivores and foraminifera (Bradshaw, 1961; Carr and Bruno, 2013), to lower grazing rates with simultaneously increased oxygen consumption (Ferreira et al., 2010), or to raise respiration to cover the costs of maintenance of protein metabolism levels (Whiteley and Faulkner, 2005).

In general, DIC-¹³C values in this experiment were very high compared to foraminiferal ¹³C. It cannot be excluded that the

foraminiferal cultures contained bacteria. Thus, a considerable amount of DIC could originate from phytodetritus remineralization by microbial activity. Several perturbation- and long term experiments on marine bacteria have also reported increased respiration due to warming (Vázquez-Domínguez et al., 2007; Hoppe et al., 2008; Wohlers et al., 2009). However, the higher temperatures trigger noticeable stress in *H. germanica*, which impacts food uptake efficiency. The interactive time/temperature 5 effect supports this argument, apparent as the higher temperatures caused convergence of pC and pN content compared to the 20°C results, particularly until day 7. This indicates an offset of nutritional ingestion performance in the two warmer approaches. It further suggests a general preference of lower environmental temperatures. Evidently, lower temperatures are more supportive to trigger reproduction in *H. germanica* (Goldstein and Alve, 2011). Interestingly, there is no significant temperature 10 effect on pC between 25°C and 30°C. This implicates a critical threshold for this species between 20°C and 25°C. Above this level, the mineralization of carbon increases, as the DIC-¹³C rises at 25°C and 30°C as a result of elevated respiratory activity. The decoupling of pC and pN in *H. germanica* over time supports this observation.

In contrast, *A. tepida* showed a trend for optimum uptake of carbon at 25°C and the steadiest carbon uptake rates at 30°C, while the pC and pN values showed similar patterns (Fig 2 and Fig 4). Laboratory experiments on *A. tepida* specimens collected at the Brouage mudflat (France) revealed an optimum temperature of grazing on bacteria at 30°C (Pascal et al., 2008b). This 15 indicates that strains of *A. tepida* feature adaptations to the higher average sea surface temperatures of the Brouage mudflat. In other laboratory experiments (Bradshaw, 1957, 1961) warm temperatures between 25°C and 30°C, offered optimum conditions for reproduction, resulted in higher growth and reproductive rates and a decrease in generation time in laboratory experiments with *A. tepida* specimen. Accordingly, higher temperatures are likely to offer competitive advantages for *A. tepida* over *H. germanica* in terms of food uptake. Considering the high expression of stress proteins at 35°C in *A. tepida* (Heinz et al., 2012) 20 in relation to the here presented results, a critical temperature limit effecting physiologic performances is likely to be found between 30°C and 35°C in this species.

Interestingly, the higher temperature treatments caused an increase of biomass C and N in *H. germanica* samples on the second sampling day, followed by a quick decrease on the fourth day. A similar trend was observed in *A. tepida*. This effect is difficult 25 to interpret and more detailed information about initial food uptake between day 0 and day 2 would be necessary. In summary, continued food C and N uptake appears to be a key factor in the strategies of *A. tepida*, according to the strong response to phytodetritus and the low effect of temperature on its uptake. Simultaneously, a high viability of the specimen in culture implies specific environmental adaptations and a strong response and sensitivity to disturbances caused by the transfer to the laboratory (Fig 2). In contrast, *H. germanica* showed a comparably high robustness and overall vitality, but an obviously strong impact of 30 temperature on mechanisms involved in carbon processing. This implicates physiological resource investment for survival by costs of increased metabolic activity.

4.2 Relationships between carbon and nitrogen uptake and cytoplasmatic C:N ratios

In both species, the relation of pC to pN (pC:pN ratio) within foraminiferal cytoplasm decreased compared to their phytodetrital C:N ratio of ~10 (Table 4) as time progressed. In theory, a comparably high ratio of detrital derived pC:pN would approach or undercut somatic consumer C:N ratio to retain homeostasis or ensure appropriate nutrition (Cross et al., 2005; Frost et al.,

2005; Sterner and Elser, 2002). The growing content of pN within foraminiferal cytoplasm over time with a remarkable increase of ^{15}N values (Table 1 and Fig 2) could be explained by the integration of a high amount of food derived nitrogen into amino acids and metabolic consumption of pC. Benthic foraminifera inhabiting Antarctic sediments have been reported to use the major amount of algae derived carbon sources to synthesize nitrogen rich proteins in relation to a lower generation of other products with lower nitrogen content (Rivkin and DeLaca, 1990). Such effects could be responsible for the relatively slight decrease in nitrogen uptake rates over time compared to the strong decrease in carbon uptake rates (Fig 4 and Fig 5). A strong decline of continued food uptake or degradation induced change of phytodetrital C:N could likewise be responsible for this effect.

Both tested species showed similar demands for food derived carbon and nitrogen, since at moderate temperatures (20°C) pC:pN ratios showed a similar trend over time (Fig 6). Again, minor temperature influences were demonstrated within the overall linear relationship of pC to pN in *A. tepida* across the samples (Fig 7). In contrast, the rising variation and highly significant difference in pC:pN (lower ratios at higher temperatures, Fig 6) resulting in the diffuse relationship of pC:pN (Fig 7) in *H. germanica* reflect the thermal stress on the level of increased carbon expenditure at steady feeding progress (barely influenced pN levels between treatments). Therefore, sediments with high dominance of *H. germanica* like river inlets or estuaries (Debenay et al., 2006; Murray and Alve, 2000) could possibly experience additional temperature induced nutrient flux variations, boosting C losses while triggering N retention.

An *in situ* experiment, monitoring population shifts of near shore benthic foraminifera to artificially heated sediments, proposed Migrations to deeper (colder) regions as a response to elevated environmental temperatures (Schafer et al., 1996). Marine species are known to Migrate linked to temperature changes (Parmesan, 2006). To maintain an optimum energy budget, expenses of metabolic maintenance could therefore be compensated by a shift of *H. germanica* populations towards the subtidal. Foraminiferal C:N ratios are known from bathyal foraminifera of the Sagami Bay and from oxygen minimum zones of the Arabian Sea, where they range from 2.6 - 6.4 (Nomaki et al., 2008, 2011; Enge et al., 2016). In our study, both species showed similar C:N ratios and natural ^{13}C and ^{15}N signatures (Table 1 and Table 4), indicating equivalent trophic levels and similar nutritional demands. During the course of the experiment, the temperature stress in *H. germanica* was even noticeable on the level of cytoplasmatic stoichiometry. The remarkable drop of the C:N ratio on the 4th day at 30°C (Table 4) and the subsequent recovery and rapid increase of pC and pN after the 4th day (Fig 2) could denote a metabolic adaptation process. However, the strong temperature influence on carbon uptake persisted.

In *A. tepida*, the occasionally significant (at 25°C) fluctuations in the C:N ratio can be related to the high uptake of phytodetritus containing high C:N ratios and therefore an influence of food source C:N on the grazer C:N (Fig 3 and Table 4). Generally, considerations of C:N ratios and intracellular pC/pN coupling also reflect the lower effects of high temperature exposure on *A. tepida* and the high impact on the feeding behaviour of *H. germanica*. These observations reflect the different strategies of the two coexisting intertidal foraminiferal species on the level of nutrient balance. It further implies shifts in population distributions of *A. tepida* and *H. germanica* at events of persisting food source and/or temperature changes. This study verifies a strong involvement of foraminifera in the turnover of intertidal POM. Depending on the POM source or dominant foraminiferal species, this turnover can be strongly influenced by increased temperature, causing a decoupling of carbon and nitrogen cycling.

However, when relating these findings to natural environments, considerations of cascading effects of temperature change e.g. switches in trophic conditions and related shifts in nutrient coupling of primary producers have to be included.

4.3 Species specific food uptake

The very high uptake rates and levels of phytodetrital carbon and nitrogen in *Ammonia tepida* (day 2 ~14% pC:C; ~10% pN:N, 5 comp. Fig 3 and Fig 4) indicate a strong response to the food source, in contrast to *H. germanica* which showed a clear, but much lower uptake (day 2: ~2% pC:C; ~1.5% pN:N). The observed decline of carbon uptake rates by time in both species suggests the highest uptake within the first two days of the experiment and indicates a fast processing of the phytodetritus source. During an in situ experiment, Moodley et al. (2000) also observed a rapid response of *Ammonia* sp. to green algae (Chlorella sp.) within hours, together with the much weaker ingestion of such food source by *Haynesina* sp.

10 The quick slowdown of food uptake rates could indicate levels of saturation especially after a strong response following the food pulse. In addition, cellular release of label is likely to exceed uptake as time progresses. Other reasons for the decrease in food uptake with increasing time (reflected in decreasing uptake rates of C and N as well as in pC and pN) could be the stage of phytodetrital decay or decreasing food availability at the subsequent sampling days. Aging or phytodetrital degeneration and its change in quality in marine benthic environments is a result of bacterial colonisation and subsequent transformation 15 and mineralization of the algal material (Middelburg et al., 2000; Moodley et al., 2002; Buhring et al., 2006; Gihring et al., 2009). Through bacterial mineralization, the degrading detritus decreases in quality with an increase in C:nutrient ratios. However, microorganisms colonizing patches of degrading detritus are likely to be incorporated simultaneously with foraminiferal phytodetritus grazing. Therefore, a fraction of the isotope label within the foraminiferal cytoplasm could be a result of indirect 20 label intake with phytodetritus associated microorganisms especially towards the end of the experiment. However, a visible layer of phytodetritus particles persisted until the end of the experiment. The influence of food availability on feeding, where high food concentrations support increased feeding rates, have been reported for foraminifera and also in other organismic groups such as macrofauna or bacteria (Pascal et al., 2008b, a; Quijón et al., 2008; Mayor et al., 2012). Fast incorporation of food pulses seems to be necessary to cover the high energy demand for reproduction and growth, since food concentrations (Dunaliella) sp. of less than 112 cells mm² (0.40 µg C cm⁻² (Heinz et al., 2002)) do not permit growth or reproduction in *A. 25 tepida*, while growth rate and reproductive activity increase with additional food (Bradshaw, 1955, 1957). Lee et al. (1966) stated, that *Ammonia beccarii* specimen preferred to feed on 'new' (Fig 1. 10 days old) cultures of living *Chlorococcum* sp. over 'old' (up to 40 days old) cultures. Following these observations, food limitation resulting from the aging of the phytodetrital food source could be present in this study. This proceeding degradation of phytodetritus appears to be particularly problematic to *A. tepida* specimen, which showed an increasing fraction of dead individuals with progressing time (Fig 1). On the other 30 hand, the health condition of the specimens could be a response to the conditions of the laboratory incubation. Compared to a previous feeding experiment by Linsky et al. (2014) with *A. tepida* fed with *D. tertiolecta* phytodetritus (614 mg C m⁻²), uptake rates of carbon of 1899 pg ind⁻¹ h⁻¹ were much higher in this study after 48h than in the other feeding experiment (149 pg ind⁻¹ h⁻¹). This might be related to general differences in feeding of the tested strains.

The low affinity of *H. germanica* to *D. tertiolecta* phytodetritus could be explained by a generally lower carbon demand, or

more likely a specialization on other food sources, for instance diatoms or secondary products of microphytobenthic biofilms (e.g. extracellular polymeric substances). The latter assumption corresponds to the preference of a diatom diet over sewage derived POM (Ward et al., 2003), the presence of a diatom cracking mechanism (Austin et al., 2005), the sequestration of chloroplasts derived exclusively from diatoms (Knight and Mantoura, 1985; Pillet et al., 2011; Cevasco et al., 2015), or the 5 correlation of the distribution of *H. germanica* populations with high abundances of *Nitzschia* sp. (Hohenegger et al., 1989). This coherence with food availability or organic matter accumulation respectively, was used to explain population distributions of a *H. germanica*/*A. beccarii* assemblages in muddy sediments of Spain (Papaspyrou et al., 2013; Diz et al., 2012). The different patterns of pC accumulation over time in the two species also reflect a different feeding behaviour. The onset of 10 declining health conditions in *H. germanica* specimen at the end of the experiment (Fig 1) was likely caused by starvation, due to the absence of an appropriate food source. In general, the quality of organic carbon or foraminiferal food sources oscillates throughout the year and includes allochthonous detritus (Heip et al., 1995), while the main source of primary production in intertidal environments are microphytobenthic diatoms, accompanied by chlorophytes or other autotrophic microorganism, 15 seasonally suppressing diatom dominance (Scholz and Liebezeit, 2012). In this experiment, *D. tertiolecta* was used as food source, because this species is easy to maintain in culture and was previously used in several feeding experiments with benthic foraminifera and serves as a representative for allochthonous detrital carbon sources. Ratios of *A. tepida* and *H. germanica* abundances are considered to be related to organic matter quality or environmental variability, due to their different feeding 20 specialisations or environmental adaptations (Scholz and Liebezeit, 2012). High amounts of pC (max. 30 % pC to C) in *A. tepida* compared to *H. germanica* (max. 2.6 % pC to C), together with the low influence on temperature on the feeding behavior of *A. tepida*, prove an opportunistic feeding behavior and generalist temperature adaptations in *A. tepida*. These findings will 25 help to interpret oscillations in abundances of these two common intertidal foraminiferal species in relation with organic carbon quality and ambient temperatures. These data report varying strategies in phytodetritus feeding by means of source exploitation and time dependence. However, laboratory studies do not always reflect natural behavior accurately and might be biased by the suppression or alteration of natural behaviors in the investigated organisms. This work is meant to help solving ecological problems and the here presented patterns of foraminiferal feeding behavior can be useful for the careful interpretation or explanation of successions in species dominances by means of food source availability and complements data from field surveys.

5 Conclusions

According to the results of this study, *H. germanica* exhibited a more vulnerable carbon retention behaviour to variations in temperature and a relatively low affinity to *D. tertiolecta* phytodetritus, suggesting other food preferences. *Ammonia tepida* 30 showed a broader tolerance range of temperature concerning carbon and nitrogen uptake and a highly effective food exploitation. The different response of the two species to the applied treatments implies different strategies and environmental adaptations. Consequently, a temperature related shift in abundances of the two species could alter carbon and nitrogen fluxes in intertidal sediments, with respect to high chlorophyte detritus uptake of *A. tepida* (*A. tepida*: pC max. 90 ng ind⁻¹, pN max.

20 ng ind⁻¹ *H. germanica*: pC max. 5 ng ind⁻¹, pN max. 2 ng ind⁻¹) and temperature sensitivity of *H. germanica*. In contrast to food derived carbon, the increasing accumulation of nitrogen was barely affected by temperature in either species. This raises the hypothesis, that rapid temperature increases in *H. germanica* dominated sediments could cause a shift in organic carbon and nitrogen cycling towards enhanced nitrogen retention and carbon losses. A general enrichment in cytoplasmatic nitrogen in 5 both species prove, that nitrogen has a higher potential to retain within the foraminiferal consumer. Further, increasing time had a prolonged negative influence on the food uptake rates from a single food pulse. A reduced availability of fresh phytodetritus sources is likely to cause such effects and appeared to be specifically limiting for *A. tepida*.

Author contributions. EAJ, PH and JW planned the study. EAJ and JW conducted the fieldwork and processed the samples in the lab. MW performed elemental and isotope analysis and adapted the method for the foraminiferal samples. JW performed data analysis and wrote the 10 manuscript. EAJ, PH and WW contributed by discussing results and critical revision of manuscript drafts.

Competing interests. The authors declare no competing interests.

Acknowledgements. Gerhard Schmiedl (Universität Hamburg) thankfully provided equipment for field sampling. Yvonne milker (Universität Leipzig) and Katharina Müller-Navarra (Universität Hamburg) supported us at the sampling location. Patrick Bukenberger helped with lab work and microscoping at the University of Vienna. This article was supported by the Open Access Publishing Fund of the University of Vienna.

References

Al-Raei, A. M., Bosselmann, K., Böttcher, M. E., Hespenheide, B., and Tauber, F.: Seasonal dynamics of microbial sulfate reduction in temperate intertidal surface sediments: controls by temperature and organic matter, *Ocean Dynamics*, 59, 351–370, 2009.

5 Allen, A., Gillooly, J., and Brown, J.: Linking the global carbon cycle to individual metabolism, *Functional Ecology*, 19, 202–213, 2005.

Alve, E. and Murray, J.: Ecology and taphonomy of benthic foraminifera in a temperate mesotidal inlet, *Journal of Foraminiferal Research*, 24, 18–27, 1994.

Alve, E. and Murray, J. W.: Temporal variability in vertical distributions of live (stained) intertidal foraminifera, southern England, *The Journal of Foraminiferal Research*, 31, 12–24, 2001.

10 Austin, H. A., Austin, W. E., and Paterson, D. M.: Extracellular cracking and content removal of the benthic diatom *Pleurosigma angulatum* (Quekett) by the benthic foraminifera *Haynesina germanica* (Ehrenberg), *Marine Micropaleontology*, 57, 68–73, 2005.

Bowser, S. S., Alexander, S. P., Stockton, W. L., and Delaca, T. E.: Extracellular matrix augments mechanical properties of pseudopodia in the carnivorous foraminiferan *Astrammina rara*: role in prey capture, *The Journal of Protozoology*, 39, 724–732, 1992.

Bradshaw, J. S.: Preliminary laboratory experiments on ecology of foraminiferal populations, *Micropaleontology*, pp. 351–358, 1955.

15 Bradshaw, J. S.: Laboratory Studies on the Rate of Growth of the Foraminifer, " *Streblus beccarii* (Linné) var. *tepida* (Cushman)", *Journal of Paleontology*, pp. 1138–1147, 1957.

Bradshaw, J. S.: Laboratory experiments on the ecology of foraminifera, Contributions from the Cushman Foundation for foraminiferal research, 7, 87 – 106, 1961.

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: Toward a metabolic theory of ecology, *Ecology*, 85, 1771–1789,

20 2004.

Buhring, S., Lampadariou, N., Moodley, L., Tselepides, A., and Witte, U.: Benthic microbial and whole-community responses to different amounts of ^{13}C -enriched algae: In situ experiments in the deep Cretan Sea (Eastern Mediterranean), *Limnology and Oceanography*, 51, 157, 2006.

Buzas, M. A.: Foraminifera as prey for benthic deposit feeders: results of predator exclusion experiments, *Journal of Marine Research*, 1978.

25 Buzas, M. A. and Carle, K. J.: Predators of foraminifera in the Indian River, Florida, *The Journal of Foraminiferal Research*, 9, 336–340, 1979.

Carr, L. A. and Bruno, J. F.: Warming increases the top-down effects and metabolism of a subtidal herbivore, *PeerJ*, 1, e109, 2013.

Cesbron, F., Geslin, E., Jorissen, F., Delgard, M., Charrieau, L., Deflandre, B., Jézéquel, D., Anschutz, P., and Metzger, E.: Vertical distribution and respiration rates of benthic foraminifera: Contribution to aerobic remineralization in intertidal mudflats covered by *Zostera noltei* meadows, *Estuarine, Coastal and Shelf Science*, 2016.

30 Cevasco, M. E., Lechliter, S. M., Mosier, A. E., and Perez, J.: Initial Observations of Kleptoplasty in the Foraminifera of Coastal South Carolina, *Southeastern Naturalist*, 14, 361–372, 2015.

Conover, W. J., Johnson, M. E., and Johnson, M. M.: A comparative study of tests for homogeneity of variances, with applications to the outer continental shelf bidding data, *Technometrics*, 23, 351–361, 1981.

35 Cook, P. L. M., Revill, A. T., Clementson, L. A., and Volkman, J. K.: Carbon and nitrogen cycling on intertidal mudflats of a temperate Australian estuary. III. Sources of organic matter, *Marine Ecology Progress Series*, 280, 55–72, doi:10.3354/meps280055, ://WOS:000225268500005, 2004.

Cross, W. F., Benstead, J. P., Frost, P. C., and Thomas, S. A.: Ecological stoichiometry in freshwater benthic systems: recent progress and perspectives, *Freshwater Biology*, 50, 1895–1912, 2005.

Culver, S. and Buzas, M.: The effects of anthropogenic habitat disturbance, habitat destruction, and global warming on shallow marine benthic foraminifera, *The Journal of Foraminiferal Research*, 25, 204–211, 1995.

de Nooijer, L. J.: Shallow water benthic foraminifera as proxy for natural versus human-induced environmental change, vol. 272, Utrecht University, 2007.

Debenay, J.-P., Bicchi, E., Goubert, E., and du Châtelet, E. A.: Spatio-temporal distribution of benthic foraminifera in relation to estuarine dynamics (Vie estuary, Vendée, W France), *Estuarine, Coastal and Shelf Science*, 67, 181–197, 2006.

Dickson, A. G. and Goyet, C.: Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water, publisher not identified, 1994.

Diz, P. and Francés, G.: Distribution of live benthic foraminifera in the Ría de Vigo (NW Spain), *Marine Micropaleontology*, 66, 165–191, 2008.

Diz, P., Barras, C., Geslin, E., Reichart, G.-J., Metzger, E., Jorissen, F., and Bijma, J.: Incorporation of Mg and Sr and oxygen and carbon stable isotope fractionation in cultured *Ammonia tepida*, *Marine Micropaleontology*, 92, 16–28, 2012.

du Châtelet, E. A., Degre, D., Sauriau, P.-G., and Debenay, J.-P.: Distribution of living benthic foraminifera in relation with environmental variables within the Aiguillon cove (Atlantic coast, France): improving knowledge for paleoecological interpretation, *Bulletin de la Société Géologique de France*, 180, 131–144, 2009.

Enge, A., Witte, U., Kucera, M., and Heinz, P.: Uptake of phytodetritus by benthic foraminifera under oxygen depletion at the Indian margin (Arabian Sea), *Biogeosciences*, 11, 2017–2026, 2014.

Enge, A. J., Nomaki, H., Ogawa, N. O., Witte, U., Moeseneder, M. M., Lavik, G., Ohkouchi, N., Kitazato, H., Kučera, M., and Heinz, P.: Response of the benthic foraminiferal community to a simulated short-term phytodetritus pulse in the abyssal North Pacific, *Marine Ecology Progress Series*, 438, 129–142, 2011.

Enge, A. J., Wukovits, J., Wanek, W., Watzka, M., Witte, U. F., Hunter, W. R., and Heinz, P.: Carbon and nitrogen uptake of calcareous benthic foraminifera along a depth-related oxygen gradient in the OMZ of the Arabian Sea, *Frontiers in microbiology*, 7, 2016.

Evrard, V., Soetaert, K., Heip, C. H., Huettel, M., Xenopoulos, M. A., and Middelburg, J. J.: Carbon and nitrogen flows through the benthic food web of a photic subtidal sandy sediment, *Marine Ecology Progress Series*, 416, 1–16, 2010.

Ferreira, V., Goncalves, A. L., Godbold, D. L., and Canhoto, C.: Effect of increased atmospheric CO₂ on the performance of an aquatic detritivore through changes in water temperature and litter quality, *Global Change Biology*, 16, 3284–3296, 2010.

Florent Baty, Christian Ritz, Sandrine Charles, Martin Brutsche, Jean-Pierre Flandrois, and Marie-Laure Delignette-Muller: A Toolbox for Nonlinear Regression in R: The Package nlstools, *Journal of Statistical Software*, 66, 1–21, 2015.

Frost, P. C., Evans-White, M. A., Finkel, Z. V., Jensen, T. C., and Matzek, V.: Are you what you eat? Physiological constraints on organismal stoichiometry in an elementally imbalanced world, *Oikos*, 109, 18–28, 2005.

Gihring, T. M., Humphrys, M., Mills, H. J., Huette, M., and Kostka, J. E.: Identification of phytodetritus-degrading microbial communities in sublittoral Gulf of Mexico sands, *Limnology and Oceanography*, 54, 1073–1083, 2009.

Goineau, A., Fontanier, C., Jorissen, F., Buscail, R., Kerhervé, P., Cathalot, C., Pruski, A., Lantoine, F., Bourgeois, S., Metzger, E., et al.: Temporal variability of live (stained) benthic foraminiferal faunas in a river-dominated shelf–Faunal response to rapid changes of the river influence (Rhône prodelta, NW Mediterranean), *Biogeosciences*, 9, 1367–1388, 2012.

Goldstein, S. T. and Alve, E.: Experimental assembly of foraminiferal communities from coastal propagule banks, *Marine Ecology Progress Series*, 437, 1–11, 2011.

Goldstein, S. T. and Corliss, B. H.: Deposit feeding in selected deep-sea and shallow-water benthic foraminifera, *Deep Sea Research Part I: Oceanographic Research Papers*, 41, 229–241, 1994.

Guillard, R. R.: Culture of phytoplankton for feeding marine invertebrates, in: *Culture of marine invertebrate animals*, pp. 29–60, Springer, 1975.

Guillard, R. R. and Ryther, J. H.: Studies of marine planktonic diatoms: I. *Cyclotella Nana* Hustedt, and *Detonula Confervacea* (CLEVE) Gran., *Canadian journal of microbiology*, 8, 229–239, 1962.

10 Haynert, K. and Schönfeld, J.: Impact of changing carbonate chemistry, temperature, and salinity on growth and test degradation of the benthic foraminifer *Ammonia aomoriensis*, *The Journal of Foraminiferal Research*, 44, 76–89, 2014.

Hayward, B. W., Grenfell, H., Cairns, G., and Smith, A.: Environmental controls on benthic foraminiferal and thecamoebian associations in a New Zealand tidal inlet, *The Journal of Foraminiferal Research*, 26, 150–171, 1996.

15 Heinz, P., Hemleben, C., and Kitazato, H.: Time-response of cultured deep-sea benthic foraminifera to different algal diets, *Deep Sea Research Part I: Oceanographic Research Papers*, 49, 517–537, 2002.

Heinz, P., Marten, R. A., Linsley, V. N., Haap, T., Geslin, E., and Köhler, H.-R.: 70 kD stress protein (Hsp70) analysis in living shallow-water benthic foraminifera, *Marine Biology Research*, 8, 677–681, 2012.

Heip, C., Goosen, N., Herman, P., Kromkamp, J., Middelburg, J., and Soetaert, K.: Production and consumption of biological particles in temperate tidal estuaries, 1995.

20 Hohenegger, J., Piller, W., and Baal, C.: Reasons for spatial microdistributions of foraminifers in an intertidal pool (northern Adriatic Sea), *Marine Ecology*, 10, 43–78, 1989.

Hoppe, H.-G., Breithaupt, P., Walther, K., Koppe, R., Bleck, S., Sommer, U., and Jürgens, K.: Climate warming in winter affects the coupling between phytoplankton and bacteria during the spring bloom: a mesocosm study, *Aquatic Microbial Ecology*, 51, 105–115, 2008.

Hunter, W., Levin, L., Kitazato, H., and Witte, U.: Macrobenthic assemblage structure and organismal stoichiometry control faunal processing 25 of particulate organic carbon and nitrogen in oxygen minimum zone sediments, *Biogeosciences*, 9, 993–1006, 2012.

Jeffreys, R. M., Burke, C., Jamieson, A. J., Narayanaswamy, B. E., Ruhl, H. A., Smith Jr, K. L., and Witte, U.: Feeding preferences of abyssal macrofauna inferred from in situ pulse chase experiments, *PloS one*, 8, e80510, 2013.

Joye, S. B., De Beer, D., and Cook, P. L. M.: Biogeochemical dynamics of coastal tidal flats., in: *Coastal wetlands: an ecosystem integrated approach.*, edited by Perillo, G. M. E., Wolanski, E., Cahoon, D. R., and Brinson, M. M., pp. 345–374, Elsevier, 2009.

30 Kester, D. R., Duedall, I. W., Connors, D. N., and Pytkowicz, R. M.: Preparation of artificial seawater, *Limnology and oceanography*, 12, 176–179, 1967.

Knight, R. and Mantoura, R. F. C.: Chlorophyll and carotenoid pigments in Foraminifera and their symbiotic algae: analysis by high performance liquid chromatography, *Marine Ecology Progress Series*, 23, 241–249, 1985.

Kroopnick, P., Weiss, R., and Craig, H.: Total CO₂, 13C, and dissolved oxygen-18O at Geosecs II in the North Atlantic, *Earth and Planetary 35 Science Letters*, 16, 103–110, 1972.

Lee, J. J. and Muller, W. A.: Trophic dynamics and niches of salt marsh foraminifera, *American Zoologist*, 13, 215–223, doi:10.2307/3881901, <http://www.jstor.org/stable/3881901>, 1973.

Lee, J. J., McEnery, M., Pierce, S., Freudenthal, H., and Muller, W.: Tracer experiments in feeding littoral foraminifera, *The Journal of Protozoology*, 13, 659–670, 1966.

Lei, Y.-L., Stumm, K., Wickham, S. A., and Berninger, U.-G.: Distributions and Biomass of Benthic Ciliates, Foraminifera and Amoeboïd Protists in Marine, Brackish, and Freshwater Sediments, *Journal of Eukaryotic Microbiology*, 61, 493–508, 2014.

Li, Z.-P., Tao, M.-X., Li, L.-W., Wang, Z.-D., Du, L., and Zhang, M.-F.: Determination of isotope composition of dissolved inorganic carbon by gas chromatography-conventional isotope-ratio mass spectrometry, *Chinese Journal of Analytical Chemistry*, 35, 1455–1458, 2007.

5 Linsley, V., Nigam, R., and Heinz, P.: Response of Shallow Water Benthic Foraminifera to a ^{13}C -Labeled Food Pulse in the Laboratory, in: *Approaches to Study Living Foraminifera*, pp. 115–131, Springer, 2014.

Lipps, J. H. and Valentine, J. W.: The role of foraminifera in the trophic structure of marine communities, *Lethaia*, 3, 279–286, 1970.

10 Mayor, D. J., Thornton, B., and Zuur, A. F.: Resource quantity affects benthic microbial community structure and growth efficiency in a temperate intertidal mudflat, *PLoS One*, 7, e38582, 2012.

McDonald, J.: Multiple comparisons: Controlling the false discovery rate: Benjamini–Hochberg procedure, *Handbook of Biological Statistics*. Baltimore, Maryland: Sparkly House Publishing, pp. 254–60, 2014.

Melis, R. and Covelli, S.: Distribution and morphological abnormalities of recent foraminifera in the Marano and Grado Lagoon (North Adriatic Sea, Italy), *Mediterranean Marine Science*, 14, 432–450, 2013.

15 Middelburg, J. J., Barranguet, C., Boschker, H. T., Herman, P. M., Moens, T., and Heip, C. H.: The fate of intertidal microphytobenthos carbon: An in situ ^{13}C -labeling study, *Limnology and oceanography*, 45, 1224–1234, 2000.

Moodley, L., Van der Zwaan, G., Herman, P., Kempers, L., and Van Breugel, P.: Differential response of benthic meiofauna to anoxia with special reference to Foraminifera (Protista: Sarcodina), *Marine Ecology Progress Series*, 158, 151–163, 1997.

20 Moodley, L., Boschker, H., Middelburg, J., Pel, R., Herman, P., De Deckere, E., and Heip, C.: Ecological significance of benthic foraminifera: ^{13}C labelling experiments, *Marine Ecology Progress Series*, 202, 289–295, 2000.

Moodley, L., Middelburg, J. J., Boschker, H. T., Duineveld, G. C., Pel, R., Herman, P. M., and Heip, C. H.: Bacteria and Foraminifera: key players in a short term deep-sea benthic response to phytodetritus, *Marine Ecology Progress Series*, 236, 23–29, 2002.

Moser, B. K. and Stevens, G. R.: Homogeneity of variance in the two-sample means test, *The American Statistician*, 46, 19–21, 1992.

25 Muller, W. A. and Lee, J. J.: Apparent indispensability of bacteria in foraminiferan nutrition, *The Journal of Protozoology*, 16, 471–478, 1969.

Murray, J. W.: *Ecology and palaeoecology of benthic foraminifera*, Routledge, 2014.

Murray, J. W. and Alve, E.: Major aspects of foraminiferal variability (standing crop and biomass) on a monthly scale in an intertidal zone, *The Journal of Foraminiferal Research*, 30, 177–191, 2000.

Nomaki, H., Heinz, P., Nakatsuka, T., Shimanaga, M., and Kitazato, H.: Species-specific ingestion of organic carbon by deep-sea benthic 30 foraminifera and meiobenthos: In situ tracer experiments, *Limnology and Oceanography*, 50, 134–146, 2005.

Nomaki, H., Heinz, P., Nakatsuka, T., Shimanaga, M., Ohkouchi, N., Ogawa, N. O., Kogure, K., Ikemoto, E., and Kitazato, H.: Different ingestion patterns of ^{13}C -labeled bacteria and algae by deep-sea benthic foraminifera, *Marine Ecology Progress Series*, 310, 95–108, 2006.

35 Nomaki, H., Ogawa, N. O., Ohkouchi, N., Suga, H., Toyofuku, T., Shimanaga, M., Nakatsuka, T., and Kitazato, H.: Benthic foraminifera as trophic links between phytodetritus and benthic metazoans: carbon and nitrogen isotopic evidence, *Marine Ecology Progress Series*, 357, 153–164, 2008.

Nomaki, H., Ohkouchi, N., Heinz, P., Suga, H., Chikaraishi, Y., Ogawa, N. O., Matsumoto, K., and Kitazato, H.: Degradation of algal lipids by deep-sea benthic foraminifera: an in situ tracer experiment, *Deep Sea Research Part I: Oceanographic Research Papers*, 56, 1488–1503, 2009.

Nomaki, H., Ogawa, N. O., Takano, Y., Suga, H., Ohkouchi, N., and Kitazato, H.: Differing utilization of glucose and algal particulate organic matter by deep-sea benthic organisms of Sagami Bay, Japan, *Marine Ecology Progress Series*, 431, 11–24, 2011.

O'Connor, M. I., Piehler, M. F., Leech, D. M., Anton, A., and Bruno, J. F.: Warming and resource availability shift food web structure and metabolism, *PLoS Biol*, 7, e1000178, 2009.

Papaspyprou, S., Diz, P., García-Robledo, E., Corzo, A., and Jimenez-Arias, J.-L.: Benthic foraminiferal community changes and their relationship to environmental dynamics in intertidal muddy sediments (Bay of Cádiz, SW Spain), *Marine Ecology Progress Series*, 490, 121–135, 2013.

Parmesan, C.: Ecological and evolutionary responses to recent climate change, *Annual Review of Ecology, Evolution, and Systematics*, pp. 637–669, 2006.

Pascal, P.-Y., Dupuy, C., Mallet, C., Richard, P., and Niquil, N.: Bacterivory by benthic organisms in sediment: Quantification using 15 N-enriched bacteria, *Journal of Experimental Marine Biology and Ecology*, 355, 18–26, 2008a.

Pascal, P.-Y., Dupuy, C., Richard, P., and Niquil, N.: Bacterivory in the common foraminifer *Ammonia tepida*: Isotope tracer experiment and the controlling factors, *Journal of Experimental Marine Biology and Ecology*, 359, 55–61, 2008b.

Petchey, O. L., Brose, U., and Rall, B. C.: Predicting the effects of temperature on food web connectance, *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 365, 2081–2091, 2010.

Pillet, L., de Vargas, C., and Pawlowski, J.: Molecular identification of sequestered diatom chloroplasts and kleptoplastid in foraminifera, *Protist*, 162, 394–404, 2011.

Quijón, P. A., Kelly, M. C., and Snelgrove, P. V.: The role of sinking phytodetritus in structuring shallow-water benthic communities, *Journal of Experimental Marine Biology and Ecology*, 366, 134–145, 2008.

R Development Core Team: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria., 2008.

Rivkin, R. and DeLaca, T.: Trophic dynamics in Antarctic benthic communities. I. In situ ingestion of microalgae by Foraminifera and metazoan meiofauna., *Marine ecology progress series*. Oldendorf, 64, 129–136, 1990.

RStudio Team: RStudio: Integrated Development Environment for R, RStudio, Inc., Boston, MA, <http://www.rstudio.com/>, 2015.

Ruxton, G. D.: The unequal variance t-test is an underused alternative to Student's t-test and the Mann–Whitney U test, *Behavioral Ecology*, 17, 688–690, 2006.

Schafer, C., Cole, F., Frobel, D., Rice, N., and Buzas, M.: An in situ experiment on temperature sensitivity of nearshore temperate benthic foraminifera, *Oceanographic Literature Review*, 9, 913, 1996.

Scholz, B. and Liebezeit, G.: Microphytobenthic dynamics in a Wadden Sea intertidal flat–Part II: Seasonal and spatial variability of non-diatom community components in relation to abiotic parameters, *European Journal of Phycology*, 47, 120–137, 2012.

Sterner, R. W. and Elser, J. J.: Ecological stoichiometry: the biology of elements from molecules to the biosphere, Princeton University Press, 2002.

Sweetman, A. K., Sommer, S., Pfannkuche, O., and Witte, U.: Retarded response by macrofauna-size foraminifera to phytodetritus in a deep Norwegian fjord, *The Journal of Foraminiferal Research*, 39, 15–22, 2009.

Taipale, S. J. and Sonninen, E.: The influence of preservation method and time on the $\delta^{13}\text{C}$ value of dissolved inorganic carbon in water samples, *Rapid Communications in Mass Spectrometry*, 23, 2507–2510, 2009.

Topping, J., Murray, J., and Pond, D.: Sewage effects on the food sources and diet of benthic foraminifera living in oxic sediment: a microcosm experiment, *Journal of Experimental Marine Biology and Ecology*, 329, 239–250, 2006.

Vázquez-Domínguez, E., Vaqué, D., and Gasol, J. M.: Ocean warming enhances respiration and carbon demand of coastal microbial plankton, *Global Change Biology*, 13, 1327–1334, 2007.

Ward, J. N., Pond, D. W., and Murray, J. W.: Feeding of benthic foraminifera on diatoms and sewage-derived organic matter: an experimental application of lipid biomarker techniques, *Marine Environmental Research*, 56, 515–530, 2003.

5 Wasserstein, R. L. and Boyer Jr, J. E.: Bounds on the power of linear rank tests for scale parameters, *The American Statistician*, 45, 10–13, 1991.

Whiteley, N. and Faulkner, L.: Temperature influences whole-animal rates of metabolism but not protein synthesis in a temperate intertidal isopod, *Physiological and Biochemical Zoology*, 78, 227–238, 2005.

10 Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York, <http://ggplot2.org>, 2009.

Wickham, H.: The Split-Apply-Combine Strategy for Data Analysis, *Journal of Statistical Software*, 40, 1–29, <http://www.jstatsoft.org/v40/i01/>, 2011.

Witte, U., Wenzhöfer, F., Sommer, S., Boetius, A., Heinz, P., Aberle, N., Sand, M., Cremer, A., Abraham, W.-R., Jørgensen, B., et al.: In situ experimental evidence of the fate of a phytodetritus pulse at the abyssal sea floor, *Nature*, 424, 763–766, 2003.

15 Wohlers, J., Engel, A., Zöllner, E., Breithaupt, P., Jürgens, K., Hoppe, H.-G., Sommer, U., and Riebesell, U.: Changes in biogenic carbon flow in response to sea surface warming, *Proceedings of the national academy of sciences*, 106, 7067–7072, 2009.

Yvon-Durocher, G., Jones, J. I., Trimmer, M., Woodward, G., and Montoya, J. M.: Warming alters the metabolic balance of ecosystems, *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 365, 2117–2126, 2010.

Zuur, A. F., Ieno, E. N., and Elphick, C. S.: A protocol for data exploration to avoid common statistical problems, *Methods in Ecology and Evolution*, 1, 3–14, 2010.