

Dear Editor,

We have carefully read all comments from the referee. We present hereafter changes made to our manuscript to address these comments. We hope that you will be satisfied by this revision and that the paper will be accepted for publication. We kindly inform you of a change in co-author ranking. This was done to pay full credit to the work invested in the whole process to reach the present manuscript.

Referee C#1 : *However, to make this article suitable for BG, the science has to be emphasized more and the data-model comparison has to be emphasized less (as specified already in my last review). As it stands now, 12 out of 19 pages are basically a model-data comparison (as indicated in the title). While this is interesting, a model evaluation is more the subject of the journal "Geoscientific Model Development", while Biogeosciences wants to "cut across the boundaries of established sciences and achieve an interdisciplinary view" of "interactions between the biological, chemical, and physical processes in terrestrial or extraterrestrial life with the geosphere, hydrosphere, and atmosphere". Within the first 12 pages, there are a lot of numbers in the text and a lot of detailed sentences about the model-data comparison, which make the paper very lengthy and partly difficult to read. I strongly suggest to not describe too much details in the text but let figures/tables stand for themselves and just provide a summary of the most important features and/or give more details in the appendix. Try to hold technical details like the data/model comparison short and focus on the scientifically important results. The authors should point out very clearly what the scientific novelty of the paper is.*

Our Answer : In the last review, the two referees asked in fact that the science had to be emphasized more and the model-data comparison had to be emphasized less. They also asked not to limit the model-data comparison to the OVIDE section and to include a model-data comparison for air-sea fluxes. All of these requests were taken into consideration so much that the number of pages dedicated to model-data comparison has not changed compared with the first version and despite the large reduction of the initial Sections. However, we admit that there are still a lot of numbers in the text and a lot of detailed sentences that make the paper very difficult to read. To address these comments:

- i) we removed all details in Sect. 3 that were not useful to discuss results on mechanisms driving changes in the North Atlantic Cant storage rate over the last four decade presented in Sect. 5. We also substituted tables for numbers in the body of the text. Section 3 "model-data comparison" is now limited to 4^{1/2} pages instead of 6 pages.
- ii) We removed ": model-data comparison" from the title to make it more representative of the paper's content.
- iii) Results from model-data description was also removed from the abstract to focus the latter on the scientific results.

A marked-up version of the old manuscript is available with this letter. All removals are crossed out with red font color. We hope that these removals are enough to match with BG requirements.

Referee C#3: Furthermore, the paper states that that the model simulates Cant storage rate and its variability and driving processes well. I would agree that Cant storage rate and variability are well simulated and strongly disagree that the driving processes are well simulated. Driving processes of the Cant storage rate are (1) anthropogenic air-sea CO₂ flux and (2) Cant transport. The model (1) overestimates the anthropogenic air-sea CO₂ flux and even simulates to wrong phasing for the seasonal cycle north of 50N and (2) underestimates the Cant transport. I think that the authors should be careful and rather specify that their model shows the key role of transport for the Cant storage rate despite its underestimation of transport. Hence, the “real-life” transport might have an even more important role, while the anthropogenic air-sea CO₂ flux might be less important than simulated.

Our Answer: We agree with the comment: the model does not reproduce correctly driving processes for the regional Cant storage rate. However, the model reproduces correctly and for “good reasons” the interannual variability of the regional Cant storage rate because this interannual variability is driven by the transport as expected from observation and despite its large under-estimation. To clarify this message, we have re-phrased the end of Sect. 3.

Referee C#4 : The authors decided to describe the period after 1995 in “Discussion and Conclusion”, but I think this should be described in section 4.3 (for consistency). Also, as the division of the time period into “before 1995” and “after 1995” is quite important, this reasoning behind that should be described in more detail.

Our Answer: To address the first comment, the description of the period after 1995 and its associated figure (Fig. 15) has been moved from Sect. 5 to Sect. 4.3. Sect. 4.3 and Fig. 13 include now both periods (before and after 1995) described in detail. For the second comment, the division of the time period into “before” and “after” 1995 has been described in more detail in Sect. 4.3. Its discussion has also been enhanced in Sect. 5.

Referee C#4 : In general, the paper would benefit from English language editing.

Our Answer: The paper has benefit from English language editing. Please note that the final editing was not done on the marked-up version for reasons of clarity.

Referee C#5 : Specific comments:

(1) Please do not use abbreviation in the abstract without introducing them.

Our Answer: All abbreviations are now explained in the abstract.

(2) In Line 37, it reads “to supply IW then NADW” – I don’t understand what the authors mean. Consider re-phrasing.

Our Answer: This was re-phrased : “North Atlantic Central Water played a key role for storing Cant in the upper layer of the subtropical region and for supplying Cant to Intermediate Water and North Atlantic Deep Water.”.

(3) As the authors do not use the pCO₂ values from the Landschützer data-base, but the air-sea CO₂-fluxes, I would prefer if they refer to “air-sea CO₂-fluxes” in Line 163 and Line 200

Our Answer: “pCO₂ data set” was changed by “air-sea CO₂-flux data set”

(4) Line 265-268: I don't understand the calculation. If the authors want to calculate how much of the incoming Cant fluxes is stored inside the region, shouldn't the equation sum up the incoming Cant fluxes and divide by the Cant storage, i.e. $(0.156+0.044+0.092)/(0.216+0.045)$

Our Answer: This detail was removed in the revised manuscript. However, to calculate how much of the incoming Cant fluxes is stored inside the region, the equation is :
Cant storage rate (= incoming – outgoing) / incoming Cant fluxes

(5) Figure 13: for north- and southward transport, the size of the arrow is in line with the volume of the transport, this is not the case for vertical arrows. Please change this.

Our Answer: The size of all arrows is now in line with the volume of the Cant transported within Class 1 at 25°N as asked by the referee. We also add the period after 1995 (Fig. 13b). This new panel was detailed as for the period before 1995 and replaced Fig. 15.

Referee C#6: Technical Comments:

The paper would benefit from English language editing. Below is a list of mainly language errors that I spotted, but I am very sure that I have not spotted all errors.

-Figure 4: Please re-structure the figure-description. Though it is a nice figure, the description is confusing and difficult to read.

-Figure 13/15: I am sure that you meant "purple" instead of "purpose"

-Line 39: "Finally, at **the** multi-decadal scale"

-Line 40: "North Atlantic Cant storage is **rather** driven by **the** increasing air-sea fluxes"

-Line 73-74: "the **yearly** 2010's, **the region undergoes there is a decline in the** NAO index"

-Line 74: "This **has** caused"

-Line 75: "and a **slowing** down"

-Line 108: "as **follows**"

-Line 108: "are **detailed described** in"

-Line 111: "**regarding model-data comparison**"

-Line 155: "The reader is **invited to referred** to ..."

-Line 159: "Observational **data** sets"

-Line 170: There is a period missing at the end of the sentence.

-Line 229-230: "of **each component diffusive, eddy and advective terms, we only derive the advective term from the offline approach only allows for calculation of the advective term.**"

-Line 276: "our simulated transport of Cant (Fig. 3) is **nevertheless**"

-Line 288-291: Please consider rephrasing the sentence to: "Moreover, the modeled magnitude of the MOC (see Sect. S1 for details of its estimation) underestimates the observational estimate of 15.5+/- 2.3 Sv for both the month June (13.4+/-0.6 Sv) and the annual average (12.7+/-0.6 Sv)"

-Line 295-296: "ORCA-PISCES increases **the in** cumulative volume transport **of by** 15 Sv instead of 25 Sv"

-Line 306: "**It follows that Hence**"

-Line 392: "**As a consequence This implies**"

-Line 394: "**next subsequently**"

-Line 492-493: "**Since From 1985 on,** "

-Line 561: "Figure 13 also reveals **a** positive anthropogenic CO2 fluxes"

Our answer: English language errors spotted by the referee were corrected. In general, all figure-descriptions and English languages were edited. Please note that the editing was not done on the marked-up version for reasons of clarity.

1 Legend of this marked-up version of old manuscript.

2

3 ~~Crossed out section with color font in red~~ : Section removed in the revised manuscript.

4 ~~Crossed out section with color font in green~~ : Old position of section moved in the revised
5 manuscript

6 Section with color font in green : New position of section moved in the revised manuscript.

7 Section with color font in blue : New section to address referee comments.

8

9

10 **Transport and storage of anthropogenic C in the Subpolar North Atlantic :**

11 **Model—Data comparison**

12 Virginie Racapé^{1,2}, Patricia Zunino³, ~~Herlé Mercier~~³, Pascale Lherminier², ~~Herlé Mereier~~³, Laurent
13 Bopp^{1,4}, Fiz F. Pérez⁵ and Marion Gehlen¹

14

15 ¹LSCE/IPSL, Laboratoire des Sciences du Climat et de l'environnement, CEA-CNRS-UVSQ, Orme
16 des Merisiers, Bât. 712, CEA/Saclay, 91190 Gif-sur-Yvette, Cedex, France

17

18 ²IFREMER, Laboratoire d'Océanographie Physique et Spatiale, UMR 6523, CNRS-IFREMER-
19 IRD-UBO, Plouzané, France

20

21 ³CNRS, Laboratoire d'Océanographie Physique et Spatiale, UMR 6523, CNRS-IFREMER-IRD-
22 UBO, Plouzané, France

23

24 ⁴Département de Géosciences, Ecole Normale Supérieure, 24 rue Lhomond 75005 Paris

25

26 ⁵Instituto de Investigaciones Marinas, CSIC, Eduardo Cabello 6, 36208 Vigo, Spain

27

28 **Corresponding author** : Virginie Racapé, virginie.racape@ifremer.fr

29

30 **Abstract**

31 The North Atlantic Ocean is a major sink region for anthropogenic carbon (C_{ant}) and a major
32 contributor to its storage. While it is in general agreed that the intensity of the meridional
33 overturning circulation (MOC) modulates uptake, transport and storage of C_{ant} in the North
34 Atlantic Subpolar Ocean, processes controlling their recent variability and 21st century evolution
35 remain uncertain. This study aims to investigate the relationship between the transport of C_{ant}, the
36 air-sea anthropogenic CO₂ fluxes and the storage of C_{ant} in the North Atlantic Subpolar Ocean over
37 the past 44 years. It relies on the combined analysis of an annual to multi-annual in situ data set
38 and output from a global biogeochemical ocean general circulation model (NEMO/PISCES) at ½°
39 spatial resolution forced by the atmospheric reanalysis Drakkar Forcing Set 4. Despite an
40 underestimation of C_{ant} transport and an overestimation of anthropogenic air-sea CO₂ fluxes in the
41 model, ~~C_{ant} storage rate, its~~ the interannual variability of the regional C_{ant} storage rate and its
42 driving processes are well simulated. ~~At the interannual time scale, this study confirms that the~~

43 ~~time rate of changes in Cant storage in NEMO/PISCES is controlled by the divergence of the~~
44 ~~northward transport of Cant between 25°N and the Greenland-Iceland-Scotland sills.~~ Our results
45 highlight the key role played by the divergence of the NACW transport to the storage of Cant in the
46 upper oceanic layer of the subtropical region and to supply IW then NADW. In addition, this study
47 shows that Cant uptake by NADW in the lower limb of the MOC mainly occurs in the OVIDE-sills
48 box and only one quarter is exported to the subtropical region. Finally, at the multi-decadal scale,
49 the long-term changes in the north Atlantic Cant storage rate is rather driven by the increasing air-
50 sea fluxes of anthropogenic CO₂.

51

52 **1. Introduction**

53 Since the start of the industrial era and the concomitant rise of atmospheric CO₂, the ocean sink and
54 inventory of anthropogenic carbon (Cant) have increased substantially (e.g. Sabine et al., 2004; Le
55 Quéré et al., 2009; 2014; Khatiwala et al., 2013). Overall, the ocean has absorbed $28 \pm 5\%$ of all
56 anthropogenic CO₂ emissions, thus providing a negative feedback to global warming and climate
57 change (Ciais et al., 2013). Uptake and storage of Cant are, however, characterized by a significant
58 variability on interannual to decadal time scales (Le Quéré et al., 2015; Wanninkhof et al., 2013)
59 and any global assessment will hide important regional differences, which hampers the detection of
60 changes in the ocean sink in response to global warming and unabated emissions (Séférian et al.,
61 2014; McKinley et al., 2016).

62

63 The North Atlantic Ocean is a key region for Cant uptake (e.g. Sabine et al., 2004; Mikaloff-
64 Fletcher et al., 2006; Gruber et al., 2009) and stores currently as much as 20% of the total oceanic
65 inventory of 155 ± 31 PgC (Khatiwala et al., 2013). Uptake and enhanced storage of Cant in this
66 region result from the combination of two processes: (1) winter deep convection in the Labrador
67 and Irminger Seas, which efficiently transfers Cant from surface waters to the deep ocean
68 (Körtzinger et al. 1999; Sabine et al., 2004; Pérez et al., 2008) and (2) the northward transport of
69 warm and Cant-laden tropical waters by the upper limb of the meridional overturning circulation
70 (MOC; e.g. Álvarez et al., 2004; Mikaloff-Fletcher., 2006; Gruber et al., 2009; Pérez et al., 2013).
71 Both terms, deep-water formation and circulation, are characterized by high temporal variability in
72 response to the leading mode of atmospheric variability in the North Atlantic, the North Atlantic
73 Oscillation (NAO). Hurrell (1995) defined the NAO index as the normalized sea-level pressure
74 difference in winter between the Azores and Iceland. A positive (negative) NAO phase is
75 characterized by a high (low) pressure gradient between these two systems coupled to strong (weak)
76 westerly winds in the subpolar region. Between the mid-1960s and the mid-1990s, the North

77 Atlantic evolved from a negative to positive NAO phase. The change in wind conditions induced an
78 acceleration of the North Atlantic Current (NAC), as well as increased heat loss and vertical mixing
79 in the subpolar gyre (e.g. Dickson et al., 1996; Curry and McCartney, 2001; Sarafanov, 2009;
80 Delworth and Zeng, 2015). Concomitant enhanced deep convection led to the formation of large
81 volumes of Labrador Sea water (LSW) with a high load of Cant (Lazier et al., 2002; Pickart et al.,
82 2003; Pérez et al., 2008; 2013). Between 1997 and the yearly 2010's, the region undergoes a decline
83 in NAO index. This has caused a reduction of LSW formation (Yashayaev, 2007; Rhein et al., 2011)
84 and a slowing down of the northward transport of subtropical water by the NAC (Häkkinen and
85 Rhines, 2004; Bryden et al., 2005; Pérez et al., 2013). As a result, the increase in the subpolar Cant
86 inventory is below that expected from rising atmospheric anthropogenic CO₂ levels alone
87 (Steinfeldt et al., 2009; Pérez et al., 2013).

88
89 Based on the analysis of a time series of physical and biogeochemical properties between 1997 and
90 2006, Pérez et al. (2013) proposed that Cant storage rates in the subpolar gyre are primarily
91 controlled by the MOC intensity. A reduction in the MOC intensity would thus lead to a decrease in
92 Cant storage and would give rise to a positive climate-carbon feedback. The importance of MOC in
93 modulating the North Atlantic Cant inventory was previously suggested by model studies. Those
94 projected a decrease in the North Atlantic Cant inventory over the 21st century in response to a
95 projected MOC slow-down under future climate warming (e.g. Maier-Reimer et al. 1996; Crueger
96 et al., 2008; Schwinger et al., 2014). Based on the same section than Pérez et al. (2013), Zunino et
97 al. (2014) extended the time window of analysis to 1997-2010 and proposed a novel proxy for Cant
98 transport. It is defined as the difference of the Cant concentration between the upper and the lower
99 limbs of the overturning circulation times MOC intensity (see section S1 in Supplement for a
100 model-based discussion of the proxy). They observed that while the multi-annual variability of
101 transport of Cant at the OVIDE section was controlled by the variability of MOC intensity, its long-
102 term change could depend on the increase in Cant concentration in the upper limb of the MOC. As
103 the latter reflects uptake of Cant through air-sea gas exchange at the atmosphere-ocean boundary, it
104 questions the dominant role of ocean dynamics in controlling Cant storage in the subpolar gyre at
105 the decadal time scale (Pérez et al., 2013). If the storage rate of Cant in the subpolar gyre is indeed
106 at first order controlled by the load of Cant in the upper limb of the MOC, the subpolar Cant
107 inventory is expected to increase along with increasing atmospheric CO₂ - albeit not necessarily at
108 the same rate - and to provide a negative feedback on rising atmospheric CO₂ levels over the 21st
109 century.

110

111 The objective of the present study is to evaluate the relationship between Cant transport, air-sea
112 fluxes and storage rate in the Subpolar North Atlantic, along with their combined evolution over the
113 past 44 years (1958-2012). It relies on the combination of an annual to multi-annual data set
114 gathered from 25°N to the Greenland-Iceland-Scotland sills over the period 2003-2011 and output
115 from the global biogeochemical ocean general circulation model NEMO/PISCES at 1/2° spatial
116 resolution forced by an atmospheric reanalysis (Bourgeois et al., 2016). The paper is organized as
117 follow: NEMO/PISCES and *in situ* data sets are detailed in Sect. 2 and compared in Sect. 3 to
118 evaluate the model performance; main results of the interannual to decadal change of the North
119 Atlantic Cant fluxes and storage rate as well as the evaluation of their main drivers are presented in
120 Sect. 4 and discussed in Sect. 5 regarding model-data comparison.

121

122 **2. Material and methods**

123 **2.1. NEMO-PISCES model**

124 This study is based on a global configuration of the ocean model system NEMO (Nucleus For
125 European Modelling of the Ocean) version 3.2 (Madec, 2008). The quasi-isotropic tripolar grid
126 ORCA (Madec and Imbard, 1996) has a resolution of 0.5° in longitude and 0.5° x cos(ϕ) in latitude
127 (ORCA05) and 46 vertical levels whereof 10 levels lie in the upper 100m. It is coupled online to the
128 Louvain-la-Neuve sea ice model version 2 (LIM2) and the biogeochemical model PISCES-v1
129 (Pelagic interaction Scheme for Carbon and Ecosystem studies; Aumont and Bopp, 2006).
130 Parameter values and numerical options for the physical model follow Barnier et al. (2006) and
131 Timmermann et al. (2005). Two atmospheric reanalysis products, DFS4.2 and DFS4.4, were used
132 for this study. DFS4.2 is based on ERA-40 (Brodeau et al., 2010) and covers the period 1958-2007
133 while DFS4.4 is based on ERAInterim (Dee et al., 2011) and covers the years 2002-2012. The
134 simulation was spun up over a full DFS4.2 forcing cycle (50 years) starting from rest and holding
135 atmospheric CO₂ constant to levels of the year 1870 (284 ppm). Temperature and salinity were
136 initialized as in Barnier et al. (2006). Biogeochemical tracers were either initialized from
137 climatologies (nitrate, phosphate, oxygen, dissolved silica from the 2001 World Ocean Atlas,
138 Conkright et al. (2002); preindustrial dissolved inorganic carbon (C_T) and total alkalinity (A_T) from
139 GLODAP, Key et al. (2004)), or from a 3000 year long global NEMO/PISCES simulation at 2°
140 horizontal resolution (Iron and dissolved organic carbon). The remaining biogeochemical tracers
141 were initialized with constant values.

142 At the end of the spin-up cycle, two 143-year long simulations were started in 1870 and run in
143 parallel. The first one, the historical simulation, was forced with spatially uniform and temporally

144 increasing atmospheric CO₂ concentrations (Le Quéré et al., 2014). In the second one, the natural
145 simulation, the mole fraction of CO₂ was kept constant in time at 284 ppm. Both runs were forced
146 by repeating 1.75 cycles of DFS4.2 interannually varying forcing over 1870 to 1957. Then DFS4.2
147 was used from 1958 to 2007. Simulations were extended from 2002 to 2012 by switching to
148 DFS4.4. No significant differences were found in tracer distributions and Cant related quantities
149 between both atmospheric forcing products during the years of overlap (2002-2007). Carbonate
150 chemistry and air-sea CO₂ exchanges were computed by PISCES following the Ocean Carbon
151 Cycle Model Intercomparison Project protocols (www.ipsl.jussieu.fr/OCMIP) and the gas transfer
152 velocity relation provided by Wanninkhof (1992). Because climate change trends and natural modes
153 of variability are part of the forcing set used to force both simulations, potential alterations of the
154 natural carbon cycle in response to climate change (e.g. rising sea surface temperature) are thus also
155 captured by the natural simulation. The concentration of anthropogenic C, as well as anthropogenic
156 CO₂ fluxes is calculated as the difference between the historical (total C = natural + anthropogenic
157 contribution) and the natural simulations following Orr et al. (2017).

158 The global ocean inventory of Cant simulated by the model in 2010 amounted to 126 PgC. It is at
159 the lower end of the uncertainty range of the estimate by Khatiwala et al. (2013) of 155±31 PgC
160 (Fig. 1). At the global scale, the error of the model is close to 6% (values excluding arctic regions
161 and marginal seas). The mismatch between the modeled Cant inventory and that of Khatiwala et al.
162 (2013) is largely explained by the difference in the starting year of integration: 1870 for this study
163 as opposed to 1765 in Khatiwala et al. (2013). The coupled model configuration is referred to as
164 ORCA05-PISCES hereafter. The reader is invited to refer to Bourgeois et al. (2016) for a detailed
165 description of the model and the simulation strategy.

166

167

168 **2.2. Observation data sets**

169 Observations used to evaluate Cant transport computed from ORCA05-PISCES in the North
170 Atlantic Ocean were collected along the Greenland-Portugal OVIDE section and at 24.5°N
171 following the tracks presented on Fig. 2. Model output of air-sea CO₂ fluxes are compared to the
172 observation-based gridded sea surface pCO₂ product of Landschützer et al. (2015a). Programs
173 and/or data sets are briefly summarized below.

174

175 OVIDE data set

176 The OVIDE program aims to document and understand the origin of the interannual to decadal
177 variability in circulation and properties of water masses in the Subpolar North Atlantic in the

178 context of climate change (<http://www.umr-lops.fr/Projets/Projets-actifs/OVIDE>). Since 2002, one
179 spring-summer cruise is run every two years (Table 1) between Greenland and Portugal (Fig. 2)
180 Dynamical (ADCP), physical (temperature, T and salinity, S) and biogeochemical (e.g. alkalinity,
181 A_T , pH, dissolved oxygen, O_2 , and nutrients) properties are sampled at full depth hydrographic
182 stations ~~spaced by 25 nautical miles (NM). The spacing is reduced to 16 NM in the Irminger sea~~
183 ~~and to 12 NM or less over steep topographic features~~. An overview of instruments, analytical
184 methods and accuracies of each parameter is summarized in Zunino et al. (2014). The concentration
185 of C_T is calculated from pH and A_T following the recommendations and guidelines from Velo et al.
186 (2010). The OVIDE data set is distributed as part of GLODAPv2 (Olsen et al., 2016) (Table 1).

187

188 24.5°N data set

189 Data were collected along 24.5°N in 2011 between January 27th and March 15th as part of the
190 Malaspina circumnavigation expedition (<https://www.expedicionmalaspina.es/>) (Table 1). ~~A total of~~
191 ~~167 full depth hydrographic stations, whereof 13 were in the Florida Straits, were sampled along the~~
192 ~~transect, spaced by 27 NM or less across the boundary currents and topographic slopes [Hernández-~~
193 ~~Guerra et al., 2014].~~ As for the OVIDE program, ADCP, T, S, A_T , pH, O_2 and nutrients were
194 sampled during the cruise and C_T was calculated from A_T and pH. For details on methods and
195 accuracies, please refer to Hernández-Guerra et al. (2014) for dynamical and physical properties
196 and to Guallart et al. (2015) for the carbonate system. This data set is made available by **GO-SHIP**
197 ~~and delivered by~~ CCHDO (Table 1).

198

199 For both data sets, C_T is combined with T, S, nutrients, O_2 and A_T to derive the C_{ant} concentration
200 following the ϕC_T method which fix the preindustrial xCO_2 in 278.8 ppm to computed the
201 preindustrial C_T (Pérez et al., 2008; Vázquez-Rodríguez et al., 2009). This data-based diagnostic
202 approach uses water mass properties of the subsurface layer between 100-200m as reference to
203 evaluate preformed and disequilibrium conditions. The random propagation of errors associated
204 with input parameters yields an uncertainty of $5.2 \mu\text{mol kg}^{-1}$ on C_{ant} values (Pérez et al., 2010). An
205 intercomparison between different methods to separate the anthropogenic component C_{ant} from the
206 background of C_T carried out in the Atlantic Ocean (Vázquez-Rodríguez et al., 2009) and along
207 24.5°N (Guallart et al., 2015) concluded on a good agreement between ϕC_T and the other methods.

208

209 pCO_2 data base

210 The gridded sea surface pCO_2 product of Landschützer et al. (2015a) was created using the
211 SOCATv2 dataset (Bakker et al., 2014) and a 2-step neural network method detailed in

212 Landschützer et al. (2015b). It consists of monthly surface ocean pCO₂ values from 1982 to 2011 at
 213 a spatial resolution of 1°x1°. Total air-sea CO₂ fluxes were derived from equation 1 where dCO_2 is
 214 defined as the difference of CO₂ partial pressures between the atmosphere and surface ocean, K_w is
 215 the gas transfer velocity and sol , the CO₂ solubility.

$$216 \quad FCO_2^{sea-air} = K_w \times sol \times dCO_2 \quad (1)$$

217 As explain in Landschützer et al. (2014), K_w was computed as a function of wind speed following
 218 Wanninkhof (1992) rescaled to a global mean gas transfer velocity of 16 cm h⁻¹ and using winds
 219 from ERA-interim (Dee et al. 2011). sol was computed following Weiss (1974) as a function of sea
 220 surface temperature (Reynolds et al., 2002 and Hadley center EN4 sea surface salinity (Good et al.
 221 2013).

222

223

224 **2.3. Diagnostic of Cant transport and budget**

225 Transport of Cant across a section

226 The simulated transport of Cant (T_{Cant}) across a section has been evaluated either from online
 227 diagnostics (computed when the simulation is performed) or offline diagnostics (obtained after the
 228 simulation is finished, and computed using model outputs of velocities and concentrations). The
 229 transport of Cant is the sum of advective, diffusive and eddy terms. These terms are integrated
 230 vertically from bottom to surface and horizontally from the beginning (A) to the end (B) of a section
 231 along a continuous line defined by zonal (y) or meridional (x) grid segment (Fig. S2). Positive
 232 values stand for northward and/or eastward transport (see Sect. S2 in Supplement for the description
 233 of section). The advective term corresponds to the product of velocities orthogonal to the section
 234 (V) times the concentration of Cant ($[Cant]$, Eq. 2).

$$235 \quad {}^m T_{CANT}^{adv} = \int_A^B \int_{bottom}^{surface} V[Cant] dx y dz \quad (2)$$

236 The diffusive term corresponds to the transport of Cant due to the horizontal diffusion. The eddy
 237 transport is based on the parameterization of Gent and McWilliams (1990). While the online
 238 approach allows quantifying the contribution of each component, we only derived the advective
 239 term from the offline approach. We diagnosed all terms of T_{Cant} over 2003-2011, which is the only
 240 period for which the online diagnostics were available, to compare simulated T_{Cant} with the
 241 observation-based estimates from 24.5°N to the Greenland-Iceland-Scotland sills (section 3.1), and
 242 verify that the advection term was the dominant one. To study the long-term variability of Cant
 243 fluxes and storage rates (section 3.2), the time window of analysis was next extended to 1958-2012
 244 and Cant transport was derived offline from yearly averaged model outputs according to equation 2.

245 ~~These estimations were completed by the heat transport along the section computed from velocities~~
246 ~~orthogonal to the section (V) and the heat term provided by the international thermodynamic~~
247 ~~equations of seawater (TEOS 2010).~~

248

249 Budget of Cant in the North Atlantic Ocean

250 The budget of Cant was computed for three North Atlantic sub-regions (see below for definition of
251 regions). The budget was defined as the balance between i) the time rate of change in Cant,
252 vertically and horizontally integrated, ii) the incoming and outgoing transport of Cant across
253 boundaries of each region and iii) the anthropogenic air-sea CO₂ exchange, spatially integrated.
254 Budget estimates were completed by the total air-sea CO₂ flux and the heat transport over 2003-
255 2011. All terms were estimated from model output either from monthly or yearly averages
256 depending on the period analyzed (monthly for 2003-2011; yearly for 1958-2012). Relationships
257 between Cant fluxes and storage rates were investigated for each individual region.

258

259 2.4: diagnostic of heat transport

260 ~~These estimations were completed by the heat transport along the section computed from velocities~~
261 ~~orthogonal to the section (V) and the heat term provided by the international thermodynamic~~
262 ~~equations of seawater (TEOS 2010).~~

263

264 **3. Model evaluation over the period 2003-2011**

265 Figure 3 summarizes the budget of Cant in the North Atlantic simulated by the model over the
266 period 2003-2011. In order to enable the comparison of the model-derived budget to previous
267 estimates (e.g. Jeansson et al., 2011; Pérez et al. 2013; Zunino et al., 2014, 2015a,b; Guallart et al.,
268 2015), we defined two boxes separated by the Greenland-Portugal OVIDE section. The first one
269 extends from 25° N to the OVIDE section and the second box extends from the OVIDE section to
270 the Greenland-Iceland-Scotland sills. Seasonality was removed beforehand using a 12-month
271 running filter.

272 ~~In the model, over one third of Cant entering in the southern box at 25° N (0.092 ± 0.016 PgC yr⁻¹) is~~
273 ~~transported across the OVIDE section (0.035 ± 0.005 PgC yr⁻¹) and leaves the domain at the~~
274 ~~Greenland-Iceland-Scotland sills (0.034 ± 0.004 PgC yr⁻¹). The outgoing flux corresponds to a net~~
275 ~~northward transport resulting from a northwards flux across the Iceland-Scotland strait~~
276 ~~(0.053 ± 0.005 PgC yr⁻¹) and a southward flux across the Denmark strait (-0.020 ± 0.014 PgC yr⁻¹).~~
277 ~~The remainder of the regional Cant storage is provided by the air to sea exchange with the largest~~
278 ~~values south of the OVIDE section (South: 0.156 ± 0.008 PgC yr⁻¹; North 0.044 ± 0.003 PgC yr⁻¹). As~~

279 ~~a consequence, 88% of the incoming Cant flux (computed as $(0.092 + 0.156 + 0.044 - 0.034) / (0.092 +$~~
280 ~~$0.156 + 0.044)$; Fig. 3) is stored inside the region every year, predominantly south of the OVIDE~~
281 ~~section (South: 0.216 ± 0.019 PgC yr⁻¹; North: 0.045 ± 0.006 PgC yr⁻¹). In the next sections, Cant~~
282 ~~transport, anthropogenic air-sea CO₂ fluxes and Cant storage rate are successively compared to~~
283 ~~published estimates and to the observations described in section 2.2 in order to evaluate the model~~
284 ~~performance and to study the long term change in Cant storage rate and its driving processes.~~

285

286 **3.1. Advective transport of Cant**

287 In the model, over one third of Cant entering in the southern box at 25° N (0.092 ± 0.016 PgC yr⁻¹) is
288 transported across the OVIDE section (0.035 ± 0.005 PgC yr⁻¹) and leaves the domain at the
289 Greenland-Iceland-Scotland sills (0.034 ± 0.004 PgC yr⁻¹). The comparison between online and
290 offline estimates of Cant transport across the OVIDE section confirms the dominant contribution of
291 advection (Fig. S3), suggested already by Tréguier et al. (2006). Compared to previous studies, our
292 simulated transport of Cant (Fig. 3) is nevertheless clearly underestimated: it is three times smaller
293 at 25° N and at the OVIDE section (Pérez et al., 2013; Zunino et al., 2014, 2015a and b) and 1.5 to
294 2 times smaller at the sills (Jeansson et al., 2011, Pérez et al., 2013). The net Cant flux entering the
295 OVIDE box through the Denmark strait is only one third of the estimation of Jeansson et al. (2011),
296 whereas the outgoing flux at the Iceland-Scotland strait is only half. The following paragraphs focus
297 on mass transport and concentration of Cant (equation 2) in order to identify the causes of the
298 significant underestimation of modeled T_{Cant} .

299

300 Mass transport across the Greenland-Portugal OVIDE section and 25°N

301 The analysis of the stream function simulated by ORCA05-PISCES along the Greenland-Portugal
302 OVIDE section reveals a general pattern that is very similar to that estimated from observation (Fig.
303 4). The model does not, however, reproduce the interannual variability present in the observations
304 (Figs. 4a and 4b). Moreover, the magnitude of MOC (see Sect. S1 for details of its estimation)
305 computed for the month of June from model output (13.4 ± 0.6 Sv), comparable to the annual
306 average values (12.7 ± 0.6 Sv), is underestimated by around 2 Sv (dated-based estimate: 15.5 ± 2.3 Sv,
307 Mercier et al., 2015; Table 2). The upper limb of the MOC, the NAC (Lherminier et al., 2010),
308 ~~flows northeastward in the Eastern part of the section (East of 1100 km; Fig. 4b), with its modified~~
309 ~~branch, the Irminger Current, in the Western part (around 700km off the Greenland Coast) in model~~
310 ~~and data as defined by Mercier et al. (2015) (Fig. 4b). The NAC is simulated with a lower~~
311 variability and weaker intensity (Fig. 4b; ORCA05-PISCES increase in cumulative volume
312 transport of 15 Sv instead of 25 Sv between 1100km and 2500km from Greenland coast). In

313 addition, the vertical stream function (Fig. 4a) reveals a stronger current between the surface and
314 the density anomaly (σ_1) 31.5 kg m^{-3} in the model, ~~only observed at the east of the Reykjanes Ridge~~
315 ~~(not show here). This overestimation of the overturning stream function in the model is likely due to~~
316 ~~a shift in the position of the Western limit of the NAC. The Western limit is identified by close to~~
317 ~~zero values for volume transport. It occurs around 1000 km off Greenland in the model, instead of~~
318 ~~1300 km in observations (Fig. 4b).~~

319 ~~The lower limb of MOC, mainly related to the Western Boundary Current (WBC), flows southward~~
320 ~~in the western part of the section (Lherminier et al., 2007; 2010; Mercier et al., 2015). Sigma 1~~
321 ~~separating both limbs of the MOC simulated by the model is lower ($32.01 \pm 0.01 \text{ kg m}^{-3}$) than~~
322 ~~estimated from *in situ* data (32.14 kg m^{-3}). It follows that the lower (upper) limb in the model takes~~
323 ~~up a bigger (smaller) volume along the section compared to the OVIDE data set (Fig. 5).~~ The model
324 underestimates the intensity of the southward transport of the WBC in the Irminger Sea, and the
325 East Reykjanes Ridge current in the Iceland basin (Fig. 4b), ~~which are the most intense currents~~
326 ~~flowing in the lower limb of the MOC.~~ It also underestimates the cumulative volume transport for
327 $\sigma_1 > 32.40 \text{ kg m}^{-3}$ ($\sigma_0 > 27.7 \text{ kg m}^{-3}$), which is close to 0 Sv in the model (Fig. 4a) as opposed to 7 Sv
328 recorded by Lherminier et al. (2007) and García-Ibáñez et al. (2015). These densest water masses
329 correspond to lower North East Atlantic Deep Water (INEADW), Denmark Strait Overflow Water
330 (DSOW) and Iceland Scotland Overflow Water (ISOW). Taken together, the misfit between
331 observation-derived estimates and modeled volume transport is largest in the Irminger and Iceland
332 basins. This suggests that the significant underestimation of volume transport in the highest density
333 classes is probably due to the close to zero contribution of overflow waters to the transport in the
334 model at the latitude of the OVIDE section.

335 At 25°N , the upper limb of the MOC, composed by North Atlantic Central Water (NACW),
336 Antarctic Intermediate Water (AAIW) and Mediterranean Water (MW) (Talley et al., 2008;
337 Hernández-Guerra et al., 2014), flows northward with an intensity of $8.99 \pm 2.28 \text{ Sv}$ in the model
338 from January through March 2011 (Fig. 56a). The lower limb, transporting southward North
339 Atlantic Deep Water (NADW) and northward Antarctic Bottom Water (AABW; Kuhlbrodt et al.,
340 2007; Talley et al., 2008; Fig. 56b), is characterized by a net maximal flux of $-10.82 \pm 2.14 \text{ Sv}$ (Fig.
341 56a) detected at the density level (σ_1) $31.95 \pm 0.00 \text{ kg m}^{-3}$. While there is a large seasonal variability
342 (Fig. 56b), the magnitude of the winter MOC ($10.82 \pm 2.14 \text{ Sv}$) is representative of the annual value
343 in 2011 (~~11.59 ± 1.86~~ ; Table 3) and over 2003-2011 (~~11.13 ± 0.80~~ ; Table 3). The intensity of simulated
344 MOC is weaker (Table 3) and the limit between upper ~~and lower limb is shallower (Fig. 7)~~ than
345 results reported by Hernández-Guerra et al., (2014) (~~$20.1 \pm 1.4 \text{ Sv}$ at $\gamma_n = 27.82$ or $\sigma_1 = 32.27$~~) for the
346 same period, as well as reported by McCarthy et al. (2012) at 26°N between 2005 and 2008

347 (18.5±1.0 Sv). ~~McCarthy et al. (2012) highlighted nevertheless a decline in MOC intensity of 30%~~
348 ~~over the period 2009-2010 mainly due to the increase in the southward upper ocean recirculation~~
349 ~~(shallower than 1100m) and the decrease in the southward transport of lower (1)NADW.~~ INADW is
350 essentially made up of Nordic overflow waters (Pickart, 1992; Smethie et al., 2000), which the
351 model fails to reproduce correctly. The preceding suggests that the underestimation of the volume
352 transport in the model is likely due to the large underestimation of dense overflow waters.

353

354 Cant distribution in the North Atlantic Ocean and along the OVIDE section and 25°N

355 Compared to the observation-based product of Khatiwala et al. (2013), ~~both minimum and~~
356 ~~maximum~~ Cant concentrations simulated by ORCA05-PISCES are relatively well represented from
357 25°N to the Greenland-Iceland-Scotland sills (Fig. 1). ~~Minimum values are found in the subtropical~~
358 ~~region whereas the maximum values are simulated in the subpolar gyre, especially in the Labrador~~
359 ~~Sea.~~ Figure 1 points nevertheless to an under-estimation of up to 40 molC m⁻² of modeled maxima.
360 The comparison between modeled and observed Cant along the Greenland-Portugal OVIDE section
361 and 25°N reveals a comparable distribution ~~with higher concentrations in surface waters and lower~~
362 ~~levels at depth~~ (Figs. 56 and 7). ~~The surface to depth gradient is more pronounced in the Eastern~~
363 ~~basin of two sections.~~ Along the OVIDE section (Figs. 5a and b), the two LSW cores, relatively rich
364 ~~in Cant, are identified on the two sides of the Reykjanes Ridge.~~ Despite the good agreement of
365 spatial patterns, modeled concentrations are lower by 6.3±0.6 μmol kg⁻¹ compared to observed-
366 based estimates (Table 2). ~~Half of this underestimation is due to the preindustrial atmospheric CO₂~~
367 ~~condition used by the model (284 ppm) compared to pCT method (278.8 ppm).~~ This deficit is more
368 pronounced in the upper limb of MOC ($\Delta\text{Cant}^{\text{model-data}} = -5.9\pm 0.7 \mu\text{mol kg}^{-1}$) than in the lower limb
369 ($\Delta\text{Cant}^{\text{model-data}} = -3.6\pm 0.6$, Table 2). The largest difference between model and data (up to -20 μmol
370 kg⁻¹, Fig. 5c) is detected in subsurface waters at the transition between East North Atlantic Central
371 Water (ENACW) and Mediterranean Water (MW) and between both limbs of the MOC.

372 The variability of the model-data difference, ~~diagnosed as its standard deviation, peaks at 10 μmol~~
373 ~~kg⁻¹~~ (Fig. 56d). It is largest at the boundary between upper and lower limbs of the MOC, mainly
374 between 700 km to 2000 km off Greenland. The higher variability in this region could be explained
375 by the variability of the NAC intensity, which is underestimated by ORCA05-PISCES.

376 Figure 5 also reveals an underestimation by the model of Cant levels in NEADW1 (below 3500m
377 depth in the western European basin) by 5 to 10 μmol kg⁻¹ which is in line with a close to zero
378 contribution of dense Cant rich overflow waters along the OVIDE section.

379 At 25°N, ~~a subsurface pool of Cant is detected in the western part of the section in both products~~
380 ~~(Figs. 7a and b) around 1500m depth, albeit with smaller concentrations in the model.~~ The model

381 underestimates the Cant concentration, especially in the lower limb of the MOC ~~with mean values~~
382 ~~of $2.89 \pm 0.09 \mu\text{mol kg}^{-1}$ compared to $12.00 \mu\text{mol kg}^{-1}$ calculated from observations~~ (Table 3). The
383 largest difference between ORCA05-PISCES and observations, up to $-30 \mu\text{mol kg}^{-1}$, is found
384 around 500m depth in the upper limb of the MOC. Finally and like along the OVIDE section, Fig. 7
385 reveals an under-estimation of Cant levels below 3500m depth by about $10 \mu\text{mol kg}^{-1}$. This water
386 mass corresponds to AABW ~~that becomes NEADW during its northward transport by mixing with~~
387 ~~INADW (Talley et al., 2008).~~

388

389 From the preceding follows that the underestimation of Cant transport in ORCA05-PISCES is likely
390 due to the underestimation of water mass transport intensity (mainly attributed to a too weak
391 contribution of dense overflow waters) and of Cant concentrations. ~~Half of this underestimation is~~
392 ~~due to the preindustrial atmospheric CO_2 condition used by the model (284 ppm) compared to ϕCT~~
393 ~~method (278.8 ppm).~~

394

395 ~~The hypothesis is supported by the analysis of the heat transported at 25°N and the OVIDE section,~~
396 ~~which is also underestimated by the model (Fig. 3) compared to Pérez et al (2013). Pérez et al.~~
397 ~~(2013) estimated a heat transport of $1.10 \pm 0.01 \text{PW}$ and $0.59 \pm 0.09 \text{PW}$ at 25°N and OVIDE,~~
398 ~~respectively, while the model yields a corresponding heat transport of $0.78 \pm 0.06 \text{PW}$ and 0.39 ± 0.02~~
399 ~~PW . The discrepancy between model and observation based estimates of heat transport is, however,~~
400 ~~not as large as for the advective transport of Cant, probably due to a better representation of~~
401 ~~temperature than Cant concentration by the model (mean model-data bias along the section:-~~
402 ~~$0.4 \pm 0.9^\circ\text{C}$ for a mean value of 5°C (8% of error) for temperature, $7 \mu\text{mol kg}^{-1}$ for a mean value of~~
403 ~~$25.4 \mu\text{mol kg}^{-1}$ (27%) for Cant).~~

404

405 **3.2. Air-sea fluxes of total and anthropogenic CO_2**

406 Estimates of modeled air-sea fluxes of total and anthropogenic CO_2 are higher than those derived
407 from *in situ* data by Pérez et al. (2013) : ~~Southern box: model = (anth) $0.156 \pm 0.008 \text{PgC yr}^{-1}$ /~~
408 ~~(total) $0.303 \pm 0.013 \text{PgC yr}^{-1}$, Pérez et al. (2013) = (anth) $0.12 \pm 0.05 \text{PgC yr}^{-1}$ / (total) 0.20PgC yr^{-1} ;~~
409 ~~Northern box: model = (anth) $0.044 \pm 0.003 \text{PgC yr}^{-1}$ / (total) $0.103 \pm 0.006 \text{PgC yr}^{-1}$, Pérez et al.~~
410 ~~(2013) = (anth) $0.016 \pm 0.012 \text{PgC yr}^{-1}$ / (total) 0.09PgC yr^{-1} . While the model overestimates CO_2~~
411 uptake, the ratio anthropogenic/natural is comparable to Gruber et al. (2009) and Schuster et al.
412 (2013). As a consequence, the model overestimates both natural and anthropogenic components
413 with quite similar proportion. To understand the large over-estimation of fluxes, simulated average
414 air-sea fluxes of total CO_2 over the period 2003-2011 are next compared to estimates by

415 Landschützer et al., (2015a), taken as a representative observation-based product from the SOCOM
416 exercise (Rödenberk et al. 2015). ~~The area extending from 25°N to the Greenland-Iceland-Scotland~~
417 ~~sills is a sink for atmospheric CO₂ in the model and the data-based product (Fig.8). Three areas~~
418 ~~present nevertheless differences from observations. The first one is located south of Newfoundland~~
419 ~~and centered at 35°W-45°N. In this region, which corresponds to the NAC path in the observations~~
420 ~~(see figure 1 in Daniault et al., 2016), modeled total air-sea CO₂ fluxes are around 0 molC m² yr⁻¹~~
421 ~~compared to values up to -3.5 molC m² yr⁻¹ reported in Landschützer et al. (2015a). The second area~~
422 ~~is found close to the Western African coast, where the model simulates a CO₂ source to the~~
423 ~~atmosphere shifted to the north and extending more to the west along 25°N than in observations.~~
424 The third zone that differs from observations is the northern box between the OVIDE section and
425 the Greenland-Iceland-Scotland sills. Here, the modeled oceanic CO₂ sink is overestimated in
426 average by a factor of 2 to 3. Panels 8c and 8d show the month of the maximum, respectively
427 minimum value of air-sea CO₂ flux for the period 2003-2011. It reveals a seasonal phase shift
428 between ORCA05-PISCES and Landschützer et al. (2015a), north of 50°N where the model over-
429 estimates strongly gas exchange. Fluxes peak in winter in observations while they are at a
430 maximum in summer in the model. According to Takahashi et al. (2002), the seasonal change in
431 surface water pCO₂ is dominated by the biological effect north of 40°N and by the temperature (or
432 thermodynamic) effect between 20°N and 40°N. The main driving process of seasonal variability of
433 air-sea CO₂ fluxes is well reproduced by the model in the subtropical region. However, the
434 dominant effect of temperature extends too far north in the model. As a result, the seasonal change
435 in CO₂ fluxes is dominated by the thermodynamic effect in the subpolar gyre. Despite the seasonal
436 phase shift noted in the subpolar gyre, the amplitude of the interannual variability of total air-sea
437 CO₂ fluxes (standard deviation of the 1982-2011 time series without seasonality, Fig. 9) is well
438 reproduced by the model over the total domain and even north of 40°N where the variability is the
439 largest.

440

441 **3.3. Storage rate of Cant**

442 ~~As a consequence, 88% of the incoming Cant flux is stored inside the region every year,~~
443 ~~predominantly south of the OVIDE section (Fig. 3). The storage rates of Cant estimated for the~~
444 ~~period 2003-2011 are close to the estimates from Pérez et al. (2013), referenced to 2004: Southern~~
445 ~~box: ~~model = -0.216 ± 0.019, Pérez et al. (2013) = 0.280 ± 0.011~~; Northern box: ~~model = -0.045 ±~~~~
446 ~~0.006 and Pérez et al. (2013) = 0.045 ± 0.004 PgC yr⁻¹.~~

447 These results suggest that there may be a compensation in the model between the underestimation
448 of Cant transport and the overestimation of anthropogenic air-sea CO₂ fluxes detailed above.

449 Next, the contribution of air-sea uptake and transport of Cant to the variability of the North Atlantic
450 Cant inventory is derived for each box from the analysis of multi-annual time series of
451 anthropogenic air-sea CO₂ fluxes, transport divergence of Cant (defined as the difference between
452 incoming and outgoing Cant fluxes at the borders of the boxes) and Cant storage rate. Time series
453 were smoothed as explained previously and the potential trends were removed. Correlation
454 coefficient (r) and p-value are summarized in table 4. Our results suggest that, over the period 2003-
455 2011, the rate of Cant storage between 25° N and the Greenland-Iceland-Scotland sills is strongly
456 correlated with a positive transport divergence of Cant (~~25° N: r=0.96, p-value=0.00; OVIDE: r=~~
457 ~~0.95, p-value=0.00~~). The dominance of Cant transport divergence over gas exchange is
458 corroborated by observation-based assessments (Pérez et al., 2013; Zunino et al., 2014; 2015a and
459 b). Despite an underestimation of Cant transport and an overestimation of anthropogenic air-sea
460 CO₂ fluxes, modeled storage rate, ~~its~~ variability and driving processes are coherent with
461 observations allowing the simulation to be used to study drivers of changes in Cant storage rate
462 since 1958.

463

464

465 **4. Cant fluxes and storage rate in the North Atlantic Ocean (North of 25°N) since 1958**

466 In this section, we present the analysis of the full period covered by our simulations (1958-2012)
467 with the objective of better understanding the interannual to decadal variability of the Cant
468 inventory in the North Atlantic Ocean as well as its driving processes. The study area, from 25°N to
469 the Greenland-Iceland-Scotland sills, is divided in 3 boxes instead of 2 in section 3: the first box
470 extends from 25°N to 36°N; the second box from 36°N to the OVIDE section and the third box is
471 between the OVIDE section and the Greenland-Iceland-Scotland sills. The section 36°N was added
472 to delimit the northern part of the subtropical region from the Subpolar gyre (Mikaloff-Fletcher et
473 al., 2003).

474

475 **4.1. Contribution of variability of circulation and Cant accumulation on Cant transport** 476 **variability**

477 Figure 10 presents annual time series (1958-2012) of the magnitude of the MOC and transports of
478 heat and Cant at 25°N, 36°N and across the OVIDE section. The heat transport and the MOC
479 intensity are strongly correlated at each section (~~25°N, r=0.92, p-value=0.00; 36° N, r=0.90, p-~~
480 ~~value=0.00; OVIDE, r=0.76, p-value=0.00~~) whereas a significant relationship between the MOC
481 strength and the Cant transport is only found at 36°N (~~25°N, r=0.30, p-value=0.02; 36°N, r=0.67,~~
482 ~~p-value=0.00; OVIDE, r=0.02, p-value=0.90~~). As expected, the circulation is thus the major

483 driver of interannual to decadal variability of heat transferred across these sections (Johns et al.,
 484 2011, Mercier et al., 2015). Its impact on the variability of Cant transport is, however, masked by
 485 several other mechanisms. The transport of Cant across the three sections is characterized by a
 486 continuous increase over the period of study (Fig. 10): ~~it increases from $0.030 \pm 0.002 \text{ PgC yr}^{-1}$ in~~
 487 ~~1958-60 to $0.095 \pm 0.024 \text{ PgC yr}^{-1}$ in 2010-12 at 25°N , from $0.009 \pm 0.001 \text{ PgC yr}^{-1}$ to 0.050 ± 0.018~~
 488 ~~PgC yr^{-1} at 36°N and from $0.008 \pm 0.001 \text{ PgC yr}^{-1}$ to $0.043 \pm 0.005 \text{ PgC yr}^{-1}$ at the OVIDE section.~~
 489 Such a large increase is observed neither on the heat transport (~~$0.0003 \pm 0.0004 \text{ PW yr}^{-1}$ at 25°N ,~~
 490 ~~$0.0016 \pm 0.0004 \text{ PW yr}^{-1}$ at 36°N and $0.0003 \pm 0.0002 \text{ PW yr}^{-1}$ at OVIDE)~~ nor on the MOC
 491 magnitude (~~$0.001 \pm 0.005 \text{ Sv yr}^{-1}$ at 25°N , $0.015 \pm 0.006 \text{ Sv yr}^{-1}$ at 36°N and $0.003 \pm 0.007 \text{ Sv yr}^{-1}$ at~~
 492 ~~OVIDE), nor on the net volume of water transported across the sections (~~$-0.000 \pm 0.000 \text{ Sv yr}^{-1}$ at~~
 493 ~~25°N , $0.001 \pm 0.001 \text{ Sv yr}^{-1}$ at 36°N and $-0.000 \pm 0.003 \text{ Sv yr}^{-1}$ at OVIDE). Following Zunino et al.
 494 (2014), we conclude that the increase in the northward transport of Cant since 1958 was mainly due
 495 to Cant accumulation in the northward flowing upper limb of the MOC. In order to isolate the effect
 496 of circulation, we removed the positive trend from Cant transport time series. The correlation (r)
 497 between the detrended Cant transport and the magnitude of the MOC increased from 0.30 (~~p -value =~~
 498 ~~0.02) to 0.74 (p -value = 0.00) at 25°N and from 0.67 (~~p -value = 0.00) to 0.70 (p -value = 0.00) at~~
 499 36°N . It did not change at the OVIDE section ($r=0.1$, p -value = 0.4). We conclude that the
 500 circulation controls the interannual to decadal variability of Cant transport but only at 25°N and
 501 36°N . ~~In the following section, we study the impact of circulation on Cant storage rate regarding the~~
 502 ~~Cant transport divergence.~~~~~~~~

504 **4.2. Interannual to decadal variability of the North Atlantic Cant inventory**

505 Figure 11 provides the budget of Cant from 1959 to 2011 for the three boxes. Each budget is
 506 composed of the Cant storage rate, the anthropogenic air-sea CO_2 flux and the transport divergence
 507 of Cant. We observe a continuous increase in the North Atlantic Cant inventory over the last 44-
 508 years, especially in box 2 (36°N -OVIDE) where the storage rate is multiplied by 3 (~~from~~
 509 ~~$0.043 \pm 0.000 \text{ PgC yr}^{-1}$ (1959-1961) to $0.127 \pm 0.010 \text{ PgC yr}^{-1}$ (2009-11))~~ and in box 1 (25°N - 36°N)
 510 where it doubled (~~from $0.039 \pm 0.000 \text{ PgC yr}^{-1}$ to $0.094 \pm 0.004 \text{ PgC yr}^{-1}$~~). Taking into account the
 511 anthropogenic perturbation in the surface layer and assuming the transient steady-state, we expected
 512 a factor of 2.9 that is in line with and validate our result in box 2. Air-sea flux of Cant and Cant
 513 transport divergence contribute equally to changes in Cant inventory in the southern box. Between
 514 36°N and the OVIDE section, the contribution of gas exchange dominates prior to 1985. Since
 515 1985, the transport divergence gained in importance, albeit with a pronounced interannual
 516 variability. In the northern box, changes in Cant inventory follow air-sea fluxes (weak contribution

517 of transport divergence limited to interannual variability).

518 The significant positive correlation (Table 56a, no trend removed) between storage rate and air-sea
519 gas exchange in all three boxes suggests the latter to be, over the past 42 years, a main control of
520 Cant storage rate on the longer time scales. Nevertheless, the transport divergence of Cant in the
521 southern box and between 36°N and the OVIDE section from 1985 onward, which increased
522 continuously over the period, also correlates with the change in Cant storage rate (Table 56a, trend
523 included). It did not however influence the long-term change in Cant inventory between the OVIDE
524 section and the Greenland-Iceland-Scotland sills (OVIDE-Sills; $r = 0.32$, p -value = 0.02; Table 56),
525 where it is close to zero (incoming $T_{\text{Cant}} =$ outgoing T_{Cant}). In this analysis (correlation with trend
526 included), the trend in response to increasing atmospheric CO₂ levels dominates the signal and the
527 correlation at the expense of interannual variability. In order to identify controls of the interannual
528 variability, the analysis was repeated with detrended time series. It reveals a strong correlation
529 between the Cant storage rate and the transport divergence of Cant for all three boxes (Table 56b),
530 as opposed to correlation with air-sea gas exchange which is either not significant or weak (Table
531 56b). The model output analysis suggests that while the long term changes in Cant storage rate are
532 controlled by anthropogenic air-sea CO₂ fluxes, its interannual variability is on the contrary driven
533 by the transport divergence of Cant. Additional analyses are made to identify which role is played
534 by the circulation in the annual evolution of Cant storage rate. In this context, we estimated for each
535 box the correlation between the detrended time series of Cant transport divergence and the
536 incoming and outgoing transport of Cant. These estimates, summarized in table 67, show that the
537 transport divergence of Cant is always correlated with the incoming transport of Cant and not with
538 the outgoing transport of Cant. Results of this section suggest that the interannual variability of the
539 North Atlantic Cant storage rate is driven by the transport of Cant coming from south latitude.
540 According to Sect. 4.1, the interannual changes of both terms at 25°N and 36°N depends on MOC
541 intensity. These results corroborate the conclusion of section 3.3 for the period 2003-2011 and are in
542 line with previous studies (Pérez et al., 2013; Zunino et al., 2014).

543

544 **4.3. Contribution of advection of water masses to the storage rate of Cant**

545 In this section, we analyze major water masses taking part to the upper and lower limb of the MOC
546 in order to identify their contributions to the regional Cant storage rate over the period 1958-2012.
547 The general circulation from 25°N to the Greenland-Iceland-Scotland sills is well documented (e.g.
548 Arhan, 1990; McCartney, 1992; Hernández-Guerra et al., 2015; Danialt et al., 2016). Based on
549 these studies and the water column distribution of zonally integrated mass transport at 25°N, 36°N,
550 OVIDE and the Greenland-Iceland-Scotland sills (Fig. 12), we identify three water classes : North

551 Atlantic Central Water (NACW, Class 1), Intermediate waters (IW; Class 2) and North Atlantic
552 Deep Water (NADW, Class 3).

553 **NACW (Class 1)** is transported by the upper ocean circulation, either northward (Class 1N) by the
554 Gulf Stream and the NAC, or southward (Class 1S) by the subtropical gyre recirculation in the
555 western European basin. The southeastward recirculation is characterized by cool and dense waters
556 (Talley et al., 2008) allowing distinction of Class 1S from Class 1N in our study (Fig. 12). NACW
557 loses heat during its northward journey and becomes denser. As a result, its density limit changes
558 with latitude (Fig. 12). Based on Fig. 12, we define the class 1N from surface to the density
559 anomaly $\sigma_1 = 29.1 \text{ kg m}^{-3}$ at 25°N , 30 kg m^{-3} at 36°N and 31 kg m^{-3} at the OVIDE section. This
560 class is not found at the Greenland-Iceland-Scotland sills. The class 1S, proper to the subtropical
561 region, is found from 29.1 kg m^{-3} to 31 kg m^{-3} at 25°N and from 30 kg m^{-3} to 31 kg m^{-3} at 36°N .
562 **IW (Class 2)** encompasses the densest water masses of the upper MOC limb, such as Antarctic
563 Intermediate Water (AAIW), Subantarctic Intermediate Water (SAIW) or Mediterranean Water
564 (MW). The class 2 circulates northward between $\sigma_1 = 31$ and 31.8 kg m^{-3} from 25°N to OVIDE and
565 between $\sigma_1 = 31$ and 31.9 kg m^{-3} through the Greenland-Iceland-Scotland sills (Fig. 12).
566 **NADW (Class 3)** supplies the lower limb of the MOC. It flows southward from the subpolar gyre
567 to the subtropical region. In the model, it is found below $\sigma_1 = 31.7\text{-}31.9 \text{ kg m}^{-3}$ (Fig. 12).

568
569 The long term changes in simulated volume and Cant transports for these three specified classes
570 across the four sections highlight two periods, before and after 1995. The distinction between these
571 two periods is based on Class 1N (northward NACW) at the OVIDE section and Class 2 (IW) at
572 36°N where both Cant and volume transport increased after 1995 (Fig. S4). No remarks are reported
573 on Cant storage rate in previous section. Based on these comment, we focus this section on the
574 period 1958-1994 to understand how each water mass contributes to Cant storage rate. The period
575 1996-2011 is discussed in Sect. 5 to understand causes of the increase in volume and Cant transports
576 after 1995. Results for the first period (1958-1994) are summarized on Fig. 13.

577
578 **Before 1995**, more than 50% of Cant transported by NACW flowing northward (Class 1N) at 25°N
579 crossed 36°N whereas 30% recirculated southward with Class 1S. At the OVIDE section, the
580 transport of Cant was equal to 12% of 25°N , whereas it is close to zero at the sills (Fig. 13). The
581 transport divergences of Cant for Class 1 in Box 1 ($0.034 \text{ PgC yr}^{-1} = 0.096 - 0.056 + 0.022 - 0.028$), Box
582 2 ($0.022 \text{ PgC yr}^{-1} = 0.056 - 0.012 - 0.022$) and Box 3 ($0.012 \text{ PgC yr}^{-1} = 0.012 - 0.000$) are positive and
583 higher than Cant storage rate (Fig. 13). Figure 13 also reveals a positive anthropogenic CO_2 fluxes
584 from atmosphere to surface Ocean. The Cant budget of Class 1 for each box suggests in fact a

585 vertical transport of Cant from Class 1 to Class 2. Our results from this section and Sect. 4.2
586 indicate that the NACW plays a key role in the Cant storage rate between 25°N and the OVIDE
587 section but also in the Cant transfer into the lower layer during its northward transport. This cross-
588 isopycnal transport evidenced between Class 1 and Class 2 during its northward journey (Fig. 13) is
589 related to a large decrease in the northward transport in Class 1 associated with a large increase in
590 the northward transport in Class 2 from 25°N to the OVIDE section (Fig. S4). This is in line with
591 results from De Boisséson et al. (2012) who highlight the densification of subtropical central water
592 by winter air-sea cooling and mixing with intermediate waters along the NAC path. Moreover, our
593 results from Cant transport (Fig. 13) also suggest that IW is enriched in Cant between 25°N and the
594 OVIDE section over the studied period. The large Cant uptake north of 36°N is explained by
595 regional winter deep convection that occurs along the NAC that mixed NACW, rich in Cant, with
596 IW, poor in Cant. Figure 13 also shows that 64% of Cant entering into Box 3 by advection and air-
597 sea gas exchange is exported southward by Class 3, 20% is stored whereas 16% is exported
598 northward through the Greenland-Iceland-Scotland sills by Class 2. In addition, the budget of Cant
599 computed for Class 2 reveals a significant vertical transport of Cant from IW to NADW, especially
600 north of the OVIDE section. NADW is thus enriched in Cant from NACW/IW essentially between
601 the OVIDE section and the Greenland-Iceland-Scotland sills, which is in agreement with results
602 from Sarafanov et al. (2012). Finally, a small fraction of Cant entering in Box 2 within Class 3
603 leaves the area across 25°N (24%, Fig. 13). The remainder is stored within Class 3 between 25°N
604 and OVIDE.

605
606 **After 1995:** , 27% of Cant entering within Class 1 at 25°N flowed northward across the OVIDE
607 section, that is two times higher than for the previous period (Fig. 13b). As revealed above, this
608 relative increase in Cant transport at OVIDE was associated with a significant increase in volume
609 transport across the section (Fig. S4a). This latter was in fact multiplied by 1.9 after 1995 to the
610 detriment of the diapycnal transport between Class 1 and Class 2 waters that decreased of 60%
611 compared to the previous period, decreasing thus Cant transferred from NACW to IW. Fig. 13b
612 shows that these results of Class 1-Box 2 were also concomitant with a relative but smaller decrease
613 in air-sea flux and in the net Cant transport across 36°N. In Class 1 of Box 3, the relative increase in
614 Cant transport at OVIDE was concomitant with a similar increase in the contribution of the vertical
615 transport of Cant to Class 2 waters as well as with a small decrease in the contribution of air-sea
616 flux. Moreover, the relative increase in Cant transferred into Class 2 (Box 3) is associated to a
617 relative increase in Cant transported within Class 2 waters throughout the Nordic sills, in Cant
618 transported vertically into Class 3 waters and in the regional Cant stored inside the box (Class 2

619 Box 3) but also to a relative decrease in the Cant transport of Class 2 at OVIDE. The excess of
620 NACW rich in Cant entering in the OVIDE-sills box was transferred into IW before being exported
621 to the Nordic regions or stored in the subpolar gyre.

622

623

624 **5. Discussion and Conclusion**

625 The model-data comparison highlights a large underestimation (by 2 or 3 times) of the Cant
626 transport by the model, resulting from an underestimation of both volume transport and Cant
627 accumulation in the water column. The underestimation of the volume transport is likely due to the
628 too small contribution of overflow waters. Their misrepresentation leads to an underestimation of
629 the intensity of the lower limb of the MOC and as a consequence, of that of the upper branch. It
630 results a smaller than expected export of Cant from the subtropical region to the subpolar gyre. The
631 insignificant southward flow of overflow waters also contributes to make the net export of Cant to
632 the Arctic region relatively large (outgoing flux at Iceland-Scotland Ridge is only divided by 2
633 while incoming flux at Denmark Strait is divided by 3 compared to observations). Our analysis also
634 reveals a strong overestimation of the modeled air-sea anthropogenic CO₂ exchange. This
635 discrepancy is associated with a larger total CO₂ uptake by the ocean north of the OVIDE section.
636 Moreover, we observe an overestimation of the modeled anthropogenic CO₂ flux. North of 40°N,
637 this overestimation of the total air-sea CO₂ flux is partially due to a seasonal change dominated by
638 the thermodynamic effect rather than the biological effect. While anthropogenic CO₂ exchange as
639 defined in the model is not impacted by the biological activity, thermodynamic mechanism affect
640 positively anthropogenic CO₂ fluxes. The overestimation of the modeled anthropogenic air-sea CO₂
641 fluxes could also be a response to the low Cant concentration in the North Atlantic surface Ocean
642 due to the model initial condition and the small Cant fraction transported inside the subpolar gyre
643 that enhanced the air-sea anthropogenic pCO₂ gradient. These results are clearly a limit of the
644 model that underestimates the contribution of Cant transport to storage rate. This is especially true
645 for the OVIDE-Sills box where we observe an unexpected transport divergence close to zero (no
646 contribution) along with an overestimation of the air-sea flux. The modeled Cant storage rate is,
647 however, in line with data-based estimates that reflect a compensation between the underestimation
648 of Cant transport and the overestimation of air-sea gas exchange. The spatial distribution of Cant
649 storage is well reproduced by the model. In line with independent studies (Sabine et al., 2004;
650 Khatiwala et al., 2013), the North Atlantic Ocean, north of 25°N, acts as a sink for the atmospheric
651 anthropogenic CO₂, a large part of which being stored between 36°N and the OVIDE section.
652 Moreover, mechanisms controlling the interannual to decadal changes in Cant storage rate as well

653 as Cant and heat transport match with data-based estimates (Pérez et al., 2013; Zunino et al., 2014,
654 2015b; Johns et al., 2011). The satisfying reproduction of interannual variability by the model
655 allowed its use to explore the interannual to multidecadal changes in the North Atlantic Cant
656 inventory and its driving processes.

657 At the interannual time scale, the time rate of change of Cant storage in the model is controlled by
658 the divergence of the northward transport of Cant in the region between 25°N and the Greenland-
659 Iceland-Scotland sills, similarly to the data-based results reported by Pérez et al. (2013) and Zunino
660 et al. (2014; 2015b). At the OVIDE section, the interannual variability of Cant transport is
661 controlled by Cant accumulation in the upper MOC limb whereas it is also influenced by the MOC
662 magnitude at 25°N and 36°N. Additional analysis of the Cant transport in density classes highlights
663 the key role played by the divergence of the NACW transport to the storage of Cant in the upper
664 oceanic layer of the subtropical region and to supply IW then NADW. These water mass
665 conversions are consistent with previous study (Sarafanov et al., 2012; De Boisséson et al., 2012;
666 Pérez et al., 2013). The Cant uptake by Class-3 in the lower limb of the MOC mainly occurs in the
667 OVIDE-sills box. ~~A significant correlation between the volumes of NADW transported across the~~
668 ~~OVIDE section and the NAO winter index is highlighted (Fig. 14; $r = 0.68$, $p\text{-value} = 0.00$).~~ A
669 positive (negative) anomaly of volume transport is associated with a positive (negative) NAO index.
670 Previous studies also reported an acceleration of the NAC during the transition phase period (e.g.
671 Dickson et al., 1996; Curry and McCartney, 2011). The increase in transport of the NAC is well
672 reproduced by the model with the anomaly of NACW mass transport being correlated with the
673 NAO winter index (Fig. 14). This study also highlights a specific period before and after 1995
674 likely to explain the lack of correlation. ~~According to Fig. 15 and S4, the period after 1995 is~~
675 ~~characterized by i) an increase in the transport of Cant and volume through the OVIDE section by~~
676 ~~NACW, ii) an increase in IW production between 25°N and 36°N but a decrease between 36°N and~~
677 ~~OVIDE associated with iii) an increase in NACW recirculation at 36°N. In the other word, since~~
678 ~~1995, we observed more Class 1 rich in Cant advected through the OVIDE section.~~ As shown in
679 Fig. 165, the subpolar gyre undergoes a warming of its mixed layer since 1995. Such warming was
680 reported by De Boisséson et al. (2012) for the year 1998. Authors explained this by an increase in
681 the inflow of subtropical water into the Iceland basin. This enhanced advection of subtropical water
682 into the subpolar gyre could explain the decreasing contribution of anthropogenic air-sea CO₂
683 fluxes to Cant storage in favor of the advective transport of Cant reported in Sect. 4.2 between 36°N
684 and the OVIDE section. Warm, alkalinity rich subtropical waters carry a relatively high load of Cant
685 and their enhanced northward advection decrease the air-sea gradient of anthropogenic pCO₂ and
686 slow down air-sea gas exchange (Thomas et al., 2008).

687 To conclude, at the multi-decadal time scale, the long term change in anthropogenic air-sea CO₂
688 fluxes over the whole domain exert the dominant control on the Cant inventory of the North
689 Atlantic subpolar gyre. The contribution of Cant transport from 25°N across the OVIDE section
690 emerges as the important driver on interannual to decadal time scales through its divergence. Our
691 model analysis suggests that assuming unabated emissions of CO₂, the storage rate of Cant in the
692 Subpolar North Atlantic is expected to increase assuming MOC fluctuations within observed
693 boundaries. However, under a future strong decrease in MOC in response to global warming (IPCC
694 projection 25%, Collins et al., 2013) the storage rate might decrease.

695

696

697 **References**

- 698 Álvarez, M., Pérez, F. F., Bryden, H., & Ríos, A. F. : Physical and biogeochemical transports
699 structure in the North Atlantic subpolar gyre, *J Geophys Res: Oceans*, 109(C3), 2004.
- 700 Arhan, M.: The North Atlantic Current and Subarctic Intermediate Water, *J. Mar. Res.*, 48(1), 109-
701 144, doi: 10.1357/002224090784984605, 1990.
- 702 Aumont, O., and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global*
703 *Biogeochem Cy*, 20(2), 2006.
- 704 Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S.,
705 Kozyr, A., Nojiri, Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl,
706 J., Barbero, L., Bates, N. R., Boutin, J., Bozec, Y., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen,
707 L., Chierici, M., Currie, K., de Baar, H. J. W., Evans, W., Feely, R. A., Fransson, A., Gao, Z.,
708 Hales, B., Hardman-Mountford, N. J., Hoppema, M., Huang, W.-J., Hunt, C. W., Huss, B.,
709 Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S. D., Jutterström, S., Kitidis, V., Körtzinger,
710 A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat, L.,
711 Metzl, N., Murata, A., Newberger, T., Omar, A. M., Ono, T., Park, G.-H., Paterson, K., Pierrot,
712 D., Rios, A. F., Sabine, C. L., Saito, S., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R.,
713 Skjelvan, I., Steinhoff, T., Sullivan, K. F., Sun, H., Sutton, A. J., Suzuki, T., Sweeney, C.,
714 Takahashi, T., Tjiputra, J., Tsurushima, N., van Heuven, S. M. A. C., Vandemark, D., Vlahos, P.,
715 Wallace, D. W. R., Wanninkhof, R., and Watson, A. J.: An update to the Surface Ocean CO₂
716 Atlas (SOCAT version 2), *Earth Syst. Sci. Data*, 6, 69–90, doi:10.5194/essd-6-69-2014, URL:
717 <http://www.earth-syst-sci-data.net/6/69/2014/>, 2014.
- 718 Barnier, B., Madec, G., Penduff, T., Molines, J. M., Treguier, A. M., Le Sommer, J., Beckmann, A.,
719 Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C.,
720 Theetten, S., Maltrud, M., McClean, J., and De Cuevas, B.: Impact of partial steps and

721 momentum advection schemes in a global ocean circulation model at eddy-permitting
722 resolution, *Ocean Dynam.*, 56, 543–567, 2006

723 Bourgeois, T., Orr, J. C., Resplandy, L., Terhaar, J., Ethé, C., Gehlen, M., and Bopp, L.: Coastal-
724 ocean uptake of anthropogenic carbon, *Biogeosciences* 13, 4167-4185, doi:10.5194/bg-13-4167-
725 2016, 2016.

726 Brodeau, L., Barnier, B., Treguier, A. M., Penduff, T. and Gulev, S.: An ERA40-based atmospheric
727 forcing for global ocean circulation models, *Ocean Modelling*, 31(3), 88-104, 2010.

728 Bryden, H. L., Longworth, H. R. and Cunningham, S. A.: Slowing of the Atlantic meridional
729 overturning circulation at 25 N, *Nature*, 438(7068), 655-657, 2005.

730 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., Defries, R.,
731 Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S and Thornton, P.:
732 Carbon and other biogeochemical cycles. In: *Climate Change 2013: The Physical Science Basis.*
733 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel*
734 *on Climate Change* [Stocker, T.F., D. Qin, G-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.
735 Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge,
736 United Kingdom and New York, NY, USA, 2013.

737 Collins, M., Knutti, R., Arblaster, J., Dufresne, J-L., Fichet, T., Friedlingstein, P., Gao, X.,
738 Gutwoski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J. and Wehner, M.:
739 Long-term Climate Change: Projections, Commitments and Irreversibility. . In: *Climate Change*
740 *2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment*
741 *Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G-K. Plattner,
742 M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)].
743 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

744 Conkright, M. E., Locarnini, R. A., Garcia, H. E., O'Brien, T. D., Boyer, T. P., Stephens, C. and
745 Antonov, J. I.: *World Ocean Database 2001: Objective analyses, data statistics and figures*, 2002

746 Crueger, T., Roeckner, E., Raddatz, T., Schnur, R. and Wetzell, P.: Ocean dynamics determine the
747 response of oceanic CO₂ uptake to climate change, *Clim dynam*, 31(2-3), 151-168, 2008.

748 Curry, R. G. and McCartney, M. S.: Ocean gyre circulation changes associated with the North
749 Atlantic Oscillation, *J Phys Oceanogr*, 31(12), 3374-3400, 2001.

750 Daniault, N., Mercier, H., Lherminier, P., Sarafanov, A., Falina, A., Zunino Rodriguez, P., Pérez,
751 F.F., Rios, A.F., Ferron, B., Huck, T., Thierry, V. and Gladyshev, S.: The northern North Atlantic
752 Ocean mean circulation in the early 21st century. *Prog Oceanogr*, 146, 142-158, doi:
753 10.1016/j.pocean.2016.06.007, 2016.

754 de Boissésou, E., Thierry, V., Mercier, H., Caniaux, G and Débruyères, D. : Origin, formation and

755 variability of the Subpolar Mode Water located over the Reykjanes Ridge. *JGR (C12005)*, 117,
756 doi: 10.1029/2011JC007519, 2012

757 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
758 Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
759 Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy,
760 S. B., Hersbach, H., Hólm, E. V, Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally,
761 A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-k., Peubey, C., de Rosnay, P., Tavolato, C.,
762 Thépaut, J.-N and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the
763 data assimilation system, *Q. J. Roy. Meteor. Soc.* 137 (656), 553-597, 2011

764 Delworth, T. L. and Zeng, F.: The impact of the North Atlantic Oscillation on climate through its
765 influence on the Atlantic Meridional Overturning Circulation, *J Climate*, 2015.

766 Dickson, R., Lazier, J., Meincke, J. and Rhines, P.: Long-term coordinated changes in the
767 convective activity of the North Atlantic. In *Decadal Climate Variability* (pp. 211-261). Springer
768 Berlin Heidelberg, 1996.

769 García-Ibáñez, M. I, Pardo, P. C., Carracedo, L., Mercier, H., Lherminier, P., Ríos, A. F. and Pérez,
770 F. F.: Structure, transports and transformations of the water masses in the Atlantic Subpolar Gyre,
771 *Prog. Oceanogr.* 135, 18-36, doi: 10.1016/j.pocean.2015.03.009, 2015.

772 Gent, P. R., and Mewilliams, J. C.: Isopycnal mixing in ocean circulation models, *J Phys Oceanogr*,
773 20(1), 150-155, 1990.

774 Guallart, E. F., Schuster, U., Fajar, N. M., Legge, O., Brown, P., Pelejero, C., Messias, M-J., Calvo,
775 E., Watson, A., Ríos, A. F. and Pérez, F. F.: Trends in anthropogenic CO₂ in water masses of the
776 Subtropical North Atlantic Ocean, *Progr. Oceanogr.*, 131, 21-32, doi:
777 10.1016/j.pocean.2014.11.006, 2015.

778 Good, S. A., Martin, M. J. and Rayner, N. A.: EN4: quality controlled ocean temperature and
779 salinity profiles and monthly objective analyses with uncertainty estimates, *J. Geophys. Res-*
780 *Oceans*, 118, 6704-6716, 2013

781 Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M.
782 J., Gerber, M., Jacobson, A.R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Muller, S.A,
783 Sarmiento, J.L. and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO₂,
784 *Global Biogeochem Cy*, 23(1), 2009.

785 Häkkinen, S., and Rhines, P. B.: Decline of subpolar North Atlantic circulation during the 1990s,
786 *Science*, 304(5670), 555-559, 2004.

787 Hernández-Guerra, A., Pelegrí, J. L., Fraile-Nuez, E., Benítez-Barrios, V., Emelianov, M., Pérez-
788 Hernández, M. D., Vélez-Belchí, P.: Meridional overturning transports at 7.5N and 24.5N in the

789 Atlantic Ocean during 1992–93 and 2010–11, *Progr. Oceanogr.*, 128, 98–114, doi:
790 10.1016/j.pocean.2014.08.016, 2014.

791 Hurrell, J. and National Center for Atmospheric Research staff (Eds)
792 IOC, SCOR and IAPSO: The international thermodynamic equation of seawater - 2010: Calculation
793 and use of thermodynamic properties. Intergovernmental Oceanographic Commission, Manuals
794 and Guides No. 56, UNESCO (English), 196 pp. Available from <http://www.TEOS-10.org>. See
795 section 3.3 of this TEOS-10 Manual, 2010.

796 Jeansson, E., Olsen, A., Eldevik, T., Skjelvan, I., Omar, A. M., Lauvset, S. K., Nilsen, J. E. Ø.,
797 Bellerby, R. G. J., Johannessen, T. and Falck, E.: The Nordic Seas carbon budget: Sources, sinks
798 and uncertainties, *Global Biogeochem. Cy.*, 25, GB4010, doi: 10.1029/2010GB003961, 2011.

799 Johns, W.E., Baringer, M.O., Beal, L.M., Cunningham, S.A., Kanzow, T., Bryden, H.L., Hirschi,
800 J.J.M., Marotzke, J., Meinen, C.S., Shaw, B., Curry, R.: Continuous, array-based estimates of
801 Atlantic Ocean heat transport at 26.5N, *J. Climate*, 24, 2429–2449, doi:
802 10.1175/2010JCLI3997.1, 2011.

803 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero,
804 F. J., Mordy, C. and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data
805 Analysis Project (GLODAP), *Global Biogeochem Cy* 18, GB4031, doi:10.1029/2004GB002247,
806 2004

807 Khatiwala, S., Primeau, F. and Hall, T.: Reconstruction of the history of anthropogenic CO₂
808 concentrations in the ocean, *Nature*, 462, 346-349, doi:10.1038/nature08526, 2009.

809 Khatiwala, S., Tanhua, T., Fletcher, S. M., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N.,
810 McKinley, G.A, Murata, A., Rios, A.F., and Sabine, C. L.: Global ocean storage of
811 anthropogenic carbon, *Biogeosciences*, 10(4), 2169-2191, 2013.

812 Körtzinger, A., Rhein, M., and Mintrop, L.: Anthropogenic CO₂ and CFCs in the North Atlantic
813 Ocean-A comparison of man-made tracers, *Geophys Res Lett*, 26(14), 2065-2068, 1999.

814 Kuhlbrodt, T., Griesel, A. Montoya, M., Levermann, A., Hofmann, M. and Rahmstorf, S.: On the
815 driving processes of the Atlantic meridional overturning circulation, *Rev. Geophys*, 45, RG2001,
816 doi: 10.1029/2004RG000166, 2007.

817 Landschützer, P., Gruber, N. and Bakker, D. C. E. and Schuster, U.: Recent variability of the global
818 ocean carbon sink, *Global Biogeochem. Cy.*, 28, 947-949, doi:10.1002/2014GB004853, 2014.

819 Landschützer, P., Gruber, N. and Bakker, D.C.E.: A 30 years observation-based global monthly
820 gridded sea surface pCO₂ product from 1982 through 2011. [http://cdiac.ornl.gov/ftp/oceans/](http://cdiac.ornl.gov/ftp/oceans/SPCO2_1982_2011_ETH_SOM_FF.N)
821 SPCO2_1982_2011_ETH_SOM_FF.N. Carbon Dioxide Information Analysis Center, Oak Ridge
822 National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi:

823 10.3334/CDIAC/OTG.SPCO2_1982_2011_ETH_SOM-FFN, 2015a.

824 Landschützer, P., Gruber, N., Haumann, F. A., Rödenbeck, C., Bakker, D.C.E., van Heuven, S.,
825 Hoppema, M., Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B. and Wanninkhof, R.: The
826 reinvigoration of the Southern Ocean carbon sink, *Science*, 349, 1221-1224. doi:
827 10.1126/science.aab2620, 2015b.

828 Lazier, J., Hendry, R., Clarke, A., Yashayaev, I. and Rhines, P.: Convection and restratification in
829 the Labrador Sea, 1990–2000, *Deep Sea Res PtI*, 49(10), 1819-1835, 2002.

830 Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G. and co-authors: Trends in the sources
831 and sinks of carbon dioxide, *Nature Geosciences*, 2(12), 831-836, 2009.

832 Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., and co-authors : Global carbon budget
833 2013, *ESSD*, 6, 235–263, doi:10.5194/essd-6-235-2014, 2014.

834 Le Quéré, C. Moriarty, R., Andrew, R.M., Peters, G.P., and co-authors: Global Carbon Budget
835 2014, 2015.

836 Lherminier, P., Mercier, H., Gourcuff, C., Alvarez, M., Bacon, S. and Kermabon, C.: Transports
837 across the 2002 Greenland-Portugal Ovide section and comparison with 1997, *J Geophys Res-*
838 *Oceans*, 112(C7), 2007.

839 Lherminier, P., Mercier, H., Huck, T., Gourcuff, C., Perez, F. F., Morin, P., Sarafanov, A.,
840 and Falina, A.: The Atlantic Meridional Overturning Circulation and the subpolar gyre observed
841 at the A25-OVIDE section in June 2002 and 2004, *Deep Sea Res Pt I* 57(11), 1374-1391, 2010.

842 Madec, G., and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, *Climate*
843 *Dy* 12(6), 381-388, 1996.

844 Madec, G.: NEMO Ocean Engine, vol. 27, pp. 1–217, Note du Pole de modélisation de l’Institut
845 Pierre-Simon Laplace, France, 2008

846 Maier-Reimer, E., Mikolajewicz, U. and Winguth, A. : Future ocean uptake of CO₂: interaction
847 between ocean circulation and biology, *Clim. Dynam.*, 12(10), 711-722, doi:
848 10.1007/s003820050138, 1996.

849 McCarthy, G., Frajka-Williams, E., Johns, W. E., Baringer, M. O., Meinen, C. S., Bryden, H. L.,
850 Rayner, D., Ducez, A., Roberts, C. and Cunningham, S. A.: Observed interannual variability of
851 the Atlantic meridional overturning circulation at 26.5°N, *Geophys. Res. Lett.*, 39, L19609,
852 doi:10.1029/2012GL052933, 2012

853 McCartney, M. S. and Talley, L. D.: The subpolar mode water of the North Atlantic Ocean, *J. Phys.*
854 *Oceanogr.*, 12(11), 1169-1188, doi : 10.1175/1520-0485(1982)012, 1982

855 McCartney, M. S.: Recirculation components to the deep boundary current of the northern North
856 Atlantic, *Prog. Oceanogr.*, 29(4), 283-383, doi: 10.1016/0079-6611(92)90006-L, 1992.

857 McKinley, G.A., Pilcher, D.J., Fay, A.R., Lindsay, K., Long, M.C. and Lovenduski, N.S.:

858 Timescales for detection of trends in the ocean carbon sink. *Nature*, 530, 469-472, 2016.

859 Mercier, H., Lherminier, P., Sarafanov, A., Gaillard, F., Danaïult, N., Desbruyères, D., Falina, A.,

860 Ferron, B., Gourcuff, C., Huck, T. and Thierry, V.: Variability of the meridional overturning

861 circulation at the Greenland–Portugal OVIDE section from 1993 to 2010, *Prog Oceanogr* 132

862 (2015) 250–261, [doi:10.1016/j.pocean.2013.11.001](https://doi.org/10.1016/j.pocean.2013.11.001), 2015

863 Mikaloff Fletcher, S. E., Gruber, N., and Jacobson, A. R.: Ocean Inversion Project How-to

864 Document Version 1.0, 18 pp. Institute for Geophysics and Planetary Physics, University of

865 California, Los Angeles, 2003

866 Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M.,

867 Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S.A. and Sarmiento,

868 J.L.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean. *Global*

869 *Biogeochem Cy* 20(2), [doi:10.1029/2005GB002530](https://doi.org/10.1029/2005GB002530), 2006.

870 Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A. ,

871 Tanhua, T. , Hoppema, M. , Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F.,

872 and Suzuki T.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) – an internally

873 consistent data product for the world ocean, *Earth Syst. Sci. Data*, 8, 297–323, [doi:10.5194/essd-](https://doi.org/10.5194/essd-8-297-2016)

874 [8-297-2016](https://doi.org/10.5194/essd-8-297-2016), 2016

875 Orr, J. C., Najjar, R. G., Aumont, O., Bopp, L., Bullister, J. L., Danabasoglu, G., Doney, S. C.,

876 Dunne, J. P., Dutay, J-C., Graven, H., Griffies, S. M., John, J. G., Joos, F., Levin, I., Lindsay, K.,

877 Matear, R., McKinley, G. A., Mouchet, A., Oschlies, A., Romanou, A., Schlitzer, R., Tagliabue,

878 A., Tanhua, T. and Yool, A.: Biogeochemical protocols and diagnostics for the CMIP6 Ocean

879 Model Intercomparison Project (OMIP), *Geosci. Model Dev.*, 10, 2169-2199, [doi: 10.5194/gmd-](https://doi.org/10.5194/gmd-10-2169-2017)

880 [10-2169-2017](https://doi.org/10.5194/gmd-10-2169-2017), 2017.

881 Pérez, F. F., Vazquez-Rodriguez, M., Louarn, E., Padín, X. A., Mercier, H., and Ríos, A. F. :

882 Temporal variability of the anthropogenic CO₂ storage in the Irminger Sea, *Biogeosciences*,

883 5(6), 1669-1679, 2008

884 Pérez, F. F., Vázquez Rodríguez, M., Mercier, H., Velo, A., Lherminier, P. and Ríos, A. F. : Trends

885 of anthropogenic CO₂ storage in North Atlantic water masses, *Biogeosciences*, 7, 1789–1807,

886 [doi:10.5194/bg-7-1789-2010](https://doi.org/10.5194/bg-7-1789-2010), 2010.

887 Pérez, F. F., Mercier, H., Vázquez-Rodríguez, M., Lherminier, P., Velo, A., Pardo, P. C., Roson,

888 G., and Ríos, A. F. : Atlantic Ocean CO₂ uptake reduced by weakening of the meridional

889 overturning circulation, *Nature Geoscience*, 6(2), 146-152, [doi: 10.1038/NGEO1680](https://doi.org/10.1038/NGEO1680), 2013

890 Pickart, R. S.: Water mass components of the North Atlantic deep western boundary current, *Deep-*

891 Sea Res. Pt A, 39(9), 1553-1572, doi: 10.1016/0198-0149(92)90047-W, 1992.

892 Pickart, R. S., Straneo, F. and Moore, G. W. K.: Is Labrador sea water formed in the Irminger
893 basin?, *Deep Sea Res Pt I* 50(1), 23-52, 2003.

894 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ
895 and satellite SST analysis for climate, *J. Climate*, 15, 1609–1625, 2002.

896 Rhein, M., Kieke, D., Hüttl-Kabus, S., Roessler, A., Mertens, C., Meissner, R., Klein, B., Böning,
897 C.W. and Yashayaev, I.: Deep water formation, the subpolar gyre, and the meridional
898 overturning circulation in the subpolar North Atlantic, *Deep-Sea Res. Pt II* 58(17), 1819-1832,
899 2011.

900 Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P.,
901 Metzl, N., Nakaoka, S., Olsen, A., Park, G-H, Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster,
902 U., Shutler, J. D., Valsala, V., Wanninkhof, R. and Zeng, J.: Data-based estimates of the ocean
903 carbon sink variability – first results of the Surface Ocean pCO₂ mapping intercomparison
904 (SOCOM), *Biogeosciences*, 12, 7251-7278, doi: 10.5194/bg-12-7251-2015, 2015.

905 Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R.,
906 Wong, C.S., Wallace, D.W.R., Tilbrook, B., Millero, F.J., Peng, T-H., Kozyr, A., Ono, T. and
907 Rios, A.F.: The oceanic sink for anthropogenic CO₂. *Science* 305(5682), 367-371, 2004.

908 Sarafanov, A.: On the effect of the North Atlantic Oscillation on temperature and salinity of the
909 subpolar North Atlantic intermediate and deep waters. *ICES Journal of Marine Science: Journal*
910 *du Conseil*, 66(7), 1448-1454, 2009.

911 Sarafanov, A., Falina, A., Mercier, H., Sokov, A., Lherminier, P., Gourcuff, C., Gladyshev, S.,
912 Gaillard, F. and Daniault, N. : Mean full-depth summer circulation and transports at the northern
913 periphery of the Atlantic Ocean in the 2000s, *J. Geophys. Res.*, 117(C01014), doi :
914 10.1029/2011JC007572, 2012.

915 Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., González-Dávila,
916 M., Gruber, N., Jones, S., Krijnen, J., Landschützer, P., Lefèvre, N., Manizza, M., Mathis, J.,
917 Metzl, N., Olsen, A., Rios, A. F., Rödenbeck, C., Santana-Casiano, J. M., Takahashi, T.,
918 Wanninkhof, R., and Watson, A. J.: An assessment of the Atlantic and Arctic sea–air CO₂ fluxes,
919 1990–2009, *Biogeosciences*, 10, 607–627, doi:10.5194/bg-10-607-2013, 2013

920 Schwinger, J., Tjiputra, J. F., Heinze, C., Bopp, L., Christian, J. R., Gehlen, M., Ilyina, T., Jones, C.
921 D., Salas-Méla, D., Segschneider, J., Séférian, R. and Totterdell, I.: Nonlinearity of ocean
922 carbon cycle feedbacks in CMIP5 Earth System Models, *J. Climate*, 27(11), 3869-3888, doi:
923 10.1175/JCLI-D-13-00452.1, 2014

924 Séférian, R., Ribes, A. and Bopp, L.: Detecting the anthropogenic influences on recent changes in

925 ocean carbon uptake: *Geophys. Res. Lett.*, 41, 5968-5977, doi: 10.1029/1999JC900274, 2014.

926 Smethie, W. M., Fine, R. A., Putzka, A. and Jones, E. P.: Tracing the flow of North Atlantic Deep
927 Water using chlorofluorocarbons, *J. Geophys. Res.*, 105(C6), 14297-14323, 2000.

928 Steinfeldt, R., Rhein, M., Bullister, J. L. and Tanhua, T.: Inventory changes in anthropogenic
929 carbon from 1997–2003 in the Atlantic Ocean between 20 S and 65 N, *Global Biogeochem Cy*
930 23(3), 2009.

931 Takahashia, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N.,
932 Wanninkhof, R., Feely, R. A., Sabine, C., Olafsson, J. and Nojiri, Y.: Global sea–air CO₂ flux
933 based on climatological surface ocean pCO₂, and seasonal biological and temperature effects,
934 *Deep-Sea Res. pt II*, 49,1601–1622, doi: 10.1016/S0967-0645(02)00003-6, 2002.

935 Talley, L. D., Pickard, G. L., Emery, W. J. and Swift, J. H.: *Descriptive physical oceanography: an*
936 *introduction*, Academic press, pp 555, 2008

937 Timmermann, R., Goosse, H., Madec, G., Fichefet, T., Ethe, C. and Duliere, V.: On the
938 representation of high latitude processes in the ORCA-LIM global coupled sea ice–ocean model.
939 *Ocean Modelling*, 8(1), 175-201, 2005.

940 Thomas, H., Prowe, F.A. E., Lima, I. D., Doney, S. C., Wanninkhof, R., Greatbach, R. J., Schuster,
941 U. and Corbière, A.: Changes in the North Atlantic Oscillation influence CO₂ uptake in the
942 North Atlantic over the past 2 decades, *Global Biogeochem. Cy.*, 22(4), doi:
943 10.1029/2007GB003167, 2008.

944 Treguier, A-M., Gourcuff, C., Lherminier, P., Mercier, H., Barnier, B., Madec, G., Molines, J-M.,
945 Penduff, T., Czeschel, L., Böning, C.W.: Internal and Forced variability along a section between
946 Greenland and Portugal in the CLIPPER Atlantic model. *Ocean Dynam* 56 (5-6), 568-580,
947 doi:10.1007/s10236-006-0069-y, 2006

948 Vázquez-Rodríguez, M., Padin, X. A., Ríos, A. F., Bellerby, R. G. J. and Pérez, F. F.: An upgraded
949 carbon-based method to estimate the anthropogenic fraction of dissolved CO₂ in the Atlantic
950 Ocean, *Biogeosciences Discuss.*, 6, 4527–4571, doi:10.5194/bgd-6-4527-2009, 2009.

951 Velo, A., Pérez, F. F., Lin, X., Key, R. M., Tanhua, T., de la Paz, M., Olsen, A., van Heuven, S.,
952 Jutterström, S. and Ríos, A. F.: CARINA data synthesis project: pH data scale unification and
953 cruise adjustments, *Earth Syst. Sci. Data*, 2, 133–155, doi:10.5194/essd-2-133-2010, 2010.

954 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *Jeophys Res*
955 97(C5), 7373-7382, 1992.

956 Wanninkhof, R., Park, G.H., Takahashi, T., Sweeney, C., Feely, R.A., Nojiri, Y., Gruber, N.,
957 Doney, S.C., McKinley, G.A., Lenton, A., Le Quere, C., Heinze, C. Schwinger, J., Graven, H.,
958 and Khatiwala, S.: Global Ocean carbon uptake: Magnitude, variability and trend,

959 Biogeosciences 10, 1983-2000, doi:10.5194/bg-10-1983-2013, 2013.

960 Weiss, R. F.: Carbon Dioxide in Water and Seawater: The Solubility of a Non-Ideal Gas, *Mar.*
961 *Chem.*, 2, 203–215, 1974.

962 Yashayaev, I.: Hydrographic changes in the Labrador Sea, 1960–2005, *Prog Oceanogr* 73(3), 242-
963 276, 2007.

964 Zunino, P., Garcia-Ibanez, M. I., Lherminier, P., Mercier, H., Ríos, A. F. and Pérez, F. F.:
965 Variability of the transport of anthropogenic CO₂ at the Greenland-Portugal OVIDE section:
966 Controlling mechanisms, *Biogeosciences*, 11, 2375–2389, doi:10.5194/bg-11-2375-2014, 2014

967 Zunino, P., Lherminier, P., Mercier, H., Padín, X. A., Ríos, A. F. and Pérez, F. F.: Dissolved
968 inorganic carbon budgets in the eastern subpolar North Atlantic in the 2000s from in situ data,
969 *Geophys Res Lett* 42(22), 9853-9861, 2015a.

970 Zunino, P., Pérez, F. F., Fajar, N. M., Guallart, E. F., Ríos, A. F., Pelegrí, J. L. and
971 Hernández-Guerra, A.: Transports and budgets of anthropogenic CO₂ in the tropical North
972 Atlantic in 1992–1993 and 2010–2011, *Global Biogeochem Cy*, 29(7), 1075-1091, 2015b.

973

974 **Acknowledgments**

975 For this work, VR was funded through the EU FP7 project CARBOCHANGE (grant 264879).
976 Simulations were made using HPC resources from GENCI-IDRIS (grant x2015010040). We are
977 grateful to Christian Ethé, who largely contributed to obtain Cant transport in online mode over the
978 period 2003-2011. We want to acknowledge HM (supported by CNRS and the ATLANTOS H2020
979 project (GA 633211)) and colleagues for leading the OVIDE project (supported by French research
980 institutions, IFREMER and CNRS/INSU), as well as Alonso Hernandez-guerra for the availability
981 of its mass transport data at 24.5°N. Other data at 24.5°N used in this paper were collected and
982 made publicly available by the International Global Ship-based Hydrographic Investigations
983 Program (GO-SHIP; <http://www.go-ship.org/>) and the national programs that contribute to it. We
984 are also grateful to N. Gruber and one anonymous reviewer for their constructive comments.

985

986 **Table captures**

987 Table 1: Summary of cruises and data set used throughout this study

OVIDE name	Month/year	Vessel	Reference	expocode
OVIDE 2002	06-07/2002	N/O Thalassa	Lherminier et al., 2007	35TH20020611
OVIDE 2004	06-07/2004	N/O Thalassa	Lherminier et al., 2010	35TH20040604
OVIDE 2006	05-06/2006	R/V Maria S. Merian	Gourcuff et al., 2011	06MM20060523
OVIDE 2008	06-07/2008	N/O Thalassa	Mercier et al. 2015	35TH20080610

OVIDE 2010	06-07/2010	N/O Thalassa	Mercier et al., 2015	35TH20100608
24.5°N-2011	01-03/2011	Sarmiento de Gamboa	Hernández-Guerra et al. 2014	29AH20110128

988

989 Table 2: Model-data comparison over the period covered by the OVIDE cruises (2002-2010).

990 Average and standard deviation (SD) for observation-based estimates (column 2) and model output
991 (columns 3 to 4). Model output: (1) June average with SD being a measure of interannual variability
992 and (2) average year with SD corresponding to the average interannual variability.

	OVIDE data set	ORCA05-PISCES	
		June only	average year
MOC σ (sv)	15.5±2.3	13.4±0.6	12.7±0.6
σ MOC (kg m ⁻³)	32.14	32.02±0.05	31.95±0.04
[Cant] _{section} (μmol kg ⁻¹)	25.4±1.8	18.4±1.1	18.4±1.1
[Cant] _{upper} (μmol kg ⁻¹)	45.2±3.0	38.9±3.0	39.4±3.0
[Cant] _{lower} (μmol kg ⁻¹)	19.4±1.6	14.8±1.0	14.9±1.0

993

994 Table 3: Model-data comparison along 25°N. Average and standard deviation (SD) for observation-
995 based estimates (column 2) and model output (columns 3 to 5). Model output: (1) January from
996 March 2011 average with SD being a measure of winter variability, (2) average 2011 year with SD
997 corresponding to the average seasonal variability and (3) average 2003-2011 year with SD being
998 representative of interannual variability.

	24.5°N data set	ORCA05-PISCES		
		Winter only	Average 2011 year	Average 2003-2011 year
MOC σ (sv)	20.1±1.4	10.82±2.14	11.59±1.86	11.13±0.80
σ MOC (kg m ⁻³)	32.27	31.95±0.00	32.02±0.03	32.00±0.03
[Cant] _{section} (μmol kg ⁻¹)	19.73	8.69±0.02	8.73±0.04	
[Cant] _{upper} (μmol kg ⁻¹)	40.36	39.15±0.01	38.86±0.90	
[Cant] _{lower} (μmol kg ⁻¹)	12.00	2.89±0.1	2.86±0.08	

999

1000 Table 4: Correlation coefficient (r) and p-value between the time rate of change (Trate), the
1001 divergence of Cant transport (DT_{Cant}) and air sea Cant fluxes (F_{Cant}) for the two boxes, 25°N-
1002 OVIDE and OVIDE-Sills, over the period 2003-2011. DT_{Cant} = incoming – outgoing Cant fluxes
1003 across the boundaries of boxes.

Box 25° N to OVIDE

Trate/DT_{Cant} : r = 0.96, p-value = 0.00

Trate/F_{Cant} : r = - 0.54, p-value = 0.00

Box OVIDE to sills

Trate/DT_{Cant} : r = 0.95, p-value = 0.00

Trate/F_{Cant} : r = - 0.71, p-value = 0.00

1004

1005

1006

1007

1008

1009

1010

Table 5: Summary of (a-b) the coefficient of correlation (with p-value) between the MOC and the transport of heat and Cant at 25°N, 36°N and the OVIDE section. The analyses were done first with the original time series (a. including trend)) and after, with the detrended Cant transport time series (b. without trend). The trend for each term as well as those of volume transport are reported in the third part of this table (c. trend).

	25°N	36°N	OVIDE
a. coefficient of correlation (p-value) including trend			
T _{heat} vs MOC	0.92 (0.00)	0.90 (0.00)	0.76 (0.00)
T _{Cant} vs MOC	0.30 (0.02)	0.67 (0.00)	0.02 (0.90)
b. coefficient of correlation (p-value) without trend			
T _{Cant} vs MOC	0.74 (0.00)	0.70 (0.00)	0.01 (0.40)
c. trend			
T _{Cant} (1958-60)	0.030±0.002 PgC yr ⁻¹	0.009±0.001 PgC yr ⁻¹	0.008±0.001 PgC yr ⁻¹
T _{Cant} (2010-12)	0.095±0.024 PgC yr ⁻¹	0.050±0.018 PgC yr ⁻¹	0.043±0.005 PgC yr ⁻¹
T _{heat}	0.0003±0.0004 PW yr ⁻¹	0.0016±0.0004 PW yr ⁻¹	0.0003±0.0002 PW yr ⁻¹
MOC	0.001±0.005 Sv yr ⁻¹	0.016±0.006 Sv yr ⁻¹	0.003±0.007 Sv yr ⁻¹
T _{vol}	-0.000±0.000 Sv yr ⁻¹	0.001±0.001 Sv yr ⁻¹	-0.000±0.003 Sv yr ⁻¹

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

Table 56: Correlation coefficient (r) and p-value between the time rate of change (Trate) of Cant storage, the divergence of Cant transport (DT_{Cant}) and air sea Cant fluxes (F_{Cant}) for the three boxes, 25°N-36°N, 36°N-OVIDE and OVIDE-Sills, over the period 1959-2011. DT_{Cant} = incoming – outgoing Cant fluxes across the boundaries of boxes. The analyses were done, first, with the original time series (left column in the table) and after, with the detrended Cant transport time series (right column in the table).

a. Including trend

Box 25° N to 36°N

Trate/DT_{Cant} : r = 0.93, p-value = 0.00

Trate/F_{Cant} : r = 0.90, p-value = 0.00

b. without trend

Box 25° N to 36°N

Trate/DT_{Cant} : r = 0.94, p-value = 0.00

Trate/F_{Cant} : r = 0.04, p-value = 0.78

Box 36°N to OVIDETrate/DT_{Cant} : r = 0.73, p-value = 0.00Trate/F_{Cant} : r = 0.97, p-value = 0.00

Box OVIDE to sillsTrate/DT_{Cant} : r = 0.32, p-value = 0.02Trate/F_{Cant} : r = 0.95, p-value = 0.00

Box 36°N to OVIDETrate/DT_{Cant} : r = 0.61, p-value = 0.00Trate/F_{Cant} : r = 0.52, p-value = 0.00

Box OVIDE to sillsTrate/DT_{Cant} : r = 0.76, p-value = 0.00Trate/F_{Cant} : r = 0.22, p-value = 0.12

1021

1022 **Table 67:** Correlation coefficient (r) and p-value between the divergence of Cant transport (DT_{Cant})1023 and the incoming (in) or outgoing (out) transport of Cant (T_{Cant}) for the three boxes, 25°N-36°N,1024 36°N-OVIDE and OVIDE-Sills, over the period 1959-2011. DT_{Cant} = incoming – outgoing Cant

1025 fluxes across the boundaries of boxes. Linear trend is removed from each times series beforehand.

Box 25° N to 36°NⁱⁿT_{Cant}/DT_{Cant} : r = 0.51, p-value = 0.00^{out}T_{Cant}/DT_{Cant} : r = -0.31, p-value = 0.03

Box 36°N to OVIDEⁱⁿT_{Cant}/DT_{Cant} : r = 0.79, p-value = 0.00^{out}T_{Cant}/DT_{Cant} : r = 0.07, p-value = 0.62

Box OVIDE to sillsⁱⁿT_{Cant}/DT_{Cant} : r = 0.68, p-value = 0.00^{out}T_{Cant}/DT_{Cant} : r = -0.05, p-value = 0.70

1026

1027 **Figures captions**1028 Fig. 1: Column inventory (molC m⁻²) of anthropogenic carbon for the year 2010: (a) model output
1029 and (b) Khatiwala et al. [2009].

1030

1031 Fig. 2: The Greenland-Portugal OVIDE and 24.5°N sections: observational data set (red points) and
1032 ORCA05-PISCES (black thick line).

1033

1034 Fig. 3: Anthropogenic C budget of the Subtropical and Subpolar North Atlantic regions over the
1035 period 2003-2011. Average values and their standard deviations were estimated from smoothed time
1036 series. The horizontal arrows show the lateral Cant transport in PgC yr⁻¹ (black font). Red numbers
1037 in the panel indicate the Cant storage rate in PgC yr⁻¹. The vertical arrows show the total (blue font)
1038 and anthropogenic (black font) air-sea CO₂ fluxes in PgC yr⁻¹. Green numbers represent the heat
1039 transport across sections in PW. Boundaries and surface area (m²) of each box are indicated below
1040 the panels.

1041

1042 Fig. 4. Cumulative volume transport in Sv. (a) Vertically integrated transport from bottom to each
1043 specific density level (σ₁ with 0.01 kg m⁻³ resolution). Note that the sign of the profile has been

1044 changed. (b) Surface-to-bottom integrated transport cumulated from Greenland to Portugal (km).
1045 Model outputs for the month of June over the period 2002-10 (continuous line for mean value;
1046 shadows for confidence interval) are compared to estimates derived from OVIDE (dashed lines). On
1047 panel (a) the black horizontal lines indicate the density of MOC maximum corresponding to the
1048 separation between the upper (red) and lower (blue) limbs of MOC, in the model ($\sigma_{\text{MOC}} =$
1049 $32.02 \pm 0.05 \text{ kg m}^{-3}$, black continuous line) and observation-based assessments ($\sigma_{\text{MOC}} = 32.14 \text{ kg m}^{-3}$,
1050 Zunino et al., 2014; black dashed line). The position of the Western limit of the NAC as observed
1051 from model simulations (dashed line) and from OVIDE data set (dashed-dotted line) as well as the
1052 Irminger current are indicated on panel (b).

1053

1054 Fig. 65: Zonally integrated volume transport (S_v) at 25°N computed either (a) for main water
1055 masses between January and March 2011 or (b) for density level (σ_1 with 0.1 kg m^{-3} resolution)
1056 over the year 2011 from model output. Main water masses identified at this latitude are North
1057 Atlantic Central Water (NACW) Antarctic Intermediate Water (AAIW) and Mediterranean Water
1058 (MW), which constitute the upper limb of the MOC (red), as well as North Atlantic Deep Water
1059 and Antarctic Bottom Water (AABW), which compose the lower limb of the MOC (blue). Results
1060 from panel (a) are compared to observation-based estimates from Hernández-Guerra et al. (2014)
1061 (hatched bar plot). On panel (b) the black horizontal lines indicate the density of MOC maximum
1062 corresponding to the separation of both limb in the model ($\sigma_1 = 32.05$ from July to September and
1063 $\sigma_1 = 21.95$ other months).

1064

1065 Fig. 56 : Water column distribution of anthropogenic C concentrations ($\mu\text{mol kg}^{-1}$) along the
1066 Greenland-Portugal OVIDE section in June 2002: (a) model output and (b) as estimated from the
1067 OVIDE data set. The mean and standard deviation of the differences between these two assessments
1068 (model – OVIDE) over the OVIDE period (June 2002-04-06-08-10) are displayed on Fig. c and d.
1069 Black continuous and dashed lines indicate the limit between the upper and the lower limbs of the
1070 MOC in the model and the OVIDE data set.

1071

1072 ~~Fig. 6: Zonally integrated volume transport (S_v) at 25°N computed either (a) for main water masses~~
1073 ~~between January and March 2011 or (b) for density level (σ_1 with 0.1 kg m^{-3} resolution) over the~~
1074 ~~year 2011 from model output. Main water masses identified at this latitude are North Atlantic~~
1075 ~~Central Water (NACW) Antarctic Intermediate Water (AAIW) and Mediterranean Water (MW);~~
1076 ~~which constitute the upper limb of the MOC (red), as well as North Atlantic Deep Water) and~~
1077 ~~Antarctic Bottom Water (AABW), which compose the lower limb of the MOC (blue). Results from~~

1078 ~~panel (a) are compared to observation based estimates from Hernández-Guerra et al. (2014)~~
1079 ~~(hatched bar plot). On panel (b) the black horizontal lines indicate the density of MOC maximum~~
1080 ~~corresponding to the separation of both limb in the model ($\sigma_1 = 32.05$ from July to September and~~
1081 ~~$\sigma_1 = 21.95$ other months).~~

1082

1083 Fig. 7: Water column distribution of anthropogenic C concentrations ($\mu\text{mol kg}^{-1}$) along 24.5°N-
1084 25°N during winter (JFM) 2011: (a) model output and (b) as estimated from the 24.5N-data set. (c)
1085 Difference between both assessments (model – observation) in 2011. Black continuous and dashed
1086 lines indicate the limit between the upper and the lower limbs of the MOC in the model and the
1087 observation data set.

1088

1089 Fig. 8: (a-b) averaged total air-sea CO_2 fluxes ($\text{mol m}^2 \text{yr}^{-1}$) and month during which (c-d) the
1090 maximum or (e-f) the minimum value is reached in the North Atlantic Ocean over the period 2003-
1091 2011 as simulated by ORCA05-PISCES (a-c-e) and compared to the observation-based product of
1092 Landschützer et al. (2015a) (b-d-f). Black lines delimitate both boxes, 25°N-OVIDE and OVIDE
1093 sills.

1094

1095 Fig. 9: Interannual variability of total air-sea CO_2 fluxes ($\text{mol m}^2 \text{yr}^{-1}$) for the period 1982-2011
1096 computed as the time series of its standard deviation: (a) ORCA05-PISCES and (b) the observation-
1097 based product of Landschützer et al. (2015a). Black lines delimitate both boxes, 25°N-OVIDE and
1098 OVIDE sills.

1099

1100 Fig. 10: Simulated annual time series of MOC magnitude ($\text{MOC}\sigma$, Sv) and transport of heat (PW)
1101 and anthropogenic C (PgC yr^{-1}) at at 25°N, 36° N and at the OVIDE section estimated over the
1102 period 1958-2012.

1103

1104 Fig. 11 : Simulated annual time series of anthropogenic carbon (Cant) budget (Pg yr^{-1}) from 25°N to
1105 36°N bottom), from 36°N to OVIDE section (middle) and from OVIDE section to Greenland-
1106 Iceland-Scotland sills (top) over the period 1959-2011. Each budget is composed by the storage rate
1107 of Cant (red line), the air-sea flux of Cant (black dashed line) and the transport divergence of Cant
1108 (black full line).

1109

1110 Fig. 12: Distribution of mass transport integrated into density (σ_1) layers with a 0.3 kg m^{-3}
1111 resolution for 25°N, 36°N, OVIDE section and the Greenland-Iceland-Scotland sills over the period

1112 1958-2012 (colorbar). Dashed lines indicate the density limits of three major oceanic water class :
1113 Class 1N = northward North Atlantic Central Water; Class 1S = southward North Atlantic Central
1114 Water ; Class 2 = Intermediate waters; Class 3 : North Atlantic Deep Water.

1115

1116 Fig. 13 : Simulated anthropogenic C budget (PgC yr⁻¹) between 25°N and the Greenland-Iceland-
1117 Scotland sills over the period 1958-1994. Horizontal arrows represent the transport of Cant by
1118 NACW (purple), IW (red) and NADW (blue) across 25°N, 36°N, OVIDE and sills. Grey vertical
1119 arrows show anthropogenic air-sea CO₂ fluxes for each box whereas orange values indicate Cant
1120 storage rate. Black vertical arrows represent the deduced vertical transport of Cant between two
1121 Classes. (b) between 1996 and 2012

1122

1123 Fig. 14: Annual time series of the anomaly of mass transport (Sv, bar plot) compared to the winter
1124 NAO over the period 1959-2011 for (a) Class 1 at 36°N ($r = 0.55$, p -value = 0.00) and (b) Class 3 at
1125 OVIDE ($r = 0.55$, p -value = 0.00). Winter NAO index is index provided by the Climate Analysis
1126 Section (Hurrell and NCAR, <https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-station-based>).
1127

1128

1129 ~~Fig. 15: Simulated annual averaged transport of Cant by NACW (purple), IW (red) and NADW~~
1130 ~~(blue) across 25°N, 36°N, the OVIDE section and the Greenland-Iceland-Scotland sills (a) before~~
1131 ~~and (b) between 1996 and 2012~~

1132

1133 Fig. 165: Simulated annual averaged temperature of mixed layer between 36°N and the OVIDE
1134 section (red line) and between the OVIDE section and the Greenland-Iceland-Scotland sills (black
1135 line) as simulated by the model over the period 1958-2012.