

Referee # 1:

General comments

1. This paper presents a study examining N sources for seasonal growth in *Quercus petraea*. Experimental measurements of N cycling phenology are limited and the manuscript represents a valuable addition to the literature. The author use a clever application of ^{15}N tracers at different times to different pools to provide a solid framework to infer N cycling processes and draw conclusions regarding the origins of N used for seasonal growth in deciduous trees. However, while the theoretical underpinnings and significance of the experiments appear sound, in my opinion there are deficiencies in the methodology and presentation of results which need to be addressed so that the conclusions of the manuscript can be trusted. While isotope labelling experiments are technically challenging and expensive to perform and analyze, the ability to generalize from the experiment severely limited as i) only two tree level replicates are used for each treatment and ii) there are no proper controls for natural variation in isotope abundance over time. This former deficiency prevents any descriptive statistics or statistical analysis in the paper, while the latter means that ^{15}N recovery is calculated using a pre-experiment baseline without any consideration as to whether background ^{15}N content may change over time.

Response: Isotope abundance was determined all along the experiment on each sampled compartments on non labelled trees growing at the same area. The results showed very weak variations of ^{15}N natural abundance (means A% (Isotopic abundance) for leaves = $0.3644 \pm 6.24 \cdot 10^{-5}$ for example). As a consequence, for all calculations, we have chosen to use the value of natural abundance just before labelling (but it could have been the mean of the temporal values).

L146-151 : The seasonal variations of the natural ^{15}N abundance of each compartments were also followed all long the season, those variations were very weak, consequently, it has been choose to use the ^{15}N natural abundance of the labelled trees just before labelling.

2. While neither of these aspects of experimental design can easily be amended an honest discussion of these methodological shortcomings is necessary in the discussion section to understand the limitations of interpretation which arise as a result.

Response: The discussion part has been completed by the following text:

L264-269: "Isotope labelling experiments are technically challenging, and as a consequence are very scarce on trees growing in natural conditions. In this paper, field labelling campaigns were conducted on 20-year-old naturally regenerated oaks. For each campaign (only) two trees were labelled. Nevertheless the similarity of the results between them suggests that the observed ^{15}N partitioning in soil and tree is a representative view of the functioning of such systems. "

3. In large part this discussion repeats some information which I think could be placed in the results (overall label recovery) and omits a critical discussion of the methodology.

Response: Discussion concerning label recovery was reduced in order to limit repetition L270-293.

4. I think that these changes should also be accompanied by improving the quality of graphs and detail in the methods (see specific comments on these aspects of the paper), so the experiment can be both correctly interpreted and repeated.

On this point I find the manuscript is vague and more detail would be very useful.

Response: Materials and methods section has been completed as described below and the quality of the graphs was optimized.

5. The differences between pairs of trees are not discussed besides being referred to as 'similar' at the start of the results (l 143); while the graphs show that, indeed, the time courses of proportional ^{15}N recovery seem similar there are no error bars representing measurement uncertainty nor clear indication of how many points are on the lines.

Response: At each sampling date 20 leaves, 20 twigs were randomly sampled on trees crown, all roots of a core were also sampled. Leaves, twigs and roots were pooled, ground in fine powder and an aliquot was analyzed (^{15}N and %N). As consequence, it was not possible to present error bars at each sampling date. However, at few dates (Day After Labelling (DAL) 1, 126, 337, 460 for leaves and twigs of L_1 ; DAL 126, 337, 460 for roots of L_1 ; DAL 227 and 350 for leaves and twigs of L_2 ; DAL 49 and 350 for roots of L_2 and DAL 40 and 166 for leaves, twigs and roots of L_3) 4 aliquots were analyzed to check the repeatability of analyzes. This type of replicate could not be done at all dates due to the excessive number of sample that would have generated. The results show a good repeatability, for these samples an average is made between the replicates, the errors bars corresponding were added on graphs (L125-129).

6. Given that there are only 15 points per series (Table 1), could these be shown on the graph to indicate periods where ^{15}N content is inferred by a fitted line rather than a measurement?

Response: The graphs have been changed in order to visualize the sampled points all long the experiment. A winter point was added to complete the temporal patterns. Both trees were distinguished with solid lines and dotted lines.

7. Likewise, it is not clear in the methods how samples were taken, how many samples were collected, and when they were taken. 'Leaves, twigs, trunk phloem and xylem and soil monoliths were sampled regularly'. What is regularly?

Response: Table 1 presents the date (Day after labelling DAL and Julian Day JD) of sampling after labelling for each compartment and each labelled tree L112.

At each date of sampling presented on the new graphs, leaves, twigs, roots, microbial biomass, rhizospheric soil were sampled and analyzed. Winter data have been added to the graphs concerning leaves, twigs, roots, and soil compartments. In the winter, xylem and phloem tissues were not sampled in order to limit damage on the trunks.

8. Were samples taken randomly and from all trees at all dates? How were the phloem and xylem sampled? How were twigs and leaves selected?

Response: This has been detailed in the material and methods section: L110-117: "Leaves, twigs, trunk phloem and xylem and soil monoliths (15 cm depth, very few fine roots were present below 15 cm deep) of each labelled trees (1, 2, 3, 4, 5, 6) were sampled regularly after labelling until the end of 2010 (Table 1). At each sampling date 20 leaves and 20 twigs were collected randomly throughout the crown. Sampling was always performed between 10:00 and 12:00 h UTC. The leaves were rinsed with distilled water to remove any excess ^{15}N . At each sampling date, two small disks of bark (14 mm diameter, 10 mm depth) were collected at 1.3 m height using a corer. Thereafter phloem and xylem tissues were separated by hand with a cutter blade."

9. Were multiple replicates taken at each time, allowing an uncertainty on each point to be calculated? Or is each individual point also a single measurement from a particular pool at a particular time? If so, how far can we trust the individual time series for each tree when individual measurements may not be representative of the actual mean of the pool in question?

Response: Due to technical and financial constraints we have analyzed at each date an aliquot of pooled leaves, pooled twigs or pooled roots. Nevertheless, as explain above (see point 5.), we have, at few dates, checked the repeatability of the analysis by analyzing for aliquots of a compartment L125-129.

10. I am also not sure if I follow the logic of the CFE extraction in the methodology. The commonly methodology of Vance (1987) should have a control extraction and a fumigation extraction otherwise treated identically, the difference of which is inferred to be the C or N contained in microbial biomass and liberated to the extractable pool by fumigation. Not only is no fumigation treatment mentioned (how long was it fumigated for, with what concentration of chloroform?) used for extraction (l116, 0.5M) is more than an order of magnitude than the concentration used for 'microbial ^{15}N abundance' (l118, 0.3M). It is not clear to me if this former is a 'control' unfumigated treatment and the latter is the ^{15}N fumigation treatment, or if a control (unfumigated) ^{15}N treatment was measured and is not reported. If the former, a 0.03 M solution may extract less N than 0.05 M, particularly for organic compounds (e.g. Makarov 2013, European Journal of Soil Science 46, 369-374) and estimates of microbial biomass N as the difference would be an underestimate. Also, ^{15}N extracts from low [N]/[^{15}N] samples such as microbial fumigation extracts are commonly concentrated using a diffusion trap method (Stark and Hart (1996). Soil Science Society of America Journal, 60, 1846–1855.). Was this performed here? If not, were ^{15}N contents high enough to be detectable on the IRMS? In my opinion, this section of the methods is weak and should either be entirely rewritten removed, along with corresponding results if the method was not robust enough for valid interpretation.

Response: All this methodology section has been completed : L134-143: "Microbial N contents of fresh soil samples were determined using the chloroform fumigation–extraction method (Vance et al., 1987). 2 fresh soil subsamples of 10 g were prepared. One subsample was fumigated for 24 h with chloroform vapour, while the other was not fumigated. Nitrogen extraction was performed using 50 mL of 0.5 M K_2SO_4 for 30 min under vigorous shaking. The extracts (fumigated and not fumigated) were filtered, then analysed for N content using an N analyser (TNM-1, Shimadzu, Champs-sur-Marne, France). The microbial ^{15}N abundance was estimated using the same procedure except that the extraction solution was 0.03 M of K_2SO_4 in order to avoid any alteration of the mass spectrometer with the K_2SO_4 salt during ^{15}N analysis".

11. Specific comments

L73 – how deep were the soil horizons (what would we expect to be sampled by the 15 cm corer later used?)

Response: Very few fine roots were present below 15 cm depth due to the edaphic properties of the site: gley mainly presents less than 15cm depth.

12. L74 – nitrogen deposition, if known, might be useful to include here as this study concerns N additions. High soil N availability may affect the origin of N for growth.

Response: N deposition on Fontainebleau forests was in average 8 kgN/ha/ year (Reneco for Data, National Network for Long-term FOrest ECOsystem Monitoring, 1998). More recent data estimated N through fall in Fontainebleau forest between 5 and 10 kgN/ha/ year in 2010 (Waldner et al. 2014). These quantities do not induce high N availability in soil.

13. L77 – how big were the trenched areas?

Response: It was mentioned L80: 5m² in average

14. What was the spacing of the trees?

Response: At least 20m, see L81.

15. L86 - were treatments applied in particular weather conditions? Logically, it would make sense to maximize uptake of foliar N by applying the N treatment on dry days so it is not immediately lost by being washed off the leaves.

Response: Treatments were applied on sunny days L87.

16. L90 – can you estimate how much of the sprayed N remained on the trees after application and how much was lost immediately, falling onto the plastic tarpaulin?

Response: We have not estimated this.

17. L92 – how long was the plastic tarpaulin in place? Was this long enough to prevent losses from leaf leaching (L237) from reaching the soil?

Response: L95-98: “The plastic tarpaulin remained on the soil during 2 weeks after labelling. Before removing the plastic tarpaulin, crowns were sprayed with distilled water in order to avoid any soil contamination after the removing of the tarpaulin”.

18. L110 – were these grounds by hand, or in a mill? Were the samples dried, e.g. in an oven, before this?

Response: This section has been completed: L120-133 : “All plant tissues and soil samples were brought to the laboratory in a cooler. Plant tissues were lyophilized and ground to a fine powder with a ball mill before analyses. At each sampling date, one aliquot of each plant powder (1 mg) was transferred into tin capsules (Elemental Microanalysis, UK, 6 x 4 mm, ref. D1006, BN/139877). At some date (Day After Labelling (DAL) 1, 126, 337, 460 for leaves and twigs of L₁; DAL 126, 337, 460 for roots of L₁; DAL 227 and 350 for leaves and twigs of L₂; DAL 49 and 350 for roots of L₂ and DAL 40 and 166 for leaves, twigs and roots of L₃), four aliquots of powder was transferred into tin caps in order to test the repeatability of the analysis. Total N concentration of plant and soil samples, was analysed by dry combustion using an N auto-analyser (Flash EA 1112 series, Thermofinnigan). ¹⁵N abundance was quantified in the same plant and soil fine powder aliquots with a mass spectrometer (PDZ Europa, University of Davis, Isotopes Facility, California)”.

19. L121 – I feel that something is needed here to justify this approach rather than having a concurrent control unlabeled set of trees.

Response: L146-151: “All ¹⁵N enrichments were corrected for the background natural abundance of this isotope, using control values determined in plants and soils just before labelling. The seasonal variations of the natural ¹⁵N abundance of each compartments were also followed all long the season, those variations were very weak, consequently, it has been choose to use the ¹⁵N natural abundance of the labelled trees just before labelling.”

20. L143 – See general comments about this statement. Also, were these similar patterns in TOTAL recovered N, or PRN? From the manuscript it appears it was the latter but the former may also be informative.

Response: The patterns of total recovered N were also similar between both trees. L169-170.

21. L237 – N remaining on leaves could also be lost by stemflow or throughfall and washed to the base of the stem. How were the plastic tarpaulins (if in place at this time) sealed around the stem?

Response: L94-95: The tarpaulin was sealed to the trunk at 50 cm height with Terostat-VII (Teroson, Henkel, Germany).

22. L241 – Maybe this needs a little more elaboration. Allocation of ^{15}N to non-harvested components is assumed as there is not a better explanation.

Later (L270) literature begins to be cited about storage of N – this could be incorporated into here to explain where the missing ^{15}N is going.

Response: This has been completed L281-283.

23. L271 – presumably leaf senescence is important for the constitution of N stores in deciduous plants rather than evergreen conifers, where seasonal N storage in leaves is driven by a mismatch of rates N uptake and photosynthetic C late in the growing season.

L274 – Should this be evident from fig. 1b? It appears from this panel and table 1 that root N was measured 2-3 months before yellowing (DAL 57), just before the yellowing event (DAL 126) and again after budburst (DAL318). Is this enough resolution to tell whether this N was stored in fine roots at leaf senescence, or if root ^{15}N increased earlier in the growing season and subsequently declined over the winter. The two replicates do not agree over the winter period – one is fairly level and one steeply declines. Admittedly the literature suggests that this is a storage pool but I think this may be over-interpreting these particular data.

Response: A point completes the series of measure in winter. At day 189, the proportion of recovered ^{15}N in roots is quite similar for both tree (17 and 21%) Indeed this point was not previously presented because at this sampling date (DAL 189) phloem tissue was not sampled in order to limit damage caused to the trunk. Now we clearly observe that there was an increase of ^{15}N recovered in fine roots in autumn followed by a slight decrease during winter L191-192, L205-207.

24. L304-305 – With no indication of uncertainty, it is rather speculative to interpret differences this small as real changes!

Response: Indeed fluctuations are very small but our hypotheses were supported by previous experiment (Barnard et al., 2006; Bloor et al., 2009).

25. L309 – with no measurements over winter, is this a reasonable interpretation? Could N continue to be taken up but also be decline prior to budburst? A brief mention of a lack of change in above-ground biomass outside the growing season (if true) could help explain this.

Response: The added winter point completes the series. The proportion of ^{15}N recovered in roots was lower in December than in October (5.5% vs. 4%) L206, which confirms the limitation of N soil uptake during winter. L344-349: “After leaf fall, trees may have a significant capacity for nitrate uptake in the fine roots in midwinter (i.e. in the absence of

leaves), as already shown in Japan oak (Ueda et al., 2010). However, in our case, N soil uptake was limited by low soil temperature, which affected the mineralization rate and root activity, since the ^{15}N recovered from roots slightly decrease between October 28 and December 2 (5.5% to 4%) and then declined to 1.75% between December 2 and April 8".

26. L344 – a reference for cessation of glutamine synthetase activity would be useful.

Response: Our team has conducted analyzes of root enzyme activities in mature oaks throughout a season, the results show a reduction in GS activity in winter, these data are published in *Trees structure and Function* : Bazot et al., 2013, L381.

27. L393 – This final sentence is unnecessary as this suggestion for further work does not feel like a natural result of the conclusions of the manuscript.

Response: It has been removed.

28. Table 1 is very confusing. DAL for different treatments are not the same thing as the labelling occurs at different times of the year. I wonder if this can be reformatted in a way that allows for easier interpretation, perhaps by playing the data for trees 3 and 4 and 5 and 6 at positions in the table so that the real-time day of year is close to equivalent horizontally or by splitting this into three tables, one for each set of trees. Also, are the "Amount of ^{15}N sprayed", and "Budburst" rows necessary, given that it is the same in all treatments? Budburst could instead be indicated by an entry in the table.

Response: Table 1 was simplified and Julian day numbers have been added in order to facilitate the reading of sampling days.

29. The figures need a clearer distinction between of trees. It would be nice to be able to tell which time series is from which tree. Axis titles could be the full, unabbreviated units as these are not particularly long phrases and are not standard terms which the reader can be assumed to already know. Additionally, the legends indicate that the dashed lines/continuous lines are for the different trees, but the figure legend suggests the dashed lines are the biomass pools. This should be checked across all graphs for consistency.

Response: The graphs were corrected according to those recommendations.

30. Technical corrections

L27 – the sentence 'the literature describes is unnecessary.

Response: It has been deleted

L75 – include authority with species name

Response: It has been done

L89 – 'on' April 20, or 'by' April 20?

L106 – the 'leaf mass area' (LMA) should be 'leaf mass per area'.

Response: It has been done

L124 – is this the same six trees as measured?

Response: No, but they are similar trees grown on the same site under the same conditions with the same size.

L129 – (PRN) would be easier to interpret if it immediately follows 'proportion of recovered ^{15}N Nitrogen'

Response: It has been done

L144 – The sentence 'That why results were expressed as the mean of both trees' is

poor English and should read 'That is why results were

Response: It has been corrected

L 147 – 'leafy season' - > 'growing season'?

Response: No, Growing season (of the trunk) ended in July, whereas leaves fall at the end of September

L232 – Substantial fraction is ambiguous

Response: Replaced by "significant"

L238 – the lack of contamination could be supported by referring to figure 1d.

Response: It has been done

L265 – this is repetitive, and along with section 4.1 could be considerably shortened.

Generally, this section is repeating something that is apparent from the results.

Response: This section was reduced

L309 – this sentence is very long – could it be split up?

Response: This sentence was split.

References – numerous cases where super- or subscript is not used in reference list

(e.g. line 412 '15N')

Response: It has been corrected

Figure legends: remove 'the' from 'the tree 1' and 'the tree 2'.

Response: It has been done

Figure 1d – the scale on this figure is different than the other graphs. This makes interpretation difficult. Could this be adjusted or measured in the legend?

Response: We have specified this different scale in the legend of the figure.

Referee #2 :

General comments

1. The paper represents a significant contribution to the elucidation of N flows in trees. There is no new concept or method in this study but the use of three different ¹⁵N-labelling periods (spring year n-1, autumn year n-1 and spring year n) and two labelling techniques (soil and leaf labelling) allows a good description of the contribution of leaves to the constitution of winter reserves and the contribution of N reserves to spring growth. It is clearly shown that the main contributor to the synthesis of new leaves is N stored during previous autumn. It is also shown that soil micro-organisms are good competitors for soil ¹⁵N but a significant part of the N is returned to the tree because of microbial turnover. This is a well-written paper, and a well-thought out analysis. In my opinion, the subject and the core-content of the ms are appropriate and relevant to Biogeosciences. The findings are reliable because the methods developed are appropriate. I have just a problem concerning xylem and phloem measurements. Nothing is mentioned concerning phloem and xylem sampling and how the contribution of these pools to ¹⁵N partitioning is estimated.

Response: At each sampling date, two small disks of bark (14 mm diameter, 10 mm depth) were collected at 1.3 m height using a corer. Thereafter phloem and xylem tissues were separated by hand with a cutter blade."L115-118.

2. Also, for obvious technical reasons, ¹⁵N allocated to coarse roots and trunk is not taken in account in this study. It is known that these organs represent a substantial pool of N reserves and this should be discussed.

Response: Discussion was complete : L281-283 : “Indeed, data currently available on woody plants show that nitrogen is re-translocated from leaves to storage sites such as old branches, trunk or coarse roots (Valenzuela Nunez et al., 2011; Bazot et al., 2013).”

3. Specific points Abstract Line 12 is this proportion (30 %) true for all labelling periods?

Response: Yes, for L₁, 32% of administered ¹⁵N was recovered L179; for L₂, 70% of administered ¹⁵N L195, and for L₃, 51.5% of administered ¹⁵N was recovered L244.

4. Material and methods Sampling. One can understand that the authors used only two replicates for each labelling for technical reasons even if it is difficult to generalize from six trees. However, the authors should be much more accurate concerning the samplings (number of sampling per tree, soil, phloem and xylem sampling) to improve this section and strengthen the validity of the conclusion.

Response: The sampling procedure was completed:

L110-118: “Leaves, twigs, trunk phloem and xylem and soil monoliths (15 cm depth, very few fine roots were present below 15 cm deep) of each labelled trees (1, 2, 3, 4, 5, 6) were sampled after labelling until the end of 2010 (Table 1). At each sampling date 20 leaves and 20 twigs were collected randomly throughout the crown. Sampling was always performed between 10:00 and 12:00 h UTC. The leaves were rinsed with distilled water to remove any excess ¹⁵N. At each sampling date, two small disks of bark (14 mm diameter, 10 mm depth) were collected at 1.3 m height using a corer. Thereafter phloem and xylem tissues were separated by hand with a cutter blade.”

L120-133: “All plant tissues and soil samples were brought to the laboratory in a cooler. Plant tissues were lyophilized and ground to a fine powder with a ball mill before analyses. At each sampling date, one aliquot of each plant powder (1 mg) was transferred into tin capsules (Elemental Microanalysis, UK, 6 x 4 mm, ref. D1006, BN/139877). At some date (Day After Labelling (DAL) 1, 126, 337, 460 for leaves and twigs of L₁; DAL 126, 337, 460 for roots of L₁; DAL 227 and 350 for leaves and twigs of L₂; DAL 49 and 350 for roots of L₂ and DAL 40 and 166 for leaves, twigs and roots of L₃), four aliquots of powder was transferred into tin caps in order to test the repeatability of the analysis. Total N concentration of plant and soil samples, was analysed by dry combustion using an N auto-analyser (Flash EA 1112 series, Thermofinnigan). ¹⁵N abundance was quantified in the same plant and soil fine powder aliquots with a mass spectrometer (PDZ Europa, University of Davis, Isotopes Facility, California)”.

5. Results Line 144-145 there is no verb in this sentence.

Response: The sentence was rewritten

6. I presume also there is a mistake, L2: 3+4 instead of 2+3 and L3: 5+6 instead of 3+4.

Response: It was corrected

7. Discussion Line 343-348. The authors should be much more careful here. I really do not know why the authors mention the Glutamine synthetase/Glutamate synthase pathway as no results shown in the paper concern amino acid metabolism. I presume this hypothesis is based on published literature which is not mentioned.

Response: Our team has conducted analyzes of root enzyme activities in mature oaks throughout a season, the results show a reduction in activity GS GOCAT in winter, these data are published in *Trees structure and Function* : Bazot et al., 2013 L381.

8. Also, the Morot-Gaudry reference is not in the reference list, and I am not sure it concerns tree physiology. I suspect there are more appropriate references concerning tree N assimilation.

Response: Morot Gaudry also presents conclusion about tree N assimilation in this book (p299). Gojon et al 1991 could be also mentioned, indeed they underlined that roots were the main site of NO₃⁻ reduction in *Prunus persica* L.

1

2 **Marked-up manuscript version**

3

4 **Contribution of previous year's leaf N and soil N uptake to current year's leaf**
5 **growth in sessile oak**

6

7 **STEPHANE BAZOT*, CHANTAL FRESNEAU, CLAIRE DAMESIN, LAURE**
8 **BARTHES**

9

10 Ecologie Systématique et Evolution, Univ-Paris-Sud, CNRS, AgroParisTech,
11 Université Paris Saclay, rue du Doyen A. Guinier, Orsay, F-91405, Orsay, France

12

13 ***Author for correspondence**

14 Stéphane Bazot

15 tel: (+33) 1 69 15 71 36

16 fax: (+33) 1 69 15 72 38

17 email: stephane.bazot@u-psud.fr

18

1 **Abstract**

2 The origin of the N which contributes to the synthesis of N reserves of *in situ* forest
3 trees in autumn, and to the growth of new organs the following spring, is currently
4 poorly documented. To characterize the metabolism of various possible N sources
5 (plant N and soil N), six distinct 20 year-old sessile oaks were ¹⁵N labelled by
6 spraying ¹⁵NH₄¹⁵NO₃: (i) on leaves in May, to label the N pool remobilized in the
7 autumn for synthesis of reserves; (ii) on soil in the autumn, to label the N pool taken
8 up from soil; (iii) on soil at the beginning of the following spring, to label the N pool
9 taken up from soil in the spring. The partitioning of ¹⁵N in leaves, twigs, phloem,
10 xylem, fine roots, rhizospheric soil and microbial biomass was followed during two
11 growing seasons. Results showed a significant incorporation of ¹⁵N in the soil-tree
12 system; more than 30% of the administered ¹⁵N was recovered. Analysis of the
13 partitioning clearly revealed that in autumn, roots' N reserves were formed from
14 foliage ¹⁵N (73%) and to a lesser extent from soil ¹⁵N (27%). The following spring,
15 ¹⁵N used for the synthesis of new leaves came first from ¹⁵N stored during the
16 previous autumn, mainly from ¹⁵N reserves formed from foliage (95%). Thereafter,
17 when leaves were fully expanded, ¹⁵N uptake from soil during the previous autumn
18 and before budburst contributed to the formation of new leaves (60%).

19 **keywords :**

20 *Quercus petraea*, N reserves, soil N, ¹⁵N labelling

21

22 1. Introduction

23 Tree carbon metabolism associated with photosynthesis, C allocation and
24 remobilization of C storage is well documented (Barbaroux et al., 2003; Dickson,
25 1989), but tree nitrogen metabolism is less known. Nevertheless, seasonal N cycling
26 is a determinant of plant fitness in perennials, particularly long-lived perennials such
27 as forest trees (Cooke and Weih, 2005). In early spring, trees' nitrogen demand for
28 growth can be satisfied either by uptake of external sources such as ammonium,
29 nitrate and organic N available from the soil (Gessler et al., 1998a), or by
30 remobilization of internal stores (Bazot et al., 2013; Coleman and Chen, 1993; Cooke
31 and Weih, 2005; El Zein et al., 2011b; Gilson et al., 2014; Millard, 1996; Taylor, 1967).
32 In many species, N remobilization for growth in spring occurs before utilization of N
33 taken up by roots, typically during the 20–30 days before the roots actively take up
34 N. These species include: deciduous species, such as *Quercus petraea* (El Zein et al.,
35 2011a), *Malus domestica* (Guak et al., 2003; Neilsen et al., 2001), *Populus*
36 *trichocarpa* (Millard et al., 2006), *Prunus avium* (Grassi et al., 2003), *Pyrus*
37 *communis* (Tagliavini et al., 1997) and *Sorbus aucuparia* (Millard et al., 2001);
38 marcescent/evergreen species, such as *Nothofagus fusca* (Stephens et al., 2001); and
39 coniferous evergreens, such as *Picea sitchensis* (Millard and Proe, 1993). In a few
40 species (e.g., *S. aucuparia*), remobilization has completely finished before any root
41 uptake of N occurs, even if trees are supplied with an adequate supply of mineral N
42 in the soil. In contrast, other species have been shown to begin taking up soil N
43 through their roots concomitantly with N remobilization. These include deciduous
44 *Juglans nigra × regia* (Frak et al., 2002), *Pyrus communis* (Tagliavini et al., 1997),
45 *Betula pendula* and evergreen *Pinus sylvestris* (Millard et al., 2001). All of these
46 studies were conducted on young trees or/and under controlled conditions. Few

Supprimé: The literature describes general patterns of seasonal tree nitrogen functioning as follows.

50 studies have applied ^{15}N -labeled mineral fertilizer to larger, undisturbed trees
51 growing in the field (El Zein et al., 2011a), and even those only evaluated the
52 contribution of spring N uptake to leaf and twig growth, while the contribution of
53 stored N was indirectly estimated. However, in autumn, the process of N storage (N
54 translocation from leaves to sink compartments), which starts concomitantly with
55 leaf yellowing (Bazot et al., 2013), is associated with a stimulation of soil nitrogen
56 uptake (Gessler et al., 1998b; Jordan et al., 2012; Kim et al., 2009). In the present
57 study we proposed to investigate the contribution of N storage and that of N taken up
58 from soil during autumn and spring, to the development of new leaves of 20 year-old
59 sessile oaks in the field, after budburst during the following spring. Does soil N or
60 foliar N contribute most to the storage of N compounds in autumn? Does soil N or
61 stored N contribute most to the synthesis of new leaves in spring? Soil ^{15}N labelling
62 is a suitable tool to quantify autumn and spring uptake of N by roots. Labelling of
63 foliage allows quantification of N remobilized from leaves to reserve compartments.
64 During three distinct labelling campaigns, 3 x 2 distinct 20-year-old sessile oaks
65 received $^{15}\text{NH}_4^{15}\text{NO}_3$ applied to their foliage (May), or on adjacent soil (September
66 and March of the following year). ^{15}N partitioning in all tree-soil compartments, i.e.
67 leaves, twigs, trunk, roots, rhizospheric soil and microbial biomass, was analysed
68 regularly. The contribution of assimilated ^{15}N to storage and remobilization was
69 investigated.

70

71 **2. Materials and methods**

72 **2.1. Site description**

73 The experiment was conducted in an area of 20-year-old naturally regenerated oak in
74 the Barbeau forest (48°29'N, 02°47'E), 60 km southeast of Paris, France, at an

75 elevation of 90 m on a gleyic luvisol. The average air temperature is 10.5 °C and the
76 annual rainfall in this temperate location is 690 mm. Six 20-year-old sessile oaks
77 (*Quercus petraea*, L.) were selected, their height ranged between 8 to 10 m and their
78 average diameter at breast height was 10 cm. In order to limit possible interference of
79 root cutting with nitrogen allocation, at least five months before labelling a 0.5-0.6 m
80 deep trench was dug around each tree, then the trench was lined with a polyethylene
81 film and backfilled. All roots and root exudates inside this perimeter therefore
82 originated from the isolated tree, and were contained in this trench volume. The area
83 delimited by the trench was about 5 m². The distance between each tree was at least
84 20 m.

Supprimé:)

86 2.2. ¹⁵N pulse-labelling

87 Three labelling campaigns were carried out: the first (L₁) on the foliage at the end of
88 May (2009/05/27); the second (L₂) on the soil at the beginning of September
89 (2009/09/09); and the third (L₃) on the soil the following March (2010/03/20). All
90 labelling campaigns were conducted on sunny days. Two oaks were labelled during
91 each campaign: trees 1 and 2 during L₁; trees 3 and 4 during L₂; and trees 5 and 6
92 during L₃. 50% of buds showing leaf unfolding (Vitasse et al., 2009), occurred in
93 those sessile oaks on April 20, 2010; this date was defined as budburst. The L₁
94 campaign consisted of homogenous spraying on all foliage of 5g ¹⁵NH₄¹⁵NO₃ (98
95 atom %), i.e. 1.82g of ¹⁵N, dissolved in 2.5 L distilled water. Prior to L₁, soil of the
96 surrounding trenches was protected with a plastic tarpaulin covering the whole area
97 of the trenched plot, to avoid soil pollution with ¹⁵N. The tarpaulin was sealed to the
98 trunk at 50 cm height with Terostat-VII (Teroson, Henkel, Germany). It was
99 remained on the soil during 2 weeks after labelling. Before removing the plastic

Supprimé: to avoid soil pollution with ¹⁵N.

103 tarpaulin, crowns were sprayed with distilled water in order to avoid any soil
104 contamination after the removing of the tarpaulin.

105 This first campaign aimed at the labelling of foliage and, subsequently, of the N
106 reserves developed from remobilization of leaf N the following autumn. The L₂
107 campaign consisted of homogenous spraying of 5g ¹⁵NH₄¹⁵NO₃ (98 atom %), i.e.
108 1.82g of ¹⁵N, dissolved in 20 L distilled water on the soil of the trench plot of two
109 other selected oak trees (3 and 4). With this procedure, N reserves developed from
110 autumnal soil N uptake were expected to be labelled. The third and last labelling
111 campaign, L₃, consisted of homogenous spraying of 5g ¹⁵NH₄¹⁵NO₃ (98 atom %), i.e.
112 1.82g of ¹⁵N, dissolved in 20 L distilled water on the soil of the trench plot of trees 5
113 and 6, thus labelling their spring N uptake.

114

115 2.3. Sampling and analytical methods

116 Leaves, twigs, trunk phloem and xylem and soil monoliths (15 cm depth, very few
117 fine roots were present below 15 cm deep) of each labelled trees (1, 2, 3, 4, 5, 6)
118 were sampled after labelling until the end of 2010 (Table 1). At each sampling date
119 20 leaves and 20 twigs were collected randomly throughout the crown. Sampling was
120 always performed between 10:00 and 12:00 h UTC. The leaves were rinsed with
121 distilled water to remove any excess ¹⁵N. At each sampling date, two small disks of
122 bark (14 mm diameter, 10 mm depth) were collected at 1.3 m height using a corer.
123 Thereafter phloem and xylem tissues were separated by hand with a cutter blade. The
124 leaf mass per area (LMA) was measured at each sampling date. Fine roots were
125 hand-picked from the soil monoliths, and washed with a 0.5 M CaCl₂ isotonic
126 solution. Soil adhering to roots was removed with a brush and sieved at 2 mm. All
127 plant tissues and soil samples were brought to the laboratory in a cooler. Plant tissues

Supprimé: regularly

Supprimé: Total N concentration of plant and soil samples, ground in fine powder

131 were lyophilized and ground to a fine powder with a ball mill before analyses. At
132 each sampling date, one aliquot of each plant powder (1 mg) was transferred into tin
133 capsules (Elemental Microanalysis, UK, 6 x 4 mm, ref. D1006, BN/139877). At
134 some date (Day After Labelling (DAL) 1, 126, 337, 460 for leaves and twigs of L₁;
135 DAL 126, 337, 460 for roots of L₁; DAL 227 and 350 for leaves and twigs of L₂;
136 DAL 49 and 350 for roots of L₂ and DAL 40 and 166 for leaves, twigs and roots of
137 L₃), four aliquots of powder was transferred into tin caps in order to test the
138 repeatability of the analysis. Total N concentration of plant and soil samples, was
139 analysed by dry combustion using an N auto-analyser (Flash EA 1112 series,
140 Thermofinnigan). ¹⁵N abundance was quantified in the same plant and soil fine
141 powder aliquots with a mass spectrometer (PDZ Europa, University of Davis,
142 Isotopes Facility, California).

143 Microbial N contents of fresh soil samples were determined using the chloroform
144 fumigation–extraction method (Vance et al., 1987). 2 fresh soil subsamples of 10 g
145 were prepared. One subsample was fumigated for 24 h with chloroform vapour,
146 while the other was not fumigated. Nitrogen extraction was performed using 50 mL
147 of 0.5 M K₂SO₄ for 30 min under vigorous shaking. The extracts (fumigated and not
148 fumigated) were filtered, then analysed for N content using an N analyser (TNM-1,
149 Shimadzu, Champs-sur-Marne, France). The microbial ¹⁵N abundance was estimated
150 using the same procedure except that the extraction solution was 0.03 M of K₂SO₄ in
151 order to avoid any alteration of the mass spectrometer with the K₂SO₄ salt during ¹⁵N
152 analysis.

154 2.4. Calculations

Supprimé: Extraction

Supprimé: of

Supprimé: .

158 All ^{15}N enrichments were corrected for the background natural abundance of this
 159 isotope, using control values determined in plants and soils just before labelling. The
 160 seasonal variations of the natural ^{15}N abundance of each compartments were also
 161 followed all long the season, those variations were very weak, consequently, it has
 162 been choose to use the ^{15}N natural abundance of the labelled trees just before
 163 labelling. The total weight of each compartment analysed (i.e. leaves, twigs, trunk
 164 phloem and xylem, and fine roots) was extrapolated from that of six equivalent trees
 165 (same size and same diameter) grown on the same site under the same conditions.
 166 Those trees were felled as follows: two in October of the first labelling year (2009);
 167 two in the following May (2010); and two the following February (2011). Total leaf
 168 biomass was corrected according to the LMA. All data were expressed as proportion
 169 of recovered ^{15}N nitrogen (PRN) in a specific compartment, using the following
 170 calculation Eq. (1):

$$171 \quad \text{PRN \%} = \frac{Q^{15}\text{N}_{\text{compartment}}}{\text{Max } Q^{15}\text{N}} \times 100$$

172 where $Q^{15}\text{N}$ was the quantity of ^{15}N recovered from a compartment on a specific
 173 date, and Max $Q^{15}\text{N}$ was the maximum quantity of ^{15}N recovered from all the
 174 sampled compartments during the experiment.

175 The % contribution of each ^{15}N source (L_1 : leaves; L_2 : autumn soil N; L_3 : spring
 176 soil N) to the ^{15}N recovered in the roots in autumn or in the leaves of the second year
 177 as determined according to the following calculation Eq. (2) :

$$178 \quad \% \text{ contribution } ^{15}\text{N}_{L_1, L_2, L_3} = \frac{(Q^{15}\text{N}_{\text{compartment}} / \text{Max } Q^{15}\text{N})_{L_1, L_2, L_3}}{\Sigma(Q^{15}\text{N}_{\text{compartment}} / \text{Max } Q^{15}\text{N})_{L_1, L_2, L_3}} \times 100$$

180 **3. Results**

Supprimé: (PRN)

Supprimé: ¶
¶

184 For each labelling, the two trees analysed displayed similar patterns of total
185 recovered ¹⁵N in each compartment (data not shown) and ¹⁵N partitioning throughout
186 the experiment. Moreover, the test of repeatability of the analysis revealed very few
187 variability of the ¹⁵N partitioning at a specific date in a specific compartment (Fig.
188 1, 2, 3). Consequently, results was expressed as the mean of both trees (L₁ : 1+2; L₂ :
189 3+4, L₃ : 5+6).

Supprimé: That why

Supprimé: 2+

Supprimé: 3+4

191 3.1. ¹⁵N partitioning within the plant-soil system during the first leafy season

192 3.1.1. After the foliar labelling in spring (L₁, May 27, 2009)

Supprimé:

193 The total balance for the administered ¹⁵N demonstrated maximum recoveries of ¹⁵N
194 within the plant-soil system of 32% one day after leaf labelling. It decreased to
195 13.5% of the administered ¹⁵N recovered in the sampled compartments at the end of
196 September (126 days after labelling) (Table 1).

197 The PRN was maximum in leaves (96%, Fig. 1a) one day after L₁, then decreased
198 continuously during the four following months (from May 27 to September 30, 2009,
199 i.e. until the 126th day after labelling) with a mean decrease of 80% between these
200 two dates (Fig. 1a). The same pattern was observed in twigs, where the PRN
201 decreased from 3% on day 1 to 0.4% on day 126 (Fig. 1a).

202 In the trunk phloem tissue and the fine roots, the PRN stayed relatively stable or
203 slightly increased until day 57 (July 24, 2009). They then increased until day 126
204 (September 30, 2009), when they reached 4.75% in the phloem and 16% in the roots
205 (Fig. 1b, c). The PRN from the rhizospheric soil and microbial biomass was less than
206 1% (Fig. 1d). During winter (December 2, 2009; day 189) the PRN reached 18.5% in
207 fine roots (Fig.1c).

208 3.1.2. After the first soil labelling (L₂, September 9, 2009)

213 The total balance for the administered ^{15}N demonstrated maximum recoveries within
214 the plant-soil systems three days after L_2 of 70%. By the end of October (49 days
215 after labelling), recoveries from the sampled compartments decreased to 22% of the
216 administered ^{15}N (Table 1).

217 Three days after labelling, 3% of the recovered ^{15}N was present from the fine roots
218 (Fig. 2c). Nine days after labelling (September 18, 2009), the PRN showed that the
219 majority of the ^{15}N was recovered from the soil, with 61% of the ^{15}N recovered from
220 the rhizospheric soil and 32.5% from the microbial biomass (Fig. 2d). During the
221 following 40 days (until October 28, 2009), the PRN from the soil decreased to 8.5%
222 in the rhizospheric soil and 9.5% in the microbial biomass (Fig. 2d). On the same
223 date, 6% of the ^{15}N was recovered from the fine roots (Fig. 2c). Less than 1% of the
224 ^{15}N was recovered from the phloem, xylem and twigs (Fig. 2a, b). In December (day
225 84) the PRN from the soil was similar to that of the previous date and 4% of the ^{15}N
226 was recovered from the fine roots (Fig. 2c, d).

227

228 **3.2. ^{15}N partitioning within plant-soil system before and after budburst**

229 Almost one year after the first labelling (L_1), and before budburst (April 8, 2010, 318
230 days after labelling), 7.5% of the ^{15}N were recovered in the sampled compartments.
231 Thereafter, recovery remained stable at around 12% until September (460 days after
232 labelling, Table 1).

233 On April 8, 2010, i.e. 318 days after L_1 , 11.5% of the recovered ^{15}N was found in
234 fine roots (Fig. 1 c). Twigs contained 4.5% of recovered ^{15}N (Fig. 1a), while phloem
235 contained 4% (Fig. 1b). Less than 0.5% of ^{15}N was recovered from the rhizospheric
236 soil and microbial biomass (Fig. 1d).

Supprimé :)

238 Eight days after budburst (April 28, i.e. 337 days after L₁), 25% of the recovered ¹⁵N
239 was observed in new leaves. By May 19, this had decreased to 17% (Fig. 1a). On
240 April 28, twigs contained 3.5% of the recovered ¹⁵N (Fig.1 a), phloem 4% (Fig. 1b)
241 and fine roots 10% (Fig.1c). From then until September (i.e. 460 days after
242 labelling), the PRN from leaves remained relatively stable (22%), whereas it largely
243 decreased in fine roots (0.35%) (Fig. 1a, b, c). Less than 0.2% of the total ¹⁵N
244 recovered over the season was from the rhizospheric soil and microbial biomass (Fig.
245 1d).

246 Just before budburst following the second labelling (L₂, April 8, 2010, 208 days after
247 labelling) 19% of the ¹⁵N administered were recovered from all the analysed
248 compartments (Table 1). Most of it was from the rhizospheric soil (14.5%, Fig. 2d).
249 The microbial biomass contained 9.5% of the recovered ¹⁵N and the fine roots 2%
250 (Fig. 2d, c). The rest of the ¹⁵N (less than 5%) was distributed between the twigs,
251 trunk phloem and xylem (Fig. 2a, b). The same pattern was observed eight days after
252 budburst (227 days after labelling): most of ¹⁵N was recovered from soil microbial
253 biomass and rhizospheric soil (12%, Fig. 2d); 2.25% was recovered from fine roots;
254 3.5% of ¹⁵N was recovered from phloem and xylem; only 0.5% was recovered from
255 new leaves (Fig. 2a).

256 From April 8 (208 days after labelling) to May 19 (247 days after labelling, and 30
257 days after budburst), the PRN decreased in soil microbial biomass and rhizospheric
258 soil (7%), but increased in fine roots (9.5%) (Fig. 2 d, c). A noticeable increase of the
259 PRN from leaves was also observed at this date (4.5%, Fig. 2a). Thereafter, the PRN
260 from soil microbial biomass and fine roots decreased slightly from May 19 to June
261 28 (i.e. 247 to 287 days after labelling), then remained stable until the end of August
262 (Fig. 2d, c). The PRN from leaves increased to 7% in June (Fig. 2a).

263 For trees whose soils were labelled in spring (L₃, March 20, 2010), the maximum
264 recovery of the administered ¹⁵N occurred 40 days later: 51.5% from the sampled
265 compartments. Recovery decreased thereafter and stabilized at 19.5% until autumn
266 2010 (Table 1).

267 Twenty days after labelling and before budburst, the soil microbial biomass
268 contained 44.5% of the recovered ¹⁵N and the rhizospheric soil 39% (Fig. 3d). The
269 remaining ¹⁵N was mainly located in the roots (2% of recovered ¹⁵N, Fig. 3c). 8 days
270 after budburst, the PRN was quite similar: 61% in microbial biomass and 32% in
271 rhizospheric soil (Fig. 3d). ¹⁵N recovered from fine roots followed a pattern similar
272 to that observed on the previous sampling occasion (Fig. 3c). However, between 8
273 and 30 days after budburst (from April 28 to May 19, 2010 i.e. from 40 to 61 days
274 after labelling), the PRN in microbial biomass and in rhizospheric soil decreased
275 sharply to 3.2% (Fig. 3d). On that date, 17% of the ¹⁵N was recovered from the fine
276 roots (Fig. 3c) and 21.2% from the leaves (Fig. 3a). The PRN from leaves remained
277 stable until the beginning of June (74 days after labelling) (Fig. 3a). From that date
278 until September the PRN from leaves and fine roots declined slightly (Fig. 3a, c).
279 The PRN from microbial biomass decreased continuously throughout the season and
280 reached 2.5% in September (day 166 after labelling) (Fig. 3d).

281

282 **4. Discussion**

283 **4.1. Efficiency of labelling**

284 Isotope labelling experiments are technically challenging, and as a consequence are
285 very scarce on trees growing in natural conditions. In this paper, field labelling
286 campaigns were conducted on 20-year-old naturally regenerated oaks. For each
287 campaign (only) two trees were labelled. Nevertheless the similarity of the results

288 between them suggests that the observed ^{15}N partitioning in soil and tree is a
289 representative view of the functioning of such systems

290 During the first labelling procedure (L_1), a significant fraction of the added
291 $^{15}\text{NH}_4^{15}\text{NO}_3$ was incorporated into the leaves of the sessile oaks. A significant

292 proportion of the ^{15}N was allocated to the leaves: more than 90% of the ^{15}N was
293 recovered from this compartment. The total balance for the administered ^{15}N

294 demonstrated maximum recoveries within the plant-soil systems of 32% one day
295 after leaf labelling. The remaining ^{15}N was probably lost by leaf leaching. However,

296 soil protection with plastic tarpaulins avoided all contamination of soil and roots, as
297 indicated by the ^{15}N recovered in the belowground compartments (Fig. 1d).

298 Thereafter, the recovery of administered ^{15}N from the sampled compartments
299 decreased to 14.5%, probably due to allocation of ^{15}N to non-harvested
300 compartments, such as old branches, coarse roots or the inner part of the trunk.

301 Indeed, data currently available on woody plants show that nitrogen is re-
302 translocated from leaves to storage sites such as old branches, trunk or coarse roots

303 (Valenzuela Nunez et al., 2011; Bazot et al., 2013). The soil $^{15}\text{NH}_4^{15}\text{NO}_3$ labelling
304 (L_2) conducted in September was also effective. Indeed, the total balance for the ^{15}N

305 applied to the soil demonstrated maximum recoveries within the plant-soil systems of
306 70%; 3 days after soil labelling. The rest of the ^{15}N was most probably lost by soil

307 leaching, (30% of the ^{15}N provide). Thereafter the recovery of administered ^{15}N from
308 the harvested compartments decreased to 22%. As with the leaf-labelling experiment

309 (L_1), this decrease was presumably due to allocation of ^{15}N to non-harvested
310 compartments. Finally, the soil $^{15}\text{NH}_4^{15}\text{NO}_3$ labelling carried out the following March

311 (L_3) was also effective, with maximum recoveries within the plant-soil systems of

Supprimé: substantial

Supprimé: then

Supprimé: .

Supprimé: Nevertheless, this labelling procedure allowed us to label foliar N used in the synthesis of new tissues in spring, and to follow the remobilisation of leaves' N in autumn, the contribution of foliar N to N storage, and the importance of N storage to the synthesis of new compartments the following spring. ¶

Supprimé: .

Supprimé: This labelling procedure allowed us to follow the contribution of autumn soil N to internal tree N storage. ¶

327 51.5%, 40 days after soil ^{15}N labelling. This recovery decreased to a mean of 19%
328 during the rest of the season. ▾

Supprimé: This labelling allowed us to follow the contribution of soil N in spring to the synthesis of new compartments from that moment until after budburst.

330 4.2. N dynamics in soil-tree systems during the first **leafy** season

Supprimé: growing

331 Following the first labelling procedure, the ^{15}N was quickly incorporated into leaves;
332 more than 90% of the ^{15}N applied was accounted for in leaves one day after
333 labelling. Thereafter this portion decreased continuously along the season. The
334 unaccounted for fraction of the ^{15}N had presumably been transferred to other
335 compartments, including those which were not sampled, i.e. branches and coarse
336 roots.

337 This important foliar N remobilisation was observed to continue in leaf-labelled trees
338 until yellowing, i.e. the end of September. Data currently available on woody plants
339 show that nitrogen is mainly re-translocated from leaves to storage sites during the
340 autumn (Coleman and Chen, 1993;Cooke and Weih, 2005;Dong et al., 2002;Taylor,
341 1967), due to the predominant role of leaf senescence in the constitution of N stores.
342 Leaf senescence leads to the breakdown of leaf proteins, the transfer of their nitrogen
343 to the perennial plant parts and the formation of N storage compounds (vegetative
344 storage proteins and amino acids) (Dong et al., 2000;Tromp, 1983). In this study, a
345 noticeable increase of percentage of recovered ^{15}N in fine roots was observed on
346 September 30 (16%). This compartment could be defined as a storage compartment
347 in young sessile oaks. Such an observation has been already reported for oaks of the
348 same pole stand (Gilson et al., 2014), and similar findings were reported for field-
349 grown adult peach trees by Tagliavini et al (1997), being typical of other young
350 deciduous trees (Millard and Proe, 1991;Salaün et al., 2005;Tromp and Ova, 1979;
351 Wendler and Millard, 1996). On this date (end of September), branches and

357 coarse roots could also have contributed significantly to N storage, as previously
358 described (Bazot et al., 2013).

359 At the same time, root uptake can also contribute directly to storage, as proposed by
360 Millard (1996). Indeed, 49 days after labelled ^{15}N had been applied to surrounding
361 soil (L_2), in September, 5.75% was recovered from the trees' fine roots. It can be
362 underlined that at the end of September, foliage ^{15}N made up 73% of the ^{15}N
363 recovered in roots, whereas soil ^{15}N uptake contributed to 27% of the ^{15}N recovered
364 in roots (eq. 2, Fig. 4). The soil N uptake in this period was mainly recovered in the
365 root system; there was little labelled N in the rest of the trees. This is consistent with
366 the results of Tagliavini et al (1997) and Jordan et al (2012), who found a significant
367 fraction of labelled N in fine root samples of peach trees supplied with ^{15}N applied
368 on soil before fruit harvest in September.

369 Concomitantly with root N uptake for storage, notably in fine roots, a strong
370 immobilization of N in microbial biomass was observed. Indeed, on October 7 (i.e.
371 28 days after labelling), when yellowing was well advanced, 12.5% of the applied
372 ^{15}N was recovered in microbial biomass and 21.5% in rhizospheric soil: there was a
373 competition for soil N between microbial N immobilization and reserve synthesis by
374 root N uptake at that time. This is consistent with the idea that soil microorganisms
375 are strong short term-competitors for soil N due to their high surface area to volume
376 ratio, wide spatial distribution in the soil and rapid growth rates, compared with
377 plants roots (Hodge et al., 2000). Thereafter, root N uptake was still efficient during
378 late yellowing (between October 7 and October 28), since ^{15}N recovered from the
379 fine roots slightly increased from 3.5% to 5.5%, whereas that recovered from
380 microbial biomass decreased from 12.5% to 10%. This could be explained by
381 microbial mortality and turnover, which releases N to the soil, combined with the

382 capacity of plants to sequester N for longer (Barnard et al., 2006;Bloor et al.,
383 2009;Hodge et al., 2000).

Code de champ modifié

384 After leaf fall, trees may have a significant capacity for nitrate uptake in the fine
385 roots in midwinter (i.e. in the absence of leaves), as already shown in Japan oak
386 (Ueda et al., 2010). However, in our case, N soil uptake was limited by low soil
387 temperature, which affected the mineralization rate and root activity, since the ¹⁵N
388 recovered from roots ~~slightly decrease between October 28 and December 2 (5.5% to~~
389 ~~4%) and then~~ declined to 1.75% between ~~December 2~~ and April 8.

Supprimé: even though

Supprimé: ,

Supprimé: from 5.5%

Supprimé: October 28

391 4.3. N dynamic in soil tree system the following spring

392 In April (before budburst), for trees with leaves labelled in the previous year (L₁), the
393 most part of ¹⁵N was recovered in their roots (11.5%). On the other hand, at the same
394 date, most of the labelled N applied to soil in September (L₂) was recovered from the
395 rhizospheric soil (14.5%). When soil (and hence spring N uptake) was labelled (L₃)
396 at the beginning of March, a month later most of the ¹⁵N was recovered from
397 microbial biomass and rhizospheric soil (81%), but a small proportion of ¹⁵N was
398 recovered from the fine roots (1.5%). The latter demonstrated a small N uptake
399 before budburst, as has previously been observed in Japan oak (Ueda et al., 2010).
400 This early N uptake from the soil could be related to sessile oak's hydraulic
401 properties. As a ring-porous species, sessile oak achieves 30% of its annual radial
402 stem growth before leaf expansion in spring (Breda and Granier, 1996). Water flow
403 pathways are then restored each spring before the onset of transpiration (Breda and
404 Granier, 1996). This enables early root N uptake from soil as soon as a threshold soil
405 temperature is reached.

Supprimé: %).

411 Eight days after budburst, most of the ^{15}N applied to leaves (L_1) was recovered from
412 new leaves (25.2%) and new twigs (mean of 3.5%). This clearly underlined that a
413 significant proportion of ^{15}N used to synthesize new leaves came from ^{15}N stored
414 during the previous autumn, as shown for *Ligustrum* (Salaün et al., 2005). Moreover,
415 this N came from foliar N of the previous year, not from soil N uptake during the
416 previous autumn. Indeed, trees labelled the previous autumn on soil (L_2) showed a
417 similar partitioning of ^{15}N in leaves and twigs before budburst (208 days after
418 labelling) and eight days after budburst (227 days after labelling), there was no
419 mobilisation of ^{15}N for the new leaves and twigs synthesis for those trees. Less than
420 1% of ^{15}N taken up from soil before budburst was recovered in leaves and twigs
421 eight days after budburst. A distinction might be made between stored N sourced
422 from leaves and that sourced from soil, stored mainly in roots. N from leaves could
423 be stored as amino acids in branches, trunk, and coarse roots, whereas N taken up
424 from soil could be stored in roots as NO_3^- . This N was not converted into amino acids
425 by Glutamine synthetase / Glutamate synthase enzymes during winter, most probably
426 due to low enzymatic activity in roots during winter, (Bazot et al., 2013). As a
427 consequence, the following spring, trees first remobilized easily circulating forms of
428 N, and N stored nearer to demands. Indeed in trees, NO_3^- is hardly transported to
429 their leaves but rather turned into amino acids in their roots (Morot-Gaudry, 1997).
430 Indeed roots were the main site of NO_3^- reduction (Gojon et al., 1991). Consequently,
431 soil ^{15}N was not the main contributor to the synthesis of new twigs and new leaves
432 during the eight first days after budburst. At this time, 95% of new leaves ^{15}N came
433 from ^{15}N -labelled reserves, 2% from soil labelled the previous autumn, and only 3%
434 from soil labelled in the current spring (Eq. 2, Fig. 4). Previous studies have also
435 found that N reserves contribute significantly to leaf expansion in young trees: in

Supprimé: .

Supprimé: . 1

438 white birch (Wendler and Millard, 1996); sycamore maple (Millard and Proe, 1991);
439 Japan oak (Ueda et al., 2009); pedunculate oak (Vizoso et al., 2008); and sessile oak
440 (El Zein et al., 2011a).

441 Considering trees whose soil had been labelled in autumn (L_2), eight days after
442 budburst the proportion of recovered ^{15}N in microbial biomass decreased slightly
443 whereas it slightly increased in fine roots compared to the previous sampling date.
444 One can suppose that the increased soil temperature and the first flux of C from plant
445 to soil (rhizodeposition) stimulated microbial biomass turnover, making ^{15}N
446 available for root uptake. Very little ^{15}N was recovered from the other compartments
447 of the trees.

448 Soil N uptake became really effective between 8 and 30 days after budburst. Indeed,
449 whatever the date of the soil labelling (autumn or the current spring), 30 days after
450 budburst, a sharp decrease in ^{15}N in the microbial biomass was observed, depending
451 on an increase of ^{15}N in fine roots and in young leaves. In June 28 (at leaf maturity),
452 40% of the ^{15}N recovered from leaves came from stored ^{15}N , 10% came from ^{15}N
453 applied to soil the previous autumn, and 40% came from ^{15}N applied on soil the
454 current March, one month before budburst (Eq. 2, Fig. 4). This pattern of
455 contribution was maintained throughout the season. Similar findings have been
456 reported for other species. For example, 20-30% of shoot leaf N was supplied by
457 spring-applied fertilizer for mature pear trees (Sanchez et al., 1990) and mature
458 almond trees (Weinbaum SA, 1984), while only 13% of a solution of nitrate-N and
459 ammonium-N applied to soil, contributed to total leaf N of apple trees (Nielsen et al.,
460 1997). *Sorbus aucuparia* had remobilized half the N from storage before any was
461 taken up by the roots (Millard et al., 2001). Finally, there is a concomitant/concurrent

462 remobilization and uptake of N from the soil by some other species, as shown for
463 scots pine (Millard et al., 2001) and walnut (Frak et al., 2002).

464

465 **5. Conclusion**

466 This paper completes knowledge of internal and external nitrogen cycles in a forest
467 ecosystem. We highlighted that in autumn, N reserves are formed from N
468 remobilized from leaves and N uptake by roots. This N is stored in roots, principally
469 most probably in the form of amino-acids and nitrate. Those reserves, especially N
470 coming from leaves, contributed significantly to new tissue synthesis the following
471 spring. Nevertheless, N uptake was also observed in spring before budburst; this N
472 was not transferred to new twigs and new leaves during the first days following
473 budburst. N uptake from soil only contributed significantly to the synthesis of new
474 tissues when leaves were fully expanded. Two months after budburst the relative
475 contributions of ^{15}N originating from leaves and ^{15}N uptake from soil were 40:60,
476 whereas they were 95:5 eight days after budburst.

477

478 **Acknowledgements**

479 Financial support was provided by Ecology Systematic Evolution lab and the French
480 National Research Agency through the “CATS Carbon Allocation in Tree and Soil
481 project”. The authors would like to acknowledge the contributions of Jérôme Ngao
482 and Daniel Berveiller to data collection and field work, and Michèle Viel and
483 Patricia Le Thuaut for technical assistance. We are grateful to the French National
484 Forest Office (ONF) for allowing us to carry out these experiments.

485

486 **Author contribution statement**

Supprimé: It will now be interesting to investigate soil N uptake, and the competition for N between tree and microorganisms (bacteria, fungi) in both autumn and spring.¶

492 SB and LB conceived and designed the experiments. SB, CF and LB conducted all
493 field and laboratory analyses. SB carried out data analysis, wrote most of the
494 manuscript and prepared the figures. CF, CD and LB contributed to the writing of the
495 manuscript.

496

497 **References**

498 Barbaroux, C., Bréda, N., and Dufrêne, E.: Distribution of above-ground and below-
499 ground carbohydrate reserves in adult trees of two contrasting broad-leaved species
500 (*Quercus petraea* and *Fagus sylvatica*), *New Phytologist*, 157, 605-615, 2003.

501 Barnard, R., Barthes, L., and Leadley, P.: Short-Term Uptake of ¹⁵N by a Grass and
502 Soil Micro-Organisms after Long-Term Exposure to Elevated CO₂, *Plant Soil*, 280,
503 91-99, 10.1007/s11104-005-2553-4, 2006.

504 Bazot, S., Barthes, L., Blanot, D., and Fresneau, C.: Distribution of non-structural
505 nitrogen and carbohydrate compounds in mature oak trees in a temperate forest at
506 four key phenological stages, *Trees-Structure and Function*, 27, 1023-1034,
507 10.1007/s00468-013-0853-5, 2013.

508 Bloor, J. M. G., Niboyet, A., Leadley, P. W., and Barthes, L.: CO₂ and inorganic N
509 supply modify competition for N between co-occurring grass plants, tree seedlings
510 and soil microorganisms, *Soil Biology and Biochemistry*, 41, 544-552,
511 <http://dx.doi.org/10.1016/j.soilbio.2008.12.013>, 2009.

512 Breda, N., and Granier, A.: Intra- and interannual variations of transpiration, leaf
513 area index and radial growth of a sessile oak stand (*Quercus petraea*), *Ann. Sci. For.*,
514 53, 521-536, 10.1051/forest:19960232, 1996.

515 Coleman, G. D., and Chen, T. H. H.: Sequence of a poplar bark storage protein gene,
516 *Plant Physiology*, 102, 1347-1348, 1993.

517 Cooke, J. E. K., and Weih, M.: Nitrogen storage and seasonal nitrogen cycling in
518 Populus: bridging molecular physiology and ecophysiology, *New Phytologist*, 167,
519 19-30, 2005.

520 Dickson, R. E.: Carbon and nitrogen allocation in trees, *Ann. Sci. For.*, 46, S631-
521 S647, 10.1051/forest:198905ART0142, 1989.

522 Dong, S., Cheng, L., Scagel, C. F., and Fuchigami, L. H.: Nitrogen absorption,
523 translocation and distribution from urea applied in autumn to leaves of young potted
524 apple (*Malus domestica*) trees, *Tree Physiology*, 22, 1305-1310,
525 10.1093/treephys/22.18.1305, 2002.

526 Dong, S. F., Cheng, L. L., and Fuchigami, L. H.: New root growth in relation to
527 nitrogen reserves of young Gala/M26 apple trees, 4th International Symposium on
528 Mineral Nutrition of Deciduous Fruit Crops, Penticton, Canada, 2000,
529 ISI:000173413200045, 365-369,

530 El Zein, R., Breda, N., Gerant, D., Zeller, B., and Maillard, P.: Nitrogen sources for
531 current-year shoot growth in 50-year-old sessile oak trees: an in situ N-15 labeling
532 approach, *Tree Physiology*, 31, 1390-1400, 10.1093/treephys/tpr118, 2011a.

533 El Zein, R., Maillard, P., Breda, N., Marchand, J., Montpied, P., and Gerant, D.:
534 Seasonal changes of C and N non-structural compounds in the stem sapwood of adult
535 sessile oak and beech trees, *Tree Physiology*, 31, 843-854, 10.1093/treephys/tpr074,
536 2011b.

537 Frak, E., Millard, P., Le Roux, X., Guillaumie, S., and Wendler, R.: Coupling sap
538 flow velocity and amino acid concentrations as an alternative method to N-15
539 labeling for quantifying nitrogen remobilization by walnut trees, *Plant Physiology*,
540 130, 1043-1053, 10.1104/pp.002139, 2002.

541 Gessler, A., Schneider, S., Weber, P., Hanemann, U., and Rennenberg, H.: Soluble N
542 compounds in trees exposed to high loads of N: a comparison between the roots of
543 Norway spruce (*Picea abies*) and beech (*Fagus sylvatica*) trees grown under field
544 conditions, *New Phytologist*, 138, 385-399, 1998a.

545 Gessler, A., Schultze, M., Schrempp, S., and Rennenberg, H.: Interaction of phloem-
546 translocated amino compounds with nitrate net uptake by the roots of beech (*Fagus*
547 *sylvatica*) seedlings, *Journal of Experimental Botany*, 49, 1529-1537, 1998b.

548 Gilson, A., Barthes, L., Delpierre, N., Dufrene, E., Fresneau, C., and Bazot, S.:
549 Seasonal changes in carbon and nitrogen compound concentrations in a *Quercus*
550 *petraea* chronosequence, *Tree Physiology*, 34, 716-729, 10.1093/treephys/tpu060,
551 2014.

552 [Gojon, A., Bussi, C., Grignon, C., and Salsac, L.: Distribution of NO₃⁻ reduction](#)
553 [between roots and shoots of peachtree seedlings as affected by NO₃⁻ uptake rate,](#)
554 [Physiologia Plantarum, 82, 505-512, 1991.](#)

555 Grassi, G., Millard, P., Gioacchini, P., and Tagliavini, M.: Recycling of nitrogen in
556 the xylem of *Prunus avium* trees starts when spring remobilization of internal
557 reserves declines, *Tree Physiology*, 23, 1061-1068, 2003.

558 Guak, S., Neilsen, D., Millard, P., Wendler, R., and Neilsen, G. H.: Determining the
559 role of N remobilization for growth of apple (*Malus domestica* Borkh.) trees by
560 measuring xylem-sap N flux, *Journal of Experimental Botany*, 54, 2121-2131,
561 10.1093/jxb/erg228|ISSN 0022-0957, 2003.

562 Hodge, A., Robinson, D., and Fitter, A.: Are microorganisms more effective than
563 plants at competing for nitrogen?, *Trends in Plant Science*, 5, 304-308,
564 [http://dx.doi.org/10.1016/S1360-1385\(00\)01656-3](http://dx.doi.org/10.1016/S1360-1385(00)01656-3), 2000.

565 Jordan, M.-O., Wendler, R., and Millard, P.: Autumnal N storage determines the
566 spring growth, N uptake and N internal cycling of young peach trees, *Trees*, 26, 393-
567 404, 10.1007/s00468-011-0600-8, 2012.

568 Kim, Y. K., Lim, C. S., Kang, S. M., and Cho, J. L.: Root storage of nitrogen applied
569 in autumn and its remobilization to new growth in spring of persimmon trees
570 (*Diospyros kaki* cv. Fuyu), *Sci. Hortic.*, 119, 193-196, 10.1016/j.scienta.2008.07.013,
571 2009.

572 Millard, P., and Proe, M. F.: Leaf demography and the seasonal internal cycling of
573 nitrogen in sycamore (*Acer pseudoplatanus* L) seedlings in relation to nitrogen
574 supply, *New Phytologist*, 117, 587-596, 10.1111/j.1469-8137.1991.tb00963.x, 1991.

575 Millard, P., and Proe, M. F.: Nitrogen uptake, partitioning and internal cycling in
576 *Picea-sitchensis* (Bong) carr as influenced by nitrogen supply, *New Phytologist*, 125,
577 113-119, 10.1111/j.1469-8137.1993.tb03869.x, 1993.

578 Millard, P.: Ecophysiology of the internal cycling of nitrogen for tree growth,
579 *Zeitschrift Fur Pflanzenernahrung Und Bodenkunde*, 159, 1-10, 1996.

580 Millard, P., Hester, A., Wendler, R., and Baillie, G.: Interspecific defoliation
581 responses of trees depend on sites of winter nitrogen storage, *Functional Ecology*, 15,
582 535-543, 10.1046/j.0269-8463.2001.00541.x, 2001.

583 Millard, P., Wendler, R., Grassi, G., Grelet, G.-A., and Tagliavini, M.: Translocation
584 of nitrogen in the xylem of field-grown cherry and poplar trees during
585 remobilization, *Tree Physiology*, 26, 527-536, 10.1093/treephys/26.4.527, 2006.

586 [Morot-Gaudry, J. F.: Assimilation de l'azote chez les plantes. Aspects physiologique,](#)
587 [biochimique et moléculaire, edited by: eds. I., INRA, Paris, 1997.](#)

588 Neilsen, D., Millard, P., Neilsen, G. H., and Hogue, E. J.: Sources of N for leaf
589 growth in a high-density apple (*Malus domestica*) orchard irrigated with ammonium
590 nitrate solution, *Tree Physiology*, 17, 733-739, 10.1093/treephys/17.11.733, 1997.

591 Neilsen, D., Millard, P., Herbert, L. C., Neilsen, G. H., Hogue, E. J., Parchomchuk,
592 P., and Zebarth, B. J.: Remobilization and uptake of N by newly planted apple
593 (*Malus domestica*) trees in response to irrigation method and timing of N application,
594 *Tree Physiology*, 21, 513-521, 2001.

595 Salaün, M., Guérin, V., Huché-Théliér, L., Charpentier, S., and Dily, F. L.: Nitrogen
596 storage and mobilisation for spring growth in *Ligustrum* cultivated in container, *Sci.*
597 *Hortic.*, 103, 461-471, <http://dx.doi.org/10.1016/j.scienta.2004.06.012>, 2005.

598 Sanchez, E. E., Righetti, T. L., Sugar, D., and Lombard, P. B.: Seasonal differences,
599 soil texture and uptake of newly absorbed nitrogen in field-grown pear trees, *Journal*
600 *of Horticultural Science*, 65, 395-400, 1990.

601 Stephens, D. W., Whitehead, D., Turnbull, M. H., and Millard, P.: Effects of nitrogen
602 supply on phenology and carbon assimilation in young *Nothofagus fusca*, *N. Z. J.*
603 *Bot.*, 39, 617-630, 2001.

604 Tagliavini, M., Quartieri, M., and Millard, P.: Remobilised nitrogen and root uptake
605 of nitrate for spring leaf growth, flowers and developing fruits of pear (*Pyrus*
606 *communis* L.) trees, *Plant Soil*, 195, 137-142, 1997.

607 Taylor, B.: Storage and mobilization of nitrogen in fruit trees: a review, *J. Aust. Inst.*
608 *Agric. Sci.*, 33, 23-29, 1967.

609 Tromp, J., and Ova, J. C.: Uptake and distribution of nitrogen in young apple trees
610 after application of nitrate or ammonium, with special reference to asparagine and
611 arginine, *Physiologia Plantarum*, 45, 23-28, 10.1111/j.1399-3054.1979.tb01657.x,
612 1979.

613 Tromp, J.: Nutrient reserves in roots of fruit trees, in particular carbohydrates and
614 nitrogen, *Plant Soil*, 71, 401-413, 1983.

615 Ueda, M., Mizumachi, E., and Tokuchi, N.: Winter nitrate uptake by the temperate
616 deciduous tree *Quercus serrata*, *J For Res*, 15, 411-414, 10.1007/s10310-010-0219-4,
617 2010.

618 Ueda, M. U., Mizumachi, E., and Tokuchi, N.: Allocation of nitrogen within the
619 crown during leaf expansion in *Quercus serrata* saplings, *Tree Physiology*, 29, 913-
620 919, 10.1093/treephys/tpp029, 2009.

621 Valenzuela Nunez, L., Gerant, D., Maillard, P., Breda, N., Gonzalez Cervantes, G.,
622 and Sanchez Cohen, I.: Evidence for 26KDa vegetative storage protein in the stem
623 sap wood of mature pedunculate oak, *Intersciencia*, 36, 142-147, 2011.

624 Vance, E. D., Brookes, P. C., and Jenkinson, D. S.: Microbial biomass measurements
625 in forest soils - determination of Kc values and tests of hypotheses to explain the
626 failure of the chloroform fumigation incubation method in acid soils, *Soil Biology &*
627 *Biochemistry*, 19, 689-696, 10.1016/0038-0717(87)90050-2, 1987.

628 Vitasse, Y., Delzon, S., Dufrene, E., Pontailler, J. Y., Louvet, J. M., Kremer, A., and
629 Michalet, R.: Leaf phenology sensitivity to temperature in European trees: Do
630 within-species populations exhibit similar responses?, *Agricultural and Forest*
631 *Meteorology*, 149, 735-744, 10.1016/j.agrformet.2008.10.019, 2009.

632 Vizoso, S., Gerant, D., Guehl, J. M., Joffre, R., Chalot, M., Gross, P., and Maillard,
633 P.: Do elevation of CO₂ concentration and nitrogen fertilization alter storage and
634 remobilization of carbon and nitrogen in pedunculate oak saplings?, *Tree*
635 *Physiology*, 28, 1729-1739, 10.1093/treephys/28.11.1729, 2008.

636 Weinbaum SA, K. I., Broadbent FE, Micke WC, Muraoka TT.: Effects of time of
637 nitrogen application and soil texture on the availability of isotopically labelled

638 fertilizer nitrogen to reproductive and vegetative growth of mature almond trees.,
639 Journal of the American Society of Horticultural Science, 109, 339-343, 1984.
640 Wendler, R., and Millard, P.: Impacts of water and nitrogen supplies on the
641 physiology, leaf demography and nitrogen dynamics of *Betula pendula*, Tree
642 Physiology, 16, 153-159, 1996.

643

644 |

645 |

646 **Table**

647 Table 1 :

648 Labelling characteristics and recovery of ¹⁵N administered in each labelling

649 campaign from the sampled compartments of each tree, on each sampling occasion

650 (DAL : Days after Labelling, JD :Julian day number).

Tree	1	2	3	4	5	6	
Labelling date	2009/05/27	2009/05/27	2009/09/09	2009/09/09	2010/03/20	2010/03/20	
DAL/JD	% of recovered ¹⁵ N		DAL/JD	% of recovered ¹⁵ N		DAL/JD	% of recovered ¹⁵ N
Year 1	<u>1/148</u>	39	25	<u>3/255</u>	68	72	
	<u>3/150</u>	31	25	<u>6/258</u>	68	50	
	<u>6/153</u>	30	24	<u>9/261</u>	68	70	
	<u>9/156</u>	22	19	<u>16/268</u>	33	38	
	<u>16/163</u>	19	16	<u>28/280</u>	31	22	
	<u>30/177</u>	17	15	<u>49/301</u>	29	15	
	<u>57/205</u>	17	14	<u>84/336</u>	<u>29</u>	<u>14</u>	
	<u>126/273</u>	15	14				
Year 2	<u>189/336</u>	<u>14</u>	<u>13</u>				
	<u>318/98</u>	8	7	<u>208/98</u>	24	14	<u>20/98</u>
	<u>337/118</u>	11	13	<u>227/118</u>	12	10	<u>40/118</u>
	<u>358/139</u>	10	13	<u>247/139</u>	16	20	<u>61/139</u>
	<u>370/152</u>	14	14	<u>260/152</u>	22	21	<u>74/152</u>
	<u>397/180</u>	11	10	<u>287/180</u>	38	18	<u>102/180</u>
	<u>460/244</u>	13	11	<u>350/244</u>	13	12	<u>166/244</u>
	<u>509/293</u>	7	5	<u>399/293</u>	10	8	<u>215/293</u>
						11	
						21	

651

652

661 **Figure captions**

662 Figure 1 : Partitioning of recovered ¹⁵N (PRN%) from the sampled compartments
663 following the first labelling campaign, i.e. from May 26, 2009 to October 20, 2010.

664 a. leaves \blacklozenge and twigs \times , b. phloem \blacktriangle , c. fine roots \circ , d. rhizospheric soil \blacksquare and
665 microbial biomass $+$ (for those compartments the Y axis was adjusted to 1). DAL:
666 Days after labelling. The two lines, continuous and dotted, correspond to tree 1 and
667 tree 2. Vertical bars indicate standard errors.

- Supprimé: .
- Supprimé: .
- Supprimé: .
- Supprimé: .
- Supprimé: for each categories (
- Supprimé: dashed)
- Supprimé: the
- Supprimé: the

669 Figure 2 : Partitioning of recovered ¹⁵N (PRN%) from the sampled compartments
670 following the second labelling campaign, i.e. from September 08, 2009 to October

671 20, 2010; a. leaves \blacklozenge and twigs \times , b. phloem \blacktriangle , c. fine roots \circ , d. rhizospheric soil \blacksquare
672 and microbial biomass $+$. DAL: Days after labelling. The two lines for each category
673 (continuous and dotted) correspond to tree 3 and tree 4. Vertical bars indicate
674 standard errors.

- Supprimé: .
- Supprimé: and xylem,
- Supprimé: .
- Supprimé:
- Supprimé: .
- Supprimé: categories
- Supprimé: dashed
- Supprimé: the
- Supprimé: the

676 Figure 3 : Partitioning of recovered ¹⁵N (PRN%) from the sampled compartments
677 following the third labelling campaign, i.e. from April 8, 2010 to October 20, 2010;

678 a. leaves \blacklozenge and twigs \times , b. phloem \blacktriangle , c. fine roots \circ , d. rhizospheric soil \blacksquare and
679 microbial biomass $+$. DAL: Days after labelling. The two lines for each category
680 (continuous and dotted) correspond to tree 5 and tree 6. Vertical bars indicate
681 standard errors.

- Supprimé:
- Supprimé: .
- Supprimé: and xylem,
- Supprimé: .
- Supprimé: .
- Supprimé: .
- Supprimé: categories
- Supprimé: dashed
- Supprimé: the
- Supprimé: the

683 Figure 4 : Conceptual scheme representing percentage contributions of ¹⁵N (Eq. 2)
684 from each labelling campaign (L₁: white, L₂: light grey, L₃: dark grey) in roots in the
685 autumn, and in new leaves in the season following the first labelling campaign.

Figure 3:

Supprimé: <sp><sp>




Figure 4 :

