

1 **Title Page**

2 **TITLE:**

3 Flower litters of alpine plants affect soil nitrogen and phosphorus rapidly in the eastern Tibetan Plateau

4  
5 **AUTHORS:**

6 Jinniu Wang<sup>1,2</sup>

7 1. Chengdu Institute of Biology, Chinese Academy of Sciences/Key Laboratory of Mountain  
8 Ecological Restoration and Bioresource Utilization, Chinese Academy of Sciences/Ecological  
9 Restoration Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu 610041, China

10 2. International Centre for Integrated Mountain Development (ICIMOD), G.P.O. Box 3226,  
11 Kathmandu, Nepal

12  
13 Bo Xu<sup>1,3</sup>

14 1. Chengdu Institute of Biology, Chinese Academy of Sciences/Key Laboratory of Mountain  
15 Ecological Restoration and Bioresource Utilization, Chinese Academy of Sciences/Ecological  
16 Restoration Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu 610041, China

17 3. University of Chinese Academy Sciences, Beijing 100049, China

18  
19 Yan Wu<sup>1</sup>

20 1. Chengdu Institute of Biology, Chinese Academy of Sciences/Key Laboratory of Mountain  
21 Ecological Restoration and Bioresource Utilization, Chinese Academy of Sciences/Ecological  
22 Restoration Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu 610041, China

23  
24 Jing Gao<sup>1,3</sup>

25 1. Chengdu Institute of Biology, Chinese Academy of Sciences/Key Laboratory of Mountain  
26 Ecological Restoration and Bioresource Utilization, Chinese Academy of Sciences/Ecological  
27 Restoration Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu 610041, China

28 3. University of Chinese Academy Sciences, Beijing 100049, China

29  
30 Fusun Shi (corresponding author)

31 Chengdu Institute of Biology, Chinese Academy of Sciences/Key Laboratory of Mountain Ecological  
32 Restoration and Bioresource Utilization, Chinese Academy of Sciences/Ecological Restoration  
33 Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu 610041, China

34 Email: [shifs@cib.ac.cn](mailto:shifs@cib.ac.cn)

35 Tel: +86-28-82890537

36 Fax: +86-28-82890288

37

38 **Flower Litters of Alpine Plants Rapidly Affect Soil Nitrogen and**  
39 **Phosphorus in the Eastern Tibetan Plateau**

40 **Abstract**

41 Litters of reproductive organs have been rarely studied despite their role in allocating  
42 nutrients for offspring reproduction. This study determines the mechanism through  
43 which flower litters efficiently increase the available soil nutrient pool. Field  
44 experiments were conducted to collect plant litters and calculate biomass production  
45 in an alpine meadow of the eastern Tibetan Plateau. C, N, P, lignin, cellulose content,  
46 and their relevant ratios of litters were analyzed to identify their decomposition  
47 features. A pot experiment was performed to determine the effects of litter addition on  
48 soil nutrition pool by comparing the treated and control samples. Litter-bag method  
49 was used to verify decomposition rates. The flower litters of phanerophyte plants  
50 were comparable with non-flower litters. Biomass partitioning of other herbaceous  
51 species accounted for 10%–40% of the aboveground biomass. Flower litter possessed  
52 significantly higher N and P levels but less C/N, N/P, lignin/N, and lignin and  
53 cellulose concentrations than leaf litter. Litter-bag experiment confirmed that the  
54 flower litters of *Rhododendron przewalskii* and *Meconopsis integrifolia* decomposes  
55 approximately three times faster than mixed litters within 50 days. Pot experiment  
56 findings indicated that flower litter addition significantly increased the available  
57 nutrient pool and soil microbial productivity. The time of litter fall significantly  
58 influenced soil available N and P, and soil microbial biomass. Flower litters fed soil  
59 nutrition pool and influenced nutrition cycling in alpine ecosystems more efficiently  
60 because of their non-ignorable production, faster decomposition rate and higher  
61 nutrient contents compared with non-flower litters. The underlying mechanism can  
62 enrich nutrients, which return to the soil, and non-structural carbohydrates, which feed  
63 and enhance the transitions of soil microorganisms.

64 **Key words** alpine ecosystem, flower litter, chemical property, decomposition rate,  
65 nitrogen, phosphorus

66

67 Plant properties directly affect the productivity and function of an ecosystem in a  
68 natural environment (Chapin et al., 1986; Chapin, 2003; Berendse and Aerts, 1987;  
69 Grime, 1998). Plants continuously lose N and P in their entire life history and even  
70 during litter production and decomposition (Laungani and Knops, 2009; Richardson  
71 et al., 2009). In cold environments, litter tends to be recalcitrant (Aerts, 1997), but  
72 reproductive tissues present chemical composition that differs from vegetative parts,  
73 resulting in a markedly faster decomposition and nutrient release, with repercussions  
74 on nutrient cycling and patchiness (Buxton and Marten, 1989; Lee et al., 2011).  
75 Although inflorescences comprise only a small fraction of plant biomass and  
76 production in Arctic and alpine vegetation, the inflorescence production can be a  
77 significant proportion of the total production of species under certain special  
78 circumstances (Martínez-Yrizar et al., 1999, Fabbro and Körner, 2004; Wookey et al.,  
79 2009). High contents of N and P exist in the reproductive organs of plants probably  
80 because of their essential roles in plant growth and formation (e.g., high protein  
81 content). The rate of decay and concentrations of nutrients in the litter determine the  
82 rate of nutrient release, which creates a positive feedback to site fertility. Hence, the  
83 chemical properties of litters from different plant organs and their correlations with  
84 decomposition rate must be determined.

85 The growth and health of plants in their life history have been considerably influenced  
86 by variations in the physical, chemical, and biological properties of soil, particularly  
87 around the rhizosphere. Nonetheless, soil properties can also be mediated by plants. N  
88 is a major constituent of several important plant substances (Vitousek and Howarth,  
89 1991). In cold life zone ecosystems, plant biomass production is limited by N  
90 (Körner, 2003). The plant residue is one principal component of soil organic matter  
91 (SOM), whose decomposition can supply available N to plants and microorganisms.  
92 Similar to N, P is closely associated with numerous vital plant processes.  
93 Nevertheless, in most circumstances, P is limited because of its small concentration in  
94 soil; this element is released slowly from insoluble P but is highly demanded by plants  
95 and microorganisms (Bieleski, 1973; Richardson et al., 2009). As decomposition is a  
96 prolonged process, plants contain concentrated nutrients comparable with soil, which

97 enhance the microbial immobilization of N when they provide C to soil  
98 microorganisms. The nature of litter determines its palatability to soil organisms,  
99 thereby influencing their composition and activity levels. Litter can also mediate the  
100 interactions between neighboring plants in infertile communities (Nilsson et al., 1999,  
101 Xiong and Nilsson, 1999), which significantly affect the biogeochemical cycle and  
102 feedback of plant–soil interaction.

103 Fast decay of N-rich litters suggests that litter decay rates increase with increasing N  
104 content. The initial rate of nutrient release is positively correlated with the initial  
105 concentrations of N or P (MacLean and Wein, 1978; Aber and Melillo, 1980; Berg  
106 and Ekbohm, 1983; Yavitt and Fahey, 1986; Stohlgren, 1988). Long-term increases in  
107 N availability have also been reported following the additions of C to forests  
108 (Groffman, 1999). In agricultural systems, addition of fresh residues can stimulate the  
109 decomposition and net release of N from indigenous SOM (Haynes, 1986; Scott et al.,  
110 1996). Recently, a common-garden decomposition experiment in a wide range of  
111 subarctic plant types demonstrated that structural and chemical traits are better  
112 predictors for several high-turnover organs than structural traits alone (Freschet et al.,  
113 2012). Decomposition rate of plant litters slightly differ because of their species-  
114 specific traits and various organs, whose chemical qualities vary in a wide range of  
115 plant types and environments.

116 Alpine ecosystems are thermally restricted and characterized by a low material  
117 turnover rate (Körner, 2003). In a high altitude region, plants grow in a harsh habitat  
118 that restricted their effective utilization of resources; in this regard, the total available  
119 resource is less compared with that of plants in other regions (Fabbro and Körner,  
120 2004; Hautier et al., 2009). In long-term evolution, the allocation of accumulated  
121 carbohydrates to reproduction is an adaptation strategy, leading to the partitioning of  
122 reproductive organs, that is, the availability and timely mobilization of adequate  
123 resources from the vegetative plant body to reproductive structures (Arroyo et al.,  
124 2013). Thus far, probably due to reproductive organs' comparatively minor biomass  
125 production and difficulty to be collected, studies on their decomposition have been  
126 limited particularly compared with those on leaf and other vegetative organs. In this

127 study, we conducted comprehensive field investigation, pot experiment of litter  
128 addition, and litter-bag experiment to address the following questions:

129 1) Should flower litter be considered in the alpine ecosystem's biogeochemical cycles  
130 for their relatively innegligible biomass production and/or allocation?

131 2) Does flower litter of higher quality and with unique traits have faster  
132 decomposition than leaf litter?

133 3) Does the time of litter fall influence soil available nutrients and soil microbial  
134 productivity of alpine meadow ecosystem?

135

## 136 **Materials and Methods**

### 137 *Study area*

138 The field site is located at the foot of Mt. KaKa, which belongs to the middle  
139 section of Minshan Mountain, eastern Tibetan Plateau (**Fig. 1**), with a mean annual  
140 precipitation of 720 mm. More than 70% of precipitation falls in summer from June to  
141 August. Snowfall usually occurs from the end of September to early May next year.  
142 Vegetation presents a typical alpine meadow with numerous and unique alpine plants.  
143 Mosses are abundant and cover most of the ground. The moss layer is dominated by  
144 *Polytrichum swartzii* and *Trematodon acutus* c. mull. Vascular plants include species  
145 mainly belonging to genus *Kobresia* and genus *Carex*. Other common species are  
146 *Festuca* spp., *Gentiana* spp., and *Leontopodium* spp. Plant roots in this ecosystem are  
147 generally confined to the surface A-horizon (2–20 cm). A few dwarf shrubs are  
148 scattered sporadically in the meadow, e.g., *Rhododendron* and *Salix*. The soil type is  
149 dominated by Mat Cry-gelic Cambisols (i.e., silty loam inceptisol, *Chinese Soil*  
150 *Taxonomy Research Group*, 1995).

### 151 *Plant litter sampling*

152 During the blooming period from the end of May until mid-June and from the end of  
153 July until early August, flower litters of 14 earlier flowering plants species and 15  
154 later flowering plants species were carefully collected in 2012 at two sites, namely,  
155 Mt. Kaka (103°42' E; 32°59' N, 3500–3900 m a.s.l.) and Bow Ridge Mountain  
156 (103°42' E; 33°1' N, 3600–3850 m a.s.l.). In the study, 4 litter traps were placed under

157 the crown of each individual shrub in different communities (5–8 individuals were  
158 chosen for the placement of litter traps), which were processed and modified based on  
159 the litterfall monitoring protocol (Muller-Landau and Wright, 2010). The litter trap  
160 was composed of 1 cloth bag and 4 support legs. Window screen (with a mesh size of  
161 0.8 mm) was used to seize the cloth bag. Its size was about 50 cm deep and 25 cm  
162 long. Four legs (made with 80 cm PVC pipe) were tied with a cloth bag and frame.  
163 The frame of the opening was made of iron wire with 3 mm diameter. After inserting  
164 it into the soil under the shrub's crown, the plant litter was collected twice per week,  
165 which was later sorted as flower litter and other types during the blooming period.  
166 Given the small size of herbaceous individuals, flowers were plucked at the end of the  
167 flowering phase, and their mass ratios to aboveground biomass were calculated.  
168 Freshly fallen leaves of different species were collected from the floor of the alpine  
169 meadow (i.e., mixed leaf litters, ca. 3950 m a.s.l.). These species were tentatively  
170 classified into five groups according to Raunkiaer's life-form system (i.e.,  
171 chamephyte, geophyte, hemicryptophyte, phanerophyte, and therophyte). Target  
172 species were first decided by visual observation. For herbaceous species, their  
173 dominances were determined using quadrat methods. Each quadrat (1 m × 1 m) was  
174 spaced at least 2 m apart from each other along the transect for recording community  
175 composition (totaling 10 quadrats along one transect, and three transects at each site).  
176 Weighted means of frequency and biomass of target species were sorted and used to  
177 assess their dominances. For shrubs, line-point intercept method was conducted to  
178 calculate targeted species' frequency, height, and cover, which are represented by  
179 "hit" (three transects at each site; a 20 m rope with ca. 1 cm diameter or a measuring  
180 tape was used), whose weighted means were sorted to determine the dominant species  
181 (Herrick et al., 2005). We also consulted an expert who has prior knowledge or  
182 research on the dominant species at the selected sites.

183 These species were divided into earlier flowering species and later flowering species  
184 groups based on blooming time (Table 1). According to Raunkiaer's life-form system,  
185 earlier flowering species mainly consisted of hemicryptophyte, geophyte, and  
186 phanerophyte, whereas more than half of later flowering species comprised

187 chamaephyte. Nearly half of the tested species were dominant or co-dominant in their  
188 respective communities. The dry matter content of flower litters in all of the species  
189 was ranked from 10% to 60%. Mixed leaf litters of alpine meadows were sampled on  
190 Mt. Kaka (3950 m. a.s.l.), and leaf litters of 13 dominant species were collected to  
191 compare their chemical properties with flower litters. Both types of litters were first  
192 spread on blotting paper for air drying. A small portion of each litter was further dried  
193 in an oven for 48 h to calculate dry matter content.

#### 194 *Experimental design*

195 Polyvinyl chloride (PVC) pots (15 cm deep, 20 cm diameter at the top, and 12 cm  
196 diameter at the bottom) were filled with 2 kg of soils, which were collected in autumn  
197 of 2011. The collected soil samples were stored at 4 °C. The samples were sieved  
198 through 2 mm mesh and then mixed thoroughly. The soil surface of each treatment  
199 was added with 5 g of flower litters or mixed litters (calculated as dry weight) on June  
200 21 (14 species, earlier flowering plants) and August 11, 2012 (15 species, later  
201 flowering plants). The surface was covered with a thin layer of soil to avoid being  
202 blown by wind. Other two additional treatments were conducted without litter  
203 addition (control) and with mixed leaf litter addition, respectively. In total, the pot  
204 experiment consisted of 33 treatments with 3 replicates, with a total number of 99  
205 pots. All of the pots were carefully buried 12 cm deep into the field to maintain the  
206 same soil temperature in the experimental field. The pots were randomly distributed,  
207 and their top edges were approximately 3 cm above the ground to prevent runoff from  
208 outside. All of the pots were rearranged every week to create a similar microclimate.  
209 After 50 days, each soil sample was collected from three points of each pot in the  
210 center and then mixed to avoid the boundary layer effect. Each soil sample from  
211 different PVC pots was mixed evenly by sieving through a 2 mm mesh. The samples  
212 were stored and marked separately in an ice box prior to chemical determination.

213

214

215 Table 1 General description of flower litters.

|                                 | Life form | Size of inflorescence (cm) | Dominant (Y/N) | Color      | Dry matter content (%) |
|---------------------------------|-----------|----------------------------|----------------|------------|------------------------|
| <i>Caragana jubata</i>          | C         | 1–1.5                      | N              | white      | 29.81                  |
| <i>Primula orbicularis</i>      | H         | 1.5                        | Y              | yellow     | 23.29                  |
| <i>Potentilla anserina</i>      | G         | 1–1.8                      | Y              | yellow     | 51.9                   |
| <i>Rhododendron capitatum</i>   | P         | 2–3                        | Y              | purple     | 32.84                  |
| <i>Viola rockiana</i>           | H         | 1                          | N              | yellow     | 25.22                  |
| <i>Myricaria squamosa</i>       | P         | 0.5–1                      | N              | pink       | 30.95                  |
| <i>Potentilla saundersiana</i>  | G         | 1–1.4                      | N              | yellow     | 54.01                  |
| <i>Taraxacum lugubre</i>        | H         | 3–4                        | Y              | yellow     | 14.97                  |
| <i>Aster tongolensis</i>        | H         | 4–5                        | N              | blue       | 28.72                  |
| <i>Cardamine tangutorum</i>     | G         | 0.8–1.5                    | N              | lavender   | 13.08                  |
| <i>Spiraea alpina</i>           | P         | 0.5–0.7                    | Y              | fallow     | 32.58                  |
| <i>Caltha scaposa</i>           | H         | 3–4                        | Y              | yellow     | 30.43                  |
| <i>Rhododendron przewalskii</i> | P         | 4–5                        | Y              | pink       | 33.33                  |
| <i>Meconopsis integrifolia</i>  | H/T       | 5–7                        | N              | yellow     | 21.79                  |
| <i>Stellera chamaejasme</i>     | C         | 0.5                        | N              | red        | 28.11                  |
| <i>Potentilla fruticosa</i>     | P         | 2–3                        | Y              | yellow     | 30.43                  |
| <i>Meconopsis punicea</i>       | H/A       | 5–8                        | N              | red        | 33.57                  |
| <i>Meconopsis violacea</i>      | H         | 4–6                        | N              | purple     | 35.70                  |
| <i>Sibiraea angustata</i>       | P         | 0.8                        | Y              | white      | 29.50                  |
| <i>Polygonum macrophyllum</i>   | H         | 0.2                        | Y              | pink       | 21.79                  |
| <i>Pedicularis megalochila</i>  | C         | 0.8–1                      | N              | red        | 33.57                  |
| <i>Ligularia virgaurea</i>      | C         | 1.5                        | N              | yellow     | 16.78                  |
| <i>Pilose Asiabell</i>          | C         | 2–2.5                      | N              | pale/green | 22.26                  |
| <i>Oxytropis ochrocephala</i>   | C         | 1                          | N              | fallow     | 28.72                  |
| <i>Pedicularis longiflora</i>   | C         | 0.8                        | N              | yellow     | 28.11                  |
| <i>Hedysarum vicioides</i>      | C         | 1                          | N              | pink       | 30.02                  |
| <i>Gentiana sino-ornata</i>     | C         | 3–5                        | Y              | purple     | 44.10                  |
| <i>Leontopodium sinense</i>     | C         | 0.2–0.5                    | Y              | white      | 56.92                  |
| <i>Cremanthodium lineare</i>    | G         | 1.2–1.7                    | Y              | yellow     | 48.93                  |

216 Note: C, H, G, P, and T represent chamaephyte, hemicryptophyte, geophyte (one of the subdivided  
 217 groups in cryptophytes), phanerophyte, and thermophile, respectively. Y and N indicate whether the  
 218 species is dominant or not in the community. The first 14 species are earlier flowering species, and the  
 219 other 15 species are later flowering species.

## 220 *Decomposition rate*

221 A litter bag with a size of 14 cm × 20 cm was used to determine the  
 222 decomposition rate of different plant litters. The bag was double faced and made from  
 223 nylon net material with above (4.5 mm × 4.5 mm mesh) and below layers (0.8 mm ×  
 224 0.8 mm mesh). The above layer with larger mesh size allowed free access for most  
 225 micro-arthropods, which dominate the soil fauna of alpine meadow in the eastern  
 226 Tibetan Plateau, whereas the below layer with smaller mesh size can reduce litter



227 spillage from the litter bags in the process. As representative species, flower litters of  
228 *Rhododendron. przewalskii* and *Meconopsis. integrifolia* and mixed litter were packed  
229 into litter bags with the edges sealed on June 21, 2012. The litterbag experiment was  
230 conducted to compare the decomposition rate of flower litters and mixed litter. Each  
231 treatment had eight replicates. After 7 weeks (August 8, 2012), the debris or mud was  
232 remove outside the litter bags carefully, then litters were taken outside, sank into  
233 small water basin for a short time, and sorted out clay and litter through 0.5 mm mesh  
234 filter. Lastly, remaining litters were dried in an oven for 48 hours (65 °C) and  
235 measured the weight on the balance (accuracy 0.001 g) for decomposition calculation.  
236 Litter decomposition rates can be determined by the following equation.

$$237 \quad \mathbf{DR = (P-R)/P \times 100}$$

238 where *DR* is the decomposition rate, *P* is primary litter mass in the litter bags, and *R*  
239 refers to residue litter before determining percentage mass loss.

240

#### 241 *Chemistry determination of soil and plant*

242 For soil samples, total dissolved N (TN) contents were determined using unsieved  
243 fresh moist soil subsamples. Soil subsamples were extracted using 2 M KCl and  
244 shaken for 1 h at room temperature (20 °C), with a soil-to-solution ratio of 1:5  
245 (weight/volume). The extracted solution was filtered through filter paper before  
246 further determination (Jones et al., 2004).  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N were analyzed with the  
247 indophenol blue colorimetric (Sah, 1994) and ultraviolet spectrophotometry methods  
248 (Norman et al., 1985), respectively. Dissolved organic nitrogen (DON) was calculated  
249 by subtracting dissolved inorganic N ( $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N) from TN. Soil solutions  
250 were extracted by centrifugal drainage, whereas the exchangeable pool was extracted  
251 with 2 M KCl by using the methods reported by Jones et al. (2004). Total phosphorus  
252 (TP) consists of phosphorus mineral and organic phosphorous compound in the soil,  
253 which can be converted into the dissolved orthophosphate. Available phosphorous (A-  
254 P) is the fragments in soil that can be absorbed by plants, which consist of water-  
255 soluble phosphorus, some adsorbed phosphorus, organic phosphorus, and precipitated  
256 phosphorus in certain soil types. Chemically, A-P is defined as the phosphorus and

257 phosphate in soil solution that can be isotope exchanged with  $^{32}\text{P}$  or can be easily  
258 extracted by some chemical reagents. TP and A-P in soils were estimated by  
259 extraction with 0.5 M sodium hydroxide sodium carbonate solution (Dalal, 1973).  
260 Microbial biomass carbon (MBC) and microbial biomass nitrogen (MBN) contents  
261 were determined through the chloroform–fumigation direct-extraction technique.  
262 Correction factors of 0.54 for N and 0.45 for C were used to convert the chloroform  
263 labile N and C to microbial N and C (Brookes et al., 1985). For plant samples, the  
264 contents of C and N were determined by dry combustion with a CHNS auto-analyzer  
265 system (Elementar Analysen Systeme, Hanau, Germany) (Brodowski et al., 2006).  
266 The content of P was obtained colorimetrically by the chloro molybdophosphoric blue  
267 color method after wet digestion in a mixture of  $\text{HNO}_3$ ,  $\text{H}_2\text{SO}_4$ , and  $\text{HClO}_4$  solution  
268 (Institute of Soil Academia Sinica, 1978). Lignin and cellulose were estimated by the  
269 method described by Melillo et al. (1989).

#### 270 ***Data analysis***

271 One-way ANOVA was applied to compare values between the treatments and the  
272 control. Post-hoc multiple comparisons were adopted when the groups were three or  
273 more. Multivariate ANOVA was conducted to determine the effects of blooming time  
274 and different addition of litters and their interactions. To simplify the comparison of  
275 soil N and P between control (without flower litter) and the treated group (with flower  
276 litter), we defined an index  $\alpha$  as  $\alpha = \text{Ln}(N_2/N_1)$ .  $\alpha > 0$ ,  $N_2 > N_1$ ;  $\alpha < 0$ ,  $N_2 < N_1$ ;  $\alpha = 0$ ,  
277  $N_2 = N_1$ .  $N_1$  is the control treatment without flower litter, and  $N_2$  indicated the  
278 nutrition value (N or P) of flower litter treatment. Descriptive analysis was operated to  
279 demonstrate the  $\alpha$  values of different N and P fragments in various species litter  
280 addition treatment. The box plots provide the distribution of the values by the medians  
281 (central line), the 25% and 75% quartiles (box), and the ranges (whiskers). Asterisks  
282 (\*) represent the distribution of extreme outliers. The values (mean,  $n=X$ ) are also  
283 stated by one-way ANOVA. For comparison of three or more groups, mean  
284 differences were tested at  $P < 0.05$  by using Tukey multiple range test in SPSS 19.0  
285 software package (SPSS Inc., Chicago, IL, USA). The normality of data was tested  
286 with one-sample K-S test and Q-Q plot. Otherwise, log-transformation was adopted to

287 meet the normality requirement. Homogeneity of variance test was also utilized  
288 during the analysis. In the figures and tables, information is presented as means and  
289 standard errors of means. All of the differences were tested at the  $P = 0.05$  level.

## 290 **Results**

### 291 **Flower litter production of dominant species and their biomass allocation**

292 Among 13 dominant species, the flower litters of phanerophyte plants, whose  
293 flower litters are comparable with non-flower litters, were calculated through  
294 comparison with non-flower litters in the process of flower litter collection (**Fig. 2**  
295 **(a)**). The dry weights of flower litters were 10–40  $\text{g m}^{-2}$ , whereas their non-flower  
296 litters were only 5–25  $\text{g m}^{-2}$ . Although neither of the flower litters of *S. angustata* nor  
297 *R. capitatum* were significantly different compared with their non-flower litters ( $P >$   
298 0.05), the difference between the two remained noticeable, whose values were  $28.03 \pm$   
299  $3.56 \text{ g m}^{-2}$  versus  $13.21 \pm 1.49 \text{ g m}^{-2}$  for *R. capitatum* and  $19.58 \pm 3.50 \text{ g m}^{-2}$  versus  
300  $12.95 \pm 0.61 \text{ g m}^{-2}$  for *S. angustata*. The production of flower litters was higher than  
301 that of non-flower litters. The other three species significantly produced more flower  
302 litters than non-flower litters (*R. przewalskii*:  $F = 15.76$ ,  $P < 0.001$ ; *P. fruticosa*:  $F =$   
303  $4.76$ ,  $P < 0.05$ ; *S. alpine*:  $F = 10.18$ ,  $P < 0.01$ ). The flower litters of the eight  
304 herbaceous species were compared with their individual aboveground biomass (**Fig. 2**  
305 **(b)**), which ranked from 10% to nearly 40%. This finding indicated that flower litter  
306 should be considered to determine the effect of plants on soil nutrition pool during the  
307 growing season.

### 308 **Comparison of chemical properties between flower and leaf litters**

309 Total C content was not significantly different between flower and leaf litters (**Fig. 3**  
310 **(a)**,  $F = 1.80$ ,  $P = 0.199$ ). However, the levels of cellulose, lignin, and structure C of  
311 leaf litter were significantly higher than those of flower litter ( $F = 6.74$ ,  $P < 0.05$ ;  $F =$   
312  $5.77$ ,  $P < 0.05$ ;  $F = 10.99$ ,  $P < 0.01$ ). Hence, flower litter probably contains more non-  
313 structure C than leaf litter.

314 Both N and P contents of flower litters were significantly higher than those of leaf  
315 litters (**Fig. 3 (b)**). N in flower litters was nearly doubled to that of leaf litter ( $23.17 \pm$   
316  $1.52$ ,  $11.87 \pm 0.77$ ;  $F = 45.70$ ,  $P < 0.001$ ). More than twice the amount of P were also

317 present in flower litters ( $2.95 \pm 0.25 \text{ g kg}^{-1}$ ) compared with that in leaf litters ( $1.12 \pm$   
318  $0.12 \text{ g kg}^{-1}$ ;  $F = 43.87$ ,  $P < 0.001$ ).

319 For the implication of the ratio of different chemical properties, C/N, N/P, and  
320 lignin/N were determined to compare flower and leaf litters. All the three indicators of  
321 leaf litter were significantly higher than those of flower litters (**Fig. 3 (c)**). As  
322 parameters used to demonstrate decomposition rate, C/N and lignin/N of leaf litter  
323 were nearly double to those of flower litter ( $39.27 \pm 4.16$ ,  $19.80 \pm 1.39$ ,  $F = 37.78$ ,  $P$   
324  $< 0.001$ ;  $21.09 \pm 2.25$ ,  $12.79 \pm 1.15$ ,  $F = 7.91$ ,  $P < 0.01$ ). Furthermore, the N/P of  
325 flower litter was significantly higher than that of leaf litter ( $8.42 \pm 0.42$ ,  $11.60 \pm 0.56$ ;  
326  $F = 20.62$ ,  $P < 0.001$ ). These findings indicated that flower litter can supply more P  
327 per unit N than leaf litter.

#### 328 **Assessing the effects of flower litter on soil N pool and P pool**

329 Earlier flowering species exerted positive effects on soil TN,  $\text{NO}_3^-$ -N, and  $\text{NH}_4^+$ -N  
330 (**Fig. 4 (a)**), with the addition of their flower litters according to their size of  $\alpha$  values.  
331 Most parameters were higher than 0, which indicated that  $N_2 > N_1$ . Flower litter  
332 increased the soil N pool. All of the minimum  $\alpha$  values of the five indices were also  
333 higher than 0 (**Table 2**, 0.42–1.29), which indicated that flower litter addition  
334 significantly increased soil N pool including different fragments ( $P < 0.001$ ). Among  
335 the later flowering species, except *G. sino-ornata* and *L. sinense*, soil N indices were  
336 significantly improved with flower litter addition, as demonstrated through  $\alpha$  values  
337 higher than 0 (**Fig. 4 (b)**, **Table 2**). Later flowering species differed from earlier  
338 flowering species, with minimum  $\alpha$  values lower than 0, which resulted from the  
339 exceptions of *G. sino-ornata* and *L. sinense*. However, all of the mean  $\alpha$  values were  
340 higher than 0, which presented general results after flower litter addition (0.36–1.49);  
341 soil N pool was significantly enhanced only after 50 days ( $P < 0.001$ ). Interactions  
342 between flowering time and litter addition for  $\text{NO}_3^-$ -N and  $\text{NH}_4^+$ -N were significant ( $F$   
343  $= 5.043$ ,  $P < 0.05$ ;  $F = 7.947$ ,  $P < 0.01$ ;  $F = 24.143$ ,  $P < 0.05$ , respectively) but not for  
344 TN ( $F = 0.470$ ,  $P = 0.496$ ). Different flowering times significantly affected  $\text{NO}_3^-$ -N,  
345 and  $\text{NH}_4^+$ -N (**Table 3**,  $P < 0.01$ ) but did not significantly influence TN ( $F = 2.80$ ,  $P =$   
346  $0.10$ ). As illustrated in **Fig. 4**, litter addition had significant effects on all of the N

347 fragments, which was in accordance with the results in **Table 3**. The interaction of  
 348 flowering time and litter addition exerted similar effects on soil N pool as well as its  
 349 N fragments with flowering time solely.

350 **Table 2**  $\alpha$  values of soil N and P pools in various species litters addition treatment (n  
 351 = 14 and n = 15 in earlier flowering species and later flowering species, respectively).  
 352 TP and A-P are total phosphorus and available phosphorus, respectively.  $\alpha$  values  
 353 indicate natural logarithm of ratio flower litter addition to non-addition control of  
 354 different soil indexes (i.e., TN, NO<sub>3</sub><sup>-</sup>-N, NH<sub>4</sub><sup>+</sup>-N, TP, and A-P; the same below).

| Flowering period  | Index                           | Mean | Std. Error | Minimum | Maximum | F      | P            |
|-------------------|---------------------------------|------|------------|---------|---------|--------|--------------|
| Earlier flowering | TN                              | 1.67 | 0.06       | 1.29    | 2.05    | 719.05 | <b>0.000</b> |
|                   | NO <sub>3</sub> <sup>-</sup> -N | 1.67 | 0.07       | 1.08    | 2.23    | 563.90 | <b>0.000</b> |
|                   | NH <sub>4</sub> <sup>+</sup> -N | 0.97 | 0.12       | 0.42    | 2.06    | 68.25  | <b>0.000</b> |
|                   | TP                              | 0.02 | 0.03       | -0.04   | 0.08    | 8.498  | <b>0.007</b> |
|                   | A-P                             | 0.31 | 0.17       | 0.67    | 0.13    | 47.39  | <b>0.000</b> |
| Later flowering   | TN                              | 1.29 | 0.21       | -0.37   | 2.40    | 38.37  | <b>0.000</b> |
|                   | NO <sub>3</sub> <sup>-</sup> -N | 1.11 | 0.18       | -0.75   | 1.55    | 37.77  | <b>0.000</b> |
|                   | NH <sub>4</sub> <sup>+</sup> -N | 0.36 | 0.05       | -0.09   | 0.72    | 60.64  | <b>0.000</b> |
|                   | TP                              | 0.03 | 0.11       | -0.20   | 0.12    | 0.97   | 0.33         |
|                   | A-P                             | 0.50 | 0.23       | 0.06    | 0.37    | 68.82  | <b>0.000</b> |

355  
 356 Flower litters exerted different effects on soil TP and A-P. Soil TP increased in  
 357 treatment with early flowering litters (**Fig. 4 (a)**, **Table 2**,  $F = 8.498$ ,  $P = 0.007$ ) but  
 358 not in later flowering litters (**Fig. 4 (b)**, **Table 2**,  $F = 0.97$ ,  $P = 0.33$ ). The minimum  $\alpha$   
 359 values were lower than 0 ( $-0.04$  and  $-0.20$ , respectively). However, the A-P of both  
 360 litter treatments was significantly positively stimulated ( $F = 47.39$ ,  $P < 0.001$ ;  $F =$   
 361  $68.82$ ,  $P < 0.001$ ), whose  $\alpha$  values were both higher than 0 ( $0.67$ – $0.13$  and  $0.06$ – $0.37$ ,  
 362 respectively). Multifactorial analysis indicated that soil TP was not significantly  
 363 different between the sample treated with flower litter and the control (**Table 3**,  $F =$   
 364  $1.07$ ,  $P = 0.37$ ). No significant interaction was evident between flowering time and  
 365 litter addition treatments on soil TP ( $F = 0.01$ ,  $P = 0.93$ ). Litter addition treatments  
 366 alone only had a marginal significant effect on soil TP ( $F = 3.17$ ,  $P = 0.08$ ). Moreover,  
 367 both minimum  $\alpha$  values were lower than 0, but TP was not significantly different  
 368 between treatments with later flowering litters and control treatment ( $F = 0.97$ ,  $P =$

369 0.33), which mainly resulted from *G. sino-ornata*, *L. sinense*, and *C. lineare*.  
 370 Nevertheless, A-P increased significantly after flower litter addition ( $F = 43.01$ ,  $P <$   
 371  $0.001$ ), with a significant interaction between flowering time and litter addition ( $F =$   
 372  $6.44$ ,  $P < 0.05$ ).

373 **Table 3** Multifactorial analysis of variance for the effects of flowering time, litter  
 374 addition, and their interactions on soil N and P pools.

| Source of variation                            | TN       |             | NO <sub>3</sub> <sup>-</sup> -N |             | NH <sub>4</sub> <sup>+</sup> -N |             | TP       |          | A-P      |             |
|--|----------|-------------|---------------------------------|-------------|---------------------------------|-------------|----------|----------|----------|-------------|
|  | <i>F</i> | <i>P</i>    | <i>F</i>                        | <i>P</i>    | <i>F</i>                        | <i>P</i>    | <i>F</i> | <i>P</i> | <i>F</i> | <i>P</i>    |
| Corrected Model                                | 59.25    | <b>0.00</b> | 69.24                           | <b>0.00</b> | 54.07                           | <b>0.00</b> | 1.07     | 0.37     | 43.01    | <b>0.00</b> |
| Flowering time                                 | 2.80     | 0.10        | 7.93                            | <b>0.01</b> | 24.36                           | <b>0.00</b> | 0.02     | 0.90     | 6.44     | <b>0.01</b> |
| Litter addition treatments                     | 173.47   | <b>0.00</b> | 194.34                          | <b>0.00</b> | 117.00                          | <b>0.00</b> | 3.17     | 0.08     | 114.14   | <b>0.00</b> |
| Flowering time ×<br>Litter addition treatments | 2.80     | 0.10        | 7.93                            | <b>0.01</b> | 24.36                           | <b>0.00</b> | 0.02     | 0.90     | 6.44     | <b>0.01</b> |

375 Note: *P* values for significant effects and interactions are in bold (at the level  $P=0.05$ ).

376

377 **Effects of flower litter addition on soil solution N pool and soil MBC and MBN**

378 Soil solution N pool has been improved noticeably from 31.46 mg g<sup>-1</sup> to 47.35 mg g<sup>-1</sup>  
 379 in flower litter treatment compared with the control, particularly in fragment of NO<sub>3</sub><sup>-</sup>-  
 380 N, which has been greatly increased (from 30.93 mg g<sup>-1</sup> to 46.8 mg g<sup>-1</sup>) (**Table 4**). In  
 381 mixed leaf litter treatment, no obvious variations were found after litter  
 382 decomposition, with 32.4 mg g<sup>-1</sup> NO<sub>3</sub><sup>-</sup>-N and 0.45 mg g<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N, respectively.  
 383 Notable differences in both MBC and MBN were found between different treatments.  
 384 Litter addition increased not only soil microbial biomass C (102.05, 68.08, and 46.25  
 385 mg kg<sup>-1</sup> for flower litter, mixed litter, and control, respectively) and MBN (73.02,  
 386 69.29, 67.13 mg kg<sup>-1</sup> for flower litter, mixed litter, and control, respectively) but also  
 387 their C/N ratios (1.40, 0.98, and 0.69 for flower litter, mixed litter, and control,  
 388 respectively).

389 **Table 4** Comparison of the mean values of soil solution pool and soil microbial  
 390 biomass between litter addition treated (flower litter and mixed leaf litter) and control.

| Treatments        | Soil solution N pool (mg g <sup>-1</sup> ) |                                 | Soil microbial biomass (mg kg <sup>-1</sup> ) |       |         |
|-------------------|--|---------------------------------|---|-------|---------|
|                   | NO <sub>3</sub> <sup>-</sup> -N            | NH <sub>4</sub> <sup>+</sup> -N | MBC   | MBN   | MBC/MBN |
| Flower litter     | 46.8                                       | 0.55                            | 102.05  | 73.02 | 1.40    |
| Mixed leaf litter | 32.4                                       | 0.45                            | 68.08   | 69.29 | 0.98    |
| Control           | 30.93                                      | 0.53                            | 46.25   | 67.13 | 0.69    |

391

392

393 **Comparison of decomposition rate between flower litter and mixed leaf litter**

394 *R. przewalskii* and *M. integrifolia* are two typical plant species widely distributed and  
395 easily collected. Both species were assessed to compare decomposition rates of their  
396 flower litter and mixed leaf litter. Differences in decomposition rate among flower  
397 litter of two species and mixed litter were supposed to be significant (**Fig. 5**,  $F =$   
398 130.34,  $P < 0.001$ ). The flower litters of *R. przewalskii* and *M. integrifolia*  
399 decomposed greatly faster than mixed leaf litter. Moreover, within only 50 days,  
400 more than 20% of *R. przewalskii* and *M. integrifolia* flower litters decomposed,  
401 whereas the decomposition rate for mixed leaf litter was approximately 6% only  
402 (i.e., the former was nearly three times faster). Moreover, no significant differences  
403 were evident in the decomposition rates of the flower litter of *R. przewalskii* and *M.*  
404 *integrifolia* ( $P = 0.371$ ).

405

406 **Discussion**

407 Plant litter decomposition is a critical step in the formation of SOM, mineralization of  
408 organic nutrients, and C balance in terrestrial ecosystems (Austin and Ballaré, 2010;  
409 Cotrufo et al., 2015). At an early stage of decomposition, there exists partly  
410 correlation between decomposed plant material and light fraction in the SOM pool at  
411 a transitional stage of humification process (Leifeld and Kögel-Knabner, 2005).  
412 Species-specific variations in plant phenology can affect production of litter fall,  
413 which is noticeable during the growing season from the aspect of nutrient cycling  
414 although the peak of litter fall happens in autumn (the Northern Hemisphere). Thus,  
415 the early litter fall of alpine plants during the study period from May to August can be  
416 a potential nutrient source when nutritional demands increase for rapid growth and  
417 development. In particular, the amount of flower fall in the study area exceeds the leaf  
418 fall during the blooming season. A previous study indicated that reproductive litter  
419 production accounted for  $< 10\%$  of the total litter in January–August and  $13\%$ – $26\%$  in  
420 September–December (Sanches et al., 2008), which was mainly triggered by rainfall  
421 variability that directly altered litter production dynamics and indirectly altered forest  
422 floor litter. In addition, the flowers are more nutritional than the leaves in terms of

423 nutrients necessary for plant growth (Lee et al., 2011). In this study, summit  
424 production of flower litters is booming during special periods for both earlier  
425 flowering and later flowering species. Flower biomass of herbaceous plants accounts  
426 for 10% to approximately 40% of total aboveground biomass. Moreover, these flower  
427 litters produced considerably earlier than other aboveground litters that dropped at the  
428 end of the growing season. Furthermore, flower litters and non-flower litters (mainly  
429 constituted of leaves) of woody plants were 10–40 and 5–25 g m<sup>-2</sup>, respectively,  
430 which clearly implies that flower litter can be a comparable decomposition substrate  
431 in alpine ecosystems even for phanerophyte plants.

432 Litter production and decomposition are controlled by biological and physical  
433 processes, such as the activity and composition of soil and litter fauna and climate  
434 variations (Meentemeyer, 1978; Cornejo et al., 1994; Wieder and Wright, 1995; Aerts,  
435 1997; Cleveland et al., 2004). An integration of index or traits has been recommended  
436 to indicate the process and rate of litter decomposition. Generally, tissues with high  
437 lignin, polyphenol, and wax contents and higher lignin/N and C/N ratios exhibit slow  
438 decomposition. Lignin/N and C/N ratios are commonly accepted as good indicators of  
439 decomposition rates under short time frames; however, there is minimal conclusive  
440 evidence that lignin is preferentially preserved in soils compared with bulk soil over  
441 long-time periods (Melillo et al., 1982; Mikutta et al., 2005; Kleber et al., 2007;  
442 Cotrufo et al., 2015). Moreover, lignin plays a dual role in plant litter decomposition  
443 when photochemical mineralization and abiotic decomposition are considered (Austin  
444 and Ballaré, 2010). Leaf litter with C/N ratios lower than 30 is known to decompose  
445 easily and yield a mull humus type, whereas C/N ratios above 30 result in N  
446 immobilization (Heal et al., 1997) and decomposition retardation. In this study, flower  
447 litter had significantly less C/N ratio ( $19.80 \pm 1.39$ , less than 30) than leaf litter ( $39.27$   
448  $\pm 4.16$ , more than 30). Structural (lignin, DMC) and chemical (N) traits are proposed  
449 to be better predictors for several high-turnover organs than structural traits alone  
450 (Freschet et al., 2012). Lignin content in flower litters was significantly less than that  
451 in leaf litters ( $211.37 \pm 8.63$  mg kg<sup>-1</sup> and  $237.88 \pm 6.89$  mg kg<sup>-1</sup>, respectively;  $F =$   
452  $5.77$ ,  $P = 0.02$ ), similar to cellulose ( $266.93 \pm 4.92$  mg kg<sup>-1</sup> and  $283.75 \pm 4.21$  mg kg<sup>-1</sup>



453 <sup>1</sup>, respectively;  $F = 6.74$ ,  $P = 0.01$ ), which is one of the major cell wall constituents.  
454 All of the results are in accordance with previous studies. Decomposition rate is  
455 negatively correlated with the concentration of lignin, which is a group of complex  
456 aromatic polymers that serves as a structural barrier impeding microbial access to  
457 labile C compounds (Swift et al., 1979; Taylor et al., 1989; Austin and Ballaré, 2010;  
458 Talbot and Treseder, 2012). Moreover, the absence of significant differences of total C  
459 content in flower litters but with significantly less structural carbohydrates than those  
460 in leaf litters indicated that greater non-structural carbohydrates existed in flower  
461 litters. This finding can be inferred from the contents of lignin and cellulose (**Fig. 3**  
462 **(a)**). Hence, flower litters can promote nutrients that easily complement soil (Parton et  
463 al., 2007) for plants in their entire life history. Decomposition rates of leaf litters have  
464 been considered recently from their lignin/N or lignin/cellulose (Talbot and Treseder,  
465 2012; Cornwell et al., 2008). Furthermore, in the present study, lignin/N was less in  
466 flower litters (almost 50% in leaf litters, i.e.,  $12.79 \pm 1.15$  and  $21.09 \pm 2.25$ ,  
467 respectively), whereas N/P was higher than that of leaf litters.

468 A litterbag experiment on two widely distributed dominant shrubs (*R. przewalskii* and  
469 *M. integrifolia*) confirmed that the decay rates of flower litters were significantly  
470 faster than that of other litters, which is in accordance with the fast decomposition of  
471 *R. pseudoacacia* flower from an experiment performed in Korea (Lee et al., 2010).  
472 Flower litters contained significantly higher N and P contents than leaf litters (**Fig. 3**  
473 **(b)**). Plant litter available to the decomposer community encompasses a broad range  
474 of issues that differ in chemical and physical properties (Swift et al., 1979). P has been  
475 regarded as essential for a long time, which leads to limited attention to mechanisms  
476 that drive P limitation and their interactions with the N cycle (Vitousek et al., 2010).  
477 In most soils, the concentration of orthophosphate in solution is low (Richardson et  
478 al., 2009). Although soil generally contains a large amount of total P, only a small  
479 proportion is immediately available for plant uptake from the soil solution. P is  
480 derived mainly from rock weathering and related biogeochemical cycle, and  
481 ecosystems begin their existence with a fixed complement of P, and even very small  
482 losses cannot be readily replenished (Walker and Syers, 1976). The present study

483 indicated that decomposition of flower litter can be one of the beneficial source of soil  
484 A-P in alpine ecosystems. Decomposition rates can be markedly affected by particle  
485 size, surface area, and mass characteristics (Angers and Recous, 1997). In addition,  
486 physical toughness (lignin, dry matter content, or C content) can be suitable predictors  
487 of decomposition across all of the organs. Nevertheless, the current study regarding  
488 the characteristics and driven mechanism of this source remains at the first stage.  
489 Variation in soil physical–chemical properties, vegetation types, and microbial  
490 activities can significantly affect chemical compositions and forms as well as the  
491 biological availability of soil P directly or indirectly.

492 Decay rates of different plant organs reflect the diversity that fruits decompose faster  
493 than leaves, which in turn decompose faster than woody plant parts (Swift et al.,  
494 1979; Kögel–Knabner, 2002). Flower litters decompose rapidly with higher N and P  
495 levels supplied to soil, particularly from  $\text{NO}_3^-$ -N in soil solution pool (**Table 4**).  
496 Histogram for  $\alpha$  values of DIN and A-P also presented soil available nutrients  
497 positively stimulated by flower litter (**Fig. 6**) for their values distributed at an interval  
498 greater than 0. The high DOC values in flower litter may influence N and P in soil  
499 through C substrate supplement for soil microorganisms to enhance N  
500 immobilization. Recent empirical studies noted that the changing microbial  
501 community composition significantly affects ecosystem processes, such as litter  
502 decomposition (Strickland et al., 2009; Ramirez et al., 2012). Shifts from bacterial-  
503 dominated to fungal-dominated decomposition happened over short (days to a few  
504 months) periods (Poll et al., 2008; McMahon et al., 2005). Although the present study  
505 did not present the precise analysis of microbial community, both MBC and MBN  
506 differed greatly between different treatments (**Table 4**). Litter addition increased them  
507 obviously, which is evident not only in microbial biomass C and N but also in their  
508 C/N ratios (1.40, 0.98, and 0.69 for flower litter, mixed litter, and control,  
509 respectively). Flower litter contains more than twice MBC (increased from 46.25 to  
510 102.05), and both MBC and MBN pools increased potentially after flower litter  
511 addition. Therefore, microbial functional groups might be changed for nutrient  
512 supplement from litters or could also be due to their faster turnover or growth, which

513 need more evidence in the further study by directly testing soil microbial community  
514 composition.

515 Several unexpected species in the experiment reduced soil available nutrients  
516 probably because their specific chemical properties, which change as a result of  
517 microbial activities and nutrient dynamics (Karmarkar and Tabatabai, 1991), may  
518 negatively affect soil microorganism biomass or activities (Wardle et al., 1998,  
519 Cipollini et al., 2012). Furthermore, soil microbial communities can be modified  
520 through time in response to allelopathic plants with known or potential effects on  
521 plant communities (Cipollini et al., 2012, Inderjit and Weiner, 2001). Soil carbon  
522 generally is divided into pools with varying intrinsic decomposition rates in turnover  
523 models, whose decomposition rates can be modified and codetermined by interaction  
524 between substrates, microbial actors, and abiotic driving variables. These factors are  
525 rationalized by assuming chemical structure is a primary controller of decomposition  
526 (Kleber et al., 2010). Most of the non-fertilizer N source needed for plant growth is  
527 SOM (Sollins et al., 2007), which consists of organic molecular fragments with wide-  
528 ranging amphiphilicity degrees, intimately contacting with mineral surfaces of  
529 variable chemical reactivity and a polar solvent. Mineralization and nitrification can  
530 be subdued by inhibitory compounds from the exudates of a certain plant species,  
531 which come from a negative aspect and mainly result from suppression of related  
532 microbes (Cipollini et al., 2012). In another positive perspective, considering  
533 “*priming effect*” once flower litter is added in moderate treatments causes strong  
534 short-term changes in the turnover of SOM, and nutrient release follows litter  
535 decomposition (Jenkinson et al., 1985; Kuzyakov et al., 2000; Blagodatskaya and  
536 Kuzyakov, 2008). Hence, N and P availability in the soil of alpine ecosystem can be  
537 maintained in part by tissue chemistry favorable to microbial decomposition and  
538 release of nutrients.

539 Flower litter influences different fractions in soil N and P pools as well as soil  
540 microbial biomass (i.e., MBC and MBN), which provided evidence that plant species,  
541 through tissue chemistry, biomass allocation, and phenology, affect local soil  
542 properties and SOM formation in alpine ecosystems. Soil has specific susceptibility to

543 decomposition of biochemical compounds in plant tissues, on a spectrum from  
544 quickly decomposed labile to relatively recalcitrant. Flower litters have intuitive  
545 benefits chemically and physically for the formation, stabilization, and mineralization  
546 process of SOM. In future studies, major scientific findings and also potential  
547 questions less studied previously should be highlighted, and scientific obstacles  
548 should be considered to further address the stabilization and destabilization of SOM in  
549 this field. In brief, under a changing climate and a steadily increasing service demand  
550 in the alpine ecosystems, it is essential to understand the mechanisms underlying  
551 SOM stabilization. Furthermore, soil carbon models would benefit from taking flower  
552 litters' decomposition with specific attribution into soil nutrition pools. Flower litters  
553 affect carbon and nutrient cycling and should be incorporated into SOM pools along  
554 with decomposition simultaneously, which should be enhanced in future studies to  
555 better understand the essentiality and fundamentality of litter decomposition.

#### 556 **Acknowledgment**

557 This study was financially supported by the International Cooperation Project of  
558 Science and Technology Department of Sichuan Province (2014HH0056), China  
559 Postdoctoral Science Foundation (2014M552385), and National Natural Science  
560 Foundation of China (31400389). We would like to acknowledge the Key Lab of  
561 Ecological Restoration and Biodiversity Conservation of Sichuan (ECORES) for their  
562 support in laboratory facilities. We all appreciate that Mr. Jiceng Xu revised the  
563 sampling sites on the map.

564 **References**

- 565 Aber JD, Melillo JM, Mcclaugherty CA: Predicting Long-Term Patterns of Mass-  
566 Loss, Nitrogen Dynamics, and Soil Organic-Matter Formation from Initial  
567 Fine Litter Chemistry in Temperate Forest Ecosystems. *Can J Bot.*, 68: 2201-  
568 2208, 1990.
- 569 Aerts R.: Climate, leaf litter chemistry and leaf litter decomposition in terrestrial  
570 ecosystems: A triangular relationship. *Oikos*, 79: 439-449, 1997.
- 571 Angers, D.A., and Recous, S.: Decomposition of wheat straw and rye residues as  
572 affected by particle size. *Plant Soil.*, 189: 197-203, 1997.
- 573 Arroyo MTK, Pacheco DA, Aguilera P.: Floral allocation at different altitudes in  
574 highly autogamous alpine *Chaetanthera euphrasioides* (Asteraceae) in the  
575 central Chilean Andes. *Alpine Bot.*, 123: 7-12, 2013.
- 576 Austin, A. T., & Ballaré, C. L.: Dual role of lignin in plant litter decomposition in  
577 terrestrial ecosystems. *Proc Natl Acad Sci.*, 107(10): 4618-4622, 2010.
- 578 Berendse F, Aerts R.: Nitrogen-use-efficiency:A Biologically Meaningful Definition.  
579 *Funct Ecol* 1: 4, 1987.
- 580 Berg B, Ekbohm G.: Nitrogen Immobilization in Decomposing Needle Litter at  
581 Variable Carbon - Nitrogen Ratios. *Ecology*, 64: 63-67, 1983.
- 582 Bielecki RL.: Phosphate Pools, Phosphate Transport, and Phosphate Availability.  
583 *Annu Rev Plant Phys.*, 24: 225-252, 1973.
- 584 Brookes PC, Landman A, Pruden G, Jenkinson DS: Chloroform fumigation and the  
585 release of soil nitrogen: a rapid direct extraction method to measure microbial  
586 biomass nitrogen in soil. *Soil Biol Biochem.*, 17: 837-842, 1985.
- 587 Buxton DR, Marten GC: Forage Quality of Plant-Parts of Perennial Grasses and  
588 Relationship to Phenology. *Crop Sci* 29: 429-435, 1989.
- 589 Chapin FS, Vitousek PM, Vancleve K.: The Nature of Nutrient Limitation in Plant-  
590 Communities. *Am Nat.*, 127: 48-58, 1986.
- 591 Chapin FS.: Effects of plant traits on ecosystem and regional processes: a conceptual  
592 framework for predicting the consequences of global change. *Ann Bot.*, 91:  
593 455-463, 2003.
- 594 Cleveland CC, Neff JC, Townsend AR, Hood E. Composition, dynamics, and fate of  
595 leached dissolved organic matter in terrestrial ecosystems: Results from a  
596 decomposition experiment. *Ecosystems*, 7: 275-285, 2004.
- 597 Cornejo FH, Varela A, Wright SJ: Tropical Forest Litter Decomposition under  
598 Seasonal Drought: Nutrient Release, Fungi and Bacteria -. *Oikos* 70: 183-190,  
599 1994.
- 600 Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H.  
601 and Parton, W.J: Formation of soil organic matter via biochemical and  
602 physical pathways of litter mass loss. *Nature Geosci.*, 2015.
- 603 Dalal RC. Estimation of Available Phosphorus in Soils by Extraction with Sodium  
604 Hydroxide Sodium Carbonate Solution. *J Aus Ins Agr Sci.*, 39: 142-143, 1973.
- 605 Fabbro T, Korner C: Altitudinal differences in flower traits and reproductive  
606 allocation. *Flora*, 199: 70-81, 2004.

607 Freschet GT, Aerts R, Cornelissen JHC: A plant economics spectrum of litter  
608 decomposability. *Funct Ecol.*, 26: 56-65, 2012.

609 Grime JP: Benefits of plant diversity to ecosystems: immediate, filter and founder  
610 effects. *J Ecol.*, 86: 902-910, 1998.

611 Groffman PM: Carbon additions increase nitrogen availability in northern hardwood  
612 forest soils. *Biol Fertil Soils.*, 29: 430-433, 1999.

613 Hautier Y, Randin CF, Stocklin J, Guisan A: Changes in reproductive investment with  
614 altitude in an alpine plant. *J Plant Ecol-Uk.*, 2: 125-134, 2009.

615 Haynes RJ: Mineral Nitrogen in the Plant-Soil System (Physiological Ecology).  
616 Academic Press Inc, 1986.

617 Heal OW, Anderson JM, Swift MJ: Plant litter quality and decomposition: an  
618 historical overview. In: Cadisch, G, Giller, KE eds. *Driven by nature. Plant  
619 litter quality and decomposition.* CAB International, Wallingford, 3-30, 1997.

620 Herrick, J.E., Van Zee, J.W., Havstad, K.M., Burkett, L.M. and Whitford, W.G.:  
621 Monitoring manual for grassland, shrubland and savanna ecosystems. Volume  
622 I: Quick Start. Volume II: Design, supplementary methods and interpretation.  
623 USDA-ARS Jornada Experimental Range, 2005.

624 Inderjit, Weiner, J.: Plant allelochemical interference or soil chemical ecology?  
625 *Perspect Plant Ecol.*, 4 (1): 3-12, 2001.

626 Jones DL, Shannon D, Murphy DV, Farrar J: Role of dissolved organic nitrogen  
627 (DON) in soil N cycling in grassland soils. *Soil Biol Biochem.*, 36: 749-756,  
628 2004.

629 Kleber, M., Sollins, P. and Sutton, R.: A conceptual model of organo-mineral  
630 interactions in soils: self-assembly of organic molecular fragments into zonal  
631 structures on mineral surfaces. *Biogeochemistry*, 85(1): 9-24, 2007.

632 Kleber, M., Nico, P.S., Plante, A., Filley, T., Kramer, M., Swanston, C. and Sollins, P.:  
633 Old and stable soil organic matter is not necessarily chemically recalcitrant:  
634 implications for modeling concepts and temperature sensitivity. *Global  
635 Change Biol.*, 17(2): 1097-1107, 2010.

636 Körner C.: *Alpine plant life: functional plant ecology of high mountain ecosystems.*  
637 2nd edition. Springer, Berlin; New York, 2003.

638 Laungani R, Knops JMH.: Species-driven changes in nitrogen cycling can provide a  
639 mechanism for plant invasions. *Proc Natl Acad Sci.*, 106: 12400-12405, 2009.

640 Leifeld, J. and Kögel-Knabner, I.: Soil organic matter fractions as early indicators for  
641 carbon stock changes under different land-use?. *Geoderma*, 124(1):143-155,  
642 2005.

643 Lee YC, Nam JM, Kim JG.: The influence of black locust (*Robinia pseudoacacia*)  
644 flower and leaf fall on soil phosphate. *Plant Soil.*, 341: 269-277, 2011.

645 MacLean DA and Wein RW.: Litter production and forest floor nutrient dynamics in  
646 pine and hardwood stands of New Brunswick, Canada. *Ecography*, 1: 1-15,  
647 1978.

648 Meentemeyer V.: Macroclimate and Lignin Control of Litter Decomposition Rates.  
649 *Ecology* 59: 465-472, 1978.

650 Melillo JM, Aber JD, Muratore JF.: Nitrogen and Lignin Control of Hardwood Leaf  
651 Litter Decomposition Dynamics. *Ecology* 63: 621-626, 1982.

652 Melillo, J.M., Aber, J.D., Linkins, A.E., Ricca, A., Fry, B. and Nadelhoffer, K.J.:  
653 Carbon and nitrogen dynamics along the decay continuum: plant litter to soil  
654 organic matter. *Plant Soil.*, 115, 189-198, 1989.

655 Mikutta, R., Kleber, M., Kaiser, K. and Jahn, R.: Review: organic matter removal  
656 from soils using hydrogen peroxide, sodium hypochlorite, and disodium  
657 peroxodisulfate. *Soil Sci Soc Am J.*, 69(1): 120-135, 2005.

658 Muller-Landau, H.C. and Wright, S.J.: Litterfall Monitoring Protocol, 2010.

659 Nilsson C, Xiong SJ, Johansson ME, Vought LBM.: Effects of leaf-litter accumulation  
660 on riparian plant diversity across Europe. *Ecology* 80: 1770-1775, 1999.

661 Norman RJ, Edberg JC, Stucki JW.: Determination of Nitrate in Soil Extracts by  
662 Dual-wavelength Ultraviolet Spectrophotometry. *Soil Sci Soc Am J.*, 49:  
663 1182-1185, 1985.

664 Richardson AE, Barea JM, McNeill AM, Prigent-Combaret C.: Acquisition of  
665 phosphorus and nitrogen in the rhizosphere and plant growth promotion by  
666 microorganisms. *Plant Soil.*, 321: 305-339, 2009.

667 Sah RN.: Nitrate-Nitrogen Determination-A Critical Review. *Commun Soil Sci Plan.*,  
668 25: 2841-2869, 1994.

669 Sanches L, Valentini CMA, Pinto OB, Nogueira JD, Vourlitis GL, Biudes MS, da  
670 Silva CJ, Bambi P, Lobo FD.: Seasonal and interannual litter dynamics of a  
671 tropical semideciduous forest of the southern Amazon Basin, Brazil. *J*  
672 *Geophys Res-Bioge.*, 113, 2008.

673 Scott NA, Cole CV, Elliott ET, Huffman SA.: Soil textural control on decomposition  
674 and soil organic matter dynamics. *Soil Sci Soc Am J.*, 60: 1102-1109, 1996.

675 Sollins, P., Swanston, C. and Kramer, M.: Stabilization and destabilization of soil  
676 organic matter—a new focus. *Biogeochemistry*, 85(1),1-7, 2007.

677 Stohlgren TJ., Litter dynamics in two Sierran mixed conifer forests. II. Nutrient  
678 release in decomposing leaf litter. *Can J For Res.*, 18:1136-1144, 1988.

679 Vitousek PM, Howarth RW.: Nitrogen Limitation on Land and in the Sea - How Can  
680 It Occur. *Biogeochemistry*, 13: 87-115, 1991.


681 Wieder RK, Wright SJ.: Tropical Forest Litter Dynamics and Dry Season Irrigation on  
682 Barro-Colorado Island, Panama. *Ecology*, 76: 1971-1979, 1995.

683 Wookey PA, Aerts R, Bardgett RD, Baptist F, Brathen KA, Cornelissen JHC, Gough  
684 L, Hartley IP, Hopkins DW, Lavorel S, Shaver GR.: Ecosystem feedbacks and  
685 cascade processes: understanding their role in the responses of Arctic and  
686 alpine ecosystems to environmental change. *Glob Chang Biol.*, 15: 1153-1172,  
687 2009.

688 Xiong SJ, Nilsson C.: The effects of plant litter on vegetation: a meta-analysis. *J Ecol*  
689 87: 984-994, 1999.


690 Yavitt JB, Fahey TJ.: Litter Decay and Leaching from the Forest Floor in Pinus-  
691 Contorta (*Lodgepole Pine*) Ecosystems. *J Ecol.*, 74: 525-545, 1986.

692



694

695 **Fig. 1** Location of study sites.




696


697 **Fig. 2** Production of flower litters and biomass allocation of representative dominant  
 698 species. (a) Production of flower litters and non-flower litters of shrubs  
 699 (phaenophyte, n=20) per unit area (m<sup>2</sup>); and (b) floral biomasses and their allocation  
 700 in the aboveground biomass.

701






702



703




704

705 **Fig. 3** Chemical composition and their comparison between flower and leaf litters.  
 706 Whiskers refer to quantiles for comparable data settings. Asterisks (\*) represent  
 707 distribution of extreme outliers. M=mean and N, which indicates data/sample number,  
 708 are analyzed and processed by one-way ANOVA (at  $P=0.05$  level).

709

710

711




712

713

714


715




716

717 **Fig. 4** Variation in soil N pool and P pool after addition of flower litters, (a) earlier  
 718 flowering species, and (b) later flowering species. Scatters represent  $\alpha$  mean values of  
 719 different indexes. Significant differences of deviations from the 0 lines are tested at  
 720  $P=0.05$  level ( $n=3$ ). TN, NO<sub>3</sub>-N, NH<sub>4</sub>-N, TP, and A-P represent total nitrogen, nitrate  
 721 nitrogen, ammonium nitrogen, total phosphorus, and available phosphorus,  
 722 respectively.

723



724  
 725 **Fig. 5** Percentage of decomposed dry mass of *M. integrifolia* and *R. przewalskii* in a  
 726 50-day litter-bag study. Column represents mean, and bar indicates standard Error (n =  
 727 8). Different lowercase letters indicate significant differences of decomposition rate  
 728 between litter materials (at  $P=0.05$  level).



729  
 730 **Fig. 6** Variation in soil nutrition pool with flower litters addition. Histogram for  $\alpha$   
 731 values of DIN (a) and A-P (b) indicates the change between treatments and control.