

Interactive comment on “Spatial variability of surface-water pCO₂ gas exchange in the world’s largest semi-enclosed estuarine system: St. Lawrence Estuary (Canada)” by Ashley Dinauer and Alfonso Mucci

Ashley Dinauer and Alfonso Mucci

ashley.dinauer@mail.mcgill.ca

Received and published: 11 February 2017

We thank the referee for his constructive comments, and time spent to review our manuscript. We are particularly grateful to this referee and his collaborators whose earlier work inspired us to undertake this study. Our response to his comments follow.

Given the physiography (funnel shape) and the large physical dimensions of the maritime portion of the St. Lawrence Estuary, we pursued an analysis of surface-water pCO₂ and CO₂ flux data in terms of space rather than salinity. With 75% of its total surface area occupied by the Lower Estuary (Tadoussac to Pointe-des-Monts, $S > \sim 21.2$),

[Printer-friendly version](#)

[Discussion paper](#)



it is more realistic to integrate/weight the fluxes spatially than to provide average fluxes binned by salinity, in order to obtain a representative whole-estuary flux. Furthermore, as you can appreciate from the appended figure (see Fig. 1), active processes along the estuary are not as clearly defined when the data are plotted as a function of surface salinity. Whereas the plot still shows that the Upper Estuary ($S < \sim 24.5$; surface salinity fluctuations are found around the Upper-Lower Estuary boundary due to tidally-induced upwelling of salty intermediate/bottom waters) is heterotrophic, it fails to highlight the autotrophic character of the Lower Estuary and the neutral character of the Gulf, the latter being substantially larger in surface area than the combined Upper and Lower Estuaries but contributing little to the net CO₂ flux.

As noted in lines 643-649, we suggest that biological production is responsible for the observed drawdown of pCO₂ in the surface waters of the Lower Estuary as well as the undersaturation observed near the Estuary-Gulf boundary (i.e., along the 300–600 km stretch). We agree with the reviewer's comment that this needs to be explicitly stated in the revised manuscript.

We will identify the red (smoothed temperature data using a moving average filter with a span of 50% of the total number of data points) and dashed (smoothed pCO₂(water) data) lines in the Fig. 9 caption, and add the appropriate reference to line 41 in the revised manuscript.

Interactive comment on Biogeosciences Discuss., doi:10.5194/bg-2017-1, 2017.

Printer-friendly version

Discussion paper






Fig. 1. Spatial distribution of surface-water pCO₂ (circles, squares) along the salinity gradient. Horizontal lines show the mean pCO₂(air) in the sampling months.

[Printer-friendly version](#)[Discussion paper](#)