



Ice Acidification: A review of the effects of ocean acidification on sea ice microbial communities

Andrew McMinn¹

5

¹Institute of Marine and Antarctic Science, University of Tasmania, Hobart 7001, Tasmania, Australia

Correspondence to: Andrew McMinn (andrew.mcminn@utas.edu.au)

10 **Abstract.** Sea ice algae are naturally exposed to a wider range of pH and CO₂ concentrations than marine phytoplankton. While climate change and ocean acidification (OA) will impact pelagic communities, their effects on sea ice microbial communities remains unclear.

Sea ice contains several distinct microbial communities, which are exposed to differing environmental conditions depending on their depth within the ice. Bottom communities mostly experience relatively benign bulk ocean properties, while interior
15 brine and surface communities experience much greater extremes.

Most OA studies have examined the impacts on single sea ice algae species in culture. Although some studies examined the effects of OA alone, most examined the effects of OA and either light, nutrients or temperature. With few exceptions, increased CO₂ concentration caused either no change or an increase in growth and/or photosynthesis. *In situ* studies of brine and surface algae also demonstrated a wide tolerance to increased and decreased pH and showed increased growth at higher
20 CO₂ concentrations. The short time period of most experiments (<10 days) together with limited genetic diversity (i.e. use of only a single strain), however, has been identified as a limitation to the broader interpretation of results.

While there have been few studies on the effects of OA on marine bacterial communities in general, impacts appear to be minimal. In sea ice also, the few reports available suggest no negative impacts on growth or community richness.

Sea ice ecosystems are ephemeral, melting and re-forming each year. Thus, for some part of each year organisms inhabiting
25 the ice must also survive outside of the ice, either as part of the phytoplankton or as resting spores on the bottom. During these times, they will be exposed to the full range of co-stressors that pelagic organisms experience. Their ability to continue to make a major contribution to sea ice productivity will depend not only on their ability to survive in the ice but also on their ability to survive the increasing seawater temperatures, changing distribution of nutrients and declining pH forecast for the water column over the next centuries.

30



1 Introduction

Sea ice is widely recognized as one of the most extreme habitable environments on earth (Thomas and Dieckmann, 2004, Martin and McMinn 2017). Organisms living within it can be exposed to temperatures of below -20°C and salinities greater than 200 for extended periods of time (Arrigo, 2014; Thomas and Dieckmann, 2004). These environments also endure long periods of darkness and extremes in nutrient concentration, dissolved gases (O_2 , CO_2) and pH (Thomas and Dieckmann 2004 McMinn et al., 2014). In spite of these conditions, some sea ice habitats are very productive and often support a dense microbial biomass (Arrigo, 2014). They also play a key supporting role in maintaining energy flow through polar marine food webs in winter and spring when sea ice covers much of the ocean surface and there is very little pelagic primary production (Arrigo and Thomas, 2004).

10

During the annual cycle of sea ice formation and melting, associated organisms are naturally exposed to a large range in gas concentrations, including CO_2 . Gases are expelled from the forming ice in autumn and winter and become concentrated in the trapped brine (Fig. 1). Consequently, there is a relationship between the CO_2 concentration of the brine and that of the original sea water. As dissolved CO_2 concentrations continue to rise with ocean acidification (OA), there will also be a commensurate increase in the CO_2 concentration in the trapped brine. This review will examine the effects of increasing CO_2 concentrations and decreasing pH on sea ice microbial communities. The review focuses on Antarctic sea ice communities as no Arctic studies have been identified.

15

2. Natural pH fluctuations

There are strong seasonal cycles in pH in the waters of the Southern Ocean due to brine drainage from forming sea ice in winter and high levels of biological activity in summer. (Miller et al., 2011). In Prydz Bay and the Ross Sea this has led to seasonal differences of up to 0.6 of a pH unit (Gibson and Trull, 1999; McNeil et al., 2010), which is greater than the 0.4 increase usually predicted for the end of the century (IPCC4). Large seasonal differences in pH have also been observed in McMurdo Sound, Antarctica (Kapsenberg et al., 2015; Matson et al., 2014), where pH was relatively high but stable during winter but was lower and dynamic in summer. Large diurnal fluctuations in pH beneath the sea ice were also observed and it was suggested that these changes could be almost entirely explained by the biological processes of photosynthesis and respiration (Matson et al., 2014).

25

During the early stages of ice formation in early winter, the concentration of CO_2 within the ice is determined solely by physical processes associated with freezing and brine rejection. While low temperatures increase the solubility of CO_2 , increasing brine salinity during sea ice growth leads to super-saturation (Papadimitriou et al., 2003). Change in temperature is subsequently the dominant factor determining CO_2 concentration (Geilfus et al., 2014). However, the incorporated sea ice

30



algae take up CO₂ for photosynthesis and, together with other components of the microbial food web such as bacteria, release it during respiration. The balance of these processes commonly produces brine CO₂ concentrations that are lower in summer and higher in winter relative to atmospheric levels (Gleitz et al., 1995; Thomas et al., 2010). Hare et al. (2013) examined the development of pH in an outdoor experimental sea ice facility in Winnipeg, Canada. Experiments were run over several weeks and ice attained a thickness of up to 22 cm. pH profiles had a characteristic ‘C’ profile with pH values greater than 9 at the surface and in the bottom 2 cm. Interior values were below that of the source water (8.4) with some as low as 7.1. The reason for this pattern was thought to be entirely due to abiotic factors as no algal growth was observed. The higher values at the surface and bottom reflected equilibrium with the atmosphere and underlying sea water, respectively, while those in the middle reflected CO₂ rejection during brine formation.

Changes in pH and CO₂ concentration in natural brines from the Weddell Sea in early summer have been documented by Papadimitriou et al. (2007). The *in situ* pH varied between 8.41 and 8.82, up more than 0.7 units from the underlying sea water. The concentration of CO₂ ranged from 3.1 μmol kg⁻¹ to 15.9 μmol kg⁻¹ and variations were consistent with biological activity (Papadimitriou et al., 2007).

As global temperatures warm, sea ice will inevitably form later in the season and melt sooner. As a consequence, average ice thickness will probably decrease. These large scale changes are already clearly evident in the Arctic (Barnhart et al. 2016). Thinner ice would result in average irradiances within and beneath the ice being higher, although this will also be modified by changes in snow thickness. However, there are still large uncertainties in predictions of future changes in precipitation over sea ice and while a warmer atmosphere might be expected to increase snow fall, there is no clear evidence of this happening so far or whether in future it will fall as snow or rain (Bracegirdle et al., 2008; Leonard and Maksym, 2011).

20

3. Biological communities

Sea ice is not a solid, uniform, homogenous layer but instead contains many different habitats that can be colonized by a large range of organisms. Each of these habitats is characterized by different physical and chemical conditions. Initially any organism present in the underlying water column has the potential to be trapped within the ice as it forms. The early process of ice formation (frazil ice formation) can concentrate the phytoplankton biomass many times above that of the surrounding water column (Garrison et al., 1983). All species present in the water column can be harvested, although some, such as *Nitzschia stellata*, have the ability to greatly increase the likelihood of their capture by the release of ice active substances (Raymond et al., 1994; Ugalde et al., 2014). Young ice typically contains a highly diverse microalgal community (Scott et al., 1994) but this diversity decreases with time as those species that are better adapted to this environment out compete those that are less well adapted. The biological communities can be differentiated by where they occur in the ice; in brine channels within the ice (brine communities), on the undersurface of the ice (bottom communities) or at the ice water interface (surface communities). Bottom communities typically inhabit a benign and equitable environment characterized by minimal changes in temperature and salinity, maximum access to nutrients but low light levels. Brine communities experience extreme

30



fluctuations in temperature and salinity, high to very low nutrient concentrations and intermediate light levels (Thomas and Dieckmann, 2004). Surface communities have access to the highest light but also experience the lowest temperature and nutrient levels. These communities also experience different CO₂/pH conditions. Bottom communities are constantly in contact with the underlying water column and typically experience the bulk ocean properties. Brine and surface communities, by contrast, experience supersaturated CO₂ concentrations during early ice formation but very low levels later in the season when the CO₂ is exhausted by photosynthesis.

Diatoms are the most studied group of sea ice microbes, but other eukaryotic and prokaryote groups are often important. While diatoms tend to dominate bottom communities (Garrison, 1991), dinoflagellates and other flagellate groups tend to dominate interior and surface communities (Stoecker et al., 1992; Thomson et al., 2006). There is also a diverse bacterial community (Mock and Thomas, 2005; Bowman et al., 2012; Bowman, 2013) and evidence of an active microbial loop (Martin et al., 2012).

4. Ocean acidification experiments

Studies of ocean acidification effects on sea ice algae and bacteria have taken a number of different approaches. Some have just examined the effects of increased CO₂ concentration in isolation (Coad et al., 2016, Torstensson et al., 2015), while other have looked at the combined effects of co-stressors such as temperature (Pančić et al., 2015; Torstensson, 2012), light (Xu et al., 2014; Heiden et al., 2016) or iron (Xu et al., 2014b). Most studies have been undertaken on cultures of single species in a laboratory (Mitchell and Beardall, 1996; Xu et al., 2014; Pančić et al., 2015; Torstensson et al., 2012; Young et al., 2015), although there has been one study of a brine community (Coad et al., 2016) and two *in situ* experiments on brine communities (McMinn et al., 2014; 2017). All but one of these studies (Torstensson et al., 2015) was conducted on a time scale of less than 30 days and most less than 10 days (Table 1).

Studied in isolation, increased CO₂ concentration seems only to have negative effects on growth and photosynthesis at levels above ~1000 μatm (Torstensson et al., 2015). When a natural diatom-dominated sea ice brine community was incubated over a CO₂ gradient from 400 μatm to 11,300 μatm for up to 18 days, only the treatment incubated at the most extreme concentration was significantly different from the control and other treatments (Coad et al., 2016). In a further experiment, they simulated future spring conditions by allowing a surface ice community to melt into seawater with the same large CO₂ gradient. Once again, only the most extreme CO₂ concentrations induced significant reductions in growth and photosynthesis. These results are consistent with most other studies of polar marine diatoms, which also show increased growth rates and photosynthetic performance at CO₂ concentrations up to at least 1000 μatm (Trimborn et al., 2013; Hoppe et al., 2013). Future increases in marine CO₂ concentration, however, are likely to be accompanied by changes in temperature, light, macro nutrient and iron concentrations (Boyd et al., 2016) and interactions between these and other factors are likely to produce unpredictable results (Boyd and Hutchins, 2012). However, not all of these changes are likely to occur within the sea ice environment.



Studies of current Southern Ocean phytoplankton communities have identified the importance of iron and irradiance as being the key drivers determining changes in community composition and productivity (Feng et al., 2010). Others have suggested that future changes in pCO₂, temperature and stratification, which effects average irradiance and nutrient availability, will also be significant (Constable et al., 2014). However, not all of these factors are directly relevant to ice algal communities. For instance, unlike the oceans, the temperature of sea ice habitats cannot rise above ~0°C. Increasing temperatures may well change the temporal and spatial distribution of sea ice but they will have only a small effect on the physical and chemical environments within the ice.

The combined effects of temperature and CO₂ concentration on the bipolar sea ice/phytoplankton species, *Fragilariopsis cylindrus* was examined in short incubations (7 days) by Pančić et al. (2015). Significant interactions between temperature and pH were identified but the two factors produced opposite effects. Growth rates increased with increasing temperatures but decreased with decreasing pH, resulting in little overall change with treatment. As a result, it was concluded that *F. cylindrus* would largely be unaffected by increasing ocean acidification. These experiments were conducted over a wide range of temperature (1°C to 8°C) and pH (7.1 to 8.0). A similar study on the sea ice/benthic diatom *Navicula directa* likewise found only minor responses to increased CO₂ with no synergistic effects with temperature, although a smaller range of temperature (0.5- 4.5°C) and pH (7.9-8.2) was used (Torstensson et al., 2012). These authors emphasized the need to examine multiple strains of a species and over longer incubation times before conclusions could be reached on its long term response. It should be noted that while both these studies used sea ice algae taxa, the cells were exposed to temperatures well above 0°C, i.e. temperatures that could never be experienced by cells living within sea ice. Although the maximum temperature of sea ice is fixed, the average temperature is still likely to rise. This could result in more intensive heterotrophic processes, via increased grazing rates and nutrient regeneration (Melnikov, 2009). However, it is probable that these processes, which already occur in spring and summer, will simply develop a little earlier.

Future changes in iron supply has been identified as likely to have a significant but unquantified effect on much of the world's phytoplankton (Shi et al., 2010) but the effects on sea ice communities are unknown; to date there have been no studies on the combined effects of increased CO₂ and iron-limitation on sea ice algae. A study of Southern Ocean phytoplankton, which contains many of the same species as sea ice communities, under co-limitation by CO₂, iron and other factors found that competition was likely to induce taxonomic changes in community composition, favoring small diatoms (Xu et al., 2014b). Algae growing at the ice water interface (bottom communities), like the phytoplankton, experience iron limiting conditions and show evidence of chronic iron stress (Pankowski and McMinn, 2008; 2009). However, the iron concentration within brine channels is one to two orders of magnitude greater than in the underlying seawater (Lannuzel et al., 2011) and sea ice brine algae have been shown not to be iron stressed at all (Pankowski and McMinn, 2008; 2009). Under these circumstances, iron cannot be considered a co-stressor.



The combined effect of CO₂ concentration and increased light on the important ice diatom, *Fragilariopsis curta* was examined by Heiden et al. (2016). Unlike most previous experiments, growth and photophysiology was not stimulated at relevant light (20 μmol photons m⁻² s⁻¹) and OA-relevant elevated pCO₂ (1000 μatm). These authors also showed that there were large variations in species-specific responses. The growth and photosynthetic response of the chlorophyte

5 *Chlamydomonas*, isolated from Antarctic sea ice, to ocean acidification and photoperiod was examined by Xu et al. (2014a) over the course of 28 days. While *Chlamydomonas* is not a common ice algal taxon, it none the less represents an organism adapted to living in the ice. Their data showed that CO₂ concentration had a minimal effect on the response to differing photoperiods.

It is well known that the physiological responses of diatoms in culture changes with time (Anderson, 2005). Some of the

10 culture experiments discussed herein used strains of diatoms that had been isolated decades earlier (Xu et al. 2014, Heiden et al. 2016). Therefore, the extreme age of the culture used needs to be taken into account when interpreting the physiological response.

There have been few *in situ* experiments with natural sea ice communities. McMinn et al. (2014, 2017) incubated

15 dinoflagellate-dominated brine algal communities *in situ* in McMurdo Sound, Antarctica. The 2014 study was conducted in spring while the 2017 study was conducted in late summer. They used sack holes to extract the brine (McMinn et al., 2009), adjusted the CO₂ concentrations of the treatments and then returned them to the same depth within the ice from where they were collected. Incubations were relatively short, up to 6 days. In both studies growth and photosynthesis were only affected when the pH fell below approximately 7.5. In a companion set of experiments in both studies the carbon system was

20 manipulated to hold the pH constant at ~8.0 while providing a gradient in CO₂ concentrations up to 2700 μatm. Growth increased by approximately 20% and remained constant at even the highest concentrations. This 20% increase in growth with increased CO₂ supply is of a similar scale to those estimated for a variety of temperate phytoplankton taxa on a similar CO₂ gradient (Rost et al., 2003; Riebesell, 2004).

25 The concentration of dissolved CO₂ in seawater is usually considered insufficient to maintain maximum growth rates in phytoplankton. As a consequence, most species have developed strategies to alleviate this stress; collectively referred to as Carbon Concentrating Mechanisms (CCM) (Raven et al., 2011). The presence of a CCM allows most marine microalgae, and diatoms in particular, to concentrate inorganic carbon as CO₂ and/or HCO₃ (Giordano et al., 2005) and thus alleviate much of this stress. Ocean acidification increases the natural CO₂ concentration and so potentially favours those species

30 without a CCM, however, it will also reduce the need to invest in this energetically demanding function (Beardall and Raven, 2004). Inorganic carbon uptake by the sea ice diatom *Nitzschia stellata* was examined by Mitchell and Beardall (1996). They demonstrated that this taxon was capable of carbon-dependent photosynthesis at rates greater than the CO₂ supply rate to the cell and that much of the carbonic anhydrase activity was associated with the cell surface; these characteristics demonstrated the presence of an active carbon concentrating mechanism (CCM). Other studies (Tortell et al.,



2013; Gibson et al., 1999; McMinn et al., 1999) have also found evidence for CCM activity from either carbon isotope analysis or carbonic anhydrase activity. So far, all sea ice species examined have shown evidence of an active CCM. However, little is still known about how these processes will respond to future changes in CO₂ concentration (Raven et al., 2011).

5

Sea ice contains an abundant and diverse, psychrophilic bacterial community (Mock and Thomas, 2005; Bowman et al., 2012; Bowman, 2013). Ocean acidification effects on marine bacterial communities in general appear to be minimal (Oliver et al., 2014; Lin et al., 2017). They are able to mitigate the effect of decreased pH by enhancing the expression of genes encoding proton pumps, such as respiration complexes, proteorhodopsin and membrane transporters (Bunce et al., 2016).

10 Likewise, in sea ice studies, bacterial growth rates have been found to increase with increasing CO₂ concentration (Torstensson et al., 2015). A pH range of 8.203 – 9.041 has also been found to have no effect on the community richness or diversity of Antarctic fast ice bacteria (Torstensson et al., 2012).

15 Sea ice ecosystems are ephemeral and in most relevant locations they melt and re-form each year. Thus, for some part of each year organisms inhabiting the ice must also survive outside of the ice, either as part of the phytoplankton or as resting spores on the bottom. During these times, they will be exposed to the full range of co-stressors that pelagic organisms experience (Xu et al., 2014c). Some sea ice taxa go into dormancy during these periods, e.g. *Polarella glacialis* (Thompson et al., 1996), while others, such as *Fragilariopsis curta* and *Fragilariopsis cylindrus* make a major contribution to phytoplankton biomass. Most sea ice microorganisms do not appear to be badly impacted by moderate increases in CO₂.

20 However, their ability to continue to make a major contribution to sea ice productivity will depend not only on their ability to survive in the ice but also on their ability to survive the increasing seawater temperatures, changing distribution of nutrients and declining pH forecast for the water column over the next centuries (Boyd and Hutchins, 2012).

Competing Interests

25 The author declares that he has no conflict of interest.

Acknowledgements

Australian Antarctic Science Grants program is acknowledged for its continuous financial support for sea ice research in general and for this review in particular.



References

- Anderson, R. A.; Algal culturing techniques, Elsevier Academic Press, Burlington MA, USA, 2005.
- Arrigo, K. R.: Sea Ice Ecosystems, *Ann. Rev. Mar. Sci.*, 6, 439-467, 2014
- 5 Arrigo KR (2014) Sea ice ecosystems. *Ann Rev Mar Sci* 6:439-467.
- Arrigo, K. R. and Thomas, D. N.: Large scale importance of sea ice biology in the Southern Ocean. *Antarctic Sci.*, 16, 471-486, 2004.
- 10 Barhhardt, K. R., Miller, C.R., Overeem, I., Kay, J.E.; Mapping the future expansion of Arctic open water. *Nature Climate Change*, 6, 280-285.
- Beardall, J, and Raven J. A.: The potential effects of global climate change on microalgal photosynthesis, growth and ecology, *Phycologia*, 43, 26-40, 2004.
- 15 Bowman, J. P.: Sea ice microbial communities, *In The Prokaryotes, Prokaryotic communities and ecophysiology*, eds Rosenberg E., DeLong E.F., Lory S., Stackbrandt Eand Thompson F, Springer-Verlag, Berlin, 139-161, 2013.
- 20 Bowman, J.S., Rasmussen, S., Blom, N., Deming, J. W., Rysgaard. S. and Sicheritz-Ponten, T.: Microbial community structure of Arctic multiyear sea ice and surface seawater by 454 sequencing of the 16S RNA gene. *ISME J.*, 6, 11-20, 2012.
- Boyd, P. W., Cornwall, C. E., Davidson, A., Doney, S. C., Fourquez F., Hurd, C.L., McMinn, A.: Environmental variability, heterogeneity, ocean biota, and climate change, *Global Change Biol.*, 22, 2633-2650, 2016.
- 25 Boyd, P. W., Dillingham, P. W., McGraw, C. M., Armstrong, E. A., Cornwall, C. E., Feng, C. L., Hurd, C. L., Gault-Ringold, M., Roleda, M.Y., Timmins-Schiffman, E., Nunn, B. L.: Physiological responses of a Southern Ocean diatom to complex future ocean conditions, *Nature Climate Change*, 6, 207-216, 2016.
- 30 Boyd, P. W. and Hutchins, D. A. Understanding the responses of ocean biota to a complex matrix of cumulative anthropogenic change, *Mar. Ecol. Prog. Ser.*, 470, 125-135, 2012.



- Bracegirdle, T. J., Connolley, W. M., and Turner, J.: Antarctic climate change over the twenty first century, *J. Geophys. Res.*, 113, *D03103*, doi:10.1029/2007JD008933, 2008.
- Bunse, C., Lundin, D., Karlsson, C. M. G., Vila-Costa, M., Palovaara, J., Akram, G., Svensson, L., Holmfeldt, K., González, J. M., Calvo, E., Pelejero, C., Marrasé, C., Dopson, M., Gasol, J. M., and Pinhassi, J.: Response of marine bacterioplankton pH homeostasis gene expression to elevated CO₂, *Nature Climate Change*, 2016.
- Coad, T., McMinn, A., Nomura, D., and Martin, A.: Effect of elevated CO₂ concentration on the photophysiology and growth of microalgae in Antarctic pack ice algal communities, *Deep Sea Research II*, 131, 160-169, 2016.
- Constable, A. J., Melbourne-Thomas, J., Corney, S. P., Arrigo, K. R., Barbraud, C., Barnes, D. K. A., Bindoff, N. L., Boyd, P. W., Brandt, A., Costa, D. P. et al. (2014). Climate change and Southern Ocean ecosystems I: how changes in physical habitats directly affect marine biota. *Global Change Biol.* 20, 3004-3025, 2014.
- Feng, Y., Hare, C. E., Rose, J. M., Handy, S. M., Ditullio, G. R., Lee, P. A., Smith, W. O., Peloquin, J., Tozzi, S., and Sun, J.: Interactive effects of iron, irradiance and CO₂ on Ross Sea phytoplankton. *Deep Sea Res. I*, 57, 368-383, 2010.
- Garrison, D. L., Ackley, S. F., and Buck, K. R.: A physical mechanism for establishing algal populations in frazil ice. *Nature* 306:363–365, 1983.
- Garrison, D. L.: Antarctic sea ice biota. *American Zoologist*, 31, 17-37, 1991.
- Geilfus, N-X., Tison, J-L., Ackley, S. F., Galley, R. J., Rysgaard S, Miller, L., Delille, B.: Sea ice pCO₂ dynamics and air-ice CO₂ fluxes during the Sea Ice Mass Balance in the Antarctic (SIMBA) experiment –Bellingshausen Sea, Antarctica. *Cryosphere* 8, 2395-2407, 2014.
- Gibson, J. A., and Trull, T. W. Annual cycle of fCO₂ under sea-ice and in open water in Prydz Bay, East Antarctica. *Mar. Chem.* 66, 187–200, 1991.
- Gleitz, M., Kukert, H., Riebesell, U., and Dieckmann, G. S.: Carbon acquisition and growth of Antarctic sea ice diatoms in closed bottle incubations, *Mar. Ecol. Prog. Ser.*, 135, 169-177, 1996.
- Hare, A. A., Wang, F., Barber, D., Geilfus, N-X., Galley, R. J., and Rysgaard, S.: pH evolution in sea ice grown at an outdoor experimental facility, *Mar. Chem.*, 154, 46-54, 2013.



- Heiden, J. P., Bischof, K., and Trimborn, S.: Light intensity modulates the response of two Antarctic diatom species to ocean acidification, *Front. Mar. Sci.* 3:260. doi: 10.3389/fmars.2016.00260, 2016.
- 5 Hoppe, C. J. M., Hassler, C. S., Payne, C. D., Tortell, P. D., Rost, B., and Trimborn, S.: Iron Limitation Modulates Ocean Acidification Effects on Southern Ocean Phytoplankton Communities, *PLoS One*, 8(11): e79890. doi:10.1371/journal.pone.0079890, 2013.
- Kapsenberg, L., Kelley, A. L., Shaw, E. C., Martz, T. R., and Hofmann, G. E. Near-shore Antarctic pH variability has
10 implications for the design of ocean acidification experiments. *Scientific Rep.*, 5:9638 doi:10.1038/srep09638, 2015.
- Lannuzel, D., Bowie, A. R., van der Merwe, P. C., Townsend, A. T., and Schoemann, V.: Distribution of dissolved and particulate metals in Antarctic sea ice. *Mar. Chem.*, 2011
- 15 Leonard, K. C., and Maksym, T.: The importance of wind-blown snow redistribution to snow accumulation on Bellingshausen Sea ice. *Annal. Glaciol.*, 52, 271-278, 2011.
- Lin, X., Huang, R., Li, Y., Wu, Y., Hutchins, D.A., Dai, M., and Gao, K.: Insignificant effects of elevated CO₂ on bacterioplankton community in a eutrophic coastal mesocosm experiment. *Biogeosciences*, doi:10.5194/bg-2017-10, 2017
20
- Martin A., McMinn A.: Sea ice, extremophiles and life in extra-terrestrial ocean worlds. *International Journal of Astrobiology*, doi.org/10.1017/S1473550416000483, 2017
- Martin, A., McMinn, A., Davey, S. K., Anderson, J. M., Miller, H. C., Hall, J. A., and Ryan, K. G.: Preliminary evidence for
25 the microbial loop in Antarctic sea ice using microcosm simulations. *Antarctic Sci.*, 24, 547-553, 2012.
- Matson, G., Washburn, L., Martz, T. R., Hofmann, G. E.: Abiotic versus biotic drivers of ocean pH variation under fast sea ice in McMurdo Sound, Antarctica, *PLoS ONE*, 9(9): e107239. doi:10.1371/journal.pone.0107239, 2014.
- 30 McMinn, A., Gradinger, R., and Nomura, D.: 3.8 Biochemical properties of Sea ice. *In* H. Eicken (Editor) *Handbook of Field techniques in sea ice research*. University of Alaska Press, USA, 2009.
- McMinn, A., Muller, M., Martin, A., and Ryan, K. G.: The response of Antarctic sea ice algae to changes in pH and CO₂, *PloS One* e86984. doi:10.1371/journal.pone.0086984, 2014.



- McMinn, A., Muller, M., Martin, A., and Ryan, K. G.: Effects of changing pH and CO₂ concentration on a late summer surface sea ice community. *Mar. Biol.*, 164, 87, 2017.
- 5 McMinn, A., Ryan, K. R., Ralph, P. J. and Pankowski, A.: Spring sea ice photosynthesis, primary productivity and biomass distribution in eastern Antarctica, 2002-2004. *Mar Biol.*, 151, 985-995, 2007,
- McMinn, A., Skerratt, J., Trull, T., Ashworth, C., and Lizotte, M.: Nutrient stress gradient in the bottom 5 cm of fast ice, McMurdo Sound, Antarctica, *Polar Biol.*, 21, 220-227, 1999.
- 10 Meiners, K. M., Vancoppenolle, M., Shanassekos, S., Dieckmann, D. S., Thomas, D. N., Tison, J-L., Arrigo K. R., Garrison, D., McMinn, A., Lannuzel, D., van der Merwe, P., Swadling, K. M., Smith Jr., W. O., Melnikov, I., Raymond, B.: Chlorophyll *a* in Antarctic sea ice from historical ice core data. *Geophys. Res. Lett.*, 39, 21602-21602, 2012.
- 15 Melnikov, I.A.: Recent sea ice ecosystem in the Arctic Ocean: a review. *In* Nihoul JCJ and Kostiano, AG (eds) Influence of climate change on the changing Arctic and sub-Arctic conditions. NATO Science for Peace and Security Series C- Environmental Security, 57-71, 2009.
- Miller, L.A., Papakyriakou, T. N., Collins, E. R., Deming, J. W., Ehn, J. K., Macdonald, R. W., Mucci, A., Owens, O., 20 Raudsepp, M., and Sutherland, N.: Carbon dynamics in sea ice: A winter flux time series. *J. Geophys. Res.*, 116, C02028, doi:10.1029/2009JC006058, 2011.
- Mitchell, C. and Beardall, J.: Inorganic carbon uptake by an Antarctic sea-ice diatom, *Nitzschia frigida*. *Polar Biol.*, 16, 95-99, 1996.
- 25 McNeil, B. I., Tagliabue, A., and Sweeney, C.: A multi-decadal delay in the onset of corrosive ‘acidified’ waters in the Ross Sea of Antarctica due to strong air-sea CO₂ disequilibrium. *Geophys. Res. Lett.*, 37, L19607, doi:10.1029/2010gl044597, 2010.
- 30 McNeil, B. I. and Matear, R. J.: Southern Ocean acidification: A tipping point at 450-ppm atmospheric CO₂. *Proc. Natl. Acad. Sci.*, 105, 18860–18864, doi:10.1073/pnas.0806318105, 2010.
- Mock, T., and Thomas, D. N.: Minireview: Recent advances in sea ice microbiology. *Environ. Microbiol.*, 7, 605-619, 2005.



- Oliver, A. E., Newbold, L. K., Whiteley, A. S., and van der Gast, C. J.: Marine bacterial communities are resistant to elevated carbon dioxide levels. *Environ. Microbiol. Rep.*, 6, 574-582, 2014.
- Pančić, M., Hansen, P. J., Tammilehto, A., and Lundholm, N.: Resilience to temperature and pH changes in a future climate change scenario in six strains of the polar diatom *Fragilariopsis cylindrus*. *Biogeosciences Disc.* 12, 4627-4556, 2015.
- Pankowski, A., and McMinn, A.: Ferredoxin and flavodoxin in eastern Antarctica pack ice. *Polar Biol.*, 31, 1153-1165, 2008.
- 10 Pankowski, A., and McMinn, A.: Iron availability regulates growth, photosynthesis and production of ferredoxin and flavodoxin in Antarctic sea ice diatoms, *Aquatic Biol.*, 4, 273-288, 2009.
- Papadimitriou, S., Thomas, D. N., Kennedy, H., Haas, C., Kuosa, H., Krell, A., and Dieckmann, G. S.: Biogeochemical composition of natural sea ice brines from the Weddell Sea during early austral summer, *Limnol. Oceanog.*, 52, 1809-1823, 15 2007.
- Raven, J. A., Giordano, M., Beardall, J., and Maberly, S.: Algal and aquatic plant carbon concentrating mechanisms in relation to environmental change, *Photosynth. Res.*, 109, 281-296, 2011.
- 20 Raymond, I. A., Sullivan, C. W., Devries, A. L.: Release of an ice-active substance by Antarctic sea-ice diatoms. *Polar Biol.*, 14, 71-77, 1994.
- Riebesell, U.: Effects of CO₂ enrichment on marine phytoplankton, *J. Oceanog.*, 60, 719-729, 2004.
- 25 Rost, B., Riebesell, U., Burkhardt, S., Sultemeyer, D.; 2003. Carbon acquisition of bloom forming marine phytoplankton. *Limnology and Oceanography*, 48, 55-67.
- Scott, P., McMinn, A., Hosie, G.: Physical parameters influencing diatom community structure in eastern Antarctic sea ice, 30 *Polar Biol.*, 14, 507-517, 1994.
- Shi, D., Xu, Y., Hopkinson, F., and Morel, M. M.: Effect of ocean acidification on iron availability to marine phytoplankton. *Science*, 327, 676-679, 2010.



- Søgaard, D. H., Hansen, P. J., Rysgaard, S., and Glud, R. N.: Growth limitation of three Arctic sea ice algal species: effects of salinity, pH, and inorganic carbon availability, *Polar Biol.*, 34, 1157-1165, 2011.
- Stoecker, D. K., Buck, K. R., and Putt, M.: Changes in the sea-ice brine community during the spring-summer transition, 5
McMurdo Sound, Antarctica. 1. Photosynthetic protists, *Mar. Ecol. Prog. Ser.*, 84, 265–278, 1992.
- Thomas, D. N. and Dieckmann, G. S.: *Sea Ice, an introduction to its physics, chemistry, biology and geology*. Blackwell, Oxford, UK, 2004.
- 10 Thomson, P. G., McMinn, A., Kiessling, I., Watson, M., and Goldsworthy, P. M.: Composition and succession of dinoflagellates and Chrysophytes in the upper fast ice of Davis Station, East Antarctica, *Polar Biol.*, 29, 337-345, 2006.
- Torstensson, A., Chierici, M., and Wulff, A.: The influence of increased temperature and carbon dioxide levels on the benthic/sea ice diatom *Navicula directa*, *Polar Biol.*, 35, 205-214, 2012.
- 15 Torstensson, A., Dinasquet, J., Chierici, M., Fransson, A., Riemann, L., and Wulff, A.: Physicochemical control of bacterial and protist community composition and diversity in Antarctic sea ice, *Environ. Microbiol.*, 17, 3869-3881, 2012.
- Torstensson, A., Hedblom, M., Bjork, M. M., Chierici, C., and Wulff, A.: Long-term acclimation to elevated $p\text{CO}_2$ alters carbon metabolism and reduces growth in the Antarctic diatom *Nitzschia lecoointei*, *Proc. R. Soc. B*, 282: 20151513.<http://dx.doi.org/10.1098/rspb.2015.1513>, 2015.
- 20 Tortell, P. D., Mills, M. M., Payne, C. D., Maldonado, M.T., Chierici, M., Fransson, A., Alderkamp, A. C., and Arrigo, K. R.: Inorganic C utilization and C isotope fractionation by pelagic and sea ice algal assemblages along the Antarctic continental shelf. *Mar. Ecol. Prog. Ser.*, 483, 47-66, 2010.
- 25 Trimborn, S., Breneis, T., Sweet, E., and Rost, B.: Sensitivity of Antarctic phytoplankton species to ocean acidification: Growth, carbon acquisition, and species interaction, *Limnol. Oceanog.*, 58, 997-1007, 2013.
- 30 Xu, D., Wang, Y., Fan, X., Wang, D., Ye, N., Zhang, X., Mou, S., Guan, Z., and Zhuang, Z. (2014b) Long-term experiment on physiological responses to synergetic effects of ocean acidification and photoperiod in the Antarctic sea ice algae *Chlamydomonas* sp. ICE-L, *Environ. Sci. Tech.*, 48, 7738-7746, 2014b



Xu, J., Gao, K., Li, Y., and Hutchins, D. A.: Physiological and biochemical responses of diatoms to projected ocean temperatures, *Mar. Ecol. Prog. Ser.*, 515, 73-81, 2014c

5 Xu, K., Fu, F., and Hutchins, D. A.: Comparative responses of two dominant Antarctic phytoplankton taxa to interactions between ocean acidification, warming, irradiance, and iron availability, *Limnol. Oceanog.*, 59, 1919-1931, 2014a



Figure Captions

Figure 1. Annual succession of nutrient concentration and pH in response to ice formation in winter and melting in spring.

On the bar at the bottom of the figure, blue represents open water and white represents the presence of ice.

5



Species/Community	Days	T°	CO ₂ (µatm)	Site	Author
<i>Navicula directa</i>	7	0.5-4.5°	380-960	Lab	Torstensson et al., 2012
Brine community	8	-2.5°	587-6066	<i>In situ</i>	McMinn et al., 2014
<i>Chlamydomonas</i> sp.	28	5°	390-1000	Lab	Xu et al. 2014
<i>Fragilariopsis cylindrus</i>	7	1-8°	pH 7.1-8.0	Lab	Panic et al., 2015
<i>Nitzschia lecointei</i>	194	-1.8°	280-960	Lab	Torstensson et al., 2015
Brine community	8	-5°	400-11300	Lab	Coad et al., 2016
<i>Fragilariopsis curta</i>	4	4°	180-1000	Lab	Heiden et al., 2016
Surface brine community	6	0°	45-4102	<i>In situ</i>	McMinn et al., 2017

Table 1. Ocean acidification experiments on sea ice microorganisms.




Figure 1.