

Soil water content drives spatio-temporal patterns of CO₂ and N₂O emissions from a Mediterranean riparian forest soil

Sílvia Poblador^{1*}, Anna Lupon^{1,2*}, Santiago Sabaté^{1,3}, Francesc Sabater^{1,3}

5 ¹ Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals (BEECA), Universitat de Barcelona. Av. Diagonal 643, 08028, Barcelona, Spain.

² Department of Forest Ecology and Management, Swedish University of Agricultural Sciences (SLU). Skogsmarksgränd 17S, 90183, Umeå, Sweden.

³ CREAM. Campus de Bellaterra Edifici C, 08193, Cerdanyola del Vallès, Spain.

10

* These authors contributed equally to the development of this work.

Correspondence to: Sílvia Poblador (spoblador@ub.edu)

Abstract.

15 Riparian zones play a fundamental role in regulating the amount of carbon (C) and nitrogen (N) that is exported from catchments. However, C and N removal via soil gaseous pathways can influence local budgets of greenhouse gases (GHG) emissions and contribute to climate change. Over a year, we quantified soil effluxes of carbon dioxide (CO₂) and nitrous oxide (N₂O) from a Mediterranean riparian forest in order to understand the role of these ecosystems on catchment GHG emissions. In addition, we evaluated the main soil microbial processes that produce GHG (mineralization, nitrification, and denitrification) and how changes in soil properties can modify the GHG production over time and space. Mediterranean riparian soils emitted larger amounts of CO₂ (1.2 – 10 g C m⁻² d⁻¹) than N₂O (0.001 – 0.2 mg N m⁻² d⁻¹) to the atmosphere attributed to high respiration and low denitrification rates. Both CO₂ and N₂O emissions showed a marked (but antagonistic) spatial gradient as a result of variations in soil water content across the riparian zone. Deep groundwater tables fueled large soil CO₂ effluxes near the hillslope, while N₂O emissions were higher in the wet zones adjacent to the stream channel. However, both CO₂ and N₂O emissions peaked after spring rewetting events, when optimal conditions of soil water content, temperature, and N availability favor microbial respiration, nitrification, and denitrification. Overall, our results highlight the role of water availability on riparian soil biogeochemistry and GHG emissions and suggest that climate change alterations in hydrologic regimes can affect the microbial processes that produce GHG as well as the contribution of these systems to regional and global biogeochemical cycles.

30 Keywords.

greenhouse gas emissions, riparian soils, denitrification, microbial respiration, soil water content.

1 Introduction

35 Riparian zones are hotspots of nitrogen (N) transformations across the landscape, providing a natural filter for nitrate (NO_3^-)
transported from surrounding lands via runoff and subsurface flow paths (Hill, 1996; Vidon et al., 2010). Although interest in
riparian zones has primarily been motivated by the benefits of these ecotones as effective N sinks, enhanced microbial activity
in riparian landscapes can play a key role on atmospheric pollution. For instance, riparian zones can account by 70% of global
(natural processes and human activities) terrestrial emissions of nitrous oxide (N_2O) to the atmosphere, a powerful greenhouse
gas (GHG) with 298 times the global warming potential of carbon dioxide (CO_2) (Audet et al., 2014; Groffman et al., 2000;
40 Hefting et al., 2003). Moreover, riparian soils can significantly contribute to global CO_2 emissions because they can hold high
rates of heterotrophic and autotrophic respiration (Chang et al., 2014). Soil respiration is the main natural carbon (C) efflux to
the atmosphere, contributing to 20% of the global emission of CO_2 (Kim and Verma, 1990; Raich et al., 2002; Rastogi et al.,
2002). Finally, riparian zones can support large methane (CH_4) fluxes that account for the 15 – 40 % of global emissions
(Audet et al., 2014; Segers, 1998). However, there are still many uncertainties regarding the magnitude and spatio-temporal
45 variability of soils GHG emissions in riparian zones, reaching contradictory results concerning the potential role of riparian
zones as sinks or sources of C and N (Bruland et al., 2006; Groffman et al., 1992; Harms et al., 2009; Walker et al., 2002).

Understanding the processes regulating GHG emissions from riparian soils is essential to quantify the role of riparian zones in
the global C and N cycles. Multiple environmental variables, such as soil temperature, soil water content, and both C and N
availability have been identified as key factors influencing the rate and variability of soil microbial activities that produce
GHG (Chang et al., 2014; Hefting et al., 2003; Mander et al., 2008; McGlynn and Seibert, 2003). Among them, riparian
50 hydrology seems to play a fundamental role on GHG production because it controls the substrate subsidies and, most
importantly, the redox conditions of riparian soils (Jacinthe et al., 2015; Vidon, 2017). Under saturated conditions, anaerobic
processes such as methanogenesis (i.e. the transformation of CO_2 to CH_4) and denitrification (i.e. the transformation of NO_3^-
to N gas (N_2) or N_2O) are the primary processes involved in the C and N cycles (Clément et al., 2002). Conversely, in dry
soils, aerobic transformations involved in the oxidation of the organic matter (i.e. respiration, mineralization, nitrification,
55 methane oxidation) dominate the riparian biogeochemistry (Harms and Grimm, 2008). From such observations, some one
would expect that there is a strong correlation between soil wetness and the relative importance of CO_2 , N_2O and CH_4 riparian
soil emissions to the total GHG fluxes. However, there are still relatively few studies that analyze the direct influence of soil
water content on several GHG effluxes simultaneously (but see Harms and Grimm, 2008; Jacinthe et al., 2015), and even less
that combine such analyses with other environmental factors and soil processes. Thus, it is still unclear under which
60 circumstances soil water content (rather than temperature or substrate availability) is the primary control factor of the riparian
functionality.

Mediterranean systems are a unique natural laboratory to understand the close link between spatio-temporal variations in
hydrology and riparian biogeochemistry because they are characterized by a marked spatial gradient of soil water content, that

65 can range from <10% in the hillslope edge to > 80% close to the stream (Chang et al., 2014; Lupon et al., 2016). Moreover, Mediterranean regions are subjected to seasonal alterations of precipitation and temperature regimes that might affect riparian hydrology as well as microbial activity in the riparian soils (Bernal et al., 2007; Bruland et al., 2006; Harms and Grimm, 2008; Harms et al., 2009). Increments in GHG emissions in riparian zones might occur following storms or flood events because sharp increments in soil water content enhance nitrification, denitrification, respiration, and methanogenesis rates (Casals et al., 2011; Jacinthe et al., 2015; Werner et al., 2014). However, because recent studies have shown that high temperatures and relatively moist soils can sustain large rates of C respiration and N mineralization in summer in the near-stream zone (Chang et al., 2014; Lupon et al., 2016), the contribution of such microbial pulses to annual CO₂ and N₂O production in Mediterranean riparian soils is still under debate. Moreover, improved understanding of interactions among hydrology, microbial processes, and gas emissions within Mediterranean riparian zones is not only fundamental to understand the temporal pattern of riparian biogeochemistry, but also necessary to estimate the contribution of these ecosystems to atmospheric GHG budgets at local and global scale.

80 In this study, soil properties, soil N processes, and CO₂ and N₂O soil emissions were measured over a year across a Mediterranean riparian forest that exhibited a strong gradient in soil water content (Fig. 1a). We did not measure CH₄ emissions because previous studies reported extremely low values in dry systems (-0.06 – 0.42 mg C m⁻² d⁻¹; Batson et al., 2015; Gómez-Gener et al., 2015). Specifically, we aimed (i) to evaluate the spatio-temporal patterns of CO₂ and N₂O emissions in Mediterranean riparian soils, (ii) to analyze under which conditions soil water content rules microbial processes and GHG over other physicochemical variables, and (iii) to provide some reliable estimates of GHG emissions from Mediterranean riparian soils. We hypothesized that the magnitude and the relative contribution of N₂O and CO₂ to total GHG emission strongly depend on soil water content and redox conditions rather than other variables during all year long (see conceptual approach in Fig. 1b). In the near-stream zone, we expected that saturated anoxic soils would enhance denitrification but constrain both respiration and nitrification. Thus, we predicted higher N₂O than CO₂ emissions in this zone. In the intermediate zone, we expected that wet (but not saturated) soils would enhance aerobic processes such as respiration, N mineralization or nitrification, and thus, we predicted high CO₂ emissions compared to N₂O. Finally, we expect that dry soils would deplete (or even inhibit) the soil microbial activity near the hillslope edge, and therefore, we predicted low GHG emissions in this zone. Because Mediterranean regions are subjected to strong intra-annual variations in soil water content, we expected that this general behavior would be maximized in summer, when only near-stream soils would keep wet. Conversely, we expected that all microbial processes would be enhanced short-after rainfall events, and thus, simultaneous pulses of CO₂ and N₂O emissions would occur in spring and fall.

2 Materials and methods

2.1 Study site

95 The research was conducted in a riparian forest of Font del Regàs, a forested headwater catchment (14.2 km², 500 – 1500 m above the sea level (a.s.l.)) located in the Montseny Natural Park, NE Spain (41°50'N, 2°30'E) (Fig. 1a). The climate is sub-humid Mediterranean; with mean temperature ranging from 5°C in February to 25°C in August. In 2013, annual precipitation (1020 mm) was higher than long-term average (925 ± 151 mm), with most of rain falling in spring (500 mm) (Fig. 2a). Total inorganic N deposition oscillates between 15 – 30 kg N ha⁻¹ yr⁻¹ (period 1983 – 2007; Àvila and Rodà, 2012).

100 We selected a riparian site (~600 m², ~30 m wide) that flanked a 3rd order stream close to the catchment outlet (536 m a.s.l., 5.3 km from headwaters). The riparian site was divided into three zones characterized by different species compositions (Fig. 1a). The near-stream zone was located adjacent to the stream (0 – 4 m from the stream edge) and was composed of *Alnus glutinosa* (45% of basal area) and *Populus nigra* (33% of basal area). The intermediate zone (4 – 7 m from the stream edge) was composed by *P. nigra* and *Robinia pseudoacacia* (29% and 71% of basal area respectively). Finally, the hillslope zone (7
105 – 30 m from the stream edge) bordered upland forests and was composed by *R. pseudoacacia* (93% of basal area) and *Fraxinus excelsior* (7% of basal area). The three riparian zones had sandy-loam soils (bulk density = 0.9 – 1.1 g cm⁻³), with a 5-cm deep organic layer followed by a 30-cm deep A-horizon. The top soil layer (0 – 10 cm depth) was mainly composed by sands (~90%) and silts (~7%) at the near-stream zone, whereas gravels (~16%) and sands (~80%) were the dominant particle sizes at the intermediate and hillslope zones. During the study period, groundwater level averaged -54 ± 14 cm below the soil surface (b.s.s.) at the near-stream zone, and decreased to -125 ± 4 and -358 ± 26 cm b.s.s. at the intermediate and hillslope zones, respectively (Fig. 1a and Fig. 2b).

2.2 Field sampling

We delimited five plots (1 x 1 m) within each riparian zone (near-stream, intermediate and hillslope) (Fig. 1a). During the year 2013, soil physicochemical properties, soil N processes, and gas emissions were measured in each plot every 2 – 3 months in
115 order to cover a wide range of soil water content and temperature conditions. On each sampling month, one soil sample (0 – 10 cm depth, including O- and A- horizons) was collected randomly from each plot to analyze soil physicochemical properties. Soil samples were taken with a 5-cm diameter core sampler and placed gently into plastic bags after carefully removing the litter layer. Close to each soil sample, we performed *in situ* soil incubations to measure soil net N mineralization and net nitrification rates (Eno, 1960). For this purpose, a second soil core (0 – 10 cm depth) was taken, placed in a polyethylene bag, and buried
120 at the same depth. Soil incubations were buried 4 days and then removed from the soil.

Gas emissions and denitrification rates were measured simultaneously and during four consecutive days (i.e. during the entire soil incubation period) in order to facilitate the direct comparison between microbial rates and gas fluxes. Soil CO₂ effluxes

were measured with a SRC-1 soil chamber attached to an EGM-4 portable infrared gas analyzer (IRGA) (PP Systems, Amesbury, MA). The EGM-4 has a measurement range of 0 – 2000 ppm ($\mu\text{mol mol}^{-1}$), with an accuracy of 1% and a linearity of 1% throughout the range. Every field day, CO_2 measurements started at 12 p.m. and were conducted consecutively at the 15 plots starting for the near stream zone. At each plot, the SCR-1 soil chamber was placed over the top soil for a 120 s incubation. Before each measurement, we carefully removed the litter layer to ensure no leaks. Furthermore, we aerated the SCR-1 between samples to ensure the accuracy of the instrument as well as to avoid contamination between samples. For each plot, CO_2 emissions rates were calculated from the best fit linear regression of the CO_2 accumulated in the head-space with incubation time (Fig. S1). CO_2 fluxes on an areal basis (F_{CO_2} , in $\mu\text{mol m}^{-2} \text{h}^{-1}$) were calculated following Healy et al. (1996):

$$F_g = \frac{dg}{dt} \times \frac{V P_0}{S R T_0} \quad (\text{Eq.1})$$

where dg/dt is the rate of change in gas concentration (in $\mu\text{mol mol}^{-1} \text{h}^{-1}$) in the chamber, V is chamber volume (in m^3), P_0 is initial pressure (in Pa), S is the soil surface area (in m^2), R is the gas constant ($8.314 \text{ Pa m}^3 \text{ K}^{-1} \text{ mol}^{-1}$), and T_0 is the initial chamber temperature (in $^\circ\text{K}$). For budgeting, moles of CO_2 and N_2O were converted to grams of C and N, respectively.

In situ denitrification rates and N_2O emissions were measured using closed cylinder (0.37 L) and open cylinder (0.314 m^2) chambers, respectively. For denitrification analyses, an intact soil core (0 – 10 cm depth) was introduced in the chamber, closed with a rubber serum stopper, amended with acetone-free acetylene to inhibit the transformation of N_2O to N_2 (10% v/v atmosphere), and placed at the same depth. For N_2O analysis, chambers were placed directly on the soil and no special treatment was carried out. Gas samples for both denitrification and N_2O chambers were taken at the same time (0h, 1h, 2h, and 4h of incubation) with a 20-mL syringe and stored in evacuated tubes. All soil and gas samples were kept at $< 4^\circ\text{C}$ until laboratory analysis (< 24 h after collection).

Soil physical properties were measured within each plot simultaneously to gas emissions. Volumetric soil water content (%) (5 replicates per plot) and soil temperature ($^\circ\text{C}$) (1 replicate per plot) were measured at 10-cm depth by using a time-domain reflectometer sensor (HH2 Delta-T Devices Moisture Meter) and a temperature sensor (CRISON 25), respectively. Soil pH and reduction potential (Eh, mV) (1 replicate per plot) were measured at 0 – 10 cm depth by water extraction (1:2.5 v/v) using a Thermo-Scientific ORION sensor (STAR 9107BNMD). Although Eh measures performed by water extraction may not be as accurate as other field techniques, these values have been previously used as a good proxy of the soil redox potential (Yu and Rinklebe, 2013).

2.3 Laboratory analyses

Pre-incubation soil samples were oven dried at 60°C , sieved, and the fraction < 2 mm was used for measuring soil chemical properties. The relative soil organic matter content (%) was measured by loss on ignition (450°C , 4 h). Total soil C and N

contents were determined on a gas chromatograph coupled to a TCD detector after combustion at 1000°C at the Scientific Technical Service of the University of Barcelona.

155 To estimate microbial N processes, we extracted 5 g of pre- and post- incubation field-moist soil samples with 50 ml of 2 M KCl (1g : 10ml, ww : v; 1 h shaking at 110 r.p.m. and 20°C). The supernatant was filtered (Whatman GF/F 0.7 µm pore diameter) and analyzed for ammonium (NH₄⁺) and nitrate (NO₃⁻). NH₄⁺ was analyzed by the salicylate-nitroprusside method (Baethgen and Alley, 1989) using a spectrophotometer (PharmaSpec UV-1700, SHIMADZU). NO₃⁻ was analyzed by the cadmium reduction method (Keeney and Nelson, 1982) using a Technicon Autoanalyzer (Technicon, 1987). For each pair of samples, net N mineralization and net nitrification were calculated as the differences between pre- and post-incubations values
160 of inorganic N (NH₄⁺ and NO₃⁻) and NO₃⁻, respectively (Eno, 1960). Pre-incubation NH₄⁺ and NO₃⁻ concentrations were further used to calculate the availability of dissolved inorganic nitrogen in riparian soils.

To estimate denitrification and natural N₂O emissions, we analyzed the N₂O of all gas samples using a gas chromatograph (Agilent Technologies, 7820A GC System) that was calibrated using certified standards (4.66 ppm N₂O; , AirLiquide). Both denitrification and N₂O emissions rates were calculated similarly to CO₂ fluxes (Fig. S1). In addition, we measured the
165 denitrification enzyme activity (DEA) for 3 soil cores of each riparian zone to determine the factors limiting denitrification. For each soil core, four sub-samples (20 g of fresh soil) were placed into 125-ml glass jars containing different treatments. The first jar (DEA_{MQ}) contained Milli-Q water (20 ml) to test anaerobiosis limitation. The second jar (DEA_C) was amended with glucose solution (4 g glucose kg soil⁻¹) to test C limitation. The third jar (DEA_{NO₃}) was amended with nitrate solution (72.22mg KNO₃ kg soil⁻¹) to test N limitation. Finally, the fourth jar (DEA_{C+NO₃}) was amended with both nitrate and glucose solutions
170 (4 g glucose kg soil⁻¹ and 72.22mg KNO₃ kg soil⁻¹) to test simultaneously C and N limitation. All jars were capped with rubber serum stoppers, made anaerobic by flushing N₂, and amended with acetone-free acetylene (10% v/v) (Smith and Tiedje, 1979). Gas samples were collected after 4 h and 8 h of incubation and analyzed following the same procedure of field DNT samples. DEA rates were calculated similarly to denitrification rates.

2.4 Statistical analysis

175 Statistical analyses were carried out using the package *lmer* and *pls* of R 2.15.1 statistical software (R Core Team, 2012). We performed linear mixed-model analysis of variance (ANOVA) to test differences in soil properties, microbial N processes, and gas emissions across riparian zones and seasons. We used riparian zone and season as fixed effects, and plot (nested within riparian zones) as a random effect. When multiple samples were taken within a plot (soil physical properties, denitrification, and gas emissions), the ANOVA was performed on plot means, with n = 75 (5 plots x 3 zones x 5 dates). For each model,
180 post-hoc Tukey contrasts were used to test which zones or seasons differed from each other. In all cases, residuals were tested for normality using a Shapiro-Wilk test, and homogeneity of variance was examined visually by plotting the predicted and residual values. In those cases that the normality assumption was unmet, data was log transformed. In all analyses, differences were considered significant when p < 0.05.

185 We used partial least squares regression (PLS) to explore how soil properties, C and N availability, groundwater level, and
soil N processes predict variation in CO₂ and N₂O emissions. PLS identifies the relationship between independent (X) and
dependent (Y) data matrices through a linear, multivariate model; and produces latent variables (PLS components) representing
the combination of X variables that best describe the distribution of observations in ‘Y space’ (Eriksson et al., 2006). We
determined the goodness of fit (R²Y) and the predictive ability (Q²Y) of the model by comparing modeled and actual Y
190 observations through a cross-validation process. Each model was refined by iteratively removing variables that had non-
significant coefficients in order to minimize the model overfitting (i.e. low Q²Y values) as well as the multicollinearity of the
explanatory variables (i.e. variance inflation factor (VIF) < 5). Furthermore, we identified the importance of each X variable
by using variable importance on the projection (VIP) scores, calculated as the sum of square of the PLS weights across all
components. VIP values > 1 indicate variables that are most important to the overall model (Eriksson et al., 2006). In all PLS
models, data was ranked and centered prior analysis.

195 3 Results

3.1 Spatial pattern of soil properties, microbial rates, and gas emissions

During the study period, all riparian zones had similar mean soil temperature (11 – 12°C), pH (6 – 7) and redox potential (170
– 185 mV) (Table 1). However, soil water content exhibited strong differences across riparian zones (Table 2), with the near-
stream zone holding wetter soils than the intermediate and the hillslope zones (Table 1). There were significant differences in
200 most of soil chemical properties (Table 1, Table 2). Both organic matter and soil C and N content were 2-fold lower in the
near-stream zone than in the intermediate and hillslope zones, though all zones exhibited similar C:N ratios (CN = 14).
Moreover, inorganic N concentrations (NH₄⁺ and NO₃⁻) were from 2- to 5-fold lower for the near-stream zone than for the
other two zones.

On annual basis, net N mineralization averaged 0.14 ± 0.40 , 0.39 ± 1.23 , and 0.22 ± 1.03 mg N kg⁻¹ d⁻¹ at the near-stream,
205 intermediate, and hillslope zones, respectively. Mean annual net nitrification rates were close to net N mineralization,
averaging 0.17 ± 0.38 , 0.25 ± 0.69 , and 0.28 ± 0.73 mg N kg⁻¹ d⁻¹ at the near-stream, intermediate, and hillslope zones,
respectively. There were no significant differences in mean annual net N mineralization and net nitrification rates among
riparian zones (in both cases: mixed-model ANOVA test, $F > F_{0.05}$, $p > 0.05$). Mean annual denitrification was higher at the
near-stream zone (2.69 ± 5.30 mg N kg⁻¹ d⁻¹) than at the intermediate (0.72 ± 1.85 mg N kg⁻¹ d⁻¹) and hillslope (0.76 ± 1.59 mg
210 N kg⁻¹ d⁻¹) zones (mixed-model ANOVA test, $F = 4.33$, $p = 0.038$). However, potential denitrification rates were lower in the
near-stream zone (0.3 – 0.6 mg N kg⁻¹ d⁻¹) compared to intermediate (1.0 – 2.4 mg N kg⁻¹ d⁻¹) and hillslope (1.3 – 3.8 mg N
kg⁻¹ d⁻¹) zones (Table 3).

Natural CO₂ and N₂O emissions differed among riparian zones, yet they showed opposite spatial patterns. Near-stream zone
exhibited lower CO₂ emissions (318 ± 195 mg C m⁻² h⁻¹) compared to the intermediate (472 ± 298 mg C m⁻² h⁻¹) and hillslope

215 (458 ± 308 mg C m⁻² h⁻¹) zones (mixed-model ANOVA test, F = 7.08, p = 0.009). Conversely, near-stream zone showed higher
N₂O emissions (0.035 ± 0.022 mg N m⁻² h⁻¹) than the other two zones (intermediate = 0.032 ± 0.025 mg N m⁻² h⁻¹; hillslope =
0.022 ± 0.012 mg N m⁻² h⁻¹) (mixed-model ANOVA test, F = 7.31, p = 0.008).

3.2 Temporal pattern of soil properties, microbial rates, and gas emissions

220 During the study period, there was a marked seasonality in most of soil physical properties, except for pH and Eh, which did
not show any temporal pattern (Table 2). Soil water content exhibited a marked seasonality, though it differed among riparian
zones (Table 2, “zone x season”). In the intermediate and hillslope zones, soil water content was maxima in November and
minima in August, while the near-stream soils were wetter during both spring (April-June) and autumn (November) (Fig. 3a).
Conversely, soil temperature showed similar seasonality but opposite values in all riparian zones (Table 2), with a maxima in
225 summer (August) and minima in winter (February) (Fig. 3b). Soil chemical properties (soil organic matter and both soil C and
N content) did not show any seasonal trend, but all riparian zones exhibited lower C:N ratios in February compared to the
other seasons (Fig. 3c). There was no seasonality in soil NH₄⁺ concentrations at any riparian zone (Table 2). However, soil
NO₃⁻ concentrations showed a marked temporal pattern, yet it differed among riparian zones (Table 2, “zone x season”). The
highest soil NO₃⁻ concentrations occurred in February at both the near-stream and hillslope zones, but in June-August at the
intermediate zone (Fig. 3d).

230 Soil N processes showed similar seasonal patterns in all riparian zones (in all cases: F_{date} < F_{0.05}, F_{interaction} > F_{0.05}). Both net N
mineralization and net nitrification rates were higher in April than February, June, and November (Fig. 4a and 4b), while
denitrification rates were higher in April and June compared to the rest of the year (Fig. 4c). In April, both net N mineralization
and net nitrification rates differed across riparian zone, with higher rates in the intermediate zone than in the near-stream one.
Net N mineralization rates also differed in August, when the intermediate zone exhibited 2-fold higher rates than the other two
235 zones. Finally, denitrification was higher at the near-stream than at the other two zones in both June and August.

Natural gas emissions showed a clear seasonal pattern (in both cases: mixed-model ANOVA test, F_{date} < F_{0.05}, p < 0.001), yet
it differed between CO₂ and N₂O emissions. In all zones, CO₂ emissions were maxima in June and minima in February (Fig.
5a), while highest N₂O emission rates occurred in April and lowest in both February and August (Fig. 5b). In spring (April
and June), CO₂ emissions were higher at the intermediate and hillslope zones compared to the near-stream one (Fig. 5a).
240 Moreover, the near-stream zone showed higher N₂O emissions than the hillslope zone in February, April, and June (Fig. 5b).

3.3 Relationship between soil properties, microbial processes, and gas emissions

PLS models extracted two components that explained the 71% and the 40% of the variance in CO₂ and N₂O emissions,
respectively (Table 4). The model predictability was high for CO₂ (Q²Y = 0.66), but weak for N₂O (Q²Y = 0.34). Moreover,
PLS models identified few variables as key predictors of GHG emissions (VIF < 2, VIP > 0.8), yet these variables differed
245 between CO₂ and N₂O emissions (Table 4). Soil temperature (PLS coefficient [coef] = +0.60), and soil water content (coef =

-0.24) explained most of the variation in CO₂ emissions (Table 4, Fig. S2a). Conversely, variations in N₂O emissions were primarily related to changes in denitrification rates (coef = +0.45), soil water content (coef = +0.21) and, to less extent, groundwater level (coef = -0.16) (Table 4, Fig. S2b).

250 **4 Discussion**

This study emphasized the role of soil water content as a main driver of riparian biogeochemistry and GHG emissions. By analyzing soil microbial processes and GHG emissions over a year in a Mediterranean riparian forest, we clearly demonstrate that soil water content has a major role in driving soil microbial processes, the spatio-temporal patterns of CO₂ and N₂O emissions and the overall role of Mediterranean riparian soils in the global C and N cycles.

255 **4.1 Microbial processes regulating GHG emissions**

Mean daily emissions of CO₂ found in the present study (1.2 – 10 g C m⁻² d⁻¹) were generally high, especially during spring and summer months. These soil CO₂ emissions were higher than those reported for temperate riparian regions (0.2 – 4.8 g C m⁻² d⁻¹; Batson et al., 2015; Bond-Lamberty and Thomson, 2010; Mander et al., 2008), although similar values have been reported in some dry forested wetlands of Europe and North America (Harms and Grimm, 2008; Oertel et al., 2016). These
260 substantially high CO₂ emissions observed in Font del Regàs may be attributed to high microbial respiration rates associated with relatively moist and organic matter enriched soils (Mitsch and Gosselink, 2007; Pacific et al., 2008; Stern, 2006). In agreement, previous studies have reported that microbial heterotrophic respiration can be an important contributor (> 60%) to CO₂ soil effluxes in water-limited riparian zones (Harms and Grimm, 2012; McLain and Martens, 2006). However, the absence
265 of a relationship between soil N processes and CO₂ emissions suggests that soil C and N cycles are decoupled in Mediterranean riparian forests, and thus, soil N mineralization may be not a good descriptor of bulk organic matter mineralization. Moreover, plant roots respiration and methane oxidation can increase the CO₂ emissions in riparian soils with deep groundwater tables such as in Font del Regàs (Chang et al., 2014).

Conversely, N₂O emissions of our riparian site (0.001 – 0.2 mg N m⁻² d⁻¹) were relatively low during the whole year. Similar N₂O emissions were reported in other water-limited riparian forests that are rarely flooded (-0.9 – 0.39 mg m⁻² d⁻¹; Bernal et al., 2003; Harms and Grimm, 2012; Vidon et al., 2016), yet these values were, on average, much lower than those found in
270 temperate riparian regions (0 – 54 mg N m⁻² d⁻¹; Burgin and Groffman, 2012; Hefting et al., 2003; Mander et al., 2008). In Font del Regàs, most N₂O was produced by denitrification, as we found an intimate link between this microbial process and N₂O emissions. Additionally, other processes such as nitrification or nitrate ammonification can contribute to N₂O emissions (Baggs, 2008; Hefting et al., 2003). However, it seems unlikely that nitrification could account for the observed N₂O emissions
275 because no relationship was found between net nitrification rates and N₂O emissions. Likewise relatively oxic conditions (Eh > 100) and low C:N ratios (C:N < 20) in Font del Regàs suggest low nitrate ammonification in riparian soils (Schmidt et al.,

2011). Currently, the influence of soil denitrification on N₂O emissions in riparian zones is still under debate (Giles et al., 2012). Nonetheless, our results suggest that performing simultaneous measurements of different soil N can contribute to disentangling the mechanisms underlying net N₂O emissions in riparian areas.

280 **4.2 Effects of soil water content on soil CO₂ effluxes**

As expected, we found higher soil CO₂ effluxes at the intermediate and hillslope zones than at the near-stream zone. This spatial pattern was negative and strongly related to soil water content (Table 4), suggesting that, as soils become less moist and more aerated, oxidizing aerobic respiration increases, ultimately stimulating CO₂ production in the top soil layer (Muller et al., 2015). In agreement, other aerobic processes, such as N mineralization were also higher in the intermediate and hillslope
285 zones. Moreover, deep groundwater tables in the hillslope zone can increase the volume of aerated soil, which can increase the area-specific soil CO₂ emissions near the hillslope edge (Chang et al., 2014). Increasing CO₂ emissions from wet to dry zones has been reported in other wetlands and riparian forests (Batson et al., 2015; Morse et al., 2012; Welti et al., 2012), pinpointing a close linkage between riparian hydrology and spatial variations in microbial respiration rates..

Nonetheless, the intra-annual variations of soil CO₂ emissions were strongly dependent on soil temperature (Table 4). Probably,
290 cold temperatures (< 4°C) limited soil respiration during winter, while warmer conditions (> 15°C) stimulated this process in June and August (Emmett et al., 2004; Suseela et al., 2012; Teiter and Mander, 2005). However, lower CO₂ emissions than expected for temperature dynamics were reported in summer at the intermediate and hillslope zones, likely because extreme soil dryness (soil water content < 20%) limited respiration rates during such period (Chang et al., 2014; Goulden et al., 2004; Wickland et al., 2010). Although the mechanisms by which soil dryness may affect microbial C demand are still poorly
295 understood, suppressed microbial respiration in summer can be attributed to a disconnection between microbes and resources (Belnap et al., 2005; Davidson et al., 2006), decreases in photosynthetic and exo-enzymatic activities (Stark and Firestone, 1995; Williams et al., 2000), or a relocation of the invested energy on growth (Allison et al., 2010). Altogether, these results suggest that soil water content may be as important as soil temperature to understand soil CO₂ effluxes, and therefore, future warmer conditions may not fuel higher CO₂ emissions, at least in those regions experiencing severe water limitation.

300 **4.3 Effects of soil water content on soil N₂O effluxes**

As occurred for CO₂ emissions, N₂O fluxes showed a clear spatial pattern associated with changes in soil water content across the riparian zone. In the near-stream zone, relatively wet conditions (SWC = 30 – 40%) likely promoted denitrification rates, while dry soils (SWC = 10 – 25%) could limit both nitrification and denitrification in the intermediate and hillslope zones
305 (Linn and Doran, 1984; Pinay et al., 2007). Such spatial pattern differed from those found in non-water limited riparian forests, where higher N₂O emissions occurred in the hillslope edge as a result of high resource supply (DeSimone et al., 2010; Dhondt et al., 2004; Hedin et al., 1998). These results suggest that riparian hydrology is the primary mechanisms controlling denitrification but, once water is unlimited, substrate availability controls the magnitude of denitrification rates. This former idea is supported by our potential denitrification results, which showed that, after adding water, denitrification rates were

310 similar to those observed in the field for the near-stream zone, but increase by 3-4 fold in the other two zones. Moreover, N₂:N₂O ratios estimated from acetylene method suggest that there was a spatial pattern in denitrification efficiency as well. During the study period, N₂:N₂O ratios were always higher at the near-stream (21.50 ± 40.32) than at the intermediate and hillslope zones (5.90 ± 16.02 and 4.23 ± 8.31, respectively), yet all values were much lower than those reported for temperate riparian forests (184 – 844; Mander et al., 2014). All together, these results support the idea that saturated soils favored the complete denitrification process to N₂ and can potentially emit less N₂O compared to less saturated soils (Giles et al., 2012).

315 Intra-annual variation in N₂O emission was also related to riparian hydrology because high rates of N₂O effluxes occurred in April, when large precipitation events (400 mm) raised the groundwater level and increased soil water content at the whole riparian plot. Such pulses of N₂O emissions short-after rewetting events can reflect the microbial use of the NO₃⁻ that has been accumulated during dry antecedent periods (Chang et al., 2014; Hefting et al., 2004; Pinay et al., 2007). In agreement, the PLS model showed a negative relation between soil water content and NO₃⁻ concentrations. Moreover, our results further suggest
320 that rewetting events promote a fast N cycle because all microbial N processes were maxima in April. Nevertheless, we also expected a fast N cycle as well as large N₂O emissions following rains in November because, similarly to spring, environmental conditions (i.e. high soil water content and increments in soil NO₃⁻ concentrations during the antecedent dry summer) should enhance microbial activity. Likely, low rates of N transformations during fall may be attributed to an increase in microbial N demand following large C inputs from litterfall (Guckland et al., 2010). Moreover, leaf litter from *R. pseudoacacia*, the main tree species in our study site, holds a high lignin content (Castro-Díez et al., 2009; Yavitt et al., 1997), which might enrich the
325 riparian soil with phenolic compounds and ultimately limit the use of N by microbes (Bardon et al., 2014). These results suggest that the response of N cycling to changes in water availability is more complex and less predictable than C cycling, likely because N processes depend on the interplay of additional ecosystem factors not included in this study.

4.4 Riparian soils as hot spots of GHG effluxes

330 There are several studies that attempt to upscale riparian GHG emissions at catchment scale, yet there are still fundamental uncertainties regarding the magnitude and sources of GHG emissions (Hagedorn, 2010; Pinay et al., 2015; Vidon and Hill, 2006). When accounting for all GHG (CO₂ + N₂O), our study suggest that our riparian soils can emit between 438 – 3650 g C m⁻² yr⁻¹. Assuming that GHG emissions (CO₂ + N₂O) from upland evergreen oak and beech soils (54% and 38% of the catchment, respectively) are similar to other Mediterranean regions (oak: 19 – 1240 g C m⁻² yr⁻¹; Asensio et al., 2007; Barba et al., 2016; Inclán et al., 2014); beech: 214 – 1182 g C m⁻² yr⁻¹; Guidolotti et al., 2013; Kesik et al., 2005), then riparian soils
335 (6% of the catchment area) can contribute between 16 – 22% to the total catchment soil GHG emissions. Although these estimates are rough (i.e. we assumed that riparian soils emit the same rate of GHG that our study site), our results clearly pinpoint that riparian soils can be potential hot spots of GHG emissions within Mediterranean catchments. These findings contrast with the common knowledge that water limited soils are powerless GHG sources to the atmosphere (Bernal et al.,

340 2007; Vidon et al., 2016) and stress the importance of simultaneously consider several GHG emissions (i.e. CO₂, N₂O, CH₄) to get a whole picture of the role of riparian soils in climate change.

5 Conclusions

345 Mediterranean riparian zones are dynamic systems that undergo spatial and temporal shifts in biogeochemical processes due to changes in both soil water content and substrate availability. In a first attempt to simultaneously quantify CO₂ and N₂O emissions from Mediterranean riparian soils, we showed that most of GHG emissions occur in form of CO₂, even in the wet soils located near the stream. In addition, our results clearly illustrate a strong linkage between riparian hydrology and the microbial processes that produce GHG. Deep groundwater tables fueled large respiration rates in the relatively dry soils near the hillslope, while denitrification mostly occurred in the wet zones located near the stream channel. As occurred at spatial scale, riparian soil water content was a primarily control of the temporal patterns of CO₂ and N₂O emissions. Soil dryness 350 diminished respiration rates during summer, while a fast soil N cycling promoted high N₂O emissions after a rewetting event in spring. Overall, our study highlights that future variations in catchment hydrology due to climate change can potentially affect the riparian functionality in Mediterranean zones, as well as their contribution to regional and global C and N cycles.

Author contributions

355 Sílvia Poblador, Santiago Sabaté, and Francesc Sabater designed the experiment. Sílvia Poblador and Anna Lupon carried them out. Sílvia Poblador performed all laboratory analysis. Anna Lupon and Sílvia Poblador analyzed the data set and prepared the manuscript, with contributions from Santiago Sabaté and Francesc Sabater.

Aknowledgments

We are thankful to Ada Pastor and Lúdia Cañas for their invaluable assistance in the field. Special thanks are extended to Núria Catalán for helpful comments on an earlier version of the manuscript. Financial support was provided by the Spanish 360 Government through the projects MONTES-Consolider (CSD2008-00040-MONTES), MEDFORESTREAM (CGL2011-30590), and MEDSOUL (CGL2014-59977-C3-2). Sílvia Poblador was supported by a FPI PhD fellowship from the Spanish Ministry of Economy and Competitiveness (BES-2012-054572). Anna Lupon was supported by a Kempe Foundation post-doctoral grant (Sweden) and the MEDSOUL project. We also thank site cooperators, including Vichy Catalan and the Catalan Water Agency (ACA) for permission to sample at the Font del Regàs catchment. Sílvia Poblador, Anna Lupon, Santiago 365 Sabaté, and Francesc Sabater are members of the research group FORESTREAM (AGAUR, Catalonia 2014SGR949).

References

2012, R. C. T.: Team RDC.R: A Language And Environment For Statistical Computing., n.d.

- Allison, S. D., Wallenstein, M. D. and Bradford, M. A.: Soil-carbon response to warming dependent on microbial physiology, *Nat. Geosci.*, 3(5), 336–340, doi:10.1038/ngeo846, 2010.
- 370 Asensio, D., Peñuelas, J., Ogaya, R. and Llusà, J.: Seasonal soil and leaf CO₂ exchange rates in a Mediterranean holm oak forest and their responses to drought conditions, *Atmos. Environ.*, 41, 2447–2455, doi:10.1016/j.atmosenv.2006.05.008, 2007.
- Audet, J., Hoffmann, C. C., Andersen, P. M., Baattrup-Pedersen, A., Johansen, J. R., Larsen, S. E., Kjaergaard, C. and Elsgaard, L.: Nitrous oxide fluxes in undisturbed riparian wetlands located in agricultural catchments: Emission, uptake and
375 controlling factors, *Soil Biol. Biochem.*, 68, 291–299, doi:10.1016/j.soilbio.2013.10.011, 2014.
- Àvila, A. and Rodà, F.: Changes in atmospheric deposition and streamwater chemistry over 25 years in undisturbed catchments in a Mediterranean mountain environment, *Sci. Total Environ.*, 434, 18–27, doi:10.1016/j.scitotenv.2011.11.062, 2012.
- Baethgen, W. E. and Alley, M. M.: A manual colorimetric procedure for measuring ammonium nitrogen in soil and plant
380 Kjeldahl digests, *Commun. Soil Sci. Plant Anal.*, 20(9–10), 961–969, doi:10.1080/00103628909368129, 1989.
- Baggs, E. M.: A review of stable isotope techniques for N₂O source partitioning in soils: recent progress, remaining challenges and future considerations, *Rapid Commun. Mass Spectrom.*, 22, 1664–1672, doi:10.1002/rcm, 2008.
- Barba, J., Curiel Yuste, J., Poyatos, R., Janssens, I. A. and Lloret, F.: Strong resilience of soil respiration components to drought-induced die-off resulting in forest secondary succession, *Oecologia*, 182(1), 27–41, doi:10.1007/s00442-016-3567-
385 8, 2016.
- Bardon, C., Piola, F., Bellvert, F., Haichar, F. el Z., Comte, G., Meiffren, G., Pommier, T., Puijalón, S., Tsafack, N. and Poly, F.: Evidence for biological denitrification inhibition (BDI) by plant secondary metabolites, *New Phytol.*, 204(3), 620–630, doi:10.1111/nph.12944, 2014.
- Batson, J., Noe, G. B., Hupp, C. R., Krauss, K. W., Rybicki, N. B. and Schenk, E. R.: Soil greenhouse gas emissions and
390 carbon budgeting in a short-hydroperiod floodplain wetland, *J. Geophys. Res. G Biogeosciences*, 120(1), 77–95, doi:10.1002/2014JG002817, 2015.
- Belnap, J., Welter, J. R., Grimm, N. B., Barger, N. and Ludwig, J. a: Linkages between Microbial and Hydrologic Processes in Arid and Semiarid Watersheds, *Ecol. Soc. Am.*, 86(2), 298–307, 2005.
- Bernal, S., Butturini, A., Nin, E., Sabater, F. and Sabater, S.: Leaf litter dynamics and nitrous oxide emission in a
395 Mediterranean riparian forest: implications for soil nitrogen dynamics., *J. Environ. Qual.*, 32(1), 191–197 [online] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12549558>, 2003.

- Bernal, S., Sabater, F., Butturini, A., Nin, E. and Sabater, S.: Factors limiting denitrification in a Mediterranean riparian forest, *Soil Biol. Biochem.*, 39(10), 2685–2688, doi:10.1016/j.soilbio.2007.04.027, 2007.
- 400 Bond-Lamberty, B. and Thomson, A.: Temperature-associated increases in the global soil respiration record., *Nature*, 464(7288), 579–582, doi:10.1038/nature08930, 2010.
- Bruland, G. L., Richardson, C. J. and Whalen, S. C.: Spatial variability of denitrification potential and related soil properties in created, restored, and paired natural wetlands, *Wetlands*, 26(4), 1042–1056, doi:10.1672/0277-5212(2006)26[1042:SVODPA]2.0.CO;2, 2006.
- 405 Burgin, A. J. and Groffman, P. M.: Soil O₂ controls denitrification rates and N₂O yield in a riparian wetland, *J. Geophys. Res. Biogeosciences*, 117(1), 1–10, doi:10.1029/2011JG001799, 2012.
- Casals, P., Lopez-Sangil, L., Carrara, A., Gimeno, C. and Nogués, S.: Autotrophic and heterotrophic contributions to short-term soil CO₂ efflux following simulated summer precipitation pulses in a Mediterranean dehesa, *Global Biogeochem. Cycles*, 25(3), 1–12, doi:10.1029/2010GB003973, 2011.
- 410 Castro-Díez, P., González-Muñoz, N., Alonso, A., Gallardo, A. and Poorter, L.: Effects of exotic invasive trees on nitrogen cycling: A case study in Central Spain, *Biol. Invasions*, 11(8), 1973–1986, doi:10.1007/s10530-008-9374-3, 2009.
- Chang, C. T., Sabaté, S., Sperlich, D., Poblador, S., Sabater, F. and Gracia, C.: Does soil moisture overrule temperature dependence of soil respiration in Mediterranean riparian forests?, *Biogeosciences*, 11(21), 6173–6185, doi:10.5194/bg-11-6173-2014, 2014.
- 415 Clément, J.-C., Pinay, G. and Marmonier, P.: Seasonal dynamics of denitrification along topohydrosequences in three different riparian wetlands, *J. Environ. Qual.*, 31(3), 1025–1037, doi:10.2134/jeq2002.1025, 2002.
- Davidson, E. A., Janssens, I. A. and Lou, Y.: On the variability of respiration in terrestrial ecosystems: Moving beyond Q₁₀, *Glob. Chang. Biol.*, 12(2), 154–164, doi:10.1111/j.1365-2486.2005.01065.x, 2006.
- 420 DeSimone, J., Macrae, M. L. and Bourbonniere, R. a.: Spatial variability in surface N₂O fluxes across a riparian zone and relationships with soil environmental conditions and nutrient supply, *Agric. Ecosyst. Environ.*, 138(1–2), 1–9, doi:10.1016/j.agee.2010.03.007, 2010.
- Dhondt, K., Boeckx, P., Hofman, G. and Cleemput, O.: Temporal and spatial patterns of denitrification enzyme activity and nitrous oxide fluxes in three adjacent vegetated riparian buffer zones, *Biol. Fertil. Soils*, 40(4), 243–251, doi:10.1007/s00374-004-0773-z, 2004.
- 425 Emmett, B. A., Beier, C., Estiarte, M., Tietema, A., Kristensen, H. L., Williams, D., Peñuelas, J., Schmidt, I. and Sowerby, A.: The response of soil processes to climate change: Results from manipulation studies of shrublands across an environmental gradient, *Ecosystems*, 7(6), 625–637, doi:10.1007/s10021-004-0220-x, 2004.

- Eno, C. F.: Nitrate production in the field by incubating the soil in polyethylene bags, *Soil Sci. Soc. Am. J.*, 24, 227–279, 1960.
- 430 Eriksson, L.; Byrne, T.; Johansson, E.; Trygg, J.; Vikström, C.: Multi- and Megavariate Data Analysis. Basic principles and applications., 2006.
- Giles, M., Morley, N., Baggs, E. M. and Daniell, T. J.: Soil nitrate reducing processes - Drivers, mechanisms for spatial variation, and significance for nitrous oxide production, *Front. Microbiol.*, 3(DEC), 1–16, doi:10.3389/fmicb.2012.00407, 2012.
- 435 Gómez-Gener, L., Obrador, B., von Schiller, D., Marcé, R., Casas-Ruiz, J. P., Proia, L., Acuña, V., Catalán, N., Muñoz, I. and Koschorreck, M.: Hot spots for carbon emissions from Mediterranean fluvial networks during summer drought, *Biogeochemistry*, 125(3), 409–426, doi:10.1007/s10533-015-0139-7, 2015.
- Goulden, M. L., Miller, S. D., Da Rocha, H. R., Menton, M. C., De Freitas, H. C., De Silva Figueira, A. M. and Dias de Sousa, C. A.: Diel and Seasonal Patterns of Tropical Forest CO₂ Exchange, *Ecol. Appl.*, 14(4), 42–54, doi:10.1890/02-6008, 2004.
- 440 Groffman, P. M., Gold, A. J. and Simmons, R. C.: Nitrate dynamics in riparian forests: Microbial studies, *J. Environ. Qual.*, 21(4), 666–671, doi:10.2134/jeq1992.00472425002100040022x, 1992.
- Groffman, P. M., Gold, A. J. and Jacinthe, P.-A. a: Nitrous oxide production in riparian zones and groundwater, *Chemosph. - Glob. Chang. Sci.*, 2, 291–299, doi:10.1023/A:1009719923861, 2000.
- 445 Guckland, A., Corre, M. D. and Flessa, H.: Variability of soil N cycling and N₂O emission in a mixed deciduous forest with different abundance of beech, *Plant Soil*, 336(1–2), 25–38, doi:10.1007/s11104-010-0437-8, 2010.
- Guidolotti, G., Rey, A., D’Andrea, E., Matteuci, G. and De Angelis, P.: Effect of environmental variables and stand structure on ecosystem respiration components in a Mediterranean beech forest, *Tree Physiol.*, 0, 1–13, doi:10.1093/treephys/tpt065, 2013.
- 450 Hagedorn, F.: Hot spots and hot moments for greenhouse gas emissions from soils, *Swiss Fed. Inst. For. Snow Landsc. Res.*, (1), 2010.
- Harms, T. K. and Grimm, N. B.: Hot spots and hot moments of carbon and nitrogen dynamics in a semiarid riparian zone, *J. Geophys. Res.*, 113(G1), 1–14, doi:10.1029/2007JG000588, 2008.
- Harms, T. K. and Grimm, N. B.: Responses of trace gases to hydrologic pulses in desert floodplains, *J. Geophys. Res. Biogeosciences*, 117(1), 1–14, doi:10.1029/2011JG001775, 2012.
- 455 Harms, T. K., Wentz, E. A. and Grimm, N. B.: Spatial heterogeneity of denitrification in semi-arid floodplains, *Ecosystems*, 12(1), 129–143, doi:10.1007/s10021-008-9212-6, 2009.

- Healy, R. W., Striegl, R. G., Russell, T. F., Hutchinson, G. L. and Livingston, G. P.: Numerical Evaluation of Static-Chamber Measurements of Soil - Atmosphere Gas Exchange : Identification of Physical Processes, *Soil Sci. Soc. Am. J.*, 60(3), 740–747, doi:10.2136/sssaj1996.03615995006000030009x, 1996.
- 460 Hedin, L. O., Fischer, J. C. Von, Ostrom, N. E., Kennedy, B. P., Brown, M. G., Robertson, G. P., Ecology, S. and Mar, N.: Thermodynamic Constraints on Nitrogen Transformations and Other Biogeochemical Processes at Soil-Stream Interfaces, , 79(2), 684–703, 1998.
- Hefting, M. M., Bobbink, R. and de Caluwe, H.: Nitrous oxide emission and denitrification in chronically nitrate-loaded riparian buffer zones., *J. Environ. Qual.*, 32(4), 1194–203 [online] Available from:
465 <http://www.ncbi.nlm.nih.gov/pubmed/12931872>, 2003.
- Hefting, M. M., Clément, J.-C., Dowrick, D., Cosandey, A. C., Bernal, S., Cimpian, C., Tatur, A., Burt, T. P. and Pinay, G.: Water table elevations controls on soil nitrogen cycling in riparian wetlands along a European climatic gradient, *Biogeochemistry*, 67, 113–134, doi:10.1023/B:BIOG.0000015320.69868.33, 2004.
- Hill, A. R.: Nitrate Removal in Stream Riparian Zones, *J. Environ. Qual.*, 25(4), 743–755,
470 doi:10.2134/jeq1996.00472425002500040014x, 1996.
- Inclán, R., Uribe, C., Sánchez, L., Sánchez, D. M., Clavero, Á., Fernández, A. M., Morante, R. and Blanco, A.: N₂O and CH₄ fluxes in undisturbed and burned holm oak, scots pine and pyrenean oak forests in central Spain, doi:10.1007/s10533-010-9520-8, 2014, 2014.
- Jacinthe, P. a., Vidon, P., Fisher, K., Liu, X. and Baker, M. E.: Soil Methane and Carbon Dioxide Fluxes from Cropland and
475 Riparian Buffers in Different Hydrogeomorphic Settings, *J. Environ. Qual.*, 44(0), 1080–1090,
doi:10.2134/jeq2015.01.0014, 2015.
- Keeney, D. R.; Nelson, D. W.: Nitrogen-inorganic forms, in *Agronomy Monograph 9*, ASA and SSSA., pp. 643–698, Madison., 1982.
- Kesik, M., Ambus, P., Baritz, R., Brüggemann, N., Butterbach-Bahl, K., Damm, M., Guyzer, J., Horváth, L., Kiese, R.,
480 Kitzler, B., Leip, A., LI, C., Pihlatie, M. and Pilegaard, K.: Inventories of N₂O and NO emissions from European forest soils,
353–375, 2005.
- Kim, J. and Verma, S. B.: Components of surface energy balance in a temperate grassland ecosystem, *Boundary-Layer Meteorol.*, 51(4), 401–417, 1990.
- Linn, D. M. and Doran, J. W.: Effect of Water-Filled Pore Space on Carbon Dioxide and Nitrous Oxide Production in Tilled
485 and Nontilled Soils, *Soil Sci. Soc. Am. J.*, 48(1961), 1267–1272, doi:10.2136/sssaj1984.03615995004800060013x, 1984.

- Lupon, A., Sabater, F., Miñarro, A. and Bernal, S.: Contribution of pulses of soil nitrogen mineralization and nitrification to soil nitrogen availability in three Mediterranean forests, *Environ. J. Soil Sci.*, 67, 303–313, doi:10.1016/j.aqpro.2013.07.003, 2016.
- 490 Mander, Ü., Well, R., Weymann, D., Soosaar, K., Maddison, M., Kanal, A., Lõhmus, K., Truu, J., Augustin, J. and Tournebize, J.: Isotopologue Ratios of N₂O and N₂ Measurements Underpin the Importance of Denitrification in Differently N - Loaded Riparian Alder Forests, *Environ. Sci. Technol.*, 48, 11910–11918, doi:dx.doi.org/10.1021/es501727h, 2014.
- Mander, Ü.; Lõhmus, K., Teiter, S., Muring, T., Nurk, K. and Augustin, J.: Gaseous fluxes in the nitrogen and carbon budgets of subsurface flow constructed wetlands, *Sci. Total Environ.*, 404(2–3), 343–353, doi:10.1016/j.scitotenv.2008.03.014, 2008.
- 495 McGlynn, B. L. and Seibert, J.: Distributed assessment of contributing area and riparian buffering along stream networks, *Water Resour. Res.*, 39(4), 1–7, doi:10.1029/2002WR001521, 2003.
- McLain, J. E. T. and Martens, D. A.: N₂O production by heterotrophic N transformations in a semiarid soil, *Appl. Soil Ecol.*, 32(2), 253–263, doi:10.1016/j.apsoil.2005.06.005, 2006.
- Mitsch, W. J. . and Gosselink, J. G.: *Wetlands*, 4th editio., edited by I. J. Wiley & Sons., 2007.
- 500 Morse, J. L. ., Ardon, M. and Benhardt, E. S.: Greenhouse gas fluxes in southeastern U.S. coastal plain wetlands under contrasting land uses, *Ecol. Appl.*, 22(1), 264–280, doi:10.1890/11-0527.1, 2012.
- Muller, D., Warneke, T., Rixen, T., Muller, M., Jahari, S., Denis, N., Mujahid, A. and Notholt, J.: Lateral carbon fluxes and CO₂ outgassing from a tropical peat-draining river, *Biogeosciences*, 12, 5967–5979, doi:10.5194/bg-12-5967-2015, 2015.
- 505 Oertel, C., Matschullat, J., Zurba, K., Zimmermann, F. and Erasmi, S.: Greenhouse gas emissions from soils - A review, *Chemie der Erde - Geochemistry*, 76, 327–352, doi:10.1016/j.chemer.2016.04.002, 2016.
- Pacific, V. J., McGlynn, B. L., Riveros-Iregui, D. A., Welsch, D. L. and Epstein, H. E.: Variability in soil respiration across riparian-hillslope transitions, *Biogeochemistry*, 91(1), 51–70, doi:10.1007/s10533-008-9258-8, 2008.
- 510 Pinay, G., Gumiero, B., Tabacchi, E., Gimenez, O., Tabacchi-Planty, a. M., Hefting, M. M., Burt, T. P., Black, V. a., Nilsson, C., Iordache, V., Bureau, F., Vought, L., Petts, G. E. and Décamps, H.: Patterns of denitrification rates in European alluvial soils under various hydrological regimes, *Freshw. Biol.*, 52(2), 252–266, doi:10.1111/j.1365-2427.2006.01680.x, 2007.
- 515 Pinay, G., Peiffer, S., De Dreuzy, J.-R., Krause, S., Hannah, D. M., Fleckenstein, J. H., Sebilo, M., Bishop, K. and Hubert-moy, L.: Upscaling Nitrogen Removal Capacity from Local Hotspots to Low Stream Orders ' Drainage Basins, *Ecosystems*, 18, 1101–1120, doi:10.1007/s10021-015-9878-5, 2015.

- 520 Raich, J. W., Potter, C. S. and Bhagawati, D.: Interannual variability in global soil respiration, 1980 - 94, *Glob. Chang. Biol.*, 8, 800–812, 2002.
- Rastogi, M., Singh, S. and Pathak, H.: Emission of carbon dioxide from soil, *Curr. Sci.*, 82(5), 510–517, 2002.
- Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. a., Kleber, M., Kögel-Knabner, I.,
520 Lehmann, J., Manning, D. a. C., Nannipieri, P., Rasse, D. P., Weiner, S. and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478(7367), 49–56, doi:10.1038/nature10386, 2011.
- Segers, R.: Methane production and methane consumption : a review of processes underlying wetland methane fluxes, *Biogeochemistry*, 41, 23–51, 1998.
- 525 Smith, M. S. and Tiedje, J. M.: Phases of denitrification following oxygen depletion in soil, *Soil Biol. Biochem.*, 11(3), 261–267, doi:10.1016/0038-0717(79)90071-3, 1979.
- Stark, J. M. and Firestone, M. K.: Mechanisms for soil moisture effects on activity of nitrifying bacteria, *Appl. Environ. Microbiol.*, 61(1), 218–221 [online] Available from: papers2://publication/uuid/0BBA43AD-EACB-4300-A741-A11EA89C2422, 1995.
- 530 Stern, S. N.: The Stern Review on the Economic Effects of climate change, *Popul. Dev. Rev.*, 32(December), 793–798, doi:10.2307/20058936, 2006.
- Suseela, V., Conant, R. T., Wallenstein, M. D. and Dukes, J. S.: Effects of soil moisture on the temperature sensitivity of heterotrophic respiration vary seasonally in an old-field climate change experiment, *Glob. Chang. Biol.*, 18(1), 336–348, doi:10.1111/j.1365-2486.2011.02516.x, 2012.
- 535 Teiter, S. and Mander, Ü.: Emission of N₂O, N₂, CH₄, and CO₂ from constructed wetlands for wastewater treatment and from riparian buffer zones, *Ecol. Eng.*, 25(5), 528–541, doi:10.1016/j.ecoleng.2005.07.011, 2005.
- Vidon, P., Allan, C., Burns, D., Duval, T. P., Gurwick, N., Inamdar, S., Lowrance, R., Okay, J., Scott, D. and Sebestyen, S.: Hot spots and hot moments in riparian zones: Potential for improved water quality management, *J. Am. Water Resour. Assoc.*, 46(2), 278–298, doi:10.1111/j.1752-1688.2010.00420.x, 2010.
- 540 Vidon, P., Marchese, S., Welsh, M. and McMillan, S.: Impact of Precipitation Intensity and Riparian Geomorphic Characteristics on Greenhouse Gas Emissions at the Soil-Atmosphere Interface in a Water-Limited Riparian Zone, *Water. Air. Soil Pollut.*, 227(8), doi:10.1007/s11270-015-2717-7, 2016.
- Vidon, P. G.: Not all riparian zones are wetlands: Understanding the limitation of the “wetland bias” problem, *Hydrol. Process*, 31(11), 2125–2127, doi:10.1002/hyp.11153, 2017.
- 545 Vidon, P. G. and Hill, A. R.: A landscape-based approach to estimate riparian hydrological and nitrate removal functions, *J. Am. Water Resour. Assoc.*, 3, 1099–1112, 2006.

Walker, J. T., Geron, C. D., Vose, J. M. and Swank, W. T.: Nitrogen trace gas emissions from a riparian ecosystem in southern Appalachia, *Chemosphere*, 49(10), 1389–1398, doi:10.1016/S0045-6535(02)00320-X, 2002.

550

Welti, N., Bondar-Kunze, E., Singer, G., Tritthart, M., Zechmeister-Boltenstern, S., Hein, T. and Pinay, G.: Large-scale controls on potential respiration and denitrification in riverine floodplains, *Ecol. Eng.*, 42, 73–84, doi:10.1016/j.ecoleng.2012.02.005, 2012.

Werner, C., Reiser, K., Dannenmann, M., Hutley, L. B., Jacobeit, J. and Butterbach-Bahl, K.: N₂O, NO, N₂ and CO₂ emissions from tropical savanna and grassland of northern Australia: An incubation experiment with intact soil cores, *Biogeosciences*, 11, 6047–6065, doi:10.5194/bg-11-6047-2014, 2014.

555

Wickland, K. P. ;, Neff, J. C. . and Harden, J. W.: The role of soil drainage class in carbon dioxide exchange and decomposition in boreal black spruce (*Picea mariana*) forest stands, *Can. J. For. Res.*, 40(11), 2123–2134, 2010.

Williams, C. J., Shingara, E. A. and Yavitt, J. B.: Phenol oxidase activity in peatlands in New York State: Response to summer drought and peat type, *Wetlands*, 20(2), 416–421, doi:10.1672/0277-5212(2000)020[0416:POAIP]2.0.CO;2, 2000.

Yavitt, J. B., Williams, C. J. and Wieder, R. K.: Production of methane and carbon dioxide in peatland ecosystems across North America: Effects of temperature, aeration, and organic chemistry of peat, *Geomicrobiol. J.*, 14(4), 299–316, 1997.

560

Yu, K. and Rinklebe, J.: Soil Redox Potential and pH Controllers, *Methods Biochem. Wetl.*, (10), 107–116, doi:10.2136/sssabookser10.c7, 2013.

Tables

565 **Table 1.** Mean annual values (\pm standard deviation) of soil water content (volumetric), soil temperature, soil pH, soil redox capacity (Eh), soil organic matter, soil molar C:N ratio, soil carbon (C) and nitrogen (N) content, and soil ammonium (NH_4^+) and nitrate (NO_3^-) concentrations for the three riparian zones. For each variable, different letters indicate statistical significant differences between riparian zones (*post-hoc* Tukey HSD test, $p < 0.05$).

	<i>Near-stream</i>	<i>Intermediate</i>	<i>Hillslope</i>
Soil water content (%)	29.58 \pm 7.55 ^A	19.36 \pm 6.00 ^B	19.81 \pm 6.24 ^B
Temperature (°C)	11.37 \pm 5.39 ^A	11.82 \pm 5.90 ^A	12.01 \pm 6.34 ^A
Eh	170 \pm 111 ^A	184 \pm 103 ^B	184 \pm 95 ^C
pH	6.66 \pm 0.42 ^A	6.31 \pm 0.50 ^A	6.68 \pm 0.53 ^A
Organic matter (%)	4.41 \pm 0.71 ^A	7.98 \pm 2.88 ^B	9.53 \pm 1.99 ^C
C:N ratio	14.25 \pm 3.64 ^A	14.09 \pm 1.78 ^A	13.63 \pm 1.18 ^A
C (mg kg⁻¹)	2004 \pm 1038 ^A	4007 \pm 1785 ^B	4923 \pm 1428 ^B
N (mg kg⁻¹)	160 \pm 44 ^A	330 \pm 135 ^B	418 \pm 107 ^C
NH₄⁺ (mg N kg⁻¹)	1.88 \pm 1.21 ^A	5.58 \pm 3.48 ^B	3.90 \pm 2.07 ^B
NO₃⁻ (mg N kg⁻¹)	0.75 \pm 0.58 ^A	4.66 \pm 4.25 ^B	5.30 \pm 4.20 ^B

570

Table 2. Results from the mixed-model analysis of variance (ANOVA) showing the effects of riparian zones and seasons on soil water content, soil temperature, soil pH, soil redox capacity (Eh), soil organic matter, soil molar C:N ratio, soil carbon (C) and nitrogen (N) content, and soil ammonium (NH₄⁺) and nitrate (NO₃⁻) concentrations. Plot was treated as a random effect in the model whereas riparian zones, seasons and their interactions were considered fixed effects. Values are *F*-values and the *p*-values are shown in brackets. *P*-values < 0.05 are shown in bold.

	<i>Riparian Zone</i>	<i>Seasons</i>	<i>Zone × Seasons</i>
Soil water content	18.6 [< 0.001]	100 [< 0.001]	13.6 [< 0.001]
Temperature	0.33 [0.721]	2117 [< 0.001]	0.42 [0.906]
pH	1.97 [0.182]	2.43 [0.060]	2.73 [0.052]
Eh	1.34 [0.247]	3.53 [0.062]	1.88 [0.084]
Organic matter	27.8 [< 0.001]	2.77 [0.053]	1.62 [0.144]
C:N ratio	0.99 [0.400]	10.9 [< 0.001]	1.72 [1.118]
C	27.1 [< 0.001]	1.86 [0.132]	0.77 [0.630]
N	39.7 [< 0.001]	1.22 [0.311]	0.63 [0.746]
NH₄⁺	12.4 [0.001]	2.71 [0.051]	1.52 [0.176]
NO₃⁻	22.4 [< 0.001]	5.63 [< 0.001]	4.09 [< 0.001]

Zone = near-stream, intermediate, hillslope.

Season = February, April, June, August and November.

580 **Table 3.** Mean values (\pm standard deviation) of potential denitrification rates (in mg N kg⁻¹ d⁻¹) after anoxia (DEA_{MQ}), carbon addition (DEA_C), nitrogen addition (DEA_{NO3}) and carbon and nitrogen addition (DEA_{C+NO3}) treatments for the three riparian zones during the study period. For each zone, different letters indicate statistical significant differences between treatments (*post-hoc* Tukey HSD test, n = 15, p < 0.01).

Potential Denitrification Rates (mg N kg⁻¹ d⁻¹)				
	DEA_{MQ}	DEA_C	DEA_{NO3}	DEA_{C+NO3}
<i>Near-stream</i>	0.31 \pm 0.41 ^A	0.26 \pm 0.27 ^A	0.42 \pm 0.42 ^A	0.63 \pm 0.85 ^A
<i>Intermediate</i>	1.01 \pm 1.12 ^A	1.88 \pm 1.59 ^A	2.28 \pm 3.57 ^A	2.40 \pm 2.45 ^A
<i>Hillslope</i>	1.34 \pm 1.33 ^A	2.35 \pm 1.97 ^{AB}	1.73 \pm 1.43 ^{AB}	3.82 \pm 2.78 ^B

585 **Table 4.** Summary of the partial least squares (PLS) models produced for CO₂ and N₂O emissions at the riparian site (n = 75). Values are the coefficients from PLS models which describe the relationship (direction and relative strength) between explanatory variables and gas emissions. The variance inflation factor (VIF) of each explanatory variable, indicative of collinearity, are shown in brackets. Bold values indicate the most influencing variables (variable importance in the projection (VIP) >1.0).



	<i>X-variable</i>	<i>Acronym</i>	<i>CO₂</i>	<i>N₂O</i>
Soil Properties	Soil water content (%)	SWC	-0.235 [1.72]	0.205 [1.32]
	Groundwater level (cm b.s.s.)	GWL	---	-0.157 [1.24]
	Temperature (C)	Tsoil	0.599 [1.45]	---
	pH	pH	---	---
	Redox potential (mV)	Eh	---	---
	Bulk density (g cm ⁻¹)	BD	---	---
	Coarse texture (%)	% Sand	---	---
	Organic matter (%)	SOM	---	---
	Total Carbon	C	---	---
	Total Nitrogen	N	---	---
	Molar C:N ratio	C:N ratio	---	---
	Ammonium	NH ₄ ⁺	0.167 [1.61]	---
	Nitrate	NO ₃ ⁻	0.066 [1.80]	-0.060 [1.47]
	Soil N processes	Net N Mineralization	NNM	---
Net Nitrification		NN	---	---
Denitrification		DNT	---	0.449 [1.09]
R²Y			0.71	0.40
Q²Y			0.66	0.34

Acronym is used in Figure S2 for PLS loading plots.


590

Figures

Figure 1. (a) Plot layout for the studied Mediterranean riparian forest showing the three riparian zones and the location of the chambers (n=5 for each riparian zone) (b) Conceptual approach of the influence of riparian hydrology on soil microbial processes across a Mediterranean riparian zone. Soil water content decreases from the near-stream to the hillslope zones due to changes in groundwater table, increasing unsaturated soil column and oxic conditions. Anaerobic processes (denitrification) occur under anoxic conditions while aerobic processes (respiration) are optimized under a moderate range of soil water content.




600 **Figure 2.** Temporal pattern of (a) mean monthly precipitation and (b) biweekly groundwater level at the studied riparian site during the year 2013. Circles are mean values of groundwater level at the near-stream (white), intermediate (grey), and hillslope (black) zones. Precipitation data was obtained from a meteorological station located at ca. 300 m from the studied riparian site. At each riparian zone, groundwater level was measured in 3 PVC piezometers (32-mm diameter, 1–3 m long) with a water level sensor (Eijkelkamp 11.03.30).




605

Figure 3. Temporal pattern of (a) soil water content, (b) soil temperature, (c) soil C:N molar ratio, and (d) soil nitrate concentration at 10-cm depth. Data is shown for the near-stream (white), intermediate (grey), and hillslope (black) zones during the study period. Circles are mean values and error bars are standard deviations.



610

Figure 4. Temporal pattern of (a) soil net N mineralization, (b) net nitrification and (c) denitrification rates at the near-stream (white), intermediate (grey), and hillslope (black) zones during the study period. Bars are mean values for each section and error bars are standard errors. For each season, different letters indicate significant differences among sections (mixed-model ANOVA, $p < 0.05$).



620 **Figure 5.** Temporal pattern of soil (a) CO₂ and (b) N₂O emissions at the near-stream (white), intermediate (grey), and hillslope (black) zones during the study period. Bars are mean values for each section and error bars are standard errors. For each season, different letters indicate significant differences among sections (mixed-model ANOVA, $p < 0.05$).

