

Dear Helge Niemann,

Thank you for your continued handling of the manuscript and compiling the second round of referee reports. We appreciate the opportunity to have completed the minor revisions as suggested by yourself and both referees. In particular, we have worked on consistency, formatting, and, sizing of the figures and tables, as well as included some further discussion regarding the limitations of our study given the shallow nature and small storage capacity of these freshwater bodies in our study region. Specific details are available in our below point-by-point response to the referees.

As previously, some of the issues with the figure formatting appear to persist due to the way that at this of submission the manuscript file (with figures) is processed as a single .pdf file which resulted in images being resized to fit on a page. The final submitted versions of these figures will have a consistent font size when submitted as individual files, assuming the final submission process allows for individual submission of vector-format figure files which can be better re-sized without loss. However, in the current uploaded manuscript file we believe we have addressed the aforementioned concerns.

Included below is a point-by-point response to the referees, in which we mark referee and editor comments with an (R) and (E), respectively, and author responses with an (A). At the conclusion of this document there is a complete marked-up manuscript with all changes highlighted.

**Editor:**

(E): The two reviewers seem relatively satisfied with your response and the revised version. Yet, both still raise a number of concerns that I agree with, and which I'd thus like to ask you to revise accordingly. Most importantly, reviewer 1 points out that the ponds you investigated were rather shallow and that it is therefore not surprising that rainfall events will modify the composition of solutes. The question that thus arises is: how representative are your study sites. This should be integrated in your MS to make the reader aware of the potential limitations of this study.

(A): We appreciate this feedback and have addressed this in further detail in the response to Referee #1. Briefly, we have integrated a discussion around the limitations of the study given the characteristics of our study ponds (very shallow basins with low storage capacity) and how this may vary from larger lake systems in the arctic or other temperate systems, generally with smaller systems being much more dynamic due to the annual flux terms representing a larger proportion of storage.

(E): Furthermore, the quality of figures/tables still needs considerable (!) improvement: Generally: use a common font type/size. Fig 3 is composed of two columns, Fig 4 of one but the font of axis labels etc is not uniform at all! Some figures are very rough (Excel?) exports in low resolution (eg Fig 4 labels are almost not discernible) and font type/size used for eg axis labels is differential across different figs. Scale figures sensibly. Do Fig 1 or 2 (for example) really need 1 page in width (I doubt it)?

(A): All Figures have been resized and scaled to ensure consistency in font size throughout (all figure text now is Times New Roman, either 12, 14 or 16 pt font for Figure text, axes text, and labels, respectively. Figure 1 and 2 have been re-sized and the label size has been enlarged in

Figure 4, while Figure 3 remains as 2 columns to highlight the differences between the two sets of chemical species, although the Figure has been re-sized and font size made consistent with other Figures. We believe the width of Figure 1 appropriate for context ito display the distribution of different sized surface water bodies in the general study region.

(E): Tab. 1: the horizontal lines are much, much too thick. Please check with a previously published BG article and with our format guidelines. Tab 2: even worse, the horizontal and vertical lines are double lines. Make this single and less, less thick (see above). Tab3 and 4: see tab 1 and 2 criticism.

(A): In all tables, the horizontal lines have been modified as to be consistent with published BG articles in this table. The vertical lines have been removed as per BG formatting guidelines.

(E): Fig. 1: There're several issues with this figure. The insert map indicates two things which I cannot really relate to the bigger map: there is a star (Churchill region) but it is unclear if the small area you investigated (the big map has ~10 km stretch of coastline at the Hudson bay) is within this Churchill area (the star has a diameter of perhaps 100 km. Is the aforesaid area that big?). Then it's very unclear what the dark region, which is basically the entire coastline of Ontario at the Hudson Bay signifies. Finally, the big map needs an x-axis (longitude) and y-axis (latitude). Without this, the figure is not acceptable because it lacks reference.

(A): This figure has been altered to reflect these suggestions, specifically: a) the star has been made into an arrow pointing to the specific location of the site, b) the dark coloured region, representing the Hudson Bay Lowland extent, has been further highlighted and clarified in the caption c) latitude and longitude has been added as axes to the larger map.

(E): Fig 6. the colour scaling of the PCA does not make sense like this as it is only possible to lets say discern day 200 from 150 but not 200 from 250.

(A): We have used a new colour ramp which we feel more clearly differentiates equally between these intervals in day of year in a rainbow legend as suggested by Referee #1.

### **Referee #1**

(R): 1) It is unclear to me how the authors decided which ponds were hydrologically controlled. I can't find the pattern in Table 5, not the text. It may be me. If not, please provide.

(A): It has been clarified in-text that it is particular chemical species which were determined to hydrologically controlled, as opposed to individual ponds (although there were some minor exceptions noted in the text and bolded within the table), as well as small clarification added to the methods section that this was a manual classification guided by the consistency of the  $\beta_2$  parameter within a pond for all chemical species.

(R): 2) Figure 6: I find it hard to differentiate the lakes (especially Left and Strange). Maybe use a rainbow legend for DOY and a white dot for Strange.

(A): We have used a new colour ramp and symbology for this figure for clarity.

(R):Page 6 Line 5: Do you mean to say 'spatial' twice here?

(A): This sentence is to clarify that each chemical measurement is associated with both a deviation from a temporal mean and deviation from a spatial mean. The sentence now reads: "Thus, each sample (n = 72; 12 samples from each of 6 ponds) is associated with both a temporal deviation score and spatial deviation score for each chemical species."

(R): Page 6 Line 27: When you say 'inlets', do you mean channels, or inflows / outflows?

(A): All mentions of 'inlet' in-text have been clarified as 'inflow'.

(R): Section 3.4: You need to reference Figure 6 in this paragraph.

(A): A reference to Figure 6 has been added.

(R): Page 8 Line 7: The two nitrogen species behaved in specific ways, as did other constituents; sometimes. Perhaps be specific about which species were most affected by landscape scale patterns.

(A): References of the behaviour of specifically which chemical species responded to landscape scale patterns (Cl<sup>-</sup>, Na<sup>+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, and dissolved organic nitrogen) have been added to the text, Page 9, Lines 12-13.

(R): Page 8 Line 24: This is not what Figure 5 shows.

(A): There were several misreferenced figures and tables in earlier version of this manuscript due to renumbering with the addition and removal of tables to supplementary information. All references to each figure and table number have been updated and checked.

(R): Page 8 Line 30: You need a segue here. Maybe start the paragraph with "The importance of biogeochemical versus hydrological processes can be informed by the location of individual ponds and chemical species in PCA space."

(A): As suggested, this sentence was slightly modified and inserted to more naturally introduce the discussion of PCA space. It now reads "The importance of biogeochemical and hydrological processes can be informed by the location of (a) clusters of samples from individual ponds and (b) ordination of chemical species in PCA space."

(R) Page 9 Lines 1 and 3: Less vs. lesser?

(A): This was a grammatical error in our text, and now reads "less".

(R): Page 9 Line 5: Maybe move this content about calcium to the previous paragraph as an example of the biogeochemical processes of importance.

(A): The discussion has been reorganized and the content regarding calcium cycling in this landscape is now contained with the discussion regarding biogeochemical processes in the context of PCA space.

(R): Page 9 Line 12: I found the content of this paragraph confusing and tangential. What study had 20 lakes? There is some information missing here. I think the authors are trying to put their work in the context of others, but that is not coming across well. Link it all better together or delete it all.

(A): This was a typo and should have included a reference to work of White et al. (2014) – the citation has been added in-text.

(R): Page 9 Line 25: Please include a statement of what impact the use of the curve fitting method may have on your results of which ponds are hydrologically driven or spatially variable (or not). Maybe refer to Figures 6 and 2 and Table 5 here. Can problems with the method explain differences between the hydrology and variability classifications?

(A): We have included a statement regarding the potential impact of the manual curve fitting method and how it may relate to discrepancies between those results and the results of the variability comparisons.

(R): Page 9 Line 33: This sentence has too many thoughts in it. Break it up.

(A): This sentence has been split into two and rewritten for clarity.

(R): Page 11 Line 11: I still maintain ‘synchronicity’ is not the word you need. Change them all to synchrony.

(A): All instances of ‘synchronicity’ have been replaced in-text with synchrony.

(R): Page 11 Line 28: When discussing pond hydrology be specific. Hydrology could mean any flux, including change in storage, evaporation etc. You really mean inflows and outflows here.

(A): We thank the referee for pointing out this ambiguity and have replaced mentions of “hydrology” when appropriate throughout with specific processes.

#### References:

White, J., Hall, R. I., Wolfe, B. B., Light, E. M., Macrae, M. L., and Fishback, L.: Hydrological connectivity and basin morphometry influence seasonal water-chemistry variations in tundra ponds of the northwestern Hudson Bay Lowlands. *Arct. Antarct. Alp. Res.*, 46(1), 218-235, 2014.

#### **Referee #2**

(R): For your notice, it seems to me that Bello and Smith, 1990 (who you refer to as describing mixing/stratification patterns in your study ponds) do not show that these water bodies are not diurnally stratified; they discuss heat exchange and assume complete mixing by wind action. I am, however, not concerned anymore about this issue knowing that these ponds are less than 50 cm deep. Stratification, if any, will be short-lasting...

(A): We appreciate the clarification regarding the reference - our sampling design used the assumption of wind-mixing and lack of stratification to guide our single sample per pond. We will clarify this in the methods section and be careful in attributing the fact that this lack of stratification is an assumption (which we believe to be a reasonable assumption).

(R): What concerns me really is how representative are such shallow ephemeric ponds to the general pond population in the area? Some other studies (e.g., Wolfe et al., 2011) have shown that ponds such as these studied here can disappear completely in a dry years. While I appreciate high resolution sampling and efforts taken to capture changes in chemistry I am not convinced

how novel are these findings...Water bodies that are so shallow and small will be intuitively very strongly affected by rainfalls, either through dilution or enrichment effects (in certain compounds delivered from the catchment). Novelty would arise if rainfall patterns in the studied area are subject to interannual variability (as they surely are) and this variability leads to higher/lower biogeochemical processing in studied ponds. On the other hand, I understand the necessity to start discussion about effects of short-term meteorological forcing on chemistry of shallow water bodies in the Arctic (Cortés et al. 2017).

This said, manuscript would really benefit from trimming toward being more process and less detail oriented.

(A): It is a point well taken that the results of this work are not necessarily surprising. Unfortunately, we did not collect data of this frequency over multiple years in order to examine the role of interannual variability of precipitation patterns on hydrochemistry, although we agree this would make a novel and interesting contribution to do so. Further, we agree the results from this study likely do not scale to larger and more chemostatic water bodies, as the water fluxes each year representing a much smaller change to the total storage than they do in these very shallow tundra ponds. We believe that (as the reference to the work of Cortés et al. suggests) the incorporation of hydrological context into sampling regimes is essential and we believe our study explores some themes around how future work could build on these techniques. However, we agree it is important to address the role of size in the water bodies in the transferability and applicability of this study to other study regions, so we have included text in the discussion regarding the limitation of the study ponds with such a low storage capacity and how results may differ in larger lake systems.

(R): *In the Abstract you say:*

Across all ponds, temporal variability (across the season and within a single rain event) exceeded spatial variability (variation among ponds) in concentrations of several major species (Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Na<sup>+</sup>).

*And in the Results you say:*

Spatial deviations did not significantly exceed temporal deviations for any chemical species

*Which one of these statements is correct?*

(A): Both of these statements are true – in the abstract we note for which chemical species that temporal variability exceeded spatial variability. The converse (Spatial deviations did not significantly exceed temporal deviations) was true for no chemical species. We have clarified this in-text.

(R): *There were few exceptions to this manual classification,*

Replace "manual" by "general"

(A): This sentence now reads "There were few exceptions to this general classification."

(R): In contrast, and campaigns that may be more temporally coarse but spatially intensive are applicable for larger, less chemodynamic lakes and ponds, in which total water inputs and outputs represent a much smaller proportion of the total volume.

*This sentence needs revision, for example.*

(A): We have revised this sentence and made several small additional copy-edits to the discussion to clarity, primarily through the breaking up of some longer sentences into components.

(R): *Table 4 bold/underline the exceptions, not italicized (not evident)*

*Tables general look better with thin border lines*

(A): In Table 4, the exceptions have been bolded. The horizontal lines have been modified as to be consistent with published BG articles in this table, and are much thinner as a result.

# Capturing temporal and spatial variability in the chemistry of shallow permafrost ponds

Matthew Q. Morison<sup>1</sup>, Merrin L. Macrae<sup>1</sup>, Richard M. Petrone<sup>1</sup>, LeeAnn Fishback<sup>2</sup>

<sup>1</sup>Department of Geography and Environmental Management, University of Waterloo, Waterloo, N2L 3G1, Canada

<sup>2</sup>Churchill Northern Studies Centre, Churchill, R0B 0E0, Canada

Correspondence to: Matthew Q. Morison ([mmorison@uwaterloo.ca](mailto:mmorison@uwaterloo.ca))

## Abstract.

Across the circumpolar north, the fate of small freshwater ponds and lakes (< 1 km<sup>2</sup>) has been the subject of scientific interest due to their ubiquity in the landscape, capacity to exchange carbon and energy with the atmosphere, and their potential to inform researchers about past climates through sediment records. A changing climate has implications for the capacity of ponds and lakes to support organisms and store carbon, which in turn has important feedbacks to climate change. Thus, an improved understanding of pond biogeochemistry is needed. To characterize spatial and temporal patterns in water column chemistry, a suite of tundra ponds were examined to determine (1) Does temporal variability exceed spatial variability? (2) If temporal variability exists, do all ponds (or groups of ponds) behave in a similar temporal pattern, linked to seasonal hydrologic drivers or precipitation events? Six shallow ponds located in the Hudson Bay Lowlands region were monitored between May-October 2015 (inclusive, spanning the entire open-water period). The ponds span a range of biophysical conditions including pond area, perimeter, depth, and shoreline development. Water samples were collected regularly, both bimonthly over the ice free season, and intensively during and following a large summer storm event. Samples were analyzed for nitrogen speciation (NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, dissolved organic nitrogen) and major ions (Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup>). Across all ponds, temporal variability (across the season and within a singlerain event) exceeded spatial variability (variation among ponds) in concentrations of several major species (Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Na<sup>+</sup>). Evapoconcentration and dilution of pond water with precipitation and runoff inputs were the dominant processes influencing a set of chemical species which are hydrologically driven (Cl<sup>-</sup>, Na<sup>+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, dissolved organic nitrogen), whereas the dissolved inorganic nitrogen species were likely mediated by processes within ponds. This work demonstrates the importance of understanding hydrologically driven chemodynamics in permafrost ponds on multiple scales (seasonal and event scale).

## 1 Introduction

Small, shallow freshwater bodies (<1 km<sup>2</sup>) across the circumpolar north are ubiquitous in the landscape (Muster et al., 2013; Dyke and Sladen, 2010; Macrae et al., 2014). In many regions, these features maintain the capacity to store and exchange carbon (C) (Macrae et al., 2004; Abnizova et al., 2012), act as habitat (Smol and Douglas, 2007), and mediate the landscape energy balance (Chapin et al., 2000). These tundra ponds are highly sensitive to climate change (Schindler and Smol, 2006), which may impact their hydrology and biogeochemistry. This has implications for their ability to support organisms and store or cycle C, which in turn has important feedbacks to climate change. Thus, an improved understanding of pond biogeochemistry is needed. Climate change may impact pond hydrobiogeochemical conditions by modifying pond hydrologic and thermal conditions, as well as pond-catchment connectivity. In

permafrost systems, recent work has focused on the impact of climatically-induced permafrost thaw on lake and pond chemistry (Vonk et al., 2015) over a timescale of decades, generally showing increasing concentrations of ions indicative of deeper flowpaths and weathering processes (Kokelj et al., 2009). Lougheed et al. (2011) examined changes in tundra pond chemistry over 40 years in Alaska and observed substantial increases in inorganic nutrients, which were attributed to a thaw-based release from shallow permafrost and increased atmospheric nitrogen (N) deposition. In addition to changing chemical contributions from thawing catchments, the water balances of ponds are sensitive to climatic changes that impact the ponds directly as well as indirectly through changing catchment hydromorphology (Walvoord and Kurylyk, 2016). Adjacent to pond edges, thermokarst expansion acts as a mechanism for lateral seepage and loss (eg, Yoshikawa and Hinzman, 2003) leading to increased connectivity, potentially modifying seasonal biogeochemical trajectories (Abbott et al., 2015). Elevated levels of nutrients and/or ions may impact the C uptake function of ponds via primary productivity, which has been shown to be nutrient limited in subarctic and arctic freshwater bodies (Levine and Whalen, 2001; Bergström et al., 2005; Bergström, 2010; Bonilla et al., 2005; Symons et al., 2012). Indeed, increased nutrient loading to ponds may either bring about an increase in pond primary productivity (Levine and Whalen, 2001; O'Brien et al., 2005) or may lead to a decrease through increased turbidity and suboptimal conditions for photosynthesis (northern Sweden, Ask et al., 2009; northern Québec, Roiha et al., 2015), linking nutrient concentrations to biological markers such as chlorophyll (Przytulska et al., 2016). Although climate change may lead to long-term changes in pond permanence and hydrologic exchange, pond hydrologic storage can also vary greatly over the course of a single season or among different hydrologic years (Morison et al., 2017, Woo and Guan, 2006). Annual changes to pond storage often reflect the highly variable local climatic conditions, and ponds may either desiccate completely (e.g. Bouchard et al., 2013; Smith et al., 2005) or spill over bankfull storage and/or coalesce (Macrae et al., 2014). The chemical concentrations of the water column in ponds is closely tied to hydrology, both directly through physical processes (i.e. dilution and evapoconcentration; Borghini et al., 2013) and indirectly through mediating environmental factors that control the rate of biogeochemical processes (e.g. variable pond depth controlling the diurnal water temperature amplitude; Smol and Douglas, 2007). Mazurek et al. (2012) sampled shallow ponds over several summer seasons in Svalbard and found ionic concentrations increasing in ponds over the summer months, coincident with pond drying. White et al. (2014) monitored ponds at three time steps over the ice-free season and related pond seasonal biogeochemical trajectories to pond morphology and pond-peatland connectivity. Given the significant control of [hydrology/hydrological processes](#) on pond biogeochemistry, and the highly variable hydrological conditions experienced by ponds within a single season, an improved understanding of seasonal patterns of pond hydrochemistry is needed to better understand potential pond responses to climate change.

The concepts of hydrochemical coherence (or synchrony; Baines et al., 2000) have been employed in temperate lake systems to determine the role of different climatic and hydrological factors in driving lake chemical variations, [employing often by examining](#) correlation [of](#) different chemical measures [across space](#) (e.g. Magnuson et al., 1990). For instance, Pace and Cole (2002) found in Northern Michigan that changes to dissolved organic carbon and colour were synchronous among lakes while total phosphorus, chlorophyll, and pH were not. Other methods, such as the use of evaporative normalization factors (e.g. Borghini et al., 2013), have allowed for comparisons among difference



chemical species and the degree to which they are driven by hydrological variation. Both of these methods allow for comparison of relative deviations of values which would otherwise be masked by absolute differences among lakes. Approaches of hydrochemical coherence have been less seldom employed in subarctic permafrost pond and lake settings.

Recent approaches to characterising shallow pond biogeochemical processes across Arctic, Antarctic, and alpine permafrost settings have used three to four water samples from each lake or pond, taken at different points of the ice-free season to represent “early-season”, “mid-season”, and “late-season” conditions. Other approaches have relied on a single annual sample to characterize various geochemical indicators of ponds, such as nutrient status, carbon dynamics, major ion and metal concentrations, often sampling many ponds and lakes (often 10 – 100) at that single time period (Supplementary Table S1). However, given the variability in pond storage and hydrological conditions over the course of an ice-free season, it is important to frame the results of pond chemical analyses in the proper hydrologic context, and further, to be prudent in generalizing pond geochemical process from samples taken in different hydrological periods. Beyond broad spatial sampling campaigns of many periods at a coarse temporal resolution, less is known about the amplitude, drivers, and total variation in seasonal and event chemographs of shallow ponds in remote northern catchments, despite their importance in driving key ecological functions across the circumpolar north. Further, as large summer storms increase in frequency and magnitude (Sauchyn and Kulshreshtha, 2008; Kaufman et al., 2009), it is important to understand how storm events control short-term water chemistry variation, which can represent a substantial proportion of the annual water balance over a period of hours to days (Macrae et al., 2014).

The objectives of this study were to characterize spatial and temporal patterns in the chemistry of a suite of tundra ponds to determine if: (1) temporal variability exceeded spatial variability throughout the open water season; (2) temporal variability existed, whether all ponds (or groups of ponds) behaved in a similar temporal pattern, linked to season or hydrology; and (3) spatiotemporal variability in pond biogeochemical signatures could be used to make inferences about processes occurring within ponds and between ponds and surrounding peatlands. This information will improve our understanding of hydrobiogeochemical processes in tundra ponds, particularly in terms of seasonal patterns, and, will aid in developing appropriate sampling designs in future. Further, this study provides a geochemical survey of six ponds in the Hudson Bay Lowlands region at two unprecedented temporal scales (bimonthly, spanning an ice-free season, in addition to diurnal samples spanning an intense rainfall-runoff event).

## **2 Methods**

### **2.1 Study site and sampling design**

This study was completed in the Hudson Bay Lowlands (HBL), approximately 20 km East of the town of Churchill, Manitoba, Canada. The regional climate is strongly influenced by the close proximity to Hudson Bay (Rouse, 1991), with average air temperatures in the region (1980-2010 normal) of -6.5 °C (annually) with a snow-free season from May-October, and maximum temperatures in July (monthly average 12.7 °C; Government of Canada, 2016).

The HBL are the largest contiguous wetland complex in North America, with thousands of lakes and ponds in the landscape (Keller et al., 2014; Dyke and Sladen, 2010) from impeded drainage due very low topographic gradients (Boudreau and Rouse, 1995) and the presence of permafrost. Ponds have been demonstrated to be stably stratified in the eastern HBL (Deshpande et al., 2015), while the very shallow (<1 m) ponds in the western HBL ~~have been shown~~ are assumed to be ~~non-stratified~~, well mixed by wind action (Bello and Smith, 1990). Ponds in the western HBL have been shown to be nutrient-limited (Eichel et al., 2014; Symons et al., 2012), restricted in surface and groundwater connection (Boudreau and Rouse, 1995), with steep banks and thick, organic sediments with dense mats of benthic blue-green, green and diatomaceous algae (Macrae et al., 2004; Gray, 1987), with the dominant diatom genus being *Denticula* (Macrae et al., 2004). Symons et al. (2012) showed the most abundant phytoplankton communities to be *Chlamydomonas* spp., *Sphaerocystis* spp., *Diatoma* spp. and *Crugienella* spp. Due to their small size and extremely alkaline conditions (8 to 9.8 pH, Macrae et al., 2004; Bos and Pellatt, 2012), ponds are devoid of fish communities with the exception of stickleback (*Gasterosteidae*). Ponds are also devoid of aquatic macrophytes with the exception of *Carex aquatilis* found along pond perimeters in zones where water is shallow. Eichel et al. (2014) demonstrated the importance of benthic organisms to the cycling of nutrients in these ponds, showing experimentally that mesocosms with sediment removed inorganic nutrients from the water column at a rate orders of magnitude greater than mesocosms without sediment.

The ponds and their catchments are underlain by continuous permafrost, with a seasonal active layer of 40 – 100 cm thickness. In the catchments, there is a peat layer of 30 – 60 cm overlying unsorted glaciomarine till. The surface is comprised of patterned polygonal peat plateaus segregated by sedge lawn and narrow fen channel networks. Dry uplands (plateau) are dominated by lichen-heath communities, with sparse tree cover (*Larix* spp. and *Salix* spp.). Transition zones (moderate moisture, sedge lawns) are comprised of hummocks and hollows, dominated by a mixture of lichen-heath communities and *Carex mitis*. Low-lying channel fens (saturated) are dominated by *Carex aquatilis*. Dwarf shrubs (*Betula glandulosa* and *Salix arctica*) are also found in some channel fens and around pond perimeters. Six study ponds were selected: Erin pond, Frisbee pond, Larch pond, Left pond, Strange pond, and Sandwich pond (Figure 1). These ponds were selected for their range in surface areas, perimeter, catchment area, and depth, because they have been the subject of long-term research in the area, and have been shown to be representative of a range of chemical and hydrological properties (Table 1; Macrae et al., 2004; White et al., 2014; Wolfe et al., 2011; Bos and Pellatt, 2012). Each pond was sampled regularly (every 14 days) throughout the ice-free season of 2015, from Day of Year (DOY) 140 to 300, May to October). Additionally, during an intensive storm during DOY 185 – 188, ponds were sampled twice per day during the storm and 24 hours following the event to determine the hydrochemical impacts of a storm event, including the inputs of both direct precipitation and runoff water. Pond stage (water column depth) data were recorded hourly with pressure transducers (HOBO U20, Onset Ltd) installed in PVC (50 mm inner diameter) slotted pipes located within 1m of the shore of each pond (with the exception of Erin pond) and corrected with a barometric logger housed in an identical PVC pipe to minimize temperature-related artifacts in pressure differences between barometric and level loggers (McLaughlin and Cohen, 2011). Precipitation data were taken from the Environment Canada 'Churchill' station (20 km West of the study site). Runoff was determined as the residual change in pond storage exceeding the input from direct precipitation during and up to 24 h. following a rain event, after

Morison et al. (2017). The perimeter and area of each pond were surveyed with a Trimble Juno SB GPS unit, with a differentially corrected horizontal accuracy of  $\pm 5\text{m}$ .

## 2.2 Water chemistry sampling and analysis

Samples of pond water were collected manually as a depth-integrated profile of the water column, approximately 3 m from the pond edge, and placed in sterile, triple-primed 50 ml polyethylene containers. Care was taken to not disturb pond sediments when samples were being collected. Samples were preserved on ice in the field and passed through a  $0.45\ \mu\text{m}$  cellulose acetate filter within two hours of collection. A 50 ml subsample was acidified to a final concentration of 0.2%  $\text{H}_2\text{SO}_4$  and stored until digestion. Non-acidified samples were immediately frozen. All samples were shipped to the Biogeochemistry Lab at the University of Waterloo for analysis. In the lab, acidified samples were digested (Kjeldahl digestions, Seal Analytical Hot Block Digestion System BD50, Seattle, USA) for the analysis of Dissolved Kjeldahl N (DKN). Digested samples and non-acidified subsamples were analysed for N species using colorimetric methods: (Bran-Luebbe AutoAnalyzer III, Seal Analytical: Methods G-102-93, (ammonium;  $\text{NH}_4^+\text{-N}$ ); G-103-93 (SRP-P), G-109-94 (nitrate;  $\text{NO}_3^-\text{-N}$ ), G-189-97 (DKN)). Results for  $\text{NH}_4^+$  and  $\text{NO}_3^-$  have a detection limit of  $0.001\ \text{mg N L}^{-1}$ . Dissolved organic nitrogen (DON) concentrations were determined as the difference between DKN and  $\text{NH}_4^+$  concentrations. Samples were analysed for other major ions: chloride ( $\text{Cl}^-$ ), sulphate ( $\text{SO}_4^{2-}$ ), sodium ( $\text{Na}^+$ ), potassium ( $\text{K}^+$ ), magnesium ( $\text{Mg}^{2+}$ ), and calcium ( $\text{Ca}^{2+}$ ) using ion chromatography (DIONEX ICS 3000, IonPac AS18 and CS16 analytical columns).

## 2.3 Statistical techniques

Due to large variability (spatial and temporal) in absolute values of concentration across the data set, values were normalized as deviations from mean values to permit their comparison (Schoenfeld and Numberger, 2007; Viner, 1984). This was accomplished by examining the departure of each sampling point from the mean of either temporal or spatial groups. More specifically, for each measurement, proportional deviations from either spatial or temporal means were determined as the absolute difference from unity of either the ratio of the measured sample concentration at a particular date and pond,  $[x]_{st}$ , to the mean concentration of the species from all ponds sampled that date,  $[\bar{x}]_t$ ; to examine spatial deviations, Eq. (1), or the ratio of the measured sample concentration to the mean concentration of the species from all samples taken from that pond over the entire study period,  $[\bar{x}]_s$ ; to examine temporal deviations, Eq. (2).

$$\left| \frac{[x]_{st}}{[\bar{x}]_t} - 1 \right| = \text{Spatial Deviation Score} \quad (\text{Eq. 1})$$

$$\left| \frac{[x]_{st}}{[\bar{x}]_s} - 1 \right| = \text{Temporal Deviation Score} \quad (\text{Eq. 2})$$

Thus, each sample ( $n = 72$ ; 12 samples from each of 6 ponds) is associated with both a temporal deviation score and spatial deviation score for each chemical species. Once a set of deviation scores was computed for each species, a Mann-Whitney test was used to determine if there was a significant difference between the medians of the set of spatial deviation scores and the set of temporal deviation scores for each chemical species.

To determine the relationship between pond volume and concentrations of different chemical species, a power function (Eq. 3) was fit to the spatially normalized concentration (the ratio of each concentration to the mean concentration  $[\bar{x}]_t$ ) and stage data to represent the geometry of conical bathymetry (based on observations and measurements of several sediment basins) in the non-linear pond stage-volume relationship.

$$[x] = \beta_1(\text{stage})^{\beta_2} \quad (\text{Eq. 3})$$

The parameter  $\beta_2$  is invariant under linear translation, such as spatial-normalization, and is expected to maintain a negative value with the magnitude depending on the bathymetry of the pond (with a theoretical value of 0 for a cylindrical flat-bottomed pond, and highly negative for a steep conically shaped pond basin). The fit of each power function was assessed ~~by~~ for each chemical species in each pond with the root mean ~~squared~~ square error, as the coefficient of determination  $R^2$  is inappropriate for the nonlinear case (Spiess and Neumeier, 2010). Since the pond mean-normalized concentrations are dimensionless and not scale-dependant, the root mean square error term is also dimensionless and can be compared across ponds and chemical species. The degree of hydrologic control on different nutrient species concentrations was manually determined by the performancegoodness of the fittingfit of a power curve to each set of normalized concentration data against pond stage for each of the five instrumented ponds. In each pond, chemical species with a constant pond-specific value of  $\beta_2$  (the slope of the power curve) with low root mean square error (S) were categorized as hydrologically driven, where species with either (1) a  $\beta_2$  value inconsistent with other hydrologically driven species or (2) poor performance (greater root mean square error) were categorized as non-hydrologically driven.

A principal component analysis (PCA; Dunteman, 1989) was employed on logarithmically-transformed hydrochemistry data, using singular value decomposition scaled to have unit variance. PCA has been used before in explorations of pond chemical trajectories over a scale of months to years (White et al., 2014, Sokal et al., 2010, Wiklund et al., 2012).

Coherence was assessed by Pearson correlation following the approach of Magnuson et al. (1990), both among lakes for each chemical species as well as for each chemical species (in all lakes combined). All statistical analyses were conducted with R software (version 3.3.2, R Development Core Team, 2016).

### 3 Results

#### 3.1 Spatial and temporal variability of pond chemistry

~~Spatial deviations did not significantly exceed temporal deviations for any chemical species (Figure 2).~~ Medians of temporal deviation scores significantly exceeded the median spatial deviation scores in several chemical species ( $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ); Figure 2) but there was no significant difference between the median spatial and temporal deviation scores for  $\text{DON}$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ , and  $\text{NO}_3^-$ . For species that were significantly more temporally variable than spatially variable, concentrations among ponds, despite differences in magnitude, tended to be strongly positively correlated with each other ( $p < 0.01$ ) with the notable exception of  $\text{DON}$  (Table 2).  $\text{NO}_3^-$  and  $\text{NH}_4^+$  were not significantly correlated to any other species ( $p > 0.01$ ) (Table 3) while all other species tended to be strongly positively

correlated with each other ( $p < 0.01$ ) although there were some exceptions. Spatial deviations did not significantly exceed temporal deviations for any chemical species.

### 3.2 Seasonal patterns in pond water chemistry

Pond water chemistry varied among ponds, but also throughout the season (Figure 3). Throughout 2015, the temporal trend of pond mean-normalized concentrations in the six ponds appeared to have a spatial coherency in DON, Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, Na<sup>+</sup>, K<sup>+</sup> and Mg<sup>2+</sup> concentrations. There was a gradual increase in concentrations of these species from DOY 140 to 260, punctuated by small decline in concentrations following a large storm on DOY 185-188, followed by a gradual decrease from DOY 260 to the end of the season. Ca<sup>2+</sup> displayed a contrasting temporal trajectory to the other chemical species, decreasing from DOY 190 to the end of the season, while NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> appeared to have no coherent pattern relative to the other chemical species. Generally, ponds were closest to their mean values during the mid-season (DOY 200 – 250) and pond-mean normalized concentrations differed the least among ponds during this time, being more divergent earlier (DOY 140 – 200) and later (DOY 250 – 300) in the season.

### 3.3 Pond hydrology and linkages with water chemistry

As was observed for water chemistry, there was also a seasonal pattern in pond hydrologic storage (Figure 4). In general, ponds were full following snowmelt, with water levels generally declining throughout the season and eventually rising again in late summer. Catchments showed synchronous runoff responses to two major rain events on DOY 166 and DOY 185 and three smaller rain events between DOY 260 and 290, although the magnitude of response varied. Following the early-season rain events, Left pond and Strange pond experienced prolonged summer drawdown while this drawdown was less pronounced in Larch, Frisbee and Sandwich ponds. At DOY 250 all ponds began a gradual increase in stage continuing to the end of season, but did not return to post-snowmelt levels. Within these five study ponds with recorded depth, there appeared to be no clear relationship between variation in stage and pond geometry (depth, shoreline development, area, or perimeter; Table 1).

Generally, DON, Cl<sup>-</sup>, Na<sup>+</sup>, K<sup>+</sup> and Mg<sup>2+</sup> were classified as generally the hydrologically driven chemical species among ponds and Ca<sup>2+</sup>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SO<sub>4</sub><sup>2-</sup> were classified as non-hydrologically driven (Table 4; Figure 5). There were few exceptions to this manual general classification, most notably, DON and K<sup>+</sup> tended to slightly poorer fits (larger S; slightly less hydrologically driven) in Sandwich pond, whereas SO<sub>4</sub><sup>2-</sup> may have been hydrologically driven in Strange pond. Slight differences in  $\beta_2$  values among ponds for hydrologically driven species are influenced by the differences in bathymetry which dictates the stage-volume relationship.

### 3.4 Principal component analysis

PCA allowed for a reduction in the redundancy of species that are highly correlated (Table 3) and the clustering of samples from each pond in PC space (Figure 6). The analysis also shows a clustering of chemical species along the PC axes. A collection of the hydrologically driven chemical species (Table 4; DON, Mg<sup>2+</sup>, K<sup>+</sup>, Na<sup>+</sup>, Cl<sup>-</sup>) closely follows the PC1 axis with slightly negative scores for PC2. Two other non-hydrologically driven chemical species have positive scores along PC1 with slightly negative (SO<sub>4</sub><sup>2-</sup>) and positive (Ca<sup>2+</sup>) scores along PC2. Inorganic nitrogen

species, ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ ), also non-hydrologically driven, have negative scores along the PC2 axis and slightly negative scores along PC1. The spread of data points for Strange Pond, Left Pond and Frisbee Pond are more elongate along the PC1 axis where the spread of data from Larch pond, Erin pond and Sandwich pond are aligned along the PC2 axis. Temporally, the data tend to move from the upper-left quadrant (positive scores for PC2, negative scores for PC1) to the lower-right quadrant (positive scores for PC1, negative scores for PC2) over the course of the season. Chemical species which were classified as hydrologically driven (Table 4) were not necessarily the same species which showed that temporal variance exceeded spatial variance (Figure 2). DON and  $\text{Mg}^{2+}$  were classified as hydrologically driven but did not have the same degree of temporal coherence displayed by  $\text{Cl}^-$ ,  $\text{Na}^+$ , and  $\text{K}^+$ . Conversely,  $\text{Ca}^{2+}$  and  $\text{SO}_4^{2-}$  display significantly greater temporal variability than spatial variability, but were not classified as hydrologically driven constituents. Strange, Left, and Frisbee ponds were elongated with respect to the PC1 axis, which corresponds closely with the group of chemical species that are hydrologically driven. Strange and Left ponds showed the greatest variation in pond stage (Table 1; Figure 4), linking the hydrological variation with the variation in concentrations of chemical species controlled by stage. Frisbee pond showed ~~lesser~~ less variation in stage but was elongate with respect to the axis of the hydrologically driven species cluster, suggesting that variation in pond stage alone does not necessarily dictate which species are most chemodynamic within a pond. Larch and Sandwich ponds, which showed lesser variation in stage, were elongated with respect to PC2, orthogonal to the cluster of hydrologically driven species along PC1. The redundancy in the original dataset is primarily due to the clustering of the hydrologically driven species along PC1, and the very close clustering of  $\text{Na}^+$  and  $\text{Cl}^-$  in PC space (indicating the marine halite influence on these ponds; Bos and Pellatt, 2012). In addition, the decoupling of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  trajectories (Figure 3) and in PC space (Figure 6) indicate the potential role of  $\text{Ca}^{2+}$  in either carbonate cycling associated with productivity and carbon dioxide intake (Macrae et al., 2004), precipitation, or cation substitution (Wetzel, 2001). Further, not only are the hydrologically driven species (Table 4) correlated among ponds (Table 2), but they are also correlated among species (Table 3). The apparent redundancy of several of these chemical trajectories is observed in the normalized time series of concentration data, with many of the species which align along PC1 axis showing similar concentration trajectories at the seasonal (Figure 3) and event (Figure 7) scales.

### 3.5 Event-scale variation in pond chemistry

In addition to being highly chemodynamic over the course of an ice-free season, pond chemistry varied greatly over the course of a single storm-runoff event (Figure 7). For samples taken during, and following the storm event in Strange Pond on DOY 185-188, concentrations of some chemical species varied by the same magnitude over the course of several days as they did over the period of the entire ice-free season (DON,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{NO}_3^-$ ); other species varied but not to the same extent ( $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ ; Supplementary Table S2). In Frisbee pond, chemical species tended to vary by a similar magnitude over the storm period as they did over the entire season and this was not restricted to DON,  $\text{K}^+$ , and  $\text{Ca}^{2+}$ .

## 4 Discussion

### 4.1 Seasonal and spatial variability in pond chemistry

Ponds in this study displayed a coherent trend in several chemical concentrations over ~~an~~ ice-free season, despite differences in pond geometry, catchment characteristics, and absolute magnitude of chemical concentrations. Landscape scale factors such as climate are controlling hydrological processes, and in turn, concentrations of chemical species (specifically,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ , and dissolved organic nitrogen). Magnuson et al. (1990) and Baines et al. (2000) reported  $\text{Ca}^{2+}$  to be temporally coherent among groups of large lakes in Wisconsin, USA, consistent with our findings, while Magnuson et al. (1990) found  $\text{K}^+$  or  $\text{SO}_4^{2-}$  to ~~be less coherent~~ lack coherence, contrasting with our results. However, the absolute magnitudes of different chemical species concentrations varied among ponds ~~suggesting despite similar trajectories of mean-normalized concentrations. This suggests~~ that influences such as runoff inputs, evapoconcentration, bathymetry, and internal processes (sediment-water interaction and biological cycling) may be driving these absolute differences while preserving seasonal trajectories relative to a pond mean value. Other authors have reported on local spatial heterogeneity of biogeochemical properties of ponds ~~and their extreme spatial variability~~, such as the conditions of nutrient limitation of primary productivity in different lakes (Hogan et al., 2014, Symons et al., 2012). This work suggests that the seasonal trajectory of concentration of at least some chemical species are not reflective of this spatial variability and instead more reflective of landscape-level ~~process~~ processes. The importance of biogeochemical and hydrological processes can be informed by the location of (a) clusters of samples from individual ponds and (b) ordination of chemical species in PCA space. In many of the same study ponds as this work, White et al. (2014) found similar clustering of  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Mg}^{2+}$  along their first PC axis (most variance explained) and another cluster including  $\text{NH}_4^+$  and  $\text{NO}_3^-$  along their second axes. In contrast, total nitrogen was approximately orthogonal to both those clusters, instead of being aligned with the other species which were identified as hydrologically driven in our method analysis. Differences between the findings of White et al. (2014) and this work may be explained by the inclusion of additional limnological measurements into the PCA, the greater number of lakes in ~~the~~ their study (20), the smaller number of samples taken at each lake per year (3), and the reporting of total N as opposed to the dissolved fraction. Sokal et al. (2012) did not report a similar clustering of chemical species in PC space, with TKN,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ , and  $\text{K}^+$  showing great variation relative to the tight clustering found in this study in the Slave River Delta and with less separation from the non-hydrologically driven chemical species in our study ( $\text{Ca}$ ,  $\text{SO}_4$ ). This could be due to larger and deeper lakes studied in the region ( $\sim 10^6 \text{ m}^2$  surface area; 1.5 m deep) in addition to additional limnological and isotopic measurements into the PCA. However, this difference in species clustering may not be region-specific as Wiklund et al. (2012) found a clustering of  $\text{Na}^+$ ,  $\text{Cl}^-$ , TKN,  $\text{K}^+$  and  $\text{Mg}^{2+}$  in PC space from lakes in the Peace Athabasca Delta region. Wiklund et al. (2012) also report approximately equal difference angles between ~~the angle for the~~ vectors of (a) the cluster of hydrologically driven chemical species and (b)  $\text{Ca}^{2+}$  and  $\text{SO}_4^{2-}$  as was found in this study.

### 4.2 Event-scale variation in pond chemistry

Short-term variation in concentrations during a storm was pronounced in Frisbee pond during the storm of DOY 185 – 188, 2015. This is likely due to the difference in the proportion of new water entering the pond relative to the volume

of old water in the pond. In Frisbee pond, the rainfall inputs represent a much greater fraction of the total pond water at the conclusion of the event (nearly one-half), compared to Strange pond, which is larger and more full of water pre-event, where event water represents a smaller proportion of the total post-event water (approximately one-quarter) despite the lesser runoff amount (Figure 4). This leads to the dampening of the storm-based concentration variability in Strange pond. ~~Among chemical species, those~~ Those species that may have elevated concentrations in event water relative to lower concentrations in pond water, such as DON (Morison et al., 2017), show a heightened response in terms of pond concentrations in both Frisbee pond and Strange pond, ~~whereas species, Species~~ Species which have event water concentrations which are similar to background pond water concentrations show a ~~lesser dampened~~ response. Although the volume of event water being added may not represent a great fraction of the total water in the pond, it is enriched in species that occur naturally in rainwater or through primary runoff pathways (shallow, organic subsurface material). This ~~is consistent~~ agrees with the findings of Abnizova (2013) which showed that the ionic composition at the event scale of runoff water to ponds in high arctic wetlands was caused by the vegetative structure of the catchment (related to dissolved organic material) and bedrock composition.

#### 4.3 Inferring hydrological and biogeochemical process from patterns in pond chemistry

This study does not aim to present direct measurements of any biogeochemical process, including the magnitude and direction of which physiochemical transformations are operating to which extent. Each chemical species measured in this study is subject to differing degrees of reactivity/conservatism. Future research is needed to elucidate the individual or coupled cycling processes of these elements in subarctic freshwater ecosystems. Chemical species which appear to be driven primarily by non-hydrological processes (Table 4) are likely those which are most reactive. However, our results can be contextualized within the existing body of knowledge of hydrobiogeochemical processes which have been measured in similar environments previously to determine if our observations are consistent with previous findings of different species.

Biogeochemical variations large in magnitude also often occur on short time scales, which are typically associated with the movement of water across geochemical boundaries (McClain et al., 2003). Previous work in tundra and high Arctic environments have shown pathways for transformations of runoff water through the catchments, including biological incorporation of nutrients (Thompson and Woo, 2009) and geochemical processes of weathering and ion exchange (Quinton and Pomeroy, 2006), which impact the quality of the runoff water arriving at the pond. Once the runoff water is delivered to the ponds, the biological factors driving responses to the incorporation and sedimentary burial of nitrogen and phosphorus on short timescales (< 72 hours of return to pre-event levels) were demonstrated experimentally by Eichel et al. (2014). These ~~observations~~ observations are consistent with the ~~delivery of nutrients~~ increase in ~~nutrient concentrations~~ followed by rapid uptake to pre-event levels which was observed during and following the storm event around DOY 185 (Figure 7) in this study. These mechanisms of runoff contributions to pond chemical variability are important for future studies that aim to use seasonal and interannual variations in pond and lake chemistries to examine changing flowpaths in thawing permafrost environments. The relative change of concentrations in ponds and lakes over events, seasons, and years will be a product of the difference in concentrations between new and old water and the degree of dilution of new water arriving at the pond. If the concentration of a



constituent in old water is not significantly different from that in new water, future work which infers understand changing flow paths from changing chemical concentrations over time may significantly underestimate the degree of change to the movement of water and solutes due to thawing permafrost (Lougheed et al., 2011). ~~or conversely~~. Conversely, it also possible to overestimate the change in flowpaths ~~in~~ delivering solutes which are radically different in concentration from old water. In this work, no clear relationship between chemical trajectory (Figure 3) and runoff (Figure 4) was observed, nor were either related specifically to total catchment area (Table 1). However, this may be a result of the variable source area concept applying in this fill-and-spill system (Spence et al., 2010; Morison et al., 2017), in which substantial portions of the catchment are not contributing runoff to the ponds for ~~some storms~~ certain storms which do not provide requisite overfilling of the catchment storage capacity threshold. This effects renders estimates of total catchment area which are delineated by topography less useful without additional knowledge of total subsurface storage capacity and effective contributing area for each runoff event.

The method of fitting curves to stage-concentration relations to manually determine the degree to which ~~pond hydrology~~ evapoconcentration and dilution dictates chemical concentrations may not capture all of the complex dynamics related to hydrological controls. For example, pond volume controls diurnal temperature fluctuation amplitude, and indirectly the rate of biogeochemical processes. In northern tundra environments, low temperatures limit the rates at which biogeochemical processes occur (Stark, 2007). Microbial decomposition remains the crucial bottleneck (rate limiting step) in nutrient cycling (Stark, 2007; Andersen et al., 2013). Temperatures are often the most important factor limiting the rate of microbial decomposition of organic matter in cold region aquatic and terrestrial ecosystems (Wallenstein et al., 2009; Vonk et al., 2015; Wickland et al., 2012). On the North shore of Alaska, Koch et al. (2014) showed that the chemical dynamics were less pronounced in ~~ponds that~~ larger ponds with greater storage capacity which maintained lower temperatures, limiting evaporation (and therefore evapoconcentration) as well as retarding biological uptake; ~~larger ponds with dilute inflows were subject to greater evaporation as a fraction. Some of pond volume~~ these processes may explain some discrepancies between the classified hydrologically-driven chemical species and greater nutrient depletion over the course of the year, linking the combined effects of hydrology both directly and indirectly impacting nutrient concentrations those species for which temporal variability exceeded spatial variability.

#### 4.4 Implications for sampling design in future studies

Generally, the current standard of infrequent measurements per annum are unlikely to be sufficient to capture the variability in highly chemodynamic systems, such as shallow ponds in which storage varies by up to 80% of the mean value over the course of the ice-free season. In contrast, and campaigns that may be more temporally coarse but spatially intensive are applicable for larger, less chemodynamic lakes and ponds, in which total water inputs and outputs represent a much smaller proportion of the total volume. Further, the synchrony in the chemical behaviour of some species ( $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{SO}_4^{2-}$ ,  $\text{Ca}^{2+}$ ) in these ponds suggests potential redundancy in the sampling of a great deals of lakes if the research goals are related to the measurement of these chemical species. This study may be subject to ~~the converse problem~~ different limitations by observing water column chemistry in only 6 lakes over the ice-free season. However, it appears these six ponds capture a representative range of chemical concentrations, spanning 1-2

Formatted: Justified, Space Before: 12 pt, After: 0 pt, Line spacing: 1.5 lines

orders of magnitude, (Supplementary Table S2) which are representative of much wider spatial surveys in much of the work outlined in Supplementary Table S1 (and more specifically, in line with the variance observed by other authors who have studied ponds in our specific region, Bos and Pellatt, 2012; White et al., 2014; Jacques et al., 2016). The fact that these ponds are closely spatially clustered may remove effects of microclimates on the scales of 10-100 kilometers which may reduce the degree of synchrony observed in chemical concentrations, due to different precipitation, runoff, and evaporative drawdown regimes. Further, the very shallow and small nature of these ponds in this work, although comprising the majority of surface water features in permafrost landscapes (Muster et al., 2013), may not reflect the chemostatic nature of larger lakes in permafrost regions. Larger lakes, with a much greater total volumetric storage capacity, would be generally subject to hydrologic fluxes (evaporation, precipitation, inflows) which represent a much smaller proportion of total water storage on both annual and individual rain event timescales. This would result in the hydrodynamics and associated changes to hydrochemical concentrations to be dampened in these larger lakes relative to our highly chemodynamic systems.

Future biogeochemical research must consider the relative terms in the water balance in a pond to determine periods of greatest variability (such as snowmelt vs. rainwater dominated ponds, e.g. Bouchard et al., 2013). A water budget approach should be incorporated into future studies which attempt to use changing solute concentrations as a proxy for changes to ensure that appropriate hydrological weighting to changes of concentrations in surface water are properly represented. Alternately, approaches characterizing total mass depletion of different ions and nutrients (e.g. Koch et al., 2014), an evaporative normalization factor (e.g. Borghini et al., 2013), the use of the Shannon index in a hydrochemical context (Thompson and Woo, 2009), hydrologic context inference through isotopic composition (e.g. White et al., 2014) may provide a hydrologic context to concentration measurements which is lacking otherwise.

## 5 Conclusions

This study reported on the degree of hydrologic control on nutrient and major ion chemistry of a set of six of small (<1 ha), shallow permafrost ponds in the Hudson Bay Lowlands. Generally speaking, temporal variability exceeds spatial variability in pond chemistry in this landscape. Five chemical species ( $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$ ) showed a temporal coherence in six different ponds over the course of an ice-free season (Figure 8). No pond demonstrated a spatial coherence in concentration trajectory which was distinct from every other pond. A set of species was determined to be hydrologically controlled ( $\text{DON}$ ,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Mg}^{2+}$ ), either directly through processes of evapoconcentration- and inflow/precipitation dilution, or indirectly through pond hydrology exerting an influence on other environmental controls such as temperature regimes, which influence biogeochemical characteristics. Approaches of PCA and coherence/correlation showed that these hydrologically driven chemical species are strongly correlated to each other and among ponds. This evidence suggests that landscape level controls on pond hydrology may reduce the spatial heterogeneity in pond chemical trajectories for the hydrologically driven chemical species, but not for those species that are not hydrologically controlled ( $\text{Ca}^{2+}$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ ). Further, at the event scale, variation over the course of days during, and following precipitation events may be on equal order as the variation over the course of months. This longer period variation is also not limited to the hydrologically driven species, indicating the importance of biogeochemically active periods coinciding with periods of water movement across

boundaries (catchment-pond interactions). This work has important implications for understanding the highly chemodynamic behavior of small, shallow permafrost ponds through over both short-term and long-term temporal scales while remaining spatially synchronous. Our findings indicate the design of future sampling regimes, which continue to rely on inference from broad spatial sampling at coarse temporal scales must consider the hydrologic context of the samples taken. In particular, the proximity of recent rain events and the trajectory of pond water balance are important considerations for interpreting pond chemical concentration data for the purpose of inferring landscape level changes to permafrost environments.

#### **Data availability**

Unprocessed data (not subject to temporal or spatial deviation normalization) from all chemical analyses are available as a supplement to this work, Table S2.

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Author contributions**

M.Q. Morison, R.M. Petrone, M.L. Macrae and L.A. Fishback designed the sampling strategy and study objectives, L.A. Fishback supervised the field sample collection, M.Q. Morison performed the laboratory analyses, analysed the data, generated the figures, and prepared the manuscript, R.M. Petrone and M.L. Macrae provided comments on the manuscript composition and figure presentation.

#### **Acknowledgments**

The funding for this research came from NSERC Discovery Grants (Macrae, Petrone), NSERC Northern Supplement (Petrone), the Northern Scientific Training Program, and Northern Research Fund (Churchill Northern Studies Centre). Field assistance with sample collection was provided by Matt Webb and Kimberly Thompson. Laboratory assistance was provided by Vito Lam. Additional logistical support was provided by the staff and volunteers of the Churchill Northern Studies Centre. The authors thank Dr. Genevieve Ali and Dr. Janina Plach for helpful discussions on statistical analyses of the data- [the editorial team of Biogeosciences led by Dr. Helge Niemann, as well as Dr. Maciej Bartosiewicz and other anonymous referees for their helpful comments and insight towards improving the final draft of the manuscript.](#)

#### **References**

Abbott, B. W., Jones, J. B., Godsey, S. E., Larouche, J. R., and Bowden, W. B.: Patterns and persistence of hydrologic carbon and nutrient export from collapsing upland permafrost. *Biogeosciences*, 12(12), 3725-3740, 2015.

Abnizova, A., Siemens, J., Langer, M., and Boike, J.: Small ponds with major impact: The relevance of ponds and lakes in permafrost landscapes to carbon dioxide emissions. *Global. Biogeochem. Cy.*, 26(2), GB2041, doi:10.1029/2011GB004237, 2012.

Abnizova, A.: Hydrology, carbon dynamics and hydrochemical properties of ponds in an extensive low gradient High Arctic wetland, Polar Bear Pass, Bathurst Island, Nunavut, Canada. Ph.D. thesis, York University, Canada, 2013.

Archer, S. D., McDonald, I. R., Herbold, C. W., Lee, C. K., Niederberger, T. S., and Cary, C.: Temporal, regional and geochemical drivers of microbial community variation in the melt ponds of the Ross Sea region, Antarctica. *Polar Biol.*, 39(2), 267-282, 2016.

Aronow, S.: *Beaches and Coastal Geology*. Springer, United States, 1982.

Ask, J., Karlsson, J., Persson, L., Ask, P., Byström, P., and Jansson, M.: Terrestrial organic matter and light penetration: Effects on bacterial and primary production in lakes. *Limnol. Oceanogr.*, 54(6), 2034-2040, 2009.

Baines, S. B., Webster, K. E., Kratz, T. K., Carpenter, S. R., and Magnuson, J. J.: Synchronous Behavior of Temperature, Calcium, and Chlorophyll in Lakes of Northern Wisconsin, *Ecology*, 81(3), 815-815, 2000.

Balasubramaniam, A. M., Hall, R. I., Wolfe, B. B., Sweetman, J. N., and Wang, X.: Source water inputs and catchment characteristics regulate limnological conditions of shallow subarctic lakes (Old Crow Flats, Yukon, Canada). *Can. J. Fish. Aquat. Sci.*, 72(7), 1058-1072, 2015.

Bello R., and Smith J.D.: The effect of weather variability on the energy balance of a lake in the Hudson Bay Lowlands, Canada. *Arctic. Alpine. Res.*, 22, 98-107, doi:10.2307/1551724, 1990.

Bergström, A. K., Blomqvist, P., and Jansson, M.: Effects of atmospheric nitrogen deposition on nutrient limitation and phytoplankton biomass in unproductive Swedish lakes. *Limnol. Oceanogr.*, 50(3), 987-994, 2005.

Bergström, A. K.: The use of TN: TP and DIN: TP ratios as indicators for phytoplankton nutrient limitation in oligotrophic lakes affected by N deposition. *Aquat. Sci.*, 72(3), 277-281, 2010.

Bonilla, S., Villeneuve, V., and Vincent, W. F.: Benthic and Planktonic Algal Communities in a High Arctic Lake: Pigment Structure and Contrasting Responses to Nutrient Enrichment. *J. Phycol.*, 41(6), 1120-1130, 2005.

Borghini, F., Colacevich, A., Loisele, S.A, and Bargagli, R.: Short-term dynamics of physico-chemical and biological features in a shallow, evaporative antarctic lake. *Polar Biol.*, 36, 1147-1160, doi:10.1007/s00300-013-1336-2, 2013.

Bos, D. G., and Pellatt, M. G.: The water chemistry of shallow ponds around Wapusk national park of Canada, Hudson Bay Lowlands. *Can. Water. Resour. J.*, 37(3), 163-175, 2012.

Bouchard, F., Turner, K. W., MacDonald, L. A., Deakin, C., White, H., Farquharson, N., Medeiros, A.S., Wolfe, B.B., Hall, R.I., Pienitz, R., and Edwards, T. W. D.: Vulnerability of shallow subarctic lakes to evaporate and desiccate when snowmelt runoff is low. *Geophys. Res. Lett.*, 40(23), 6112-6117, 2013.

Boudreau, L. D., and Rouse, W. R.: The role of individual terrain units in the water balance of wetland tundra. *Climate Res.*, 5(1), 31-47, 1995.

Breton, J., Vallières, C., and Laurion, I.: Limnological properties of permafrost thaw ponds in northeastern Canada. *Canadian Can. J. Fish. Aquat. Sci.*, 66(10), 1635-1648, 2009.

Chapin, F.S., III, McGuire, A.D., Randerson, J., Pielke, R., Baldocchi, D., Hobbie, S.E., Roulet, N., Eugster, W., Kasischke, E., and Rastetter, E.B.: Arctic and boreal ecosystems of western North America as components of the climate system. *Glob. Chang. Biol.*, 6, 211–223, 2000.

Deshpande, B. N., MacIntyre, S., Matveev, A., and Vincent, W. F.: Oxygen dynamics in permafrost thaw lakes: Anaerobic bioreactors in the Canadian subarctic. *Limnol. Oceanogr.*, 60(5), 1656-1670, 2015.

Dunteman, G. H. (1989). *Principal components analysis* (No. 69). Sage.

Dyke, L. D., and Sladen, W. E.: Permafrost and peatland evolution in the northern Hudson Bay Lowland, Manitoba Arctic, 429-441, 2010.

Eichel, K. A., Macrae, M. L., Hall, R. I., Fishback, L., and Wolfe, B. B.: Nutrient uptake and short-term responses of phytoplankton and benthic algal communities from a subarctic pond to experimental nutrient enrichment in microcosms. *Arct. Antarct. Alp. Res.*, 46(1), 191-205, 2014.

Government of Canada: *Canadian Climate Normals*. Meteorological Service of Canada, available at: [http://climate.weather.gc.ca/climate\\_data/daily\\_data\\_e.html?StationID=48969](http://climate.weather.gc.ca/climate_data/daily_data_e.html?StationID=48969), 2016.

Gray I.M. Factors Regulating Phytoplankton Populations at Churchill, Manitoba. M.Sc. thesis, University of Windsor, Canada, 280 pp, 1987.

Hinkel, K. M., Arp, C. D., Townsend-Small, A., and Frey, K. E.: Can Deep Groundwater Influx be Detected from the Geochemistry of Thermokarst Lakes in Arctic Alaska?. *Permafrost. Periglac.*, doi: 10.1002/ppp.1895, 2016.

Hogan, E. J., McGowan, S., and Anderson, N. J.: Nutrient limitation of periphyton growth in arctic lakes in south-west Greenland. *Polar. Biol.*, 37(9), 1331-1342, 2014.

Houben, A. J., French, T. D., Kokelj, S. V., Wang, X., Smol, J. P., and Blais, J. M.: The impacts of permafrost thaw slump events on limnological variables in upland tundra lakes, Mackenzie Delta region. *Fund. Appl. Limnol.*, 189(1), 11-35, 2016.

Jacques, O., Bouchard, F., MacDonald, L. A., Hall, R. I., Wolfe, B. B., and Pienitz, R.: Distribution and diversity of diatom assemblages in surficial sediments of shallow lakes in Wapusk National Park (Manitoba, Canada) region of the Hudson Bay Lowlands. *Ecol. Evol.*, 6(13), 4526-4540, 2016.

Kaufman, D.S., Schneider, D.P., McKay, N.P., Ammann, C.M., Bradley, R.S., Briffa, K.R., and Lakes, A.: Recent warming reverses long-term Arctic cooling. *Science*, 325, 1236-1239, 2009.

Keller, W. B., Paterson, A. M., Rühland, K. M., and Blais, J. M.: Introduction—environmental change in the Hudson and James Bay Region. *Arct. Antarct. Alp. Res.*, 46(1), 2-5, 2014.

Koch, J. C., Gurney, K., and Wipfli, M. S.: Morphology-Dependent Water Budgets and Nutrient Fluxes in Arctic Thaw Ponds. *Permafrost. Periglac.*, 25(2), 79-93, 2014.

Kokelj, S. V., Zajdlik, B., and Thompson, M. S.: The impacts of thawing permafrost on the chemistry of lakes across the subarctic boreal-tundra transition, Mackenzie Delta region, Canada. *Permafrost. Periglac.*, 20(2), 185-199, 2009.

Larsen, A. S., O'Donnell, J. A., Schmidt, J. H., Kristenson, H. J., and Swanson, D. K.: Physical and chemical characteristics of lakes across heterogeneous landscapes in arctic and subarctic Alaska. *J. Geophys. Res.-Biogeo.*, 122, doi:10.1002/2016JG003729, 2017.

Levine, M. A., and Whalen, S. C.: Nutrient limitation of phytoplankton production in Alaskan Arctic foothill lakes. *Hydrobiologia*, 455(1-3), 189-201, 2001.

Lim D.S.S., Douglas M.S.V., Smol J.P., Lean D.R.S.: Physical and chemical limnological characteristics of 38 lakes and ponds on Bathurst Island, Nunavut, Canadian High Arctic. *Int. Rev. Hydrobiol.*, 86(1), 1–22, 2001.

Lougheed, V. L., Butler, M. G., McEwen, D. C., and Hobbie, J. E.: Changes in tundra pond limnology: Re-sampling Alaskan ponds after 40 years. *Ambio*, 40(6), 589-599, 2011.

Lyons, W. B., Welch, K. A., Gardner, C. B., Jaros, C., Moorhead, D. L., Knoepfle, J. L., and Doran, P. T.: The geochemistry of upland ponds, Taylor Valley, Antarctica. *Antarct. Sci.*, 24(01), 3-14, 2012.

MacDonald, L. A., Farquharson, N., Hall, R. I., Wolfe, B. B., Macrae, M. L., and Sweetman, J. N.: Avian-driven modification of seasonal carbon cycling at a tundra pond in the Hudson Bay Lowlands (northern Manitoba, Canada). *Arct. Antarct. Alp. Res.*, 46(1), 206-217, 2014.

MacLeod, J., Keller, W., Paterson, A.M., Dyer, R.D., and Gunn, J.M.: Scale and watershed features determine lake chemistry patterns across physiographic regions in the far north of Ontario, Canada. *J. Limnol.*, doi: 10.4081/jlimnol.2016.1553, 2016.

Macrae, M. L., Bello, R. L., and Molot, L. A.: Long-term carbon storage and hydrological control of CO<sub>2</sub> exchange in tundra ponds in the Hudson Bay Lowland. *Hydrol. Process.*, 18(11), 2051-2069, 2004.

Macrae, M. L., Brown, L. C., Duguay, C. R., Parrott, J. A., and Petrone, R. M.: Observed and projected climate change in the Churchill region of the Hudson Bay Lowlands and implications for pond sustainability. *Arct. Antarct. Alp. Res.*, 46(1), 272-285, 2014.

Magnuson, J. J., Benson, B. J., and Kratz, T. K.: Temporal coherence in the limnology of a suite of lakes in Wisconsin, U.S.A. *Freshwater. Biol.*, 23, 145-159, 1990.

Mallory, M. L., Fontaine, A. J., Smith, P. A., Wiebe Robertson, M. O., and Gilchrist, H. G.: Water chemistry of ponds on Southampton Island, Nunavut, Canada: effects of habitat and orithogenic inputs. *Arch. Hydrobiol.*, 166(3), 411-432, 2006.

Manasypov, R. M., Pokrovsky, O. S., Kirpotin, S. N., and Shirokova, L. S.: Thermokarst lake waters across the permafrost zones of western Siberia. *Cryosphere*, 8(4), 1177-1193, 2014.

Mazurek, M., Paluszkiwicz, R., Rachlewicz, G., and Zwoliński, Z.: Variability of water chemistry in tundra lakes, Petuniabukta coast, Central Spitsbergen, Svalbard. *Scientific World Journal*, doi:10.1100/2012/596516, 2012.

McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M., Hart, S.C., Harvey, J.W., Johnston, C.A., Mayorga, E., McDowell, W.H., and Pinay, G.: Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems. *Ecosystems*, 6(4), 301-312, 2003.

McLaughlin D.L., and Cohen M.J.: Thermal artifacts in measurements of fine-scale water level variation. *Water. Resour. Res.*, 47, W09601, doi: 10.1029/2010WR010288, 2011.

Michelutti, N., Douglas, M. S., Lean, D. R., and Smol, J. P.: Physical and chemical limnology of 34 ultra-oligotrophic lakes and ponds near Wynniatt Bay, Victoria Island, Arctic Canada. *Hydrobiologia*, 482(1), 1-13, 2002.

Morison, M. Q., Macrae, M. L., Petrone, R. M., and Fishback, L.: Seasonal dynamics in shallow freshwater pond-peatland hydrochemical interactions in a subarctic permafrost environment. *Hydrol. Process.*, 31, 462-475, 2017.

Muster, S., Heim, B., Abnizova, A., and Boike, J.: Water body distributions across scales: a remote sensing based comparison of three arctic tundra wetlands. *Remote Sensing*, 5(4), 1498-1523, 2013.

Niu, F., Lin, Z., Liu, H., and Lu, J.: Characteristics of thermokarst lakes and their influence on permafrost in Qinghai–Tibet Plateau. *Geomorphology*, 132(3), 222-233, 2011.

O'Brien, W., Barfield, M., Betez, N., Hershey, A. E., Hobbie, J. E., Kipphut, G., Kling, G., and Miller, M. C.: Long-term response and recovery to nutrient addition of a partitioned arctic lake. *Freshwater. Biol.*, 50(5), 731-741, 2005.

Pace, M. L., and Cole, J. J.: Synchronous variation of dissolved organic carbon and color in lakes. *Limnol. Oceanogr.*, 47(2), 333–342, 2002.

Paterson, A. M., Keller, W., Rühland, K. M., Jones, F. C., and Winter, J. G.: An exploratory survey of summer water chemistry and plankton communities in lakes near the Sutton River, Hudson Bay Lowlands, Ontario, Canada. *Arct. Antarct. Alp. Res.*, 46(1), 121-138, 2014.

Pienitz, R., Smol, J. P., and Lean, D. R.: Physical and chemical limnology of 59 lakes located between the southern Yukon and the Tuktoyaktuk Peninsula, Northwest Territories (Canada). *Can. J. Fish. Aquat. Sci.*, 54(2), 330-346, 1997.

Pokrovsky, O. S., Shirokova, L. S., Kirpotin, S. N., Audry, S., Viers, J., and Dupré, B.: Effect of permafrost thawing on organic carbon and trace element colloidal speciation in the thermokarst lakes of western Siberia. *Biogeosciences*, 8(3), 565-583, 2011.

Przytulska, A., Comte, J., Crevecoeur, S., Lovejoy, C., Laurion, I., and Vincent, W.F.: Phototrophic pigment diversity and picophytoplankton in permafrost thaw lakes. *Biogeosciences*, 13, 13-26, 2016.

Quinton, W. L., and Pomeroy, J. W.: Transformations of runoff chemistry in the Arctic tundra, Northwest Territories, Canada. *Hydrol. Process.*, 20(14), 2901-2919, 2006.

R Development Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria; available at: <http://www.R-project.org>, 2016.

Rautio, M., Dufresne, F., Laurion, I., Bonilla, S., Vincent, W. F., and Christoffersen, K. S.: Shallow freshwater ecosystems of the circumpolar Arctic. *Écoscience*, 18, 204–222, 2011.

Roiha, T., Laurion, I., and Rautio, M.: Carbon dynamics in highly heterotrophic subarctic thaw ponds. *Biogeosciences*, 12(23), 7223-7237, 2015.

Sauchyn, D., and Kulshreshtha S.: Prairies. In *From Impacts to Adaptation: Canada in a Changing Climate 2007*. Government of Canada: Ottawa, 2008.

Schindler, D. W., and Smol, J. P.: Cumulative effects of climate warming and other human activities on freshwaters of Arctic and subarctic North America. *Ambio*, 35(4), 160-168, 2006.

Smith, L. C., Sheng, Y., MacDonald, G. M., and Hinzman, L. D.: Disappearing arctic lakes. *Science*, 308(5727), 1429-1429, 2005.

Smol, J. P., and Douglas, M. S.: Crossing the final ecological threshold in high Arctic ponds. *P. Natl. Acad. Sci.*, 104(30), 12395-12397, 2007.

Sokal, M. A., Hall, R. I., and Wolfé, B. B.: Inter-annual and seasonal effects of flooding on the water chemistry, diatom phytoplankton communities and macrophyte biomass of lakes in the Slave River Delta (Northwest Territories, Canada). *Ecohydrology*, 3: 41–54, 2010.

Spence, C., Guan, X.J., Phillips, R., Hedstrom, N., Granger, R., and Reid, B.: Storage dynamics and streamflow in a catchment with a variable contributing area. *Hydrol. Process.*, 24, 2209-2221, 2010.

Spiess, A. N., and Neumeier, N.: An evaluation of  $R^2$  as an inadequate measure for nonlinear models in pharmacological and biochemical research: a Monte Carlo approach. *BMC Pharmacology*, 10(1), 6, 2010.

Stark, S.: Nutrient cycling in the tundra. In *Nutrient Cycling in Terrestrial Ecosystems*. Springer, Berlin, Germany, 309-331, 2007.

Symons, C. C., Amott, S. E., and Sweetman, J. N.: Nutrient limitation of phytoplankton communities in subarctic lakes and ponds in Wapusk National Park, Canada. *Polar. Biol.*, 35(4), 481-489, 2012.

Thompson, D. K., and Woo, M. K.: Seasonal hydrochemistry of a high Arctic wetland complex. *Hydrol. Process.*, 23(10), 1397-1407, 2009.

Viner, A. B.: Laboratory experiments on the effects of light and temperature on the uptake of nutrients by Lake Rotongaio phytoplankton. *New. Zeal. J. Mar. Fresh.*, 18(3), 323-340, 1984.

Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot, M., Billet, M. F., Canário, J., Cory, R. M., Deshpande, B. N., Helbig, M., Jammot, M., Karlsson, J., Larouche, J., MacMillan, G., Rautio, M., Walter Anthony, K. M., and Wickland, K. P.: Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems. *Biogeosciences*, 12(23), 7129-7167, 2015.

Wallenstein, M.D., McMahon, S.K., and Schimel, J.P.: Seasonal variation in enzyme activities and temperature sensitivities in Arctic tundra soils. *Glob. Chang. Biol.*, 15(7), 1631-1639, 2009.

Walvoord, M. A., and Kurylyk, B. L.: Hydrologic impacts of thawing permafrost—A review. *Vadose Zone J.*, 15(6), 2016.

Wetzel, R. G.: *Limnology: lake and river ecosystems*. Gulf Professional Publishing, United States, 2001.

White, J., Hall, R. I., Wolfé, B. B., Light, E. M., Macrae, M. L., and Fishback, L.: Hydrological connectivity and basin morphometry influence seasonal water-chemistry variations in tundra ponds of the northwestern Hudson Bay Lowlands. *Arct. Antarct. Alp. Res.*, 46(1), 218-235, 2014.

Wickland, K. P., Aiken, G. R., Butler, K., Dornblaser, M. M., Spencer, R. G. M., and Striegl, R. G.: Biodegradability of dissolved organic carbon in the Yukon River and its tributaries: Seasonality and importance of inorganic nitrogen. *Global. Biogeochem. Cy.*, 26(4), GB0E03, doi:10.1029/2012GB004342, 2012.

Wiklund, J. A., Hall, R. I., and Wolfé, B. B.: Timescales of hydrolimnological change in floodplain lakes of the Peace-Athabasca Delta, northern Alberta, Canada. *Ecohydrology*, 5: 351–367, 2012.

Wolfé, B. B., E. M. Light, M. L. Macrae, R. I. Hall, K. Eichel, S. Jasechko, J. White, L. Fishback, and Edwards, T. W. D. Divergent hydrological responses to 20th century climate change in shallow tundra ponds, western Hudson Bay Lowlands. *Geophys. Res. Lett.*, 38(23), L23402, doi:10.1029/2011GL049766, 2011.

Woo, M. K., and Guan, X. J.: Hydrological connectivity and seasonal storage change of tundra ponds in a polar oasis environment, Canadian High Arctic. *Permafrost. Periglac.*, 17(4), 309-323, 2006.



Yoshikawa, K., and Hinzman, L. D.: Shrinking thermokarst ponds and groundwater dynamics in discontinuous permafrost near Council, Alaska. *Permafrost. Periglac.*, 14(2), 151-160, 2003.

**Tables**

**Table 1. Physical characteristics and locations of study ponds near Churchill, Manitoba.**

Pond	Area (m <sup>2</sup> )	Perimeter (m)	Shoreline development* (m <sup>1</sup> m <sup>-1</sup> )	Average depth, $\mu \pm \sigma$ (cm)	Catchment area (m <sup>2</sup> )	Coordinates
Erin	8479	448	1.37	No data	108160	58°43'14.3"N 93°50'20.2"W
Left	752	130	1.34	30.0 ± 6.8	5737	58°44'44.2"N 93°49'20.6"W
Strange	6307	424	1.51	22.2 ± 10.0	20441	58°43'36.8"N 93°50'20.6"W
Sandwich	17146	548	1.18	19.6 ± 4.8	34638	58°43'24.2"N 93°50'34.4"W
Frisbee	4416	258	1.10	14.2 ± 2.8	31128	58°43'35.5"N 93°50'33.9"W
Larch	4066	333	1.47	14.2 ± 4.9	15228	58°43'26.9"N 93°50'41.5"W

Formatted Table

\*Shoreline development,  $D_L$ , is the ratio of measured shoreline perimeter, P, of a given pond to the shoreline perimeter of a perfectly circular pond of equal area, A, such that  $D_L = \frac{P}{2\sqrt{\pi A}}$  (Aronow, 1982)

Table 2. Correlation coefficients among ponds for all species. Values are bolded when the correlation is significant at the  $p < 0.01$  level.

Species/Pond	Erin	Frisbee	Larch	Left	Sandwich
<b>DON</b>					
▲ Frisbee	<b>0.761</b>				
▲ Larch	<b>0.802</b>	<b>0.687</b>			
▲ Left	<b>0.794</b>	<b>0.912</b>	<b>0.846</b>		
▲ Sandwich	<b>0.860</b>	<b>0.689</b>	<b>0.965</b>	<b>0.829</b>	
▲ Strange	<b>0.763</b>	<b>0.904</b>	0.659	<b>0.900</b>	<b>0.688</b>
<b>Cl<sup>-</sup></b>					
▲ Frisbee	<b>0.859</b>				
▲ Larch	<b>0.762</b>	<b>0.751</b>			
▲ Left	<b>0.875</b>	<b>0.787</b>	<b>0.884</b>		
▲ Sandwich	<b>0.649</b>	0.539	<b>0.941</b>	<b>0.851</b>	
▲ Strange	<b>0.774</b>	<b>0.680</b>	<b>0.751</b>	<b>0.889</b>	<b>0.746</b>
<b>SO<sub>4</sub><sup>2-</sup></b>					
▲ Frisbee	<b>0.745</b>				
▲ Larch	-0.056	-0.007			
▲ Left	<b>0.740</b>	<b>0.836</b>	-0.271		
▲ Sandwich	-0.415	-0.388	0.134	-0.285	
▲ Strange	<b>0.731</b>	<b>0.728</b>	-0.133	<b>0.958</b>	-0.318
<b>Na<sup>+</sup></b>					
▲ Frisbee	<b>0.863</b>				
▲ Larch	<b>0.667</b>	<b>0.625</b>			
▲ Left	<b>0.875</b>	<b>0.765</b>	<b>0.818</b>		
▲ Sandwich	<b>0.777</b>	<b>0.643</b>	<b>0.920</b>	<b>0.948</b>	
▲ Strange	<b>0.843</b>	<b>0.682</b>	<b>0.711</b>	<b>0.963</b>	<b>0.897</b>
<b>K<sup>+</sup></b>					
▲ Frisbee	<b>0.847</b>				
▲ Larch	0.561	0.490			
▲ Left	0.401	0.261	<b>0.584</b>		
▲ Sandwich	0.428	0.562	<b>0.794</b>	<b>0.523</b>	
▲ Strange	<b>0.614</b>	0.563	<b>0.834</b>	<b>0.734</b>	<b>0.725</b>
<b>Mg<sup>2+</sup></b>					
▲ Frisbee	<b>0.935</b>				
▲ Larch	<b>0.830</b>	<b>0.881</b>			
▲ Left	<b>0.912</b>	<b>0.944</b>	<b>0.878</b>		
▲ Sandwich	<b>0.904</b>	<b>0.905</b>	<b>0.946</b>	<b>0.952</b>	
▲ Strange	<b>0.929</b>	<b>0.905</b>	<b>0.831</b>	<b>0.979</b>	<b>0.939</b>
<b>Ca<sup>2+</sup></b>					
▲ Frisbee	0.138				
▲ Larch	<b>0.700</b>	0.330			
▲ Left	0.278	0.571	0.102		
▲ Sandwich	0.226	0.433	0.354	0.274	
▲ Strange	<b>0.772</b>	0.419	<b>0.723</b>	0.340	0.566
<b>NH<sub>4</sub><sup>+</sup></b>					
▲ Frisbee	<b>0.715</b>				
▲ Larch	<b>0.605</b>	<b>0.619</b>			
▲ Left	-0.135	-0.250	0.261		
▲ Sandwich	0.000	-0.003	-0.256	-0.103	
▲ Strange	-0.449	-0.040	-0.277	-0.024	-0.325

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted Table

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

	$\text{NO}_3^-$				
Frisbee	0.000				
Larch	-0.087	0.320			
Left	0.404	-0.346	-0.366		
Sandwich	0.136	-0.011	0.295	-0.289	
Strange	-0.294	-0.133	0.353	-0.504	0.292

Table 3. Correlation coefficients among species for all ponds. Values are bolded when the correlation is significant at the  $p < 0.01$  level.

Species	DON	$\text{Cl}^-$	$\text{SO}_4^{2-}$	$\text{Na}^+$	$\text{K}^+$	$\text{Mg}^{2+}$	$\text{Ca}^{2+}$	$\text{NH}_4^+$	$\text{NO}_3^-$
$\text{Cl}^-$	<b>0.710</b>								
$\text{SO}_4^{2-}$	<b>0.594</b>	<b>0.827</b>							
$\text{Na}^+$	<b>0.615</b>	<b>0.971</b>	<b>0.802</b>						
$\text{K}^+$	0.276	<b>0.504</b>	<b>0.462</b>	<b>0.595</b>					
$\text{Mg}^{2+}$	<b>0.563</b>	<b>0.892</b>	<b>0.787</b>	<b>0.940</b>	<b>0.585</b>				
$\text{Ca}^{2+}$	<b>0.605</b>	<b>0.408</b>	<b>0.531</b>	<b>0.285</b>	0.000	0.166			
$\text{NH}_4^+$	-0.097	0.034	0.039	0.000	-0.163	0.027	-0.082		
$\text{NO}_3^-$	-0.134	-0.201	-0.143	-0.207	-0.076	-0.158	-0.078	0.218	

Table 4. Values of  $\beta_2$  and root mean square error (S) from the curve  $[x] = \frac{[x_{\text{gr}}]}{[x_s]} = \beta_1(\text{stage})^{\beta_2}$  for each set of normalized concentration data  $[x]$ . Chemical species were manually classified as hydrologically driven if they shared a common (approximate)  $\beta_2$  value within a pond with a low S relative to other species in the same pond. Notable exceptions to the general classification in the table heading are italicized.

	Hydrologically driven					Not hydrologically driven			
	DON	$\text{Cl}^-$	$\text{Na}^+$	$\text{K}^+$	$\text{Mg}^{2+}$	$\text{Ca}^{2+}$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$
Frisbee									
$\beta_2$	-2.61	-2.63	-2.01	-2.52	-2.15	-0.38	-0.23	0.23	-1.11
S	0.27	0.16	0.15	0.22	0.23	0.25	0.70	0.29	0.24
Larch									
$\beta_2$	-0.35	-0.28	-0.29	-0.38	-0.44	0.39	-0.73	-0.18	0.08
S	0.16	0.06	0.09	0.09	0.08	0.14	0.65	0.29	0.52
Left									
$\beta_2$	-0.70	-1.20	-1.07	-0.21	-1.05	-0.12	-1.44	-0.29	-1.30
S	0.28	0.19	0.14	0.19	0.16	0.15	0.52	0.38	0.35
Sandwich									
$\beta_2$	<b>-0.47</b>	-0.43	-0.53	<b>-0.28</b>	-0.63	0.36	0.82	-0.35	0.85
S	<b>0.19</b>	0.08	0.05	<b>0.15</b>	0.07	0.30	0.78	0.28	0.66
Strange									
$\beta_2$	-0.16	-0.75	-0.55	-0.23	-0.57	0.42	-0.08	0.07	<b>-0.71</b>
S	0.37	0.18	0.11	0.11	0.15	0.24	0.45	0.13	<b>0.18</b>

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, English (United Kingdom)

Formatted Table

Formatted Table

Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)


Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)


Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)


Formatted: Font: (Default) Times New Roman, 10 pt, Bold, Not Italic, English (United Kingdom)

Figures






**Figure 1.** Site map of the general near-coastal tundra region with the catchment areas containing Frisbee, Larch, Strange, Sandwich, Left, and Erin ponds. Inset map highlights the entire Hudson Bay Lowlands (dark blue coloured region to the south of Hudson Bay) and the Churchill study region (star indicated with a red dot and an arrow). Map data: Google, Digital Globe, 2017.




**Figure 2.** Boxplots of spatial and temporal proportional deviations for each chemical species. Significant differences between medians ( $p < 0.01$ ) are present for all species on the left side of the panel ( $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ), and no significant difference is present for species on the right ( $\text{DON}$ ,  $\text{Mg}^{2+}$ ,  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ).

— Erin    △ Frisbee    ▲ Larch    ● Left    ○ Sandwich    □ Strange

Temporal variation  
 exceeds spatial variation



Temporal variation does  
 not exceed spatial  
 variation







Figure 3. Pond normalized concentrations of all measured chemical species in each pond and precipitation for the 2015 snow free season. In chemical species on left side of the panel (Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>) temporal variation exceeds spatial variation, and no significant difference is present for species on the right (DON, Mg<sup>2+</sup>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>).








Figure 4: Precipitation (vertical bars, top panel), as well as depth of water column (continuous line) and runoff depths (vertical bars) in each of the instrumented study ponds over the 2015 ice-free season.





Figure 5: Comparison of the stage-concentration relationship in a) hydrologically driven chemical species ( $Cl^-$ , in Frisbee pond), and b) a non-hydrologically driven species ( $NH_4^+$ , in Larch pond).







Figure 6. Principal Components Analysis of chemical concentrations in all ponds throughout the 2015 snow-free season. The sampling date corresponds to the colour ramp with earlier dates in blue and later dates in red. PC1 primarily corresponds with the hydrologically driven chemical species while PC2 corresponds to inorganic nitrogen species (NO<sub>3</sub>, NH<sub>4</sub><sup>+</sup>). The hollow symbols are highlighted with solid points.









Figure 7. Short term variation in DON, Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, and NO<sub>3</sub><sup>-</sup> concentrations in (a) Frisbee pond and (b) Strange pond during a rain event on July 3<sup>rd</sup> to 6<sup>th</sup>, 2015 (DOY 185-188, highlighted with red bars).



(a) Pond concentrations of DON, Cl<sup>-</sup>, Na<sup>+</sup>, K<sup>+</sup> and Mg<sup>2+</sup> are hydrologically controlled



(c) Short-term variability of hydrochemistry during and following storm events may exceed seasonal-term variability



(b) Adjacent ponds experience synchronous variation of Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>2+</sup>




Figure 8. Graphical summation of major findings from this study; (a) factors driving hydrological control of particular chemical species in ponds, (b) hypothesised factors driving temporal coherence in chemical species in ponds, and (c) short-term variation in hydrochemistry can exceed seasonal-scale variation despite the large difference in timescales.

Formatted: Justified, Space After: 10 pt, Line spacing: single