

Dr. Denise M. Akob  
Associate Editor, *BG*

Dear Dr. Akob,

Thank you for considering our manuscript for publication in Biogeosciences. Please find the revised version of our manuscript "Plant *n*-alkane production from litterfall altered the diversity and community structure of alkane degrading bacteria in litter layer in lowland subtropical rainforest in Taiwan", which had been resubmitted through the online submitting system. We have corrected the typo, adjusted the terminology, and response to reviewers' comments. The details of the changes are showed in the following paragraphs.

Thank you again.

Sincerely,

Cheng-Wei Fan

## **The point-by-point reply to the comments**

### **Responses to Reviewer #1**

Since most of my previous questions have been addressed in the previous report, in this opportunity I will share some comments according to some of the facts given in the paper.

1. The title is broad when speaking about the diversity and community structure of alkane degrading bacteria. In the work you spoke about diversity at the Phylum level and I strongly believe this should be addressed in the title as well. If the title wants to remain as is, please include some Genus, or species information in the figures/analysis. This takes me to my next comment...

[Reply]

We have added information of taxonomy prediction with an estimated probability greater than 90% for all alkB-lineage-like bacteria OTUs at table S-1 at levels including genus, family, and order. It was well accepted that the majority of bacteria weren't identified and unknown. Therefore, it is reasonable that some OTUs didn't have a taxonomy prediction with an estimated probability greater than 90% at genus level. We would like to keep the title as it is.

2. The metagenome data that has been mentioned has great potential and should be exploited a bit more. A small analysis that would include assembly and binning of contigs would be required but would give some information on WHO the organisms are. Only with this information can the DIVERSITY and community structure of ALKANE degrading bacteria be concluded. The data you have shown (figure 8) does not show that it is actually these organisms that change. In other words, the WHOLE community and diversity is changing but no conclusions can be drawn about only alkane degrading bacteria since your gene has low variation.

[Reply]

Community amplicon sequencing analyses including assembly and binning of contigs of metagenome data have been performed in this study. More detail information about the community structure and diversity changes of alkB-lineage bacteria was showed at Table S-1, Figure S-5 and Figure S-6. The Figure S-5 and Figure S-6 provide supportive evidence that natural habitats harbor alkB-lineage-like bacteria with some species existing in very low copy numbers.

3. The introduction does not include enough information about litterfall and why it is important to study this environment. It is not clear why the study was performed with this specific environment, unless you read other studies.

[Reply]

To address the comment regarding of litterfall, we revised the text as following (pg. 3, l. 14-19).

“Besides the litterfall plays an essential role in sustaining the microorganism food chain in the forest ecosystem. The litter leaf, the main component of litterfall, similarly provide the major sources of plant-derived alkanes. The significant effects of litterfall derived alkanes on the bacterial alkane degradation potential have also been observed (Giebler et al. 2013 and Schulz et al., 2012). For the litterfalls in the habitats from the Nanjenshan Reserve were different, investigations into the effects of litterfall on various habitats and microbial communities in these habitats are possible.”

4. In the conclusions would it be appropriate to suggest some alkane bacteria activity assays from the environmental samples (RNA)?

[Reply]

A sentence has been added to the paragraph of conclusions (pg. 6, l. 13). It is:

“Practical testing of alkB gene concentration by semiquantitative PCR with primer coverage over Proteobacteria and Actinobacteria reveals a consistent result with the bioinformatics data, suggesting this method is helpful for future study on alkane bacteria activity assays from the environmental samples.”

5. I believe you have a typo in page 6, line 28: MEGA program instead of MAGA?

[Reply]

The typo has been corrected (pg. 7, l. 2 and 3).

6. Figure 2 can be in the supplementary data?

[Reply]

Figure 2 has been removed as supplementary data Figure S-2.

## Responses to Editor

Additional comments:

1. Reviewer #1 is also correct that the program is MEGA (pg. 6, l. 27-28).

[Reply]

The typo has been corrected (pg. 7, l. 2 and 3).

2. Pg. 6, l. 5 and elsewhere: Metagenomics is not the best choice of terminology in this case. Although, metagenomics in its strictest definition includes amplicon sequencing the prevailing use of the term in microbiology is in reference to genomic sequencing of multiple organisms in a single community. Change this throughout the paper to something like community amplicon sequencing.

[Reply]

Thanks for the comment. The change has been made throughout the manuscript.

3. Pg. 9, l. 19: change to bacterial community composition

[Reply]

The change has been made (pg. 19, l. 25).

4. Pg. 12, L. 3 (and elsewhere): NGS is a tool but not a technique. Change to microbial community characterization.

[Reply]

Thanks for the comment. The change has been made throughout the manuscript.

5. I agree with reviewer #1 that Figure 2 could be moved to the supplement

[Reply]

Figure 2 has been removed as supplementary data Figure S-2.

6. Figure 4: correct the legend to remove "\_" and for correct use of italics (Novel and Candidate Division are not taxonomic names)

[Reply]

The marks of "\_" have been removed and the use of italics have been corrected (Figure 3).

7. Figure 5: can you make the legend in the figure bigger so that you can see what the pattern is?

[Reply]

The legend in the figure 5 (now Figure 4) has been modified to bigger and clearer.

#### References

**Giebler, J., Wick, L. Y., Chatzinotas, A. & Harms, H.** (2013). Alkane-degrading bacteria at the soil-litter interface: comparing isolates with T-RFLP-based community profiles. *FEMS Microbiol Ecol* **86**, 45-58.

**Schulz, S., Giebler, J., Chatzinotas, A., Wick, L. Y., Fetzer, I., Welzl, G., Harms, H. & Schloter, M.** (2012). Plant litter and soil type drive abundance, activity and community structure of alkB harbouring microbes in different soil compartments. *ISME J* **6**, 1763-74.

# Plant *n*-alkane production from litterfall altered the diversity and community structure of alkane degrading bacteria in litter layer in lowland subtropical rainforest in Taiwan

Tung-Yi Huang<sup>1</sup>, Bing-Mu Hsu<sup>1\*</sup>, Wei-Chun Chao<sup>2</sup>, Cheng-Wei Fan<sup>1</sup>

5 <sup>1</sup> Department of Earth and Environmental Sciences, National Chung Cheng University, Chiayi County, Taiwan

<sup>2</sup> Department of Forestry and Natural Resources, National Chiayi University, Chiayi City, Taiwan

\*Tung-Yi Huang and Bing-Mu Hsu contributed equally to this paper.

Correspondence to: Cheng-Wei Fan (cwfan@mail.eq.ccu.edu.tw)

10 **Abstract.** *n*-Alkane and alkane-degrading bacteria have long been used as crucial biological indicators of paleoecology, petroleum pollution, and oil and gas prospecting. However, the relationship between *n*-alkane and alkane-degrading bacteria in natural forests is still poorly understood. In this study, long-chain *n*-alkane (C<sub>14</sub>-C<sub>35</sub>) concentrations in litterfall, litter layer, and topsoil and the diversity and abundance of *n*-alkane-degrading bacterial communities in litter layers were investigated in three habitats across a lowland subtropical rainforest in southern Taiwan: ravine, windward, and leeward

15 habitats in Nanjenshan. Our results demonstrate that the litterfall yield and productivity of long-chain *n*-alkane were highest in the ravine habitats. However, long-chain *n*-alkane concentrations in all habitats were decreased drastically to a similar low level from the litterfall to the bulk soil, suggesting a higher rate of long-chain *n*-alkane degradation in the ravine habitat. Operational taxonomic unit (OTU) analysis using next-generation sequencing data revealed that the relative abundances of microbial communities in the windward and leeward habitats were similar and different from that in the ravine habitat.

20 ~~Metagenomic d~~Data mining of community amplicon sequencing using the NCBI database revealed that alkB-gene-associated bacteria (95 % DNA sequence similarity to alkB-containing bacteria) were most abundant in the ravine habitat. Empirical testing of litter layer samples using semiquantitative polymerase chain reaction for determining alkB gene levels confirmed that the ravine habitat had higher alkB gene levels than the windward and leeward habitats. Heat map analysis revealed parallels in pattern color between the plant and microbial species compositions of the habitats, suggesting a causal

25 relationship between the plant *n*-alkane production and microbial community diversity. This finding indicates that the diversity and relative abundance of microbial communities in the litter layer are affected by *n*-alkane plant composition in the litterfall.

## 1 Introduction

Alkanes are saturated hydrocarbons that account for a small fraction of the total organic carbon, with a concentration range

30 in natural habitats of not more than a couple of thousand parts per million by weight (Gomez-Coca et al., 2016; Rojo,

2010). In noncontaminated habitats, long-chain *n*-alkanes with an odd/even predominance are produced by plants or algae to act as chemoattractants or agents that protect against microbial invasion or water loss (Feakins et al., 2016; Koch et al., 2009). Alkanes are exceptionally stable molecules that can survive in soil and sediment, which make them useful as biomarkers in paleoecology (Sachse et al., 2012). Because alkanes are the major components in petroleum, natural gas, and diesel fuel, the presence of environmental alkanes also serves as an index of oil contamination (Afzal et al., 2013). Although alkanes are very stable, most can be degraded by several microorganisms. Alkane-degrading bacteria are commonly upregulated when the concentration of alkanes is increased; it has thus been suggested that these bacteria be used as a biological indicator for oil and gas prospecting (Xu et al., 2013; Rasheed et al., 2012).

Extensive knowledge on the degradation of *n*-alkane by microorganisms has been acquired in the past decades. The first step of enzymatic reaction is oxidization of *n*-alkane to the corresponding primary alcohol by alkane terminal hydroxylases. This step is crucial because activation of the alkane molecule requires an enzyme system that is not widespread (Rojo, 2010). The products can be further oxidized into a corresponding aldehyde and fatty acid, which are digested by most microorganisms. Although fungi and yeasts can degrade alkanes, most research has focused on the role of bacteria in the degradation of alkanes (Rojo, 2009, 2010; Singh et al., 2012). To date, alkane monooxygenase gene (*alkB*) hydroxylases are the most commonly found alkane hydroxylases in both Gram-negative and Gram-positive bacteria. More than 60 genera of aerobic bacteria and 5 genera of anaerobic bacteria are known to degrade *n*-alkanes (Nie et al., 2014). Although alkanes can be metabolized anaerobically, the growth of anaerobic alkane degraders is significantly slower than that of aerobic ones, especially in the litter layer environment where the oxygen supply is adequate (Wentzel et al., 2007). The *alkB*-encoded membrane-bound homologous protein is an inducible integral membrane-bound alkane hydroxylase that plays a vital role in aerobic alkane degradation (Beilen et al., 2003). Evidence has revealed that the number of *alkB* bacteria dynamically changes in response to a fluctuation in environmental alkanes. In soil or water environments, when polluted with crude oil, the alkane-degrading bacteria were upregulated (Afzal et al., 2013). For example, a 10-fold increase in the relative abundance of oil degraders in contaminated versus clean sand was estimated for a Gulf of Mexico beach (Kostka et al., 2011). In oil and gas reservoirs, it has been shown that the abundance of *alkB* bacteria was substantially higher in the surface soils with no significant detection of soil alkane (Xu et al., 2013). The dynamic relationship between alkane and *alkB* bacteria in soil from an agriculture research farm and a laboratory maintained under controlled conditions has been previously studied. Giebler et al. (2013) reported that in a controlled environment, sizeable communities of alkane-degrading bacteria, especially *Proteobacteria* and *Actinobacteria*, were detected at the soil–litter interface when it was enriched with artificial alkane supplements. The researchers demonstrated that many of the alkane-degrading bacteria were not detected before the alkanes were supplemented, suggesting that some alkane degraders may have been present in very low copy numbers and below the limits of detection. It is reported that the substrate was discovered to modulate the biomass of microorganisms based on its initial concentration (Schmidt, 1992). Thus, different alkane yields of litterfall in various natural habitats may upregulate the growth of various alkane degraders. The relationship between *alkB* bacteria abundance and alkane substrate is more complicated in natural habitats. The dynamic changes in *alkB*-degrading bacteria populations were demonstrated to be driven by numerous factors, such as the original source of the alkane and soil type. Although

knowledge of the regulatory mechanisms regarding environmental plant litter on soil bacterial levels and of the dynamics of alkane-degrading bacteria in natural habitats has been beneficial and important in many aspects, studies on their dynamic changes are currently lacking and should be conducted (Schulz et al., 2012).

5 The Nanjenshan Reserve, a lowland subtropical rainforest in southern Taiwan, consists of several dynamic forest plots for which forest structure, vegetation pattern, climate, and topography data have been collected at well-established permanent study sites. Four dynamic forest plots and a transect zone in the habitat have been documented since 1989 (Chao et al., 2010; Chao et al., 2007; Chao et al., 2008; Chen and Huang, 1986; Hsieh et al., 2000; Tsui et al., 2004; Fan et al., 2005). Among them, three plots captured our attention, namely plot I (120° 50' 51" E, 22° 04' 54" N), plot II (120° 50' 36" E, 22° 04' 52" N), and the Lanjenchi plot (120° 51' 38" E, 22° 03' 23" N). A ravine habitat is located across plots I and II, whereas leeward and windward habitats are located on plot III. Surveys conducted over the past decade have demonstrated that the annual yield of litterfall in the ravine habitat was higher than in those in the windward and leeward habitats. Moreover, studies have shown that in the leeward habitat, plant height was higher and plant density was lower when compared with the windward habitat (8.41±1.73 m vs. 4.63±0.88 m; 7,505 trees ha<sup>-1</sup> vs. 20,065 trees ha<sup>-1</sup>). The plant height and density in the ravine habitat were 9.45±1.35 m and 4257 trees ha<sup>-1</sup> (Chin, 2008). Besides the litterfall plays an essential role in sustaining the microorganism food chain in the forest ecosystem. The litter leaf, the main component of litterfall, similarly provide the major sources of plant-derived alkanes. The significant effects of litterfall derived alkanes on the bacterial alkane degradation potential have also been observed (Giebler et al. 2013 and Schulz et al., 2012). For the litterfalls in the habitats from the Nanjenshan Reserve were different, investigations into the effects of litterfall on various habitats and microbial communities in these habitats are possible.~~The alkane distributions and concurrent alkane-degrading bacteria have yet been uncovered. The litterfall plays an essential role in sustaining the microorganism food chain. Because the litterfalls in the habitats were different, investigations into the effects of litterfall on various habitats and microbial communities in these habitats are possible.~~

Environmental microbial communities are highly complex and diverse and difficult to culture under laboratory conditions. Next-generation sequencing (NGS) is a powerful tool for high throughput sequencing technologies. With this recent advent ~~platform of next-generation sequencing (NGS) technology~~ and computational methods, we can conduct genome studies of microbes in these habitats, which have numerous variables such as annual litterfall productivity and plant vegetation (Degnan and Ochman, 2012). We investigated the productivity of long-chain *n*-alkane in the pristine natural habitats of the Nanjenshan Reserve and explored their relationship with microbial communities, particularly emphasizing the correlation between *n*-alkane production and bacteria containing the *alkB* gene. During the present 1 year follow-up study, gas chromatography with a flame ionization detector (GC-FID) was used to determine *n*-alkane (C<sub>14</sub>-C<sub>35</sub>) concentrations. Bioinformatics analyses and ~~metagenomic community amplicon sequencing~~ data mining were performed to identify microbial communities and read numbers of bacterial lineages that contained *alkB* genes in the three rainforest habitats. Semiquantitative polymerase chain reaction (PCR) was performed to test the results of the bioinformatics analysis.

## 2 Materials and methods

### 2.1 Sample description

A map of the sampling sites is provided in the supplementary file (Fig. ~~ure~~ S-1). Twenty-two custom-made aluminum stands with frames (0.71 m × 0.71 m) covered by a nylon mesh (1 mm i.d.) were used to collect litterfall. Twelve stands were located in the ravine habitat, whereas ten stands were located in the Lanjenchi plot. Of those on the Lanjenchi plot, six stands were placed in the windward habitat and four in the leeward habitat. The horizontal distance between stands was approximately 50 m. The number of stands in the three habitats of the Nanjenshan Reserve was mainly selected based on the stands available in this area. Each nylon mesh containing litterfall was collected and replaced monthly from early October 2012 to late September 2013. After their collection, the litterfall samples were sorted into litter-leaf, litter-branches, litter-flowers, litter-fruits, and miscellaneous fractions. The sorted samples were weighed after being oven-dried at 40 °C for 14 days (Conklin-Brittain et al., 2006). The litter-leaves were further separated into genus and species. Samples of the litter layer and underlying soils from the three habitats were collected in 2015 and 2016, respectively. The collected samples, including litter-leaf, litter layer, and soils, were subjected to *n*-alkane analysis, and litter layer samples from each habitat were further subjected to DNA extraction and consequential assays.

The annual yield (litterfall or litter-leaf) in each stand was used to calculate the productivity of each habitat. The annual litterfall and litter-leaf yields in each habitat were calculated by averaging the annual stand weights. Statistically significant differences between habitats were evaluated through Student's *t*-test using the *t*-test calculator (format SD).

### 2.2 Extraction of *n*-alkanes

All samples from the litter-leaf (42 plant species) and litter layer and all soils were dried and well homogenized. Approximately 0.5 g of sample powder was dissolved in dichloromethane (250 ml) and Soxhlet-extracted for 16 hours. A surrogate standard of 5 $\alpha$ -cholestane (Sigma) was added to each sample before extraction. The extracts were concentrated into 5–10 mL volumes using a rotary evaporator with a water bath and further concentrated to reduce the volume to approximately 1 mL, after which they were fractionated using silica gel chromatography (Silica gel 60, 3%). The *n*-alkane fraction was eluted with 4 mL of hexane and reduced to a 1 mL volume under a stream of nitrogen. An internal standard of squalene (Acros) was added to the concentrated 1 mL of hexane fraction prior to instrumental GC analysis. Quantification of the *n*-alkanes was performed using a PerkinElmer Clarus 500 gas chromatograph equipped with an autosampler, PerkinElmer Elite-5 CB fused silica capillary column (30 m in length, 0.32 mm i.d., film thickness 0.25  $\mu$ m), and flame ionization detector (FID). The GC oven temperature was programmed to increase from 70 °C to 310 °C at a rate of 4 °C min<sup>-1</sup>. The *n*-alkanes between C<sub>14</sub> and C<sub>35</sub> were identified and quantified using external standards that contained known concentrations of all *n*-alkanes of interest (Dr. Ehrenstorfer), 5 $\alpha$ -cholestane, and squalene.

The *n*-alkane concentrations in 42 litter-leaf samples were measured and then used to estimate the *n*-alkane concentration from the litter-leaf and to assess litter-leaf productivity in the three habitats. The *n*-alkane concentration in the litter-leaf



from each stand ( $C_{stand}$ , in  $\mu\text{g g}^{-1}$ ) was summed by weighting the concentration in one species in the litter-leaf by that species' proportion of the total litter-leaf collected in the stand as following [equation formula](#):

$$C_{stand} = \sum_{i=1}^{42} \frac{W_i}{W_t} \cdot C_i$$

where  $C_{stand}$  is  $n$ -alkane concentration in the litter-leaf from each stand ( $\mu\text{g g}^{-1}$ ),  $W_i$  is the dry weight of litter-leaf of each species (g),  $W_t$  is the dry weight of total litter-leaf (g), and  $C_i$  is the  $n$ -alkane concentrations ( $\mu\text{g g}^{-1}$ ) of 42 litter-leaf samples. Calculation of the litter-leaf  $n$ -alkane concentration in each habitat was then performed by averaging the  $n$ -alkane litter-leaf concentrations in all the stands. The annual  $n$ -alkane productivity in each stand was calculated by multiplying the  $n$ -alkane litter-leaf concentration in each stand by the annual yield of that stand. The  $n$ -alkane litter-leaf productivities of the stands in a habitat were then averaged to estimate the annual  $n$ -alkane productivity in the three habitats, which was evaluated using Student's t-test.

### 2.3 DNA extraction

For DNA extraction in the litter layer, several pieces of leaves from the litter layer were randomly selected, cut into 2 mm  $\times$  2 mm chips, and well blended. The DNA extraction procedure is described in detail in the user manual, *Genomic DNA from soil (Macherey-Nagel)*. Briefly, a bulk mixture (approximately 0.1–0.2 g) was transferred to a tube containing ceramic beads. Lysis buffer (lysis buffer SL1, 700  $\mu\text{L}$ ) and enhancer SX (150  $\mu\text{L}$ ) were added to each sample and vortexed for 5 min at room temperature. Precipitate contaminants were removed through centrifugation for 2 min at 11,000 g. DNA supernatant was transferred to a new tube, 150  $\mu\text{L}$  of SL3 was added, and the mixture was vortexed for 5 s before precipitation by centrifugation for 1 min at 11,000 g. The supernatant inhibitors were removed using an inhibitor removal column. The column that bound the DNA was washed four times and dried before DNA elution. DNA elution was performed by centrifugation for 30 s at 11,000 g after incubation with Buffer SE (50  $\mu\text{L}$ ). A final DNA extraction (approximately 30–100  $\text{ng uL}^{-1}$  in 100  $\mu\text{L}$  of Tris buffer) was eluted.

### 2.4 Sequencing using an Illumina MiSeq platform

The genomic DNA (12.5 ng) obtained from each habitat was used in amplicon PCR experiments, which were performed in triplicate. We performed PCR amplification of 16S rRNA gene sequences at the V3–V4 regions using Illumina's MiSeq system to create paired-end sequencing data. The experimental protocol was modified from the Illumina manual. The target sequence was amplified using PCR with mixed forward and reverse primers. The sequences of forward primers included an adapter overhang nucleotide sequence and V3–V4 primer pairs, which were 5'-TCGTC GGCAG CGTCA GATGT GTATA AGAGA CAGCC TACGG GNGGC WGCAG, 5'-TCGTC GGCAG CGTCA GATGT GTATA AGAGA CAGAC CTACG GGNGG CWGCA G, 5'-TCGTC GGCAG CGTCA GATGT GTATA AGAGA CAGTD CCTAC GGGNG GCWGC AG, and 5'-TCGTC GGCAG CGTCA GATGT GTATA AGAGA CAGGD RCCTA CGGGN

GGCWG CAG. The sequences of the reverse primers were 5'-GTCTC GTGGG CTCGG AGATG TGTAT AAGAG ACAGG ACTAC HVGGG TATCT AATCC, 5'-GTCTC GTGGG CTCGG AGATG TGTAT AAGAG ACAGT GACTA CHVGG GTATC TAATC C, 5'-GTCTC GTGGG CTCGG AGATG TGTAT AAGAG ACAGA CGACT ACHVG GGTAT CTAAT CC, and 5'-GTCTC GTGGG CTCGG AGATG TGTAT AAGAG ACAGG TTGAC TACHV GGGTA TCTAA TCC. After separation using agarose gel electrophoresis, PCR products with expected sizes were purified from the matrix. The Illumina Nextera XT index kit was used in second-stage PCR for the addition of the index. Capillary electrophoresis followed by a fluorescence-based method was employed to qualify and quantify libraries, respectively. Nucleotides were sequenced using the MiSeq sequencer for 18 dark cycles and 350 read cycles in the forward read, and 18 dark cycles and 250 read cycles in the reverse read. The data of forward and reverse reads were aligned using the analysis platform from (Genomic Workbench v.8.5) CLC bio with Q20 as a threshold to generate output FASTA files.

## **2.5 ~~Metagenomics library~~The construction and analysis of community amplicon sequencing library**

FASTA files were further processed using the sequence analysis tool, USEARCH. We merged all sequence files, removed duplicates, and clustered the sequences into operational taxonomic units (OTUs) at 97 % pairwise identity with a minimum cluster size of two to construct an OTU reference library. Then, a comparison between samples and the reference library at a 97 % sequence identity level was performed to create OTU tables, which contained the number of DNA sequence reads of each OTU. A 16s UTAX reference database was employed as a blast (Basic Local Alignment Search Tool) library. Finally, these data were combined to determine the relative abundances and subsequent visual patterns of a heat map. Principal coordinate analysis (PCoA) was used to identify the relative distance between OTUs. To plot PCoA figures, the `calc_distmx` command together with a `phylip_lower_triangular` parameter was used to calculate the distance of the OTUs. The axes file contained plotting coordinates and was derived from mothur using the PCoA command. The PCoA figures were plotted using R program.

## **2.6 Proportion of alkB-gene-containing bacteria in the habitats estimated using the NCBI database**

~~Metagenomic d~~Data mining for community amplicon sequencing was performed to search for microbial communities that may have had *alkB* genes. Because bacteria in the same genus may have similar gene sequences and functions, nucleotide blasting was performed to search for homogeneous sequences in the OTUs. Bacteria with DNA sequences 95 % or more similar to a known gene were defined as lineage-like bacteria. The NCBI database was used to construct an *alkB* gene reference library. We collected representative 16s gene sequences of all known *alkB*-gene-containing bacteria in the phyla *Actinobacteria* and *Proteobacteria* from the current NCBI database to investigate the *alkB* gene family populations in different habitats. The *alkB* gene sequences in bacterial 16s rRNA were downloaded to a reference nucleotide file and combined with the OTU libraries of *Actinobacteria* and *Proteobacteria*.

To search the alkB family gene-like lineage in the current OTU database, sequence alignment and phylogenetic analysis of the alkB nucleotide sequences were conducted using the Molecular Evolutionary Genetics Analysis 7 (~~MAGA-MEGA~~ 7) program. ~~MAGA-MEGA~~ 7 used parsimony, neighbor-joining, and maximum likelihood analyses to create a 16S rRNA phylogenetic gene tree. The NCBI nucleotide blast program was employed to test the similarity of DNA sequences between OTUs and adjacent references. DNA sequences with a similarity of more than 95 % were manually selected to determine the alkB-lineage OTU DNA sequence numbers read in the library. The Shannon–Wiener diversity index was used to calculate the plant diversity, overall bacteria communities, and alkB-lineage bacteria and was determined using the PRIMER-5 software (Plymouth Routines in Multivariate Ecological Research).

### 2.7 Semiquantitative PCR to determine alkB gene levels in litter layer samples

Because alkB genes are highly abundant and diverse, a standard method for alkB gene quantification was not available. We employed semiquantitative PCR for measuring environmental alkB in a method that was modified from previous studies because the alkB-targeting primer covers *Proteobacteria* and *Actinobacteria* (Kloos et al., 2006; Jurelevicius et al., 2013). Litter layer samples from three habitats in Nanjenshan were subjected to semiquantitative PCR to quantify the DNA levels of alkB genes in the natural habitats. Litter layer samples from different stands in each habitat were randomly selected for sample collection at three time points. After collection, litter layer samples were sent immediately to the laboratory at room temperature. Litter layer patches were randomly selected and cut into approximately 2 mm × 2 mm chips. Samples (250 mg) were taken for DNA extraction. Following DNA elution, 100 µL of buffer solution and 2 µL of eluted DNA were taken for semiquantitative PCR. The DNA was mixed with an alkB gene primer set (forward primer: 5'- AAY ACN GCN CAY GAR CTN GGN CAY AA -3', reverse primer: 5'- GCR TGR TGR TCN GAR TGN CGY TG -3', 1 µL, 0.4 µM), 5 µL Fast-Run Taq Master Mix with Dye, and 16 µL ddH<sub>2</sub>O with the final reaction volume of 25 µL. Thermal cycling conditions for PCR were an initial temperature of 95 °C for 5 min, followed by 30, 27, or 24 cycles of 95 °C for 30 s, 55 °C for 30 s, 72 °C for 1 min, and finally 72 °C for 5 min. The expected amplicons of the PCR were 548 bps. Following PCR, aliquots (5 µL) of each amplicon were confirmed using electrophoresis on 1.5 % agarose gel followed by 0.5 µg/mL SYBR Safe DNA staining and imaging using an ultraviolet transillumination system. The band density average levels were compared with a windward habitat density setting of 100 %. Statistically significant differences were evaluated using Student's t-test.

## 3 Results and Discussion

### 3.1 Quantitative estimates of annual litterfall in the Nanjenshan Reserve

The annual average litterfall was estimated from the litterfall at the sampling sites. The annual yields of litterfall in the three habitats in Nanjenshan are illustrated in Fig. 1. As shown in the records from the past decade, the litterfall was higher in the ravine habitat than in the other habitats. Our results on the litterfall of Nanjenshan indicated that the annual

weight productivity in the ravine habitat was approximately 30 % higher than in the leeward or windward habitats, which is consistent with previous studies. According to records from 1999 to 2007, seasonal litterfall in Nanjenshan ranged from 2–7 (ton ha<sup>-1</sup>); the increases of litterfall were contributed by typhoons, precipitation, and the effects of monsoon seasons. In this study, the annual litterfall was discovered to be approximately 7–10 (ton ha<sup>-1</sup>), which is considered a high-volume when compared with the records. The primary reason for the large amount of litterfall during this study was the occurrence of typhoons. Because typhoons increase litterfall in the three habitats equally, the ranking in annual litterfall volume was not altered by the advent of typhoons. One of the reasons that the ravine habitat had more annual litterfall may be the topography of the ravine habitat, which is more suitable for plant growth than that of the other habitats. The average height and diameter of cross sections of plants in the ravine habitat were the largest of those in all the habitats. Because litter-leaf was the major part of litterfall and average *n*-alkane resources, we focused on *n*-alkane levels in the litter-leaf in these habitats.

### 3.2 Example of *n*-alkane measurements in the Nanjenshan Reserve

The *n*-alkane concentrations in litter-leaf from 42 plants were measured using GC-FID. Figure S-2 presents representative GC chromatograms of the *Ilex rotunda* leaves in each habitat. The recovery rates from the surrogate of 5 $\alpha$ -cholestane ranged from 70 to 108 % and averaged at 99  $\pm$  7 %, suggesting satisfactory extraction efficiency. The *n*-alkane concentrations (ranging from 123  $\mu$ g g<sup>-1</sup> to 2,694  $\mu$ g g<sup>-1</sup>) calculated in this study were not corrected by the recovery rate. All samples exhibited an odd/even carbon-number predominance, with maximums at C<sub>31</sub>, C<sub>29</sub>, and C<sub>33</sub>. *n*-Alkanes of C<sub>14</sub> to C<sub>26</sub> were present in only relatively low amounts. Further distribution analyses of the *n*-alkane litter-leaf components, such as the carbon preference index (CPI) and C<sub>29</sub>/C<sub>33</sub> ratio, for each species were needed to further identify the relationships between plant species and *n*-alkane distribution patterns. This paper is the first to reveal the *n*-alkane levels of plant vegetation in Nanjenshan.

### 3.3 Annual litter-leaf productivity and *n*-alkane productivity rankings

A total of 42 plant species were identified in this study. Data from the plant species revealed the total amount of leaves in the annual litterfall, and the resultant ranking of leaf *n*-alkane concentrations in these plants is presented in Table 1. The leaf *n*-alkane concentration and dry weight data were employed to make a gross estimate of the *n*-alkane productivity of the litter-leaf in the Nanjenshan Reserve. The *n*-alkane concentration was discovered to change during leaf development and was affected by environmental parameters (Hoffmann et al., 2013; Jetter and Schäffer, 2001; Kahmen et al., 2011). For example, temperature and relative humidity affected the composition of *n*-alkanes in both *Acacia* and *Eucalyptus* in Australia (Hoffmann et al., 2013). In our pilot study, the *n*-alkane concentration variation among plant species was less than 15 % of the standard error, as determined through comparison of several samples from different stands and habitats for the same species. The ranges of leaf *n*-alkane concentrations between species were from 2,694 ( $\mu$ g g<sup>-1</sup>) in *Ilex rotunda*

to 123 ( $\mu\text{g g}^{-1}$ ) in *Alniphyllum pterospermum*. These data demonstrated that intraspecies variation in *n*-alkane concentration could be ignored compared with the interspecies variation. Therefore, using a representative plant *n*-alkane concentration was reasonable for estimating the overall litter-leaf *n*-alkane production in a habitat. When comparing the plant vegetation and leaf *n*-alkane concentration, the plants with leaves that had a high *n*-alkane concentration tended to grow in the ravine habitat; *Ilex rotunda* and *Celastrus kusanoi*, for example.

### 3.4 Estimates of annual *n*-alkane productivity in the three habitats

Figure 3-2 presents the *n*-alkane concentrations from the litter-leaf to the bulk soil and the average litter-leaf *n*-alkane productivity in the three habitats. The average litter-leaf *n*-alkane concentration was highest in the ravine habitat. Moreover, the *n*-alkane concentration in different layers formed a steep gradient from the litterfall to the bulk soil. The *n*-alkane flux in the litterfall in the ravine habitat was approximately twice that in the other habitats. We concluded that the ravine habitat had higher *n*-alkane input than the other habitats. This characteristic offers us an example of higher *n*-alkane input in the natural habitat to investigate the consequence of microbial community.

The balance between litterfall yield and decomposition affects the development of organic carbon in soil layers. This study and previous reports have indicated that the litterfall in ravine habitats is higher than that in other habitats. However, a study of these habitats demonstrated that the total organic carbon in the litter layer and bulk soil in the ravine habitat was equal to or less than that in the windward and leeward habitats (Guo, 2010). Therefore, decomposition effects may be stronger in ravine habitats. Figure 32(a) reveals that the *n*-alkane concentration decreased significantly from litter-leaf to surface soil, which could be due to the effects of bacteria. It is plausible that other organic compounds were degraded in addition to the *n*-alkanes. Several effects such as leaching, erosion, and sediment transportation could play roles in governing organic compound accumulation; therefore, we cannot rule out the diffusion effects of other physical factors (Kirkels et al., 2014; Quinton et al., 2006). We performed NGS-based analysis and bioinformatics studies to discover the roles of microbial communities in *n*-alkane degradation.

### 3.5 Bacterial community composition in the Nanjenshan Reserve

NGS-based study was performed to reveal the microbial communities in the litter layer from the three habitats. Figure 4-3 shows the relative abundance in community amplicon sequencing metagenomic data of OTUs that were grouped by phylum. More detailed information about the relative abundance in class, order, and family levels is presented in Figure-Fig\_S-45. Interestingly, a similarity was identified in microbial community structure between the windward and leeward habitats. Because the windward and leeward habitats are located on the same plot of a different orography, several parameters such as plant vegetation and soil properties were similar between the habitats. As mentioned previously, the most considerable differences between the windward and leeward habitats were the average height of plants and average number of trees per

hectare. The physical parameters that affect plant growth in mountainous areas did not appear to affect the relative abundance of microbial communities. Conversely, the relative abundance of microbial communities in the ravine habitat was different from that in the windward and leeward habitats, with more members of the *Actinobacteria* phyla and fewer members of the *Proteobacteria* and *Acidobacteria* phyla. The more abundant microbial communities of the *Actinobacteria* phyla may have been because of higher *n*-alkane productivity in the ravine habitat. Although many phyla have been identified as carrying the *alkB* gene, our data revealed that the only microbial communities carrying the *alkB* gene in our study were *Proteobacteria* and *Actinobacteria*.

### 3.6 Prediction of bacteria carrying *alkB* genes in Nanjenshan Reserve

The relative abundance of bacteria containing the *alkB* gene in the *Actinobacteria* and *Proteobacteria* phyla from different habitats is illustrated in [Figure-Fig. 54](#). The phylogenetic trees of OTUs in the *Proteobacteria* and *Actinobacteria* phyla with *alkB* reference strains are shown in [Figures-Fig. S-3](#) and [S-4](#). The nucleotide sequence similarities of OTUs and the adjacent *alkB* reference strains were manually blasted. Table S-1 reveals the nomenclature of OTUs in *Proteobacteria* and *Actinobacteria* with nucleotide sequences that had similarity of higher than 95 % when compared with adjacent *alkB* reference strains. [Figure 5-4](#) illustrates that a significant number of reads in the OTUs whose sequences were similar to those of *alkB*-gene-containing bacteria was identified for the ravine habitat. Although the relative abundance of *Proteobacteria* in the ravine habitat was 20 % less than that in the other habitats ([Figure-Fig. 43](#)), the number of DNA sequence reads of *alkB*-lineage-like bacteria in the ravine habitat was greater than 1.5 times that in the windward and leeward areas ([Figure-Fig. 54](#)). To summarize, there were twice as many DNA sequence reads of *alkB*-lineage-like bacteria in the ravine habitat than in the other habitats. Despite the relative abundance of DNA sequence reads in many OTUs of the *alkB*-lineage-like bacteria, *Proteobacteria*, and *Actinobacteria* phyla were limited in the leeward and windward habitats ([Figure-Fig. 54](#)); these two habitats had small amounts of *alkB*-lineage-like bacteria (approximately 0.1 % relative abundance) in most OTUs (Table S-1). Our data provide evidence that there is a universal presence of alkane degraders in natural habitats.

### 3.7 PCoA of bacteria carrying *alkB* genes

PCoA was used to visualize the similarities of DNA sequences in *Proteobacteria* and *Actinobacteria* from the three habitats. The biodiversity index in OTUs of *alkB*-lineage-like bacteria and other sample types is shown in Table S-2. [Figure 6-5](#) displays the distribution of 240 OTUs in the *Proteobacteria* and *Actinobacteria* phyla. The majority of OTUs (>90%) are present in two or three habitats. An initial capital letter in [Figure-Fig. 6-5](#) was used to denote the OTUs with the most read numbers for the habitats. The nomenclature of the *alkB*-lineage-like bacteria OTUs and relative locations are presented in the figure. Thirty-three OTUs contained *alkB*-lineage-like bacteria in this study. The numbers of OTUs that contained *alkB*-lineage-like bacteria in the ravine, windward, and leeward habitats were 30, 30, and 25, respectively. The Shannon-

Weiner index of OTUs that contained alkB-lineage-like bacteria in the ravine, windward, and leeward habitats was 2.51, 2.39, and 2.46, respectively. Although the relative abundance of OTUs that contained alkB-lineage-like bacteria in the ravine habitat was more than twice that in the windward and leeward habitats (Figure-Fig. 54), the Shannon–Weiner indices of the three habitats were similar, suggesting that the effective species numbers of alkB-lineage-like bacteria were almost identical. The results indicate the pre-existence of diverse alkB-gene-containing bacteria in natural habitats, with some bacteria so small in number that they are undetectable using normal PCR. They could, however, proliferate in the habitats with the appropriate substrate supplements. This finding is consistent with the results from a previous study using a laboratory controlled system and an agricultural research farm (Schulz et al., 2012; Giebler et al., 2013).

### 3.8 Empirical testing of bacteria carrying alkB gene

In the bacterial metagenomics-community amplicon sequencing study, equal amounts of DNA were amplified to estimate the bacterial communities in each habitat. To compare the productivity of the alkB-gene-containing bacteria in different habitats, we conducted semiquantitative PCR to determine relative alkB gene concentrations. Figure 76(a) shows a triplicate study of DNA staining of amplicons from the three habitats after various semiquantitative PCR cycles. No bands were detected if fewer than 24 amplification cycles were performed. Marginal PCR products could be detected by increasing the cycle number to 27. Figure 76(b) shows the statistical results regarding alkB gene levels in the three Nanjenshan habitats and confirms that bacterial alkB numbers were higher in the ravine habitat than in the other two habitats. We have thus provided evidence that litterfall and plant vegetation increase the abundance of alkane-degrading bacteria communities in natural habitats.

### 3.9 Exploring the relationship between plant vegetation and microbial communities using heat map analysis

Heat map analysis of plant species and read-number of OTU versus habitat were performed to investigate the relationship between plant vegetation patterns and microbial communities in different habitats. The heat map presented in Figure-Fig. 87 was based on the microbial community abundance and plant productivity of litter-leaf from different habitats, with the color intensity indicating the number of reads or weight of litter-leaf (in logarithmic form). In Figure-Fig. 87(b), the color patterns of litter-leaf yields show that microbial abundance was similar in the windward and leeward habitats, demonstrating the similarity between the microbial community compositions in these two habitats. Furthermore, the litterfall data were also similar between the two habitats, which suggests a connection between plant vegetation species and microbial composition. The pattern similarities in the plant and bacterial species can be easily explained by substrate-specific-induced growth responses in the natural habitats. The data provided supportive evidence that diversity of bacteria in all OTUs was affected by plant vegetation, not only in quantity but also in species. In Figures-Fig. 32, 54, and 76, an increase in *n*-alkane productivity is demonstrated to increase the number of alkB-gene-containing bacteria. In Figure-Fig. 87, we showed that a substrate-specific relationship may exist between plant vegetation and microbial communities. Thus,

a leaf in a natural habitat may upregulate some alkB-gene-containing bacteria while downregulating other types of alkB-gene-containing bacteria. A future study could include a litter-bag study that would enable better understanding of the relationship between individual factors within plant and bacterial species; for instance, one could study the interaction of a specific leaf species with a particular genus of *n*-alkane-degrading bacteria or microbial community in different habitats.

## 5 4 Conclusions

We confirmed that the annual litterfall productivity of the ravine habitat was approximately 30 % higher than that of the windward and leeward habitats, whereas the long-chain *n*-alkane productivity in the ravine habitat was twice that in the other habitats in a lowland subtropical rainforest of Taiwan. Plant vegetation not only affects *n*-alkane productivity but also changes litter layer microbial communities. In the ravine habitat, high litter-leaf *n*-alkane productivity resulted in an increase in the *n*-alkane-degrading bacteria in the litterfall. Increases in alkB-gene-containing bacteria were identified primarily in the *Proteobacteria* and *Actinobacteria* phyla, with NGS-based analysis and semiquantitative PCR employed for determining alkB gene concentrations. Biodiversity analyses showed that the diversity of bacteria containing the alkB gene was similar among the different habitats, suggesting that natural habitats harbor diverse alkB-lineage-like bacteria with some species existing in very low copy numbers. [Practical testing of alkB gene concentration by semiquantitative PCR with primer coverage over \*Proteobacteria\* and \*Actinobacteria\* reveals a consistent result with the bioinformatics data, suggesting this method is helpful for future study on alkane bacteria activity assays from the environmental samples.](#) Cross analyses regarding the quantity and quality of plant vegetation and structures of the microbial communities revealed that *n*-alkane-induced increases in alkB-gene-containing bacteria might have been substrate specific. To our knowledge, this is the first report on the regulation of *n*-alkane-degrading bacteria in litterfall in natural habitats.

## Acknowledgements

This manuscript was edited by Wallace Academic Editing.

## References

- 25 Afzal, M., Yousaf, S., Reichenauer, T. G., and Sessitsch, A.: Ecology of Alkane-Degrading Bacteria and Their Interaction with the Plant, *Molecular Microbial Ecology of the Rhizosphere*, 92, 975-989, 10.1002/9781118297674.ch92, 2013.
- Beilen, J. B. v., Li, Z., Duetz, W. A., Smits, T. H. M., and Witholt, B.: Diversity of Alkane Hydroxylase Systems in the Environment, *Oil & Gas Science and Technology*, 58 427-440, 2003.



- Chao, K.-J., Chao, W.-C., Chen, K.-M., and Hsieh, C.-F.: Vegetation Dynamics of a Lowland Rainforest at the Northern Border of the Paleotropics at Nanjenshan, Southern Taiwan, *Taiwan Journal of Forest Science*, 25, 29-40, 2010.
- Chao, W.-C., Chao, K.-J., Song, G.-Z. M., and Hsieh, C.-F.: Species Composition and Structure of the Lowland Subtropical Rainforest at Lanjenchi, Southern Taiwan, *TAIWANIA*, 52(3):, 253-269, 2007.
- 5 Chao, W.-C., Wu, S.-H., Fan, S.-W., Lin, H.-Y., Hsieh, C.-F., and Chao, K.-J.: Distribution Patterns of Tree Species in a Lowland Rainforest at Nanjen Lake, Southern Taiwan, *Taiwania*, 53, 124-133, 2008.
- Chen, Y.-F., and Huang, T.-C.: Vegetation analysis in Nanjenshan area, *Ann. Taiwan Mus.* , 29:, 189-258. , 1986.
- Chin, C.-C.: Study on the Spatial and Temporal Change of Litterfall in Nanjenshan Lowland Rain Forest, master thesis, 2008. [National Pingtung University of Science and Technology, Taiwan.](#)
- 10 Conklin-Brittain, N. L., Knott, C. D., and Wrangham, R. W.: Feeding Ecology in Apes and Other Primates. Ecological, Physical and Behavioral Aspects, , Energy intake by wild chimpanzees and orangutans: methodological considerations and a preliminary companson, 17, edited by: Hohmann, G., Robbins, M. M., and Boesch, C., Cambridge University Press, 445-465 pp., 2006.
- Degnan, P. H., and Ochman, H.: Illumina-based analysis of microbial community diversity, *ISME J*, 6, 183-194, 15 10.1038/ismej.2011.74, 2012.
- Fan, S.-W., Chao, W.-C., and Hsieh, C.-F.: Woody Floristic Composition, Size Class Distribution and Spatial Pattern of a Subtropical Lowland Rainforest at Nanjen Lake, Southernmost Taiwan, *Taiwania*, 50, 307-326, 2005.
- Feakins, S. J., Peters, T., Wu, M. S., Shenkin, A., Salinas, N., Girardin, C. A. J., Bentley, L. P., Blonder, B., Enquist, B. J., Martin, R. E., Asner, G. P., and Malhi, Y.: Production of leaf wax *n*-alkanes across a tropical forest elevation transect, 20 *Organic Geochemistry*, 100, 89-100, 10.1016/j.orggeochem.2016.07.004, 2016.
- Giebler, J., Wick, L. Y., Chatzinotas, A., and Harms, H.: Alkane-degrading bacteria at the soil-litter interface: comparing isolates with T-RFLP-based community profiles, *FEMS Microbiol Ecol*, 86, 45-58, 10.1111/1574-6941.12097, 2013.
- Gomez-Coca, R. B., Perez-Camino Mdel, C., and Moreda, W.: Saturated hydrocarbon content in olive fruits and crude olive pomace oils, *Food Addit Contam Part A Chem Anal Control Expo Risk Assess*, 33, 391-402, 25 10.1080/19440049.2015.1133934, 2016.
- Guo, J. H.: The Estimation of Total Soil Organic Carbon Storage under three Habitat Types in Nanjenshan Lowland Rain Forest, master thesis, 2010.
- Hoffmann, B., Kahmen, A., Cernusak, L. A., Arndt, S. K., and Sachse, D.: Abundance and distribution of leaf wax *n*-alkanes in leaves of Acacia and Eucalyptus trees along a strong humidity gradient in northern Australia, *Organic* 30 *Geochemistry*, 62, 62-67, 10.1016/j.orggeochem.2013.07.003, 2013.
- Hsieh, C.-F., Sun, I.-F., and Yang, C.-C.: Species Composition and Vegetation Pattern of a Lowland Rain Forest at the Nanjenshan L TER Site, Southern Taiwan, *Taiwania*, 45, 107-119, 2000.
- Jetter, R., and Schäffer, S. S.: Chemical Composition of the *Prunus laurocerasus* Leaf Surface. Dynamic Changes of the Epicuticular Wax Film during Leaf Development, *Plant Physiol*, 126, 2001.

- Jurelevicius, D., Alvarez, V. M., Peixoto, R., Rosado, A. S., and Seldin, L.: The Use of a Combination of *alkB* Primers to Better Characterize the Distribution of Alkane-Degrading Bacteria, *PLoS One*, 8, e66565, 10.1371/journal.pone.0066565, 2013.
- Kahmen, A., Dawson, T. E., Vieth, A., and Sachse, D.: Leaf wax *n*-alkane  $\delta D$  values are determined early in the ontogeny of *Populus trichocarpa* leaves when grown under controlled environmental conditions, *Plant Cell Environ*, 34, 1639-1651, 10.1111/j.1365-3040.2011.02360.x, 2011.
- Kirkels, F. M. S. A., Cammeraat, L. H., and Kuhn, N. J.: The fate of soil organic carbon upon erosion, transport and deposition in agricultural landscapes — A review of different concepts, *Geomorphology*, 226, 94-105, 10.1016/j.geomorph.2014.07.023, 2014.
- Kloos, K., Munch, J. C., and Schloter, M.: A new method for the detection of alkane-monoxygenase homologous genes (*alkB*) in soils based on PCR-hybridization, *J Microbiol Methods*, 66, 486-496, 10.1016/j.mimet.2006.01.014, 2006.
- Koch, K., Bhushan, B., and Barthlott, W.: Multifunctional surface structures of plants: An inspiration for biomimetics, *Progress in Materials Science*, 54, 137-178, 10.1016/j.pmatsci.2008.07.003, 2009.
- Kostka, J. E., Prakash, O., Overholt, W. A., Green, S. J., Freyer, G., Canion, A., Delgardio, J., Norton, N., Hazen, T. C., and Huettel, M.: Hydrocarbon-degrading bacteria and the bacterial community response in gulf of Mexico beach sands impacted by the deepwater horizon oil spill, *Appl Environ Microbiol*, 77, 7962-7974, 10.1128/AEM.05402-11, 2011.
- Nie, Y., Chi, C. Q., Fang, H., Liang, J. L., Lu, S. L., Lai, G. L., Tang, Y. Q., and Wu, X. L.: Diverse alkane hydroxylase genes in microorganisms and environments, *Sci Rep*, 4, 4968, 10.1038/srep04968, 2014.
- Quinton, J. N., Catt, J. A., Wood, G. A., and Steer, J.: Soil carbon losses by water erosion: Experimentation and modeling at field and national scales in the UK, *Agriculture, Ecosystems & Environment*, 112, 87-102, 10.1016/j.agee.2005.07.005, 2006.
- Rasheed, M. A., Kalpana, M. S., Veena Prasanna, M., Lakshmi, M., Madhavi, T., Devleena Mani, T., Patil, D. J., Dayal, A. M., and Raju, S. V.: Geo-microbial and light gaseous hydrocarbon anomalies in the near surface soils of Deccan Syneclise Basin, India: Implications to hydrocarbon resource potential, *Journal of Petroleum Science and Engineering*, 84-85, 33-41, 10.1016/j.petrol.2012.01.010, 2012.
- Rojo, F.: Degradation of alkanes by bacteria, *Environ Microbiol*, 11, 2477-2490, 10.1111/j.1462-2920.2009.01948.x, 2009.
- Rojo, F.: Enzymes for Aerobic Degradation of Alkanes, *Handbook of Hydrocarbon and Lipid Microbiology*, 781-797, 10.1007/978-3-540-77587-4\_59, 2010.
- Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R., McInerney, F. A., van der Meer, M. T. J., Polissar, P., Robins, R. J., Sachs, J. P., Schmidt, H.-L., Sessions, A. L., White, J. W. C., West, J. B., and Kahmen, A.: Molecular Paleohydrology: Interpreting the Hydrogen-Isotopic Composition of Lipid Biomarkers from Photosynthesizing Organisms, *Annual Review of Earth and Planetary Sciences*, 40, 221-249, 10.1146/annurev-earth-042711-105535, 2012.
- Schmidt, S. K.: A substrate-induced growth-response method for estimating the biomass of microbial functional groups in soil and aquatic systems, *FEMS Microbiology Ecology*, 101, 197-206, 1992.

- Schulz, S., Giebler, J., Chatzinotas, A., Wick, L. Y., Fetzer, I., Welzl, G., Harms, H., and Schloter, M.: Plant litter and soil type drive abundance, activity and community structure of *alkB* harbouring microbes in different soil compartments, *ISME J*, 6, 1763-1774, 10.1038/ismej.2012.17, 2012.
- Singh, S. N., Kumari, B., and Mishra, S.: Microbial Degradation of Alkanes, 439-469, 10.1007/978-3-642-23789-8\_17, 5 2012.
- Tsui, C.-C., Chen, Z.-S., and Hsieh, C.-F.: Relationships between soil properties and slope position in a lowland rain forest of southern Taiwan, *Geoderma*, 123, 131-142, 10.1016/j.geoderma.2004.01.031, 2004.
- Wentzel, A., Ellingsen, T. E., Kotlar, H. K., Zotchev, S. B., and Throne-Holst, M.: Bacterial metabolism of long-chain *n*-alkanes, *Appl Microbiol Biotechnol*, 76, 1209-1221, 10.1007/s00253-007-1119-1, 2007.
- 10 Xu, K., Tang, Y., Ren, C., Zhao, K., and Sun, Y.: Diversity and abundance of *n*-alkane-degrading bacteria in the near-surface soils of a Chinese onshore oil and gas field, *Biogeosciences*, 10, 2041-2048, 10.5194/bg-10-2041-2013, 2013.

Ranking of Leaf <i>n</i> -alkane concentration	Plant species	Relative annual Litter-leaf production (by dry weight)		
		Ravine	Windward	Leeward
1	<i>Ilex rotunda</i>	++++	++	+
2	<i>Ficus benjamina</i> L.	+++++		
3	<i>Ficus microcarpa</i> <del><i>benjamina</i></del> L. f.	+++++	+	
4	<i>Diospyros eriantha</i>	++		++
5	<i>Reevesia formosana</i>	+++		
6	<i>Celastrus kusanoi</i>	+++++	+	
7	<i>Erycibe henryi</i>	+++		+
8	<i>Cyclobalanopsis longinux</i>	+	+	+++++
9	<i>Archidendron lucidum</i>		+++++	+++++
10	<i>Celtis sinensis</i> Pers.	+++		
11	<i>Vitis kelungensis</i> Moriyama	+	++++	++
12	<i>Ilex cochinchinensis</i>		+++	+++
13	<i>Rhaphiolepos indica</i>		+++++	+
14	<i>Castanopsis fabri</i> Hance		+	+++++
15	<i>Symplocos caudata</i> Wall.		++++	+
16	<i>Cryptocarya hainanensis</i>	++++	+	+
17	<i>Podocarpus macrophyllus</i>		+++++	+++
18	<i>Sapium discolor</i>	++	+	++++
19	<i>Illicium arborescens</i>	+	+++	+
20	<i>Syzygium formosanum</i>		+++++	
21	<i>Syzygium kususense</i>	+	+++	+
22	<i>Machilus thunbergii</i>	+++++		
23	<i>Cyclobalanopsis pachyloma</i>	+	+++	+
24	<i>Lithocarpus formosanus</i>	+++++	+	+
25	<i>Bischofia javanica</i>	++++	+	+
26	<i>Machilus zuihoensis</i>	++		+++++
27	<i>Ficus aurantiaca</i>	++++	+	
28	<i>Helicia formosana</i>	+++	++++	++++
29	<i>Cyclobalanopsis championii</i>		+++++	++
30	<i>Aglaia formosana</i>		+++	+++++
31	<i>Turpinia ternata</i>	+	++++	++++
32	<i>Aglaia elliptifolia</i>	++	++	+
33	<i>Castanopsis indica</i>	+++	+	+++
34	<i>Psychotria rubra</i>	++++	++++	+++
35	<i>Castanopsis carlesii</i>			+++
36	<i>Schima superba</i>	+	+	++
37	<i>Antidesma hiranense</i>	+	++	+
38	<i>Lithocarpus amygdalifolius</i>	+	++	+
39	<i>Schizostachyum diffusum</i>	+	+	++++
40	<i>Beilschmiedia fordii</i>	+	+	++++
41	<i>Beilschmiedia erythrophloia</i>	+	+	++
42	<i>Alniphyllum pterospermum</i>	++	++	+

Table 1: The resultant ranking of *n*-alkane concentration of 42 litter-leaf samples with the nomenclature and the semiquantitative results of annual litter-leaf productivity in three habitats. The plus sign (+) was used to denote the annual productivity rankings of litter-leaf in each habitat. Five plus signs (+++++) denoted the litter-leaf productivity rankings from 1 to 5; four plus signs (++++) denoted the productivity rankings from 6 to 10; three plus signs (+++) denoted the rankings from 11 to 15; two plus signs (++) denoted the rankings from 16 to 20; one plus sign (+) denoted the rankings after 21. A blank space denoted no litter-leaf was yielded all the year around in the habitat.