## Differential response of carbon cycling to long-term nutrient input and altered hydrological conditions in a continental Canadian peatland

Sina Berger<sup>1,2,3</sup>, Leandra Praetzel<sup>1,2</sup>, Marie Goebel<sup>1,2</sup>, Christian Blodau<sup>1,2,†</sup>, Klaus-Holger Knorr<sup>1</sup>

<sup>1</sup> University of Muenster, Institute of Landscape Ecology, Ecohydrology and Biogeochemistry Group, Heisenbergstraße 2,
 48149 Muenster, Germany

<sup>2</sup> University of Guelph, School of Environmental Sciences, 50 Stone Road East, Guelph, Ontario, N1G 2W1, Canada

<sup>3</sup> Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research (IMK-IFU), Kreuzeckbahnstraße 19, 82467 Garmisch-Partenkirchen, Germany

10 Correspondence to: Sina Berger (gefleckterschierling@gmx.de), Klaus-Holger Knorr (kh.knorr@uni-muenster.de)

**Abstract.** Peatlands play an important role in global carbon cycling, but their responses to long-term anthropogenically changed hydrologic conditions and nutrient infiltration are not well known. While experimental manipulation studies, e.g. fertilization or water table manipulations, exist on the plot scale, only few studies have addressed such factors under *in-situ* conditions. Therefore, an ecological gradient from center to periphery of a continental Canadian peatland bordering a eutrophic

- 15 water reservoir, as reflected by increasing nutrient input, enhanced water level fluctuations, and increasing coverage of vascular plants, was used for a case study of carbon cycling along a sequence of four differently altered sites. We monitored carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) surface fluxes and dissolved inorganic carbon (DIC) and CH<sub>4</sub> concentrations in peat profiles from April 2014 through September 2015. Moreover, we studied bulk–peat and pore–water quality and we applied  $\delta^{13}$ C–CH<sub>4</sub> and  $\delta^{13}$ C–CO<sub>2</sub> stable isotope abundance analyses to examine dominant CH<sub>4</sub> production and emission pathways during the
- 20 growing season of 2015. We observed differential responses of carbon cycling at the four sites, presumably driven by abundances of plant functional types and vicinity to the reservoir. A shrub dominated site in close vicinity to the reservoir, was a comparably weak sink for CO<sub>2</sub> (in 1.5 years:  $-1093 \pm 794$ , in 1 year:  $+135 \pm 281$  g CO<sub>2</sub> m<sup>-2</sup> (=net release)) as compared to two graminoid–moss dominated sites and a moss dominated site (in 1.5 years: -1552 to -2260 g CO<sub>2</sub> m<sup>-2</sup>, in 1 year: -896 to -1282 g CO<sub>2</sub> m<sup>-2</sup>). Also, the shrub dominated site featured notably low DIC pore–water concentrations as well as comparably
- <sup>13</sup>C enriched CH<sub>4</sub> ( $\delta^{13}$ C–CH<sub>4</sub>: -57.81 ±7.03 ‰) and depleted CO<sub>2</sub> ( $\delta^{13}$ C–CO<sub>2</sub>: -15.85 ±3.61 ‰) in a more decomposed peat, suggesting a higher share of CH<sub>4</sub> oxidation and differences in predominant methanogenic pathways. In comparison to all other sites, the graminoid–moss dominated site in closer vicinity to the reservoir featured a ~30 % higher CH<sub>4</sub> emission (in 1.5 years: +61.4 ±32, in 1 year: +39.86 ±16.81 g CH<sub>4</sub> m<sup>-2</sup>). Low  $\delta^{13}$ C–CH<sub>4</sub> signatures (-62.30 ±5.54 ‰) indicated only low mitigation



of  $CH_4$  emission by methanotrophic activity here. Pathways of methanogenesis and methanotrophy appeared to be related to the vicinity to the water reservoir: the importance of acetoclastic  $CH_4$  production apparently increased toward the reservoir, whereas the importance of  $CH_4$  oxidation increased toward the peatland center. Plant mediated transport was the prevailing  $CH_4$  emission pathway at all sites even where graminoids were rare. Our study thus illustrates accelerated carbon cycling in a

5 strongly altered peatland with consequences for  $CO_2$  and  $CH_4$  budgets. However, our results suggest that long-term excess nutrient input does not necessarily lead to a loss of the peatland carbon sink function.

#### **1** Introduction

- Since the last deglaciation, northern peatlands have played an important role in global carbon (C) cycling by storing atmospheric carbon dioxide (CO<sub>2</sub>) as peat, but also emitting significant amounts of C as methane (CH<sub>4</sub>) (Succow and Joosten,
  2012). C sequestration and CO<sub>2</sub> and CH<sub>4</sub> release are driven by numerous processes and the accumulation of peat results from only a small imbalance of photosynthetic C uptake over respiratory losses. CO<sub>2</sub> can be released through autotrophic and heterotrophic respiration under aerobic and anaerobic conditions (Limpens et al., 2008). Heterotrophic respiration has been intensively studied and controls are manifold, as e.g. reviewed by Blodau (2002).
- Methanogenesis is strictly limited to anaerobic conditions (Conrad, 2005). Due to thermodynamic controls, CH<sub>4</sub> production is
   only competitive upon depletion of alternative, energetically more favorable electron acceptors for anaerobic respiration, such as nitrate, iron, sulfate or oxidized humics (Blodau, 2002; Klüpfel et al., 2014). CH<sub>4</sub> is predominantly produced via two pathways: hydrogenotrophic and acetoclastic methanogenesis. During hydrogenotrophic methanogenesis CO<sub>2</sub> is reduced to CH<sub>4</sub>, while during acetoclastic methanogenesis acetate is split into CH<sub>4</sub> and CO<sub>2</sub>. These pathways differ with respect to their discrimination against the heavier <sup>13</sup>C–isotopes due to the kinetic isotope effect (Hoefs, 1987). Differences in the isotopic
- 20 composition are thereby commonly presented as  $\delta^{13}$ C values, expressed as:  $\delta^{13}$ C = (R<sub>sample</sub>/R<sub>standard</sub> 1) 1000 [‰], where R is the ratio of heavy isotope to light isotope of the samples and the VPDB- standard. Acetoclastic methanogenesis results in  $\delta^{13}$ C-CH<sub>4</sub> values of -65 to -50 ‰, while hydrogenotrophic methanogenesis discriminates stronger against the heavier C isotope and results in  $\delta^{13}$ C-CH<sub>4</sub> values of -110 to -60 ‰ and considerably <sup>13</sup>C enriched CO<sub>2</sub> compared to the acetoclastic pathway (Whiticar et al., 1986). Specific patterns have been observed in terms of spatial and temporal occurrence of the major CH<sub>4</sub>
- 25 production pathways, with acetoclastic methanogenesis typically increasing in contribution towards the surface or within the rhizosphere (Holmes et al., 2015). On the other hand, an assignment of methanogenic pathways based on <sup>13</sup>C signatures of CH<sub>4</sub> can be biased by microbial oxidation of CH<sub>4</sub>. This can in particular be the case in transition to the unsaturated profile where oxygen can enter by diffusion, or in the rhizosphere where plants deliver oxygen through aerenchyma to their roots (Chasar et al., 2000). Upon oxidation of CH<sub>4</sub> into CO<sub>2</sub>, the residual CH<sub>4</sub> gets enriched in <sup>13</sup>C compared to the source CH<sub>4</sub> (Teh
- 30 et al., 2006), a process which yields similar  $\delta^{13}$ C–CH<sub>4</sub> signatures as observed upon CH<sub>4</sub> production via the acetoclastic pathway.

 $CH_4$  is released to the atmosphere by three different processes: i) diffusion through the acrotelm, which is a relatively slow process, ii) ebullition, i.e. fast evasion of  $CH_4$  bubbles, and iii) fast diffusion or pressurized throughflow convection through aerenchymatous tissues of vascular plants (Morris et al., 2011; Schütz et al., 1991; Whiting and Chanton, 1996; van den Berg et al., 2016; Hornibrook et al., 2009). Due to the slow diffusion of  $CH_4$  in peat, up to 100 % of diffusive  $CH_4$  is oxidized in the

- 5 acrotelm before it reaches the atmosphere, while the other processes effectively bypass oxidation and thus contribute a major fraction to observed fluxes (Whalen et al., 1990; Whalen, 2005). Therefore, a change in vascular plant cover or changes in the peat structure and degree of decomposition can be expected to affect CH<sub>4</sub> emissions.
- C cycling is strongly linked to nitrogen (N) cycling in peatlands. For these normally nutrient-limited ecosystems, eutrophication is a major threat, as demonstrated in several long-term fertilization experiments. A decade of fertilizer applications to bogs in Canada (Mer Bleue), in the UK (Whim Bog), in Sweden (Degerö Stormyr) and seven years of high nitrogen deposition to a bog in the Italian Alps caused a loss of mosses and an increase in vascular plant biomass (Bubier et al., 2007; Wang et al., 2016; Sheppard et al., 2013; Wiedermann et al., 2007; Bragazza et al., 2012). In the Mer Bleue bog, nutrient addition mostly increased abundance of shrubs, whereas at Whim bog, Degerö Stormyr and at an Italian mire both shrub or graminoid cover increased. A number of studies supported that an increase of vascular plant cover can reduce the
- 15 productivity of peat mosses and, in addition, can potentially promote the decomposition of organic matter by affecting the stoichiometry of soil enzymatic activity (Bragazza et al. 2013, Bragazza et al. 2015). This could lead to a decreasing ability of peatlands to sequester CO<sub>2</sub> from the atmosphere (Bubier et al., 2007), and to higher decomposition of peat (Rydin and Jeglum, 2013). Altered plant communities in peatlands were repeatedly shown to alter CO<sub>2</sub> and CH<sub>4</sub> fluxes: Maximum net ecosystem exchange (NEE) was found to be reduced after long-term fertilization and a concomitantly promoted vascular plant community
- 20 in the Mer Bleue bog (Bubier et al., 2007); Increased CH<sub>4</sub> emissions were observed at Degerö Stormyr from plots with an increased vascular plant coverage after a decade of excess nutrient supply (Eriksson et al., 2010). Indeed, selective removal of plant functional types (PFTs) and vegetation changes can have a strong impact on gas exchange (Larmola et al., 2013; Ward et al., 2013; Kuiper et al., 2014; Robroek et al., 2015). While such plot based manipulation experiments as reported above revealed clear patterns, there is still a gap of knowledge in terms of long-term consequences of excess nutrients supply to a
- 25 peatland and the resulting interactions and feedbacks between plants and peat, especially under *in-situ* conditions. There is only a poor understanding of the interplay of PFTs, substrate quality, and anoxic-oxic conditions, and of how exchange of  $CO_2$  and  $CH_4$  at the soil/atmosphere-interface would eventually be affected.

To address this research gap, we investigated C cycling of the once oligotrophic Wylde Lake peatland, which since 1954 is exposed to infiltration of nutrients and strongly pronounced water level fluctuations as induced by a nearby water reservoir. A particular recent finding at this site was that even after decades of excess nutrient supply (currently  $5.9 \pm 0.1$  to  $4.35 \pm 0.3$  g m<sup>-</sup>

- 30 particular recent finding at this site was that even after decades of excess nutrient supply (currently  $5.9 \pm 0.1$  to  $4.35 \pm 0.3$  g m<sup>-2</sup> y<sup>-1</sup> of N input and input of several other macro nutrients), the peatland still featured high peat accumulation rates of ~200 to ~300 g C m<sup>-2</sup> y<sup>-1</sup> (Berger et al., 2017). However, a strong gradient in vascular plant cover was apparent. As pointed out by Berger et al. (2017), lateral nutrient influx through repeated inundation events cannot be easily compared to sites subjected to deposition from the atmosphere. Nevertheless, an apparently intact peatland system, i.e. an intact mire with active C
  - 3

accumulation, despite such serious anthropogenic impacts would contradict several findings from above mentioned studies; According to these studies, already after one decade of N fertilization increased decomposition and peat degradation would be expected. Moreover, the particular scenario in our study, namely the impact of inundation on nearby ecosystems, is gaining increasing importance as there is a worldwide increase of impoundment area (Tranvik et al., 2009) and therefore serious effects

- 5 on peatland C cycling are likely (Ballantyne et al., 2014; Kim et al., 2015). The objective of this study was therefore to extend our findings from the existing study on nutrient impact, vegetation, and net C accumulation (Berger et al., 2017), comparing effects on C cycling and methanogenesis in more detail. To this end, we assessed current CO<sub>2</sub> and CH<sub>4</sub> exchange, peat quality and pore–water chemistry along a transect, ranging from a shrub dominated site (Site 4, 200 m distance to the reservoir; greatest nutrient input), over two graminoid–moss dominated sites
- 10 (Sites 3 and 2, 400 and 550 m distance to the reservoir; intermediate nutrient input) to a moss dominated site (Site 1, 800 m distance to the reservoir; smallest nutrient input) in the Wylde Lake peatland in Ontario, Canada. Moreover, to address changes in methanogenic pathways and to study predominant pathways of  $CH_4$  emission, we assessed seasonal variation in  $\delta^{13}C$  of  $CH_4$  and  $CO_2$  in peat profiles and in  $CH_4$  surface fluxes.

We hypothesized that 1) hydrologically altered and nutrient enriched peripheral sites feature accelerated C cycling and more decomposed peat, due to input of labile litter from productive vascular vegetation, 2) increased abundance of vascular plants

15 decomposed peat, due to input of labile litter from productive vascular vegetation, 2) increased abundance of vascular plants can increase  $CO_2$  uptake but also change patterns of  $CH_4$  production and emission, in particular if graminoids with aerechymatous roots dominate, and that 3) long-term nutrient enrichment in combination with hydrologically altered conditions may therefore cause differential responses of C cycling and does not necessarily cause a loss of the C-sink function of peatland ecosystems.

#### 20 2 Methods

25

#### 2.1 Description of the study area and study sites

Wylde Lake peatland has been described in detail by Berger et al. (2017). In brief, it is located in southeastern Ontario, 80 km northwest of Toronto (43.920361° N, 80.407167° W) (Fig. 1), as part of the Luther Lake Wildlife Management Area. Climate is cool temperate, average July temperature is 19.1 °C, average January temperature is -8.0 °C and the mean annual temperature is about 6.7 °C. Annual precipitation amounts to 946 mm, with the major portion falling in summer (1981 to 2010, Fergus Shand Dam, National Climate Data and Information Archive, 2014). Peat formation started about 9000 years before

present on calcareous limnic sediments and the total peat depth today is about 5 meters. For flood control and water management, the "Luther Lake" reservoir, neighboring Wylde Lake peatland, had been created in 1954. Through flooding of the reservoir, Wylde Lake peatland has been exposed to altered hydrological conditions in a way

30 that the water reservoir enhanced water level fluctuations in a large part of the site: in summer or under dry conditions, water is released from the reservoir, draining water out of the peatland; under wet conditions, increased water table levels of the reservoir push water into the peatland. Sites in closer vicinity to the reservoir are presumably more affected than sites further away (Berger et al., 2017).

Four intensively investigated sites (Fig. 1) were arranged along a transect stretching from nearby the shoreline of the reservoir about ~1 km south into the central, treed bog area. Each site featured an individual mosaic of hummocks, hollows, and lawns;

- 5 however, for comparability, all measurements of this study were taken in and all samples were collected from hollows. Site 4 was located about 200 m away from the reservoir in an area overgrown by *Myrica gale* where *Sphagnum* mosses were in retreat. Site 3 and site 2 were in the open poor fen - bog transition area with site 2 being further away from the reservoir (550 m) than site 3 (400 m). These intermediate sites 2 and 3 were dominated by *Sphagnum* mosses and graminoids with only few shrubs. These sites featured a variety of arenchymatous graminoid species, such as *Eriophorum* spp. at the sites 3 and 2,
- 10 and *Dulichium arundinaceum* at site 3. Site 1 (~800 m away from the reservoir) accommodated equal shares of few graminoids and shrubs above dominant *Sphagnum* mosses. The four sites also differed with respect to their most abundant *Sphagnum* species, reflecting increasingly minerotrophic conditions towards the lake. While *S. capillifolium*, an ombrotrophic to slightly minerotrophic hummock species (Laine et al., 2011), was abundant at the sites 1, 2 and 3, its abundance decreased towards site 4. Moreover, site 1 featured the abundant *S. magellanicum* (another ombrotrophic to weakly minerotrophic hummock
- 15 species (Laine et al., 2011)), site 2 featured the abundant *S. angustifolium* (tolerating ombrotrophic to minerotrophic conditions (Laine et al., 2011)) and site 3 featured the abundant *S. girgensohnii*, a minerotrophic hollow species (Laine et al., 2011). The two most abundant *Sphagnum* species at site 4 were *S. fuscum* (mostly on hummocks but also in hollows, an ombrotrophic species (Laine et al., 2011), with a great ability to recover from desiccation (Nijp et al., 2014) and again the minerotrophic hollow species *S. girgensohnii*. See Table 1 for a detailed overview of the vegetation at the sites and see Fig. S1 for
- 20 photographs of the sites.

As presented in Berger et al. (2017), the study area is subject to apparent nutrient infiltration from the "Luther Lake" water reservoir as indicated by increasing concentrations of N, phosphorus (P), sulfur, potassium (K), calcium (Ca), magnesium (Mg), iron, copper, and zinc as well as other metals in peat mostly toward the peatland periphery. N input rates of  $5.9 \pm 0.1$  g N m<sup>-2</sup> y<sup>-1</sup> were reported for site 4 and  $4.35 \pm 0.3$  g N m<sup>-2</sup> y<sup>-1</sup> for site 1; moreover, C/P and N/P ratios of surface peat suggested

- 25 P limitation typical of fens, C/Ca and C/Mg ratios indicated Ca and Mg limitation, while C/K ratios indicated higher K availability as compared to typical values for bogs presented in Wang et al. (2015). The peatland periphery appeared to act as a buffer for nutrients in a way that site 4 received the highest loads of nutrients but also areas further away were to some extent affected. Nevertheless, surface peat accumulation rates of ~200 to ~300 g C m<sup>-2</sup> y<sup>-1</sup> at all sites revealed great recent C sequestration rates.
- 30 The impact of anthropogenic activities, in particular the formation of the reservoir, is evident from peat quality found at the sites: quite similar peat quality was found at all sites for those depths that had accumulated before dam construction at Wylde Lake peatland. A clear difference before and after dam construction was evident, reflected by the enrichment in nutrients in the upper depths and the concomitantly altered vegetation.

#### 2.2 Determination of organic matter quality of peat and pore-water

Peat samples were taken in July, 2014, in depths of 5, 10 and 20 cm below the living *Sphagnum* layer by manual cutting. Peat from 75 cm depth was taken with a Russian peat corer. Peat was filled in jars avoiding any headspace and closed air–tight to maintain anoxic conditions as far as possible during transport to the laboratory.

- 5 To collect *in-situ* pore-water, suction samplers (Macro Rhizons, Eijkelkamp, Giesbeck, The Netherlands; pore size ~ 0.2 μm) were inserted into the peat at 5, 10 and 20 cm depth. Sampling was done by applying vacuum and collecting water with syringes; syringes were covered with aluminum foil and peat to avoid exposure to light. Pore-water from 75 cm depth was pumped from 75 cm deep piezometers that were emptied one day prior to sampling to ensure sampling of fresh pore-water. Samples from piezometers were filtered using Macro Rhizons in the laboratory to ensure similar treatment of pore-water of all depths. All samples were taken and analyzed as three replicates.
- Prior to Fourier-transform infrared spectroscopic (FTIR) analysis, oven–dried (70 °C) bulk peat samples were ground with a ball mill. Pore–water samples were oven–dried (70 °C) to complete dryness; afterwards 2 mg of the remaining organic matter were ground in a mortar with 200 mg of potassium bromide (KBr) and pressed to pellets for analysis. We recorded spectra on an FT–IR Spectrometer (Varian 660, Palo Alto, USA) over a scan range of 4000 to 650 cm<sup>-1</sup> with a resolution of 2 cm<sup>-1</sup> and
- 15 32 scans per sample. A KBr background was subtracted from the spectra and spectra were baseline corrected. We identified spectral peaks (average location +/- 30 cm<sup>-1</sup>) and related them to functional moieties as described in Niemeyer et al. (1992). As absorbance values do not give quantitative information on absolute abundance of functional groups, we related peaks of around 1620 cm<sup>-1</sup> to 1610 cm<sup>-1</sup> (aromatic C=C compounds/aromatic moieties) to polysaccharide peaks at 1170 cm<sup>-1</sup> to 950 cm<sup>-1</sup> wavenumbers (Niemeyer et al., 1992). A relative increase in ratios thus indicates a relative decrease in the labile
- 20 polysaccharide moieties and thus an increase in the degree of decomposition in regard of a residual enrichment of refractory aromatics (Broder et al., 2012).

Pore–water samples were analyzed by absorption spectroscopy in the ultra violet and visible range (UV–VIS-spectroscopy; Varian UV 1006 M005 spectrometer, Palo Alto, USA). We recorded UV–VIS spectra over a range of 200 to 800 nm with a resolution of 0.5 nm using a 1 cm quartz cuvette. Prior to measurement, a blank spectrum of ultrapure water was recorded and

- 25 subtracted from each sample. We additionally recorded fluorescence properties of dissolved organic matter (DOM) on a fluorescence spectrometer (Varian Cary Eclipse, Palo Alto, USA) at a scan rate of 600 nm/min. Excitation wavelengths (ex) were 240 to 450 nm in 5 nm steps, emission wavelengths (em) 300 to 600 nm in 2 nm steps to obtain excitation–emission–matrices (EEMs). Repeated blanks were run to ensure cleanliness of the cuvette. Raman spectra of a blank were recorded each day to check analytical drift and to normalize fluorescence to Raman units (Murphy et al., 2010).
- 30 To evaluate DOM quality, we calculated commonly used indices, such as specific ultraviolet absorbance SUVA<sub>254</sub> (as a proxy for aromaticity, Weishaar et al., 2003) and the E2:E3 ratio (the ratio of UV absorbance at 250 nm divided by absorbance at 365 nm) providing information about molecular weight of organic matter (Peuravouri and Pihlaja, 1997) from UV–VIS data. From fluorescence data, we calculated a humification index HIX (Ohno, 2002) (see Table S3 for equations used).



#### 2.3 Measurements of environmental variables

Air temperature and photosynthetically active radiation (PAR) were recorded about 1 km south of site 1 in an open area by a HOBO U30 weather station (U30-NRC-SYS-B, Onset, Bourne, MA, USA) at a temporal resolution of 5 min. Water table depth below surface (wtd), water temperature ( $T_{water}$ ) and air pressure were measured in 30–min intervals using one pressure

5 transducer (Solinst Levelogger Edge) in a monitoring well at each site, corrected for barometric pressure (Barologger Gold at site 2; Solinst Ltd., Georgetown, Canada). On each day of closed chamber measurements, an extra PAR sensor (Smart Sensor, Onset; Part # S-LIA-M003) and an extra temperature sensor (Temperature Smart Sensor, Onset; Part # S-TMB-M0XX) recorded PAR and air temperature at a temporal resolution of 10 secs at the site were chamber measurements were being taken.

#### 2.4 Determination of CO<sub>2</sub> and CH<sub>4</sub> fluxes

- 10 In the hollows of sites 1–4, a set of six collars for chamber measurements were established in April 2012. The collars installed 0.1-0.15 m into the soil were cylindrical, had a diameter of 0.4 m and a total height of 0.2 m. Through object based image analysis (OBIA) based on aerial imagery obtained from UAV flights, the spatial coverages of PFTs at each site were obtained (data summarized in Table 1). Accordingly, the locations for chamber measurement collars of our study sites were defined to proportionally reflect the actual distribution of PFTs.
- 15 Closed chamber measurements were performed following Burger et al. (2016). Measurements were taken every 10 to 30 days at each site from April 20th, 2014 through September 22nd, 2015. In total, 19 to 23 daily courses per site could be accomplished. Cylindrical Plexiglas chambers were used for the flux measurements: a transparent chamber to measure net ecosystem exchange (NEE), a chamber covered with reflective insolation foil for ecosystem respiration (R<sub>eco</sub>). Chamber closure time was 180 secs.
- Air was circulated between the chamber and a trace gas analyzer (Ultraportable Greenhouse Gas Analyzer 915-001, Los Gatos Research Inc., Mountain View, USA) through 2 mm inner diameter polyethylene tubing, recording concentrations of CO<sub>2</sub> and CH<sub>4</sub> at a temporal resolution of 1 sec. According to the manufacturer, the reproducibility of CH<sub>4</sub> and CO<sub>2</sub> is < 2 ppb and < 300 ppb, respectively. The analyzer was factory–calibrated before the campaign. Stability of the calibration was checked repeatedly during summer of 2014. In January and July 2015, the analyzer was again re–calibrated. If CH<sub>4</sub> concentrations
- 25 increased sharply within the first 60 secs of the measurement due to  $CH_4$  bubble release caused by the positioning of the chamber, the measurement was discarded and repeated.

During each measurement day, each collar was monitored several times with the transparent and dark chamber at different times (typically between 5 am and 8 pm) and different PAR levels (typically 5 to 2000  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>) throughout the day. Unfortunately, due to the remoteness of our study site, measurements at night were not possible.

30

Gas fluxes were determined by Eq. 1:

 $F_{chamber} = \frac{\Delta c}{\Delta t A} \cdot \frac{P V}{R T},$ 

based on the changes of concentration over time inside the chamber ( $\Delta c$ ), applying the ideal gas law with the ideal gas constant R, and correcting for atmospheric pressure P and temperature inside the chamber T. The chamber volume V and basal area A

- 5 were calculated from the chamber's physical dimensions, taking into account each collar's vegetation volume, determined in May, July and October 2014 as well as in April and August 2015 and extrapolated for the other campaigns. The concentration change over time was derived from the slope of a linear regression of concentration vs. time. The first 40 secs after chamber deployment were discarded to account for the analyzer's response time. If the slope was not significantly different from 0 (tested with an F–test,  $\alpha = 0.05$ ), the flux was set to zero.
- 10 An empirical description of the measured NEE fluxes of each site was accomplished applying a hyperbolic light response model (Owen et al., 2007). The non-linear least squares fit of the data to the model was done according to Eq. 2:

$$NEE = \frac{\alpha\beta Q}{\alpha\beta} + y$$

20

15 where *NEE* is in mol m<sup>-2</sup> s<sup>-1</sup>,  $\alpha$  is the initial slope of the light response curve (in mol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup> per mol photon m<sup>-2</sup> s<sup>-1</sup>),  $\beta$  is the maximum NEE in mol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, Q is the photosynthetic active radiation in mol photon m<sup>-2</sup> s<sup>-1</sup>, and  $\gamma$  is an estimate of the average R<sub>eco</sub>. Integration of NEE over the course of one day gave net daily ecosystem production (NEP). Gross primary productivity (GPP) was retrieved by subtracting R<sub>eco</sub> from NEP.

Average  $CH_4$  fluxes for measurement days of each site were obtained and lastly, cumulative emissions of  $CO_2$  and  $CH_4$  were calculated likewise according to Tilsner et al. (2003).

To determine isotopic signatures of  $CH_4$  fluxes, we carried out additional chamber flux measurements once a month from May to September 2015 using a shrouded chamber and the setup described above. The chamber was closed until  $CH_4$  concentrations reached > 10 ppm for analysis of isotopic composition, but not more than 30 min. Samples for isotopic analysis were extracted from the chamber with 60 ml syringes and filled into 40 ml crimp vials that had before been flushed with nitrogen (N<sub>2</sub>) and

sealed with rubber stoppers. To correct isotopic values of CH<sub>4</sub> for background isotopic signature in the chamber, we collected six air samples at each site on every sampling day. Isotope analysis was carried out as outlined for dissolved gases (see below).

#### 2.5 Sampling of dissolved gases in the peat

Concentrations of CH<sub>4</sub> and dissolved inorganic carbon (DIC/ΣCO<sub>2</sub>) along peat profiles were analyzed in 5, 15, 25, 35, 45 and 55 cm depth with three replicates at each site using diffusive equilibration samplers made of permeable silicone tubes
(Kammann et al., 2001) equipped with valves. Samples were taken with 10 ml syringes every two to three weeks from June 2014 through September 2015. Samples were stored overnight at 5 °C and analyzed the next day.

To determine temporal dynamics of isotopic signatures of  $CH_4$  and  $CO_2$  in the peat, we installed a separate set of silicone tubes in 5, 15, 25 and 35 cm depth with three replicates each per site. Silicone tubes for isotope sampling had an inner diameter of 1 or 0.5 cm, corresponding to a volume of 20 or 5 ml. The larger samplers were installed in 5 cm depth and the smaller samplers below, as close to the surface larger volumes of samples were necessary in order to obtain sufficiently high concentrations (>

- 5 2.5 ppm) for isotope analysis. Equilibrium of gases such as N<sub>2</sub>O, CH<sub>4</sub>, and CO<sub>2</sub> at the silicone membrane has been shown to adjust within hours to days and isotopic fractionation can be expected to be negligible (Nielsen et al., 1997; Panikov et al., 2007; Pack et al., 2015). All samplers were installed one month prior to the first sampling. Samples were taken monthly from May 2015 to September 2015 using 10 and 60 ml syringes and filled into 10 respectively 40 ml crimp vials that had been flushed with N<sub>2</sub> and sealed with rubber stoppers. After sampling, silicone samplers were refilled with N<sub>2</sub> to avoid intrusion of
- 10 oxygen.

To obtain high resolution depth profiles of concentration and isotopic signatures of  $CH_4$  and DIC, pore–water peepers of 60 cm length and a 1 cm resolution (Hesslein, 1976) were inserted on three occasions in June, July and September 2015 and allowed to equilibrate for four weeks. As results of pore–water peepers generally confirmed the data of the silicone samplers, results are not presented here but described in the supporting information (see Fig. S6).

#### 15 2.6 Analyses of CO<sub>2</sub> and CH<sub>4</sub> concentrations and $\delta^{13}$ C-CO<sub>2</sub> and $\delta^{13}$ C-CH<sub>4</sub>-values

Gaseous CO<sub>2</sub> and CH<sub>4</sub> concentrations were analyzed with a gas chromatograph (SRI 8610 C, SRI Instruments, Torrance, US) equipped with a Flame Ionization Detector (FID) and a Methanizer. Samples from pore–water peepers were analyzed by measuring the headspace concentration in the vials.

- Ratios of  $\delta^{13}$ C of CO<sub>2</sub> and CH<sub>4</sub> were determined by Cavity Ringdown Spectroscopy (CRDS; Picarro G2201-*i*, Picarro Inc., 20 Santa Clara, US), simultaneously determining <sup>13</sup>C isotopic composition of CO<sub>2</sub> and CH<sub>4</sub> with a precision of <0.16 ‰ for  $\delta^{13}$ C– CO<sub>2</sub> and <1.15 ‰ for  $\delta^{13}$ C–CH<sub>4</sub>. The analyzer was calibrated before each measurement with two working standards of CO<sub>2</sub> (1000 ppm, -31.07 ‰) and CH<sub>4</sub> (1000 ppm, -42.48 ‰). Standard deviation for  $\delta^{13}$ C–CO<sub>2</sub> was below 2 ‰ and below 4 ‰ for  $\delta^{13}$ C–CH<sub>4</sub>. For CO<sub>2</sub>, additional in–house standards with a  $\delta^{13}$ C value of -26.61 ‰, -0.19 ‰ and -15,16 ‰ were used, validated by IRMS certified reference materials. Isotopic signatures were expressed in the  $\delta$ –notation in ‰ versus VPDB-
- 25 Standard according to Eq. 3:

 $\delta^{13}C = (R_{sample}/R_{standard} - 1) \cdot 1000 \, [\%]$ 

where  $R_{Sample}$  is the <sup>13</sup>C/<sup>12</sup>C ratio of the sample and  $R_{Standard}$  is the <sup>13</sup>C/<sup>12</sup>C ratio of the standard.

30 As the accuracy of  $\delta^{13}$ C–CO<sub>2</sub> values was affected by high CH<sub>4</sub> concentrations present in the samples, we established a correction to revise  $\delta^{13}$ C–CO<sub>2</sub> values. This was necessary for molar concentration ratios of CO<sub>2</sub>:CH<sub>4</sub> between 0.3 and 1.5. Samples with CO<sub>2</sub>:CH<sub>4</sub> ratios < 0.3 could not be corrected and were discarded; samples with ratios > 1.5 did not need correction. Additionally,  $\delta^{13}$ C–CO<sub>2</sub> values had to be corrected for a storage effect. As samples were stored for several weeks,

 $CO_2$  was lost from the vials and isotopic signatures increased by 0.056 ‰ per day. There was no such effect detectable for  $CH_4$ .

Dissolved concentration of  $CO_2$  and  $CH_4$  were recalculated from partial pressures inside the silicon samplers applying Henry's Law according to Eq. 4:

#### 5

10

 $c = K_H * p$ 

where *c* is the concentration in  $\mu$ mol/L, *p* is the pressure in atm and *K*<sub>H</sub> is the *in–situ* temperature corrected Henry–constant in mol L<sup>-1</sup> atm<sup>-1</sup> (Sander, 1999). Speciation of aqueous CO<sub>2</sub> was considered using equilibrium constants from Stumm and Morgan (1996) to calculate total DIC.

DIC and  $CH_4$  concentrations in samples from pore–water peepers were recalculated from gas concentrations in the headspace, applying the ideal gas law and temperature corrected Henry–constants.

To gain information about the dominant CH<sub>4</sub> production pathway, the isotope fractionation factor  $\alpha_C$  (for 35 cm depth) was calculated according to Eq. 5 after Whiticar et al. (1986):

15

 $\alpha_{\rm C} = (\delta^{13}{\rm C-CO_2} + 1000) / (\delta^{13}{\rm C-CH_4} + 1000).$ 

#### 2.7 Statistical analysis

Statistics software R i386 version 3.1.0 was used to verify differences in organic matter quality between depths and sites. Data was tested for normal distribution (Shapiro-Wilk-Test,  $\alpha = 0.05$ ) and homogeneity of variance (Levene-Test,  $\alpha = 0.05$ ). In case

- 20 both requirements were met, we carried out a one-way ANOVA (Analysis of Variance) ( $\alpha = 0.05$ ) with a post-hoc Tukey's Honest Significant Difference (HSD) test ( $\alpha = 0.05$ ) to identify which depths or which sites differed significantly from each other. If either normal distribution or homogeneity of variance were not met, a Kruskal-Wallis test ( $\alpha = 0.05$ ) with a multiple comparison test after Kruskal-Wallis ( $\alpha = 0.05$ ) as post-hoc test was applied.
- Using *RStudio* Version 0.99.902 as well as R i386 3.2.3 we examined differences in  $\delta^{13}$ C values of CO<sub>2</sub> and CH<sub>4</sub>, CO<sub>2</sub> and CH<sub>4</sub> concentrations and cumulative emissions between the sites. Means were compared with t-Tests (if data was normally distributed) respectively Kruskal-Wallis and post hoc Wilcoxon-Mann-Whitney-Test (if data was not normally distributed), with confidence levels of  $\alpha = 0.05$  for all statistical tests. Normality was tested with Shapiro-Wilk-Test ( $\alpha = 0.05$ ) and homogeneity of variance was confirmed with a Levene-Test ( $\alpha = 0.05$ ). Correlations between environmental variables and fluxes, concentrations and isotopic signatures were determined with Pearson's product-moment correlation for normally
- 30 distributed data or with Spearman's rank correlation if data was not normally distributed. With ANOVA ( $\alpha = 0.05$ ), the effect of categorical variables on CH<sub>4</sub> fluxes and  $\delta^{13}$ C values was computed.

#### **3 Results**

#### 3.1 Organic matter quality of peat and pore-water

The highest degree of bulk peat decomposition, as indicated by the highest 1618.5/1033.5 absorption ratios, was found at site 4 between 5 and 20 cm depth (p < 0.05 in 10 and 20 cm depth, Fig 2 (a)). The 1618.5/1033.5 ratios of the sites 1–3 were not

- 5 significantly different. Pore–water samples' 1618.5/1033.5 ratios of site 3 were smallest between 5 and 20 cm depth as compared to all other sites (p < 0.05), indicating the lowest degree of decomposition of DOM here (Fig 2 (b)). Aromaticity as determined with *SUVA*<sub>254</sub> (Fig 2 (c)) did not significantly differ between sites in pore–water samples (exception: site 1 and site 3 in 20 cm depth (p = 0.033), where site 1 SUVA<sub>254</sub> was significantly higher than site 3 SUVA<sub>254</sub>). The degree of humification, as expressed by HIX (Fig 2 (d)), was significantly lowest in site 3 pore–water (5 cm site 3 and 4: p = 0.026; 10 cm site 1 and
- 10 3: p = 0.014; 20 cm site 3 and 4: p = 0.020). The slope ratio E2:E3 (Fig 2 (e)), indicative of molecular size and aromaticity,
  - did not significantly differ between sites.

#### 3.2 Development of wtd and $T_{water}$ during the study period

During our study period, hollow wtd showed strong seasonal fluctuations. Maximum wtd (i.e. highest water table levels) throughout the study period were reached during snowmelt in spring 2014 (site 1: 6.94 cm, site 2: 4.99 cm, site 3: 16.26 cm,

- 15 site 4: 23.18 cm above hollow surface). Minimum wtd (i.e. lowest water table levels) were reached during the summer of 2015 (site 1: 32.5 cm, site 2: 31.75 cm, site 3: 13.34, site 4: 19.11 cm below hollow surface). All sites showed similar courses of wtd, however, site 3 and site 4 water levels were generally higher than site 1 and 2 water levels (p < 0.05). The amplitude between maximum and minimum wtd at all sites was overall similar (site 1: ~39.5 cm, site 2: ~36.7 cm, site 3: ~30 cm (logger failure when water levels were lowest), site 4: ~42.3 cm). T<sub>water</sub> varied between ~2 °C in winter and ~16 °C in summer. Detailed
- 20 courses of wtd and  $T_{water}$  are presented in Fig. 3 (a) and (b).

# 3.3 Fluxes of $CO_2$ and $CH_4$ at the soil/atmosphere interface and concentrations of $CH_4$ and DIC along peat profiles during the study period

Fluxes of CH<sub>4</sub> and CO<sub>2</sub> (Fig. 3, panels (c) – (f)) showed strong annual variability. Greatest CH<sub>4</sub> emission (Fig. 3 (c)) occurred during the growing season, minor fluxes were detected during the dormant season. In general, site 3 emitted more and site 4 less CH<sub>4</sub> than the other sites with an exception on August 16th, 2015 when a mean flux of  $0.76 \pm 0.58$  g CH<sub>4</sub> m<sup>-2</sup> d<sup>-1</sup> was

- 25 less CH<sub>4</sub> than the other sites with an exception on August 16th, 2015 when a mean flux of  $0.76 \pm 0.58$  g CH<sub>4</sub> m<sup>-2</sup> d<sup>-1</sup> was detected at site 4, exceeding the fluxes measured at all other sites. During the entire study period, site 3 released significantly (p < 0.001) more CH<sub>4</sub> (61.4 ± 32 g CH<sub>4</sub> m<sup>-2</sup>) than the sites 1 (41.8 ± 25.4 g CH<sub>4</sub> m<sup>-2</sup>), 2 (44.6 ± 13.7 g CH<sub>4</sub> m<sup>-2</sup>), and 4 (46.1 ± 35.2 g CH<sub>4</sub> m<sup>-2</sup>); see also Fig. S5. Annual cumulative CH<sub>4</sub> emissions from May 2014 to May 2015 were 22.18 ± 8.96 at site 1, 30.66 ± 7.63 at site 2, 39.86 ± 16.81 at site 3, and 12.53 ± 11.38 g CH<sub>4</sub> m<sup>-2</sup> at site 4. Thus, site 3 emitted significantly (p <
- 0.05) more CH<sub>4</sub> than site 4, but CH<sub>4</sub> emission at sites 3 and 4 were not different from emissions at the sites 1 and 2. Site 3 had the highest negative NEP, indicating greatest CO<sub>2</sub> net uptake, whereas site 4 had the lowest negative, sometimes even positive

NEP, indicating little net uptake if not a net emission of  $CO_2$  (Fig. 3 (d)). Regarding  $R_{eco}$  (Fig. 3 (e)), patterns were similar at all sites. In accordance with the NEP results, GPP (Fig. 3 (f)) was lowest at site 3, indicating highest photosynthetic uptake here, whereas site 4 had the highest GPP, indicating smallest uptake. Between May, 2014 and September, 2015 site 4 accumulated significantly less CO<sub>2</sub> ( $-1093 \pm 794$  g CO<sub>2</sub> m<sup>-2</sup>, p < 0.05) than the other three sites (-1552 to -2260 g CO<sub>2</sub> m<sup>-2</sup>),

5 while there were no significant differences in terms of CO<sub>2</sub> uptake for the sites 1, 2 and 3. Between May 2014 and May 2015 NEP of the sites 1, 2 and 3 was strongly negative (-896 to -1282 g CO<sub>2</sub> m<sup>-2</sup>) compared to site 4 NEP ( $+135 \pm 281$  g CO<sub>2</sub> m<sup>-2</sup>). p < 0.05).

Site 4 CH<sub>4</sub>, NEP and GPP fluxes differed notably between the growing seasons of 2014 and 2015. This was particularly caused by two plots, which in 2015 dramatically increased productivity and CH<sub>4</sub> emissions as compared to the previous year (data not shown).

- Concentrations of  $CH_4$  along depth profiles (Fig. 4, top panels) of all sites varied strongly throughout the year: they generally increased during the growing season, reached maximum values in the winter season 2014/2015 and comparably decreased during snowmelt in spring. A similar pattern was observed for DIC concentrations along depth profiles (Fig. 4, lower panels). Maximum DIC concentrations were observed below 20 cm depth in autumn 2014 and winter 2014/2015. Minimum
- 15 concentrations were observed during snowmelt in March and April 2015. Site 4 DIC concentrations at all depths were overall lower and significantly decreased (p < 0.05) in comparison to all other sites from February 23rd through April 4th, 2015. Moreover, site 4 DIC concentrations were significantly (p < 0.05) lower than site 3 DIC concentrations on August 6th, 2014 and between April 19th through July 18th, 2015. Concentrations in the uppermost depths of both CO<sub>2</sub> and CH<sub>4</sub> were strongly affected by fluctuations of wtd, with strong decreases upon water table decline and vice versa (see table S4 for statistical
- 20 results).

25

10

### 3.4 Temporal and spatial variability of $\delta^{13}$ C-CO<sub>2</sub> and $\delta^{13}$ C-CH<sub>4</sub> -values in peat pore-gas profiles during the growing season in 2015

Values of  $\delta^{13}$ C of CH<sub>4</sub> in the peat ranged from -78.74 to -26.77 ‰,  $\delta^{13}$ C of CO<sub>2</sub> ranged from -25.81 to +4.03 ‰ (see Fig. 5). Highest  $\delta^{13}$ C–CH<sub>4</sub> and CO<sub>2</sub> values were measured at site 1 in 5 respectively 35 cm depth in September. Lowest  $\delta^{13}$ C–CH<sub>4</sub> and CO<sub>2</sub> values were detected at site 1 in 15 cm depth in June and at site 2 in 15 cm depth in August respectively.

- Overall,  $\delta^{13}$ C–CH<sub>4</sub> values showed an increasing trend with time from June to August in all depths. Average signatures in 5 to 35 cm depth differed significantly between sampling dates at all sites except between August and September (p < 0.05). Concomitant to a decline in water table levels in August and September,  $\delta^{13}$ C–CH<sub>4</sub> signatures shifted to less negative values in the upper 5 cm at sites 1–3; this shift was most distinctive at site 1 and least distinctive at site 3. At site 4, such shift occurred 30
- down to 15 cm depth.

For  $\delta^{13}$ C–CO<sub>2</sub> signatures, significant differences between some sampling dates were found at sites 1, 2 and 4 for average values in 5 to 35 cm depth. At sites 1 and 2, signatures in August and September were higher than in June and July, paralleling the trend in  $\delta^{13}$ C–CH<sub>4</sub>. At sites 3 and 4, such significant shifts could not be observed.

At the sites 1 and 2, δ<sup>13</sup>C-CH<sub>4</sub> signatures apparently increased with depth in June and July, no trend was observable at sites 3
and 4. In August and September, δ<sup>13</sup>C-CH<sub>4</sub> signatures seemed to decrease with depth except for site 4. Values of δ<sup>13</sup>C of CO<sub>2</sub> increased with depth except at site 1 in July and at site 2 in July and August.

Mean signatures of  $\delta^{13}$ C–CH<sub>4</sub> at site 4 (-57.81 ±7.03 ‰) differed significantly from those at the other sites (site 1: -61.48 ±10.71 ‰, site 2: -60.28 ±5.57 ‰, site 3: -62.30 ±5.54 ‰) for the whole sampling period (p < 0.01, p < 0.05, p < 0.001).

Values of  $\delta^{13}$ C of CO<sub>2</sub> at site 3 were significantly higher than at the other sites in July (p < 0.05, p < 0.01, p < 0.01). Overall, highest mean values were found at site 1 (12.05 + 8.23 %) whereas site 4 revealed lowest  $\delta^{13}$ C CO signatures (15.85 +

10 highest mean values were found at site 1 (-12.05  $\pm$  8.23 ‰) whereas site 4 revealed lowest  $\delta^{13}$ C-CO<sub>2</sub> signatures (-15.85  $\pm$  3.61 ‰).

Isotopic composition of  $CH_4$  and  $CO_2$  as determined from pore–water peepers confirmed results obtained from the silicone gas samplers. Data is presented in the Fig. S5 in the supplemental information.

- Fractionation factors  $\alpha_{C}$  to characterize methanogenic pathways (according to Whiticar et al. (1986)) were calculated for water 15 saturated, presumably anoxic conditions at -35 cm depth only (Table 2). Frequent or prevailing unsaturated conditions above this depth would favor methanotrophy and thus bias the interpretation of  $\alpha_{C}$ . Given that  $\alpha_{C}$  values between 1.04 and 1.055 indicate the prevalence of the acetoclastic CH<sub>4</sub> production pathway, whereas  $\alpha_{C}$  values higher than 1.065 support a shift towards the hydrogenotrophic pathway, the acetoclastic pathway was apparently favored in July and August at the sites 1 and 2, in August at site 3 and in July, August and September at site 4. A shift towards a higher contribution of the hydrogenotrophic
- 20 pathway was observed in June and September at site 1, and in June at site 2

#### 3.5 $\delta^{13}$ C signatures of emitted CH<sub>4</sub> during summer 2015

Values of  $\delta^{13}$ C of emitted CH<sub>4</sub> ranged from  $-81.87 \pm 3.81$  to  $-55.61 \pm 1.20$  ‰ (see Fig. 6, panel (a) to (d)). Thereby,  $\delta^{13}$ C-CH<sub>4</sub> signatures increased from July to August and slightly decreased again in September. This pattern was thus related to the course of the wtd. Significant differences were only found at sites 3 and 4 between July and August (p < 0.01, p < 0.05),

- 25 however, from visual inspection of the panels (a) (d) of Fig. 6  $\delta^{13}$ C–CH<sub>4</sub> values seemed to increase between May and September at site 3, while they appeared to decrease at site 4, with distinctly low values in August. There was no such pattern at sites 1 and 2. Taking  $\delta^{13}$ C–CH<sub>4</sub> signatures from all sites, isotopic signatures in July differed significantly from those in August and September (p < 0.05). In September, isotopic signatures of the CH<sub>4</sub> flux at site 2 differed significantly from those at the other sites (p < 0.05).
- 30 Comparing isotopic signatures of dissolved, strongly <sup>13</sup>C depleted  $CH_4$  in the upper 35 cm of the peat and emitted, less <sup>13</sup>C depleted  $CH_4$ , plant mediated transport was the dominant  $CH_4$  emission pathway during the summer 2015 at all sites and all sampling dates according to Hornibrook (2009) (Fig. 6 (e)).

#### 4 Discussion

As expected from our studied transect ranging from a strongly altered to an only slightly altered site in terms of nutrient supply, hydrological conditions, and coverage of PFTs, we observed pronounced differences in gas fluxes and peat quality. Besides

5 all differences, the dominant CH<sub>4</sub> emission pathway was plant-mediated transport at all sites. We are aware that effects of anthropogenic impact are much more difficult to constrain in an *in-situ* study as ours, compared to well defined ecosystem manipulation studies. Nevertheless, our results support an obvious interplay of processes, fluxes, and vegetation that can be related to the observed impacts of nutrient enrichment and altered hydrology, as discussed in the following paragraphs.

#### 4.1 Long-term insights into carbon cycling at the sites

- 10 Long-term plant community changes were recently shown to affect peatland organic matter composition (Hodgkins et al., 2014), while such an effect was not identified in a short-term study (Robroek et al., 2015). Along our transect of study sites, we observed the highest degree of bulk peat decomposition in the upper peat layers of site 4, which was located in closest vicinity to the water reservoir and which was the most altered one among our four sites (Fig 2 (a)). Our initial hypothesis 1 that peripheral sites feature accelerated C cycling, reflected in more decomposed peat, could thus only partly be verified: the
- 15 fact that we did not find a gradual decrease in terms of degree of bulk peat decomposition with increasing distance from the reservoir, but observed significant differences only for site 4, suggests that the observed differences could also be primarily induced by the shift to a predominance of shrubs. Shrubs contain more woody parts and thus have higher lignin contents and more phenolic groups than graminoids or mosses and they are also more productive than mosses and graminoids (Bragazza et al., 2007). In recent studies, an increasing ericaceous shrub cover was associated with increasing polyphenol content in plant
- 20 litter and pore-water, as well as increasing phenol oxidase in litter of ericaceous shrubs. Also, a higher release of labile C from vascular plant roots was observed. Increases in shrub cover, observed along an altitudinal gradient reflecting altered temperature regimes, were accompanied by a decreasing *Sphagnum* productivity (Bragazza et al., 2013; Bragazza et al., 2015). Even though at site 4 we primarily dealt with eutrophication, rather than warming, a similar explanation may apply to our observations: Shrubs outcompete *Sphagnum* mosses after long-term nutrient infiltration and a reduced recalcitrance of the
- 25 peat arising from shrub litter can result in a reduced C storage, i.e. peat accumulation (Turetsky et al., 2012; Larmola et al., 2013; Ward et al., 2013). This explanation is further suggested by the lowest observed CO<sub>2</sub> uptake and lowest DIC concentrations along peat profiles throughout the study period at that particular site. Pore-water DOM quality indices at site 3 revealed a significantly lower share of aromatic compounds, and thus suggested a

lower degree of humification and comparably increased molecular weight at that site (Fig 2 (c) – (e)). This more labile nature

30 of DOM compared to otherwise similar bulk peat quality suggested either an input from the vegetation (Robroek et al., 2015) or some inflow of water and solutes from the nearby reservoir. Given that the site 4 pore–water DOM characteristics differed strongly from those at site 3, as did predominant PFTs, the distinctive features of the site 3 pore–water were probably also induced by the vegetation. However, the fact that the vegetational composition of site 3 and site 2 were rather similar, whereas the pore–water DOM quality was again significantly different, suggested that DOM properties were likely affected by both vegetation, i.e. photosynthetic productivity and concomitantly higher input of labile compounds, and by inflow of DOM from the reservoir.

5 The nature of our results does of course not allow for an unambiguous conclusion in terms of whether it is the vicinity to the reservoir or the plant community composition, which drives C cycling and peat accumulation at the sites. However, since peatland plant community compositions are known to be remarkably stable over time but experience changes in relative abundances (Rydin and Barber, 2001, Bragazza et al., 2006), we suggest that it was probably the vicinity to the reservoir that shaped the plant community composition at the sites over time, whereas the plant community actually drives C cycling.

#### 10 4.2 Seasonal development of carbon fluxes

Different PFTs were recently shown to have a strong impact on peatland ecosystem  $CO_2$  fluxes (Ward et al., 2013; Kuiper et al., 2014). This could be confirmed by our results: shrub dominated site 4 showed the lowest cumulative  $CO_2$  uptake, whereas, at the graminoid–moss dominated sites 3 and 2, and at the moss dominated site 1, very high  $CO_2$  uptake rates were observed. The  $CO_2$  uptake rates of our sites 1, 2 and 3 exceeded reported  $CO_2$  uptake rates of bogs by far: for instance Teklemariam et

- al. (2010) reported a net ecosystem exchange of the ombrotrophic, continental Mer Bleue bog of -140 to -20 g C m<sup>-2</sup> while hollow CO<sub>2</sub> uptake rates of our study were notably higher, rather comparable to uptake rates reported for fens (Lund et al., 2010). These latter values also compare well with the surface peat accumulation rates of ~200 to ~300 g C m<sup>-2</sup> observed at our site (Berger et al. (2017)). NEP of site 4 was significantly lower compared to the other sites fluxes indicating less CO<sub>2</sub> uptake. In the light of the strong alterations in terms of vegetation cover and the most decomposed surface peat at site 4, our findings
- 20 from an *in-situ* transect support earlier findings of reduced net  $CO_2$  uptake and a concomitantly promoted vascular plant community in a controlled long-term fertilization experiment at the Mer Bleue bog (Bubier et al., 2007). Partitioning of NEP into R<sub>eco</sub> and GPP further illustrated that the observed differences in  $CO_2$  fluxes between sites were predominantly driven by varying GPP, while R<sub>eco</sub> of all sites was in a comparable range.

With regard to CH<sub>4</sub> emissions, site 3 exceeded the other three sites by on average 30 %. In existing studies, greatest emissions

- were similarly found in wetter habitats dominated by graminoids (Levy et al., 2012, Gray et al. 2013). Given that  $CH_4$  emissions of site 2 were significantly smaller than those from site 3, even though the two sites featured a very similar graminoid–moss dominated vegetation cover, the differences in  $CH_4$  fluxes could either be attributed to i) the wetter conditions at site 3 or ii) a greater nutrient supply at site 3, stimulating greater  $CH_4$  production and emissions (Eriksson et al., 2010) or iii) a mixture of both effects. Interestingly, site 4, which experienced similar water table fluctuations like site 3, but featured a notably different
- 30 vegetation cover, emitted CH<sub>4</sub> in a similar range like sites 1 and 2. High CH<sub>4</sub> production due to input of labile organic matter nearby the reservoir was probably outweighed by less plant–mediated CH<sub>4</sub> transport and therefore emission due to a lower graminoid cover.

In our study NEP, GPP and  $CH_4$  emissions were negatively correlated with  $CH_4$  and DIC concentrations in the uppermost 50 cm of the profiles at the sites 1, 2 and 3. Such a decoupling of  $CO_2$  and  $CH_4$  fluxes from pools in the peat was already observed in previous studies: Graminoids are known to be important facilitators of  $CH_4$  emissions because they can transport  $CH_4$  from deeper, water–saturated layers of the peat into the atmosphere via aerenchymatous tissue and bypass the zone of  $CH_4$  oxidation

- Shannon and White, 1994; Marushchak et al., 2016). Moreover, they supply exudates via their roots, stimulating microbial activity and accordingly methanogenesis (Bubier et al., 1995). Through their deeper rooting system, graminoids may thus have connected the CH<sub>4</sub> pools of deeper layers below our studied profile, i.e. below 50 cm depth, to fuel the observed surface fluxes. This effect is supported by the observed differences in isotopic signatures of CH<sub>4</sub> in the peat and CH<sub>4</sub> emitted (see below). The decreasing concentrations of CH<sub>4</sub> in near surface layers due to a decrease in water table levels and partial aeration did thus not
- 10 translate in lower fluxes, a similar effect as suggested by Strack et al. (2006). Moreover, at low water tables and unsaturated conditions diffusivity for  $CO_2$  increases, leading to notably higher diffusive fluxes despite low concentrations (Knorr et al., 2008). It is striking that DIC concentrations at site 4 were notably lower as compared with other sites. A reasonable explanation is a lower peat quality resulting from repeated peat oxygenation upon water table fluctuations of the reservoir, stimulating microbial decomposition in the presence of deciduous shrubs (Bragazza et al., 2016), which are apparently promoted in closer
- vicinity to the eutrophic water reservoir. Such effect of aeration might appear contradictory, as wetter conditions would be expected near the water reservoir. However, repeated water table fluctuations driven by management of the reservoir could effectively recharge electron acceptor pools to support ongoing decomposition, as e.g. shown in water table manipulation experiments (Blodau et al., 2004; Knorr et al., 2009). Moreover, near the reservoiran advective redistribution and removal of CO<sub>2</sub> and CH<sub>4</sub> through advective flow cannot be excluded.

#### 20 4.3 Methane production, methanotrophy and pathways of CH<sub>4</sub> emissions as inferred from stable isotopes

Distinguishing CH<sub>4</sub> production pathways in peatlands using  $\delta^{13}$ C–signatures along depth profiles is a common approach (e.g. Holmes et al., 2015; McCalley et al., 2014; Hodgkins et al., 2014; Kotsyurbenko et al., 2004; Chasar et al., 2000). However, methanogenesis is a strictly anaerobic process and thus saturated, anoxic conditions are a prerequisite for an unbiased differentiation of pathways using <sup>13</sup>C only (Conrad, 1996). Methanotrophy would otherwise bias the interpretation of <sup>13</sup>C

- 25 isotopic signatures of CH<sub>4</sub>, as residual CH<sub>4</sub> gets enriched in <sup>13</sup>C, mimicking values as observed under methanogenic conditions predominated by the acetoclastic pathway (Whiticar, 1999; Alstad and Whiticar, 2011). Indeed, summer wtd at all study sites dropped down to -32.5 cm (site 1), -31.8 cm (site 2), -13.3 (site 3) and -19.1 cm (site 4) below surface and we could thus only assume saturated, anoxic conditions below that depth. We will limit the discussion of CH<sub>4</sub> production pathways accordingly. For shallower depths, effects of under such conditions much more favorable methanotrophy can be expected to
- 30 predominate: If the proportion of methanogenesis vs. methanotrophy is comparatively shifted toward methanogenesis, a relative  ${}^{13}C$ -CH<sub>4</sub> depletion would be detected, and if the proportion of methanogenesis vs. methanotrophy is comparatively shifted toward methanotrophy, a relative  ${}^{13}C$  enrichment in CH<sub>4</sub> would be detected.

 $CH_4$  oxidation was accordingly observed in the top -5 to -15 cm along our study transect during the summer months, with least negative  $\delta^{13}C$ - $CH_4$  values at 5 cm depth of site 1. Moreover,  $\delta^{13}C$ - $CH_4$  signatures at 5 cm depth of different sampling dates appeared to be most variable at the sites 1 and 2, which were also found to be drier than the sites 3 and 4, where less pronounced shifts of  $\delta^{13}C$ - $CH_4$  signatures occurred throughout the sampling period. However, also at the latter sites, variations

- 5 in δ<sup>13</sup>C–CH<sub>4</sub> were apparently driven by fluctuations of the water table levels, suggesting that CH<sub>4</sub> oxidation must have been an important factor throughout the dry season in summer. Another interesting finding was the strong δ<sup>13</sup>C–CH<sub>4</sub> signal pointing to notable CH<sub>4</sub> oxidation at 15 cm depth of site 4 in August 2015 (–39.10 ‰) as compared to more <sup>13</sup>C depleted CH<sub>4</sub> (–57.73 ‰) in 5 cm depth. This was probably due to the particularly low CH<sub>4</sub> concentrations, suggesting an input of atmospheric CH<sub>4</sub> (~-55 ‰) into the surface peat. Site 4 also featured the most enriched δ<sup>13</sup>C–CH<sub>4</sub> signatures in general, suggesting either least
- 10 CH<sub>4</sub> production or most CH<sub>4</sub> oxidation here. Site 3, showed the smallest variations in  $\delta^{13}$ C–CH<sub>4</sub> signatures throughout the sampling period, suggesting least modification of  $\delta^{13}$ C–CH<sub>4</sub> from oxidation here, which corresponds well with greatest CH<sub>4</sub> emissions and in general highest water levels measured at that site.

Overall lowest  $\delta^{13}$ C–CO<sub>2</sub> values were found at site 4, where simultaneously least negative values of  $\delta^{13}$ C–CH<sub>4</sub> were observed, suggesting a higher share of CO<sub>2</sub> from increased CH<sub>4</sub> oxidation. CO<sub>2</sub> generally got more enriched in <sup>13</sup>C with depth at all sites

- and sampling dates, as expected from ongoing fractionation by methanogenesis. Great shifts in  $\delta^{13}$ C–CO<sub>2</sub> values of the drier sites 1 and 2 during the entire sampling period could again be explained by increased exchange of peat derived CO<sub>2</sub> with atmospheric CO<sub>2</sub> under unsaturated conditions with dropping water tables in August. Regarding observed ranges of  $\alpha_{c}$  values at –35 cm depth at the sites, also a gradient in terms of the CH<sub>4</sub> production pathway
- along the transect of study sites became apparent. The sites 1 and 2, which experienced the lowest water tables during the summer, and which were located in farthest distance to the water reservoir, featured a distinct shift from mostly hydrogenotrophic CH<sub>4</sub> production in June to acetoclastic CH<sub>4</sub> production in July and August and another shift back to hydrogenotrophic CH<sub>4</sub> production in September, with these shifts being more pronounced at site 1. This could be related to increased vascular plant activity in the growing season and concomitant substrate supply to methanogens, e.g. through exudation; an increased share of acetoclastic methanogenesis within the rhizosphere has previously been reported (Chasar et
- al., 2000; Hornibrooket al., 2007). At the sites 3 and 4, such obvious shifts of  $CH_4$  production pathways could not be observed, though;  $\alpha_C$  values indicated either acetoclastic  $CH_4$  production or a co-occurrence of acetoclastic and hydrogenotrophic  $CH_4$ production. As acetoclastic methanogenesis is in particular supported in minerotrophic peatlands in presence of vascular plants (Alstad and Whiticar, 2011; Chasar et al., 2000), predominance of that pathway –in particular in closer vicinity to the reservoir– is not a surprising finding for Wylde Lake peatland. Indeed, under predominance of sedges, which supply labile organic matter
- 30 through roots, aceotclastic CH<sub>4</sub> production prevailed (Bellisario et al., 1999; Popp et al., 1999; Strom et al., 2003). However, the fact that CH<sub>4</sub> production pathways at the sites 3 and 4 (different vegetation), were similar, whereas CH<sub>4</sub> production pathways at the sites 2 and 3 (similar vegetation) were different, suggested that variation of  $\alpha_C$  would rather reflect the impact of the reservoir, by either a) sustaining higher water tables, or b) increased nutrient input, than the presence of sedges at the sites.

The emitted CH<sub>4</sub> (see Fig. 6 (a) – (d)) was in general depleted in <sup>13</sup>C compared to the CH<sub>4</sub> in (see Fig. 5) all sampled peat layers. This suggests that the emitted CH<sub>4</sub> must have been produced in the deeper peat layers (Marushchak et al., 2016), where  $\delta^{13}$ C–CH<sub>4</sub> signatures were probably more depleted and during transport through plant aerenchyma, the lighter CH<sub>4</sub> could bypass oxidation. Moreover, plant–mediated transport also slightly discriminates against the <sup>13</sup>C–CH<sub>4</sub> (Chanton, 2005),

- 5 favoring more negative values of  $\delta^{13}$ C in emitted CH<sub>4</sub>. Interestingly, plant mediated transport was the dominant CH<sub>4</sub> emission pathway even at the sites 1 and 4, where graminoid cover accounted only for about 10 %. We suggest that this is due to the great CH<sub>4</sub> oxidation in the upper peat layers and rather high concentrations at greater depth, facilitating plant–mediated transport and ebullition. From visual inspection of the panels (a) – (d) of Fig. 6 and Fig. 5 we suggest that the emitted CH<sub>4</sub> originated from at least –35 cm depth or below.
- 10 Hypothesis 2 stating that increased abundance of vascular plants can increase  $CO_2$  uptake but also change patterns of  $CH_4$ production and emission, in particular if graminoids dominate, can only partly be accepted. If increased vascular plant cover translated into increased  $CO_2$  uptake, we should have observed increasing uptake in the order of site 1 < 2 = 3 < 4, but in fact we observed only significantly decreased uptake at site 4. The  $CO_2$  uptake at site 1 (the site with the least coverage of vascular plants) was not statistically different from the cumulative NEP observed at the sites 2 and 3. Moreover, we cannot directly
- 15 state that  $CH_4$  production and emission was increased where graminoids dominated. Although greatest  $CH_4$  emission was observed at site 3, cumulative  $CH_4$  emission at site 2 was significantly lower, despite relatively similar vegetation. Besides by PFTs,  $CH_4$  production appeared to be affected by the vicinity of the water reservoir, whereas plant–dominated  $CH_4$  emission was the dominant  $CH_4$  emission pathway at all study sites, even where graminoid coverage accounted for only 10 %. So, we conclude that an interplay of nutrient input, water table depth, and vegetational composition shaped  $CO_2$  uptake,  $CH_4$
- 20 production and emissions and there was likely no unique driver in our *in-situ* study, compared to well defined manipulation experiments.

#### **5** Concluding remarks

Our study and earlier work at this particular site confirm that despite long-term increased nutrient supply, peatland ecosystem functioning in terms of C sequestration was largely maintained. However, along a sequence of study sites it became apparent

- 25 that the affected sites responded differently to the altered conditions after dam construction in 1954. Shrub dominated site 4, which was in closest vicinity to the reservoir and accordingly faced greatest nutrient input and most pronounced water level fluctuations, indeed showed indications of degradation, such as most decomposed bulk peat, least atmospheric CO<sub>2</sub> uptake, and reduced coverage of *Sphagnum* mosses. However, even here, overall net CO<sub>2</sub> uptake still exceeded net CO<sub>2</sub> release. The two graminoid–moss dominated sites and the moss dominated site featured very high CO<sub>2</sub> uptake rates despite apparent impact
- 30 of nutrients and altered hydrology. Therefore, as hypothesized, our case study supports that long-term nutrient enrichment in combination with hydrologically altered conditions may cause differential responses of C cycling and do not necessarily cause a loss of the C-sink function of peatland ecosystems.

Moreover, methanogenesis and methanotrophy featured a pattern which appeared to be related not predominantly to vegetation, but primarily to the vicinity to the reservoir and thus nutritional status and hydrologic regime. On the other hand, plant-mediated transport was determined to be the dominant  $CH_4$  emission pathway at all sites, even if graminoid cover was only 10 %. All surface peat layers indicated high methanotrophic activity, mitigating  $CH_4$  emission through diffusion.

- 5 Lastly, our results suggest that with regard the overall C budget, a graminoid-moss dominated peatland site can obviously withstand eutrophication in combination with frequent inundation better than a shrub dominated peatland site. Straightforward results from manipulation experiments of individual factors (e.g. fertilization or water table changes) may therefore not be easily transferred to complex *in-situ* conditions. We suggest that there could be a tipping point, when a peatland system shifts from a net C sink –even though experiencing eutrophic conditions– to decreasing productivity, which might be related to an
- 10 expansion of shrub dominated vegetation, decreasing overall C uptake.

#### 6 Data availability

The data can be accessed by email request to the corresponding authors.

15 Author contributions. Christian Blodau, Klaus-Holger Knorr and Sina Berger designed the experiments; Sina Berger, Leandra Praetzel and Marie Goebel conducted field work and analyses with help of Klaus-Holger Knorr. Sina Berger prepared the manuscript with contributions from Klaus-Holger Knorr, Leandra Praetzel and Marie Goebel.

The authors declare that they have no conflict of interest.

20

30

*Acknowledgements*. This study was funded by the Deutsche Forschungsgemeinschaft (German Research Foundation - DFG) (BL563/21-1) We thank Martin Neumann from the Grand River Conservation Authority for the permission to carry out research in the Luther Marsh Wildlife Management Area and we thank Claudia Wagner-Riddle, Peter Smith and Linda Wing for their help on organizational issues. We also thank Inge-Beatrice Biro, Magdalena Burger, Ines Spangenberg, Niclas Kolbe,

Eike Esders, Michael Rammo, Niels Vickus, Fabian Benninghoff, Leonie Fröhlich, Jörg Rostek and Cornelia Mesmer for their support in the field. Analyses of CO<sub>2</sub> and CH<sub>4</sub> concentrations,  $\delta^{13}$ C abundance and spectral analyses of various samples were carried out at the institutional lab or the Institute of Landscape Ecology, University of Münster. We thank Stefanie Holm, Ronya Wallis and Madelaine Supper for assistance during analysis of numerous samples in the laboratory.

This paper is dedicated to the memory of Christian Blodau, who led the Wylde Lake peatland project until he tragically passed away in July 2016.

We thank two anonymous reviewers for their thoughtful comments that helped to improve this manuscript.

#### References

Alstad, K. P., Whiticar, M. J.: Carbon and hydrogen isotope ratio characterization of methane dynamics for Fluxnet peatland ecosystems, Org Geochem, 42(5), 548–558, 2011.

5 Ballantyne, D. M., Hribljan, J. A., Pypker, T. G., Chimner, R. A.: Long- term water table manipulations alter peatland gaseous carbon fluxes in Northern Michigan, Wetl Ecol Manag, 22, 35–47, 2014.

Berger, S., Gebauer, G., Blodau, C., Knorr, K.-H.: Peatlands in a eutrophic world – assessing the state of a poor fen-bog transition in southern Ontario, Canada, after long term nutrient Input and altered hydrological conditions, Soil Biol Biochem, 114, 131–144, 2017.

Bellisario, L. M., Bubier, J. L., Moore, T. R.: Controls on CH<sub>4</sub> emissions from a northern peatland, Global Biogeochem Cycles, 13(1), 81–91, 1999.

15 Biester, H., Knorr, K.-H., Schellekens, J., Basler, A., and Hermanns, Y.-M.: Comparison of different methods to determine the degree of peat decomposition in peat bogs, Biogeosciences, 11, 2691–2707, 2014.

Blodau, C.: Carbon cycling in peatlands - A review of processes and controls, Environ Rev, 10, 111-134, 2002.

20 Bragazza, L.: A decade of plant species changes on a mire in the Italian Alps: vegetation-controlled or climate-driven mechanisms?, Clim Change, 77, 415–429, 2006.

Bragazza, L., Siffi, C., Lacumin, P., Gerdol, R.: Mass loss and nutrient release during litter decay in peatland: The role of microbial adaptability to litter chemistry, Soil Biol Biochem, 39(1), 257–267, 2007.

10

Bragazza, L., Buttler, A., Habermacher, J., Brancaleoni, L., Gerdol, R., Fritze, H., Hanjík, P., Laiho, R., Johnson, D.: High nitrogen deposition alters the decomposition of bog plant litter and reduces carbon accumulation, Glob Chang Biol, 18, 1163–1172, 2012.

30 Bragazza, L., Parisod, J., Buttler, A., Bardgett, R. D.: Biogeochemical plant-soil microbe feedback in response to climate warming in peatlands, Nat Clim Change, 3, 273–277, 2013.

<sup>25</sup> 

Bragazza, L., Bardgett, R. D., Mitchell, E. A. D., Buttler, A.: Linking soil microbial communities to vascular plant abundance along a climate gradient, New Phytol, 205, 1175–1182, 2015.

Bragazza, L., Buttler, A., Robroek, B., Albrecht, R., Zaccone, C., Jassey, V., Signarbieux, C.: Persistent high temperature and
low precipitation reduce peat carbon accumulation, Glob Change Biol, 22, 4114–4123, 2016.

Broder, T., Blodau, C., Biester, H., Knorr, K.-H.: Peat decomposition records in three pristine ombrotrophic bogs in southern Patagonia, Biogeosciences, 9, 1479–1491, 2012.

10 Bubier, J. L., Moore, T. R., Bellisario, L., Comer, N. T., Crill, P. M.: Ecological controls on methane emissions from a Northern Peatland Complex in the zone of discontinuous permafrost, Manitoba, Canada, Glob Biogeochem Cycles, 9(4), 455–470, 1995.

Bubier, J. L., Moore, T. R., Bledzki, L. A.: Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog, Glob Chang Biol, 13, 1168–1186, 2007.

#### 15

Burger, M., Berger, S., Spangenberg, I., Blodau, C.: Summer fluxes of methane and carbon dioxide from a pond and floating mat in a continental Canadian peatland, Biogeosciences, 13, 3777–3791, 2016.

Chanton, J. P.: The effect of gas transport on the isotope signature of methane in wetlands, Org Geochem, 36 (5), 753–768, 20 2005.

Chasar, L. S., Chanton, J. P., Glaser, P. H., Siegel, D. I.: Methane Concentration and Stable Isotope Distribution as Evidence of Rhizospheric Processes: Comparison of a Fen and Bog in the Glacial Lake Agassiz Peatland Complex, Ann Bot, 86, 655–663, 2000.

#### 25

Conrad, R.: Quantification of methanogenic pathways using stable carbon isotopic signatures. A review and a proposal, Org Geochem, 36(5), 739–752, 2005.

Eriksson, T., Öquist, M. G., Nilsson, M. B.: Production and oxidation of methane in a boreal mire after a decade of increased temperature and nitrogen and sulfur deposition, Glob Chang Biol, 16, 2130–2144, 2010.

Gray, A., Levy, P. E., Cooper, M. D. A., Jones, T., Gaiawyn, J., Leeson, S. R., Ward, S. E., Dinsmore, K. J., Sheppard, L. J., Ostle, N. J., Evans, C. D., Burden, A., Zielinski, P.: Methane indicator values for peatlands: a comparison of species and functional groups, Glob Change Biol, 19, 1141–1150, 2013.

Heidel, D.: Isotopenfraktionierungseffekte bei der Beprobung von Bodenluft und gelösten Gasen, unpublished Bacherlor's Thesis, 2009.

5 Hesslein, R. H.: Insitu Sampler for close interval pore water studies, Limnol Oceanogr, 21(6), 912–914, 1976.

Hodgkins, S. B., Tfaily, M. M., McCalley, C. K., Logan, T. A., Crill, P. M., Saleska, S. R., Rich, V. I., Chanton, J. P.: Can changes in peat chemistry associated with permafrost thaw increase greenhouse gas production, PNAS, 111(16), 5819–5824, 2014.

10

Hoefs, J.: Stable Isotope Geochemistry, 3<sup>rd</sup> ed., Springer, Berlin, 1987.

Holmes, M. E., Chanton, J. P., Tjaily, M. M., Ogram, A.: CO<sub>2</sub> and CH<sub>4</sub> isotope compositions and production pathways in a tropical peatland, Glob Biochem Cycles, 29, 1–18, 2015.

15

Hornibrook, E. R. C., Bowes, H. L.: Trophic status impacts both the magnitude and stable carbon isotope composition of methane flux from peatlands, Geophys Res Lett, 34(21), 2007.

Hornibrook, E. R. C.: The stable carbon isotope composition of methane produced and emitted from northern peatlands.
Andrew J. Baird, Lisa R. Belyea, Xavier Comas, A. S. Reeve und Lee D. Slater (Ed.): Carbon Cycling in Northern Peatlands, Bd.184. Washington, D. C.: American Geophysical Union (Geophysical Monograph Series), pp. 187–203, 2009.

Kammann, C., Grünhage, L., Jäger, H.-J.: A new sampling technique to monitor concentrations of CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> in air at well-defined depths in soils with varied water potential, Eur J Soil Sci, 52, 297–303, 2001.

25

Kim, Y., Ullah, S., Roulet, N. T., Moore, T. R.: Effect of inundation, oxygen and temperature on carbon mineralization in boreal ecosystems, Sci Total Environ, 511, 381–392, 2015.

Klüpfel, L., Piepenbrock, A., Kappler, A., Sander, M.: Humic substances as fully regenerable electron acceptors in recurrently 30 anoxic environments, Nat Geosci, 7, 195–200, 2014.

Knorr, K.-H., Oosterwoud, M. R., Blodau, C.: Experimental drought alters rates of soil respiration and methanogenesis but not carbon exchange in soil of a temperate fen, Soil Biol Biochem, 40(7), 1781–1791, 2008.

Kotsyurbenko, O. R., Chin, K.-J., Glagolev, M. V., Stubner, S., Simankova, M. V., Nozhevnikova, A. N., Conrad, R.: Acetoclastic and hydrogenotrophic methane production and methanogenic populations in an acidic West-Siberian peat bog, Environ Microbiol, 6 (11), 1159–1173, 2004.

5 Kuiper, J. J., Mooij, W. M., Bragazza, L., Robroek, B. J. M.: Plant functional types define magnitude of drought response in peatland CO<sub>2</sub> exchange, Ecology, 95(1), 123–131, 2014.

Laine, J., Harju, P., Timonen, T., Laine, A., Tuittila, E.-S., Minkkinen, K., Vasander, H.: The Intricate Beauty of *Sphagnum* mosses – a Finnish Guide to Identification, 2<sup>nd</sup> amended edition, 2011.

10

Larmola, T., Bubier, J. L., Kobyljanec, C., Basiliko, N., Juutinen, S., Humphreys, E., Preston, M., Moore, T. R.: Vegetation feedbacks of nutrient addition lead to a weaker carbon sink in an ombrotrophic bog, Glob Change Biol, 19, 3729–3739, 2013.

Levy, P. E., Burden, A., Cooper, M. D. A., Dinsmore, K. J., Drewer, J., Evans, C., Fowler, D., Gaiawyn, J., Gray, A., Jones, S. K., Jones, T., Mcnamara, N. P., Mills, R., Ostle, N., Sheppard, L. J., Skiba, U., Sowerby, A., Ward, S. E., Zielinski, P.:

15 S. K., Jones, T., Mcnamara, N. P., Mills, R., Ostle, N., Sheppard, L. J., Skiba, U., Sowerby, A., Ward, S. E., Zielinski, P.: Methane emissions from soils: synthesis and analysis of a large UK data set, Glob Change Biol, 18, 1657–1669, 2012.

Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H., Schaepman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications – a synthesis, Biogeosciences, 5, 1475–1491, 2008.

20

Lund, M., Lafleur, P. M., Roulet, N. T., Lindroth, A., Christensen, T. R., Aurela, M., Chojnicki, B. H., Flanagan, L. B., Humphreys, E. R., Laurila, T., Oechel, W., Olejnik, J., Rinne, J., Schubert, P., Nilsson, M. B.: Variability in exchange of CO<sub>2</sub> across 12 northern peatland and tundra sites, Glob Change Biol, 16, 2436–2448, 2010.

25 Marushchak, M. E., Friborg, T., Biasi, C., Herbst, M., Johannson, T., Kiepe, I., Liimatainen, M., Lind, S. E., Martikainwn, P. J., Virtanen, T., Soegaard, H., Shurpali, N. J.: Methane dynamics in the subarctic tundra: combining stable isotope analyses, plot- and ecosystem-scale flux measurements, Biogeosciences, 13, 597–608, 2016.

McCalley, C., Woodcroft, B., Hodgkins, S., Wehr, R., Kim, E.-H., Mondav, R., Crill, P., Chanton, J., Rich, V., Tyson, G.,
Saleska, S.: Methane dynamics regulated by microbial community response to permafrost thaw, Nature, Vol. 514, no 7523, 478–481, 2014.

Morris, P. J., Waddington, J. M., Benscoter, B. W., Turetsky, M. R.: Conceptual frameworks in peatland ecohydrology: looking beyond the two-layered (acrotelm–catotelm) model, Ecohydrology, 4, 1–11, 2011.

Murphy, K. R., Butler, K. D., Spencer, R. G., Stedmon, C. A., Boehme, J. R., Aiken, G. R.: Measurement of dissolved organic matter fluorescence in aquatic environments: an interlaboratory comparison, Environ Sci Technol, 44, 9405–9412, 2010.

5 National Climate Data and Information Archive, 2014. Canadian Climate Normals. Dataset for the climate station Fergus, Shand Dam, 1981 to 2010. URL http://climate.weather.gc.ca/climate normals/index e.html (accessed November 18th, 2015).

Niemeyer, J., Chen, Y., Bollag, J. M.: Characterization of humic acids, composts, and peat by diffuse reflectance Fourier-Transform Infrared Spectroscopy, Soil Sci Soc Am J, 56, 135–140, 1992.

10

Nijp, J. J., Limpens, J., Metselaar, K., van der Zee, S. E. A. T. M., Berendse, F., Robroek, B. J. M.: Can frequent precipitation moderate the impact of drought on peatmoss carbon uptake in northern peatlands?, New Phytol, 203, 70–80, 2014.

Ohno, T.: Fluorescence inner-filtering correction for determining the humification index of dissolved organic matter, Environ 15 Sci Technol, 36, 742–746, 2002.

Owen, K. E., Tenhunen, J., Reichstein, M., Wang, Q., Falge, E., Geyer, R., Xiao, X. M., Stoy, P., Amman, C., Arain, A.,
Aubinet, M., Aurela, M., Bernhofer, C., Chojnicki, B. H., Grainer, A., Gruenwald, T., Hadley, J., Heinsch, B., Hollinger, D.,
Knohl, A., Kutsch, W., Lohila, A., Meyers, T., Moors, E., Moureaux, C., Pilegaard, K., Saigusa, N., Verma, S., Vesala, T.,
Vogel, C.: Linking flux network measurements to continental scale simulations: ecosystem carbon dioxide exchange capacity
under non-water-stressed conditions, Glob Change Biol, 13(4), 734–760, 2007.

Pack, M.A., Xu, X., Lupascu, M., Kessler, J.D., Czimczik, C.I.: A rapid method for preparing low volume CH<sub>4</sub> and CO<sub>2</sub> gas samples for <sup>14</sup>C AMS analysis, Org Geochem, 78, 89–98. 2015.

25

20

Popp, T. J., Chanton, J. P., Whiting, G. J., Grant, N.: Evaluation of methane in the rhizosphere of a Carex dominated fen in north central Alberta, Canada, Biogeochemistry, 51, 259–281, 2000.

R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, 30 2015.

Robroek, B., Jassey, V. E. J., Kox, M., A. R., Berendsen, R. L., Mills, R. T. E., Cécillon, L., Puissant, J., Meima-Franke, M., Bakker, P. A. H. M., Bodelier, P. L. E.: Peatland vascular plant functional types affect methane dynamics by altering microbial community structure, J Ecol, 103, 925–934, 2015.

Rydin, H., Barber, K. E.: Long-term and fine-scale coexistence of closely related species, Folia Geobot, 36, 53-61, 2001.

Rydin, H., Jeglum, J. K.: The Biology of Peatlands, 2<sup>nd</sup> ed., Oxford University Press, 2013.

5

Sander, R.: Compilation of Henry's Law Constants for Inorganic and Organic Species of Potential Importance in Environmental Chemistry, Max Planck Institute of Chemistry, Mainz, 1999. Online at http://www.henrys-law.org/henry-3.0.pdf.

Schütz, H., Schröder, P., Rennenberg, H.: Role of plants in regulating the methane flux to the atmosphere, Sharkey, T.D., Holland, E.A., Mooney, H.A. (Eds.), Trace Gas Emission by Plants, Academic Press, NY, pp. 29–69, 1991.

Shannon, R. D., White, J. R.: A Three-Year Study of Controls on Methane Emissions from Two Michigan Peatlands, Biogeochemistry, 27(1), 35–60, 1994.

15

Sheppard, L. J., Leith, I. D., Leeson, S. R., vam Dijk, N., Field, C., Levy, P.: Fate of N in a peatland, Whim bog: immobilisation in the vegetation and peat, leakage into pore water and losses as N2O depend on the form of N, Biogeosciences, 10, 149–160, 2013.

20 Strack, M., Waller, M. F., Waddington, J. M.: Sedge succession and peatland methane dynamics: A potential feedback to climate change, Ecosystems, 9(2), 278–287, 2006.

Strom, L., Ekberg, M., Mastepanov, M., Christensen, T. R.: The effect of vascular plants on carbon turnover and methane emissions from a tundra wetland, Glob Change Biol, 9, 1185–1193, 2003.

25

Stumm, W., Morgan, J. J.: Aquatic chemistry. Chemical Equilibria and Rates in Natural Waters. 3rd ed. Hoboken: Wiley (Environmental Science and Technology), 1996.

Succow, M., Joosten, H.: Landschaftsökologische Moorkunde. Zweite, völlig neu bearbeitete Auflage. 2. Auflage. E.
Schweizerbart`sche Verlagsbuchhandlung, Stuttgart. 2012.

Teh, Y. A., Silver, W. L., Conrad, M. E., Borglin, S. E., Carlson, C. M.: Carbon isotope fractionation by methane-oxidizing bacteria, J Gephys Res.: Biogeosci, 111, G01003.

Tilsner, J., Wrage, N., Lauf, J., Gebauer, G.: Emission of gaseous nitrogen oxides from an extensively managed grassland in NE Bavaria, Germany. I. Annual budgets of  $N_2O$  and  $NO_x$  emissions, Biogeochemistry, 63, 229–247, 2003.

Teklemariam, T. A., Lafleur, P. M., Moore, T. R., Roulet, N. T., Humphreys, E. R.: The direct and indirect effects of inter-

5 annualmeteorological variability on ecosystem carbon dioxide exchange at a temperate ombrotrophic bog, Agric For Meteorol, 150, 1402–1411.

Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., McCallister, L., McKnight, D. M., Melack,
J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., von Wachenfeldt, E., Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, Limnol Oceanogr, 54(6, part 2), 2298–2314, 2009.

Turetsky, M. R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frolking, S., McGuire, M. R., Tuittila, E.-S.: The resilience and functional role of moss in boreal and arctic ecosystems, New Phytol, 196(1), 49–67, 2012.

van den Berg, M., Ingwersen, J., Lamers, M., Streck, T.: The role of *Phragmites* in the CH<sub>4</sub> and CO<sub>2</sub> fluxes in a minerotrophic peatland in southwest Germany, Biogeosciences, 13, 6107–6119, 2016.

20 Ward, S. E., Ostle, N. J., Oakley, S., Quirk, H., Henrys, P. A., Baedgett, R. D.: Warming effects on greenhouse gas fluxes in peatlands are modulated by vegetation composition, Ecol Lett, 16, 1285–1293, 2013.

Wang, M., Moore, T. R., Talbot, J., Riley, J. L.: The stochiometry of carbon and nutrients in peat formation, Glob Biochem Cycles, 29, 113–121, 2015.

25

Wang, M., Larmola, T., Murphy, M. T., Moore, T. R., Bubier, J.: Stoichiometric response of shrubs and mosses to long-term nutrient (N, P and K) addition in an ombrotrophic peatland, Plant Soil, 400, 403–416, 2016.

Weishaar, J. L., Aiken, G. R., Bergamischi, B. A., Fram, M. S., Fujii, R., Mopper, K.: Evaluation of specific ultraviolet
absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon, Environ Sci Technol, 37, 4702–4708, 2003.

Whalen, S. C., Reeburgh, W. S., Sandbeck, K. A.: Rapid methane oxidation in a landfill cover soil, Appl Environ Microbiol, 56, 3405–3411, 1990.

Whalen, S. C.: Biogeochemistry of methane exchange between natural wetlands and the atmosphere, Environ Eng Sci, 22, 73–94, 2005.

5 Whiticar, M. J., Faber, E., Schoell, M.: Biogenic methane formation in marine and freshwater environments. CO<sub>2</sub> reduction vs. acetate fermentation—Isotope evidence, Geochim Cosmochim Acta, 50 (5), 693–709, 1986.

Whiting, G. J., Chanton, J. P.: Control of the diurnal pattern of the methane emission from emergent aquatic macrophytes by gas transport mechanisms, Aquat Bot, 54, 237–253, 1996.

Wiedermann, M. M., Nordin, A., Gunnarsson, U., Nilsson, M. B., Ericson, L.: Global Change Shifts vegetation and plantparasite interactions in a boreal mire, Ecology, 88(2), 454–464, 2007.

<sup>10</sup> 



Figure 1: Location of Wylde Lake peatland complex in North America (left), and sampling sites (black dots) within Wylde Lake 5 peatland complex (right). Source: ArcGIS.



Figure 2: FTIR ratios 1618.5/1033.5 in bulk peat (a) and pore-water (b) as well as SUVA<sub>254</sub>, indicating aromaticity, (c), HIX, humification index, (d) and E2:E3, indicative of molecular size and aromaticity, (e) for pore-water samples of the sites 1 to 4. n = 3. Error bars indicate +/- standard deviation.



Figure 3: Development of (a)  $T_{water}$  [°C], (b) wtd [cm], (c)  $CH_4$  fluxes [g  $CH_4$  m<sup>-2</sup> d<sup>-1</sup>] and (d) – (f)  $CO_2$  fluxes (NEP partitioned into  $R_{eco}$  and GPP) [g  $CO_2$  m<sup>-2</sup> d<sup>-1</sup>], ± 1 SD (n=6) in hollows of the sites 1–4 from April 1st, 2014 through September 22nd, 2015. Negative  $CO_2$  and  $CH_4$  fluxes indicate uptake, positive fluxes indicate a release to the atmosphere. The dashed grey line in the NEP graph indicates a 0-flux.



Figure 4: Development of mean CH<sub>4</sub> and mean DIC concentrations [ $\mu$ mol L<sup>-1</sup>], in hollows of the sites 1–4 from April 1st, 2014 through September 22nd, 2015. Concentrations were interpolated based on biweekly sampling at depths of 5, 15, 25, 35, 45 and 55 cm.



Figure 5: Profiles of  $\delta^{13}$ C-CH<sub>4</sub> (left) and  $\delta^{13}$ C-CO<sub>2</sub> (right) signatures at sites 1-4 in the peat in 5-35 cm depth at different points in 5 time. Squares = June (06/11), circles = July (07/08), triangles = August (08/27), diamonds = September (09/17). Graphs show mean values and standard deviations from three replications at each site. n = 1-3.



Figure 6: δ<sup>13</sup>C-CH<sub>4</sub> signatures (‰) of CH<sub>4</sub> fluxes from May to September for the sites 1 (a), 2 (b), 3 (c) and 4 (d). n = 1-4. In July 2015, sampling at site 2 was not possible. Bold lines are the median, boxes show the 25 and 75 percentile, whiskers indicate minima and maxima within 1.5 times the interquartile range. Single points show outlier. (e): dominant flux pathway of CH<sub>4</sub> according to (Hornibrook, 2009). Empty circles = site 1, circles with diagonal lines = site 2, circles with crosses = site 3, circles with vertical lines = site 4. Dashed line represents transport via ebullition or active plant transport without any isotopic fractionation. Values are means of pore-gas samples from 5-35 cm depth and chamber flux measurements. Graphs show mean values and standard deviations from three replications at each site. n = 1-3.

Table 1: Overview of the four sites' distances to the water reservoir and of the composition of the vegetation in terms of coverage [%] of plant functional types (PFT) in hollows and abundances of plant species (vascular plants and mosses, excluding liverworts and hornworts). Abundances are abbreviated as follows: "*d*" means "dominant" (> 75 %), "*a*" means "abundant" (51-75 %), "*f*" means "frequent" (26-50 %), "*o*" means "occasional" (11-25 %), and "*r*" means "rare" (1-10 %). Because *Sphagnum* mosses were very hard to distinguish in the field, we only determined the abundance of the most abundant *Sphagnum* species of each site.

Site 1	Site 2		Site 3		Site 4			
Distance to reservoir [m]							-	
800		550		400		200		
Coverage of PFT [%]								
Sphagnum spp.: 100		100		100		60		
Graminoids: 10		30		30		10		
Shrubs: 8		5		4		30		
<u>Plant species</u> <u>Sphagna</u>								
S. magellanicum a		S. magellanicum		S. magellanicum		S. magellanicum		
S. capillifolium a		S. capillifolium	а	S. capillifolium	а	S. capillifolium		
S. fuscum		S. fuscum		S. fuscum		S. fuscum	f	
S. squarrosum		S. wulfianum		S. girgensohnii	а	S. girgensohnii	f	
S. angustifolium		S. angustifolium	а	S. squarrosum		S. wulfianum		
				S. wulfianum				
				S. angustifolium				
				S. cuspidatum				
Other mosses								
Polytrichum spp.	0	Polytrichum spp.	о	Polytrichum spp.		Polytrichum spp.	r	
Rhytidiadelphus triquestrus	r	Polytrichum spp.	0	Polytrichum spp. o				
<u>Graminoids</u>								
Carex disperma	f	Scheuchzeria palustris	о	C. disperma	о	C. disperma	о	
Carex oligosperma	r	C. disperma	о	C. magellanica	0	E. angustifolium	r	
Eleocharis palustris	r	C. oligosperma	r	Dulichium arundinaceum	f	E. vaginatum	о	
Eriophorum angustifolium	r	Carex limosa	r	r <i>E. palustris</i> o		E. virginicum	о	
Eriophorum vaginatum	0	Carex magellanica	r	E. angustifolium	о	J. effusus	r	
Eriophorum virginicum	0	E. palustris	о	E. vaginatum	f			
Juncus effusus	r	E. angustifolium	о	E. virginicum	f			
		E. vaginatum	f	J. effusus	о			
		E. virginicum	f					
		J. effusus	r					

r	Myrica gale	r	M. gale L.		M. gale	а
r	A. glaucophylla	r	A. glaucophylla r A. g		A. glaucophylla	r
о	C. calyculata	o <i>C. calyculata</i> o		C. calyculata	о	
0	K. polifolia	r	<i>K. polifolia</i> r		K. polifolia	r
0	R. groenlandicum	r	R. groenlandicum	r	R. groenlandicum	о
r					V. oxycoccos	о
r						
r	L. laricina	r	P. strobus	r	B. pumila	r
r	P. strobus	r				
r	B. pumila	r				
r						
r	S. purpurea	r	Maianthemum trifolium	r	S. purpurea	r
r	D. rotundifolia	r	S. purpurea	r	D. rotundifolia	r
			D. rotundifolia	r		
	r o o r r r r r r r r	r Myrica gale r A. glaucophylla o C. calyculata o K. polifolia o R. groenlandicum r r L. laricina r P. strobus r B. pumila r r S. purpurea r D. rotundifolia	r Myrica gale r r A. glaucophylla r o C. calyculata o o K. polifolia r o R. groenlandicum r r r L. laricina r r P. strobus r r B. pumila r r r D. rotundifolia r	r Myrica gale r M. gale L. r A. glaucophylla r A. glaucophylla o C. calyculata o C. calyculata o K. polifolia r K. polifolia o R. groenlandicum r R. groenlandicum r r r L. laricina r P. strobus r P. strobus r r B. pumila r r r D. rotundifolia r S. purpurea D. rotundifolia	r Myrica gale r M. gale L. o r A. glaucophylla r A. glaucophylla r o C. calyculata o C. calyculata o o K. polifolia r K. polifolia r o R. groenlandicum r R. groenlandicum r r r r L. laricina r P. strobus r r P. strobus r r B. pumila r r r D. rotundifolia r S. purpurea r D. rotundifolia r	r Myrica gale r M. gale L. o M. gale r A. glaucophylla r A. glaucophylla r A. glaucophylla o C. calyculata o C. calyculata o C. calyculata o K. polifolia r K. polifolia r K. polifolia o R. groenlandicum r R. groenlandicum r R. groenlandicum r L. laricina r P. strobus r B. pumila r P. strobus r r B. pumila r r D. rotundifolia r S. purpurea r D. rotundifolia r S. purpurea r D. rotundifolia r

Table 2:  $\alpha_C$  values and standard deviation obtained from silicone samplers in 35 cm depth at sites 1 to 4 from June to September.  $\alpha_C$  values between 1.04 and 1.055 indicate the prevalence of the acetoclastic CH<sub>4</sub> production pathway,  $\alpha_C$  values higher than 1.065 indicate the hydrogenotrophic pathway.

5

Site	<u>1</u>		<u>2</u>		<u>3</u>		<u>4</u>	
	αc	std.dev	α	std.dev	α <sub>c</sub>	std.dev	α <sub>c</sub>	std.dev
June	1.068		1.064	0.004	1.061	0.004	1.056	0.004
July	1.042		1.044		1.058	0.001	1.048	
August	1.043		1.046	0.001	1.052	0.004	1.045	0.002
September	1.066	0.007	1.057	0.002	1.058	0.003	1.051	0.002