

Associate Editor Decision: Publish subject to minor revisions (Editor review) (14 Sep 2017) by Paul Stoy

Comments to the Author:

Responses to referees were comprehensive and thank you for the attention to detail. There are a few areas where the manuscript would benefit from copy-editing or minor editorial improvements that would improve accuracy and readability.

We thank the associate editor for reviewing the manuscript and for pointing out the errors in some details we overlooked. We are happy to apply the suggestions for improving the manuscript.

The following is a partial list of minor revisions that could be made to improve the manuscript.

1. I recommend a couple more careful reads by the authors before recommending the manuscript for publication.

We have gone through the manuscript and filtered out many mistakes we overlooked in the previous versions of the manuscript.

2. For the title I might recommend "in Jambi province"

We have adopted the suggested change. The title is now:

Expansion of oil palm and other cash crops causes an increase of the land surface temperature in the Jambi province in Indonesia.

3. "Surface roughness affect the transferred sensible and latent heat" on line 88 is an example of verb/subject agreement that needs to be changed.

We thank the associate editor for pointing us to this error. We corrected the sentence.

4. The statement on line 92 would benefit from a reference.

We added a reference to underline the statement in this line and rewrote the sentence to “.. LST is a major land surface parameter and as climatic factor it is regarded a main driver of diversity gradients related to the positive relationships between temperature and species richness (Wang et al., 2016).”

Wang, J., Pan, F., Soininen, J., Heino, J. and Shen, J.: Nutrient enrichment modifies temperature-biodiversity relationships in large-scale field experiments, Nat. Commun., 7, 13960, doi:10.1038/ncomms13960, 2016.

5. On line 97, this is true "all else being equal" because soil and ecosystem heat flux and net radiation of course changes as well with a change in vegetation.

We rewrote this paragraph completely and added the condition as suggested in the new paragraph. See also #6.

6. The paragraph beginning line 105 is a bit vague and out of place. It could be instead integrated into the previous paragraphs to provide context.

We rewrote this paragraph completely and integrated it with the previous paragraph. See also #5. The restructured paragraph is:

“The replacement of natural vegetation also changes evapotranspiration (ET) (Boisier et al., 2014) and LST because the surface biophysical variables (i.e. surface albedo, LST, emissivity and indirectly Leaf Area Index (LAI) and Normalized Difference Vegetation Index (NDVI)) are interconnected through the surface radiation balance. When ET decreases for example, surface temperatures and sensible heat (H) fluxes increase; on the other hand, when ET increases, the increased LE fluxes lower surface temperatures and decrease H fluxes (Mahmood et al., 2014) under equal net radiation conditions because with a change in vegetation, soil and ecosystem heat flux and net radiation also change due to an alteration of the biophysical variables. Vegetation structure, represented by NDVI, LAI and vegetation height, is in this respect an important determinant of the resistances or conductivities to heat, moisture, and momentum transfer between the canopy and the atmosphere (Bright et al., 2015) facilitating the amounts/ratios of sensible heat to water vapour dissipation away from the surface (Hoffmann and Jackson, 2000).”

7. Equation 2 should have subscripts.

8. On 302 the degree symbol is not necessary before K

We thank the associate editor for pointing to these details. We added the subscripts (comment #7) and removed the degree symbol before K (comment #8).

9. Equation 5 would be more accurate as sky temperature rather than near surface air temperature. That being said in this case the difference wouldn't be very large.

We changed the term to ‘sky temperature’. The term sky temperature is more appropriate because we used the temperature measured 2 m above the ground.

10. Please use the multiplication sign rather than the dot (which can mean dot product) in equation 6 and elsewhere and please also use standard parentheses rather than curly braces.

We thank the associate editor for pointing to this detail. We changed the equations as suggested.

11. 415: relatively hot and cool.

We changed the description of the images as suggested.

12. I find 4 significant digits for the values starting line 440 to be a bit too accurate given that these are from remote sensing platforms. 3 significant digits is better but still probably a bit generous.

We applied this suggestion and use now 3 significant digits. We applied this to other parts in the manuscript too.

13. capital 'Atmosphere' on line 603.

We corrected this.

14. 816 could be more specific.

We removed the general statement of this first sentence of the conclusion, rewrote it added specific information related to the study area. We rewrote the first sentence of the conclusion as:

“In summary, we studied the effects of land use and land cover changes on the surface biophysical variables in Jambi and explained the underlying mechanisms of the surface temperature regulation.”

I. R. Paterson (russell.paterson@deb.uminho.pt)

1. I am unclear how the data in Paterson et al. (2015) supported LULC change in Indonesia as was stated in the manuscript. The authors should clarify.

Line 718 (in the discussion): Paterson et al. (2015) mention that the LST in Malaysia increased over the last four decades of 2.7 – 4.0 degrees C per 100 years. This is an example of LST change in another region close to Indonesia. However, this is not from first hand observation, data or research but results from the Malaysian Meteorological Department. We admit that this reference is misplaced here and removed the reference to Paterson et al. (2015) from the discussion.

2. They could also write LULC in full at first with the abbreviation in brackets afterwards.

We realize that the abbreviation LULC is introduced in line 600 and that it was not defined anywhere else. We added the full words followed by the abbreviation in line 600 as “LST patterns across different land use and land cover (LULC) types.

3. The English could be tightened up in parts of the paper.

a. E.g. they start sentences with "We" on lines 31 and 34. Also, this is not the passive voice. Similarly, "Our" lines 40 and 43.

We have tried to improve the English in the manuscript, removing the repeated words in the same paragraph, such as the “we” or “our” in the indicated lines. We have especially tightened the abstract, but kept it in active voice, as the Biogeosciences’ guidelines do not explicitly mention the preferred style of writing.

b. Line 57: "in the past decades" is imprecise and clumsy.

We changed this to “in the past two and a half decades” to refer to our study period 2000 - 2015

c. Lines 66-69: Three "and"s are used in this sentence. Not the best English.

The biophysical variables (albedo, emissivity and surface roughness) affect gas and energy exchange processes between the land surface and the atmosphere. We did not find any obsolete “and” in this construction.

d. Line 75: Replace "rise" with "increase".

We changed the verb as suggested.

e. Line 94-96: You should not have to explain what the "vice versa" is.

We removed the “vice versa” and connected the remaining sentence to the previous. We wanted to describe both situations of ET increase and ET decrease: what happens when ET increases and when ET decreases. The rephrased sentence is now: “In case ET is decreased, surface temperatures and fluxes of sensible heat (H) increase. On the other hand, when ET increases, increased LE fluxes lower surface temperatures and decrease H fluxes (Mahmood et al., 2014).”

f. Anyway, I will stop there as its too much work to point out other such smallish things. Maybe you consider these unimportant...

We thank Dr. Paterson for his comments and suggestions for improvements. We are happy to incorporate the suggested language corrections. Additionally, we reread the entire manuscript and made further English correction.

II. A. RIVAL (alain.rival@cirad.fr)

1. Given the nature, impact and extend of results presented the paper submitted by Sabajo et al, the present title of article clearly appears as unapropriate. Indeed results do not show any pivotal role for oil palm cultivation which should support the assumption that it is a key driver of phenomena observed. The title should stick only to facts and findings, which tend to evidence a difference between native forest and cultivated land in terms of land surface temperature.

We agree with Dr. Rival that the title is too general. Our study shows results from Jambi province/Indonesia only, thus we now mention this specifically in the title. We however think that the evidence from our study is sufficient to link the observed increase of land surface temperature to the expansion of oil palm and other cash crops as the observed temperature increase at provincial level is in line with the observed temperature differences across land use type and the expansion of oil palm and cash crops over the last two and a half decades in the Jambi province. The area cultivated with oil palm grew faster than the area cultivated with rubber plantations between 1990 and 2011 (Clough et al. 2016). The title of our article is now: "Expansion of oil palm and other cash crops cause an increase of the surface temperature of the Jambi province in Indonesia"

III. Anonymous Referee #1

General: The authors investigate the effect of land cover change (from forests to 'other' and mostly oil palm plantations) on regional land surface temperatures. They use remote sensing to determine LST (and albedo, NDVI, and ET). They conclude that conversion from forests had led to a ~ 1 degree C temperature (positive) change after accounting for albedo. They also conclude that this is a positive feedback to climate warming.

1. I only suggest some minor edits and (if the authors can) and expansion of the discussion of what these LST changes might translate to in the atmosphere?

The reviewer brings up an important issue of land surface – atmosphere feedback. A recent study by Tölle et al. (2016)[#] showed for SE Asia as a whole that land use change at large scale impacts the boundary layer structure, cloud-cover regime and other aspects of local and regional weather and climate. Particularly, land clearings can amplify the response to climatic extreme events such as El Nino Southern Oscillation. Analyzing this kind of effects require however a regional climate model, which is beyond the scope of our study. We now discuss this aspect in the manuscript and added the following sentence:

Line 773: "A recent study by Tölle et al. (2017) showed for SE Asia that land use change at large scale may increase not only surface temperature but also impact other aspects of local and regional weather and climate occurring also in regimes remote from the original landscape disturbance. Land clearings can amplify the response to climatic extreme events such as El Nino Southern Oscillation."

[#]Tölle, M. H., Engler, S., and Panitz, H. 2017: Impact of Abrupt Land Cover Changes by Tropical Deforestation on Southeast Asian Climate and Agriculture. *Journal of Climate*, **30(7)**, 2587 – 2600, doi: 10.1175/JCLI-D-16-0131.1.

2. How much larger of a region will they affect? How would you determine this?

This study focuses on local and effects at the provincial level. Estimating or predicting the effects at a larger regional scale also requires a regional climate model as used in the study of

Tölle et al. (2017). Tölle et al. (2017) show that the effects of land use changes occur in remote regions other than where the land use changes occur. These effects are caused by the impacts the land cover change has on El Niño/La Niña episodes thereby enhancing wetter conditions in other regions, whereas in other regions wetter conditions decrease.

Introduction: nicely written and I appreciate the well thought out definitions.

3. Line 96: missing an “as” after “such”.

We added the missing word as suggested.

4. Methods: Could you describe the study sites in a little more detail (rather than the reference Drescher).

We added the following information:

*L. 186: “Previously logged rainforests in the Jambi province have been converted into intensively managed agro-industrial production zones as well as into smallholder farms to grow cash crop trees of rubber (*Hevea brasiliensis*) and oil palm (*Elaeis guineensis*) or fast-growing tree species such as *Acacia mangium* for pulp production (Drescher et al., 2016). The area cultivated with oil palm grew faster than the area cultivated with rubber plantations between 1990 and 2011 (Clough et al. 2016).”*

5. ET calculations: I’m familiar with the use of satellite data for all of the variables except for ET. Did you compare ET with the tower sites? How well does it work?

I see that you added this to the supplement, but it would be nice to have a validation of this method explained in the main text.

We have ET and LE estimates from eddy covariance measurements for two oil palm plantations in Jambi Province (young and mature oil palm plantation). Our SEBAL based LE estimates are within the variability range of LE measured from eddy covariance under similar meteorological conditions.

We added the following text: “The SEBAL based LE estimates are within the variability range of LE measurements using the eddy covariance technique under similar meteorological conditions (see SI).”

6. Results: Line 405-406: Hot = red? And cool = Blue colors. Can you please specify this?

In our description of the figure we added the matching colors as suggested. Line 405 – 406: “the hot areas (red) correspond to the known clear-cut areas, urban areas or other sparsely vegetated areas, the cooler areas (blue) correspond to vegetated areas such as forest, plantation forests and mature oil palm plantations.”

7. Discussion: Line 668: When I look at the figures, there also seems to be a high correlation between NDVI and ET (simply because the response pattern, the pattern of the changes, look very similar). Can you explain this? Is it because of the ET calculation?

Yes, ET and NDVI are highly correlated on one hand because the NDVI is used in the calculation of ET. On the other hand, another input for ET is LST, which is calculated from the raw thermal band (L6). L6 and NDVI are also highly correlated ($r = -0.87$) (see table 2, Line 494 – 502) even though NDVI and L6 are derived/measured from independent satellite bands. Thus, it come as no surprise that there is a correlation between NDVI and ET.

8. Line 763: “concurrent to” should be “concurrent with”

We changed the sentence with the correct prepositions as suggested.

9. Line 768: “governmental” should be “government”

We changed this and used the correct word.

Final remarks: This is a well-written, well-organized manuscript. I support publication in Biogeosciences.

We thank the anonymous referee for reviewing the manuscript and for the suggestions to improve the manuscript.

IV. Anonymous Referee #2

General comments: Sabajo et al. evaluates the impact of land use changes on land surface temperatures in Indonesia over the MODIS timespan (1999-2015). The study is well written and provides a good, long-term observational analysis clearly showing the impact of regional deforestation on increasing land surface temperature across an entire region.

We thank the anonymous referee for reviewing the manuscript and for the suggestions to improve the manuscript.

1. The only general comment I have is that it would be good to include a **seasonality analysis** showing how deforestation has changed land surface temperature in both wet and dry season. I know that satellite remote sensing is more challenging during the wet season, but I think evaluating the impact of land changes with seasonality would be useful. This could also highlight likely reductions in ET with land change (and shallower rooting zones) during the dry season. The dry season is also when heat impacts (including wildfires) could be more significant.

We agree that a seasonality analysis might show differences between the wet and dry season. We now made a seasonality analysis. Overall, the relationships in the dry season are stronger than for the wet season as we have much more usable data during the dry season. We found significant differences between LST of the dry and wet season. At 10:30 am the LST increased 0.09 ± 0.02 °C per year during the dry season, while the increase during the wet season was lower (0.06 ± 0.02 °C per year) (Fig. S10.1). Around 1:30 pm the LST increased 0.08 ± 0.03 °C per year, against 0.03 ± 0.02 °C increase per year during the wet season. At 10:30 pm the LST increased 0.03 ± 0.01 °C per year in the dry season, compared to a LST increase of 0.02 ± 0.01 °C in the wet season. At 1:30 am, the LST increased 0.05 ± 0.02 °C in the dry season, while the LST during the wet season increased 0.05 ± 0.03 °C. The increase of the LST at 1:30 pm, 10:30 pm and 1:30 am in the wet season was not significant ($p = 0.12$, $p = 0.06$ and $p = 0.11$, respectively). The significant increase of the LST during the dry season at all 4 times of observations suggests that the warming is more pronounced during the dry season compared to the wet season, which is reasonable as we have more incoming radiation during the dry season. Nevertheless, we prefer to pool the data from the dry and the wet season in order to get more statistically robust relationships.

In our analysis of the MODIS LST data we have not come across anomalous LST that could be attributed to forest fires. This is caused by the mask we applied in selecting the best quality pixels which mostly also removed pixels covered by smoke. A seasonality analysis is not possible with Landsat data because there is not enough data.

We added the following sentence to the manuscript (line 755):

“We like to point out that our MODIS analysis has a larger proportion of data from the dry season compared from the wet season, as there were more cloud free conditions during the dry season. Thus, our reported warming effect reflects cloud free conditions. During cloudy conditions, particularly in the wet season, the warming effect is expected to be lower.”

We also added the seasonal analysis to the supplementary information (S10).

We thank the anonymous referee for reviewing the manuscript and for the suggestions to improve the manuscript.





Fig. S10.1 (from manuscript, extended): Wet and dry season are separated.

Wet season: All months except June – September/October; Dry season: June – September/October (Meijide et al., 2017 & Drescher et al., 2016).

This figure has been added to the Supplementary Information (S10)

2. Specific comments: For Figure 1 (and text in the manuscript related to Figure 1), it might be good to describe the general atmospheric circulation for wet and dry seasons (are winds from the east or from the west). This would help the reader evaluate whether there are substantial land use changes upwind of the forest plots that are used as the baseline “control” to evaluate land surface temperature changes to due land use changes and not overall global climate change. *We include a wind rose from one of our reference meteorological stations in the area, (see Drescher et al., 2016), for data collected between October 2013 to May 2016. Based on the climate diagram for the region (obtained from data from 1991-2011) we considered as the dry season the months of June-September and the rest was considered as wet season. See the wind roses for the dry (left) and wet (right) seasons below:*



During the dry season counts by winds were predominantly from the SE, whereas during the wet season winds were predominantly from the NW. The SE vs. NW shift in wind directions is in line with the regional monsoonal circulation. The landscape in the lowland of Jambi province is, however, very patchy with small-scale mosaics of different land uses. While we cannot fully rule out that advection from upwind land use changes may play a role, but it seems unlikely to have a systematic bias given the typical patchiness of the landscape. Also warm air advection would mean that the “climate change” warming of the forested “control” site is overestimated, thus making the land-use change effect even larger.

V. List of major changes

1. We have changed the title to: “*Expansion of oil palm and other cash crops cause an increase of the surface temperature of the Jambi province in Indonesia*”. The first title was too general, the results apply to the Jambi province and not to the whole of Indonesia.
2. We removed the reference of Paterson et al. (2015). After re-reading we concluded that this reference was not correctly cited and misplaced.
3. We added a new reference of Tölle et al. (2017). This reference was a new publication that complemented our results with a modelling approach.
4. We added 1 section to the supporting information (S10). S10 contains a seasonality analysis as suggested by anonymous referee #2. We also add a short sentence in the discussion and refer to the S10 for the results of the analysis.
5. Equations 9 – 11 have been renumbered from 10 – 12, due to a mistake in the equation 9 (which was by accident numbered as 1, while in fact that had to be equation 9).

6. We changed figure 5: we only adjusted the legend and paid attention to the rounding of the numbers in the equations.

1 **Expansion of oil palm and other cash crops causes an increase of the land surface**
2 **temperature of in the the Jambi province in Indonesia**

3
4 Clifton R. Sabajo^{1,2†}, Gueric le Maire³, Tania June⁴, Ana Meijide¹, Olivier Roupsard^{3,5},
5 Alexander Knohl^{1,6}

6
7 ¹ University of Goettingen, Bioclimatology, 37077 Göttingen, Germany

8 ² AgroParisTech – Centre de Montpellier, Agropolis International, 648 rue Jean-François
9 Breton, 34093 Montpellier, France

10 ³ CIRAD, UMR Eco&Sols, F-34398 Montpellier, France

11 ⁴ Agrometeorology Laboratory Department of Geophysics and Meteorology,

12 Faculty of Mathematics and Natural Sciences, Bogor Agricultural University (IPB), Indonesia

13 ⁵ CATIE (Centro Agronómico Tropical de Investigación y Enseñanza / Tropical Agriculture
14 Centre for Research and Higher Education), 7170 Turrialba, Costa Rica

15 ⁶ University of Goettingen, Centre of Biodiversity and Sustainable Land Use (CBL), 37073
16 Goettingen, Germany

17
18 † Correspondence: Clifton R. Sabajo, University of Goettingen, Bioclimatology, Büsgenweg 2,
19 37077 Göttingen, Germany. E-mail: csabajo@uni-goettingen.de

20 Telephone: +49 (0) 551 39 12114

21
22
23 **Abstract**

24
25 Indonesia is currently one of the regions with the highest transformation rate of the land surface
26 worldwide due-related to the expansion of oil palm plantations and other cash crops replacing
27 forests on large scales. Land cover changes, which modify land surface properties, have a direct

28 effect on the land surface temperature (LST), a key driver for many ecological functions.
29 Despite the large historic land transformation in Indonesia toward oil palm and other cash crops
30 and governmental plans for future expansion, this is the first study so far to quantify the impacts
31 of land transformation on the LST in Indonesia ~~on LST~~. We analyse LST from the thermal band
32 of a Landsat image and produce a high-resolution surface temperature map (30m) for the
33 lowlands of the Jambi province in Sumatra (Indonesia), a region which suffered large land
34 transformation towards oil palm and other cash crops over the past decades. The comparison of
35 LST, albedo, Normalized Differenced Vegetation Index (NDVI), and evapotranspiration (ET)
36 between seven different land cover types (forest, urban areas, clear cut land, young and mature
37 oil palm plantations, acacia and rubber plantations) shows that forests have lower surface
38 temperatures than the otherse land cover types, indicating a local warming effect after forest
39 conversion. LST differences were up to $10.09-1 \pm 2.6$ °C (mean \pm SD) between forest and clear-
40 cut land. The differences in surface temperatures are explained by an evaporative cooling effect,
41 which offsets the albedo warming effect. Our analysis of the LST trend of the past 16 years
42 based on MODIS data, shows that the average daytime surface temperature ~~of in~~ the Jambi
43 province increased by 1.05 °C, which followed the trend of observed land cover changes and
44 exceed the effects of climate warming. This study provides evidence that the expansion of oil
45 palm plantations and other cash crops leads to changes in biophysical variables, warming the
46 land surface and thus enhancing the increase of the in-air temperature ~~due to~~because of climate
47 change.

48

49

50 *Keywords:* Land surface temperature, albedo, NDVI, evapotranspiration, biophysical variables,
51 oil palm, remote sensing, Landsat, MODIS, Indonesia, land-use / land cover change

52

53

54 **1 Introduction**

55

56 Indonesia is one of the regions where the expansion of cash crop monocultures such as acacia
57 (timber plantations), rubber, oil palm plantations and smallholder agriculture has drastically
58 reduced the area of primary forest in the last two and a half decades (Bridhikitti and Overcamp,
59 2012; Drescher et al., 2016; Marlier et al., 2015; Miettinen et al., 2012; Verstraeten et al., 2005).

60 This large scale conversion of rainforest for agricultural use has been observed on the island of
61 Sumatra, which has experienced the highest primary rainforest cover loss in all of Indonesia
62 (Drescher et al., 2016; Margono et al., 2012; Miettinen et al., 2011). Forest cover in the
63 Sumatran provinces of Riau, North Sumatra and Jambi, declined from 93 to 38% of provincial
64 area between 1977 and 2009 (Miettinen et al., 2012). These large scale transformations,
65 observed as land cover change, and land-use intensification have led to substantial losses in
66 animal and plant diversity, ~~and~~ ecosystem functions and changed microclimatic conditions
67 (Clough et al., 2016; Dislich et al., 2016; Drescher et al., 2016). Additionally, these changes
68 directly alter vegetation cover and structure ~~ands well as~~ land surface properties such as albedo,
69 emissivity, and surface roughness which affect gas and energy exchange processes between the
70 land surface and the atmosphere (Bright et al., 2015).

71

72 Replacing natural vegetation with another land cover modifies the surface albedo, which affects
73 the amount of solar radiation that is absorbed or reflected and consequently alters net radiation
74 and local surface energy balance. A lower or higher albedo results in a smaller or greater
75 reflection of shortwave radiation. As a result, the higher or lower amounts of net radiation
76 absorption may increase or decrease the surface temperature and change evapotranspiration
77 (Mahmood et al., 2014).

78

79 Changes in land cover also alter surface emissivity, i.e. the ratio of radiation emitted from a
80 surface to the radiation emitted from an ideal black body at the same temperature following the
81 Stefan–Boltzmann law. Emissivity of vegetated surfaces varies with plant species, density,
82 growth stage, water content and surface roughness (Snyder et al., 1998; Weng et al., 2004). A
83 change of emissivity affects the net radiation because it determines the emission of longwave
84 radiation that contributes to radiative cooling (Mahmood et al., 2014).

85

86 Water availability, surface type, soil humidity, local atmospheric and surface conditions affect
87 the energy partitioning into latent (LE), sensible (H) and ground heat (G) fluxes (Mildrexler et
88 al., 2011). Surface roughness affects the transferred sensible and latent heat by regulating
89 vertical mixing of air in the surface layer (van Leeuwen et al., 2011) thereby regulating land
90 surface temperature (LST). Through its association with microclimate, net radiation and energy
91 exchange (Coll et al., 2009; Sobrino et al., 2006; Voogt and Oke, 1998; Weng, 2009; Zhou and
92 Wang, 2011), LST is a major land surface parameter and as climatic factor it is regarded a main
93 driver of diversity gradients related to the positive relationships between temperature and
94 species richness (Wang et al., 2016).~~that also influences habitat quality and thus the distribution~~
95 ~~of plants and animals and biodiversity.~~

96

97 The replacement of natural vegetation also changes evapotranspiration (ET) (Boisier et al.,
98 2014) and LST because the surface biophysical variables (i.e. surface albedo, LST, emissivity
99 and indirectly Leaf Area Index (LAI) and Normalized Difference Vegetation Index (NDVI))
100 are interconnected through the surface radiation balance. When ET decreases, surface
101 temperatures and fluxes of sensible heat (H) increase. On the other hand, when ET increases,
102 the increased LE fluxes lower surface temperatures and decrease H fluxes (Mahmood et al.,
103 2014). When ET decreases for example, surface temperatures and sensible heat (H) fluxes
104 increase; on the other hand, when ET increases, the increased LE fluxes lower surface

105 temperatures and decrease H fluxes (Mahmood et al., 2014) under equal net radiation conditions
106 because with a change in vegetation, soil and ecosystem heat flux and net radiation also change
107 due to an alteration of the biophysical variables. Vegetation structure, represented by ~~as~~
108 ~~reflected by parameters such as the Normalized Difference Vegetation Index (NDVI)~~NDVI,
109 ~~Leaf Area Index (LAI)~~LAI and vegetation height, is in this respect an important determinant
110 of the resistances or conductivities to heat, moisture, and momentum transfer between the
111 canopy and the atmosphere (Bright et al., 2015) facilitating the amounts/ratios of sensible heat
112 to water vapour dissipation away from the surface (Hoffmann and Jackson, 2000).

113

114 ~~Surface albedo, surface temperature, surface emissivity, and indirectly LAI and NDVI are~~
115 ~~interconnected through the surface radiation balance. When the land surface is changed,~~
116 ~~feedback mechanisms involving these biophysical variables control the radiation balance and~~
117 ~~the surface temperature.~~

118 To understand the effects of land cover changes on LST, the associated biophysical variables
119 must be evaluated. This can be done through the surface radiation budget and energy
120 partitioning which unites these biophysical variables directly or indirectly: albedo as direct
121 determinant of the net solar radiation, NDVI as a vegetation parameter determining the
122 emissivity, which in turn determines the amount of reflected and emitted longwave radiation,
123 LST directly affecting the amount of emitted longwave radiation from the surface and ET,
124 which affects the amount of energy that is used for surface cooling via evaporating of water.

125

126 The effect of land cover change on LST is dependent on the scale, location, direction and type
127 of the change (Longobardi et al., 2016). Several studies showed an LST increase ~~of the LST~~
128 after forest conversion to built-up areas, ~~and~~ agricultural land (Zhou and Wang, 2011), ~~and to~~
129 crop land and pasture lands (Peng et al., 2014) in China. Similar ~~findings observations~~ were
130 reported for South American ecosystems: low vegetation such as grasslands in Argentina were

131 warmer than tall tree vegetation (Nosetto et al., 2005). In Brazil, the surface temperature
132 increased after the conversion of natural Cerrado vegetation (a savanna ecosystem) into
133 crop/pasture (Loarie et al., 2011a). Similar effects were also ~~shown~~observed for other South
134 American biomes (Salazar et al., 2016). In a global analysis, Li et al. (2015) showed that the
135 cooling of forests is moderate at mid latitudes ~~and that~~but Northern boreal forests are ~~even~~
136 warmer, an indication that the effect of land cover change on LST varies with the location of
137 the land cover change (Longobardi et al., 2016). Similar studies on the Indonesian Islands are
138 lacking but ~~increases in~~ surface temperature increases are expected as an effect of ~~the expansion~~
139 ~~of~~ oil palm and cash crop land expansion in the recent decades.

140

141 Measuring LST changes ~~in LST~~ is critical for understanding the effects of land cover changes,
142 but challenging. LST can be monitored with LST products retrieved from thermal infrared
143 (TIR) remote sensing data e.g. the use of the thermal bands of the Moderate Resolution Imaging
144 Spectrometer (MODIS) onboard the Terra and Aqua satellite (Sobrino et al., 2008), the thermal
145 band of the Thematic Mapper (TM) onboard the LANDSAT-5 platform (Sobrino et al., 2004,
146 2008) or Enhanced Thematic Mapper (ETM+) onboard the LANDSAT-7 platform. The
147 advantage of MODIS data is the availability of readily processed products at high temporal
148 resolution (daily) at medium (250 – 500 m) to coarse (1000 – 5000 m) spatial resolution (~~1000~~
149 ~~– 5000 m~~) scale; MODIS LST product (MOD11A1/MYD11A1) for example is provided at a
150 daily temporal resolution with a spatial resolution of 1 km. Landsat data are provided at a higher
151 spatial resolution (30 m), but ~~its~~the temporal resolution is however limited to 16 days and the
152 retrieval of LST requires the correction of the satellite observed radiances for atmospheric
153 absorption and emission (Coll et al., 2009). Besides LST, the connected biophysical variables
154 of the energy and radiation budget can be derived from the visible and near-infrared (VIS-NIR)
155 bands of ~~either~~ MODIS or Landsat, making integrated monitoring of the biophysical variables
156 related to changing land surface possible. In Indonesia, a large proportion of the land use

157 changes is driven by smallholders (Dislich et al., 2016), thus a combination of Landsat (for a
158 fine spatial resolution) and MODIS (for temporal developments) seems desirable.

159

160 The modification of the physical land surface properties ~~of the land surface~~ influences climate
161 and local microclimatic conditions via biogeochemical and biophysical processes. Therefore,
162 given Indonesia's history of large scale agricultural land conversion and governmental plans to
163 substantially expand the oil palm production (Wicke et al., 2011), it is important to study the
164 effects of the expansion of cash crop areas on the biophysical environment, especially on LST
165 as a key land surface parameter. These effects have been poorly studied in this region and
166 according to our knowledge this is the first study to quantify the effects of land use change on
167 LST in Indonesia. We focus on the Jambi province ~~of Jambi~~ ~~(on Sumatra/Indonesia)~~ as it
168 experienced large land transformation towards oil palm and other cash crops such as rubber
169 plantations in the past and it may serve as an example of future changes in other regions.

170

171 Our main objective is to quantify the differences in LST across different land cover types and
172 to assess the impact of cash crop expansion on the surface temperature ~~of in the~~ Jambi province
173 ~~(on Sumatra / Indonesia)~~ in the past decades (2000 – 2015). With this study we aim to (1)
174 evaluate the use of Landsat and MODIS satellite data as sources for a reliable surface
175 temperature estimation ~~of the surface temperature~~ in a tropical region with limited satellite data
176 coverage by comparing the surface temperatures retrieved from both satellite sources to each
177 other and against ground observations, (2) to quantify the LST variability across different land
178 cover types and (3) to assess the long term effects of land transformation on the surface
179 temperature against the background of climatic changes and (4) to identify the mechanisms that
180 explain ~~changes of~~ the surface temperature changes caused by alterations of ~~through changes~~
181 ~~in other~~ biophysical variables. In this study we compare the surface temperatures of different
182 land cover types that replace forests (i.e. oil palm, rubber and acacia plantations, clear cut land

183 and urban areas) by using high resolution Landsat and medium resolution MODIS satellite data
184 and discuss the differences by taking into account other biophysical variables such as the
185 albedo, NDVI and evapotranspiration (ET).

186

187 **2 Materials and methods**

188

189 **2.1 Study area**


190

191 The study was carried out in the lowlands (approx. 25 000 km²) of the Jambi province (total
192 area 50 160 km²) on Sumatra, Indonesia, between latitudes 0°30'S and 2°30'S and longitudes
193 101°E and 104°30'E (Fig. 1). This region has undergone large land transformation towards oil
194 palm and rubber plantations s over the past decades and thus may serve as an example of expected
195 changes in other regions of Indonesia (Drescher et al., 2016). The area has a humid tropical
196 climate with a mean annual temperature of 26.7 ± 0.2 °C (1991 – 2011, annual mean \pm SD of
197 the annual mean), with little intra-annual variation. Mean annual precipitation was 2235 ± 381
198 mm and a dry season with less than 120 mm monthly precipitation usually occurred between
199 June and September (Drescher et al., 2016). Previously logged rainforests in the Jambi province
200 have been converted to intensively managed agro-industrial production zones as well as and into
201 smallholder farms to grow cash crops ~~tree~~-of rubber (*Hevea brasiliensis*) and oil palm (*Elaeis*
202 *guineensis*) or fast-growing tree species such as *Acacia mangium* for pulp production (Drescher
203 et al., 2016). The area cultivated with oil palm grew faster than the area cultivated with rubber
204 plantations between 1990 and 2011 (Clough et al., 2016).

205

206 For this study, we used two data sets of different plot sizes. For the first data set, we delineated
207 28 large plots (ranging from 4 to 84 km²) of 7 different land cover types (Forest (FO), Rubber
208 (RU), Acacia Plantation Forest (PF), Young oil palm plantation (YOP), Mature Oil Palm

209 Plantation (MOP), Urban area (UB) and Clear-Cut areas (CLC)) (Fig. 1). The delineation was
 210 based on visual interpretation in combination with field observations, information from field
 211 work, which were as carried out conducted between October – December 2013. The large size
 212 of the plots was necessary to make a comparison between MODIS and Landsat images (see
 213 section satellite data). For the second data set, we selected 49 smaller plots within and outside
 214 these 28 large plots (between 50 × 50 m and 1000 × 1000 m) (Fig. 1) which allowed us to
 215 increase the number of plots to use when analysing Landsat images. These small plots were
 216 used to extract the surface temperature (LST), Normalized Difference Vegetation Index
 217 (NDVI), albedo (α) and evapotranspiration (ET) from a high resolution Landsat satellite image
 218 (see section satellite data) for the 7 different land cover types of interest.



219
 220 **Fig. 1** Geographic location of the study area. Jambi province on the Sumatran Island of
 221 Indonesia (Figs. 1a and 1b). The background of the map (Fig. 1c) is a digital elevation model,
 222 showing that the plots are located in the lowlands of the Jambi province. The large rectangles
 223 are the 28 different land cover types (Forest, Young and Mature Oil palm, Rubber, Urban area,

224 Acacia Plantation Forest and Clear-Cut land), the small squares are the locations of the 49 small
225 plots of the 7 different land cover types. Abbreviations: CLC = Clear-cut land, UB = Urban
226 area, YOP = Young oil palm plantation, MOP = Mature Oil Palm plantation, PF = Acacia
227 plantation forest, RU = Rubber plantation, FO = Forest.

228

229 2.2 Meteorological data

230

231 Air temperature and relative air humidity were measured at four reference meteorological
232 stations located in open areas within the ~~study area-of study~~ (Drescher et al., 2016), with
233 thermohygrometers (type 1.1025.55.000, Thies Clima, Göttingen, Germany) placed at 2m
234 height. Measurements were ~~taken-recorded~~ every 15 s and then averaged and stored in a DL16
235 Pro data logger (Thies Clima, Göttingen, Germany) as 10 min mean, from February 2013 to
236 December 2015. We used the air temperature from the meteorological stations to compare to
237 MODIS air temperatures (MOD07_L2). The relative air humidity was used as an input
238 parameter for NASA's online atmospheric correction (ATCOR) parameter tool to derive
239 parameters to correct Landsat thermal band for atmospheric effects (see Satellite data). We also
240 used air temperature and relative humidity from two eddy covariance flux towers located in the
241 study area (Meijide et al., 2017): one in a young oil palm plantation (two years old, S
242 01°50.127', E 103°17.737'), and the other one in a mature oil palm plantation (twelve years old,
243 S 01°41.584', E 103°23.484'). At these flux towers, air temperature and relative humidity were
244 measured above the canopy respectively with the same instruments as in the reference
245 meteorological stations (see Meijide et al., (2017), for a description of methodology). ~~In At~~ the
246 flux tower located in the mature oil palm plantation, we also measured the surface canopy
247 temperature between August 2014 and December 2015, which was compared to MODIS LST
248 estimates from the same period. ~~Measurements-of e~~The canopy temperature was measured were
249 ~~performed~~ with two infrared sensors (IR100) connected to a data logger, (CR3000) both from
250 Campbell Scientific Inc. (Logan, USA). For a regional coverage we used ERA Interim daily air

251 temperature grids (<http://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/>; (Dee et
252 al., 2011) from 2000 – 2015 at 0.125 degrees resolution to study the annual air temperature
253 trend in this period.

254

255 **2.3 Satellite data**

256

257 A Landsat 7 ETM+ VIS/TIR 30 m resolution surface reflectance image with low cloud cover,
258 acquired at 10:13 hours (local time) on 19 June 2013 covering the lowland area of the Jambi
259 province (path 125, row 61) was used in this study. Like all Landsat 7 ETM+ images acquired
260 after 31 May 2003, the image we used was affected by a scan line error causing a data loss of
261 about 22% (http://landsat.usgs.gov/products_slcoffbackground.php). Most selected plots were
262 located in the center of the image and thus not affected by the data loss, e.g. the forest plots
263 located at the edges of the scan line error zone faced minimal data loss because they were large
264 enough.

265 We also downloaded the tile h28v09 of the MODIS Terra (MOD) and Aqua (MYD) daily 1km
266 Land Surface Temperature and Emissivity products (MOD11A1 and MYD11A1 Collection-5)
267 and MODIS 16-days 500 m Vegetation Indices NDVI/EVI product (MOD13A1 Collection-5)
268 from 05 March 2000 till 31 December 2015 for Terra data and from 8 July 2002 till 31
269 December 2015 for Aqua data. We downloaded other supporting satellite data such as the
270 MODIS Atmospheric Profile product (MOD07_L2) and the MODIS Geolocation product
271 (MOD03). All MODIS data were reprojected to WGS84, UTM zone 48 South ~~using~~with the
272 MODIS Reprojection Tool (MRT). The quality of the MODIS data was ~~checked~~examined
273 ~~using~~with the provided quality flags and only pixels with the highest quality flag were used in
274 the analysis.

275

276 2.4 Retrieval of biophysical variables from Landsat 7 ETM+ VIS/TIR images

277

278

- 279 • *NDVI*

280

281 NDVI was derived using-from the reflectances corrected for atmospheric effects in the red
282 (ρ_{RED} , band 3 Landsat 7 ETM+) and near infrared (ρ_{NIR} , band 4 Landsat 7 ETM+) bands,
283 with:

284

$$285 \quad NDVI = \frac{\rho_{NIR} - \rho_{RED}}{\rho_{NIR} + \rho_{RED}} \quad (1)$$

286

- 287 • *Surface albedo*

288

289 The surface albedo (α) was computed using-with the equation of Liang (2000) for estimating
290 broadband albedo from Landsat surface reflectance bands, with:

291

$$292 \quad \alpha = 0.3141 \rho_1 + 0.1607 \rho_3 + 0.369 \rho_4 + 0.1160 \rho_5 + 0.0456 \rho_7 - 0.0057 \quad (2)$$

293

294 where ρ_1 , ρ_3 , ρ_4 , ρ_5 and ρ_7 are the Landsat 7 ETM+ surface reflectance bands (corrected for
295 atmospheric effects).

296

- 297 • *Surface temperature (LST)*

298

299 LST was derived following the method proposed by Bastiaanssen (2000), Bastiaanssen et al.
300 (1998a), Coll et al. (2010) and Wukelic et al. (1989) for computing the surface temperature

301 from the thermal infrared band (TIR, band 6) of Landsat (Supporting information, S1). The
302 thermal infrared band (TIR, band 6) was first converted to thermal radiance (L6, W/m²/sr/μm)
303 and then to atmospherically corrected thermal radiance (Rc, W/m²/sr/μm) ~~following the~~
304 ~~methodas~~ described by Wukelic et al. (1989) and Coll et al. (2010), and ~~using-with~~ the
305 atmospheric parameters obtained on NASA's online Atmospheric Correction Calculator (Barsi
306 et al., 2003, 2005) (supporting information, S2). The surface temperature (LST, K) was
307 computed ~~withthrough~~ the following equation similar to the Planck equation, as in Coll et al.
308 (2010) and Wukelic et al. (1989):

309

$$310 \quad LST = \frac{k_2}{\ln\left(\frac{\epsilon_{NB} \cdot k_1}{Rc} + 1\right)} \quad (3)$$

311

312 where εNB is the emissivity of the surface obtained from the NDVI (Supporting information,
313 Table S1), k1 (= 666.09 mW/cm²/sr/μm) and k2 (= 1282.71 K) are sensor constants for
314 converting the thermal radiance obtained from band 6 of Landsat 7 to surface temperature.

315 The surface temperature derived from Landsat thermal band was compared with ~~a-the~~ MODIS
316 LST product that was acquired on the same day at 10:30 am local time. ~~For this, t~~The Landsat
317 LST image was first resampled to MODIS resolution to enable a pixel to pixel comparison,
318 followed by extracting the average LST of 7 land cover types ~~using-with~~ the data set containing
319 the large delineated plots (Fig. 1).

320

- 321 • *Evapotranspiration (ET)*

322

323 Based on the Surface Energy Balance Algorithm for Land (SEBAL) (Bastiaanssen, 2000;
324 Bastiaanssen et al., 1998a, 1998b) we estimated ET (mm/hr) from latent heat fluxes (LE, W/m²)

325 which were computed as the residual from sensible (H, W/m²) and ground (G, W/m²) heat
326 fluxes subtracted from net radiation (Rn, W/m²) as:

327

$$328 \quad LE = Rn - G - H \quad (4)$$

329

330 We calculated Rn as the sum of incoming shortwave and longwave radiation, minus the
331 reflected shortwave and longwave radiation and the emitted longwave radiation (equation 5).

332 The surface albedo, surface emissivity and surface temperature determine the amounts of
333 incoming and reflected radiation:

334

$$335 \quad Rn = (1 - \alpha) S_{d\downarrow} + \epsilon_a \sigma T_a^4 - (1 - \epsilon_0) \epsilon_a \sigma T_a^4 - \epsilon_0 \sigma LST^4 \quad (5)$$

336

337 Where $S_{d\downarrow}$ is the incoming shortwave solar radiation (W/m²) at the surface; α is the surface
338 albedo (equation 2); ϵ_0 is the surface emissivity (-); ϵ_a is the atmospheric emissivity (-); σ is the
339 Stephan-Boltzmann constant (5.67×10^{-8} W/m²/K⁴); LST is the surface temperature (K,
340 equation 3); T_a is the ~~near-surface-air~~sky temperature (K). The surface emissivity (ϵ_0) is derived
341 from the NDVI and is described in the supporting information (Table S1). The average
342 atmospheric emissivity (ϵ_a) is estimated with the model of Idso and Jackson, (1969):

343

$$344 \quad \epsilon_a = 1 - 0.26 \exp \left\{ \frac{-7.77 \times 10^{-4} (273.15 - T_a)^2}{1} \right\} \quad (6)$$

345

346
347 Ground heat fluxes (G, W/m²) were derived as a fraction of Rn from an empirical relationship
348 between LST, α , and NDVI (Bastiaanssen, 2000) as:

349

350 $G = R_n \times \frac{LST - 273.15}{\alpha} \times (0.0038\alpha + 0.0074\alpha^2) \times (1 - 0.98NDVI^4)$ (7)

351

352 In SEBAL Sensible heat flux (H , W/m^2) was calculated as:

353

354 $H = \rho C_p \frac{\Delta T}{r_{ah}} = \rho C_p \frac{a LST + b}{r_{ah}}$ (8)

355

356 Where ρ is the air density (1.16 kg/m^3); C_p is the specific heat of air at constant pressure (1004
 357 J/kg/K); r_{ah} is the aerodynamic resistance to heat transport (s m^{-1}); a and b are regression
 358 coefficients which are determined by a hot extreme pixel (where $LE = 0$ and H is maximum)
 359 and a cold extreme pixel (where $H = 0$ and LE is maximum). The aerodynamic resistance to
 360 heat transport, r_{ah} , is calculated through an iterative process with air temperature measured at 2
 361 m as input. SEBAL is described in Bastiaanssen (2000) and Bastiaanssen et al. (1998a, 1998b).
 362 The application of SEBAL in this research is briefly described in the supporting information
 363 (S3: ET from satellite images).

364

365 2.5 Local short term differences between different land cover types

366

367 From the created LST, NDVI, Albedo and ET images we extracted the average values of the
 368 different land cover classes. ~~For this we used with~~ the data set containing the small 49 delineated
 369 plots covering 7 different land cover types (Fig. 1). The average effect of land transformation,
 370 i.e. the change from forest to another non-forest land cover type, on the surface temperature
 371 was evaluated as (cf. Li et al. (2015)) :

372

373 $\Delta LST = LST_{\text{non-forest}} - LST_{\text{forest}}$ (9)

374

375 A negative ΔLST indicates a cooling effect and positive ΔLST indicates a warming effect of
376 the non-forest vegetation compared ~~to~~with forest. The same procedure was applied in
377 evaluating the effect of land transformation on the NDVI, albedo and ET.

378

379 **2.6 Effects of land cover change on the provincial surface temperature in the past decades**

380

381 To analyse the long-term effects on the provincial scale we used the MODIS daily LST time
382 series (MOD11A1 and MYD11A1) from 2000 – 2015. MOD11A1 provides LST for ~~two times~~
383 ~~of the day:~~ 10:30 am and 10:30 pm and we used the times series between 2000 and 2015.
384 MYD11A1 provides LST for 1:30 am and 1:30 pm and is available from 8 July 2002; we used
385 complete years in our analysis and therefore used the MYD11A1 time series from 2003 – 2015.
386 We calculated the mean annual LST at four different times of the day (10:30 am, 1:30 pm,
387 10:30 pm and 1:30 am) between 2000 and 2015 for the lowland of ~~the~~Jambi from the MODIS
388 daily LST time series (MOD11A1 and MYD11A1). ~~To do so~~First, (1) we calculated for each
389 pixel the average LST pixel value using only the best quality pixels for every year; (2) from
390 these pixels we made a composite image (n = 16, one for each year) for the province and (3)
391 from each composite image we calculated the mean annual lowland provincial temperature as
392 the average of all the pixels that are enclosed by a zone delineating the lowland of the Jambi
393 province. We performed the same analysis with the MODIS 16-day NDVI product (2000 –
394 2015) and the ERA daily temperature grid (2000 – 2015) to compare the annual trends of LST,
395 NDVI and air temperature of the province. The average provincial LST and NDVI were
396 compared ~~to~~with the mean LST and NDVI of a selected forest that remained undisturbed forest
397 during the 2000 – 2015 period.

398

399 2.7 Statistical analysis

400

401 For [a](#) comparison of the Landsat derived LST and the MODIS LST we analyzed the statistical
402 relationships with the coefficient of determination (R^2), the root mean square error (RMSE),
403 the mean absolute error (MAE) and the bias (Bias):

404

$$405 \text{ RMSE} = \sqrt{\frac{\sum_{i=1}^N (E_i - O_i)^2}{N}} \quad (10)$$

406

$$407 \text{ Bias} = \frac{\sum_{i=1}^N (E_i - O_i)}{N} \quad (11)$$

408

$$409 \text{ MAE} = \frac{\sum_{i=1}^N |E_i - O_i|}{N} \quad (12)$$

410

411 Where O_i is MODIS LST, E_i is the Landsat surface temperature, and N is the number of pixels
412 compared. Model type 2 linear regression was applied for fitting the relation between MODIS
413 LST and Landsat LST.

414 We tested the relation between the biophysical variables LST (or L6 and Rc, both as pre- or
415 intermediate products before obtaining LST), albedo (α), NDVI and ET with [a](#) correlation
416 analysis and a multiple linear regression was applied to analyse the effects of the biophysical
417 variables on the LST. We used the model: $\text{LST (or Rc or L6)} \sim \alpha + \text{NDVI} + \text{ET}$, and used R^2
418 and standardized β -coefficients to evaluate the strength of the biophysical variables in
419 predicting the LST.

420

421 3 Results

422

423 3.1 Landsat LST compared to MODIS LST


424

425 Landsat and MODIS images showed similar spatial LST patterns ~~of LST~~ (Fig. 2). In both
426 images the relatively hot areas (red) correspond to the known clear cut areas, urban areas or
427 other sparsely vegetated areas, the relatively cooler areas (blue) correspond to vegetated areas
428 such as forest, plantation forests and mature oil palm plantations. The coarse resolution scale
429 of MODIS (1000 m for LST) allows a large regional coverage of the study area but does not
430 allow to retrieve detailed information on small patches (smaller than 1 km²). On the other hand,
431 the Landsat 7 image allows a detailed study of patches that are small enough (as small as 30 x
432 30 m²), but is affected by the scan line error causing data loss at the edges of the image. In both
433 MODIS and Landsat images clouds and cloud shadows were removed and therefore lead to
434 data gaps in the images.

435

436

437



438
439


440

441 **Fig. 2** MODIS LST image (top) compared with Landsat LST image (bottom). Cloud cover and
 442 cloud shadow cover resulted in data gaps (No data). The difference in acquisition time between
 443 the images is 15 minutes. The square in the MODIS image is the area that is covered by the
 444 Landsat tile (path 125, row 61). Both satellite images were acquired on 19 June 2013.

445

446 Landsat derived LST correlated well with MODIS LST ($R^2 = 0.82$; $p < 0.001$; Fig. 3) with a
 447 RMSE of 1.83 °C. The 7 land cover types had distinctive LSTs and the observed differences

448 between these land cover types were consistent in both images. The non-vegetated surfaces
449 (Clear cut land (CLC) and Urban areas (UB)) had higher surface temperatures than the
450 vegetated surface types (FO, YOP, MOP, PF and RU). Clear cut land had the highest surface
451 temperature of all compared land cover types, followed by urban areas whereas the vegetated
452 land cover types had lower surface temperatures: $LST_{CLC} (39.74 \pm 2.01 \text{ } ^\circ\text{C}) > LST_{UB} (35.79 \pm 1.26 \text{ } ^\circ\text{C}) > LST_{YOP} (30.31 \pm 0.72 \text{ } ^\circ\text{C}) > LST_{PF} (30.25 \pm 0.67 \text{ } ^\circ\text{C}) > LST_{MOP} (28.29 \pm 0.75 \text{ } ^\circ\text{C}) > LST_{RU} (27.78 \pm 0.89 \text{ } ^\circ\text{C}) > LST_{FO} (27.57 \pm 1.41 \text{ } ^\circ\text{C})$ (Landsat LST, Fig.
453 3). The same trend was derived from the MODIS image but with higher surface temperatures,
454 except for CLC: $LST_{CLC} (37.67 \pm 1.75 \text{ } ^\circ\text{C}) > LST_{UB} (36.33 \pm 1.57 \text{ } ^\circ\text{C}) > LST_{YOP} (31.73 \pm 0.85 \text{ } ^\circ\text{C}) > LST_{MOP} (30.67 \pm 0.88 \text{ } ^\circ\text{C}) > LST_{PF} (29.92 \pm 0.93 \text{ } ^\circ\text{C}) > LST_{RU} (29.60 \pm 0.36 \text{ } ^\circ\text{C}) > LST_{FO} (29.21 \pm 0.40 \text{ } ^\circ\text{C})$ (MODIS LST, Fig. 3).



459

460 **Fig. 3** Average surface temperature (LST) and standard deviation (SD) of 7 land cover types

461 derived from [a](#) Landsat thermal image compared with the mean and SD of MODIS LST.

462 CLC = Clear cut land, UB = Urban areas, YOP = young oil palm plantation, PF = Acacia

463 Plantation Forest, MOP = Mature Oil palm plantation, FO = Forest, RU = Rubber plantation.

464 The dashed line is the theoretical 1:1 line, the solid lines are the Linear Model type 2 regression

465 line (black) and the confidence limits of the regression line (red). The Landsat and MODIS

466 images were acquired on 19 June 2013, Landsat at 10:13 am local time, MODIS at 10:30

467 am local time respectively. Landsat pixels (30 m) were resampled to MODIS pixel resolution

468 (926 m) to make a pixel to pixel comparison between the two sources possible. RMSE is the
469 root mean squared error, MAE is the mean absolute error.

470

471 **3.2 Local short term differences between different land cover types**

472

473 The Δ LST between RU, MOP, PF, YOP, UB and CLC land cover types and FO were all
474 positive, meaning that ~~the all~~ other land cover types were warmer than forests (Fig. 4a &
475 Supporting Information S4 and S5). RU and MOP were 0.4 ± 1.5 °C and 0.8 ± 1.2 °C warmer
476 than forest, respectively. PF and YOP were much warmer than forests (Δ LST_{PF-FO} = 2.3 ± 1.1
477 °C, Δ LST_{YOP-FO} = 6.0 ± 1.9 °C). The largest Δ LSTs were between forest and the non-vegetated
478 land cover types, i.e. UB (Δ LST = 8.5 ± 2.1 °C) and CLC (Δ LST = 10.9 ± 2.6 °C). The LST
479 differences were significant ($p < 0.05$, post-hoc Tukey's HSD test), except between RU and FO
480 ($p = 0.78$, post-hoc Tukey's HSD test (Supporting Information S6, Table S6.1 & table S6.2).

481

482 Similar differences were found for the Δ NDVI between forest and ~~the~~ other land covers (Fig.
483 4b). The negative Δ NDVI indicates that the non-forest land cover types had lower NDVI than
484 forest. Δ NDVI between FO and RU, MOP, PF and YOP were small (between -0.01 ± 0.02
485 (Δ NDVI_{MOP-FO}) and -0.12 ± 0.06 (Δ NDVI_{YOP-FO}). The largest Δ NDVIs were between forest
486 and the non-vegetated land cover types, i.e. UB and CLC (Δ NDVI = -0.42 ± 0.11 and -0.41
487 ± 0.08 , respectively). All Δ NDVIs were significant ($p < 0.05$, post-hoc Tukey's HSD test).


488

489 The difference in albedo (Δ Albedo) between forest and the other land covers was very small
490 (Fig. 4c), with Δ Albedo values between -0.03 ± 0.01 (Δ Albedo_{PF-FO}) and 0.03 ± 0.02
491 (Δ Albedo_{YOP-FO}). These differences were significant ($p < 0.05$, post-hoc Tukey's HSD test).
492 PF had a lower albedo than forest (Δ Albedo_{PF-FO} = -0.03 ± 0.01), while the other land cover
493 types had a higher albedo than forest.

494

495 All compared land covers had lower ET than forest. RU, MOP and PF had slightly lower ET
496 than FO ($\Delta ET_{RU-FO} = -0.03 \pm 0.04$, $\Delta ET_{MOP-FO} = -0.03 \pm 0.03$ mm/hr, $\Delta ET_{PF-FO} = -0.04 \pm$
497 0.03 mm/hr) (Fig. 4d). YOP, UB and CLC had much lower ET values than forests: ΔET_{YOP-FO}
498 $= -0.18 \pm 0.04$ mm/hr, $\Delta ET_{UB-FO} = -0.23 \pm 0.04$ mm/hr, $\Delta ET_{CLC-FO} = -0.26 \pm 0.06$ mm/hr).
499 The ΔET s were significant ($p < 0.05$, post-hoc Tukey's HSD test). The SEBAL based LE
500 estimates were within the variability range of LE measurements from eddy covariance
501 measurements under similar meteorological conditions (see SI 3).

502



503
 504 **Fig. 4** Differences (mean \pm SD) in surface temperature (Δ LST), normalized difference
 505 vegetation index (Δ NDVI), Albedo (Δ Albedo) and Evapotranspiration (Δ ET) between other
 506 land covers (RU, MOP, PF, YOP, UB and CLC) and forest (FO) in the Jambi province, derived
 507 from [athe](#) Landsat LST image acquired on 19 June 2013 at 10:13 am local time.

508
 509 Albedo had a weaker influence on the LST ($\rho = 0.25$, $p < 0.05$) (Table 2) than NDVI and ET.
 510 As the thermal radiance band (L6) and the atmospherically corrected thermal band (Rc) were
 511 the basis for the LST calculation, the high correlation between L6 and NDVI ($\rho = -0.87$, $p <$

512 0.05) and between L6 and ET ($\rho = -0.98$, $p < 0.05$) resulted in a high correlation between LST
 513 and NDVI ($\rho = -0.88$) and between LST and ET ($\rho = -0.98$). The analysis showed that albedo,
 514 NDVI and ET were all significant predictors of LST ($F_{(3, 41586)} = 1 \times 10^6$, $p < 0.05$). ET was the
 515 strongest predictor of LST (stand. $\beta = -1.11$, $p < 0.05$). Albedo (stand. $\beta = -0.19$, $p < 0.05$;
 516 ~~resp.~~) and NDVI (stand. $\beta = -0.19$, $p < 0.05$) were weaker predictors of LST.

517
 518 **Table 2** Statistical analysis between biophysical variables (albedo (α), NDVI and ET) and
 519 Spectral Radiance band (L6), corrected thermal band (Rc) and Landsat surface temperature
 520 (LST).

Model		ρ	R^2	β	Stand. β	Model fit (R^2)	F-statistics
L6 ~ α + NDVI + ET	α	0.26	0.05	-2.94	-0.19	0.99	F (3, 41586) = 1.10×106, ***
	NDVI	-0.87	0.10	0.23	0.11		
	ET	-0.98	1.13	-4.00	-1.16		
Rc ~ α + NDVI + ET	α	0.25	0.05	-4.88	-0.20	0.99	F (3, 41586) = 1.79×106, ***
	NDVI	-0.88	0.04	0.16	0.05		
	ET	-0.98	1.00	-6.21	-1.10		
LST ~ α + NDVI + ET	α	0.25	0.05	-34.01	-0.19	0.99	F(3, 41586) = 2.3×106, ***
	NDVI	-0.88	0.05	1.30	0.05		
	ET	-0.98	1.00	-43.53	-1.11		

521 ***: $p = 2 \times 10^{-16}$

522 LM: Multiple linear regression analysis between LST (or L6 or Rc) and 3 biophysical variables:
 523 Albedo (α), NDVI and ET. ρ = correlation coefficient; R^2 : R-squared of the components; β =
 524 regression coefficient of the component; stand. β = standardized β ; Model fit (R^2): overall model
 525 fit of the multiple linear regression. ~~The values in brackets are for the analysis between the~~
 526 ~~biophysical variables and the corrected thermal band (Rc).~~

527
 528 A separate analysis (Table S6.3, Supporting information S6) showed that ET was a strong
 529 predictor of LST for each land cover type in this study and that NDVI and albedo were minor
 530 predictors of LST.

531

532 3.3 Effects of land-use change on the provincial surface temperature in the past decades

533

534 The average annual LST of ~~the province~~Jambi was characterized by a fluctuating but increasing
535 trend during daytimes (Fig. 5a and 5b) between 2000 and 2015. The average morning LST
536 (10:30 am) increased by 0.07 °C per year ($R^2 = 0.59$; $p < 0.001$), the midday afternoon LST
537 (13:30 local time) increased by 0.13 °C per year ($R^2 = 0.35$; $p = 0.02$) between 2003 and 2015.
538 While the daytime LST showed a clear increase, the night and evening LST (10:30 pm and 1:30
539 am, Fig. 5c and 5d) trends ~~were small~~show~~ed~~ing a small decrease of -0.02 °C ($R^2 = 0.29$; $p =$
540 0.02) and -0.01 °C ($R^2 = 0.05$; $p = 0.504$) per year, respectively. The observed LST trends
541 resulted in a total LST increase of 1.05 °C and 1.56 °C in the morning (10:30 am) and afternoon
542 (1:30 pm) respectively and a total decrease of the ~~province~~-LST of 0.3 °C (10:30 pm) and 0.12
543 °C (1:30 am) at night over the period from 2000 to 2015 in Jambi.

544

545 ~~In order to~~To separate the effect of land use change from global climate warming, we used a site
546 constantly covered by forest over that period (from the forest sites we used in this study) as a
547 reference that was not directly affected by land cover changes. That site showed small~~less~~
548 changes in LST than the entire province: only the mean morning LST (10:30 am) had a
549 significant but small trend with an increase by of 0.03 °C per year ($R^2 = 0.21$, $p < 0.05$) resulting
550 in a total LST increase ~~of the province~~in Jambi of 0.45 °C between 2000 and 2015 (Fig. 5a).
551 This LST warming is much smaller than the overall warming at provincial level of 1.05 °C. The
552 LST time series at other times showed no significant trends: the mean afternoon LST (1:30 pm)
553 increased by with -0.05 °C per year ($R^2 = 0.01$, $p = 0.31$) (Fig. 5b), the night and evening LST
554 by with 0.01 °C per year (Fig. 5c and 5d, $p = 0.19$ and $p = 0.605$, respectively).


555

556 The mean annual NDVI ~~of in the province~~Jambi decreased by 0.002 per year, ~~which result~~ing
557 in a total NDVI decrease of 0.03 ($R^2 = 0.34$; $p = 0.01$; Fig. 5e). The NDVI of the forest showed

558 a small but not significant increase of 0.001 per year ($R^2 = 0.04$, $p = 0.23$) (Fig. 5e) fluctuating
 559 around an NDVI of 0.84.

560

561 The mean annual midday air temperature (at 1:00 pm, local time, Fig. 5f) and the mean annual
 562 night air temperature (at 1:00 am, local time) increased every year by 0.05 °C and 0.03 °C,
 563 respectively, resulting in a total air temperature increase of 0.75 °C ($R^2 = 0.66$, $p < 0.001$) and
 564 0.45 °C ($R^2 = 0.32$, $p = 0.014$) between 2000 and 2015 (Fig. 5f).



565

566 **Fig 5.** Mean annual LST (a – d), mean annual NDVI (e) and mean annual air temperature trends

567 (f) in the Jambi province between 2000 and 2015 derived from MODIS LST (5a. 10:30 am, 5b.

568 1:30 pm, 5c. 10:30 pm and 5d. 1:30 am, local time), MODIS NDVI and ERA Interim Daily air

569 temperature (1:00 am and 1:00 pm, local time) data sets respectively. Grey-shaded areas are the
570 confidence intervals of the means, blue-shaded areas are the confidence intervals of the
571 regression lines. MODIS LST time series for 1:30 pm and 1:30 am were available from the mid
572 of 2002; for this reason we used the complete years from 2003 till 2015.

573

574 **4 Discussion**

575

576 **4.1 Landsat LST compared to MODIS LST**

577

578 In our study we retrieved the surface temperature from a Landsat image and compared this with
579 MODIS LST. Our results showed a good agreement between both LSTs (Fig. 3), which is
580 comparable to other studies and thus gives confidence in our analysis. Bindhu et al. (2013)
581 found also a close relationship between MODIS LST and Landsat LST by using the same
582 aggregation resampling technique as our method and found a R^2 of 0.90, a slope of 0.90, and
583 an intercept of 25.8 °C for LST, compared to our R^2 of 0.8, slope of 1.35 and intercept of –
584 11.58 °C (Fig. 3). Zhang and He (2013) validated Landsat LST with MODIS LST and also
585 found good agreements (RMSD 0.71 – 1.87 °C) between the two sensors, whereas we found a
586 RMSE of 1.71 °C. Nevertheless, there still are differences and slope versatility between the two
587 satellite sources. These differences are typically caused by differences between the MODIS and
588 Landsat sensors in terms of (a) different sensor properties e.g. spatial and radiometric resolution
589 and sensor calibration; (b) geo-referencing and differences in atmospheric corrections (Li et al.,
590 2004); and (c) emissivity corrections i.e. the use of approximate equations to derive the
591 emissivity from the NDVI from Landsat's Red and NIR bands. Li et al. (2004) and Vlassova et
592 al. (2014) identified these same factors in their comparison of ASTER LST with MODIS LST
593 and Landsat LST with MODIS LST, respectively. Vlassova et al. (2014) found good
594 agreements between MODIS and Landsat LST and – obtain ed ing higher LST_s with MODIS

595 than with Landsat, which they attributed to the delay of 15 minutes in acquisition time between
596 MODIS and Landsat. MODIS LST is measured 15 minutes later and our results showed that
597 MODIS LSTs were indeed higher than Landsat LST. A comparison of MODIS LST with
598 locally measured canopy surface temperatures during the overpass time of MODIS also showed
599 agreement (Supporting information S7, Figure S7.1). The slope was possibly ~~due-related~~ to
600 differences in instrumentation and emissivity corrections and to scale issues, still this
601 comparison could corroborate the quality check of MODIS LST.

602 As the MODIS LST product is proven to be accurate within 1 °C (Silvério et al., 2015; Wan et
603 al., 2004) and has been intensively validated, the use of MODIS LST was a proper way to assess
604 the quality of our Landsat LST.

605

606 The errors from the different sources (such as atmospheric correction, emissivity correction,
607 resampling Landsat to MODIS resolution) are difficult to quantify. When we tested the impact
608 of atmospheric correction and emissivity errors on the LST from Landsat retrieval we found
609 that: (a) the overall patterns across different land use types did not change, (b) emissivity was
610 the most important factor, but the effects on LST retrieval were small and (c) errors ~~due-related~~
611 to atmospheric correction parameters were small because there were minor differences between
612 ~~the default Atmospheric-atmospheric~~ correction (ATCOR) parameters and ~~the~~ ATCOR
613 parameters derived with actual local conditions (relative humidity (RH), air pressure and air
614 temperature). Following the method of Coll et al. (2009) and Jiang et al. (2015) we show that
615 the use of the online atmospheric correction parameter calculator is a good option provided that
616 RH, air temperature and air pressure measurements are available. We additionally compared
617 locally measured air temperatures with MODIS air temperature and found a good agreement
618 (Supporting information S8, Figure S8.1), which served as a verification that we used a correct
619 air temperature for the atmospheric correction parameter calculator.

620 Overall, our comparison of ~~LST from~~ Landsat LST ~~with~~ ~~against~~ MODIS ~~LST from~~ ~~MODIS~~ and
621 against ground observations suggests that we are able to retrieve meaningful spatial and
622 temporal patterns of LST in the Jambi province.

623

624 **4.2 LST patterns across different land use and land cover (LULC) types**

625

626 The land cover types in our study covered a range of land surface types that develop after forest
627 conversion. This is the first study in this region that includes oil palm and rubber as land use
628 types that develop after forest conversion. The coolest temperatures were at the vegetated land
629 cover types while the warmest surface temperatures were on the non-vegetated surface types
630 like urban areas and bare land. Interestingly, the oil palm and rubber plantations were only
631 slightly warmer than the forests whereas the young oil palm plantations had clearly higher LST
632 than the other vegetated surfaces. For other parts of the world, Lim et al. (2005, 2008), Fall et
633 al. (2010) and Weng et al. (2004) also observed cooler temperatures for forests and the highest
634 surface temperatures for barren and urban areas.

635 In Indonesia, land transformation is often not instantaneous from forest to oil palm or rubber
636 plantation, but can be associated with several years of bare or abandoned land in-between (Sheil
637 et al., 2009). Oil palm plantations typically have a rotation cycle of 25 years, resulting in
638 repeating patterns with young plantations (Dislich et al., 2016). Given the large LST differences
639 ~~in LST~~ between forests and bare soils or young oil palm plantations that we observed, a
640 substantial warming effect of land transformation at regional scale is expected.

641

642 **4.3 Drivers of local differences between different land cover types**

643

644 All the land cover types (except Acacia Plantation Forests) had a higher albedo than forest,
645 indicating that these land cover types absorbed less incoming solar radiation than forests.

646 Nevertheless, these land cover types were warmer than forests, suggesting that the albedo was
647 not the dominant variable explaining the LST. Indeed, the statistical analysis showed that $ET \sim$
648 LST had a higher correlation than $albedo \sim LST$. The ΔET s were significant, underlying that
649 despite their higher albedo, all land cover types had higher LSTs than forests due-related to
650 lower ET rates than forests. ~~Vice-versa~~On the other hand, forests that absorb more solar
651 radiation ~~due-to~~because of the lower albedo, have lower LST ~~due-to~~because of the higher ET
652 they exhibit, hereby identifying evaporative cooling as the main determinant of regulating the
653 surface temperature of all vegetation cover types (Li et al., 2015).

654

655 Both observational and modeling studies carried out in other geographic regions and with other
656 trajectories support our observations. Observational studies in the Amazonia by Lawrence and
657 Vandecar (2015) on the conversion of natural vegetation to crop or pasture land showed a
658 surface warming effect. Salazar et al. (2015) provided additional evidence that conversion of
659 forest to other types of land use in the Amazonia caused significant reductions in precipitation
660 and increases in surface temperatures.

661 Alkama and Cescatti (2016) and earlier studies by Loarie et al. (2011a, 2011b) showed that
662 tropical deforestation may increase the LST. Croplands in the Amazonian regions were also
663 warmer than forests through the reduction of ET (Ban-Weiss et al., 2011; Feddema et al., 2005)
664 and that the climatic response strongly depends on changes in energy fluxes rather than on
665 albedo changes (Loarie et al., 2011a, 2011b). A study by Silvério et al. (2015) indeed found
666 that tropical deforestation changes the surface energy balance and water cycle and that the
667 magnitude of the change strongly depends on the land uses that follow deforestation. They
668 found that the LST was 6.4 °C higher over croplands and 4.3 °C higher over pasture lands
669 compared to the forest they replaced, as a consequence of energy balance shifts. Ban-Weiss et
670 al. (2011) and Davin and de Noblet-Ducoudré (2010) added that in addition to the reduction of
671 ET, the reduction of surface roughness most likely enhanced the substantial local warming.

672

673 Also for non-Amazonian regions, the replacement of forests by crops caused changes
674 comparable with our observations. In temperate Argentina, Houspanossian et al. (2013) found
675 that the replacement of dry forests by crops resulted in an increase of albedo ~~and~~ but still forests
676 exhibited cooler canopies than croplands. The cooler canopies were a result of a higher
677 aerodynamic conductance that enhanced the capacity of tree canopies to dissipate heat into the
678 atmosphere, and to both latent and sensible heat fluxes operating simultaneously to cool forest
679 canopies.

680

681 In a global analysis Li et al. (2015) showed that tropical forests generally have a low albedo,
682 but still the net energy gain caused by solar energy absorption is offset by a greater latent heat
683 loss via higher ET and that in the tropical forests the high ET cooling completely offsets the
684 albedo warming. For China, this cooling effect was also shown by Peng et al. (2014) who
685 compared LST, albedo and ET of plantation forests, grassland and cropland with forests.

686

687 For the USA, Weng et al. (2004) and for China, Yue et al. (2007), using NDVI as an indicator
688 of vegetation abundance, also found that areas with a high mean NDVI had a lower LST than
689 areas with a low mean NDVI, therefore suggesting that vegetation abundance is an important
690 factor in controlling the LST through higher ET rates. Our result support their assumptions by
691 showing the high correlation between NDVI – LST and ET – LST.

692

693 Our findings are also supported by modelling studies. Beltrán-Przekurat et al. (2012) found for
694 the Southern Amazon that conversion of wooded vegetation to soy bean plantations caused an
695 increase of the LST due to decreased latent heat and increased sensible heat fluxes. Climate
696 models also show the same warming trends and land surface modelling also projects an increase
697 in surface temperatures following deforestation in the Brazilian Cerrado (Beltrán-Przekurat et

698 al., 2012; Loarie et al., 2011b). In a global analysis, Pongratz et al. (2006) showed a LST
699 increase of forest to cropland or pasture transitions, which was driven by a reduced roughness
700 length and an increased aerodynamic resistance, and that the temperature response is intensified
701 in forest to clear/bare land transitions (1.2 – 1.7 °C increase). Similar to observational studies,
702 the modelling results of Bathiany et al. (2010) show that ET is the main driver of temperature
703 changes in tropical land areas.

704

705 ~~In order to~~To understand the effects of deforestation on biophysical variables in Indonesia, our
706 study identifies the following mechanisms: (a) reduction of ET decreases surface cooling, (b)
707 reduced surface roughness reduces air mixing in the surface layer and thus vertical heat fluxes,
708 (c) changes in albedo change the net radiation, and (d) changes in energy partitioning in sensible
709 and latent heat and heat storage. The effect is an increase of the mean temperatures leading to
710 warming effects in all tropical climatic zones (Alkama and Cescatti, 2016). We point here that
711 our study (1) included a ground heat flux, but did not take into account the storage of heat in
712 the soil and the release of stored heat out of the soil during the daily cycle and (2) that the
713 Landsat satellite image was obtained under cloud free conditions with high shortwave radiation
714 input and low fraction of diffuse radiation. Therefore, the LST retrieved on cloud free days
715 might be overestimated compared to cloudy days, as the differences in LST between land uses
716 are supposed to be lower when diffuse radiation increases.

717

718 Our study is the first to include the oil palm and rubber expansion in Indonesia. In Indonesia,
719 smallholders take 40% of the land under oil palm cultivation for their account (Dislich et al.,
720 2016). ~~Since Because~~ the landscape in ~~the~~Jambi ~~province~~ is characterized by a small-scale
721 smallholder-dominated mosaic including rubber and oil palm monocultures (Clough et al.,
722 2016), studies using medium to coarse resolution data are not able to capture the small scale
723 changes and processes at the small-scale level. By using high resolution Landsat data we were

724 ~~also~~ able to ~~also~~ include the effects of land use change on biophysical variables and the
725 underlying processes of the small scale holder agriculture.

727 **4.4 Effects of land use change on the provincial surface temperature in the past decades**

729 The increases ~~in-of the~~ mean surface temperature ~~of-thein~~ Jambi ~~province~~ were stronger during
730 the morning (10:30 am) and afternoon (1:30 pm) than during the evening (10:30 pm) and night
731 (1:30 am). Given that our results show a decrease of the NDVI in the same period, this suggests
732 that the observed increased trend of the day time ~~province~~-LST can be attributed to the land
733 cover changes that occurred. Our assumption that the observed decreasing NDVI trend is caused
734 by land conversions is supported by two different studies which reported that in ~~the~~-Jambi
735 ~~province~~, between 2000 and 2011 (Drescher et al., 2016) and between 2000 and 2013 (Clough
736 et al., 2016), the forest area decreased and that the largest increases were for rubber, oil palm,
737 and agricultural and tree crop areas. The class ‘other land use types’, which includes urban
738 areas, showed a minor increase (around 1%), suggesting that the decrease in NDVI was most
739 likely caused by forest cover loss and not by urban expansion (see Supporting information,
740 Table S9). The same observations on LULC change in Indonesia were also ~~supported-done~~ by
741 Lee et al. (2011), Margono et al. (2012, 2014) and Luskin et al. (2014). Luskin et al. (2014)
742 showed that in ~~the~~-Jambi-~~province~~, during the period 2000 – 2010, forests decreased by 17%
743 while oil palm and rubber area increased by 85% and 19%, respectively.

745 Given these trends in LULC changes, the observed LST trends were most likely caused by
746 gradual decrease of forest cover loss at the expense of agriculture and croplands. Our
747 assumptions are supported by findings of Silvério et al. (2015), Costa et al. (2007), Oliveira et
748 al. (2013), Spracklen et al. (2012) and Salazar et al. (2015) which indicate that land use
749 transitions in deforested areas likely have a strong influence on regional climate. Alkama and

750 Cescatti's (2016) ~~analysis~~ show that biophysical effects of forest cover changes ~~in forest cover~~
751 can substantially affect the local climate by altering the average temperature, which is consistent
752 with our observations and can be related to the observed land use change in the Jambi province.
753 As Indonesia has undergone high rates of forest cover loss from 2000 to 2012 (Margono et al.,
754 2014), these findings support our assumptions that the observed LST increase in the Jambi
755 province was most likely caused by the observed land use changes.

756

757 To separate the effect of global warming from land-use change induced warming, we
758 considered areas with permanent and large enough forests as a reference where changes are
759 mainly ~~due to~~ because of global warming. We find that LST of forests show either no significant
760 trends (at 1:30 pm, 10:30 pm, 1:30 am) or just a clearly smaller increase of 0.03 °C per year at
761 10:30 am. The difference between the LST trend of the province and of the forest at 10:30 am
762 was 0.04 °C per year, resulting in a Δ LST of 0.6 °C between the province and forest in the
763 period 2000 and 2015. We point out that our MODIS analysis has a larger proportion of data
764 from the dry season compared from the wet season, as there were more cloud free conditions
765 during the dry season. Thus, our reported warming effect reflects cloud free conditions. During
766 cloudy conditions, particularly in the wet season, the warming effect is expected to be lower. A
767 seasonality analysis showed that the relationships in the dry season are stronger than for the wet
768 season (see Supporting information S10, fig. S10.1) which suggests that the warming is more
769 pronounced during the dry season compared to the wet season, which is reasonable as we have
770 more incoming radiation during the dry season.

771

772 Using With the warming effects we found between forest and other land cover types (Δ LST,
773 Fig. 4a) and the observed land cover changes by Clough et al. (2016), Drescher et al. (2016)
774 (Supporting Information S9, table S9.1 and S9.2) we estimated the contribution of all land cover
775 types (except forest) to the Δ LST of the province between 2000 and 2015 to be 0.51°C out of

776 ~~the observed~~ 0.6°C ~~observed above~~, which also supports our assumption that the LST increase
777 ~~in Jambi of the province LST~~ was ~~by for~~ 85% driven by land cover changes (see Supporting
778 Information 9, Table S9.1 & S9.2: Land use change analysis), with clear cut areas having a
779 large contribution as they have the largest warming effect.

780

781 The observed small, but significant increase in LST of forests of 0.03 °C per year at 10:30 am
782 reflects a LST change independent ~~to of~~ land cover changes, as the forest remained unchanged
783 over that time period. A potential driver of that LST increase is the general global air
784 temperature trend ~~due to because of~~ changes in radiative forcing or border effects (advection
785 from warmer land uses), which is similar to the 1994 - 2014 time series analysis of Kayet et al.
786 (2016), who showed a LST increase for all land cover types ranging from wasted land,
787 agriculture land, open forest, dense forest, water bodies and built up areas.

788

789 The observed trends of ~~the province-provincial~~ air temperature (Fig. 5f) were significant,
790 suggesting that a general warming due to global and regional effects contributes to the observed
791 warming at ~~the province-provincial~~ level during day and night time, but that it is smaller than
792 the land cover change induced effects (Supporting Information S9, Table S9.1 & S9.2) at ~~the~~
793 provincial level (Fig. 5a and 5b).

794

795 In our long-term analysis on the regional effects of land use change we observed an increase in
796 the mean LST and mean air temperature in the 2000 - 2015 period, concurrent with a decrease
797 of the NDVI. The warming observed from MODIS LST data and from the air temperature
798 obtained from the independent ERA Interim Reanalysis in the Jambi province are most likely
799 caused by the observed decrease of the forest area and an increase ~~of~~ oil palm, rubber and other
800 cash crop areas in the same period, with other effects such as radiative forcing changes and
801 additional natural effects playing a smaller role. Given the plan of the Indonesian government

802 to substantially expand oil palm production with a projected additional demand of 1 to 28 Mha
803 in 2020 (Wicke et al., 2011), the strong warming effect we show for Jambi ~~province~~ may serve
804 as an indication of future LST changes ~~in LST~~ for other regions of Indonesia that will undergo
805 land transformations towards oil palm plantations.

806 A recent study by Tölle et al. (2017) showed that for Southeast Asia, land use change at large
807 scale may increase not only surface temperature but also may impact other aspects of local and
808 regional weather and climate occurring also in regions remote from the original landscape
809 disturbance. Their results also indicate that land clearings can amplify the response to climatic
810 extreme events such as El Niño Southern Oscillation (ENSO). The observed effects of land use
811 change on the biophysical variables may have implications for ecosystem services in the Jambi
812 province beyond a pure warming effect. The high precipitation in this region in combination
813 with the reduced vegetation cover of bare land and young oil palm plantations impose risks of
814 soil erosion caused by surface run off. Less water infiltrationes into the soil, thereby decreasing
815 the soil water storage ~~that~~ may lead to low water availability in the dry season (Dislich et al.,
816 2016; Merten et al., 2016). High surface temperatures in combination with low water
817 availability may make the vegetation and the surroundings more vulnerable to fires.

818

819 **5 Conclusion**

820

821 In summary, we studied the effects of land use and land cover changes on the surface
822 biophysical variables in Jambi and explained the underlying mechanisms of the surface
823 temperature regulation. showed the importance of forests in regulating the local and regional
824 climate. We derived biophysical variables from satellite data, analyzed the biophysical impacts
825 of deforestation and on a local scale we found a general warming effect after forests are
826 transformed to cash or tree croplands (oil palm, rubber, acacia) in the Jambi province of
827 Sumatra. The warming effect after forest conversion results from the reduced evaporative

828 cooling, which was identified as the main determinant of regulating the surface temperature.
829 On a regional scale, we saw that the effects of land cover changes are reflected back in changes
830 of the LST, NDVI and air temperature ~~of the in~~ Jambi ~~province~~. The warming effect induced by
831 land cover change clearly exceeded the global warming effect. Understanding the effects of
832 land cover change on the biophysical variables may support policies regarding conservation of
833 the existing forests, planning and expansion of the oil palm plantations and possible
834 afforestation measures.

835 **Supporting Information**

836

837 Supporting information to this article is arranged as follows:

838

839 **S1. Surface temperature retrieval from Landsat thermal images**

840 **Table S1.1.** Steps in the retrieval of the surface temperature from Landsat TIR band

841 **Table S1.2.** LMIN and LMAX values for Landsat 7 ETM+

842 **Table S1.3.** Mean solar exo-atmospheric irradiance ($ESUN_{\lambda}$) for Landsat 7 ETM+

843

844 **S2. Atmospheric correction of the thermal band**

845 **Table S2.1.** Input and output parameters for/from NASA's online atmospheric correction
846 parameter calculator

847

848 **S3. ET from satellite images with SEBAL**

849 **Fig. S3.1** Analysis of the steps involved in deriving the input for deriving ET from Landsat
850 images with SEBAL

851 **Fig. S3.2** Comparison of ET derived from upper anchor and lower anchor pixels.

852 **Table S3.1.** u^* , rah , LE and H measured at a young and mature oil palm plantation

853

854 **S4. Mean LST, NDVI, Albedo and NDVI extracted for 7 land cover types**

855 **Fig. S4.1** Mean LST, NDVI, Albedo and NDVI extracted from Landsat LST images for 7
856 land cover types

857

858 **S5. Difference in LST, NDVI, albedo and ET between Forest (FO) and 6 other land cover
859 types**

860 **Fig. S5.1** Differences in LST (ΔLST), NDVI ($\Delta NDVI$), Albedo ($\Delta Albedo$) and
861 Evapotranspiration (ΔET) between other land covers (RU, MOP, PF, YOP, UB and CLC) and
862 forest (FO) in the Jambi province

863

864 **S6. Statistical analysis**

865 **Table S6.1** ANOVA statistics

866 **Table S6.2** Post-hoc Tukey HSD test statistics

867 **Table S6.3** The relation LST-Albedo-NDVI-ET separated by land cover type

868

869 **S7. Comparison of MODIS LST to in situ measured canopy LST**

870 **Fig. S7.1** MODIS LST compared ~~to~~-with in situ measured canopy surface temperature.

871

872 **S8. Comparison of MODIS Air temperature with locally measured air temperature**

873 **Fig. S8.1** MODIS Air temperature compared with in situ measured air temperatures

874

875 **S9. Land use change analysis for the Jambi province for 2000 – 2010**

876 **Table S9.1** Land use change (1990) – 2000 – 2010

877 **Table S9.2** Contribution of land cover change to total LST increase

878

879 **S10. Seasonality analysis**

880 **Fig S10.1** Mean annual LST in the Jambi province between 2000 and 2015 derived from
881 MODIS LST during the wet and dry season.

882

883 *Author contributions.* Clifton R. Sabajo conducted the research, fieldwork an analysis and
884 prepared the manuscript, which was reviewed by Guerric le Maire, Tania June, Ana Meijide,
885 Olivier Roupsard and Alexander Knohl. Ana Meijide and Alexander Knohl provided the
886 meteorological data.

| *Competing interests.* The authors declare ~~that they~~o have no conflict of interest.

887 *Acknowledgements.* This research was funded by the Erasmus Mundus Joint Doctorate
888 Programme Forest and Nature for Society (EMJD FONASO) and the German Research
889 Foundation (DFG) through the CRC 990 “EForTS, Ecological and Socioeconomic Functions
890 of Tropical Lowland Rainforest Transformation Systems (Sumatra, Indonesia)” (subproject
891 A03). A special thanks to Huta Julu Bagus Putra, a.k.a. Monang, for his assistance and
892 translation during the field work in Indonesia. ~~The authors state to have no conflict of interest.~~

893 **References**

894

895 Alkama, R. and Cescatti, A.: Biophysical climate impacts of recent changes in global forest
896 cover, *Science*, 351(6273), 600–604, doi:10.1126/science.aac8083, 2016.

897 Ban-Weiss, G. A., Bala, G., Cao, L., Pongratz, J. and Caldeira, K.: Climate forcing and response
898 to idealized changes in surface latent and sensible heat, *Environ. Res. Lett.*, 6(3), 34032, 2011.

899 Barsi, J. A., Barker, J. L. and Schott, J. R.: An Atmospheric Correction Parameter Calculator
900 for a Single Thermal Band Earth-Sensing Instrument, *Geosci. Remote Sens. Symp. 2003*
901 *IGARSS 03 Proc. 2003 IEEE Int.*, 5, 3014–3016 vol.5, doi:10.1109/IGARSS.2003.1294665,
902 2003.

903 Barsi, J. A., Schott, J. R., Palluconi, F. D. and Hook, S. J.: Validation of a web-based
904 atmospheric correction tool for single thermal band instruments, in *Proc. SPIE, Earth Observing*
905 *Systems X*, vol. 5882, San Diego, California, USA., 2005.

906 Bastiaanssen, W. G. .: SEBAL-based sensible and latent heat fluxes in the irrigated Gediz
907 Basin, Turkey, *J. Hydrol.*, 229(1–2), 87–100, doi:10.1016/S0022-1694(99)00202-4, 2000.

908 Bastiaanssen, W. G. M., Menenti, M., Feddes, R. A. and Holtslag, A. A. M.: A remote sensing
909 surface energy balance algorithm for land (SEBAL) - 1. Formulation, *J. Hydrol.*, 212(1–4),
910 198–212, doi:10.1016/s0022-1694(98)00253-4, 1998a.

911 Bastiaanssen, W. G. M., Pelgrum, H., Wang, J., Ma, Y., Moreno, J. F., Roerink, G. J. and van
912 der Wal, T.: A remote sensing surface energy balance algorithm for land (SEBAL): Part 2:
913 Validation, *J. Hydrol.*, 212–213, 213–229, doi:10.1016/S0022-1694(98)00254-6, 1998b.

914 Bathiany, S., Claussen, M., Brovkin, V., Raddatz, T. and Gayler, V.: Combined biogeophysical
915 and biogeochemical effects of large-scale forest cover changes in the MPI earth system model,
916 *Biogeosciences*, 7(5), 1383–1399, doi:10.5194/bg-7-1383-2010, 2010.

917 Beltrán-Przekurat, A., Pielke Sr, R. A., Eastman, J. L. and Coughenour, M. B.: Modelling the
918 effects of land-use/land-cover changes on the near-surface atmosphere in southern South
919 America, *Int. J. Climatol.*, 32(8), 1206–1225, doi:10.1002/joc.2346, 2012.

920 Bindhu, V. M., Narasimhan, B. and Sudheer, K. P.: Development and verification of a non-
921 linear disaggregation method (NL-DisTrad) to downscale MODIS land surface temperature to
922 the spatial scale of Landsat thermal data to estimate evapotranspiration, *Remote Sens. Environ.*,
923 135, 118–129, doi:10.1016/j.rse.2013.03.023, 2013.

924 Boisier, J. P., de Noblet-Ducoudré, N. and Ciais, P.: Historical land-use-induced
925 evapotranspiration changes estimated from present-day observations and reconstructed land-
926 cover maps, *Hydrol. Earth Syst. Sci.*, 18(9), 3571–3590, doi:10.5194/hess-18-3571-2014, 2014.

927 Bridhikitti, A. and Overcamp, T. J.: Estimation of Southeast Asian rice paddy areas with
928 different ecosystems from moderate-resolution satellite imagery, *Agric. Ecosyst. Environ.*,
929 146(1), 113–120, doi:10.1016/j.agee.2011.10.016, 2012.

930 Bright, R. M., Zhao, K., Jackson, R. B. and Cherubini, F.: Quantifying surface albedo and other
931 direct biogeophysical climate forcings of forestry activities, *Glob. Change Biol.*, 21(9), 3246–
932 3266, doi:10.1111/gcb.12951, 2015.

- 933 Clough, Y., Krishna, V. V., Corre, M. D., Darras, K., Denmead, L. H., Meijide, A., Moser, S.,
934 Musshoff, O., Steinebach, S., Veldkamp, E., Allen, K., Barnes, A. D., Breidenbach, N., Brose,
935 U., Buchori, D., Daniel, R., Finkeldey, R., Harahap, I., Hertel, D., Holtkamp, A. M., Hörandl,
936 E., Irawan, B., Jaya, I. N. S., Jochum, M., Klarner, B., Knohl, A., Kotowska, M. M.,
937 Krashevskaya, V., Kreft, H., Kurniawan, S., Leuschner, C., Maraun, M., Melati, D. N.,
938 Opfermann, N., Pérez-Cruzado, C., Prabowo, W. E., Rembold, K., Rizali, A., Rubiana, R.,
939 Schneider, D., Tjitrosoedirdjo, S. S., Tjoa, A., Tschardtke, T. and Scheu, S.: Land-use choices
940 follow profitability at the expense of ecological functions in Indonesian smallholder landscapes,
941 *Nat. Commun.*, 7, 13137, 2016.
- 942 Coll, C., Wan, Z. and Galve, J. M.: Temperature-based and radiance-based validations of the
943 V5 MODIS land surface temperature product, *J. Geophys. Res.*, 114(D20), 2009.
- 944 Coll, C., Galve, J. M., Sanchez, J. M. and Caselles, V.: Validation of Landsat-7/ETM+ Thermal-
945 Band Calibration and Atmospheric Correction With Ground-Based Measurements, *Geosci.*
946 *Remote Sens. IEEE Trans. On*, 48(1), 547–555, doi:10.1109/TGRS.2009.2024934, 2010.
- 947 Costa, M. H., Yanagi, S. N. M., Souza, P. J. O. P., Ribeiro, A. and Rocha, E. J. P.: Climate
948 change in Amazonia caused by soybean cropland expansion, as compared to caused by
949 pastureland expansion, *Geophys. Res. Lett.*, 34(7), doi:10.1029/2007GL029271, 2007.
- 950 Davin, E. L. and de Noblet-Ducoudré, N.: Climatic Impact of Global-Scale Deforestation:
951 Radiative versus Nonradiative Processes, *J. Clim.*, 23(1), 97–112,
952 doi:10.1175/2009JCLI3102.1, 2010.
- 953 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
954 Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
955 Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
956 Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M.,
957 McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P.,
958 Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and
959 performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597,
960 doi:10.1002/qj.828, 2011.
- 961 Dislich, C., Keyel, A. C., Salecker, J., Kisel, Y., Meyer, K. M., Auliya, M., Barnes, A. D.,
962 Corre, M. D., Darras, K., Faust, H., Hess, B., Klasen, S., Knohl, A., Kreft, H., Meijide, A.,
963 Nurdiansyah, F., Otten, F., Pe'er, G., Steinebach, S., Tarigan, S., Tölle, M. H., Tschardtke, T.
964 and Wiegand, K.: A review of the ecosystem functions in oil palm plantations, using forests as
965 a reference system, *Biol. Rev.*, doi:10.1111/brv.12295, 2016.
- 966 Drescher, J., Rembold, K., Allen, K., Beckschäfer, P., Buchori, D., Clough, Y., Faust, H., Fauzi,
967 A. M., Gunawan, D., Hertel, D., Irawan, B., Jaya, I. N. S., Klarner, B., Kleinn, C., Knohl, A.,
968 Kotowska, M. M., Krashevskaya, V., Krishna, V., Leuschner, C., Lorenz, W., Meijide, A., Melati,
969 D., Nomura, M., Pérez-Cruzado, C., Qaim, M., Siregar, I. Z., Steinebach, S., Tjoa, A.,
970 Tschardtke, T., Wick, B., Wiegand, K., Kreft, H. and Scheu, S.: Ecological and socio-economic
971 functions across tropical land use systems after rainforest conversion, *Philos. Trans. R. Soc.*
972 *Lond. B Biol. Sci.*, 371(1694), doi:10.1098/rstb.2015.0275, 2016.
- 973 Fall, S., Niyogi, D., Gluhovsky, A., Pielke, R. A., Kalnay, E. and Rochon, G.: Impacts of land
974 use land cover on temperature trends over the continental United States: assessment using the
975 North American Regional Reanalysis, *Int. J. Climatol.*, 30(13), 1980–1993,
976 doi:10.1002/joc.1996, 2010.

- 977 Feddema, J. J., Oleson, K. W., Bonan, G. B., Mearns, L. O., Buja, L. E., Meehl, G. A. and
 978 Washington, W. M.: The Importance of Land-Cover Change in Simulating Future Climates,
 979 *Science*, 310(5754), 1674, doi:10.1126/science.1118160, 2005.
- 980 Hoffmann, W. A. and Jackson, R. B.: Vegetation–Climate Feedbacks in the Conversion of
 981 Tropical Savanna to Grassland, *J. Clim.*, 13(9), 1593–1602, doi:10.1175/1520-
 982 0442(2000)013<1593:VCFITC>2.0.CO;2, 2000.
- 983 Houspanossian, J., Nosoetto, M. and Jobbágy, E. G.: Radiation budget changes with dry forest
 984 clearing in temperate Argentina, *Glob. Change Biol.*, 19(4), 1211–1222,
 985 doi:10.1111/gcb.12121, 2013.
- 986 Idso, S. B. and Jackson, R. D.: Thermal radiation from the atmosphere, *J. Geophys. Res.*,
 987 74(23), 5397–5403, doi:10.1029/JC074i023p05397, 1969.
- 988 Jiang, Y., Fu, P. and Weng, Q.: Assessing the Impacts of Urbanization-Associated Land
 989 Use/Cover Change on Land Surface Temperature and Surface Moisture: A Case Study in the
 990 Midwestern United States, *Remote Sens.*, 7(4), doi:10.3390/rs70404880, 2015.
- 991 Kayet, N., Pathak, K., Chakrabarty, A. and Sahoo, S.: Spatial impact of land use/land cover
 992 change on surface temperature distribution in Saranda Forest, Jharkhand, *Model. Earth Syst.*
 993 *Environ.*, 2(3), 1–10, doi:10.1007/s40808-016-0159-x, 2016.
- 994 Lawrence, D. and Vandecar, K.: Effects of tropical deforestation on climate and agriculture,
 995 *Nat. Clim. Change*, 5(1), 27–36, 2015.
- 996 Lee, X., Goulden, M. L., Hollinger, D. Y., Barr, A., Black, T. A., Bohrer, G., Bracho, R., Drake,
 997 B., Goldstein, A., Gu, L., Katul, G., Kolb, T., Law, B. E., Margolis, H., Meyers, T., Monson,
 998 R., Munger, W., Oren, R., Paw U, K. T., Richardson, A. D., Schmid, H. P., Staebler, R., Wofsy,
 999 S. and Zhao, L.: Observed increase in local cooling effect of deforestation at higher latitudes,
 1000 *Nature*, 479(7373), 384–387, doi:10.1038/nature10588, 2011.
- 1001 van Leeuwen, T. T., Frank, A. J., Jin, Y., Smyth, P., Goulden, M. L., van der Werf, G. R. and
 1002 Randerson, J. T.: Optimal use of land surface temperature data to detect changes in tropical
 1003 forest cover, *J. Geophys. Res. Biogeosciences*, 116(G2), doi:10.1029/2010JG001488, 2011.
- 1004 Li, F., Jackson, T. J., Kustas, W. P., Schmugge, T. J., French, A. N., Cosh, M. H. and Bindlish,
 1005 R.: Deriving land surface temperature from Landsat 5 and 7 during SMEX02/SMACEX, 2002
 1006 *Soil Moisture Exp. SMEX02*, 92(4), 521–534, doi:10.1016/j.rse.2004.02.018, 2004.
- 1007 Li, Y., Zhao, M., Motesharrei, S., Mu, Q., Kalnay, E. and Li, S.: Local cooling and warming
 1008 effects of forests based on satellite observations, *Nat. Commun.*, 6 [online] Available from:
 1009 <http://dx.doi.org/10.1038/ncomms7603>, 2015.
- 1010 Liang, S.: Narrowband to broadband conversions of land surface albedo I: Algorithms, *Remote*
 1011 *Sens. Environ.*, 76(2), 213–238, doi:10.1016/S0034-4257(00)00205-4, 2000.
- 1012 Lim, Y.-K., Cai, M., Kalnay, E. and Zhou, L.: Observational evidence of sensitivity of surface
 1013 climate changes to land types and urbanization, *Geophys. Res. Lett.*, 32(22),
 1014 doi:10.1029/2005GL024267, 2005.
- 1015 Lim, Y.-K., Cai, M., Kalnay, E. and Zhou, L.: Impact of Vegetation Types on Surface
 1016 Temperature Change, *J. Appl. Meteorol. Climatol.*, 47(2), 411–424, 2008.

- 1017 Loarie, S. R., Lobell, D. B., Asner, G. P., Mu, Q. and Field, C. B.: Direct impacts on local
1018 climate of sugar-cane expansion in Brazil, *Nat. Clim. Change*, 1(2), 105–109,
1019 doi:10.1038/nclimate1067, 2011a.
- 1020 Loarie, S. R., Lobell, D. B., Asner, G. P. and Field, C. B.: Land-Cover and Surface Water
1021 Change Drive Large Albedo Increases in South America, *Earth Interact.*, 15(7), 1–16, 2011b.
- 1022 Longobardi, P., Montenegro, A., Beltrami, H. and Eby, M.: Deforestation Induced Climate
1023 Change: Effects of Spatial Scale, *PLoS ONE*, 11(4), e0153357,
1024 doi:10.1371/journal.pone.0153357, 2016.
- 1025 Luskin, M. S., Christina, E. D., Kelley, L. C. and Potts, M. D.: Modern Hunting Practices and
1026 Wild Meat Trade in the Oil Palm Plantation-Dominated Landscapes of Sumatra, Indonesia,
1027 *Hum. Ecol.*, 42(1), 35–45, doi:10.1007/s10745-013-9606-8, 2014.
- 1028 Mahmood, R., Pielke, R. A., Hubbard, K. G., Niyogi, D., Dirmeyer, P. A., McAlpine, C.,
1029 Carleton, A. M., Hale, R., Gameda, S., Beltrán-Przekurat, A., Baker, B., McNider, R., Legates,
1030 D. R., Shepherd, M., Du, J., Blanken, P. D., Frauenfeld, O. W., Nair, U. S. and Fall, S.: Land
1031 cover changes and their biogeophysical effects on climate, *Int. J. Climatol.*, 34(4), 929–953,
1032 doi:10.1002/joc.3736, 2014.
- 1033 Margono, B. A., Turubanova, S., Zhuravleva, I., Potapov, P., Tyukavina, A., Baccini, A., Goetz,
1034 S. and Hansen, M. C.: Mapping and monitoring deforestation and forest degradation in Sumatra
1035 (Indonesia) using Landsat time series data sets from 1990 to 2010, *Environ. Res. Lett.*, 7(3),
1036 34010, doi:10.1088/1748-9326/7/3/034010, 2012.
- 1037 Margono, B. A., Potapov, P. V., Turubanova, S., Stolle, F. and Hansen, M. C.: Primary forest
1038 cover loss in Indonesia over 2000–2012, *Nat. Clim Change*, 4(8), 730–735, 2014.
- 1039 Marlier, M. E., DeFries, R., Pennington, D., Nelson, E., Ordway, E. M., Lewis, J., Koplitz, S.
1040 N. and Mickley, L. J.: Future fire emissions associated with projected land use change in
1041 Sumatra, *Glob. Change Biol.*, 21(1), 345–362, doi:10.1111/gcb.12691, 2015.
- 1042 Meijide, A., Röhl, A., Fan, Y., Herbst, M., Niu, F., Tiedemann, F., June, T., Rauf, A., Hölscher,
1043 D. and Knohl, A.: Controls of water and energy fluxes in oil palm plantations: Environmental
1044 variables and oil palm age, *Agric. For. Meteorol.*, 239, 71–85,
1045 doi:10.1016/j.agrformet.2017.02.034, 2017.
- 1046 Merten, J., Röhl, A., Guillaume, T., Meijide, A., Tarigan, S., Agusta, H., Dislich, C., Dittrich,
1047 C., Faust, H., Gunawan, D., Hein, J., Hendrayanto, Knohl, A., Kuzyakov, Y., Wiegand, K. and
1048 Hölscher, D.: Water scarcity and oil palm expansion: social views and environmental processes,
1049 *Ecol. Soc.*, 21(2), doi:10.5751/ES-08214-210205, 2016.
- 1050 Miettinen, J., Shi, C. and Liew, S. C.: Deforestation rates in insular Southeast Asia between
1051 2000 and 2010, *Glob. Change Biol.*, 17(7), 2261–2270, 2011.
- 1052 Miettinen, J., Hooijer, A., Wang, J., Shi, C. and Liew, S. C.: Peatland degradation and
1053 conversion sequences and interrelations in Sumatra, *Reg. Environ. Change*, 12(4), 729–737,
1054 doi:10.1007/s10113-012-0290-9, 2012.

- 1055 Mildrexler, D. J., Zhao, M. and Running, S. W.: A global comparison between station air
1056 temperatures and MODIS land surface temperatures reveals the cooling role of forests, *J.*
1057 *Geophys. Res. Biogeosciences*, 116(G3), doi:10.1029/2010JG001486, 2011.
- 1058 Nosetto, M. D., Jobbágy, E. G. and Paruelo, J. M.: Land-use change and water losses: the case
1059 of grassland afforestation across a soil textural gradient in central Argentina, *Glob. Change*
1060 *Biol.*, 11(7), 1101–1117, doi:10.1111/j.1365-2486.2005.00975.x, 2005.
- 1061 Oliveira, L. J. C., Costa, M. H., Soares-Filho, B. S. and Coe, M. T.: Large-scale expansion of
1062 agriculture in Amazonia may be a no-win scenario, *Environ. Res. Lett.*, 8(2), 24021, 2013.
- 1063 Paterson, R. R. M., Kumar, L., Taylor, S. and Lima, N.: Future climate effects on suitability for
1064 growth of oil palms in Malaysia and Indonesia, *Sci. Rep.*, 5, 14457, 2015.
- 1065 Peng, S.-S., Piao, S., Zeng, Z., Ciais, P., Zhou, L., Li, L. Z. X., Myneni, R. B., Yin, Y. and
1066 Zeng, H.: Afforestation in China cools local land surface temperature, *Proc. Natl. Acad. Sci.*,
1067 111(8), 2915–2919, 2014.
- 1068 Pongratz, J., Bounoua, L., DeFries, R. S., Morton, D. C., Anderson, L. O., Mauser, W. and
1069 Klink, C. A.: The Impact of Land Cover Change on Surface Energy and Water Balance in Mato
1070 Grosso, Brazil, *Earth Interact.*, 10(19), 1–17, 2006.
- 1071 Salazar, A., Baldi, G., Hirota, M., Syktus, J. and McAlpine, C.: Land use and land cover change
1072 impacts on the regional climate of non-Amazonian South America: A review, *Glob. Planet.*
1073 *Change*, 128, 103–119, doi:10.1016/j.gloplacha.2015.02.009, 2015.
- 1074 Salazar, A., Katzfey, J., Thatcher, M., Syktus, J., Wong, K. and McAlpine, C.: Deforestation
1075 changes land–atmosphere interactions across South American biomes, *Glob. Planet. Change*,
1076 139, 97–108, doi:10.1016/j.gloplacha.2016.01.004, 2016.
- 1077 Sheil, D., Casson, A., Meijaard, E., Van Noordwijk, M., Gaskell, J., Sunderland-Groves, J.,
1078 Wertz, K. and Kanninen, M.: The impacts and opportunities of oil palm in Southeast Asia: What
1079 do we know and what do we need to know?, Center for International Forestry Research
1080 (CIFOR), Bogor, Indonesia., 2009.
- 1081 Silvério, D. V., Brando, P. M., Macedo, M. N., Beck, P. S. A., Bustamante, M. and Coe, M. T.:
1082 Agricultural expansion dominates climate changes in southeastern Amazonia: the overlooked
1083 non-GHG forcing, *Environ. Res. Lett.*, 10(10), 104015, 2015.
- 1084 Snyder, W. C., Wan, Z., Zhang, Y. and Feng, Y.-Z.: Classification-based emissivity for land
1085 surface temperature measurement from space, *Int. J. Remote Sens.*, 19(14), 2753–2774,
1086 doi:10.1080/014311698214497, 1998.
- 1087 Sobrino, J. A., Jiménez-Muñoz, J. C. and Paolini, L.: Land surface temperature retrieval from
1088 LANDSAT TM 5, *Remote Sens. Environ.*, 90(4), 434–440, doi:10.1016/j.rse.2004.02.003,
1089 2004.
- 1090 Sobrino, J. A., Jiménez-Muñoz, J. C., Zarco-Tejada, P. J., Sepulcre-Cantó, G. and de Miguel,
1091 E.: Land surface temperature derived from airborne hyperspectral scanner thermal infrared data,
1092 *Remote Sens. Environ.*, 102(1–2), 99–115, doi:10.1016/j.rse.2006.02.001, 2006.
- 1093 Sobrino, J. A., Jimenez-Muoz, J. C., Soria, G., Romaguera, M., Guanter, L., Moreno, J., Plaza,
1094 A. and Martinez, P.: Land Surface Emissivity Retrieval From Different VNIR and TIR Sensors,

- 1095 Geosci. Remote Sens. IEEE Trans. On, 46(2), 316–327, doi:10.1109/TGRS.2007.904834,
1096 2008.
- 1097 Spracklen, D. V., Arnold, S. R. and Taylor, C. M.: Observations of increased tropical rainfall
1098 preceded by air passage over forests, *Nature*, 489(7415), 282–285, doi:10.1038/nature11390,
1099 2012.
- 1100 Tölle, M. H., Engler, S. and Panitz, H.-J.: Impact of Abrupt Land Cover Changes by Tropical
1101 Deforestation on Southeast Asian Climate and Agriculture, *J. Clim.*, 30(7), 2587–2600,
1102 doi:10.1175/JCLI-D-16-0131.1, 2017.
- 1103 Verstraeten, W. W., Veroustraete, F. and Feyen, J.: Estimating evapotranspiration of European
1104 forests from NOAA-imagery at satellite overpass time: Towards an operational processing
1105 chain for integrated optical and thermal sensor data products, *Remote Sens. Environ.*, 96(2),
1106 256–276, doi:10.1016/j.rse.2005.03.004, 2005.
- 1107 Vlassova, L., Perez-Cabello, F., Nieto, H., Martín, P., Riaño, D. and de la Riva, J.: Assessment
1108 of Methods for Land Surface Temperature Retrieval from Landsat-5 TM Images Applicable to
1109 Multiscale Tree-Grass Ecosystem Modeling, *Remote Sens.*, 6(5), doi:10.3390/rs6054345,
1110 2014.
- 1111 Voogt, J. A. and Oke, T. R.: Effects of urban surface geometry on remotely-sensed surface
1112 temperature, *Int. J. Remote Sens.*, 19(5), 895–920, doi:10.1080/014311698215784, 1998.
- 1113 Wan, Z., Zhang, Y., Zhang, Q. and Li, Z.-L.: Quality assessment and validation of the MODIS
1114 global land surface temperature, *Int. J. Remote Sens.*, 25(1), 261–274,
1115 doi:10.1080/0143116031000116417, 2004.
- 1116 Wang, J., Pan, F., Soininen, J., Heino, J. and Shen, J.: Nutrient enrichment modifies
1117 temperature-biodiversity relationships in large-scale field experiments, *Nat. Commun.*, 7,
1118 13960, doi:10.1038/ncomms13960, 2016.
- 1119 Weng, Q.: Thermal infrared remote sensing for urban climate and environmental studies:
1120 Methods, applications, and trends, *ISPRS J. Photogramm. Remote Sens.*, 64(4), 335–344,
1121 doi:10.1016/j.isprsjprs.2009.03.007, 2009.
- 1122 Weng, Q., Lu, D. and Schubring, J.: Estimation of land surface temperature–vegetation
1123 abundance relationship for urban heat island studies, *Remote Sens. Environ.*, 89(4), 467–483,
1124 doi:10.1016/j.rse.2003.11.005, 2004.
- 1125 Wicke, B., Sikkema, R., Dornburg, V. and Faaij, A.: Exploring land use changes and the role
1126 of palm oil production in Indonesia and Malaysia, *Land Use Policy*, 28(1), 193–206, 2011.
- 1127 Wukelic, G. E., Gibbons, D. E., Martucci, L. M. and Foote, H. P.: Radiometric calibration of
1128 Landsat Thematic Mapper thermal band, *Remote Sens. Environ.*, 28(0), 339–347,
1129 doi:10.1016/0034-4257(89)90125-9, 1989.
- 1130 Yue, W., Xu, J., Tan, W. and Xu, L.: The relationship between land surface temperature and
1131 NDVI with remote sensing: application to Shanghai Landsat 7 ETM+ data, *Int. J. Remote Sens.*,
1132 28(15), 3205–3226, doi:10.1080/01431160500306906, 2007.
- 1133 Zhang, Z. and He, G.: Generation of Landsat surface temperature product for China, 2000–
1134 2010, *Int. J. Remote Sens.*, 34(20), 7369–7375, doi:10.1080/01431161.2013.820368, 2013.

1135 Zhou, X. and Wang, Y.-C.: Dynamics of Land Surface Temperature in Response to Land-
1136 Use/Cover Change, *Geogr. Res.*, 49(1), 23–36, doi:10.1111/j.1745-5871.2010.00686.x, 2011.

1137