

Dear Editor,

We deeply appreciate you for giving us an opportunity to improve our manuscript. We would like to thank all of you and the two reviewers for the thoughtful and valuable suggestions on our manuscript entitled “**Divergence of dominant factors on soil microbial communities and functions in forest ecosystems along a climatic gradient**” [ID: bg-2017-243]. According to the comments and suggestions, we have carefully revised our manuscript. We have followed the formatting requirements as presented in the Guide for Authors. We have uploaded the document of the responses to reviewer and a clean manuscript. Here are the point-to-point responses (color-coded blue) to Editor’s and **Reviewer’s** comments (color-coded black). The page and line numbers mentioned here refer to our latest revision of our manuscript simultaneously submitted.

Reviewer #1:

Interactive comment on “Divergence of dominant factors on soil microbial communities and functions in forest ecosystems along a climatic gradient” by Zhiwei Xu et al. Anonymous Referee #1 Received and published: 28 September 2017 Divergence of dominant factors on soil microbial communities and functions in forest ecosystems along a climatic gradient is a investigation paper. Authors chose 12 forests along three climate zones to investigate the variation of soil activities and microbe structures among these forests along three climate zones. The results showed that soil enzyme activities and microbial PLFAs differed with forest types along climatic zones. Both climate and forest type had significant effects on soil enzyme activities and microbial communities. Litter nutrients made an important effect to variations in the soil microbial communities and enzyme activities in temperate zones, while soil micro-climate and nutrients were the main effect factors on the soil microbial community structure and enzymatic activities in warm temperate and subtropical zones. The C1 BGD Interactive comment Printer-friendly version Discussion paper has valuable to be published in this journal. However, the following points should be considered to revise.

(1) Abstract: line 44-45 "Our results indicate that the main controls on soil microbes and functions vary across forest ecosystems in different climatic zones, and that the effects of soil moisture content, soil temperature, and the soil N/P ratio were considerable." was the results, not indications. Instead, please give a general summery about reasons of the variation.

AN: we have improved this part as "Our results showed that the main controls on soil microbes and functions vary in different climatic zones, and that the effects of soil moisture content, soil temperature, clay content, and the soil N/P ratio were considerable." (P2, Line 49-52).

(2) Materials and Method: The investigation was conducted in July and August in three climate zones, It is better to illustrate the climate information of the investigation month and detail investigation date in each site. This is because the activities of microbe is very sensitive to the climates, especially the moisture and temperature.

AN: We illustrated the climate information of the investigation month in the text. The average temperature of the sampling month was 21.3 °C, 17.4°C, 27.3°C with the relative humidity of 78%, 60-65%, 83.5% in LS, TY, and DH, respectively. The sampling dates are Jul.5 2013, Jul.28 2013, Aug.15 2013 in LS, TY, and DH, respectively. (P5, Line136-139).

(3) Results: In the 3.1 section, the activities of four enzymes did not be described carefully. Most information were ignored, for example, there were not comparison between forest types in the same climate zone. And there were not comparison between different climate zones, for example, the LAP activities of microbes in worm temperate zone were much higher than that in temperate zone.

AN: We have added necessary description about the enzyme activities. The soil BG and NAG activities were much higher in the coniferous forest than in

the conifer broad-leaved mixed forests and the broad-leaved forests (Table S2). The soil AP enzyme activities were highest in the conifer broad-leaved mixed forests and lowest in the coniferous forests (Table S2). (P8, Line208-211).

The soil BG, NAG, and LAP activities were much higher in the warm temperate zone than in the temperate and the subtropical climate zones (Table S2). The AP activities were highest in the subtropical climate zone (Table S2). (P8, Line 213-215)

(4) In the 3.2, same as 3.1, no comparison among three climate zones. Although there were difference among forest types in the same zone, authors should compare the similar quality forest such as SCB along three climate zones. If the results of these comparison could be reported, more mechanism of divergences among zones and forest types could be understood very well.

AN: We have added necessary description about the microbial communities. We compare the microbial PLFAs among the three different climate zones and three forest types (conifer broad-leaved mixed forest, broad-leaved forest, and coniferous forest), respectively.

The forest type had a significant effect on the soil bacteria, fungi, gram-positive bacteria (G<sup>+</sup>), and gram-negative bacteria (G<sup>-</sup>) PLFAs (Table 2). The soil total PLFAs, bacteria, G<sup>+</sup>, G<sup>-</sup>, and actinomycete were much higher in the conifer broad-leaved mixed forests than in the coniferous forests and the broad-leaved forests (Table S2). The soil fungi was highest in the broad-leaved forest and lowest in the coniferous forest (Table S2). (P8, Line224-229).

With the exception of the soil G<sup>+</sup>/ G<sup>-</sup>, the effects of the combination of climate and forest type on all soil PLFAs were significant, and were stronger than the individual effects of either climate or forest type (Table 2, Table S2). Climate had a significant effect on the total PLFAs, fungi, and G<sup>-</sup> (P<0.0001) (Table 2). The soil total PLFAs, bacteria, G<sup>+</sup>, and G<sup>-</sup> were much higher in the temperate zone than in the warm temperate and the subtropical zones (Table

S2). The fungi, F/B, and G<sup>+</sup>/G<sup>-</sup> were highest in the subtropical zone (Table S2). (P9, Line 230-235)

(5) In conclusion part, we would like to see the conclusion what changes along the climate zone could be found

AN: We have added description in the abstract ((P2, Line 36-44), result and conclusions part about the variations in microbial communities and enzyme activities along the climate zone.

The soil BG, NAG, and LAP activities were much higher in the warm temperate zone than in the temperate and the subtropical climate zones (Table S2). The AP activities were highest in the subtropical climate zone (Table S2). (P8, Line 213-215)

With the exception of the soil G<sup>+</sup>/G<sup>-</sup>, the effects of the combination of climate and forest type on all soil PLFAs were significant, and were stronger than the individual effects of either climate or forest type (Table 2, Table S2). Climate had a significant effect on the total PLFAs, fungi, and G<sup>-</sup> (P<0.0001) (Table 2). The soil total PLFAs, bacteria, G<sup>+</sup>, and G<sup>-</sup> were much higher in the temperate zone than in the warm temperate and the subtropical zones (Table S2). The fungi, F/B, and G<sup>+</sup>/G<sup>-</sup> were highest in the subtropical zone (Table S2). (P9, Line 230-235)

Conclusion: Except AP, soil enzyme activities were highest in warm temperate zone. Soil tPLFAs, bacteria, G<sup>-</sup> increased from temperate zone to subtropical zone, but fungi was in reverse. (P15, Line 407-409).

(6) Discussion: 4.1, It is unclear what the response of soil enzyme activities and microbial plfas to variation of forest types is. Authors should clearly discuss the variation pattern and formation reason.

AN: We have improved this part. Forests in the same climate zone developed similar microbe functions which confirmed the result that the effect of climate

on soil enzyme activities were stronger than the forest type and their interactive effect. However, there were still differences among the enzyme activities in different forest types of the same climate zone. Soil microorganisms are usually considered to be C limited, and the litter inputs with high C/N ratio of PCB in the temperate zone will stimulate microbes to grow and secrete more enzymes (Table 1). Therefore, all enzyme activities were highest in PCB in the temperate zone. (P10, Line 267-273).

The high soil BG enzyme activities in the LOw forest in the warm temperate zone reflect the litter inputs with low C. Because that soil enzyme activities will not continuously increase or decrease as nutrient availability increases or decreases. When the soil nutrients are short in supply, microbes will potentially increase production of nutrient-acquiring enzymes, because they are expected to optimize the allocation of their resource reserves by acquiring the resource that is most limiting (Bloom et al., 1985). (P10, Line 273-278).

The interactive effect of climate and forest type were more important than the individual effect of them. Therefore, the soil microbial communities of the 12 forests were separated from each other. Vegetation transfers substrate material of varying quality to microbes through litter fall. Fungi are more suitable for life in environments containing higher C/N ratios and low soil pH (Nilsson et al., 2012). The four broadleaved forests were high in litter C/N ratio (Table 1). Therefore, fungi were dominated in this harsh nutrient environments and highest in broadleaved forests. The litter and soil from conifer broad-leaved mixed forest were high in C, N, and P, and promotes the propagation of bacteria that favor high-nutrient soil (Priha and Smolander, 1997; Priha et al., 2001). Therefore, the structures and functions of the soil microbial communities that developed in the different types of forest were unique. (P10, Line 280-286; P11, Line 287-289)

To avoid the repetition with the 4.2 and 4.3, some more detail reasons of the variations were discussed later.

(7) 4.2, How to compare the common effect and key effect? if there is obvious difference between two effects, could you explain the identification method of two effects.

AN: The common effect refers to the same environmental variables which are significantly correlated with the RDA1 in the three biplots of the three climate zones ( $P < 0.05$ ). The key effect refers to the environmental variables that were more important in determining soil microbial communities and functions of the individual climate zones ( $P < 0.01$ ).

In addition, we have done a new RDA again by putting the data of 12 forests in the three climate zones together to observe the variations in soil enzyme activities (Fig.S1) and microbial communities (Fig.S2) among different forest types and climate zones.

(8) Conclusion: Authors should address the main conclusion of the variation of enzyme activities and microbial community among forest types along the three zones in the suitable part of the paragraph.

AN: We have added the main conclusion of the variations of enzyme activities and microbial community among forest types in the result and conclusion.

The soil total PLFAs, bacteria,  $G^+$ ,  $G^-$ , and actinomycete were much higher in the conifer broad-leaved mixed forests than in the coniferous forests and the broad-leaved forests. The soil BG and NAG activities were much higher in the coniferous forest than in the conifer broad-leaved mixed forests and the broad-leaved forests. Except AP, soil enzyme activities were highest in warm temperate zone. Soil tPLFAs, bacteria,  $G^-$  increased from temperate zone to subtropical zone, but fungi was in reverse. (P15, Line 404-409)

Minor mistakes

(9) line 135. authors should give detail information about collection such as which samples were collected in July?

AN: The average temperature of the sampling month was 21.3 °C, 17.4°C, 27.3°C with the relative humidity of 78%, 60-65%, 83.5% in LS, TY, and DH, respectively. The sampling dates are Jul.5 2013, Jul.28 2013, Aug.15 2013 in LS, TY, and DH, respectively. (P5, line 136-139; Table 1).

(10) SCB in temperate zone was not same as it in Subtropical forest, it is better abbreviated as SCBt SCBs

AN: DONE (Table 1, Figure 1, Figure 2 and Figure 4).

(11) Fig. 2 ABCD was represent different enzyme activities, please check them.

AN: DONE (P24, Figure 1).

(12) The format of some references did not fit with the format of this journal such as New Physiologist which did not was abbreviated.

AN: DONE (P16, Line 439). We have checked all through the text and made necessary variations.

Reviewer 2

(13) The authors present a comprehensive study of soil microbial communities and extracellular enzyme activities in different forests along a climatic gradient. The methods are technically sound. This paper clearly elucidates the dominant factors controlling microbial communities and enzyme activities in each climatic zone. The authors also attempt to emphasize the importance of climatic zones in addition to forest types. However, it's unclear for readers why different dominant factors exhibit in different climatic zones. For example, the authors

state that “soil clay content had most influence on the soil enzyme activities in subtropical forests” (Line 353). However, the following discussion is very general and does not explain why this is only found in the subtropics.

AN: We have improved this part. Therefore, soil enzyme activities and microbial PLFAs were highest in the SCBs forest with finely texture. Except SCBt in the temperate zone and PT in the warm temperate zone, the soil clay content were not significant different among other three forest types. However, the soil clay contents of the four forest types in the subtropical zone were significant different from each other and important for variations in microbial communities and functions (Table 1). (P14, Line 379-384).

(14) Here is another example, soil nutrients (N, P) are more important in warm temperate and subtropical forests than in temperate forests, because nutrients are more likely limiting factors in warm temperate and subtropical forest. This kind of comparison between different climatic zones should be expanded in Discussion and could add value to this study.

AN: We have improved this part as “The soil TN and TP were lower in the warm temperate and subtropical zone than in the temperate zone in our study (Table 1), and these two kinds of nutrients were more likely limiting factors in warm temperate and subtropical forest (DeForest et al., 2012; Xu et al., 2017).



Therefore, soil TN and TP are more important in warm temperate and subtropical forests than in temperate forests.” (P13, Line 357-360).

(15) I have a few more suggestions to improve the presentation of this study: In Conclusions, soil clay fraction is identified as an important predictor in subtropical zones. However, “soil clay” is not mentioned in Abstract.

AN: We have improved the abstract. Our results showed that the main controls on soil microbes and functions vary in different climatic zones, and that the effects of soil moisture content, soil temperature, clay content, and the soil N/P ratio were considerable. (P2, Line 49-52).

(16) Line 266-268: I don’t understand the logic here. The authors are talking about microbial/enzyme responses to forest types in Section 4.1. The concluding sentence addresses “climatic region may be more important than forest types” without any expanded discussion, though I understand “climatic effects” may be indirectly discussed in Section 4.3.

AN: We have moved this sentence to the section 4.2 and improved it as “This was also demonstrated by the stronger effect of climate on soil enzyme activities and the combined interaction effect of climate and forest type on soil microbial communities. Other studies have reported that precipitation and mean

annual temperature played important roles in explaining on the large-scale distribution of soil microbial community composition and functions (de Vries et al., 2012; Xu et al., 2017).” (P11, Line 314-315; P12, Line 316-319).

(17) Line 298-300: This clause does not explain why there are more Gram-negative bacteria, less Gram-positive bacteria, and (less?) bacteria PLFAs under increasing pH.

AN: We have improved this part as “Soil G<sup>+</sup>/G<sup>-</sup> ratios were highest in the subtropical forest where G<sup>-</sup> bacteria PLFAs were least abundant, which may reflect microbial growth strategies. The G<sup>+</sup> bacteria are primarily K-strategists that can survive over long periods in the soil under harsh conditions with lower soil pH (Andrews & Hall, 1986). Increased pH causes an increase in bacterial diversity and a shift in the bacterial community to more G<sup>-</sup> and fewer G<sup>+</sup> bacteria PLFAs (Wu et al., 2009; Shen et al., 2013). “(P12, Line 324-329).

(18) Line 210-212: please spell out G<sup>-</sup> (Gram-negative bacteria) and G<sup>+</sup> (Gram-positive bacteria) when they are first introduced.

AN: DONE (P8, Line 225-226).

(19) Line 241-243: The causal explanation herein is not specifically related to the results in Section 3.3 and Fig. 4a. Does the “higher inputs of mixed litter” mean higher litter C/N and lower litter TN? To my understanding, from Fig.4a, BG/NGC/LAP activities are positively correlated with litter C/N and negatively correlated with litter TN. The following explanation for the warm temperate zone is more informative.

AN: We have improved this part as “Soil microorganisms are usually considered to be C limited, and the litter inputs with high C/N ratio of PCB in the temperate zone will stimulate microbes to grow and secrete more enzymes (Table 1). Therefore, all enzyme activities were highest in PCB in the temperate zone.”

(P10, Line 270-273).

(20) Line 262: please spell out SLA and LDMC.

AN: We have deleted this part.

(21) Line 328: please spell out F/B ratio.

AN: DONE (P13, Line 345).

Thanks again to you and the two reviewers for the thoughtful and thorough comments.

We hope that our revisions will be satisfactory, and we are very happy to work with you and the reviewers to resolve any remaining problems.

Yours sincerely,

Zhiwei Xu, Guirui Yu, Xinyu Zhang, Nianpeng He, Qiufeng Wang,

Shengzhong Wang, Xiaofeng Xu, Ruili Wang, Ning Zhao

## **List of all relevant changes made in the manuscript**

1. Change to the affiliations of the first author.
2. Changes to the abstract.
3. Change to section 2.1(study area) (line 136-139).
4. Change to result section (3.1 and 3.2).
5. Change to discussion section 4.1-4.4.
6. Change to conclusion.
7. Change to Acknowledgements
8. Change to reference.
9. Change to Table 1.
10. Deleted the Fig.1. (Distribution of typical forest ecosystems)
11. Change to Fig.1, Fig.3, Fig.4.
12. Added Fig.S1 and Fig.S1 in the supporting information.

1 **Divergence of dominant factors on soil microbial**  
2 **communities and functions in forest ecosystems along a climatic**  
3 **gradient**  
4

5 Zhiwei Xu <sup>1,2</sup>, Guirui Yu <sup>3,4,\*</sup>, Xinyu Zhang <sup>3,4,\*</sup>, Nianpeng He <sup>3,4</sup>, Qiufeng Wang <sup>3,4</sup>, Shengzhong  
6 Wang <sup>1,2</sup>, Xiaofeng Xu<sup>5</sup>, Ruili Wang<sup>6</sup>, Ning Zhao<sup>7</sup>

7  
8 1 Institute for Peat and Mire Research, College of Geographical Sciences, Northeast Normal  
9 University, Changchun, 130024, China

10 2 Jilin Provincial Key Laboratory for Wetland Ecological Processes and Environmental Change in  
11 the Changbai Mountains, Changchun, 130024, China

12 3 Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic  
13 Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

14 4 College of Resources and Environment, University of Chinese Academy of Sciences, Beijing,  
15 100190, China

16 5 Biology Department, San Diego State University, San Diego, CA 92182, USA

17 6 College of Forestry, Northwest A&F University, Yangling, Shaanxi Province 712100, China

18 7 Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of  
19 Sciences, Lanzhou 730000, China

20  
21  
22 \* Corresponding author: Key Laboratory of Ecosystem Network Observation and Modeling,  
23 Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences,  
24 Beijing 100101, China. No. 11A, Datun Road, Chaoyang District, Beijing, 100101, China. Tel.:  
25 +86-10-64889268; fax: +86 10 64889432.

26 E-mail: [yugr@igsnr.ac.cn](mailto:yugr@igsnr.ac.cn) (G.Y.), [zhangxy@igsnr.ac.cn](mailto:zhangxy@igsnr.ac.cn) (X.Z.)

27

1 **Abstract.** Soil microorganisms play an important role in regulating nutrient cycling in terrestrial  
2 ecosystems. Most of the studies conducted thus far have been confined to a single forest biome or  
3 have focused on one or two controlling factors, and few have dealt with the integrated effects of  
4 climate, vegetation, and soil substrate availability on soil microbial communities and functions  
5 among different forests. In this study, we used phospholipid-derived fatty acid (PLFA) analysis to  
6 investigate soil microbial community structure, and extracellular enzymatic activities to evaluate  
7 the functional potential of soil microbes of different types of forests in three different climatic zones  
8 along the North-South transect in eastern China (NSTEC). Both climate and forest type had  
9 significant effects on soil enzyme activities and microbial communities with considerable  
10 interactive effects. Except soil acid phosphatase (AP), other three enzyme activities were much  
11 higher in the warm temperate zone than in the temperate and the subtropical climate zones. The soil  
12 total PLFAs and bacteria were much higher in the temperate zone than in the warm temperate and  
13 the subtropical zones. The soil  $\beta$ -glucosidase (BG) and N-acetylglucosaminidase (NAG) activities  
14 were much highest in the coniferous forest. Except the soil fungi and fungi/bacteria (F/B), the  
15 different group of microbial PLFAs were much higher in the conifer broad-leaved mixed forests  
16 than in the coniferous forests and the broad-leaved forests. In general, soil enzyme activities and  
17 microbial PLFAs were higher in primary forests than in secondary forests in temperate and warm  
18 temperate regions. In the subtropical region, soil enzyme activities were lower in the primary forests  
19 than in the secondary forests and microbial PLFAs did not differ significantly between primary and  
20 secondary forests. Different compositions of the tree species may cause variations in soil microbial  
21 communities and enzyme activities. Our results showed that the main controls on soil microbes and  
22 functions vary in different climatic zones, and that the effects of soil moisture content, soil  
23 temperature, clay content, and the soil N/P ratio were considerable. This information will add value  
24 to modeling of microbial processes and will contribute to carbon cycling in large-scale carbon  
25 models.  
26

## 1 **1 Introduction**

2 There is a growing awareness that above- and below-ground interactions make an essential  
3 contribution to ecosystem function (van Dam and Heil, 2011). Variations in soil microbial diversity  
4 and community structure have a strong influence on soil organic matter turnover and may impact  
5 on the function of a given ecosystem (Baumann et al., 2013). For example, mycorrhizal fungi and  
6 nitrogen (N) fixing bacteria are responsible for 80% of all N, and up to 75% of phosphorus (P), that  
7 is acquired by plants annually (van der Heijden et al., 2008). Therefore, it is important to study the  
8 composition and enzyme activities of soil microbial communities to obtain an improved  
9 understanding of the mechanisms that control soil organic carbon dynamics in different forest  
10 ecosystems.

11 Vegetation composition may alter soil physicochemical properties by changing the quantity  
12 and quality of plant litter, which further influence microbial community composition and function  
13 (Ushio et al., 2010). There is increasing evidence that vegetation types influence the structure and  
14 functions of the soil microbial community (Zheng et al., 2015). Differences in microbial  
15 communities, as represented by PLFAs, have also been reported among adjacent maple, beech,  
16 hornbeam, lime, and ash forests in Germany (Scheibe et al., 2015) and among forests of four conifer  
17 species in coastal British Columbia (Grayston and Prescott, 2005). From a functional perspective,  
18 both soil acid phosphatase and b-glucosidase activities were higher in a monsoon evergreen  
19 broadleaf forest than in a Masson pine forest (Zheng et al., 2015). However, vegetation type does  
20 not always have an effect on the composition of the soil microbial community. Hannam et al. (2006)  
21 reported that the microbial community composition of a white spruce-dominated forest differed  
22 substantially from that of an aspen-dominated stand, but was similar to that of a mixed stand with  
23 equivalent proportions of deciduous and coniferous trees. Most of the studies conducted thus far  
24 have been confined to a single forest biome or have focused on one or two controlling factors (Ultra  
25 et al., 2013), and few have dealt with the integrated effects of climate, vegetation, and soil substrate  
26 availability on soil microbial communities and functions in different forest biomes.

27 Soil microbial communities and enzyme activities can be influenced by an array of factors,  
28 such as climate (Xu et al., 2015), vegetation types (Urbanová et al., 2015), plant diversity (Li et al.,  
29 2015), and physico-chemical soil properties (Tripathi et al., 2015). The links between the diversity

1 of plant and soil microbial communities and enzyme activities are widely acknowledged (Chung et  
2 al., 2007). The composition of the vegetation species can be used to successfully predict the soil  
3 microbial community (Mitchell et al., 2010). Soils with different vegetation types develop distinct  
4 physico-chemical properties that will have pronounced effects on the structure and function of the  
5 soil microbial community (Priha and Smolander, 1997). Soil organic matter is related to the  
6 variations in microbial activities and community function (Brockett et al., 2012). Soil pH (Shen et  
7 al., 2013), elemental stoichiometric ratios (Högberg et al., 2007), and nutrient status (Lauber et al.,  
8 2008) have also been identified as determinants of microbial community structure. However, we  
9 still do not know which mechanisms control the variability in the structure and functions of soil  
10 microbial communities within different groups of plant species (broadleaved and coniferous trees)  
11 on similar soil types within the same climatic region.

12 Forest soil microbial community structures and enzyme activities are influenced by different  
13 factors in different climatic zones. For example, Högberg et al. (2007) found that the soil microbial  
14 community composition in a boreal forest was strongly influenced by the soil carbon to nitrogen  
15 ratio (C/N) and the soil pH. Studies in temperate forests have shown that dehydrogenase and urease  
16 were closely related to the mean air temperature, litter production, and nutrient availability (Kang  
17 et al., 2009). In addition, Hackl et al. (2005) reported that soil water availability was responsible for  
18 variability in the microbial community structure of temperate forests. Precipitation and soil moisture  
19 may be important controls on the structure of soil fungal communities of tropical forests (McGuire  
20 et al., 2012). However, there is a lack of well-defined information about the factors that influence  
21 the structure and functions of soil microbial communities in forests with different plant species  
22 (broadleaved and coniferous trees) across a range of climates and soils.

23 The North-South Transect of Eastern China (NSTEC) represents a latitudinal and climatic  
24 gradient. It is a unique belt in which vegetation ranges from boreal forest to tropical rain forest,  
25 depending on the local temperature and precipitation conditions. In this study we examined  
26 variations in the soil microbial communities and their functions in forests comprising different  
27 species (broadleaved and coniferous trees) in temperate, warm temperate, and tropical forest biomes  
28 along the NSTEC. The temperature and precipitation are different in these three climatic zones. We  
29 used information about the soil physico-chemical properties, microbial community structure, and



1 hydrolytic enzyme activities involved in C, N, and P transformations to explore how soil microbial  
2 communities and enzyme activities differed among different forest types in different climatic zones,  
3 and to determine the influence of different environmental variables on the soil microbial  
4 communities and enzyme activities in different climatic zones.

## 5 **2 Materials and methods**

### 6 **2.1 Study area and soil sampling**

7 We chose three study sites, namely Liangshui in Northeast China, Taiyue Mountain in North China,  
8 and Dinghu Mountain in South China, along the North-South Transect in Eastern China (NSTEC)  
9 for field measurements and soil sampling. Both the air temperature and precipitation decrease from  
10 south to north along the NSTEC (Table 1).

11 We examined all the representative forest species in each climatic zone. In Liangshui, on the  
12 Xiao Xing'an Mountain, we sampled primary conifer broad-leaved mixed forest (PCB), secondary  
13 conifer broad-leaved mixed forest (SCBt), and two coniferous plantations, one of which was mainly  
14 *Pinus koraiensis* (PK) while the other was *Larix olgensis* (LOt). On Taiyue Mountain, we sampled  
15 primary deciduous broad-leaved forest (PDB), secondary deciduous broad-leaved forest (SDB), and  
16 two coniferous plantations, one of which was comprised mainly of *Pinus tabulaeformis* (PT) while  
17 the other was mainly *Larix olgensis* (LOw). On Dinghu Mountain, we sampled a primary evergreen  
18 broadleaved forest (*Castanopsis chinensis*, *Cryptocarya chinensis*, *Cryptocarya concinna*,  
19 *Erythrophleum fordii*, and *Cyathea podophylla*), secondary conifer and broadleaf mixed forest  
20 (*Pinus massoniana*, *Schima superba*), a coniferous plantation (*Pinus massoniana*), and an evergreen  
21 broadleaved plantation (*Erythrophleum fordii*) along a successional stage, hereafter referred to as  
22 PEB, SCBs, PM, and EF, respectively. The average temperature of the sampling month was 21.3 °C,  
23 17.4 °C, 27.3 °C with the relative humidity of 78%, 60-65%, 83.5% in LS, TY, and DH, respectively.  
24 The sampling dates are Jul.5 2013, Jul.28 2013, Aug.15 2013 in LS, TY, and DH, respectively. The  
25 primary forests are zonal forests that reflect the regional climate and the others are zonal forests that  
26 reflect the extreme site conditions. Information about the climate, soil classification (Soil Survey  
27 Staff 2010), and soil properties at each site is provided in Table 1.

28 Soil samples were collected at nine sampling sites along the NSTEC in July and August 2013.  
29 Each site had four independent plots in well-drained areas, which covered an area of 30 m × 40 m,

1 and were at least 10 m apart. The vegetation composition of the four plots at each site was similar.  
2 Samples of mineral soil were collected from a depth of 0–10 cm at between 30 and 50 points in each  
3 plot along an S-shape using a custom-made coring device with a diameter of 6 cm. The above-  
4 ground standing biomass, dead plant parts, and litter were removed from each sampling point. These  
5 samples were pooled together as a composite sample. Visible roots and residues were removed and  
6 then the soil fractions of each sample were homogenized.

7 We stored the samples at 4 °C in a portable refrigerator during field sampling. Once returned  
8 to the laboratory, samples were stored at 4 °C before analysis. Soils were analyzed for enzyme  
9 activities and PLFAs in September 2013. The fresh soil samples were sieved through a 2-mm mesh  
10 and were subdivided into three subsamples. One subsample was stored at 4 °C until analyzed for  
11 soil enzyme activities and physical and chemical properties. The second was stored at –20 °C before  
12 analysis for microbial community structures. The third was air dried, and then sieved through a 0.25  
13 mm mesh before SOC, TN, and TP analysis. The soil temperatures were measured *in situ* at the time  
14 of sampling. Soil moisture content (SMC) was measured gravimetrically on 20 g fresh soil that was  
15 oven-dried at 105 °C to constant weight immediately on arrival at the laboratories at the study sites  
16 (Liu et al., 2012).

## 17 **2.2 Soil chemical analyses**

18 Soil pH was measured at a soil-to-water ratio of 1:2.5. Soil total N (TN) concentrations were  
19 determined by dry combustion of ground samples (100-mesh) in a C/N analyzer (Elementar, Vario  
20 Max CN, Germany). The soil organic carbon (SOC) concentrations were determined by dichromate  
21 oxidation and titration with ferrous ammonium sulfate (Huang et al., 2014). The litter total C (litter  
22 TC) and total N (litter TN) were determined with the same method that was used for soil TN. Total  
23 phosphorus (TP) was determined with a flow injection auto-analyzer following digestion with  
24 H<sub>2</sub>SO<sub>4</sub>-HClO<sub>4</sub> (Huang et al., 2011). The soil clay fraction (hereafter referred to as Clay, comprised  
25 of particles <53 µm) was separated by wet-sieving and then freeze-dried (Six, Elliott & Paustian  
26 2000).

## 27 **2.3 Phospholipid fatty-acid and enzyme activity analysis**

28 Samples were analyzed for phospholipid fatty-acids (PLFA) using the method described by Bååth  
29 & Anderson (2003). After mild alkaline methanolysis to form fatty acid methyl esters (FAMES),

1 samples were then dissolved in hexane and analyzed with a DB-5 column in a gas chromatography  
2 mass spectroscopy (GCMS) system (Thermo TRACE GC Ultra ISQ). Total amounts of the different  
3 PLFA biomarkers were used to represent the different groups of soil micro-organisms (Table S1).  
4 Taken together, the combination of bacterial, fungal and actinomycic PLFAs biomarkers represented  
5 the total PLFAs of the soil microbial community.

6 The activities of  $\beta$ -glucosidase (BG), N-acetylglucosaminidase (NAG), acid phosphatase (AP),  
7 and leucine aminopeptidase (LAP) were measured as outlined by Saiya-Cork, Sinsabaugh & Zak  
8 (2002). The microplates were incubated in the dark at 20 °C for 4 h. During the incubation, the  
9 incubation plates were shaken every hour to ensure the reaction mixtures were homogenous.  
10 Fluorescence was measured using a microplate fluorometer with 365-nm excitation and 450-nm  
11 emission filters (Synergy<sup>H4</sup> Hybrid Reader, Synergy<sup>H4</sup> BioTek, USA).

12 **2.4 Statistical analysis**

13 One-way analysis of variance (ANOVA) with a post-hoc Tukey HSD test was used to test the  
14 differences between the soil and microbial properties in the various forests of the three climatic  
15 zones. All data were normality distributed. Two-way analysis was used to test the effect of climate  
16 and vegetation on the soil microbial properties. All ANOVA and two-way analysis were performed  
17 using SPSS 19.0 for Windows. Figures were generated using the Origin 8.0 package. Data are  
18 reported as the mean  $\pm$  SE.

19 Redundancy analysis (RDA) was used to examine the relationships between the litter factors  
20 (litter TC, litter TN, litter C/N), soil biochemical variables (soil temperature (ST), soil moisture  
21 content (SMC), pH, C/N, soil carbon to phosphorus ratio (C/P), soil nitrogen to phosphorus ratio  
22 (N/P), SOC, TN, TP), soil texture (Clay), and the soil microbial community compositions and  
23 enzyme activities. Before redundancy analysis, we conducted forward selection of the  
24 environmental variables that were significantly correlated with variations in the microbial  
25 communities and enzyme activities using stepwise regression and the Monte Carlo Permutation Test  
26 that was similar to the multiple regression analysis. Stepwise regression and RDA were processed  
27 using CANOCO software 4.5 (Ter Braak & Smilauer 2002). The vectors of greater magnitude that  
28 formed smaller angles with an axis were more strongly correlated with that axis.

29 **3 Results**

### 1 3.1 Soil enzyme activities in different vegetation types

2 The soil enzyme activities were generally higher in the primary forests than in the secondary forests  
3 in temperate and warm temperate climatic zones (Fig. 1). However, in the subtropical climatic zone,  
4 soil enzyme activities were higher in the SCBs forest than in the PEB forest. The BG, NAG, and  
5 AP enzymes in the two soils of the PT and LOw in the warm temperate zone were significantly  
6 different (Fig. 1(A, B, D)). The soil BG and NAG activities were much higher in the coniferous  
7 forest than in the conifer broad-leaved mixed forests and the broad-leaved forests (Table S2). The  
8 soil AP enzyme activities were highest in the conifer broad-leaved mixed forests and lowest in the  
9 coniferous forests (Table S2).

10 Climate, a significant influence on the variations of soil enzyme activities ( $P < 0.0001$ ), had  
11 more influence than forest type. The soil BG, NAG, and LAP activities were much higher in the  
12 warm temperate zone than in the temperate and the subtropical climate zones (Table S2). The AP  
13 activities were highest in the subtropical climate zone (Table S2). The effects of climate and forest  
14 type interactions were only significant for soil NAG ( $P < 0.0001$ ) and AP activities ( $P = 0.035$ ) (Table  
15 2, Table S2). Forests within the same climatic zones had similar soil enzyme activities (Fig. S1).

### 16 3.2 Soil microbial community composition in different vegetation types

17 Soil PLFAs were higher in the primary forest in the temperate and warm temperate zones than in  
18 the secondary forest. In the temperate zones, soil PLFAs were higher in the PCB forest than in the  
19 SCBt, PK, and LOt (Fig. 2A). In the warm temperate forests, total soil microbial PLFAs were  
20 highest in the LOw forest (Fig. 2B). In the subtropical zone, total, bacterial, and actinomycic PLFAs  
21 were higher in the PEB and SCBs forests than in the PM and EF forests (Fig. 2C). The forest type  
22 had a significant effect on the soil bacteria, fungi, gram-positive bacteria ( $G^+$ ), and gram-negative  
23 bacteria ( $G^-$ ) PLFAs (Table 2). The soil total PLFAs, bacteria,  $G^+$ ,  $G^-$ , and actinomycete were much  
24 higher in the conifer broad-leaved mixed forests than in the coniferous forests and the broad-leaved  
25 forests (Table S2). The soil fungi was highest in the broad-leaved forest and lowest in the coniferous  
26 forest (Table S2).

27 With the exception of the soil  $G^+ / G^-$ , the effects of the combination of climate and forest type  
28 on all soil PLFAs were significant, and were stronger than the individual effects of either climate or  
29 forest type (Table 2, Table S2). Climate had a significant effect on the total PLFAs, fungi, and  $G^-$

1 ( $P<0.0001$ ) (Table 2). The soil total PLFAs, bacteria,  $G^+$ , and  $G^-$  were much higher in the temperate  
2 zone than in the warm temperate and the subtropical zones (Table S2). The fungi, F/B, and  $G^+/G^-$   
3 were highest in the subtropical zone (Table S2). The soil microbial communities in the different  
4 forests in the three climate zones were generally unique (Fig.4, Fig.S2).

### 5 **3.3 Relationships between soil enzyme activities and soil properties**

6 The variations in the soil enzyme activities in the 12 forests were significantly and positively  
7 correlated with soil nutrient ratios (C/P and N/P), ST, and litter TN ( $P=0.002$ ), but were negatively  
8 correlated with soil pH and TP ( $P=0.002$ ) (Fig.S1). The litter C/N, litter TN, and SMC ( $P=0.002$ )  
9 were the most important influences on the soil enzyme activity variations in the temperate forests,  
10 followed by ST, soil N/P, and soil TP (Fig. 3(A)). In the warm temperate forests, the variations in  
11 the soil enzyme activities were significantly and positively correlated with ST and soil pH ( $P=0.002$ ),  
12 but were negatively correlated with SMC and soil nutrients (TN and SOC) (Fig. 3(B)). In the  
13 subtropical forests, soil enzyme activities were significantly and positively correlated with clay,  
14 SMC, soil TN, and TP ( $P=0.002$ ), followed by soil nutrient ratios (Fig. 3(C)). These results indicate  
15 that the litter inputs, soil micro-climate, and soil texture were the main drivers of variations in the  
16 soil enzyme activities in the temperate, warm temperate, and subtropics, respectively, with ST, pH,  
17 SMC, and soil N/P as additional influences.

### 18 **3.4 Relationships between PLFA profiles and measured soil properties**

19 The variations in the soil microbial communities in the in 12 forests were significantly and positively  
20 correlated with ST, clay content, and soil nutrient ratios (C/P and N/P), TN ( $P=0.002$ ), but were  
21 negatively correlated with litter TC ( $P=0.002$ ) (Fig.S2). In the temperate forests, the variations in  
22 the soil microbial community structure were strongly affected by the litter TN, litter TC, litter C/N,  
23 soil TP, and ST ( $P=0.002$ ) (Fig. 4(A)). In the warm temperate forests, the first axis of the RDA plot  
24 of the soil microbial community structure was significantly and positively correlated with ST  
25 ( $P=0.002$ ), but was negatively correlated with soil N/P, soil TN, soil C/P, and SOC ( $P=0.002$ ) (Fig.  
26 4(B)). In subtropical forests, the variations in the soil microbial community structure were  
27 significantly and positively correlated with litter TC and ST ( $P=0.002$ ), but negatively correlated  
28 with SMC, soil C/P, soil N/P, and soil C/N ( $P=0.002$ ), followed by the soil TN and clay contents  
29 (Fig. 4(C)). The litter C/N was the main influences on the variations in the soil microbial

1 communities in the temperate, and the soil N/P was the main influences in the warm temperate and  
2 subtropical forests. The microbial communities were also influenced by ST, pH, SMC.

### 3 **4 Discussion**

#### 4 **4.1 Response of soil enzyme activities and microbial PLFAs to variations in forest type**

5 Forests in the same climate zone developed similar microbe functions which confirmed the result  
6 that the effect of climate on soil enzyme activities were stronger than the forest type and their  
7 interactive effect. However, there were still differences among the enzyme activities in different  
8 forest types of the same climate zone. Soil microorganisms are usually considered to be C limited,  
9 and the litter inputs with high C/N ratio of PCB in the temperate zone will stimulate microbes to  
10 grow and secrete more enzymes (Table 1). Therefore, all enzyme activities were highest in PCB in  
11 the temperate zone. The high soil BG enzyme activities in the LOw forest in the warm temperate  
12 zone reflect the litter inputs with low C. Because that soil enzyme activities will not continuously  
13 increase or decrease as nutrient availability increases or decreases. When the soil nutrients are short  
14 in supply, microbes will potentially increase production of nutrient-acquiring enzymes, because they  
15 are expected to optimize the allocation of their resource reserves by acquiring the resource that is  
16 most limiting (Bloom et al., 1985). (Table 1). The soil enzyme activities were highest in the SCBs  
17 forest, reflecting the higher soil nutrient concentrations in subtropical zones.

18 The interactive effect of climate and forest type were more important than the individual effect  
19 of them. Therefore, the soil microbial communities of the 12 forests were separated from each other.  
20 Vegetation transfers substrate material of varying quality to microbes through litter fall. Fungi are  
21 more suitable for life in environments containing higher C/N ratios and low soil pH (Nilsson et al.,  
22 2012). The four broadleaved forests were high in litter C/N ratio (Table 1). Therefore, fungi were  
23 dominated in this harsh nutrient environments and highest in broadleaved forests. The litter and soil  
24 from conifer broad-leaved mixed forest were high in C, N, and P, and promotes the propagation of  
25 bacteria that favor high-nutrient soil (Priha and Smolander, 1997; Priha et al., 2001). Therefore, the  
26 structures and functions of the soil microbial communities that developed in the different types of  
27 forest were unique.

#### 28 **4.2 Common influences on soil enzyme activities and microbial communities**

29 Many other studies have reported how different factors determine the response of the soil microbial

1 community and function to variations in forests (Högberg et al., 2007; McGuire et al., 2012). Mostly  
2 limited to one climatic zone, these studies were quite diverse and featured a range of microbial  
3 methods, sampling times, and environmental properties, which means it is difficult to compare the  
4 results. In this study, we collected the samples at the same times and used the same methods to  
5 analyze the soil microbial communities and enzyme activities. We found that ST, SMC, soil pH, and  
6 soil N/P ratio influenced, but perhaps did not dominate, the responses of the soil microbial  
7 community structures and enzyme activities in the different forest types across the three climatic  
8 zones.

9 Temperature can influence enzyme activity directly and indirectly by modifying the enzyme  
10 kinetics and influencing the proliferation of microbes, respectively (Kang et al., 2009). By changing  
11 the quality and quantity of the substrate on which microbes function, soil moisture is an important  
12 driver of the overall microbial composition and soil microbial function (Hackl et al., 2005). The  
13 responses of soil enzyme activities and microbial communities in the various forest types were all  
14 significantly influenced by the SMC in the three climatic zones. Increases in soil moisture can  
15 enhance both the release and the diffusion rates of enzymes, substrates, and reaction products (Burns  
16 et al., 2013), and our results showed that soil enzyme activities and microbial PLFAs increased as  
17 the SMC increased in the warm temperate and subtropical zones. However, water-logged conditions  
18 are not suitable for microbes and are not beneficial for the release of soil enzymes (Lucas-Borja et  
19 al., 2012), and, similar to other studies, soil enzyme activities and SMC were negatively correlated  
20 in the temperate zone forests (Brockett et al., 2012). As the SMC increases, the bacterial PLFAs  
21 increase (Myers et al., 2001) and fungal PLFAs decrease (Staddon et al., 1998), which indicates that  
22 the soil microbial communities and enzyme activities in the different climatic zones were all  
23 influenced by the soil micro-climate. This was also demonstrated by the stronger effect of climate  
24 on soil enzyme activities and the combined interaction effect of climate and forest type on soil  
25 microbial communities. Other studies have reported that precipitation and mean annual temperature  
26 played important roles in explaining on the large-scale distribution of soil microbial community  
27 composition and functions (de Vries et al., 2012; Xu et al., 2017).

28 Soil pH directly affects the activities of extracellular enzymes immobilized in the soil matrix,  
29 and the effect of soil pH on the soil microbial community and function reflects the influence of

1 vegetation through changes in soil chemistry. Every enzyme has a well-defined optimal soil pH  
2 value (Sinsabaugh et al., 2008) that results from different levels of soil enzyme activities under  
3 different soil pH conditions. Soil  $G^+/G^-$  ratios were highest in the subtropical forest where  $G^-$   
4 bacteria PLFAs were least abundant, which may reflect microbial growth strategies. The  $G^+$  bacteria  
5 are primarily K-strategists that can survive over long periods in the soil under harsh conditions with  
6 lower soil pH (Andrews & Hall, 1986). Increased pH causes an increase in bacterial diversity and a  
7 shift in the bacterial community to more  $G^-$  and fewer  $G^+$  bacteria PLFAs (Wu et al., 2009; Shen et  
8 al., 2013).

9 **4.3 Key influences on soil enzyme activities and microbial communities**

10 Our results showed that the most important controls on the responses of soil microbial communities  
11 and enzyme activities to vegetation types varied across climatic zones. The litter quality and quantity  
12 contribute to the maintenance of soil fertility in forest ecosystems (Wang et al., 2011). In our study,  
13 and the C/N ratios were highest, in litter from PCB stands (Table 1), which shows that the soil in  
14 the PCB was more N-limited than the other soils because of litter inputs with high C/N ratios (Table  
15 1). Therefore, the microbial N demand was highest in soil in the PCB forest, which resulted in higher  
16 NAG and LAP values. Plant litter has a strong influence on soil microbial composition and activity,  
17 as the litter decomposition process provides nutrients for microorganism growth through inputs of  
18 leaf litter (Attiwill and Adams, 1993), dying roots (Silver and Miya, 2001), and root secretion  
19 (Grayston et al., 1997). The litter from the mixed forests, represented in our study by PCB, is more  
20 diverse than that from the pure forests, and so a wider variety of soil microbes participate in the  
21 decomposition process, so that the soil organic matter is richer, and there are more soil microbial  
22 PLFAs, than in the other forest types. Fungi typically dominate N-limited environments and the  
23 fungal biomass is positively related to the C/N ratio (Nilsson et al., 2012). The fungi/bacteria ratio  
24 (F/B ratio) was therefore highest in the PCB forest where the litter C/N values were highest.

25 Microbes obtain the nutrients they need to construct biomass by decomposing soil organic  
26 matter. Wallenius et al. (2011) found that the soil bacterial biomass was higher in forests where the  
27 soil organic matter concentrations were higher than in forests with low soil organic matter  
28 concentrations, and Xu et al. (2017) found positive relationships between soil enzyme activities and  
29 SOC and TN concentrations along the NSTEC. In line with the resource limitation model, and also



1 confirmed by several other studies (Brockett et al., 2012), Schimel and Weintraub (2003) suggested  
2 that increases in N and C substrate availability might favor enzyme synthesis. Soil microorganisms  
3 however did not grow when the available P concentrations in soil were less than  $0.7 \text{ mg kg}^{-1}$  and  
4 were stimulated by P additions (Zheng et al., 2009). Other studies have reported that P additions  
5 stimulated the different PLFA microbial groups in soils (Dong et al., 2015). The soil TN and TP  
6 were lower in the warm temperate and subtropical zone than in the temperate zone in our study  
7 (Table 1), and these two kinds of nutrients were more likely limiting factors in warm temperate and  
8 subtropical forest (DeForest et al., 2012; Xu et al., 2017). Therefore, soil TN and TP are more  
9 important in warm temperate and subtropical forests than in temperate forests.

10 The soil N/P ratio was the most important influence on the soil microbial communities and  
11 enzyme activities in the warm temperate and subtropical zone, which is consistent with the results  
12 of previous studies (Shen et al., 2013; Högberg et al., 2007). Soil stoichiometric C, N, and P ratios  
13 reflect the nutrient limitations of the ecosystems (Sterner and Elser, 2002) and should indicate soil  
14 organic matter mineralization and sequestration (Gundersen et al., 1998). Soil microorganisms  
15 obtain C, N, and P in such a way that enzyme release corresponds with the soil stoichiometric ratios  
16 of C, N, and P. When supplies of N or P are limited, the activities of the enzymes that are responsible  
17 for nitrate or phosphate mineralization will be higher. Consistent with this discussion, soil enzyme  
18 activities in subtropical forests (DH) responded positively to the soil C/N and N/P ratios.

19 Soil texture is a key property that affects the accessibility of organic matter to microbes, and  
20 is an important determinant of soil moisture, and nutrient availability and retention (Veen and  
21 Kuikman, 1990). Consistent with our results, Lagomarsino et al. (2012) reported that the activities  
22 of soil BG, AP, and NAG were higher in silt and clay fractions than in coarser fractions. This may  
23 be attributed to the presence of clay-humus-enzyme complexes in the finest soil fractions, and  
24 implies that physical protection affects soil enzyme activities. In addition, fine textured soils with  
25 higher silt and clay contents are known to be more conducive to bacterial growth than coarser soils  
26 because they have a greater water-holding capacity, higher nutrient availability, and offer better  
27 protection against bacterial grazers (Carson et al., 2010). Therefore, soil enzyme activities and  
28 microbial PLFAs were highest in the SCBs forest with finely texture. Except SCBt in the temperate  
29 zone and PT in the warm temperate zone, the soil clay content were not significant different among

1 other three forest types. However, the soil clay contents of the four forest types in the subtropical  
2 zone were significant different from each other and important for variations in microbial  
3 communities and functions (Table 1).

#### 4 **4.4 Implications for ecosystem modeling**

5 There is increasing recognition that, to improve climate models, microbial processes should be  
6 simulated (DeLong et al., 2011). As such, this study has three important implications. First,  
7 microbial datasets that have information about enzyme activities and soil microbial properties  
8 contribute to improved parameterization of ecosystem models (Xu et al., 2017). Information about  
9 the spatial patterns of, and factors that control, microbial properties and enzymatic activities can  
10 enrich the datasets that are used to parameterize models of microbial processes (Wang et al., 2013).

11 Secondly, knowledge about microbial community structure and its environmental controls can give  
12 a better understanding of how microbes adapt to changing environments, which is the main direction  
13 of model development (Schimel and Schaeffer, 2012). Information about edaphic controls on  
14 microbial processes is critical for developing new modeling frameworks with improved links with  
15 field experimental data (Abramoff et al., 2017). Finally, the information generated in this study  
16 about the divergence of the dominant factors that control soil microbial properties across forests is  
17 extremely valuable for improving our understanding of soil microbial ecology and forest  
18 management.

#### 19 **5 Conclusions**

20 In this study, we characterized the soil microbial communities and enzyme activities and factors that  
21 controlled them in various forest types across three different climatic zones. We found that forest  
22 types with specific soil conditions supported the development of distinct soil microbial communities  
23 with variable functions. The soil total PLFAs, bacteria, G<sup>+</sup>, G<sup>-</sup>, and actinomycete were much higher  
24 in the conifer broad-leaved mixed forests than in the coniferous forests and the broad-leaved forests.  
25 The soil BG and NAG activities were much higher in the coniferous forest than in the conifer broad-  
26 leaved mixed forests and the broad-leaved forests. Except AP, soil enzyme activities were highest  
27 in warm temperate zone. Soil tPLFAs, bacteria, G<sup>-</sup> increased from temperate zone to subtropical  
28 zone, but fungi was in reverse. The litter TN, soil temperature, and soil clay contents were important  
29 predictors of the variance in soil enzyme activities in temperate, warm temperate, and subtropical

1 zones, respectively, while litter and soil nutrient ratios were significant predictors of the variance in  
2 soil microbial communities. We also found that SMC, soil temperature, soil pH, and the soil N/P  
3 ratio were common drivers of variations in the soil microbial community structure and enzyme  
4 activities across the different forest types in the three climatic zones. Forests within the same  
5 climatic zones had similar soil microbial communities and enzyme activities, and these patterns  
6 were mainly determined by the litter input, soil micro-environment, and soil nutrient ratios. The data  
7 in this study is extremely valuable for improving our understanding of soil microbial ecology and  
8 forest management.

9 *Data accessibility.* Requests for data and materials should be addressed to N.H. (henp@igsnr.ac.cn) and G.Y.  
10 ([yugr@igsnr.ac.cn](mailto:yugr@igsnr.ac.cn)).

11  
12 *Author contributions.* Z.W.X., G.R.Y. and X.Y.Z. planned and designed the research. Z.W.X., N.P.H., R.L.W., N.Z.,  
13 C.C.J., and C.Y.W. conducted fieldwork. Z.W.X., G.R.Y., X.Y.Z. Q.F.W., S.Z.W. and X.F.X wrote the manuscript.  
14 All authors contributed critically to the drafts and gave final approval for publication.

15  
16 *Competing interests.* The authors declare that they have no conflict of interest.

17  
18 *Acknowledgements.* This research was jointly supported by the Key Program of the National Natural Science  
19 Foundation of China (31290221), the National Natural Science Foundation of China (41601084, 41571251), and the  
20 Fundamental Research Funds for the Central Universities (2412016KJ029). X.X. was grateful for the financial  
21 support from the San Diego State University and the Oak Ridge National Laboratory.

22

## 23 **References**

- 24 Abramoff, R., Xu, X., Hartman, M., O'Brien, S., Feng, W. T., Davidson, E. A., Finzi, A., Moorhead, D. L., Schimel,  
25 J. P., Torn, M. S., and Mayes, M. A.: The millennial model: in search of measurable pools and transformations  
26 for modeling soil carbon in the new century, *Global Biogeochem. Cy.* ( In review), 2017.  
27 Andrews, J. H., Hall, R. F., 1986. R- and K-selection and microbial ecology. *Advan. Microb. Ecol.* 9, Springer US,  
28 pp 99-147.  
29 Attiwill, P.M. & Adams, M.A., 1993. Nutrient cycling in forests. *New Phytol.*, 124, 561-582.  
30 Bååth, E. and Anderson, T.H.: Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and  
31 PLFA-based techniques, *Soil Biol. Biochem.*, 35, 955-963, 2003.  
32 Baumann, K., Dignac, M-F., Rumpel, C., Bardoux, G., Sarr, A., Steffens, M., and Maron, P. A.: Soil microbial  
33 diversity affects soil organic matter decomposition in a silty grassland soil, *Biogeochemistry*, 114, 201-212, 2013.  
34 Bloom, A., Chapin, F.I., Mooney, H.: Resource limitation in plants-an economic analogy, *Annu. Rev. Ecol. S.*, 16,  
35 363-392, 1985.

1 Brockett, B.F.T., Prescott, C.E., and Grayston, S.J.: Soil moisture is the major factor influencing microbial  
2 community structure and enzyme activities across seven biogeoclimatic zones in western Canada, *Soil Biol.*  
3 *Biochem.*, 44, 9-20, 2012.

4 Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein, M.D., Weintraub, M.  
5 N., and Zoppini, A.: Soil enzymes in a changing environment: Current knowledge and future directions, *Soil*  
6 *Biol. Biochem.*, 58, 216-234, 2013.

7 Carson, J.K., Gonzalez-Quinones, V., Murphy, D.V., Hinz, C., Shaw, J.A., and Gleeson, D.B.: Low pore connectivity  
8 increases bacterial diversity in soil, *Appl. Environ. Microbiol.*, 76, 3936-3942, 2010.

9 Chung, H., Zak, D.R., Reich, P.B., and Ellsworth, D.S.: Plant species richness, elevated CO<sub>2</sub>, and atmospheric  
10 nitrogen deposition alter soil microbial community composition and function, *Global Change Biol.*, 13, 980-989,  
11 2007.

12 van Dam, N.M. and Heil, M.: Multitrophic interactions below and above ground: en route to the next level, *J. Ecol.*,  
13 99, 77-88, 2011.

14 DeLong, E. F., Harwood, C. S., Chisholm, P. W., Karl, D. M., Moran, M. A., Schmidt, T. M., Tiedje, J. M., Treseder,  
15 K. K., and Worden, A. Z.: Incorporating microbial processes into climate models, *Am. Acad. Microbiol.*, 2011.

16 Grayston, S.J. and Prescott, C.E.: Microbial communities in forest floors under four tree species in coastal British  
17 Columbia, *Soil Biol. Biochem.*, 37, 1157-1167, 2005.

18 Grayston, S.J., Vaughan, D., and Jones, D.: Rhizosphere carbon flow in trees, in comparison with annual plants: the  
19 importance of root exudation and its impact on microbial activity and nutrient availability, *Appl. Soil Ecol.*, 5,  
20 29-56, 1997.

21 Gundersen, P., Callesen, I., and Vries, W.D.: Nitrate leaching in forest ecosystems is related to forest floor C/N ratios,  
22 *Environ. Pollut.*, 102, 403-407, 1998.

23 Hannam, K., Quideau, S., and Kishchuk, B.: Forest floor microbial communities in relation to stand composition  
24 and timber harvesting in northern Alberta, *Soil Biol. Biochem.*, 38, 2565-2575, 2006.

25 Högberg, M.N., Högberg, P., and Myrold, D.D.: Is microbial community composition in boreal forest soils  
26 determined by pH, C-to-N ratio, the trees, or all three, *Oecologia*, 150, 590-601, 2007.

27 Hackl, E., Pfeffer, M., Donat, C., Bachmann, G., and Zechmeister-boltenstern, S.: Composition of the microbial  
28 communities in the mineral soil under different types of natural forest, *Soil Biol. Biochem.*, 37, 661-671, 2005.

29 van der Heijden, M.G., Bardgett, R.D., and van Straalen, N.M.: The unseen majority: soil microbes as drivers of  
30 plant diversity and productivity in terrestrial ecosystems, *Ecol. Lett.*, 11, 296-310, 2008.

31 Huang, X., Liu, S., Wang, H., Hu, Z., Li, Z., and You, Y.: Changes of soil microbial biomass carbon and community  
32 composition through mixing nitrogen-fixing species with *Eucalyptus urophylla* in subtropical China, *Soil Biol.*  
33 *Biochem.*, 73, 42-48, 2014.

34 Huang, Z., Clinton, P.W., Baisden, W.T., and Davis, M.R.: Long-term nitrogen additions increased surface soil  
35 carbon concentration in a forest plantation despite elevated decomposition, *Soil Biol. Biochem.*, 43, 302-307,  
36 2011.

37 Lagomarsino, A., Grego, S., and Kandeler, E.: Soil organic carbon distribution drives microbial activity and  
38 functional diversity in particle and aggregate-size fractions, *Pedobiologia*, 55, 101-110, 2012.

39 Lauber, C.L., Strickland, M.S., Bradford, M.A., and Fierer, N.: The influence of soil properties on the structure of  
40 bacterial and fungal communities across land-use types, *Soil Biol. Biochem.*, 40, 2407-2415, 2008.

41 Li, H., Wang, X., Liang, C., Hao, Z., Zhou, L., Ma, S., Li, X. B., Yang, S., Yao, F., and Jiang, Y.: Aboveground-  
42 belowground biodiversity linkages differ in early and late successional temperate forests, *Sci Rep-UK*, 5, 12234,  
43 2015.

44 Liu, L., Gundersen, P., Zhang, T., and Mo, J. M.: Effects of phosphorus addition on soil microbial biomass and  
45 community composition in three forest types in tropical China, *Soil Biol. Biochem.*, 44, 31-38, 2012.

46 Lucas-Borja, M.E., Candel Pérez, D., López Serrano, F.R., Andrés, M., and Bastida, F.: Altitude-related factors but  
47 not *Pinus* community exert a dominant role over chemical and microbiological properties of a Mediterranean  
48 humid soil, *Eur. J. Soil Sci.*, 63, 541-549, 2012.

49 McGuire, K.L., Fierer, N., Bateman, C., Treseder, K.K., and Turner, B.L.: Fungal community composition in  
50 neotropical rain forests: the influence of tree diversity and precipitation, *Microb. Ecol.*, 63, 804-812, 2012.

51 Mitchell, R.J., Hester, A.J., Campbell, C.D., Chapman, S.J., Cameron, C.M., Hewison, R.L., and Potts, J. M.: Is  
52 vegetation composition or soil chemistry the best predictor of the soil microbial community, *Plant Soil*, 333, 417-  
53 430, 2010.

54 Myers, R.T., Zak, D.R., White, D.C., and Peacock, A.: Landscape-level patterns of microbial community  
55 composition and substrate use in upland forest ecosystems, *Soil Sci. Soc. Am. J.*, 65, 359-367, 2001.

56 Nilsson, L.O., Wallander, H., and Gundersen, P.: Changes in microbial activities and biomasses over a forest floor  
57 gradient in C-to-N ratio, *Plant Soil*, 355, 75-86, 2012.

58 Priha, O., Grayston, S.J., Hiukka, R., Pennanen, T., and Smolander, A.: Microbial community structure and  
59 characteristics of the organic matter in soils under *Pinus sylvestris*, *Picea abies* and *Betula pendula* at two forest  
60 sites, *Biol. Fert. Soils*, 33, 17-24, 2001.

61 Priha, O. and Smolander, A.: Microbial biomass and activity in soil and litter under *Pinus sylvestris*, *Picea abies* and  
62 *Betula pendula* at originally similar field afforestation sites, *Biol. Fert. Soils*, 24, 45-51, 1997.

63 Saiya-Cork, K.R., Sinsabaugh, R.L., and Zak, D.R.: The effects of long term nitrogen deposition on extracellular  
64 enzyme activity in an *Acer saccharum* forest soil, *Soil Biol. Biochem.*, 34, 1309-1315, 2002.

65 Scheibe, A., Steffens, C., Seven, J., Jacob, A., Hertel, D., Leuschner, C., and Gleixner, G.: Effects of tree identity  
66 dominate over tree diversity on the soil microbial community structure, *Soil Biol. Biochem.*, 81, 219-227, 2015.

67 Schimel, J. P. and Schaeffer, S. M.: Microbial control over carbon cycling in soil, *Front. Microbiol.*, 3, 1-11, 2012.

- 1 Schimel, J.P. and Weintraub, M.N.: The implications of exoenzyme activity on microbial carbon and nitrogen  
2 limitation in soil: a theoretical model, *Soil Biol. Biochem.*, 35, 549-563, 2003.
- 3 Shen, C., Xiong, J., Zhang, H., Feng, Y., Lin, X., Li, X., Liang, W. J., and Chu, H. Y.: Soil pH drives the spatial  
4 distribution of bacterial communities along elevation on Changbai Mountain, *Soil Biol. Biochem.*, 57, 204-211,  
5 2013.
- 6 Silver, W.L. and Miya, R.K.: Global patterns in root decomposition: comparisons of climate and litter quality effects,  
7 *Oecologia*, 129, 407-419, 2001.
- 8 Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C. Contosta, A.R., Cusack,  
9 D., Frey, S., Gallo, M. E., Gartner, T. B., Hobbie, S. E., Holland, K.i, Keeler, B. L., Powers, J. S., Stursova,  
10 Takacs-Vesbach, C., Waldrop, M.P., Wallenstein, M. D., Zak, D. R., and Zeglin, L. H.: Stoichiometry of soil  
11 enzyme activity at global scale, *Ecol. Lett.*, 11, 1252-1264, 2008.
- 12 Six, J., Elliott, E.T., and Paustian, K.: Soil structure and soil organic matter: II. A normalized stability index and the  
13 effect of mineralogy, *Soil Sci. Soc. Am. J.*, 64, 1042-1049, 2000.
- 14 Soil Survey Staff: Keys to Soil Taxonomy, 11th ed. Natural Resources Conservation Service, Washington, DC, 2010.
- 15 Staddon, W.J., Trevors, J.T., Duchesne, L.C., and Colombo, C.A.: Soil microbial diversity and community structure  
16 across a climatic gradient in western Canada, *Biodivers. Conserv.*, 7, 1081-1092, 1998.
- 17 Sterner, R. W. and Elser, J. J.: Ecological stoichiometry: the biology of elements from molecules to the biosphere,  
18 Princeton University Press, 2002.
- 19 Ter Braak, C.J.F., and Smilauer, P.: CANOCO Reference manual and CanoDraw for Windows User's guide:  
20 Software for Canonical Community Ordination (Version 4.5), 2002.
- 21 Tietema, A.: Microbial carbon and nitrogen dynamics in coniferous forest floor material collected along a European  
22 nitrogen deposition gradient, *Forest Ecol. Manag.*, 101, 29-36, 1998.
- 23 Tripathi, B.M., Kim, M., Tateno, R., Kim, W., Wang, J., Lai-Hoe, A, Shukor, N.A., Rahim, R.A., Go, R., and Adams,  
24 J. M.: Soil pH and biome are both key determinants of soil archaeal community structure, *Soil Biol. Biochem.*,  
25 88, 1-8, 2015.
- 26 Ultra, V. U., Han, S.-H., and Kim, D.-H.: Soil properties and microbial functional structure in the rhizosphere of  
27 *Pinus densiflora* (S. and Z.) exposed to elevated atmospheric temperature and carbon dioxide, *J. Forest Res.*, 18,  
28 149-158, 2013.
- 29 Urbanová, M., Šnajdr, J., and Baldrian, P.: Composition of fungal and bacterial communities in forest litter and soil  
30 is largely determined by dominant trees, *Soil Biol. Biochem*, 84, 53-64, 2015.
- 31 Ushio, M., Kitayama, K., and Balsler, T.C.: Tree species effects on soil enzyme activities through effects on soil  
32 physicochemical and microbial properties in a tropical montane forest on Mt. Kinabalu, Borneo, *Pedobiologia*,  
33 53, 227-233, 2010.
- 34 Veen, J.A.V. and Kuikman, P.J.: Soil structural aspects of decomposition of organic-matter by micro-organism.  
35 *Biogeochemistry*, 11, 213-233, 1990.
- 36 Wang, C., Han, G., Jia, Y., Feng, X., Guo, P., and Tian, X.: Response of litter decomposition and related soil enzyme  
37 activities to different forms of nitrogen fertilization in a subtropical forest, *Ecol. Res.*, 26, 505-513, 2011.
- 38 Wang, G., Post, W. M., and Mayes, M. A.: Development of microbial-enzyme-mediated decomposition model  
39 parameters through steady-state and dynamic analyses, *Ecol. Appl.*, 23, 255-272, 2013.
- 40 Wu, Y., Ma, B., Zhou, L., Wang, H., Xu, J., Kemmitt, S., and Brookes, P. C.: Changes in the soil microbial community  
41 structure with latitude in eastern China, based on phospholipid fatty acid analysis, *Appl. Soil Ecol.*, 43, 234-240,  
42 2009.
- 43 Xu, X., Thornton, P. E., and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus  
44 in terrestrial ecosystems, *Global Ecol. Biogeogr.*, 22, 737-749, 2013.
- 45 Xu, Z., Yu, G., Zhang, X., Ge, J., He, N., Wang, Q., and Wang, D.: The variations in soil microbial communities,  
46 enzyme activities and their relationships with soil organic matter decomposition along the northern slope of  
47 Changbai Mountain, *Appl. Soil Ecol.*, 86, 19-29, 2015.
- 48 Xu, Z.W., Yu, G.R., Zhang, X.Y., He, N.P., Wang, Q.F., Wang, S.Z., Wang, R.L., Zhao, N., Jia, Y.L., and Wang, C.Y.:  
49 Soil enzyme activity and stoichiometry in forest ecosystems along the North-South Transect in eastern China  
50 (NSTEC), *Soil Biol. Biochem.*, 104, 152-16, 2017.
- 51 Zheng, M., Huang, J., Chen, H., Wang, H., and Mo, J.: Responses of soil acid phosphatase and beta-glucosidase to  
52 nitrogen and phosphorus addition in two subtropical forests in southern China, *Eur. J. Soil Biol.*, 68, 77-84, 2015.

## 1 Figure captions

2 **Figure 1.** Soil enzyme activities under different forest types in different climatic zones. BG, b-1, 4-glucosidase; NAG,  
3 b-1,4-N-acetylglucosaminidase; LAP, leucine aminopeptidase; AP, acid phosphatase. **The capital letters A, B, C,**  
4 **and D represent the variations in the enzyme activities of BG, NAG, LAP and AP, respectively.** Different lowercase  
5 letters indicate significant differences between forests in the same climatic zone. The abbreviations of the sampling  
6 sites are shown in Table 1.

7 **Figure 2.** The PLFA contents, Fungi:Bacteria ratios, and  $G^+/G^-$  for different forest types in different climatic zones  
8 (A. Liangshui; B. Taiyue; C. Dinghu). Different lowercase letters indicate significant differences among forests in  
9 the same climatic zone. F/B, fungi/bacteria;  $G^+/G^-$ , Gram-positive bacteria/ Gram-negative bacteria. The  
10 abbreviations of the sampling sites are shown in Table 1.

11 **Figure 3.** Redundancy analysis (RDA) ordination biplot of soil enzyme activities and environmental properties for  
12 the different forest types in different climatic zones (A. Liangshui; B. Taiyue; C. Dinghu). Only the environmental  
13 variables that were significantly correlated with RDA1 are shown. The dotted lines and solid lines represent the  
14 environmental variables and enzyme activities. The variables in this table were abbreviated as follows: TC(litter) =  
15 litter total carbon; TN(litter) = litter total nitrogen; C/N(litter) = litter total carbon/nitrogen; ST = soil temperature;  
16 SMC = soil moisture content; Clay = soil clay content; SOC = soil organic carbon; TN = soil total nitrogen; TP =  
17 soil total phosphorus; C/N = soil carbon/nitrogen; C/P = soil carbon/phosphorus, and N/P = soil nitrogen/phosphorus.

18 **Figure 4.** Redundancy analysis (RDA) ordination biplot of soil microbial community structure and environmental  
19 properties for different forest types in different climatic zones (A. Liangshui; B. Taiyue; C. Dinghu). Only the  
20 environmental variables that were significantly correlated with RDA1 are shown. The dotted lines and solid lines  
21 represent the environmental variables and lipid signatures. The abbreviations of the variables included in this figure  
22 are shown in Figure 4.

## 23 Supporting Information

24 **Table S1.** The PLFA biomarkers used to represent the different groups of soil micro-organisms (Frostegård *et*  
25 *al.*1996).

26 **Table S2.** Average values of soil enzyme activities and microbial PLFAs in the three different climatic zones and  
27 three different forest types, respectively.

28 **Figure S1.** Redundancy analysis (RDA) ordination biplot of soil enzyme activities and environmental properties for  
29 the 12 forests.

30 **Figure S2.** Redundancy analysis (RDA) ordination biplot of soil microbial community structure and environmental  
31 properties for the 12 forests.

1 Tables  
2

**Table 1.** Stand characteristics and soil properties under different forest types in the three climatic zones

| Areas <sup>a</sup>           | XiaoXing'an Mountain (LS) |         |         |         | Taiyue Mountain (TY) |         |         |         | Dinghu Mountain (DH) |         |         |         |
|------------------------------|---------------------------|---------|---------|---------|----------------------|---------|---------|---------|----------------------|---------|---------|---------|
| Sampling date                | Jul.5 2013                |         |         |         | Jul.28 2013          |         |         |         | Aug.15 2013          |         |         |         |
| Latitude(°)                  | 47.19                     |         |         |         | 36.70                |         |         |         | 23.17                |         |         |         |
| Longitude(°)                 | 128.90                    |         |         |         | 112.08               |         |         |         | 112.54               |         |         |         |
| Climatic zone                | Temperate                 |         |         |         | Warm temperate       |         |         |         | Subtropical          |         |         |         |
| MAT (°C)                     | 0.3                       |         |         |         | 6.2                  |         |         |         | 20.9                 |         |         |         |
| MAP (mm)                     | 676                       |         |         |         | 662                  |         |         |         | 1927                 |         |         |         |
| Altitude (m)                 | 401                       |         |         |         | 1668                 |         |         |         | 240                  |         |         |         |
| Soil type                    | Cryumbrept                |         |         |         | Eutrochrepts         |         |         |         | oxisol               |         |         |         |
| Vegetation type <sup>b</sup> | PCB(M)                    | SCBt(M) | PK(C)   | LOt(C)  | PDB(B)               | SDB(B)  | PT(C)   | LOw(C)  | PEB(B)               | SCBs(M) | PM(C)   | EF(B)   |
| pH                           | 6.17a                     | 5.68b   | 6.01a   | 6.28a   | 6.85c                | 7.70a   | 7.20b   | 6.78c   | 5.43a                | 5.38a   | 5.21b   | 5.07b   |
| ST (°C)                      | 15.87a                    | 15.11b  | 15.33b  | 16.13a  | 16.00b               | 24.04a  | 16.37a  | 15.33b  | 24.40b               | 24.59b  | 25.34a  | 25.39a  |
| SMC (%)                      | 46.94c                    | 69.97a  | 50.7b   | 57.95c  | 36.01a               | 22.66c  | 27.89b  | 34.87a  | 37.84b               | 44.76a  | 26.67b  | 30.20b  |
| Clay (%)                     | 63.98a                    | 55.92b  | 64.57a  | 64.30a  | 49.39a               | 52.13a  | 35.69b  | 53.90a  | 49.74b               | 76.05a  | 45.05d  | 52.31c  |
| SOC (g kg <sup>-1</sup> )    | 62.08a                    | 75.23a  | 61.47a  | 57.10a  | 41.34a               | 17.87b  | 42.72a  | 42.15a  | 28.47c               | 40.03a  | 26.83c  | 37.99b  |
| TN (g kg <sup>-1</sup> )     | 4.59a                     | 4.57a   | 4.01a   | 4.54a   | 2.43b                | 1.41c   | 3.09a   | 2.79a   | 1.77b                | 2.55a   | 1.26c   | 1.83b   |
| TP (g kg <sup>-1</sup> )     | 0.59b                     | 0.78a   | 0.83a   | 0.94a   | 0.52b                | 0.51b   | 0.56a   | 0.52b   | 0.20c                | 0.26a   | 0.23b   | 0.22b   |
| Litter TC                    | 460.50b                   | 489.66a | 476.48b | 414.26c | 507.47a              | 456.64b | 509.65a | 435.00c | 422.65c              | 451.69b | 521.11a | 520.51a |
| Litter TN                    | 10.87c                    | 20.23a  | 14.86b  | 16.10b  | 10.38b               | 12.23a  | 9.59b   | 13.97a  | 14.1c                | 16.38b  | 17.25a  | 17.38a  |
| Litter C/N                   | 43.11a                    | 24.03c  | 31.96b  | 25.54c  | 48.56a               | 37.82b  | 53.16a  | 30.82c  | 28.67a               | 27.06a  | 30.31a  | 29.85a  |

3 <sup>a</sup> PCB, SCBt, PK, and LOt represent primary conifer broad-leaved mixed forest, secondary conifer broad-leaved mixed forest, *Korean pine* forest and *Larix olgensis* forest, respectively. PDB,  
4 SDB, PT, and LOw represent primary deciduous broad-leaved forest, secondary deciduous broad-leaved forest, *Pinus tabulaeformis* forest and *Larix olgensis* forest, respectively. PEB, SCBs,  
5 PM, and EF represent primary evergreen broadleaved forest, secondary conifer and broadleaf mixed forest, *Pinus massoniana* forest and *Erythrophleum fordii* forest, respectively. The letters in  
6 the bracket after the vegetation type represent M, conifer broad-leaved mixed forest; C, coniferous forest; B, broad-leaved forest. MAT and MAP indicate mean annual air temperature and mean  
7 annual precipitation, respectively; ST, soil temperature; SMC, soil moisture content; SOC, soil organic carbon; TN, soil total nitrogen; TP, soil total phosphorus; Clay, soil clay content; litter  
8 C/N, total carbon/total nitrogen of litter.

1


**Table 2.** The effect of forest types and climate on the soil enzyme activities and PLFAs

| Treatment       | Climate                         |                | Forest type       |               | Climate × Forest type |               |                   |
|-----------------|---------------------------------|----------------|-------------------|---------------|-----------------------|---------------|-------------------|
|                 | <i>F</i>                        | <i>P</i>       | <i>F</i>          | <i>P</i>      | <i>F</i>              | <i>P</i>      |                   |
| Enzyme activity | BG                              | <b>30.487</b>  | <b>&lt;0.0001</b> | <b>6.852</b>  | <b>0.003</b>          | 3.105         | 0.056             |
|                 | NAG                             | <b>32.793</b>  | <b>&lt;0.0001</b> | 5.183         | 0.10                  | <b>3.635</b>  | <b>0.035</b>      |
|                 | LAP                             | <b>171.864</b> | <b>&lt;0.0001</b> | <b>16.364</b> | <b>&lt;0.0001</b>     | 1.813         | 0.176             |
|                 | AP                              | <b>95.070</b>  | <b>&lt;0.0001</b> | <b>48.117</b> | <b>&lt;0.0001</b>     | <b>22.446</b> | <b>&lt;0.0001</b> |
| PLFAs           | tPLFA                           | <b>7.764</b>   | <b>0.001</b>      | 2.697         | 0.079                 | <b>8.666</b>  | <b>0.001</b>      |
|                 | Bacteria                        | 2.796          | 0.073             | <b>4.921</b>  | <b>0.012</b>          | <b>8.357</b>  | <b>0.001</b>      |
|                 | Fungi                           | <b>8.002</b>   | <b>0.001</b>      | <b>21.255</b> | <b>&lt;0.0001</b>     | <b>25.023</b> | <b>&lt;0.0001</b> |
|                 | Actinomycetes                   | 0.533          | 0.591             | 2.979         | 0.062                 | <b>3.500</b>  | <b>0.040</b>      |
|                 | F/B                             | <b>3.731</b>   | <b>0.032</b>      | <b>15.502</b> | <b>&lt;0.0001</b>     | <b>6.378</b>  | <b>0.004</b>      |
|                 | G <sup>+</sup>                  | 0.603          | 0.552             | <b>3.395</b>  | <b>0.043</b>          | <b>5.934</b>  | <b>0.005</b>      |
|                 | G <sup>-</sup>                  | <b>12.503</b>  | <b>&lt;0.0001</b> | <b>6.890</b>  | <b>0.003</b>          | <b>11.106</b> | <b>&lt;0.0001</b> |
|                 | G <sup>+</sup> / G <sup>-</sup> | 1.662          | 0.202             | 0.069         | 0.933                 | 2.257         | 0.117             |

2

The abbreviations of the variables included in this table are shown in Figure 2 and 3.





1

2

**Figure 1.** Soil enzyme activities under different forest types in different climatic zones. BG, b-1, 4-glucosidase;

3

NAG, b-1,4-N-acetylglucosaminidase; LAP, leucine aminopeptidase; AP, acid phosphatase. **The capital letters A,**

4


**B, C, and D** represent the variations in the enzyme activities of BG, NAG, LAP and AP, respectively. Different

5

lowercase letters indicate significant differences between forests in the same climatic zone. The abbreviations of

6

the sampling sites are shown in Table 1.




2

3 **Figure 2.** The PLFA contents, Fungi:Bacteria ratios, and G<sup>+</sup>/G<sup>-</sup> for different forest types in different climatic zones


4 (A. Liangshui; B. Taiyue; C. Dinghu). Different lowercase letters indicate significant differences among forests in

5 the same climatic zone. F/B, fungi/bacteria; G<sup>+</sup>/G<sup>-</sup>, Gram-positive bacteria/ Gram-negative bacteria. The

6 abbreviations of the sampling sites are shown in Table 1.



1  
 2 **Figure 3.** Redundancy analysis (RDA) ordination biplot of soil enzyme activities and environmental properties for the different forest types in different climatic zones (A. Liangshui; B. Taiyue;  
 3 C. Dinghu). Only the environmental variables that were significantly correlated with RDA1 are shown. The dotted lines and solid lines represent the environmental variables and enzyme activities.  
 4 The variables in this table were abbreviated as follows: TC(litter) = litter total carbon; TN(litter) = litter total nitrogen; C/N(litter) = litter total carbon/nitrogen; ST = soil temperature; SMC = soil  
 5 moisture content; Clay = soil clay content; SOC = soil organic carbon; TN = soil total nitrogen; TP = soil total phosphorus; C/N = soil carbon/nitrogen; C/P = soil carbon/phosphorus, and N/P =  
 6 soil nitrogen/phosphorus.



2 **Figure 4.** Redundancy analysis (RDA) ordination biplot of soil microbial community structure and environmental properties for different forest types in different climatic zones (A. Liangshui;  
 3 B. Taiyue; C. Dinghu). Only the environmental variables that were significantly correlated with RDA1 are shown. The dotted lines and solid lines represent the environmental variables and lipid  
 4 signatures. The abbreviations of the variables included in this figure are shown in Figure 4.