

Dear Jack Middelburg,

Thank you for considering our manuscript for publication after minor revisions. We also thank Isaac Santos and Referee #2 for their time and effort reviewing this manuscript. Please find our answers to the comments, detailed changes and new manuscript, where changes are shown in red.

Referee #1 (Isaac Santos)

Comment 1: Page 1, Line 2, abstract: “lower”. Add a short note on how much lower.

Answer 1: The radium flux of our study is around 20% lower than the inventory of radium 228 in the Atlantic by Moore et al. [2008] and the global estimate of Kwon et al. [2014], which is significant. This is now stated in the introduction (page 1 line 7) and in the conclusion (page 16 line 8)

Comment 2: Page 2, Line 33: “raw assumptions”. I suggest the authors spell out the major raw assumptions here, or just omit this early criticism.

Answer 2: This sentence is ambiguous. No assumption Interactive is required to perform box averages or interpolations. The “raw assumptions” are the very fact that box averages or linear interpolations can produce a precise and unbiased estimate of the total flux, which is not true when data are sparse. It is has rephrased as: “These direct approaches suffer from strong potential biases when the data are sparse, because the total amount of ^{228}Ra in the ocean is then roughly estimated, using regional averages (Rodellas et al., 2015) of observations or linear interpolations (Moore et al., 2008). Therefore, they are suitable only in regions with dense sampling, such as the Atlantic basin.”

Comment 3: Page 7, Line 17: Why the Indian and Pacific Basins are so high? I looked for a C1 BGD Interactive comment Printer-friendly version Discussion paper comment on that later in the discussion but could not find. I encourage the authors to add a paragraph (probably in the discussion) offering some thoughts explaining the spatial distribution.

Answer 3: This is a very interesting question. I think there are two points here: a) At page 7 line 17, a comparison is made between radium fluxes, a proxy of SGD, and rivers. My point is that these two sources of water and nutrients are distributed in a very different way. But, given the size of the basins, rivers may be more biased than radium/SGD. The Indo-Pacific basin accounts for 52% of continental shelves and 63% of radium fluxes, compared to 29% and 27% for the Atlantic. In order to address a comment by referee #2, this comparison will be moved to 3.4, transformed into a comparison between rivers and SGD, and the previous remark will be added among other improvements. b) The distribution of fluxes is more complex than “high in the Indian and Pacific, low in the Atlantic and Arctic”. Differences within each basin are larger. Except for a few cases where the explanation is obvious, for instance the very low fluxes in the

eastern Pacific, where the shelves are very thin, it is difficult to identify the reasons why some regions emit more radium than others. This is probably related to the local geology: high fluxes may be due to more permeable sediments, to more wave, tide or geothermal energy inducing larger SGD, but also to a higher content in radium-228 and its parent radionuclide, thorium-232. We are not specialists of this field and the literature we know so far does not provide the global scale pattern of these parameters. For instance, it does not explain what the main differences between regions like the Bay of Bengal or the Gulf of Guinea are. Previous radium and SGD studies do not explain the reason for the source distribution either. Relating differences in radium-228 fluxes to a specific geological parameter is not possible. However, a few explanations has been added in the discussion (page 13 lines 18-25): “The spatial distribution of the fluxes in this study is consistent with Kwon et al. (2014), with two thirds of radium-228 flowing to the Indian and Pacific Basins and also quite high fluxes in the western North Atlantic. It turns out that these basins have the 20 highest riverine sediment loads (Milliman, 2001). However, we have shown that rivers could account for 7% to 20% of ^{228}Ra fluxes only, which does not exclude an indirect impact on the other sources through the geology of the continental shelf. A difference among basins in the quantity of radium diffusion, which is poorly known, is also possible. If it is not the case, then the SGD are significantly higher in East Asia than in other regions. SGD can be driven by as many factors as storms, waves, tides and thermal gradients, and depend on the permeability and structure of coastal and shelf sediments (Moore, 2010a). 25 Comparative data on these factors in several basins will be necessary to explain the origin of the observed differences.”

Comment 4: What depth of the upper ocean was used to integrate the radium observations? How does it compare to previous studies? What sort of extrapolation was made in terms of depth integration for locations with no data in deeper waters?

Answer 4: There may be a misunderstanding. No integration (averaging) on the water column is performed. So no extrapolation has to be made on concentrations between observations. This is the strength of inverse modeling. We just minimize the differences between observed concentrations and model concentrations at all observation points shallower than 200m, without assuming anything about radium concentration elsewhere or regional averages. This will be made clearer in the introduction, where some sentences of the current manuscript are ambiguous, especially on page 3 from lines 2 to 6, which now read as : “The advantage of this method is that no arbitrary averaging or interpolation is required: observed concentrations are simply compared to model concentrations at the same points. It is expected to be robust and consistent, since the model is based on physical considerations.”

Comment 5: Page 10, Line 21: “A fraction” can be replaced by “Nearly all”.

Answer 5: Done

Comment 6: Page 10, Line 30: A number of diffusion studies are briefly cited. Considering the emphasis on diffusion, I encourage the authors to add more information about those studies. Maybe a summary table with the source of diffusion data and how it was estimated.

Answer 6: A sixth table was added, citing studies on diffusion from fine-grained sediments (those who contribute the most), their locations, their methods and the results. We have put in bold those which might have accidentally included SGD into the diffusive and bioturbative flux. Explaining the methods in full detail would be long and is not possible here, but we have shown the references.

Comment 7: Page 11, Line 20 and elsewhere: The comparison between SGD and river flow is appropriate and should be kept since it puts results in perspective. However, radium-derived SGD is likely to be saline water, while rivers are a source of fresh water. I encourage the authors to add a note qualifying those differences using the literature.

Answer 7: Page 11 line 20: Indeed, SGD is mainly recirculated saline water, as already explained in the introduction. According to Taniguchi [2007], there is only 2600 km³/yr of fresh SGD, between 2% and 20% of the total SGD. The comparison with rivers is made because they both contain nutrients and trace elements. The end of 3.4 has been changed to explain all this more clearly, using the literature: “According to Taniguchi et al. (2007), there is only $2.6 \times 10^{12} \text{ m}^3 \text{ yr}^{-1}$ of fresh groundwater discharge, representing between 2% and 20% of the SGD, and the remaining part is recirculated seawater.”

Comment 8: Similar to the previous comment, the comparison to seepage meters on the last paragraph of page 12 may need to be qualified. Many seepage meter deployments are made in very shallow nearshore waters and may capture fresher SGD, while Ra-228 covers a much larger scale. Consider using radium studies to build this comparison.

Answer 8: Indeed, seepage meters are not representative of total SGD, as they are concentrated close to the coast, where both fresh and saline SGD are expected to be higher than the continental shelf average. This is why the fluxes of most local studies are one order of magnitude higher than ours. This is now explained more clearly by several small changes in 4.1. In the previous version of the manuscript, the only mentioned weakness of seepage meters was spatial variability, which was incomplete. However, seepage meters are interesting because they are independent from radium and are not affected by uncertainties on radium diffusion or radium concentration in groundwater. Not mentioning them would be strange. Radium studies, at a larger scale, are already the object of the previous paragraphs.

Referee #2

The manuscript was proofread again, and a few errors were corrected, including the use of numbers and abbreviations at the beginning of sentences.

Comment 1: I do not think the title accurately represents the main focus of the paper, which is an effort to improve the application of the inverse modeling technique to ^{228}Ra , as opposed to an interpretation of an updated ^{228}Ra flux estimate or SGD estimates. Either the title should be changed to better reflect the fact that the main contributions of this work are improvements to the model, or more discussion should be added on the interpretation of the results of the model and how they change our understanding of ^{228}Ra and/or SGD. If the latter is chosen, the authors should be sure to highlight how their contributions are unique from those of Kwon et al (2014), aside from simply reporting the newly calculated Arctic and Southern Ocean ^{228}Ra fluxes.

Answer 1: The title should reflect three aspects: a) We have shown that parameters such as the number of source regions and the cost function matter a lot in inverse modeling, b) By using appropriate parameters, we produce a more precise estimate than previous studies, c) However, this is not immediately usable for SGD studies as this source still have to be separated from diffusion. The current title might not emphasize the first aspect enough but finding a better one is difficult. The title has been changed to “Improving the inverse modeling of a trace isotope: how precisely can radium- 228 fluxes toward the ocean and Submarine Groundwater Discharge be estimated?”

Comment 2: There is a short discussion about the sources of Ra, and it is mentioned that dust inputs are small compared to the other sources used in the model. The dissolved riverine source should also be introduced here instead of later in the paper, because this is where the sources/sinks are first introduced. You state later on that the dissolved component of the riverine flux is negligible, but it would be better to approximate the relative contribution as you have done for the dust source in line 20.

Answer 2: Riverine dissolved radium-228 is less than 1% of the total radium-228 flux. This figure, similar to that of dust, has been added to the introduction (Page 2 line 23). More explanations are given at page 10, using the annual river discharge and the average concentration in dissolved radium: “According to Moore et al. (2008), the average dissolved ^{228}Ra activity in rivers is $0.65\text{--}1.95 \times 10^5 \text{ atoms.l}^{-1}$. As rivers annually discharge 35000 km^3 of freshwater (Milliman, 2001), the dissolved riverine source lies between 2.3 and $6.8 \times 10^{21} \text{ atoms.yr}^{-1}$, which is less than 1% of the total flux.”

Comment 3: Page 7 lines 17 – 18: Why is the flux of ^{228}Ra (in units of atoms y^{-1}) compared to river discharge (I’m assuming this is units of $\text{m}^3 \text{ y}^{-1}$)? Why not compare the flux of SGD to river discharge, so that it is a volume-to-volume comparison? I’m not sure if this is intended to make the point that the largest flux of ^{228}Ra does not correlate with the greatest river discharge, but the

authors argue that rivers do not carry much dissolved ^{228}Ra , so in that case they shouldn't correlate anyway.

Answer 3: It would indeed make more sense to compare river discharge to the other large fluxes of water and nutrients to the ocean, Submarine Groundwater Discharge (SGD), rather than to radium-228 fluxes. The problem is that SGD suffers from high uncertainty, as highlighted in the article. Average basin-wide SGD fluxes can be compared, but this is implicitly assuming that diffusion per unit of surface is nearly the same in both basins, which is far from certain. This issue will be moved to the end of 3.4, where we now compare the distribution of river discharge with both radium fluxes and SGD, and explain why the latter is imprecise, although more interesting. The ratios are very close anyway. Page 11 lines 30-35 now read as: “In total, 63% of the ^{228}Ra flows into the Indian and Pacific basins, which is more than their share of the continental shelf (52%). As shown in Fig.8, the estimated contributions of SGD to ^{228}Ra fluxes are in very similar proportions, with 68% going to the Indian and Pacific basins. This represents more than 70% of all the groundwater discharge when the differences in groundwater ^{228}Ra concentration are considered. However, uncertainties on SGD distribution are high, because diffusion from sediment might not be identical in all basins. These results contrast with the other source of water, nutrients and trace elements to the ocean: 60% of the global river discharge flows to the Atlantic and Arctic basins (Milliman, 2001).”

Comments 4 and 5: Indeed, I had called the wrong figures. This is now corrected

Comment 6: Conclusion lines 14 – 15: be specific about the other cost functions that were tested, instead of saying “. . .the other cost functions. . .”. That way if a reader skips to the conclusion, they will still understand your specific results. The same sentence ends by saying “. . .more realistic assumptions on error statistics”; again, be specific about what the more realistic assumptions were. A large part of the paper is dedicated to testing assumptions, so the conclusion should point out the results of those tests.

Answer 6: The difference between the three cost functions and between the underlying assumptions on error statistics is now described explicitly in the conclusion, so that a reader skipping to the conclusion can understand what it is about and why it is important. Page 16 lines 2-6 now read as: “These precise estimates are obtained by minimizing the squared differences between model and observed concentrations on a logarithmic scale. We think this cost function is more realistic than a linear least-squares one because it assumes that the error standard deviation is proportional to ^{228}Ra concentration, rather than constant, which proves to be more accurate. It is also better than a "proportional" cost function, weighted by the inverse of the observed concentration, which tends to produce underestimations of concentrations and fluxes.”

Comment 7: Conclusion lines 24 – 25: The sentence that begins with “Therefore, besides estimating the sources. . .” should not be included in the conclusion, as it is not one of the main

points emphasized in the paper. It is more appropriate to move this line to the end of the “model biases” section or another part of the manuscript.

Answer 7: This sentence will be removed from the conclusion, as it is not a main focus of the paper. This possibility of improving a circulation model using the residuals of inversions is already mentioned in 3.2 and 4.2 and there is not much to add.

Comment 8: The conclusion should have a stronger ending and remind the reader why it is important to better resolve the ^{228}Ra model. Line 17 is a good reminder of why determining the shelf fluxes is important, but the conclusion should also highlight why it is important to separate the SGD and diffusion sources. This could also be discussed in more detail earlier in the paper, but should at least be mentioned again in the conclusion.

Answer 8: Evaluating the SGD is the goal of several studies and radium-228 has often been used as a tracer for this purpose. One important conclusion of this article is that using radium-228 in an inventory or in inverse model is very imprecise because SGD and diffusion are poorly separated. This is now stated more explicitly by changes in 3.4 (page 11 lines 18-29: “As long as diffusion cannot be separated from SGD more efficiently and is not shown to be negligible, no precise SGD estimate based on ^{228}Ra can be produced.”) and in the conclusion (“These results mean that all the studies which have used or will use ^{228}Ra to estimate SGD will suffer from a very high uncertainty until diffusion is properly estimated or proven to be negligible.”).

Comment 9: Page 16 line 2: include the actual NEMO website

Answer 9: Done (page 16 line 23)

Comment 10: Full references for the data sources used in the model are not included in the reference list; is this because they are only listed in the supplementary material? I suggest adding a supplementary reference list for these data sources. Also, in the Arctic references, the year for the Rutgers van der Loeff et al. 2013 reference should be 2012, not 2013.

Answer 10: All the references used in our radium-228 dataset have been included in the supplement. The Rutgers van der Loeff et al. 2012 and two other mistakes have been corrected.

Comment 11: On Figure 9, the cruise across the Fram Strait has been completed, so this should be in yellow instead of red.

Answer 11: Corrected

Improving the inverse modeling of a trace isotope: how precisely can radium-228 fluxes toward the ocean and Submarine Groundwater Discharge be estimated ? ~~An updated estimate of radium 228 fluxes toward the ocean : how well does it constrain the submarine groundwater discharge ?~~

Guillaume Le Gland¹, Laurent Mémery¹, Olivier Aumont², and Laure Resplandy³

¹LEMAR, Institut Universitaire Européen de la Mer, Plouzané, France

²LOCEAN, Institut Pierre Simon Laplace, Paris, France

³Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

Correspondence to: Guillaume Le Gland (guillaume.legland@univ-brest.fr)

Abstract. Radium-228 (^{228}Ra), an almost conservative trace isotope of the ocean, supplied from the continental shelves and removed by a known radioactive decay ($T_{1/2} = 5.75 \text{ yr}$), can be used as a proxy to constrain shelf fluxes of other trace elements, such as nutrients, iron, or rare earth elements. In this study, we perform inverse modeling of a global ^{228}Ra dataset (including GEOSECS, TTO and GEOTRACES programs, and, for the first time, data from the Arctic and around the Kerguelen islands) to compute the total ^{228}Ra fluxes toward the ocean, using the ocean circulation obtained from the NEMO 3.6 model with a 2° resolution. We optimized the inverse calculation (source regions, cost function) and find a global estimate of the ^{228}Ra fluxes of $8.01 - 8.49 \times 10^{23} \text{ atoms yr}^{-1}$, **more precise and around 20% lower**~~lower and more precise~~ than previous estimates. The largest fluxes are in the western North Atlantic, the western Pacific and the Indian Ocean, with roughly two thirds in the Indo-Pacific basin. A first estimate in the Arctic Ocean is assessed ($0.43 - 0.50 \times 10^{23} \text{ atoms yr}^{-1}$). Local misfits between model and data in the Arctic, the Gulf Stream and the Kuroshio regions could result from flaws of the ocean circulation in these regions (resolution, atmospheric forcing). As radium is enriched in groundwater, a large part of the ^{228}Ra shelf sources comes from submarine groundwater discharge (SGD), a major but poorly known pathway for terrestrial mineral elements, including nutrients, to the ocean. In contrast to the ^{228}Ra budget, the global estimate of SGD is rather unconstrained, between 1.3 and $14.7 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$, due to high uncertainties on the other sources of ^{228}Ra , especially diffusion from continental shelf sediments. Better precision on SGD cannot be reached by inverse modeling until a proper way to separate the contributions of SGD and diffusion at a global scale is found.

1 Introduction

Trace Elements and Isotopes (TEI) are low concentration components of the ocean, but they contain decisive information for our understanding of its dynamics. The international program GEOTRACES has been designed to improve our knowledge on the TEIs concentrations and the oceanic processes controlling their distribution, by means of observations, modeling and

laboratory experiments. Since 2006, GEOTRACES cruises have been mapping the global distribution of tens of these TEIs. Some of them are studied because they constitute micronutrients for living organisms, like iron (Fe), or **are** pollutants, like lead (Pb) and cadmium (Cd). Others are proxies of ocean dynamics or of biogeochemical processes: For instance, neodymium (Nd) is a proxy of the exchanges between seabed and seawater (Jeandel et al., 2007), and thorium 234 (^{234}Th) is related to the biological carbon pump (Clegg and Whitfield, 1991; Buesseler et al., 1992; Henson et al., 2011). Radium (**Ra**) isotopes are of particular interest as they are proxies of all TEI fluxes from sediments and continents toward the ocean. More specifically, radium isotopes have been used to estimate a still poorly known pathway to the ocean : Submarine Groundwater Discharge (SGD).

SGD is defined as the flux of water from coastal aquifers to the ocean, regardless of its composition and origin. Part of it is meteoric freshwater, but the largest part is infiltrated seawater (Burnett et al., 2006; Moore, 2010b). In these aquifers, water gets enriched in nutrients and trace elements, released from soils and rocks or coming from land pollution, before flowing back to the ocean. It has traditionally been considered that the main coastal sources of terrestrial mineral elements to the ocean are rivers, but there is growing evidence that SGD is in fact a source of nutrients of the same order of magnitude, affecting the biogeochemistry at all scales, from coastal regions (Hwang et al., 2005; Kim et al., 2005) to ocean basins (Moore et al., 2008; Rodellas et al., 2015). SGD is a suspected cause of algal blooms, including harmful algal blooms (LaRoche et al., 1997), and a pathway for contamination. In spite of this, **its**~~their~~ contribution is still much less precisely known than the river inputs. Because of the strong heterogeneity in its distribution and intensity, properly estimating SGD by direct methods requires an intense sampling, which is far from being fulfilled (Burnett et al., 2006; Moore, 2010a). Indirect methods should then be used: Radium (**Ra**) isotopes offer a great potential due to their relation to SGD and their simple chemistry.

All the four natural radium isotopes, ^{223}Ra ($T_{1/2} = 11.4$ d), ^{224}Ra ($T_{1/2} = 3.6$ d), ^{226}Ra ($T_{1/2} = 1602$ yr) and ^{228}Ra ($T_{1/2} = 5.75$ yr), are produced within the rocks by the radioactive decay of thorium. Since radium is far more soluble in water than thorium, its main source is not the decay of dissolved thorium in the ocean, but dissolution from lithogenic material. Therefore it is used as a tracer of boundary fluxes. This element is released into the ocean by three main sources located on the continental shelf: ~~desorption~~**dissolution** from riverine particles, diffusion from seabed sediments, and SGD, which are highly enriched. Dust inputs **and dissolved riverine** ^{228}Ra **each** account for less than 1% of all inputs (Moore et al., 2008). In the ocean, **radium**~~Ra~~ is almost conservative. It is removed by radioactive decay and scavenging. Scavenging is associated with a residence time of approximately 500 yr (Moore and Dymond, 1991), making it negligible for all Ra isotopes except ^{226}Ra . Then the distributions of the other three isotopes, whose radioactive sinks are known, depend only on the source distribution and transport by the ocean circulation. Since their half-life time scales are small relative to the time scale of basin-wide horizontal mixing, typically a few years to a few decades, ^{223}Ra and ^{224}Ra are unsuitable for large scale studies. **Radium-228** ~~^{228}Ra~~ , whose half-life of 5.75 yr is short enough to neglect scavenging but long enough to consider the global ocean, is suitable for global scale analyses and is thus considered here.

A simple way to use the information provided by this isotope is to make an observation-based inventory of the ocean ^{228}Ra . At steady state, the supply of ^{228}Ra must balance the loss from disintegration, i.e. 12% every year. According to Charette et al. (2016), these total ^{228}Ra fluxes can be used to estimate fluxes of nutrients, iron and rare earth elements. **Radium-228** ~~^{228}Ra~~

fluxes are also a way to estimate the SGD fluxes by subtracting the contribution from rivers, diffusion and bioturbation, and dividing the remaining flux by the mean ^{228}Ra concentration in groundwater. By this method SGD has been estimated at $0.03 - 0.48 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$ in the Mediterranean Sea (Rodellas et al., 2015) and $2 - 4 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$ in the Atlantic Ocean (Moore et al., 2008). These direct approaches suffer from strong potential biases when the data are sparse, because the total amount of ^{228}Ra in the ocean is roughly estimated, using regional averages (Rodellas et al., 2015) of observations or linear interpolations (Moore et al., 2008). Therefore, they are suitable only in regions with dense sampling, such as the Atlantic basin. These direct approaches suffer from strong potential biases associated with the relative sparsity of observations: raw assumptions have to be made in order to estimate regional averages or to interpolate scattered observations in space. Therefore, it is suitable only in regions with dense sampling, such as the Atlantic basin.

Inverse modeling techniques represent an alternative and powerful approach to estimate the fluxes, providing that the ocean circulation is known with sufficient accuracy. Inverse modeling is based on three elements: observations of the physical quantity of interest (here oceanic ^{228}Ra concentration), a forward prognostic model simulating the same quantity as a function of the unknown variables to be assessed (here the shelf fluxes), and an algorithm finding the values of the unknowns minimizing the misfit between model and observations. tracer (Ra) observations, a forward prognostic model which simulates the tracer distribution as a function of the fluxes to be assessed (here the spatial distribution and intensity of the shelf fluxes), and an algorithm optimizing the fluxes of the model in order to minimize the misfit between the observations and the model results. The advantage of this method is that no arbitrary averaging or interpolation is required: observed concentrations are simply compared to model concentrations at the same points. It is expected to be robust and consistent, since the model is based on physical considerations. The advantage of this method is that the ocean model is used as an interpolator, based on physical considerations, which is expected to be robust and consistent.

Kwon et al. (2014) used such an inverse modeling approach to produce a global estimate of ^{228}Ra fluxes of $9.1 - 10.1 \times 10^{23} \text{ atoms yr}^{-1}$ between 60°S and 70°N , corresponding to $9 - 15 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$ of SGD. Their study is based on a data-constrained global ocean circulation model (DeVries and Primeau, 2011), considering 50 source regions on the continental shelf, and minimizing an ordinary least-squares cost function. In this study, we estimate the radium fluxes from all continental shelves around the world and localize the most intense sources, using an inverse modeling technique with more data than previous studies (Kwon et al., 2014). The data set has been augmented with data from two recent GEOTRACES cruises and from the Southern Ocean, the North Pacific, the Mediterranean Sea and the Indonesian Seas. It also contains data from the Arctic, a basin absent from Kwon's study. The forward model is built on the Ocean General Circulation Model (OGCM) NEMO (Nucleus for European Modeling of the Ocean). Our main improvement is a careful analysis of sensitivity and errors, which reveals that the result depends on the model mathematical parameters, such as the cost function and the number of regional sources that are considered. We have performed several inversions and analyzed the residuals and uncertainties, to determine the most appropriate mathematical parameters and evaluate the precision of the flux estimates. This paper is organized as follows. In section 2, we describe the different aspects of the inversion technique, e.g. the global dataset, the forward model based on the NEMO OGCM, the different cost functions, the choice of the source regions related to SGDs, and the inverse method. Section 3 presents the main results, e.g. the global and regional estimates of ^{228}Ra supply, and the sensitivity of these estimates to several parameters of our approach, such as the cost function or the

number of coastal ^{228}Ra sources. Section 4 compares our results with results obtained in previous studies, and discusses issues associated with such an approach, with an emphasis on SGD.

2 Methods

2.1 ^{228}Ra dataset

5 Since the late 1960's (Moore, 1969; Kaufman et al., 1973) tens of oceanographic cruises have carried out measurements of ^{228}Ra (e.g. articles listed in Table S1). The dataset used in this study includes, among others, observations from three international programs sampling trace elements. Data from the Indian Ocean cruise of GEOSECS (**GEochemical Ocean SECTIONS Study**), in 1977-1978, are included. From 1981 to 1989, the TTO (Transient Tracers in the Oceans) program produced a considerable number of ^{228}Ra measurements in the Atlantic, from 80°N to 60°S , at all depths, making **of** the Atlantic Ocean the
10 best sampled ocean by far. Currently, new data from all oceans are being produced by GEOTRACES. **In total**, 6059 data from all basins are used **in-total**, of which 2789 are shallower than 10 m, 1107 are located between 10 and 200 m deep, 606 between 200 and 600 m deep, and 1557 deeper than 600 m. Our data set comprises 1359 more measurements than in Kwon et al. [2014]: Two GEOTRACES sections, GA03 (United States – Cape Verde – Portugal) and GP16 (Ecuador – French Polynesia), and data
15 present study (See Table S1). These supplementary observations extend the data coverage to regions north of 70°N and around the Kerguelen islands. In a near future, other sections from the GEOTRACES program will complete the global covering and may help in studying deeper sources.

For the purpose of our study, data have been averaged in each model grid cell (see section 2.2 for more details on the model configuration), leading to the 3076 cell averages shown on Fig. 1. The density of the measurement is noticeably uneven. The
20 Atlantic Ocean, north of 20°S , and the Arctic, are the most densely covered basins and the only regions with a significant number of data at depth deeper than 10 m. Other regions are sparsely sampled, leaving wide areas with no or very few measurements, like the western Indian Ocean, the equatorial Pacific or the Pacific sector of the Southern Ocean.

Data are expressed in concentration or activity units, with the following conversion factor: $1 \text{ dpm m}^{-3} = 4.36 \times 10^6 \text{ atoms m}^{-3}$. They range from 0.04 to 724.5 dpm m^{-3} . The highest concentrations are found in the Bay of Bengal and the coastal seas of
25 eastern Asia, the lowest values are located in the Southern Ocean. Concentrations are generally higher than 10 dpm m^{-3} in the Indian Ocean, the Atlantic and the Pacific north of 30°N , lower in the rest of the Pacific. In the Atlantic, west of a line running from the Amazon delta to Newfoundland, most concentrations are higher than 30 dpm m^{-3} .

2.2 Forward tracer model

The second requirement of the inversion technique is a ^{228}Ra transport model, allowing to link in situ observations to the
30 boundary conditions or shelf sources of ^{228}Ra . The transport equation of tracer A_i (originating from the source region i) is

given by:

$$\frac{\partial A_i}{\partial t} = -\mathbf{U} \cdot \nabla A_i + \nabla \cdot (K \nabla A_i) - \lambda A_i + S_i \quad (1)$$

\mathbf{U} and K are the velocity field and the eddy diffusivity coefficient respectively. The two first terms on the right together constitute transport. They are derived from NEMO 3.6 model (~~Nucleus for European Modeling of the Ocean~~) using OPA
5 (Madec, 2015) as a general circulation component, coupled with the sea-ice model LIM3 (Vancoppenolle et al., 2009), with an ORCA2 global configuration. The model has an horizontal resolution of $2^\circ \times 2^\circ \cos \phi$ (where ϕ is the latitude) enhanced to 0.5° near the equator. The mesh is tripolar in order to overcome singularities, the North Pole being replaced by two inland poles in the Northern Hemisphere. It has 31 vertical levels, ranging from the surface to 6000 m deep, the upper layer covering the first 10 m. The simulation is forced by a seasonal climatological dataset, based on NCEP/NCAR reanalysis and satellite
10 data.

λ is the radioactive decay constant, 0.12 yr^{-1} , given by the half-life of ^{228}Ra which is 5.75 yr. Decay is the sole sink. It is known and does not depend on environmental parameters, leaving the source term as the only unknown to be determined by the inversion technique.

S_i is the source term specific to the i^{th} region, representing riverine inputs, sedimentary diffusive fluxes and groundwater
15 discharge fluxes of ^{228}Ra from region i . In this study, sources are assumed to be only on the continental shelf, defined as the seabed shallower than 200 m. This depth range, spanning 16 model levels, is chosen because it is where most groundwater discharge outflows. However, diffusion from sediments also occurs at higher depths (Hammond et al., 1990).

Equation (1) shows that the A_i fields depend linearly on S_i . That means that any ^{228}Ra distribution can be written as a linear combination of the A_i fields. It is important to emphasize that the circulation is supposed to be "perfect", e.g. no correction
20 of the \mathbf{U} field is looked for. Nevertheless, the simulated circulation obviously suffers from deficiencies, and that point has to be kept in mind when interpreting the results. From now, what we refer to as "model concentration", $[^{228}\text{Ra}]^{\text{mod}}$, is a linear combination of the tracer final concentrations A_i , the coefficients being the source intensities x_i :

$$[^{228}\text{Ra}]^{\text{mod}} = \sum_{i=1}^n A_i x_i \quad (2)$$

A non-optimal model concentration was computed by assuming a uniform constant flux per unit of surface everywhere, with
25 a global fit using the average concentration estimated from the observations: This defines the first guess estimate before the inversion. The inversion undertaken in this study aims at optimizing the parameters x_i in order to minimize the total difference between the observations and the model ^{228}Ra distribution.

As a consequence of its coarse horizontal resolution, continental shelves are only poorly resolved by the ORCA2 grid. The emitting surface is underestimated and some regions with narrow continental shelves would be completely omitted. To over-
30 come that deficiency, sub-model grid scale bathymetric variations are accounted for by comparing the model grid to a global $2'$ resolution bathymetry ETOPO2 of the National Geophysical Data Center (NGDC). The algorithm is detailed in Aumont and Bopp (2006). According to this method, the total surface of continental shelf is $2.73 \times 10^{13} \text{ m}^2$, 73% higher than the $1.58 \times 10^{13} \text{ m}^2$ obtained with the coarser bathymetry.

The ocean – continent interface, including the Arctic and the Antarctic, is divided into 38 regions (Fig. 2). This first guess takes into account the sampling coverage (very low in the Antarctic for instance, and higher in the North Atlantic Basin or Bay of Bengal) and differences in the tracer distributions A_i , which should be large enough to give independent information. Delimitation is done by trials and errors, using the posterior covariance matrix of the inversion (see section 2.3): The number of sources is minimized by merging regions associated with negligible fluxes with close highly correlated regions. Most islands are ignored, because the areas of their continental shelf and thus their expected contributions to the ^{228}Ra balance are small. In the inversion process, islands can give rise to spurious fluxes to accommodate for other types of errors. The only islands considered in this study are the Kerguelen and Crozet Islands, in the Southern Ocean, because many samples have been taken in their surroundings which make it possible to constrain their contributions. Because of the lack of measurements and the coarse model resolution, the Persian Gulf, the Red Sea, the Baltic Sea, the North Sea and the Hudson Bay are not taken into account. The flux per unit of surface is assumed to be constant on each of the 38 emitting regions. Model simulations last for the equivalent of 100 years in order to reach a quasi steady state. It is more than 17 times larger than the half-life of ^{228}Ra , so that the total amount of ^{228}Ra does vary by less than 0.001%. As there are not enough data to study global inter-annual or seasonal variations, we do not take seasonal variations of ^{228}Ra concentrations into account. We implicitly assume that radium concentration is constant over time, and work with average concentrations over the 100^{th} year of simulation.

2.3 Inverse method

The last requirement of the inversion technique is to define a cost function measuring the misfit between the data and the model. This cost is then minimized by a method already used to assess air-sea gas fluxes (Gloor et al., 2001; Mikaloff Fletcher et al., 2006; Jacobson et al., 2007) and oceanic heat fluxes (Resplandy et al., 2016). If η represents the residuals, e.g. the discrepancy between model results and observations:

$$\mathbf{A}x = [^{228}\text{Ra}]^{obs} + \eta \quad (3)$$

Both sides are vectors, representing radium concentrations in each model cell containing data, their size being the number of data. \mathbf{A} is the matrix of footprints, representing the circulation model and composed of the A_i at each data point. x is the vector of unknowns, the flux of ^{228}Ra per unit of surface of each region. As only shelf sources are modeled and as data coverage below 200 m is sparse, only data shallower than 200 m are considered.

The distance between model concentrations and data is summed up in a scalar, the cost function $C(x)$. We look for the optimal flux vector x_{opt} minimizing the cost function. Different choices for C are possible, depending on the assumed probability distribution for the prior error on data and model footprints \mathbf{A} . Errors due to biased sampling are not considered here. All errors are supposed to be uncorrelated. In this study, three different cost functions have been tested and minimized. They all correspond to the sum of squares of a specific type of residuals. Their respective equations are listed in Table 1. The first one is an ordinary least-squares cost function, C_{ols} . According to the Gauss-Markov theorem, its minimization produces the best linear unbiased estimator when prior errors have no correlation, zero expectation, and the same variance. It is the simplest least-squares method, chosen in the study by Kwon et al. (2014). This function gives the same weight to all observations.

However, the hypothesis of the homogeneity of the variance is questionable: Far offshore, where concentrations are lower and less sensitive to small changes in coastal sources, observations and model errors can be expected to be lower. Neglecting this fact means these data are not fully exploited, as their contribution to the cost function is relatively small. Two other cost functions with a higher weight for smaller values are then considered for comparison. They are assuming heteroscedastic data, with higher variances for higher concentrations. In the proportional least-squares cost function, C_{prp} , the error standard deviation is supposed to be proportional to the observed concentrations. The logarithmic least-squares cost function, C_{log} , works differently and assumes the logarithms of concentrations have the same error variance. It is less sensitive than C_{prp} to model overestimations and more to underestimations. It is the only cost function which is not a quadratic function of the sources.

The residuals after inversion indicate what the inverse model cannot fit. In a "perfect" inversion, these residuals should be assimilated to noise, e.g. small and without structure, due for instance to coarse resolution. In most inversions, that is not the case, and the distribution of residuals emphasizes biases or errors either in the chosen hypotheses (such as a perfect circulation) or in the setting of the inversion (number and choice of the regions). The posterior uncertainties on radium fluxes and correlations between regions are computed following the method described in Appendix A. A regional flux has a large uncertainty when it is constrained by few data or correlated to other regions (Gloor et al., 2001), and two regions are strongly correlated when the ^{228}Ra emitted by each is transported to the same places and are then harder to differentiate. The computed posterior uncertainties are precise only if all the preceding assumptions on prior errors are correct. The coherence of error assumptions with the results has to be checked (see 3.2). Along with the main inversion considering 38 regions, we performed 4 other inversions with a higher or lower number of regions in order to estimate the sensitivity to this parameter (see 3.3).

3 Results

3.1 ^{228}Ra fluxes

The ^{228}Ra fluxes from each of the 38 regions, deduced by minimizing each of the three cost functions, are shown on Fig. 3 with their confidence intervals, and compared with the prior estimates. The global fluxes for each method are also shown. As they are sums of local fluxes, their standard deviations are proportionally lower.

The global ^{228}Ra flux within one standard deviation is $8.01 - 8.49 \times 10^{23} \text{ atoms yr}^{-1}$ according to the C_{log} inversion. As we will explain in section 3.2., this estimate is the most accurate of the three. Fluxes are found to be comparatively high in the North Atlantic (regions 5 to 16), in the western Pacific (22 to 27) and in the Indian Ocean (28 to 34), together accounting for 62.6% of the continental shelf and 82.8% of the global flux of ^{228}Ra . The highest fluxes are located in the China Seas (23 and 25) and in the eastern Indian Ocean (29 to 32), where the inversion process produces the largest increase compared to prior estimates, and to a lesser extent on the east coast of North America (10 and 13). Highest fluxes are located on the east coast of North America (10 and 13), in the China Seas (23 and 25) and in the eastern Indian Ocean (29 to 32), where the inversion process produces the largest increase compared to prior estimates. Conversely, inversion significantly reduces the prior estimates in the Arctic Ocean (35 to 37), in the Bering Sea (21) and in the eastern Pacific Ocean (17 to 20). Fluxes are also quite

low in the Southern Ocean (1 and 38) and in the South Atlantic (2 to 4). The newly estimated Arctic and Antarctic sources are in the range of 0.43 to 0.50 and 0.31 to 0.37 $\times 10^{23}$ atoms yr⁻¹ respectively, accounting for 5% to 6.2% and 3.6% to 4.6% of the total sources. ~~In total, roughly two thirds of the ²²⁸Ra flows into the Indian and Pacific basins, contrasting with the 60% of the global river discharge flowing into the Atlantic and Arctic basins~~ (Milliman, 2001).

- 5 Although having the same order of magnitude, uncertainties are generally lower than fluxes. They are highest in the western Pacific and Indian Oceans (regions 22 to 34), because of data sparsity. It is lower in better sampled oceans: The Arctic (35 to 37) and the Atlantic (2 to 16) Oceans, except for region 13 (Cape Hatteras to Newfoundland). The eastern Pacific (17 to 21) also has low uncertainties in absolute values, probably due to the low concentrations and prior errors there.
- The two other inversions produce roughly similar results, although fluxes are generally lower when derived from C_{prp} and generally have higher uncertainties when derived from C_{ols} . The global ²²⁸Ra flux is estimated to 7.16 – 8.14 $\times 10^{23}$ atoms yr⁻¹ with C_{ols} and 4.96 – 5.28 $\times 10^{23}$ atoms yr⁻¹ with C_{prp} . The three inversions agree on which basins and continents have the largest and smallest sources. Yet, local disagreements occur. Regions 5 (Amazon delta), 21 (Alaska and Bering Sea) and 33 (Arabian Sea) have higher fluxes with C_{ols} than with C_{log} , with non-overlapping confidence intervals, whereas the contrary is true for regions 12 (Mediterranean), 26 (Indonesian Seas) and 34 (East Africa). C_{prp} fluxes are generally lower, and in eight regions their confidence intervals overlap with none of the other inversions. Discrepancies happen because the fitting of the model to each observation implies different and possibly opposite effects on the source intensities. Each inversion uses different weights, which translates into different flux corrections. Fluxes from each of the three inversions are most dissimilar for regions where observations impose most dissimilar constraints, where the model fails to reconstruct the pattern of the data and has to choose between fitting some data or others in priority. In such cases, all the results should be considered carefully. When the confidence intervals between the different inversion techniques fail to overlap, it is likely that one or several estimates are incorrect. As algorithms are built by assuming a prior error statistics, it is likely that some rely on wrong assumptions.

3.2 Model concentrations and residuals

- The residuals, i.e. the differences between model concentrations and observations, determine how well the model reproduces the observations and quantify the improvements in the tracer distribution provided by the inversion. They are also a basic tool to identify biases in the model and to assess the quality of the assumptions.

- Radium fluxes obtained by the inverse method largely improve the model match to observations compared to the prior radium flux (Fig. 4 and Fig. 5). The improvement is quantified by the increase in the model–data correlations (Table 2) and the decrease in the root mean square of the residuals (Table 3), a proxy of the cost function. The correlation coefficient is increased from 0.383 to 0.813 on a linear scale and from 0.809 to 0.902 on a logarithmic scale. The correlation is higher on a logarithmic scale because it is less sensitive to the few very high residuals associated with the highest concentrations (see Fig. 67). On average, the inversion is able to reduce the ordinary residuals (C_{ols}) by a factor 2, logarithmic residuals (C_{log}) by 1.4 and proportional residuals (C_{prp}) by 3.5.

In spite of being smaller, the order of magnitude of the residuals remains comparable to the data (Fig. 5). On the one hand, in all oceans, positive and negative residuals are observed with no clear patterns at the scale of a few grid cells. Because of

the rather low model resolution (2° , which is not sufficient to reproduce medium and small scale processes) and of issues associated with temporal (seasonal or higher frequency) variability of the data, this kind of "noise" is expected. It is consistent with the assumption of independent errors used when computing the error variances on fluxes. On the other hand, in several regions, residuals can display coherent large scale patterns, which cannot be attributed to noise. These areas may suffer from systematic overestimations, like in the Gulf Stream region, the western Pacific between 20°N and 40°N , and off Eastern Siberia, or underestimations, such as in the center of the North Atlantic Gyre. These residuals point out to possible flaws in the model circulation. For instance, ^{228}Ra is quite homogeneously distributed in the western North Atlantic according to data, but in the model, the gradients are stronger, and no combination of sources manages to reduce these gradients. This is probably caused by a bias in the Atlantic circulation, with a too low exchange rate between inshore and offshore waters. In a 2° resolution model, mesoscale eddies are not represented and cannot transport ^{228}Ra south and east of the Gulf Stream or north of the North Brazil Current. Inversions minimize the misfit by increasing the fluxes from regions 5 (North Brazil), 8 (Caribbean) and 10 (southern East Coast of the US), making the model concentration too high close to the coast while still too low in the gyre. Such large scale biases are not consistent with the assumption of no prior error correlation, which may lead to underestimation of flux uncertainties around these basins.

Having assumed specific prior error statistics when choosing the cost functions, we need to check that there is no a posteriori contradiction. Figure 6 displays the residuals and model concentrations as a function of the observations. If the residuals depend on the observed concentrations, it means some observations are more precise than others, contain more information, and should be given a higher weight in order to obtain the best linear unbiased estimate. Figure 7 shows the probability density functions (PDF) of residuals after all three inversions and compares them with a Gaussian curve representing the expected distribution given the root mean square of residuals. Each PDF should look like a Gaussian curve for the computed posterior uncertainties to be relevant descriptors of errors.

Figure 7a emphasizes that the ordinary residuals do not follow a Gaussian distribution. On the contrary, most residuals are very close to zero ~~while~~~~and only~~ a small number of them is much higher than the standard deviation. Figure 6a shows that these high residuals occur at high concentrations only, and that error variance is not homogeneously distributed. Then high and low concentrations should not be given the same weights, as in C_{ols} , but the highest concentrations should be given the lowest weights, as in C_{log} and C_{prp} . Flux estimates based on C_{ols} are biased because the cost function puts more emphasis on high concentrations, and this method then tries to fit more specifically the misfits at high concentrations. C_{ols} also produces very large error bars because the error variance is assumed to be constant and its computed value, influenced by a few very large residuals, is larger than the actual error variances of the vast majority of data. Figure 6c and Fig. 7c show that the proportional residuals are not normally distributed either. They are not even symmetrical, as they cannot be smaller than -1 but they do not have an upper limit. This asymmetry produces a bias in the flux estimate. The algorithm based on C_{prp} is more sensitive to positive residuals because underestimations are never associated to proportional residuals lower than -1 whereas overestimations can produce residuals higher than 1. As a consequence, this method tends to reduce the fluxes. The hypothesis of constant variance is more realistic although the highest residuals occur at low concentrations. Finally, the distribution of the

logarithmic residuals displayed on 6b) and 7b) is much closer to a Gaussian curve and much less dependent on concentrations, which makes the logarithmic cost function more relevant for this study.

3.3 Sensitivity to the number of regions

The choice of the regions (and their number) has been made rather subjectively, although several criteria have been used (spatial distribution of the observations, independence of the A_i fields). The global ^{228}Ra should ideally not depend upon the number of regions. Therefore, alternative region geometries have been tested for comparison. The ^{228}Ra fluxes are shown on Table 4 and the root mean square of their residuals is presented on Table 5. Case 1 inversion uses 52 regions: It was the original distribution of regions before some of them were merged to define the 38 standard regions of this study. It includes more regions in undersampled areas such as the Arctic, the South Atlantic, the western Indian or the equatorial Pacific. Case 5 has just one emitting region for each of the following ocean basins: Southern, South Atlantic, North Atlantic, South Pacific, North Pacific, Indian and Arctic. Case 3 and Case 4 are intermediate cases with source regions built by merging regions from Case 2. The root mean square of residuals is a proxy of the cost function. On the one hand, this parameter should be as low as possible. Increasing the number of regions always decreases it because the number of degrees of freedom increases, which tends to improve the fit to the observations. In this inversion, the largest decrease is found between 7 and 12 source regions. Further increases in the number of regions have smaller impacts. On the other hand, too many source regions may produce spurious results. Some regional fluxes, with too few observations nearby to constrain them, would be computed using observations farther away, already used by other fluxes. Because of the lower sensitivity of the concentrations at these farther-off locations, this process can create extreme fluxes, positive or even negative. The presence of physically impossible negative values, set to zero by the constraint of positivity, necessarily means such poor constraints exist. When 52 fluxes are computed, 5 to 7 of them, according to the cost function, are so poorly constrained that their fluxes have been set to zero to prevent them from being negative. This number is reduced to 1 with C_{ols} and zero with C_{log} and C_{prp} when there are 38 regions, and completely disappears with 19 or fewer regions. Regions with fluxes within the error estimate are also very poorly constrained by the observations and the circulation model: their number is also reduced from 26 out of 52 to 11 out of 38 with C_{ols} , from 13 to 3 with C_{prp} and from 9 to 3 with C_{log} . All these fluxes have a low impact on the cost function, but make the global ^{228}Ra flux less precise. This analysis shows that Case 1 (with 52 regions) is not constrained enough and that Cases 3 to 5 display too large residuals. Therefore, Case 2 (38 regions) is considered to be the best choice.

The C_{ols} -based global ^{228}Ra flux is varying in a non-monotonic way, with a difference of 23% between the highest and the lowest. ~~The flux with 38 regions is lower than the fluxes computed with either more or fewer regions~~ ~~With 38 regions, it is lower than fluxes computed with both a higher and a lower number of regions.~~ This high sensitivity may be related to the high uncertainty associated with this inversion, certainly linked to the relatively poor data coverage. All the confidence intervals within one standard deviation from Case 1 to Case 4 overlap. The C_{prp} -based global fluxes are always lower than the other fluxes, and they decrease as the number of regions decreases. This is consistent with our previous hypothesis on C_{prp} . This cost function tends to fit the lowest data in priority. Larger regions suffer more from this bias because they are constrained by more data, likely to be more dispersed. The confidence intervals within one standard deviation based on C_{prp} fail to overlap.

Only the logarithmic least-squares method produces very similar fluxes whatever the number of regions, with all confidence intervals overlapping. The global flux based on C_{log} again seems to be the most reliable.

3.4 Submarine groundwater discharge estimates

The shelf fluxes after inversion combine groundwater discharge, riverine particles, diffusion from sediments and bioturbation.

- 5 Here we deduce the contribution from groundwater discharge by using existing estimates of the other sources of radium. Rivers are poor in dissolved ^{228}Ra and transport ^{228}Ra mainly with the sediments they carry (Moore and Shaw, 2008). According to Moore et al. (2008), the average dissolved ^{228}Ra activity in rivers is $0.65 - 1.95 \times 10^5 \text{ atoms.l}^{-1}$. As rivers annually discharge 35000 km^3 of freshwater (Milliman, 2001), the dissolved riverine source lies between 2.3 and $6.8 \times 10^{21} \text{ atoms.yr}^{-1}$, which is less than 1% of the total flux. Nearly all the particulate radium is desorbed in the mixing zone,
- 10 because of the salinity increase. According to various studies (Key et al., 1985; Moore et al., 1995; Krest et al., 1999), the amount of ^{228}Ra desorbed per gram of sediment lies in the range of 2.9 to $8.7 \times 10^6 \text{ atoms}$. In this study, we follow Milliman (2001) who proposed a global river sediment flux of $1.8 \times 10^{16} \text{ g.yr}^{-1}$, divided into fluxes from six basins: South Atlantic, North Atlantic, South Pacific, North Pacific, Indian Ocean and Arctic. This leads to a global ^{228}Ra river flux estimate of $0.53 - 1.60 \times 10^{23} \text{ atoms.yr}^{-1}$, significant but not dominant.
- 15 **Radium-228** is released from the sediments by diffusion, bioturbation and advection, the latter being associated with the SGD. Like Moore et al. (2008), we assume that the ^{228}Ra fluxes by diffusion and bioturbation from relict sands, composing 70% of the total continental shelf area, is weak, typically of the order of $10^9 \text{ atoms m}^{-2} \text{ yr}^{-1}$ (Colbert, 2004; Hancock et al., 2006). Fluxes from continental shelf muds, which correspond to the remaining 30%, have been estimated by several studies using different methods (Table 6), such as inventories (Moore et al., 1995), benthic chambers (Hancock et al., 2000), sediment
- 20 profiles (Hancock et al., 2000) or modeling (Hancock et al., 2006). But some of them were done at a time when SGD were not considered to be an important source of radium and may have included them in their estimates. These estimates should then be considered as an upper limit. Furthermore, their locations are often very close to the coast. Some recent studies addressing diffusive fluxes (from muds or broader regions containing both types) separately, in order to estimate local groundwater discharge (Crotwell and Moore, 2003; Kim et al., 2005) used the simplified equation of diffusion from Krest et al. (1999)
- 25 (using only water diffusion, sediment porosity, decay constants and production rates) and pointed out to significantly lower values, but may not be representative of all continental shelves. So far, it is not possible to be very precise and a wide range has to be considered: From the low recent values, close to $5 \times 10^9 \text{ atoms.m}^{-2} \text{ .yr}^{-1}$, to the higher values typically ranging from 25 to $75 \times 10^9 \text{ atoms m}^{-2} \text{ yr}^{-1}$ (Moore et al., 2008). The full range is then $5 - 75 \times 10^9 \text{ atoms.m}^{-2} \text{ .yr}^{-1}$. As the continental shelf area in our model is $2.73 \times 10^{13} \text{ m}^2$, this means a total flux due to diffusion and bioturbation ranging from 0.43
- 30 to $6.51 \times 10^{23} \text{ atoms.yr}^{-1}$. Then, using the logarithmic cost function, the SGD ^{228}Ra flux estimate varies between 0.62 and $6.82 \times 10^{23} \text{ atoms.yr}^{-1}$: these two fluxes are within the same range.

Figure 8 shows the ^{228}Ra fluxes for seven basins corresponding to the Southern Ocean and the six basins used by Milliman (2001) to define sediment inputs by rivers. The largest ^{228}Ra fluxes are found in the North Atlantic, the North Pacific and the Indian Ocean, in roughly equal proportions, whereas those from the South Atlantic, the South Pacific and the Arctic Ocean

are far smaller. Fluxes from the three latter regions are low enough in fact to be within the confidence intervals of riverine and diffusive sediment ^{228}Ra fluxes, so that the SGD confidence intervals include negative values, which are physically impossible. SGD ^{228}Ra end-member concentration varies considerably from one aquifer to another, ranging from 0.04 to 125×10^6 atoms m^{-3} (Moore et al., 2008). Measurements so far have shown aquifer concentrations in the Atlantic Ocean higher than the average by typically 30% (Kwon et al., 2014). Following Kwon et al. (2014), we assume that the aquifer concentrations are log-normally distributed with an average of 0.98 to 1.15×10^3 dpm m^3 . Taking a geometric mean rather than an arithmetic mean is implicitly assuming that aquifers characterized by high ^{228}Ra concentrations are emitting less water. It also suffers from wells being concentrated in developed countries. Using the results from the inversion technique based on the logarithmic cost function, the total SGD flux is estimated to $1.3 - 14.7 \times 10^{13}$ $\text{m}^3 \text{yr}^{-1}$, to be compared with the global river flow of 3.5×10^{13} $\text{m}^3 \text{yr}^{-1}$ (Milliman, 2001). As long as diffusion cannot be separated from SGD more efficiently and is not shown to be negligible, no precise SGD estimate based on ^{228}Ra can be produced. According to Taniguchi et al. (2007), there is only 2.6×10^{12} $\text{m}^3 \text{yr}^{-1}$ of fresh groundwater discharge, representing between 2% and 20% of the SGD, and the remaining part is recirculated seawater. In total, 63% of the ^{228}Ra flows into the Indian and Pacific basins, which is more than their share of the continental shelf (52%). As shown in Fig 8, the estimated contributions of SGD to ^{228}Ra fluxes are in very similar proportions, with 68% going to the Indian and Pacific basins. This represents more than 70% of all the groundwater discharge when the differences in groundwater ^{228}Ra concentration are considered. However, uncertainties on SGD distribution are high, because diffusion from sediment might not be identical in all basins. These results contrast with the other source of water, nutrients and trace elements to the ocean: 60% of the global river discharge flows to the Atlantic and Arctic basins (Milliman, 2001).

4 Discussion

4.1 Comparisons with previous studies

Our inversion estimates are in good agreement with previous regional studies of ^{228}Ra based on inventories. The ^{228}Ra inventory of the Mediterranean Sea computed by Rodellas et al. (2015) led to a total flux due to rivers, sediments and groundwater discharge of $1.86 - 2.48 \times 10^{22}$ atoms yr^{-1} . This is compatible with our estimate based on C_{log} , $1.96 - 2.28 \times 10^{22}$ atoms yr^{-1} . Other cost functions produce an underestimation in the model by a factor 2.5 (ols) or 3.8 (prp). In the Yellow Sea, Kim et al. (2005) estimated a flux of 3.3×10^{15} dpm yr^{-1} , or 1.4×10^{22} atoms yr^{-1} . Expressed per unit of surface, it corresponds to 3.6×10^{10} atoms $\text{m}^{-2} \text{yr}^{-1}$. This is slightly lower than our flux for the larger region 23 (Sea of Japan, Yellow Sea, East China Sea) of $4.2 - 5.6 \times 10^{10}$ atoms $\text{m}^{-2} \text{yr}^{-1}$ based on C_{log} , both these fluxes being larger than the global average. At larger scale, Moore et al. (2008) estimated the total ^{228}Ra flux over the Atlantic between 50°S and 80°N to be $2.8 - 4.2 \times 10^{23}$ atoms yr^{-1} . Restricting the inversion to the Atlantic with C_{log} yields $2.64 - 2.92 \times 10^{23}$ atoms yr^{-1} , in the lower range but compatible. At least two reasons can explain the difference. Moore et al. (2008) have included data down to 1000 m, which allowed them to estimate ^{228}Ra release from sediments down to that depth. The authors have estimated the sources between 200 m and 1000 m deep to $0.13 - 0.37 \times 10^{22}$ atoms yr^{-1} . Furthermore, in order to compute the total ^{228}Ra content of the Atlantic, they have performed a linear interpolation of the data, potentially leading to errors, especially in areas where measurements are sparse.

At the global scale, Kwon et al. (2014) have used a method which is quite similar to ours and they have computed a total global ^{228}Ra flux of $9.1 - 10.1 \times 10^{23} \text{ atoms yr}^{-1}$, of which between 4.2 and $7 \times 10^{23} \text{ atoms yr}^{-1}$ are released by SGD. This corresponds to a global SGD flux of $9 - 15 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$. The flux of ^{228}Ra estimated by the present study is thus significantly lower than the estimates by Kwon et al. (2014), although our results include the Arctic and Antarctic sources ($0.43 - 0.50$ and $0.28 - 0.35 \times 10^{23} \text{ atoms yr}^{-1}$ respectively). Kwon et al. (2014) minimized an ordinary least-squares cost function with 50 regions. We have shown here that both a high number of source regions and the use of the ordinary least-squares cost function concur to produce a higher estimate. However, this is at the expense of a higher uncertainty and of producing unrealistic negative fluxes in some source regions. Additional differences in Kwon's study may explain their higher estimates such as a different ocean circulation model, a coarser vertical resolution, and a bathymetry re-gridded onto the model domain. Finally, they added dust deposition and removed data higher than 140 dpm m^{-3} , but these two factors should rather tend to reduce their fluxes.

As recently proposed by Charette et al. (2016), shelf ^{228}Ra fluxes can be used as gauges of shelf fluxes of trace elements and isotopes, including nutrients, iron, and rare earth elements. ^{228}Ra is particularly relevant because it is chemically conservative and integrates information over annual to decadal timescales. At first approximation, the flux of TEIs is deduced from the ^{228}Ra flux and the ratio of the nearshore gradients. Limited for now due to the lack of nearshore measurements, this method could be more common in the future. As they are based on realistic assumptions on error statistics and have low uncertainties, our radium fluxes are able to improve the current estimates of all elements originating from the continental shelf.

On the contrary, our uncertainties on groundwater discharge are large, even when compared to previous estimates. These larger uncertainties stem from the poor knowledge of the non SGD sources of radium that we subtract from the total flux. As diffusion and bioturbation are expressed in flux per area, the mean SGD fluxes and their uncertainties depend to a large extent on the radium emitting area we consider. Based on a more realistic refined bathymetry than Kwon et al. (2014) ($2.73 \times 10^{13} \text{ m}^2$ compared to $1.5 \times 10^{13} \text{ m}^2$), our study also has a larger sedimentary flux, with an upper range close to the total ^{228}Ra flux. Thus, the lower range of SGD fluxes, $1.3 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$, is very low, while the upper range, $14.7 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$, is similar to other studies. The use of this model provides an upper estimate but cannot precisely compute the global SGD flux, and no inverse model can if the surface diffusive flux and the area emitting radium are not clarified. Our expectation is that the lower range of sedimentary flux is more likely, because it comes from studies where SGD is also considered and because it does never produce negative regional fluxes. By contrast, the higher range is very similar to the total bottom flux, as expected when no distinction is made between diffusion and SGD.

Comparisons with local direct estimates of submarine groundwater discharge based on seepage meters and piezometers are also possible but less conclusive because of the high spatial variability of SGD. Our global average SGD fluxes lie between 0.5 and 5.5 m yr^{-1} . Values from seepage meters and piezometers in the upper 200 m reported by Taniguchi et al. (2002) and Knee and Paytan (2011) range from 0.03 m yr^{-1} in the Tokyo Bay (Taniguchi et al., 2002) to 1790 m yr^{-1} near Mauritius (Burnett et al., 2006). Most measurements have been performed in shallow coastal areas, less than 10m deep, the upper 10 meters and range from 1 to 50 m yr^{-1} , one order of magnitude higher than our averages on all the seafloor down to 200m deep. Thus, this range is very wide and many local studies span several orders of magnitude. At a global scale, Taniguchi et al. (2009) produced

an estimate of the global SGD flux, ~~based on seepage measurements of~~ $6.1 - 12.8 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$, in the upper range of our global estimate, ~~by extrapolating seepage measurements~~. However, it suffers from most measurements being concentrated in developed countries.

The spatial distribution of the fluxes in this study is consistent with Kwon et al. (2014), with two thirds of radium-228 flowing to the Indian and Pacific Basins and also quite high fluxes in the western North Atlantic. It turns out that these basins have the highest riverine sediment loads (Milliman, 2001). However, we have shown that rivers could account for 7% to 20% of ^{228}Ra fluxes only, which does not exclude an indirect impact on the other sources through the geology of the continental shelf. A difference among basins in the quantity of radium diffusion, which is poorly known, is also possible. If it is not the case, then the SGD are significantly higher in East Asia than in other regions. SGD can be driven by as many factors as storms, waves, tides and thermal gradients, and depend on the permeability and structure of coastal and shelf sediments (Moore, 2010a). Comparative data on these factors in several basins will be necessary to explain the origin of the observed differences.

4.2 Model biases

Our model of ^{228}Ra is based on a circulation model and assumptions on the cycle of this isotope. Both are potential sources of errors.

Dust has not been included in the model. The model then replaces them with other sources, potentially leading to an overestimation. At global scale, it represents $1.7 \times 10^{21} \text{ atoms yr}^{-1}$ (Kwon et al., 2014), less than 0.2% of the total flux and less than most individual source regions, which cannot create large biases. Nevertheless, the largest dust deposition is associated to Saharan dust transported to the North Atlantic, in the Canary Upwelling Region (Mahowald et al., 2005). In this region, dust may have some impact on the ^{228}Ra distribution, since it brings this isotope directly to the open ocean. Its absence in our study might explain the overestimation by our model near the coast of region 6 and underestimation offshore. Yet, the exclusion of dust deposition in our analysis cannot explain the largest bias in the North Atlantic: The overestimation in the Gulf Stream coupled with an underestimation just south east (see 3.2), because the area where the model cannot transport radium is located west of the maximum of dust input and displays higher concentration.

Scavenging is a neglected potential sink in this study. As the residence time related to scavenging is approximately 500 yr (Moore and Dymond, 1991), it accounts for between 1% and 2% of all sinks. Thus, its inclusion in our computations would increase the source intensity necessary to maintain global balance. Contrary to radioactivity, scavenging is highly heterogeneous, most intense where primary productivity and particle concentrations are highest. Fluxes from the high latitudes of the North Atlantic are thus potentially more underestimated than fluxes in other regions, since they are areas of intense biological activity during spring blooms. As the actual total lifetime of ^{228}Ra is overestimated when scavenging is not taken into account, the gradient between coast and open ocean could be too low. However, as the horizontal mixing time scale of the ocean is a few decades, the relative overestimation of open ocean concentrations is less than 10%.

The contribution of rivers to radium fluxes is considered when estimating the SGD, but only at a basin-wide scale. As most riverine ^{228}Ra travels attached to particles and is desorbed in the mixing zone, we have based our computation on the sediment loads of Milliman (2001), which are basin-wide. Although NEMO 3.6 takes rivers into account for their impact on salinity

(Madec, 2015), for now no information at the model grid scale is available on their sediment loads. If this information existed, we would have been able to estimate the contribution of rivers, and consequently sediments and SGD, in the ^{228}Ra fluxes for each of the 38 regions. Some local SGD fluxes close to large rivers (Amazon, Congo, Yellow River, etc.) could then appear to be significantly lower than their shares of the total ^{228}Ra fluxes suggest.

- 5 The other part of the model is the circulation model. The climatological circulation of NEMO 3.6 was not optimized in this study. However, the residuals after inversion show that some regions are associated with spatially structured residuals. There are good reasons to incriminate the ocean circulation. Because of the low resolution of the model (2°), isopycnal diffusion has been used to parameterize sub grid processes and mixing (Redi, 1982) with a constant eddy diffusivity of $2000 \text{ m}^2 \text{ s}^{-1}$. Nevertheless, in very energetic regions, such as the Western Boundary Currents (WBCs, e.g. the Gulf Stream and the Kuroshio),
- 10 higher eddy diffusivity might be needed to enhance the exchanges with the open ocean, possibly improving the fit with the observations. Furthermore, it is well known that the mean currents are also dependent on this low resolution (impacting for instance the position and intensity of the WBCs). Although it cannot solve all the flaws of the circulation, improvements could thus be brought by an increase of the resolution towards eddy resolving models. However, as 100 years of simulation are needed at a global scale for each source region, this would be computationally expensive.
- 15 Other sources of errors are the four statistical assumptions on the errors: errors are assumed to have zero expectation, no correlation, a normal distribution and variances depending on the concentrations in a way specific to each cost function. Systematic biases on data or model are not corrected by least-squares algorithms. They increase or decrease values without leaving clues. However, the model conserves mass: the quantity of radium present in the ocean from each model tracer is precisely known and if concentration is too high at some place, it will be too low elsewhere. As for the measurements, their uncertainty is
- 20 generally around 10% or lower, and cruises take them independently from each other, making the assumption of zero expectation on the observational errors reasonable. The second assumption is the absence of prior correlation. If prior uncertainties of neighboring data are correlated, it means that the errors are likely to be of the same sign whatever the solution, and that multiplying measurements in this area does not multiply information proportionally. Where measurements are dense, with residuals far from the expected white noise, for instance in the North Atlantic, there may be correlation and uncertainties may
- 25 be underestimated. The last assumptions are about the structure of variance. We have shown that logarithmic residuals were almost normally distributed and independent from concentrations (see 3.2.), justifying the choice of C_{log} as a cost function. C_{ols} leads to higher uncertainties, especially for small sources, and C_{prp} to systematic underestimation, but both are useful for comparison, in order to identify regions where physical assumptions are inaccurate (see 3.1.).

4.3 What new data would be most useful ?

- 30 Observations are not evenly distributed. Some coastal regions cannot be constrained properly because ^{228}Ra data are lacking. Improving the coverage would increase the quality of the inversion in two ways: It would reduce the uncertainties and make it possible to divide wide regions into smaller regions as long as they have distinct footprints. For instance, the Philippines, Papua, or the Gulf of Guinea, whose footprints are very different from the Indonesian Seas and the southwestern coast of Africa, could be considered as independent regions. More samples in the Indian Ocean, the South Atlantic (south of 30°S), the

Southern Ocean and the western equatorial Pacific are priorities. All these regions will be sampled by upcoming GEOTRACES cruises shown on Fig. 98. At the same time, deep samples will be taken outside of the Atlantic, enabling a more comprehensive global inversion with extra source regions at greater depths. This would improve our knowledge of the contribution of deeper sediments.

- 5 Most direct submarine groundwater discharge measurements have been performed in developed countries, with a focus on the North Atlantic and the Mediterranean Sea (Taniguchi et al., 2002). At the same time, measurements are completely lacking over large regions. More SGD studies in areas where they are potentially the highest, namely the Bay of Bengal, the Indonesian Seas and the China Seas, would produce more representative estimates of the ^{228}Ra content of SGD around the world and direct estimates of local SGD magnitude to be compared with regional inversion results. They have begun more recently (Knee and
- 10 Paytan, 2011) and are still sparse.

Information contained in ^{228}Ra might be completed with ^{226}Ra concentrations, measured during the same campaigns and for this reason available with a similar coverage. Associated with the same source as ^{228}Ra , but with a much longer half-life, 1602 yr, ^{226}Ra would constrain deeper sources and would help in assessing the quality of the thermohaline circulation and deep ventilation of the circulation model. However, inverting ^{226}Ra data would require a precise modeling of scavenging,

15 which is not negligible at these longer time scales, as well as a much longer integration duration in order to get steady state distribution.

5 Conclusions

- Based on inverse modeling, we have computed a global ^{228}Ra flux from continental shelves of $8.01 - 8.49 \times 10^{23}$ atoms yr^{-1} , with the largest sources in the western Pacific, the western North Atlantic and the Indian Oceans and the smallest sources in the
- 20 eastern Pacific. The Arctic and Antarctic sources have been estimated for the first time, accounting for $0.43 - 0.50$ and $0.28 - 0.35 \times 10^{23}$ atoms yr^{-1} respectively. These precise estimates are obtained by **minimizing the squared differences between model and observed concentrations on a logarithmic scale. We think this cost function is more realistic than a linear least-squares one because it assumes that the error standard deviation is proportional to ^{228}Ra concentration, rather than constant, which proves to be more accurate. It is also better than a "proportional" cost function, weighted by the inverse of the observed**
- 25 **concentration, which tends to produce underestimations of concentrations and fluxes.**~~minimizing the squares of logarithmic concentrations between model and data, a cost function we think is more realistic than the other functions we have tested because it is based on more realistic assumptions on error statistics.~~ Given the number of available measurements, we were able to constrain 38 regional fluxes. The shelf fluxes produced using these optimal parameters are **around 20%** lower than previous estimates. In a near future, they will enable to quantify continental shelf fluxes of trace elements and isotopes to the
- 30 oceans at any place where nearshore gradients are measured (Charette et al., 2016). The estimated global SGD ~~flux~~ is far less precise, ranging between 1.3 and 14.7×10^{13} $\text{m}^3 \text{yr}^{-1}$, because of the very large uncertainty on the two other sources of ^{228}Ra , i.e. riverine particles and most of all diffusion from bottom sediments, also located on the continental shelf. Only the upper range is compatible with previous estimates. **These results mean that all the studies which have used or will use**

²²⁸Ra to estimate SGD will suffer from a very high uncertainty until diffusion is properly estimated or proven to be negligible.

After inversion, we were able to reproduce the basin scale patterns of ²²⁸Ra distribution with nevertheless systematic biases in several regions, especially in the Arctic, and west of the sub-tropical gyres. Shortcomings in the circulation model are the most probable explanation of these biases (too weak exchanges between continental shelves and open ocean). ~~Therefore,~~

5 ~~besides estimating the sources and sinks of tracers, interesting information about the ocean model can be obtained by Ra like tracer inversions.~~ Extensive regions are lacking observations, mostly in the Southern Hemisphere (Pacific, Indian, and mostly Southern Ocean, as well as western equatorial Pacific Ocean), and better coverage in basins where SGDs are known to be influential and to produce large horizontal gradients is also needed (such as in South Asia). But the main impediment to achieve precise estimates of global Ra SGD fluxes comes from the very poor knowledge of diffusive sedimentary fluxes:
10 without a proper way to separate diffusion and SGD, inversions can compute the total bottom flux but are not able to precisely evaluate these two components.

6 Code and Data availability

The source code of NEMO is available on the NEMO website (<http://www.nemo-ocean.eu>). Studies providing data are listed in Table S1 in the supplementary materials. The inversion code and data used in this study can be obtained directly by contacting
15 the authors.

Appendix A: Error Statistics

This section describes the way regional ²²⁸Ra flux estimates and their error bars are computed. The inverse problem is the following one : using p measurements of oceanic ²²⁸Ra concentrations, their n sources must be traced back, by means of a circulation model. The model produces n ²²⁸Ra concentration fields, one from each of the source regions. It is run through
20 steady state, the ith field reaching concentration A_i . The model concentration is a linear combination of the latter:

$$[^{228}\text{Ra}]^{\text{mod}} = \sum_{i=1}^n A_i x_i$$

Inversion requires fitting the model concentration to the observations. As the expression of model concentration is linear in terms of the unknowns, the problem can be written:

$$\mathbf{Ax} = \mathbf{b} + \boldsymbol{\eta}$$

25 where $\mathbf{b} \in \mathcal{M}_{p,1}$ is the vector of the p measured radium concentrations, $\mathbf{x} \in \mathcal{M}_{n,1}$ is the unknown and the vector of the n source intensities, $\mathbf{A} \in \mathcal{M}_{p,n}$ is the footprint matrix, composed of the A_i at each data point, and $\boldsymbol{\eta} \in \mathcal{M}_{p,1}$ is the vector of residuals, accounting for the misfits, which are inevitable in our over-constrained problem ($p > n$).

We want to minimize the distance between data and model concentrations, summed up in a cost function. The ordinary least-squares cost function we first used is given by:

$$30 \quad C_{ols}(\mathbf{x}) = (\mathbf{Ax} - \mathbf{b})^\top (\mathbf{Ax} - \mathbf{b})$$

The cost function has to be minimal at the optimal x , called x_{opt} , implying, in the absence of inequality constraints:

$$\nabla C_{ols}(x_{opt}) = 2\mathbf{A}^\top (\mathbf{A}x - \mathbf{b}) = \mathbf{0}$$

As the n ^{228}Ra concentration fields are independent from each other, $\mathbf{A}^\top \mathbf{A}$ is invertible. Then:

$$x_{opt} = \mathbf{A}_{inv} \mathbf{b} \tag{A1}$$

5

$$\mathbf{A}_{inv} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top$$

\mathbf{A}_{inv} is the pseudo-inverse of \mathbf{A} , transforming the consequences into their most probable causes and justifying the word "inversion". Equation (A1) corresponds to (2.95) in Wunsch (2006). In practice, fluxes are constrained to be positive. This is managed by just reducing the number of source regions if necessary, without changing the principle of the inversion.

10 Matrix \mathbf{H} ~~issi~~ defined by :

$$\mathbf{H} = \mathbf{A}(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top = \mathbf{A} \mathbf{A}_{inv}$$

As $\mathbf{H}^2 = \mathbf{H}^\top = \mathbf{H}$, \mathbf{H} is an orthogonal projector on a subspace of dimension n . As $\mathbf{A}x_{opt} = \mathbf{H}\mathbf{b}$, the model concentration is the projection of the observations on this subspace whereas the vector of residuals is the projection on the complementary subspace.

15 Simple algebraic transformations yield:

$$\mathbf{H}\mathbf{A} = \mathbf{A}(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{A} = \mathbf{A}$$

$$\mathbf{A}_{inv} \mathbf{A} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{A} = \mathbf{I}_n$$

$$\mathbf{A}_{inv} \mathbf{A}_{inv}^\top = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{A} (\mathbf{A}^\top \mathbf{A})^{-1} = (\mathbf{A}^\top \mathbf{A})^{-1} \tag{A2}$$

Uncertainties on x resulting from uncertainties on data (\mathbf{b}) and model (\mathbf{A}) must now be evaluated. By principle of the ordinary

20 least-squares cost function, the error covariance of $(\mathbf{A}x - \mathbf{b})$ is considered to be a constant diagonal matrix:

$$\langle (\mathbf{A}x - \mathbf{b})(\mathbf{A}x - \mathbf{b})^\top \rangle = \sigma^2 \mathbf{I}_p$$

The posterior covariance matrix of x is:

$$\mathbf{V}_{xx} = \langle (x - x_{opt})(x - x_{opt})^\top \rangle$$

$$= \langle (\mathbf{A}_{inv} \mathbf{A}x - \mathbf{A}_{inv} \mathbf{b})(\mathbf{A}_{inv} \mathbf{A}x - \mathbf{A}_{inv} \mathbf{b})^\top \rangle$$

25

$$= \mathbf{A}_{inv} \langle (\mathbf{A}x - \mathbf{b})(\mathbf{A}x - \mathbf{b})^\top \rangle \mathbf{A}_{inv}^\top$$

$$= \sigma^2 \mathbf{A}_{inv} \mathbf{A}_{inv}^\top \tag{A3}$$

Equation (A3) corresponds to (2.102) in Wunsch (2006).

σ^2 is related to the root mean square of residuals v_η the following way:

$$\begin{aligned}
v_\eta &= \langle (\mathbf{A}\mathbf{x}_{opt} - \mathbf{b})^\top (\mathbf{A}\mathbf{x}_{opt} - \mathbf{b}) \rangle \\
&= \langle (\mathbf{H}\mathbf{b} - \mathbf{b})^\top (\mathbf{H}\mathbf{b} - \mathbf{b}) \rangle \\
5 \quad &= \langle ((\mathbf{I}_p - \mathbf{H})(\mathbf{A}\mathbf{x} - \mathbf{b}) + (\mathbf{H}\mathbf{A} - \mathbf{A})\mathbf{x})^\top ((\mathbf{I}_p - \mathbf{H})(\mathbf{A}\mathbf{x} - \mathbf{b}) + (\mathbf{H}\mathbf{A} - \mathbf{A})\mathbf{x}) \rangle \\
&= \langle (\mathbf{A}\mathbf{x} - \mathbf{b})^\top (\mathbf{I}_p - \mathbf{H})(\mathbf{A}\mathbf{x} - \mathbf{b}) \rangle \\
&= \sigma^2 \frac{p-n}{p}
\end{aligned} \tag{A4}$$

Then, using Eqs. (A2), (A3) and (A4), we can compute \mathbf{V}_{xx} :

$$\mathbf{V}_{xx} = \frac{p}{p-n} v_\eta (\mathbf{A}^\top \mathbf{A})^{-1} \tag{A5}$$

- 15 The diagonal terms of \mathbf{V}_{xx} are the squared uncertainties on \mathbf{x} whereas the other terms are covariances, indicating source regions with similar footprints.

The result can be extended to cases where the cost function is weighted, such as the proportional least-squares inversion, by just normalizing matrix \mathbf{A} and vector \mathbf{b} . Extension to logarithmic inversion is done by linearizing $\log(\mathbf{A}\mathbf{x}) - \log(\mathbf{b})$ to $\mathbf{F}\mathbf{x} - \beta$ with $\beta = \log(\mathbf{b}) + \mathbf{F}\mathbf{x}_{opt} - \log(\mathbf{A}\mathbf{x}_{opt})$, assuming a small error.

- 15 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. The authors thank all the scientists who produced the data used in this article. We thank Matt Charette, Eun Young Kwon, Virginie Sanial and Pieter van-Beek for helping us putting the data together. We also thank Olivier Marchal for a discussion about algorithms. This work is part of the first author's PhD, supported by the "Laboratoire d'Excellence" LabexMER (ANR-10-LABX-19) and co-funded by a grant from the French government under the program "Investissements d'Avenir", and by a grant from the Regional Council

- 20 of Brittany.

References

- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochemical Cycles*, 20, doi:10.1029/2005gb002591, <https://doi.org/10.1029/2005GB002591>, 2006.
- Buesseler, K. O., Bacon, M. P., Cochran, J. K., and Livingston, H. D.: Carbon and nitrogen export during the JGOFS North Atlantic Bloom
5 estimated from ^{234}Th : ^{238}U disequilibria, *Deep Sea Research Part I: Oceanographic Research Papers*, 39, 1115–1137, doi:10.1016/0198-0149(92)90060-7, [https://doi.org/10.1016/0198-0149\(92\)90060-7](https://doi.org/10.1016/0198-0149(92)90060-7), 1992.
- Burnett, W., Aggarwal, P., Aureli, A., Bokuniewicz, H., Cable, J., Charette, M., Kontar, E., Krupa, S., Kulkarni, K., Loveless, A., and et al.: Quantifying submarine groundwater discharge in the coastal zone via multiple methods, *Science of The Total Environment*, 367, 498–543, doi:10.1016/j.scitotenv.2006.05.009, <https://doi.org/10.1016/j.scitotenv.2006.05.009>, 2006.
- 10 Charette, M. A., Lam, P. J., Lohan, M. C., Kwon, E. Y., Hatje, V., Jeandel, C., Shiller, A. M., Cutter, G. A., Thomas, A., Boyd, P. W., and et al.: Coastal ocean and shelf-sea biogeochemical cycling of trace elements and isotopes: lessons learned from GEOTRACES, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374, 20160076, doi:10.1098/rsta.2016.0076, <https://doi.org/10.1098/rsta.2016.0076>, 2016.
- Clegg, S. L. and Whitfield, M.: A generalized model for the scavenging of trace metals in the open ocean—II. Thorium scavenging,
15 *Deep Sea Research Part I: Oceanographic Research Papers*, 38, 91–120, doi:10.1016/0198-0149(91)90056-L, [https://doi.org/10.1016/0198-0149\(91\)90056-L](https://doi.org/10.1016/0198-0149(91)90056-L), 1991.
- Colbert, S. L.: Ra Isotopes in San Pedro Bay, CA: Constraint on Inputs and Use of Nearshore Distribution to Compute Horizontal Eddy Diffusion Rates., Ph.D. thesis, University of Southern California, 2004.
- Crotwell, A. M. and Moore, W. S.: Nutrient and Radium and Fluxes from Submarine and Groundwater Discharge to Port Royal Sound,
20 *South Carolina, Aquatic Geochemistry*, 9, 191208, doi:10.1023/b:aqua.0000022954.89019.c9:10.1023/b:aqua.0000022954.89019.c9, <https://doi.org/10.1023/b:aqua.0000022954.89019.c9>, 2003.
- DeVries, T. and Primeau, F.: Dynamically and Observationally Constrained Estimates of Water-Mass Distributions and Ages in the Global Ocean, *Journal of Physical Oceanography*, 41, 2381–2401, doi:10.1175/jpo-d-10-05011.1, <https://doi.org/10.1175/JPO-D-10-05011.1>, 2011.
- 25 Gloor, M., Gruber, N., Hughes, T. M. C., and Sarmiento, J. L.: Estimating Net Air-sea Fluxes from Ocean Bulk Data, *Global Biogeochemical Cycles*, 15, 767–782, doi:10.1029/2000GB001301, <https://doi.org/10.1029/2000GB001301>, 2001.
- Hammond, D. E., Marton, R. A., Berelson, W. M., and Ku, T.-L.: Radium 228 Distribution and Mixing in San Nicolas and San Pedro Basins, Southern California Borderland, *Journal of Geophysical Research*, 95, 3321–3335, doi:10.1029/jc095ic03p03321, <https://doi.org/10.1029/jc095ic03p03321>, 1990.
- 30 Hancock, G. J., Webster, I. T., Ford, P. W., and Moore, W. S.: Using Ra isotopes to examine transport processes controlling benthic fluxes into a shallow estuarine lagoon, *Geochimica et Cosmochimica Acta*, 64, 3685–3699, doi:10.1016/s0016-7037(00)00469-5, [https://doi.org/10.1016/s0016-7037\(00\)00469-5](https://doi.org/10.1016/s0016-7037(00)00469-5), 2000.
- Hancock, G. J., Webster, I. T., and Stieglitz, T. C.: Horizontal mixing of Great Barrier Reef waters: Offshore diffusivity determined from radium isotope distribution, *Journal of Geophysical Research*, 111, 1–14, doi:10.1029/2006jc003608, <https://doi.org/10.1029/2006JC003608>, 2006.
35

- Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F., and Quartly, G. D.: A reduced estimate of the strength of the oceans biological carbon pump, *Geophysical Research Letters*, 38, L04 606, doi:10.1029/2011gl046735, <https://doi.org/10.1029/2011GL046735>, 2011.
- Hwang, D.-W., Kim, G., Lee, Y.-W., and Yang, H.-S.: Estimating submarine inputs of groundwater and nutrients to a coastal bay using radium isotopes, *Marine Chemistry*, 96, 61–71, doi:10.1016/j.marchem.2004.11.002, <https://doi.org/10.1016/j.marchem.2004.11.002>, 2005.
- Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global Biogeochemical Cycles*, 21, GB1019, doi:10.1029/2005gb002556, <https://doi.org/10.1029/2005GB002556>, 2007.
- Jeandel, C., Arsouze, T., Lacan, F., Techine, P., and Dutay, J.: Isotopic Nd compositions and concentrations of the lithogenic inputs into the ocean: A compilation, with an emphasis on the margins, *Chemical Geology*, 239, 156–164, doi:10.1016/j.chemgeo.2006.11.013, <https://doi.org/10.1016/j.chemgeo.2006.11.013>, 2007.
- Kaufman, A., Trier, R. M., and Broecker, W. S.: Distribution of ^{228}Ra in the World Ocean, *Journal of Geophysical Research*, 78, 8827–8848, doi:10.1029/jc078i036p08827, <https://doi.org/10.1029/jc078i036p08827>, 1973.
- Key, R. M., Stallard, R. F., Moore, W. S., and Sarmiento, J. L.: Distribution and Flux of ^{226}Ra and ^{228}Ra in the Amazon River Estuary, *Journal of Geophysical Research*, 90, 6995–7004, doi:10.1029/jc090ic04p06995, <https://doi.org/10.1029/jc090ic04p06995>, 1985.
- Kim, G., Ryu, J.-W., Yang, H.-S., and Yun, S.-T.: Submarine groundwater discharge (SGD) into the Yellow Sea revealed by ^{228}Ra and ^{226}Ra isotopes: Implications for global silicate fluxes, *Earth and Planetary Science Letters*, 237, 156–166, doi:10.1016/j.epsl.2005.06.011, <https://doi.org/10.1016/j.epsl.2005.06.011>, 2005.
- Knee, K. and Paytan, A.: Treatise and on Estuarine and Coastal and Science, chap. Submarine Groundwater Discharge: A Source of Nutrients, Metals, and Pollutants to the Coastal Ocean, pp. 206–228, Elsevier, 2011.
- Krest, J. M., Moore, W. S., and Rama: ^{226}Ra and ^{228}Ra and in the mixing zones of the Mississippi and Atchafalaya Rivers and indicators of groundwater input, *Marine Chemistry*, 64, 129–152, doi:10.1016/s0304-4203(98)00070-x, [https://doi.org/10.1016/s0304-4203\(98\)00070-x](https://doi.org/10.1016/s0304-4203(98)00070-x), 1999.
- Kwon, E. Y., Kim, G., Primeau, F., Moore, W. S., Cho, H.-M., DeVries, T., Sarmiento, J. L., Charette, M. A., and Cho, Y.-K.: Global estimate of submarine groundwater discharge based on an observationally constrained radium isotope model, *Geophysical Research Letters*, 41, 8438–8444, doi:10.1002/2014gl061574, <https://doi.org/10.1002/2014GL061574>, 2014.
- LaRoche, J., Nuzzi, R., Waters, R., Wyman, K., Falkowski, P. G., and Wallace, D. W. R.: Brown tide blooms in Long Island’s coastal waters linked to inter-annual variability in groundwater flow, *Global Change Biology*, 3, 397–410, doi:10.1046/j.1365-2486.1997.00117.x, <https://doi.org/10.1046/j.1365-2486.1997.00117.x>, 1997.
- Madec, G.: NEMO ocean engine, Institut Pierre-Simon Laplace, 2015.
- Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochemical Cycles*, 19, 1–15, doi:10.1029/2004GB002402., <https://doi.org/10.1029/2004GB002402.>, 2005.
- Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., and et al.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean, *Global Biogeochemical Cycles*, 20, doi:10.1029/2005gb002530, <https://doi.org/10.1029/2005GB002530>, 2006.
- Milliman, J.: Encyclopedia of Ocean Sciences, chap. River Inputs, pp. 2419–2427, Elsevier BV, doi:10.1006/rwos.2001.0074, <https://doi.org/10.1006/rwos.2001.0074>, 2001.

- Moore, W. S.: Oceanic concentrations of ^{228}Ra , *Earth and Planetary Science Letters*, 6, 437–446, doi:10.1016/0012-821x(69)90113-7, [https://doi.org/10.1016/0012-821x\(69\)90113-7](https://doi.org/10.1016/0012-821x(69)90113-7), 1969.
- Moore, W. S.: The Effect of Submarine Groundwater Discharge on the Ocean, *Annu. Rev. Marine. Sci.*, 2, 59–88, doi:10.1146/annurev-marine-120308-081019, <https://doi.org/10.1146/annurev-marine-120308-081019>, 2010a.
- 5 Moore, W. S.: A reevaluation of submarine groundwater discharge along the southeastern coast of North America, *Global Biogeochemical Cycles*, 24, GB4005, doi:10.1029/2009GB003747, <https://doi.org/10.1029/2009GB003747>, 2010b.
- Moore, W. S. and Dymond, J.: Fluxes of ^{226}Ra and barium in the Pacific Ocean: The importance of boundary processes, *Earth and Planetary Science Letters*, 107, 55–68, doi:10.1016/0012-821x(91)90043-h, [https://doi.org/10.1016/0012-821x\(91\)90043-h](https://doi.org/10.1016/0012-821x(91)90043-h), 1991.
- Moore, W. S. and Shaw, T. J.: Fluxes and behavior of radium isotopes, barium, and uranium in seven Southeastern US rivers and estuaries, *Marine Chemistry*, 108, 236–254, doi:10.1016/j.marchem.2007.03.004, <https://doi.org/10.1016/j.marchem.2007.03.004>, 2008.
- 10 Moore, W. S., Astwood, H., and Lindstrom, C.: Radium isotopes in coastal waters on the Amazon shelf, *Geochimica et Cosmochimica Acta*, 59, 4285–4298, doi:10.1016/0016-7037(95)00242-r, [https://doi.org/10.1016/0016-7037\(95\)00242-r](https://doi.org/10.1016/0016-7037(95)00242-r), 1995.
- Moore, W. S., Sarmiento, J. L., and Key, R. M.: Submarine groundwater discharge revealed by ^{228}Ra distribution in the upper Atlantic Ocean, *Nature Geosci*, 1, 309–311, doi:10.1038/ngeo183, <https://doi.org/10.1038/ngeo183>, 2008.
- 15 Rama and Moore, W. S.: Using the radium quartet for evaluating groundwater input and water exchange in salt marshes, *Geochimica et Cosmochimica Acta*, 60, 4645–4652, doi:10.1016/S0016-7037(96)00289-X, [https://doi.org/10.1016/S0016-7037\(96\)00289-X](https://doi.org/10.1016/S0016-7037(96)00289-X), 1996.
- Redi, M. H.: Oceanic isopycnal mixing by coordinate rotation, *Journal of Physical Oceanography*, 12, 1154–1158, doi:10.1175/1520-0485(1982)012<1154:oimber>2.0.co;2, <http://journals.ametsoc.org/doi/pdf/10.1175/1520-0485%281982%29012%3C1154%3AOIMBCR%3E2.0.CO%3B2>, 1982.
- 20 Resplandy, L., Keeling, R. F., Stephens, B. B., Bent, J. D., Jacobson, A., Rödenbeck, C., and Khatiwala, S.: Constraints on oceanic meridional heat transport from combined measurements of oxygen and carbon, *Climate Dynamics*, 47, 3335–3357, doi:10.1007/s00382-016-3029-3, <https://doi.org/10.1007/s00382-016-3029-3>, 2016.
- Rodellas, V., Garcia-Orellana, J., Masqué, P., Feldman, M., and Weinstein, Y.: Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea, *Proceedings of the National Academy of Sciences of the United States of America*, 112, 3926–3930, doi:10.1073/pnas.1419049112, <https://doi.org/10.1073/pnas.1419049112>, 2015.
- 25 Santschi, P. H., Li, Y.-H., and Bell, J.: Natural radionuclides in the water of Narragansett Bay, *Earth and Planetary Science Letters*, 45, 201–213, doi:10.1016/0012-821x(79)90121-3, [https://doi.org/10.1016/0012-821x\(79\)90121-3](https://doi.org/10.1016/0012-821x(79)90121-3), 1979.
- Taniguchi, M., Burnett, W. C., Cable, J. E., and Turner, J. V.: Investigation of submarine groundwater discharge, *Hydrological Processes*, 16, 2115–2129, doi:10.1002/hyp.1145, <https://doi.org/10.1002/hyp.1145>, 2002.
- 30 Taniguchi, M., Ishitobi, T., Burnett, W. C., and Shimida, J.: Comprehensive evaluation of the groundwater–seawater interface and submarine groundwater discharge, in: *A new focus on groundwater-seawater interactions*, pp. 86–92, http://iahs.info/uploads/dms/13930.15-86-92-312-31_Taniguchi.pdf, 2007.
- Taniguchi, M., Ishitobi, T., and Burnett, W. C.: *From Headwaters to the Ocean: Hydrological Change and Watershed Management*, chap. Global assessment of submarine groundwater discharge, pp. 613–617, CRC Press, 2009.
- 35 Vancoppenolle, M., Fichefet, T., Goosse, H., Bouillon, S., Madec, G., and Maqueda, M. A. M.: Simulating the mass balance and salinity of Arctic and Antarctic sea ice. 1. Model description and validation, *Ocean Modelling*, 27, 33–53, doi:10.1016/j.ocemod.2008.10.005, <https://doi.org/10.1016/j.ocemod.2008.10.005>, 2009.
- Wunsch, C.: *Discrete inverse and state estimation problems*, Cambridge University Press, 2006.

Table 1. Cost functions

Cost function	Formula	Residuals
Ordinary least-squares (ols)	$C_{ols}(x) = \sum_{j=1}^p ([^{228}\text{Ra}]_i^{mod} - [^{228}\text{Ra}]_i^{obs})^2$	$\eta_{ols}(x) = [^{228}\text{Ra}]^{mod} - [^{228}\text{Ra}]^{obs}$
Logarithmic least-squares (log)	$C_{log}(x) = \sum_{j=1}^p (\log[^{228}\text{Ra}]_i^{mod} - \log[^{228}\text{Ra}]_i^{obs})^2$	$\eta_{log}(x) = \log[^{228}\text{Ra}]^{mod} - \log[^{228}\text{Ra}]^{obs}$
Proportional least-squares (prp)	$C_{prp}(x) = \sum_{j=1}^p \left(\frac{[^{228}\text{Ra}]_i^{mod} - [^{228}\text{Ra}]_i^{obs}}{[^{228}\text{Ra}]_i^{obs}} \right)^2$	$\eta_{prp}(x) = \frac{[^{228}\text{Ra}]^{mod} - [^{228}\text{Ra}]^{obs}}{[^{228}\text{Ra}]^{obs}}$

Table 2. Correlation between data and model concentration fields

Model	Linear correlation	Logarithmic correlation
Prior	0.383	0.809
ols	0.813	0.883
log	0.797	0.902
prp	0.754	0.877

Table 3. Root mean square of residuals before and after inversion

Model	Ordinary residuals rms (dpm.m ⁻³)	Logarithmic residuals rms (no unit)	Proportional residuals rms (no unit)
Prior	70.7	0.892	1.91
ols	36.6	0.703	1.10
log	37.9	0.636	1.02
prp	47.4	0.940	0.558

Table 4. Model global ²²⁸Ra fluxes (10²³ atoms yr⁻¹) with different numbers of source regions

Cost function	Case 1: 52 regions	Case 2: 38 regions	Case 3: 19 regions	Case 4: 12 regions	Case 5: 7 regions
ols	7.81 – 9.93	7.16 – 8.14	7.51 – 8.49	7.36 – 8.16	8.98 – 9.78
log	8.09 – 8.61	8.01 – 8.49	7.90 – 8.34	7.85 – 8.27	7.93 – 8.37
prp	5.15 – 5.47	4.96 – 5.28	4.70 – 4.98	4.25 – 4.49	3.75 – 4.01

Table 5. Root mean square of residuals after inversions with different numbers of source regions

Cost function	Case 1: 52 regions	Case 2: 38 regions	Case 3: 19 regions	Case 4: 12 regions	Case 5: 7 regions
ols	35.8	36.6	37.6	38.2	42.9
log	0.623	0.636	0.656	0.671	0.781
prp	0.545	0.558	0.581	0.597	0.665

Table 6. Estimates of radium-228 diffusive fluxes ($10^9 \text{ atoms m}^{-2} \text{ yr}^{-1}$) from fine-grained sediments (muds) by different methods. The studies which do not separate SGD and diffusion are in bold.

Reference	Study area	Method	Flux
Santschi et al. (1979)	Narragansett Bay	Inventory	52
Moore et al. (1995)	Amazon shelf	Inventory	110
Rama and Moore (1996)	North Inlet	Sediment profile	5
Hancock et al. (2000)	Zecks Lagoon	Benthic chamber	80
		Sediment Profile	80
Crotwell and Moore (2003)	Port Royal Sound Marsh	Krest Formula	9
Hancock et al. (2006)	Greet Barrier Reef	Modeling	26




Figure 1. Observed concentrations of ^{228}Ra in the global ocean, averaged when available on the ORCA2 cells used for inversion. Data plotted on subfigures b), c) and d) are also vertically averaged in order to take all layers into account.




Figure 2. Model ^{228}Ra source regions




Figure 3. ^{228}Ra annual flux from each source within one standard deviation, after minimization of three cost functions. Prior estimates, proportional to shelf surfaces, are shown for comparison.




Figure 4. Model surface ^{228}Ra concentration after minimization of three cost functions. Prior estimate is shown for comparison.




Figure 5. Surface ^{228}Ra logarithmic residuals after minimization of three cost functions. Prior estimate is shown for comparison.




Figure 6. (a) Residuals, (b) logarithmic residuals, (c) proportional residuals and (d) model ^{228}Ra concentration as a function of observed ^{228}Ra concentration. Lines of zero residuals are drawn in black.




Figure 7. Probability Density Functions (PDF) of ^{228}Ra (a) ordinary residuals, (b) logarithmic residuals and (c) proportional residuals after inversion, compared with a Gaussian PDF (black full line) based on their standard deviations.




Figure 8. ^{228}Ra sources by basin, divided into riverine input, diffusion from sediment and SGD. Only logarithmic least-squares results are shown. Fluxes are divided the following way: Regions 1 and 38: Southern; 2 – 4: South Atl; 5 – 16: North Atl; 17 – 18 and 26 – 27: South Pac; 19 – 25: North Pac; 28 – 34: Indian; 35 – 37: Arctic




Figure 9. map of GEOTRACES cruises (from <http://www.geotraces.org/cruises/cruise-summary>). Planned sections are in red, completed sections in yellow, International Polar Year sections in black. Names in rectangles correspond to cruises potentially bringing the largest extra information on ^{228}Ra fluxes because they are performed in places lacking measurements.