



Assimilating bio-optical glider data during a phytoplankton bloom in the southern Ross Sea

Daniel E. Kaufman¹, Marjorie A. M. Friedrichs¹, John C. P. Hemmings^{2,3}, Walker O. Smith Jr.¹

¹Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, VA, USA

5 ²Wessex Environmental Associates, Salisbury, UK

³now at Met Office, Exeter, UK

Correspondence to: Daniel E. Kaufman (dekaufman@vims.edu) and Marjorie A. M. Friedrichs (marjy@vims.edu)

Abstract. The Ross Sea is a region characterized by high primary productivity in comparison to
10 other Antarctic coastal regions, and its productivity is marked by considerable variability both
spatially (1-50 km) and temporally (days to weeks). This variability presents a challenge for
inferring phytoplankton dynamics from observations that are limited in time or space, which is
often the case due to logistical limitations of sampling. To better understand the spatiotemporal
variability of Ross Sea phytoplankton dynamics and determine how restricted sampling may
15 skew dynamical interpretations, high-resolution bio-optical glider measurements were
assimilated into a one-dimensional biogeochemical model adapted for the Ross Sea. Assimilation
of glider data using the micro-genetic and local search algorithms in the Marine Model
Optimization Testbed improves model-data fit by ~50%, generating rates of integrated primary
production of $104 \text{ g C m}^2 \text{ y}^{-1}$ and export at 200 m of $27 \text{ g C m}^2 \text{ y}^{-1}$. Assimilating glider data from
20 three different latitudinal bands and three different longitudinal bands results in minimal changes
to the simulations, improves model-data fit with respect to unassimilated data by ~35%, and
confirms that analyzing these glider observations as a time series via a one-dimensional model is
reasonable on these scales. Whereas assimilating the full glider data set produces well-
constrained simulations, assimilating subsampled glider data at a frequency consistent with
25 cruise-based sampling, results in a wide range of primary production and export estimates. These
estimates depend strongly on timing of the assimilated observations, due to the presence of high
mesoscale variability in this region. Assimilating surface glider data subsampled at a frequency
consistent with available satellite-derived data results in 40% lower carbon export, primarily
resulting from optimized rates generating more slowly sinking diatoms. This analysis highlights
30 the need for strategic consideration of impacts of data frequency, duration, and coverage when



combining observations with biogeochemical modeling in regions with strong mesoscale variability.



1 Introduction

35 Phytoplankton blooms in the Ross Sea are responsible for some of the highest rates of
productivity in the Southern Ocean (Arrigo et al., 2008), and yet the phytoplankton assemblage
exhibits considerable spatiotemporal variability (DiTullio and Smith, 1996; Hales and Takahasi,
2004; Smith et al., 2010). This heterogeneity, and the spatial/temporal limitations of observations
due to logistical challenges of sampling, may affect the inferred phytoplankton dynamics and
40 produce biases in productivity or export estimates. The magnitude of the underlying ecosystem
variability that contributes to these potential biases is not well understood, nor is it well known
how the use of different observational platforms in the Ross Sea might affect the inferred
dynamics. Acquiring data with an appropriate resolution is important for assessing
phytoplankton variability in the Ross Sea (Hales and Takahashi, 2004).

45 Over the past several decades, biogeochemistry in the Ross Sea has been observed by
ship and satellite, providing data at different temporal and spatial resolutions. Since Ross Sea
phytoplankton became a focus of scientific research in the late 1970s, water column
measurements have primarily come from research vessels (e.g., El-Sayed et al., 1978; Smith and
Nelson, 1985; Vaillancourt et al., 2003). Typically, sampling stations are separated by tens of
50 kilometers (Hales and Takahashi, 2004), and although vessels may return to resample a station,
they typically do not return more than once or twice in a single year. During the 1990s, the use of
remote sensing was expanded to look more closely at the Ross Sea bloom (Arrigo and McClain,
1994), and satellite retrievals have continued to provide valuable insights into characteristics of
the phytoplankton assemblage (Arrigo et al., 1998; Arrigo and van Dijken, 2004; Peloquin and
55 Smith, 2007; Schine et al., 2015). Satellite observations offer a synoptic view of spatial regions
at frequencies that are within the time scale of biological changes (e.g. growth); however, the
presence of sea ice and clouds often obscures remote-sensing measurements in the Ross Sea
(Arrigo et al., 1998).

60 At the mesoscale (days-weeks, 1-10 km), gliders are a relatively new and effective means
to characterize phytoplankton variability, and the development of ice-avoidance algorithms has
enabled the use of gliders in the Ross Sea for these purposes. For example, a glider equipped
with bio-optical sensors was directed along a section near 76° 40' S in austral summer 2010 -
2011 and provided valuable estimates of biomass variability on short time scales (Kaufman et al.,
2014). Estimates of the POC:Chl ratio from the glider optical sensors suggested a transition from



65 a *Phaeocystis antarctica* to a diatom-dominated assemblage over several days (Kaufman et al.,
2014; Thomalla et al., 2017). Moreover, Jones and Smith (2017) used glider observations from
austral summer 2012-2013 to distinguish three phases of the Ross Sea bloom and identified high
frequency (hours) associations between wind-driven mixing and biomass. A perennial challenge
when using glider data (as well as ship-based data), however, is separating the effects of time and
70 space (Kaufman et al., 2014; Little, 2016).

Numerical models are another approach for examining phytoplankton variability in the
remote Ross Sea, providing an effective means for coordinating knowledge and understanding
the underlying system complexities (Leonelli, 2009; Vallverdú, 2014). Furthermore, numerical
simulations offer the ability for experimental manipulations that would be impractical or
75 impossible in the real system. Such manipulations were implemented in the scenario experiments
described by Kaufman et al. (2017a) to investigate how projected climate changes might alter the
dynamics of the phytoplankton assemblage. These experiments showed that earlier availability of
low light resulting from sea ice reduction was the primary driver of projected increases in
production and export and composition change over the next century.

80 Data assimilation, which refers to methodologies that systematically combine a
mathematical model with observations, is often used in biogeochemical applications (Hofmann
and Friedrichs, 2001, 2002) to improve estimates of model parameters that are frequently poorly
known (Lawson et al., 1995, 1996; Matear, 1995; Fennel et al., 2001; Friedrichs, 2002; Schartau
and Oschlies, 2003; Hemmings et al., 2004; Doron et al., 2013; Xiao and Friedrichs, 2014a,b;
85 Melbourne-Thomas et al., 2015; Gharamti et al., 2017, Schartau et al., 2017). This entails a
smoothing or optimization procedure, in which elements of the model are adjusted to minimize
differences between the model output and the observations. Typically, an aggregate measure of
the differences between observations and model output is provided by calculation of a cost
function, defined as the model-data misfit, and an optimization algorithm searches for model
90 parameters that minimize the value of this cost function.

In this study, data assimilation is used to obtain an optimal representation of Ross Sea
lower trophic levels. Specifically, observations from an autonomous glider are assimilated into a
biogeochemical model of the Ross Sea (Kaufman et al., 2017a) to better understand the spatial
and temporal variability of phytoplankton in this region. Assimilation experiments also examine



95 how the space and time characteristics of observational sampling frequency impacts the ability of
observations to produce optimal system representations.

2 Methods

2.1 One-dimensional biogeochemical model

Numerical experiments were conducted with the Model of Ecosystem Dynamics, nutrient
100 Utilisation, Sequestration and Acidification for the Ross Sea (MEDUSA-RS; Kaufman et al.,
2017a), a regionally adapted version of MEDUSA-1.0 (Yool et al., 2011). Three phytoplankton
groups are represented in the MEDUSA-RS model: colonial *P. antarctica*, solitary *P. antarctica*,
and diatoms. Phytoplankton growth in the model is temperature dependent as well as limited by
light and nutrient availability. Colonial *P. antarctica*, diatoms, and detritus all sink at distinct
105 rates. The model handles the sinking of large detrital particles implicitly as a fast-sinking group
to avoid issues related to the scale of the model time step and to avoid the need for an additional
tracer. A ballast scheme is used to allow inorganic materials to “protect” a variable fraction of
the sinking organic material from degradation. A full description of the model and its set-up
within the Marine Model Optimization Testbed (MarMOT; Hemmings and Challenor, 2012), as
110 well as the physical forcings derived from glider observations, are documented in Kaufman et al.
(2017a,b).

2.2 Data for assimilation

In situ observations used for the assimilation experiments came from an iRobot Seaglider
equipped with a Wet Labs ECO Puck sensor and are available in the BCO-DMO data repository
115 (<http://www.bco-dmo.org/dataset/568868>). Glider dives from 22 November 2012 to 01 February
2013 covered a horizontal area spanning 76.83 - 77.44 °S and 168.9 - 171.97 °E (Fig. 1). Data
spanning the upper 200 m of the water column were binned into hourly, 5-m vertical bins.
Concentrations of chlorophyll (Chl) and particulate organic carbon (POC) were derived,
respectively, from fluorescence and optical backscatter counts measured by the sensor and
120 converted using regression equations (Kaufman et al., 2017a). These bio-optical quantities were
used for calculating model-data misfits during assimilation.

2.3 Cost function

The ‘cost function’ (J), defined as a measure of misfit between a particular model
simulation and observational data, is computed as a weighted average of the squared differences
125 between simulated and observed values:



$$J = \frac{1}{N} \sum_{i=1}^N \left(\frac{1}{\sigma_{chl}^2} (x_{i,chl} - y_{i,chl})^2 + \frac{1}{\sigma_{poc}^2} (x_{i,poc} - y_{i,poc})^2 \right)$$

where N is the number of observation points, σ is the standard deviation of each variable, x_i is the simulated value of either chlorophyll or POC at the i th observation point and y_i is its observed value. Model parameters were optimized in MarMOT by finding the minimum of this cost function through a combination of the micro-genetic algorithm and Powell's non-gradient direction set algorithm, as described below.

2.4 Implementation of micro-genetic algorithm and direction set algorithm

Genetic algorithms, including the micro-genetic algorithm [μ GA], are a subtype of computational methods known as evolutionary algorithms, so-called because of their inspiration from, and metaphoric relationship to, biological evolution. Described using this metaphor, a genetic algorithm procedure modifies a population of candidate solutions over successive generations by variation and selection processes to converge on a single solution or solution area. GAs have several advantages for optimization, including their intrinsic parallelism, suitability for systems with multiple local minima, and their generalizability (Bajpai and Kumar, 2010; Ward et al., 2010). The μ GA uses three steps to transition from one generation to the next, described following the biological metaphor as: selection, crossover, and resampling (Krishnakumar, 1990; Črepinšek et al., 2013). An advantage of the μ GA is its reduced risk of premature convergence, resulting from reinitializing after each convergence, and generating new random populations while maintaining the best fit individual from the previous set (Schmitt, 2001).

In this μ GA implementation, optimizations begin with a population of five individual parameter sets randomly generated for the first μ GA generation. An evaluation of the cost function for each model solution indicates the 'fitness' of each individual. A binary tournament procedure is then followed to select parents from this population for the next generation. The most-fit individuals (i.e., those with the lowest cost function values) are paired with one another and undergo recombination of the bits representing parameter values. After each generation, the proportion of bits differing from those of the fittest individual is calculated to determine whether the population can be deemed converged (though this does not necessarily indicate closeness in parameter space). After the threshold for convergence has been achieved, the population is reinitialized to random individuals, although the fittest individual is maintained. The μ GA is



terminated upon the first convergence occurring after a minimum number of generations has
155 been reached.

Once convergence has been achieved after a minimum number of μ GA generations,
Powell's non-gradient direction set algorithm performs a local search using the μ GA solutions as
starting points. The direction set method performs sequential minimizations in iterative
directions, updating the search direction after each iteration (Powell, 1964; Press et al., 1992).
160 Although the μ GA is well suited for global search problems partly because of its stochasticity,
Powell's direction set algorithm is well suited to searching for a local optimum. Brent's method,
which combines root-bracketing with secant and inverse quadratic interpolation (Brent, 1973), is
used to numerically locate cost minima between neighboring function evaluations along each
direction identified by the Powell algorithm. The direction set algorithm stops when a cost
165 function minimum is located or when a maximum number of iterations is reached. The optimized
parameter values are those that generated the cost function minimum.

2.5 Selection of parameters to be optimized

Even though it is tempting to try to optimize all parameters in a model, uncertainty in the
parameter estimates increases as the number of parameters optimized increases (Friedrichs et al.,
170 2007; Ward et al., 2010). Although the optimization of more parameters generally lowers the
assimilated cost, the increasing potential for equifinality with more parameters means the
optimization may find equivalent low-cost solutions with substantially different parameter
values. Therefore, before assimilating observations and optimizing parameters, a subset of "free"
or "optimizable" model parameters must be chosen. In this study, the parameters to be optimized
175 are selected through a three-step process: defining a range of permitted values for every
parameter (Sect. 2.5.1), identifying the parameters to which model outputs are most sensitive
(Sect. 2.5.2), and evaluating how many of these sensitive parameters can be reasonably
optimized when assimilating the available data (Sect. 2.5.3). Initial values for each parameter,
prior to the assimilation, were set to values identified in Kaufman et al. (2017a).

180 2.5.1 Parameter ranges

Upper and lower bounds of the allowable range for each free parameter were defined
loosely following Hemmings et al. (2015). Bounds were set to be geometrically symmetric
(factor of four for rates; factor of five for half-saturation concentrations) around the initial
values. For fractional parameter values, limits were set to ± 0.25 their initial values, although



185 not allowed to exceed 0.05 or 0.95. Ranges for parameters not expressed as fractions were log-
transformed for sampling purposes.

2.5.2 Sensitivity Analysis

Parameters to which model outputs are highly sensitive are important and useful to
optimize. In contrast, it is futile to optimize parameters to which model outputs of interest are not
190 sensitive; no amount of varying these parameters will result in improved model performance.
Therefore, the first criterion used to designate a parameter as optimizable was the sensitivity of
model outputs to the values of that parameter. Model sensitivities were evaluated for assimilated
variables (Chl and POC) and carbon fluxes of interest (primary production (PP) and carbon
export at 200 m). To quantify the sensitivities of these outputs to each of the 80 parameters in the
195 model, a series of runs were conducted following the approach of Hemmings et al. (2015). Each
run used a unique sample of parameter values drawn from within the specified parameter ranges
(Section 2.5.1) using a Latin hypercube. This approach provides more even coverage of the
parameter space than Monte Carlo sampling methods that can result in clustered values and
unsampled regions. One thousand values were drawn from sequential intervals throughout the
200 range for each parameter. Using this technique, unique parameter sets were constructed such that
over the course of all runs, the full range of values for each parameter was represented.

The model was run 1000 times, each time using one of the unique parameter sets
resulting from Latin hypercube sampling of the full parameter space. Sensitivity was quantified
by evaluating the amount of variance in the output diagnostics explained by each parameter (i.e.,
205 by computing the coefficient of determination (r^2) between each parameter and each of the four
output variables of interest; Fig. 2). All four model outputs (Chl, POC, PP, and export) were
most sensitive ($r^2 \geq 0.01$) to attenuation of blue-green light by phytoplankton pigments, diatom
maximum growth rate, and C:Chl ratio for solitary *P. antarctica*. Three additional parameters
(maximum growth rate of *P. antarctica* colonies, maximum growth rate of solitary *P. antarctica*,
210 and microzooplankton maximum grazing rate) exhibited $r^2 \geq 0.01$ for both chlorophyll and POC.
The 21 parameters with $r^2 \geq 0.01$ (Fig. 2) were selected for further evaluation (Sect. 2.5.3).

2.5.3 Using twin experiments to select optimizable subset

After selecting the 21 potentially optimizable parameters, Numerical Twin Experiments
(NTEs) were conducted to evaluate the extent to which known values of sensitive parameters
215 could be recovered given the data available for assimilation. The implementation of NTEs



involves four primary steps (Hofmann and Friedrichs, 2001). First, the chosen model is run forward in time to create a simulation using a known, “true” parameter set. Second, output from this simulation is sub-sampled to create a so-called “synthetic” data set. Third, the synthetic dataset is then assimilated to optimize model parameters. Fourth, the optimized parameter set is compared to the true parameter set. The assimilation is successful if the optimized values recover the true parameters used to generate the assimilated synthetic data.

The procedure followed here for determining the subset of optimizable parameters is similar to that used by Friedrichs et al. (2007). First, a reference simulation was generated using the initial parameter set, and chlorophyll and POC estimates from this reference simulation were subsampled to generate a synthetic data set. Starting with a parameter space defined by the set of 21 parameters deemed sensitive in the Latin hypercube tests (Fig. 2), a series of sequential NTEs was then performed with a progressively smaller number of optimized parameters: after each NTE, the optimized parameter that was most different from its ‘true’ value was removed from the optimizable parameter set. Thus, after each NTE the number of optimized parameters was reduced by one. The series of NTEs was evaluated to identify the largest parameter set for which the original parameter values were recoverable and the cost function remained essentially zero. From this analysis (Fig. 3), it was determined that optimizing eight parameters would be ideal (Table 1), because values of these eight parameters were recovered much better than larger parameter sets and model-data misfit (cost) remained low.

2.6 Assimilation experiments

The μ GA optimization procedure was used to assimilate glider data in two sets of experiments that explored aspects of spatiotemporal variability and data availability. Estimates of depth- and time-integrated PP and time-integrated carbon export at 200 m were computed from the full model simulation in each experiment.

2.6.1 Experiment #1

The first set of experiments examined the differences in model simulations resulting from assimilating Chl and POC data from different spatial regions. In Experiment #1a, glider observations were assimilated from the upper 50 m of the full temporal and spatial domain, referred to hereafter as the “Full Assimilation” case (Table 2). (Comparisons showed only minor differences between assimilating data from the upper 50 m vs. the full upper 200 m). Observations from different spatial areas of the glider track were also assimilated. Observations



from the glider track were divided into three latitudinal bands (Northern, Central, and Southern bands) as well as into three longitudinal areas constituting Eastern, Central, and Western bands. Glider data from each of these three latitudinal and longitudinal bands were assimilated in
250 Experiment #1b and #1c, respectively (Table 2, Fig. 4), resulting in three cost functions for each of these experiments.

2.6.2 Experiment #2

The second set of experiments investigated the assimilation of data at different resolutions mimicking different data sources. In Experiment #2a, glider data were subsampled at
255 ~12-hour intervals (Table 2). The subsampling was repeated 12 times, with each iteration offset from the previous by +1, 2, 3... 11 hours, to generate a series of 12 glider observation sets. The assimilation of these 12 time series yields the “Glider Assimilation” case. In Experiment #2b, glider data were subsampled at a reduced temporal resolution similar to cruise sampling (Table 2). Sampling during cruise missions often takes place for a few days in one location before
260 moving elsewhere, and the ship sometimes returns to the first location after a number of weeks. To roughly mimic this sampling pattern, daily vertical profiles (again down to 50 m) were assimilated for three days in a row, starting from the first day of available glider data (22 Nov), and then three days of data were assimilated two weeks later. Shifting this pattern forward one week at a time generated a series of eight cruise-based observation sets for assimilation in this
265 “Cruise Based Assimilation” case. In Experiment #2c, glider data were assimilated only from the upper 5 m surface layer to produce a data set resembling satellite-derived data. These data were then subsampled at two-week intervals, to represent typical data return from remote sensing observations of ocean color in the Ross Sea, where the availability of satellite image retrieval is frequently limited by excessive, though variable, cloud cover (Arrigo and van Dijken, 2004). The
270 two-week subsampling pattern covered the entire period of glider data (22 Nov - 1 Feb), and was sequentially shifted forward one day at a time to generate a series of 14 satellite-based observation sets for assimilation in this “Satellite-Based Assimilation” case (Table 2).

2.7 Predictive Cost Assessment

In addition to the assimilative cost (J_A) calculated during the optimization procedure
275 using assimilated data, a predictive cost (J_P) was calculated to assess model-data misfit computed using the unassimilated data in each experiment. Because predictive costs represent model-data misfit from unassimilated data only (Friedrichs et al., 2006; Ward et al., 2010), it is



an objective measure of the skill of an optimized model in reproducing observations at different points in time or space (Gregg et al., 2009). In this case, the aim of these experiments is to assess the skill of each optimized simulation regardless of which subset of the available data is assimilated. By computing the mean and median predictive cost for each experiment (other than the Full Assimilation case), the skill of the resulting simulations can be compared directly with one another.

3 Results

3.1 Experiment #1

Assimilation of the glider data over the full temporal and spatial domain (Full Assimilation case) improves the model-data fit of both Chl and POC (Fig. 5a,b) and reduces the cost by nearly half (47%) compared to the a priori simulation without assimilation (Table 3). Average Chl and POC concentrations in the upper 50 m are both slightly lower (8% and 12%, respectively) in the optimized simulation. The contribution of each phytoplankton group to total chlorophyll remains similar to the No Assimilation case (Fig. 6a,c), but colonial *P. antarctica* carbon is lower and diatom carbon is higher in December and early January (Fig. 6b,d). Compared to the No Assimilation case, PP is only slightly lower (7%), whereas export flux is nearly 50% higher (Table 3; Fig. 7). Compared to their initial values, colonial *P. antarctica* parameters change the most as a result of the optimization, with reductions between 40-70% for the colonial *P. antarctica* maximum growth rate, maximum sinking rate and C:Chl ratio (Table 4). In contrast, the diatom maximum growth rate and C:Chl ratio increased (~10% and 20% respectively).

Chlorophyll and POC time-series exhibit only minor differences between latitudinal band experiments when data from the northern, central, and southern sections are assimilated independently (Fig. 5c,d) or when data from the eastern, central western sections are assimilated (Fig. 5e,f). Specifically, the optimal simulations for Chl and POC exhibit similar seasonal cycles across the three latitudinal and longitudinal bands, with only slightly higher Chl and POC concentrations when assimilating data from the southern band (Fig. 5c,d) and higher Chl from the western band (Fig. 5e,f). Mean costs are much lower for the latitudinal and longitudinal experiments than for the No Assimilation case, and only slightly higher than the Full Assimilation case (Table 3). This indicates that data sampled from within only one spatial band improved the match between modeled and observed variables in the unassimilated areas as well.



310 Average estimates of PP and export in both the latitudinal and longitudinal experiments are only slightly less (< 5%) than the Full Assimilation estimate (Fig. 7, Table 3).

3.2 Experiment #2

Assimilation of data subsampled at a frequency one-twelfth that of the original glider data (Expt. 2a) results in twelve model simulations, all of which are similar to the Full Assimilation case, with Chl and POC time series closely following the observed seasonal pattern (Fig. 8a,b).
315 Mean assimilative and predictive costs in the Glider Assimilation case are close to the cost of the Full Assimilation case (Table 3). Mean PP and export estimates are also close to estimates from the Full Assimilation case. The mean optimal parameter values obtained from the Glider Assimilation case are generally within one standard deviation of the optimal values from the Full Assimilation case (Table 4).

320 Assimilation of data subsampled with a frequency typical of cruise observations (Expt. 2b) results in a wide range of solutions, with several Chl and POC time series exhibiting markedly different peak bloom timings (Fig. 8c,d). Two of the solutions yield substantially higher concentrations of POC in November, and Chl peaks range from mid-November to early January. The mean predictive cost from this experiment (1.24) is roughly three times the
325 assimilative cost for the Full Assimilation case (0.41) and three times the predictive cost for the Glider Assimilation case (0.43; Table 3). The PP estimates from the Cruise-based Assimilation case span a broad range (92 to 156 g C m⁻² y⁻¹) around the Full Assimilation estimate but are generally higher (Fig. 7a). This experiment similarly yields a very large range of export estimates (11 to 33 g C m⁻² y⁻¹) encompassing the results from Experiment #1 (Fig. 7b). Optimal parameter
330 values obtained from the Cruise-based Assimilation case are generally less well constrained (higher standard deviations) than the Glider Assimilation case (Table 4).

Assimilation of data subsampled as satellite-based observations from the surface layer (Expt. 2c) results in Chl and POC concentrations generally higher than the Full Assimilation case (Fig. 8e,f). The predictive costs are similar on average to those of the Cruise-based Assimilation
335 experiment; however, there is less variation (Table 3). The median integrated production is higher (9%) than the Full Assimilation estimate and the Cruised-based Assimilation estimate (Fig. 7a; Table 3); however, the range of PP estimates for this Satellite-based Assimilation case is smaller than those for the Cruise-based Assimilation case (Fig. 7a). Most notably, despite generally higher PP and higher POC concentrations, carbon export from the Satellite-based



340 Assimilation case is substantially lower (41%) than the Full Assimilation estimate (Fig. 7b;
Table 3). In fact, export estimates from individual runs in this experiment are all lower (-19% to
-56%) than the Full Assimilation estimate (Fig. 7b). Again, the range of export estimates is
smaller for the Satellite-based Assimilation than for the Cruise-based Assimilation. When
assimilating data at a resolution similar to that of satellite-based observations, mean optimal
345 parameter values were similar to those obtained in the Glider Assimilation and Cruise-based
Assimilation cases, with the exception of the fast detritus sinking fraction for diatoms, which was
significantly lower in the Satellite-based Assimilation case (0.62 ± 0.14) than in the other
experiments (Glider Based Assimilation Case: 0.86 ± 0.05). In contrast to this sinking parameter
for mortality from diatoms, the mean maximum sinking rate of colonial *P. antarctica* in the
350 Satellite-based case was not significantly different than its value in either the Full Assimilation
or Cruise-based cases (Table 4). Standard deviations of optimal parameters for the Satellite-
based Assimilation case were generally similar to or lower than those for the Cruise-based
Assimilation case, except for the C:Chl ratio for diatoms, which produced a very high optimal
value and was particularly poorly constrained (375 ± 187 gC gChl⁻¹; Table 4).

355 4 Discussion

4.1 Ross Sea simulation resulting from assimilation of glider data

Data assimilation is a valuable tool for efficiently utilizing limited observational data in
remote regions like the Ross Sea. In this study, glider data consisting of both fluorescence-
derived chlorophyll and backscatter-derived POC were assimilated into a one-dimensional
360 marine biogeochemical model developed for the Ross Sea. Eight ecosystem parameters,
including phytoplankton rates and C:Chl ratios, were optimized resulting in a simulation with a
50% reduced model-data misfit. This optimal simulation yielded lower *P. antarctica* carbon
concentrations and higher diatom carbon concentrations, resulting in higher carbon export
compared to those generated by the initial hand-tuned simulation (Kaufman et al., 2017a),
365 despite slightly lower estimates of overall annual primary production. Changes in chlorophyll
concentrations of diatoms and *P. antarctica* were minor. This optimal simulation was obtained
largely via changes in the C:Chl ratios: the colonial *P. antarctica* ratio of C:Chl was lower and
the diatom C:Chl was higher than in the original simulation. Although modified from their initial
values, the relative differences between these optimized C:Chl ratios for *P. antarctica* and
370 diatoms are consistent with shipboard measurements of C:Chl ratios, which found higher C:Chl



in diatom-dominated waters compared to *P. antarctica*-dominated waters: ~200 vs. 90 g C g Chl⁻¹ (DiTullio and Smith, 1996), and ~50-100 vs. 20-50 g C g Chl⁻¹ (Mathot et al., 2000). Although the authors are not aware of any specific estimates in the literature for the fraction of diatom mortality that becomes fast-sinking detritus, other optimal rate parameters are consistent with those previously reported in the literature. For example, the optimized growth rates (0.29 - 0.4 m d⁻¹) are similar to measured values in the Ross Sea (Smith and Gordon, 1997; Smith et al., 1999; Mosby and Smith, 2015), and the optimized sinking rate of *P. antarctica* colonies (14 m d⁻¹) is similar to previous estimates (Asper and Smith, 1999; Asper and Smith, 2003; Smith et al., 2011).

4.2 Spatial variation within the glider track

Phytoplankton in the Ross Sea exhibit both spatial and temporal variability. Cruise transects across the continental shelf show a marked spatial variability in both the east-west and north-south direction over short periods of time (Smith et al., 2013). Within the Ross Sea Polynya, ship-based observations show biochemical gradients that suggest patchiness of phytoplankton dynamics on the mesoscale (Hales and Takahashi, 2004; Smith et al., 2017). Nutrient pools have been found to exhibit gradients from both north to south and east to west (DiTullio and Smith, 1996; Sedwick et al., 2011; Smith et al., 2013; Marsay et al., 2014), and phytoplankton assemblage composition is not necessarily uniform across longitudes (DiTullio and Smith, 1996; Garrison et al., 2003; Smith et al., 2013). In addition, cold and fresh eddies have been observed along the ice shelf edge potentially reshaping the phytoplankton assemblage on short time (<10 days) and space (<20 km) scales (Li et al., 2017).

When analyzing glider data in regions characterized by high mesoscale variability, it is often not apparent whether observed patterns represent spatial or temporal variability. As Rudnick (2016) discusses, “Because gliders can occupy lines, their data can be viewed as traditional sections, such as those measured from a ship. However, because high-frequency variability is projected onto a spatial structure, it is sometimes more convenient to think of the data as a time series from a mooring.” This ambiguity led Kaufman et al. (2014) to concede “both spatial and temporal gradients may have played a role in the observed variability” when analyzing physical-biological relationships from glider data in the southern Ross Sea.

Although both temporal and spatial gradients may be present, observations can be presented as either primarily spatial or temporal patterns with simple tests guiding the decision.



For example, a comparison of means and standard deviations across spatial sections and time periods was previously used to identify time as the dominant dimension of variability in the 2012-2013 glider observations (Jones and Smith, 2017). In this study, a similar conclusion was
405 reached, using a very different methodology. The assimilation of glider data from nine different sub-areas of the study region (separated latitudinally or longitudinally by ~20 km) indicated that the seasonal cycle is similar in phase throughout the region of the glider track. The assimilation of glider data from each of the nine regions yielded similar estimates of POC and Chl, generally within the variance of the glider observations (gray areas of Fig. 5c-f), and similar estimates of
410 temporally averaged primary productivity and export. This further supports the approach of using the glider data as a time series and suggests that temporal patterns at this scale play a greater role than spatial patterns in structuring variability of the phytoplankton assemblage. Moreover, the similarity between predictive and assimilative costs when assimilating the latitudinal and longitudinal bands of data suggests that the parameters are not being over-fit for
415 these experiments. Thus, temporally resolved observations in any of these regions might be expected to provide similar constraints on modeled temporal patterns of the phytoplankton.

4.3 Differences between assimilating glider, satellite-derived, and cruise-based data

Results from experiments that assimilated data at different spatial and temporal resolutions suggest that assimilating only surface observations, as are typically available from
420 remote-sensing platforms, underestimates carbon export and more weakly constrains estimates of productivity relative to assimilation of depth-resolved glider data. The lower estimates of carbon export occurred because the optimal diatom fraction for fast-sinking detritus obtained via the assimilation of surface-only data (0.62 ± 0.14) was significantly lower than that obtained via the assimilation of data throughout the upper 50 m (Expt. 2a: 0.86 ± 0.05 ; Expt. 2b: 0.86 ± 0.11).
425 Experimental results also indicate that the assimilation of satellite-derived data provides a weaker constraint on productivity estimates, as seen by the larger range of estimates (114 ± 11 $\text{gC m}^{-2} \text{y}^{-1}$), as compared to the assimilation of glider data (104 ± 2 $\text{gC m}^{-2} \text{y}^{-1}$). Although not statistically significant, the higher productivity estimates generated by the assimilation of satellite-derived data is consistent with those of Gregg (2008), who found that assimilation of
430 satellite-based chlorophyll estimates into a three-dimensional global biogeochemical model overestimated primary production. In contrast, results from assimilating satellite-derived



chlorophyll concentrations into a one-dimensional model in the equatorial Pacific produced underestimates of primary productivity compared to in situ observations (Friedrichs, 2002).

Although both chlorophyll and POC were assimilated in the present study, chlorophyll
435 alone has been the dominant satellite data product used in biogeochemical assimilation, although
other data types are available and can impact the assimilation results. For instance, a study
investigating the assimilation of different types of satellite-derived data, including POC and size-
fractionated chlorophyll, found that assimilation of satellite-derived POC estimates worsened the
model estimates of chlorophyll, whereas the assimilation of chlorophyll did not substantially
440 impact the POC estimates (Xiao and Friedrichs, 2014b). Additionally, satellite-based sampling
bias could be reduced by concurrently assimilating export flux data derived from sediment trap
measurements (Friedrichs et al., 2007), or by assimilating satellite measurements such as remote-
sensing reflectance directly (Jones et al., 2016). It is also worth noting that when assimilating
actual satellite data, the biases suggested by this study resulting from assimilation of only surface
445 data would be compounded with biases inherent in the satellite retrieval algorithms (Saba et al.,
2011; Stukel et al., 2015).

Assimilating cruise-based data in the highly variable Ross Sea may also yield potentially
large errors in primary production, as well as in carbon export estimates, depending on which
specific days are sampled. Estimates of bloom timing from the assimilation of cruise-based
450 observations may also vary substantially (Fig. 8c,d). This echoes the results of a series of
reduced resolution data interpolations, from which Hales and Takahashi (2004) reported that
cruise-based observations in the Ross Sea were likely able to capture average conditions well,
but miss some mesoscale phenomena. Likewise, a subsampling analysis of physical-biological
correlations from 2010 Ross Sea glider data demonstrated the possibility of lower resolution data
455 obscuring or biasing biogeochemical interpretations (Kaufman et al., 2014). The results provided
by the data assimilative study described here can be used to help guide decisions of when and
how long to sample certain locations in the Ross Sea; this is especially important given the
limitations of ship-based sampling in such a remote region (Smith et al., 2014). In fact, the use of
data collection from other sampling platforms may decrease the pressure to conduct repeated
460 transects by ship, and allow limited vessel-time to be used for more thorough process-based
investigations uniquely-suited for research vessels.



5 Summary and Conclusions

A series of experiments investigating spatiotemporal variability of the phytoplankton assemblage and potential effects of assimilating data from different observation platforms highlighted the benefits and challenges of combining data and biogeochemical models in the Ross Sea. The assimilation of glider data reduced model-data misfit by 50%, and resulted in reduced depth-integrated primary production and higher carbon export at 200 m. Additional experiments for different spatial regions reduced predictive costs with respect to unassimilated data by ~35%, suggested that the model parameters were well constrained, and implied that using glider data as time series in these local studies is a reasonable approach. This may further suggest the value of using moorings or buoys, or even deploying gliders in a “virtual mooring” mode. However, the effects of mesoscale variability were apparent when assimilating data at a frequency characteristic of cruise-based sampling, which resulted in a wide range of primary production and export estimates depending on the sampling times. Results of assimilating data characteristic of satellite-based sampling suggest that assimilating satellite-derived data will result in underestimated carbon export. These findings can be used to help avoid potential sources of error when using ship-based or satellite-based observations alongside the development, calibration, or running of Ross Sea biogeochemical models. The combination of high-resolution glider data and modeling in this study underscores the importance of considering how the timing at which observations are collected affect the subsequent interpretations, and it is believed that further comparison of data from additional autonomous gliders throughout the Ross Sea and between years will provide greater clarity of the spatiotemporal variability on multiple scales.



485 *Data Availability.* Data from the autonomous glider are available from the BCO-DMO data repository (<http://www.bco-dmo.org/dataset/568868>), and other data to support this article will be made available at W&M Publish (<http://doi.org/xxxx>) and upon request from the authors (dekaufman@vims.edu, marjy@vims.edu).

490 *Competing Interests.* The authors declare that they have no conflict of interest.

495 *Acknowledgements.* This material is based upon work supported by the U.S. National Science Foundation's Office of Polar Programs (NSF-ANT-0838980). The authors thank Drs. Elizabeth A. Canuel, Eileen E. Hofmann, and Elizabeth H. Shadwick for constructive comments. Additional thanks go to Michael S. Dinniman for helping with model forcings. This work was performed (in part) using computational facilities at the College of William and Mary which were provided by contributions from the National Science Foundation, the Commonwealth of Virginia Equipment Trust Fund and the Office of Naval Research. This paper is contribution XXXX of the Virginia Institute of Marine Science, College of William and Mary.

500



Literature Cited

- 505 Arrigo, K. R., and McClain, C. R.: Spring phytoplankton production in the western Ross Sea, *Science*, 266(5183), 261–263, doi:10.1126/science.266.5183.261, 1994.
- Arrigo, K. R., Robinson, D. H., Worthen, D. L., Schieber, B., and Lizotte, M. P.: Bio-optical properties of the southwestern Ross Sea, *J. Geophys. Res. Ocean.*, 103(C10), 21683–21695, doi:10.1029/98JC02157, 1998.
- 510 Arrigo, K. R., and van Dijken, G. L.: Annual changes in sea-ice, chlorophyll a, and primary production in the Ross Sea, Antarctica, *Deep Sea Res. II*, 51(1–3), 117–138, doi:10.1016/j.dsr2.2003.04.003, 2004.
- Arrigo, K. R., van Dijken, G. L., and Bushinsky, S.: Primary production in the Southern Ocean, 1997–2006, *J. Geophys. Res.*, 113(C08004), 1–27, doi:10.1029/2007JC004551, 2008.
- 515 Asper, V. L., and Smith, W. O.: Particle fluxes during austral spring and summer in the southern Ross Sea, Antarctica, *J. Geophys. Res.*, 104(C3), 5345–5359, doi:10.1029/1998JC900067, 1999.
- Asper, V. L., and Smith, W. O.: Abundance, distribution and sinking rates of aggregates in the Ross Sea, Antarctica, *Deep. Res. Part I Oceanogr. Res. Pap.*, 50(1), 131–150, doi:10.1016/S0967-0637(02)00146-2, 2003.
- 520 Bajpai, P., and Kumar, M.: Genetic algorithm—an approach to solve global optimization problems, *Indian J. Comput. Sci. Eng.*, 1(3), 199–206, 2010.
- Brent, R. P.: “An algorithm with guaranteed convergence for finding a minimum of a function of one variable.” *Algorithms for Minimization Without Derivatives*, pp. 61–80, Prentice-Hall, Inc, 1973.
- 525 Črepinšek, M., Liu, S.-H. and Mernik, M.: Exploration and Exploitation in Evolutionary Algorithms: A Survey, *ACM Comput. Surv.*, 45(3), 1–33, doi:10.1145/2480741.2480752, 2013.
- DiTullio, G. R., and Smith, W. O.: Spatial patterns in phytoplankton biomass and pigment distributions in the Ross Sea, *J. Geophys. Res.*, 101(C8), 18467–18477. doi:
530 10.1029/96JC00034, 1996.
- Doron, M., Brasseur, P., Brankart, J.-M., Losa, S. N., and Melet, A.: Stochastic estimation of biogeochemical parameters from Globcolour ocean colour satellite data in a North Atlantic



- 3D ocean coupled physical–biogeochemical model, *J. Mar. Syst.*, 117–118, 81–95, doi:10.1016/j.jmarsys.2013.02.007, 2013.
- 535 El-Sayed, S. Z., Biggs, D. C., Stockwell, D., Warner, R., and Meyer, M.: Biogeography and metabolism of phytoplankton and zooplankton in the Ross Sea, Antarctica, *Antarct. J. U.S.*, 131–133, 1978.
- Fennel, K., Losch, M., Schröter, J., and Wenzel, M.: Testing a marine ecosystem model: Sensitivity analysis and parameter optimization, *J. Mar. Syst.*, 28(1–2), 45–63, doi:10.1016/S0924-7963(00)00083-X, 2001.
- 540 Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W.,
- 545 Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langle, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C.,
- 550 and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *Cryosph.*, 7(1), 375–393, doi:10.5194/tc-7-375-2013, 2013.
- Friedrichs, M. A. M.: A data assimilative marine ecosystem model of the central equatorial Pacific: Numerical twin experiments, *J. Mar. Res.*, 59, 859–894. doi:10.1357/00222400160497544, 2001.
- 555 Friedrichs, M. A. M.: Assimilation of JGOFS EqPac and SeaWiFS data into a marine ecosystem model of the central equatorial Pacific Ocean, *Deep-Sea Res. II*, 49, 289–319. doi:10.1016/S0967-0645(01)00104-7, 2002.
- Friedrichs, M. A. M., Hood, R. R., and Wiggert, J. D.: Ecosystem model complexity versus physical forcing: Quantification of their relative impact with assimilated Arabian Sea data, *Deep-Sea Res. II*, 53(5–7), 576–600, doi:10.1016/j.dsr2.2006.01.026, 2006.
- 560 Friedrichs, M. A. M., Dusenberry, J. A., Anderson, L. A., Armstrong, R. A., Chai, F., Christian, J. R., Doney, S. C., Dunne, J., Fujii, M., Hood, R., McGillicuddy Jr., D. J., Moore, J. K., Schartau, M., Spitz, Y. H., Wiggert, J. D.: Assessment of skill and portability in regional



- marine biogeochemical models: Role of multiple planktonic groups, *J. Geophys. Res. Ocean.*, 112(C08001), 1–22, doi:10.1029/2006JC003852, 2007.
- 565 Garrison, D. L., Gibson, A., Kunze, H., Gowing, M. M., Vickers, C. L., Mathot, S., and Bayre, R. C.: The Ross Sea Polynya Project: Diatom- and Phaeocystis-dominated phytoplankton assemblages in the Ross Sea, Antarctica, 1994–1996, in *Biogeochemistry of the Ross Sea, Antarctic Research Series*, vol. 78, pp. 53–76, 2003.
- 570 Gharamti, M. E., Samuelsen, A., Bertino, L., Simon, E., Korosov, A., and Daewel, U.: Online tuning of ocean biogeochemical model parameters using ensemble estimation techniques: Application to a one-dimensional model in the North Atlantic, *J. Mar. Syst.*, 168, 1–16, doi:10.1016/j.jmarsys.2016.12.003, 2017.
- Gregg, W. W.: Assimilation of SeaWiFS ocean chlorophyll data into a three-dimensional global ocean model, *J. Mar. Syst.*, 69, 205–225, doi:10.1016/j.jmarsys.2006.02.015, 2008.
- 575 Gregg, W. W., Friedrichs, M. A. M., Robinson, A. R., Rose, K. A., Schlitzer, R., Thompson, K. R., and Doney, S. C.: Skill assessment in ocean biological data assimilation, *J. Mar. Syst.*, 76(1–2), 16–33, doi:10.1016/j.jmarsys.2008.05.006, 2009.
- Hales, B., and Takahashi, T.: High-resolution biogeochemical investigation of the Ross Sea, Antarctica, during the AESOPS (U. S. JGOFS) Program, *Global Biogeochem. Cycles*, 18, 580 1–24, doi:10.1029/2003GB002165, 2004.
- Hemmings, J. C. P., Srokosz, M. A., Challenor, P., and Fasham, M. J. R.: Split-domain calibration of an ecosystem model using satellite ocean colour data, *J. Mar. Syst.*, 50(3–4), 141–179, doi:10.1016/j.jmarsys.2004.02.003, 2004.
- 585 Hemmings, J. C. P., and Challenor, P. G.: Addressing the impact of environmental uncertainty in plankton model calibration with a dedicated software system: the Marine Model Optimization Testbed (MarMOT 1.1 alpha), *Geosci. Model Dev.*, 5(2), 471–498, doi:10.5194/gmd-5-471-2012, 2012.
- Hemmings, J. C. P., Challenor, P. G., and Yool, A.: Mechanistic site-based emulation of a global ocean biogeochemical model (MEDUSA 1.0) for parametric analysis and calibration: an application of the Marine Model Optimization Testbed (MarMOT 1.1), *Geosci. Model Dev.*, 590 8(3), 697–731, doi:10.5194/gmd-8-697-2015, 2015.
- Hofmann, E. E., and Friedrichs, M. A. M.: Biogeochemical Data Assimilation, in *Encyclopedia of Ocean Sciences*, edited by J. H. Steele et al. pp. 364–370, Elsevier, 2001.



- 595 Hofmann, E. E., and Friedrichs, M. A. M.: Predictive Modeling for Marine Ecosystems, in *The Sea, Volume 12: Biological-Physical Interactions in the Sea*, edited by A. R. Robinson, J. J. McCarthy, and B. J. Rothschild, pp. 537–565, John Wiley & Sons, Inc., New York, 2002.
- Jones, E. M., Baird, M. E., Mongin, M., Parslow, J., Skerratt, J., Lovell, J., Margvelashvili, N., Matear, R. J., Wild-Allen, K., Robson, B., Rizwi, F., Oke, P., King, E., Schroeder, T.,
600 Steven, A., Taylor, J.: Use of remote-sensing reflectance to constrain a data assimilating marine biogeochemical model of the Great Barrier Reef, *Biogeosciences*, *13*, 6441–6469, doi:10.5194/bg-13-6441-2016, 2016.
- Jones, R. M., and Smith, W. O.: The influence of short-term events on the hydrographic and biological structure of the southwestern Ross Sea, *J. Mar. Syst.*, *166*, 184–195,
605 doi:10.1016/j.jmarsys.2016.09.006, 2017.
- Kaufman, D. E., Friedrichs, M. A. M., Smith, W. O., Queste, B. Y., and Heywood, K. J.: Biogeochemical variability in the southern Ross Sea as observed by a glider deployment, *Deep Sea Res. I*, *92*, 93–106, doi:10.1016/j.dsr.2014.06.011, 2014.
- Kaufman, D. E., Friedrichs, M. A. M., Smith, W. O., Hofmann, E. E., Dinniman, M. S., and
610 Hemmings, J. C. P.: Climate change impacts on southern Ross Sea phytoplankton composition, productivity, and export, *J. Geophys. Res. Ocean.*, *122*(3), 2339–2359, doi:10.1002/2016JC012514, 2017a.
- Kaufman, D.E., Friedrichs, M.A.M., Smith, W.O., Jr., Hofmann, E.E., Dinniman, M.S., Hemmings, J.C.P.: Associated Dataset: Climate change impacts on southern Ross Sea
615 phytoplankton composition, productivity and export. Virginia Institute of Marine Science. College of William and Mary. <http://doi.org/10.21220/V5PC71>, 2017b.
- Krishnakumar, K.: Micro-genetic algorithms for stationary and non-stationary function optimization, *Proc. SPIE. 1196, Intell. Control Adapt. Syst.*, *289*, doi:10.1117/12.969927, 1990.
- 620 Lawson, L. M., Spitz, Y. H., Hofmann, E. E., and Long, R. B.: A data assimilation technique applied to a predator-prey model, *Bull. Math. Biol.*, *57*(4), 593–617, doi:10.1007/BF02460785, 1995.
- Lawson, L. M., Hofmann, E. E., and Spitz, Y. H.: Time series sampling and data assimilation in a simple marine ecosystem model, *Deep-Sea Res. II*, *43*(2–3), 625–651, doi:10.1016/0967-
625 0645(95)00096-8, 1996.



- Leonelli, S.: “The Impure Nature of Biological Knowledge and the Practice of Understanding.”
Scientific Understanding: Philosophical Perspectives, Pittsburgh University Press, 1–27,
2009.
- Li, Y., McGillicuddy Jr., D. J., Dinniman, M. S., and Klinck, J. M.: Processes influencing
630 formation of low-salinity high-biomass lenses near the edge of the Ross Ice Shelf, *J. Mar.
Syst.*, *166*, 108–119, doi:10.1016/j.jmarsys.2016.07.002, 2017.
- Little, H.: Quantifying spatial and temporal scales of phytoplankton variability in the Sub-
Antarctic Ocean using a high-resolution glider dataset, University of Cape Town, Thesis,
2016.
- 635 Löptien, U., and Dietze, H.: Constraining parameters in marine pelagic ecosystem models – is it
actually feasible with typical observations of standing stocks?, *Ocean Sci.*, *11*(4), 573–590,
doi:10.5194/os-11-573-2015, 2015.
- Marsay, C. M., Sedwick, P. N., Dinniman, M. S., Barrett, P. M., Mack, S. L., and McGillicuddy
Jr., D. J.: Estimating the benthic efflux of dissolved iron on the Ross Sea continental shelf,
640 *Geophys. Res. Lett.*, *41*, 7576–7583, doi:10.1002/2014GL061684, 2014.
- Matear, R. J.: Parameter optimization and analysis of ecosystem models using simulated
annealing: A case study at Station P, *J. Mar. Res.*, *53*, 571–607,
doi:10.1357/0022240953213098, 1995.
- Mathot, S., Smith, W. O., Carlson, C. A., Garrison, D. L., Gowing, M. M., and Vickers, C. L.:
645 Carbon partitioning within *Phaeocystis antarctica* (Prymnesiophyceae) colonies in the Ross
Sea, Antarctica, *J. Phycol.*, *36*, 1049–1056, doi:10.1046/j.1529-8817.2000.99078.x, 2000.
- Melbourne-Thomas, J., Wotherspoon, S., Corney, S., Molina-Balari, E., Marini, O., and
Constable, A.: Optimal control and system limitation in a Southern Ocean ecosystem
model, *Deep. Res. Part II Top. Stud. Oceanogr.*, *114*, 64–73,
650 doi:10.1016/j.dsr2.2013.02.017, 2015.
- Mosby, A., and Smith, W. O.: Phytoplankton growth rates in the Ross Sea, Antarctica, *Aquat.
Microb. Ecol.*, *74*, 157–171, doi:10.3354/ame01733, 2015.
- Peloquin, J. A., and Smith, W. O.: Phytoplankton blooms in the Ross Sea, Antarctica:
Interannual variability in magnitude, temporal patterns, and composition, *J. Geophys. Res.*,
655 *112*(C08013), 1–12, doi:10.1029/2006JC003816, 2007.



- Powell, M. J. D.: An efficient method for finding the minimum of a function of several variables without calculating derivatives, *Comput. J.*, 7(2), 155–162, doi:10.1093/comjnl/7.2.155, 1964.
- 660 Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.: Numerical recipes in C: the art of scientific computing, 2nd ed., Cambridge University Press, 1992.
- Rudnick, D. L.: Ocean Research Enabled by Underwater Gliders, *Ann. Rev. Mar. Sci.*, 8(1), 519–541, doi:10.1146/annurev-marine-122414-033913, 2016.
- Saba, V. S., Friedrichs, M. A. M., Antoine, D., Armstrong, R. A., Asanuma, I., Behrenfeld, M., J. Ciotti, A. M., Dowell, M., Hoepffner, N., Hyde, K. J. W., Ishizaka, J., Kameda, T.,
665 Marra, J., Mélin, F., Morel, A., O'Reilly, J., Scardi, M., Smith, W. O., Smyth, T. J., Tang, S., Uitz, J., Waters, K., and Westberry, T. K.: An evaluation of ocean color model estimates of marine primary productivity in coastal and pelagic regions across the globe, *Biogeosciences*, 8(2), 489–503, doi:10.5194/bg-8-489-2011, 2011.
- Schartau, M., and Oschlies, A.: Simultaneous data-based optimization of a 1D-ecosystem model
670 at three locations in the North Atlantic: Part II - Standing stocks and nitrogen fluxes, *J. Mar. Res.*, 61(6), 795–821, doi:10.1357/002224003322981156, 2003.
- Schartau, M., Wallhead, P., Hemmings, J., Löptien, U., Kriest, I., Krishna, S., Ward, B. A., Slawig, T., and Oschlies, A.: Reviews and syntheses : parameter identification in marine planktonic ecosystem modelling, *Biogeosciences*, 14, 1647–1701, doi:10.5194/bg-14-1647-
675 2017, 2017.
- Schine, C. M. S., van Dijken, G., and Arrigo, K. R.: Spatial analysis of trends in primary production and relationships with large-scale climate variability in the Ross Sea, Antarctica (1997-2013), *J. Geophys. Res. Ocean.*, 120, 1–19, doi:10.1002/2015JC011014, 2015.
- Schmitt, L. M.: Theory of genetic algorithms, *Theor. Comput. Sci.*, 259(1–2), 1–61,
680 doi:10.1016/S0304-3975(00)00406-0, 2001.
- Sedwick, P. N., Marsay, C. M., Sohst, B. M., Aguilar-Islas, A. M., Lohan, M. C., Long, M. C., Arrigo, K. R., Dunbar, R. B., Saito, M. A., Smith, W. O., and DiTullio, G. R.: Early season depletion of dissolved iron in the Ross Sea polynya: Implications for iron dynamics on the Antarctic continental shelf, *J. Geophys. Res.*, 116(C12019), 1–19,
685 doi:10.1029/2010JC006553, 2011.



- Smith, W. O., and Nelson, D. M.: Phytoplankton bloom produced by a receding ice edge in the Ross Sea: spatial coherence with the density field, *Science*, 227(8), 163–166, doi:10.1126/science.227.4683.163, 1985.
- 690 Smith, W. O., and Gordon, L. I.: Hyperproductivity of the Ross Sea (Antarctica) polynya during austral spring, *Geophys. Res. Lett.*, 24(3), 233–236, doi:10.1029/96GL03926, 1997.
- Smith, W. O., Nelson, D. M., and Mathot, S.: Phytoplankton growth rates in the Ross Sea, Antarctica, determined by independent methods: temporal variations, *J. Plankton Res.*, 21(8), 1519–1536. doi:10.1093/plankt/21.8.1519, 1999.
- 695 Smith, W. O., Dinniman, M. S., Tozzi, S., DiTullio, G. R., Mangoni, O., Modigh, M., and Saggiomo, V.: Phytoplankton photosynthetic pigments in the Ross Sea: Patterns and relationships among functional groups, *J. Mar. Syst.*, 82(3), 177–185, doi:10.1016/j.jmarsys.2010.04.014, 2010.
- Smith, W. O., Shields, A. R., Dreyer, J. C., Peloquin, J. A., and Asper, V.: Interannual variability in vertical export in the Ross Sea: Magnitude, composition, and environmental correlates, 700 *Deep-Sea Res. I*, 58, 147–159, doi:10.1016/j.dsr.2010.11.007, 2011.
- Smith, W. O., Tozzi, S., Long, M. C., Sedwick, P. N., Peloquin, J. A., Dunbar, R. B., Hutchins, D. A., Kolber, Z., and DiTullio, G. R.: Spatial and temporal variations in variable fluorescence in the Ross Sea (Antarctica): Oceanographic correlates and bloom dynamics, *Deep-Sea Res. I*, 79, 141–155, doi:10.1016/j.dsr.2013.05.002, 2013.
- 705 Smith Jr., W. O., Goetz, K. T., Kaufman, D. E., Queste, B. Y., Asper, V., Costa, D. P., Dinniman, M. S., Friedrichs, M. A. M., Hofmann, E. E., Heywood, K. J., Klinck, J. M., Kohut, J. T., and Lee, C. M.: Multiplatform, multidisciplinary investigations of the impacts of Modified Circumpolar Deep Water in the Ross Sea, Antarctica, *Oceanography*, 27(2), 180–185, doi:10.5670/oceanog.2014.36, 2014.
- 710 Smith, W. O., McGillicuddy, D. J., Olson, E. B., Kosnyrev, V., Peacock, E. E., and Sosik, H. M.: Mesoscale variability in intact and ghost colonies of *Phaeocystis antarctica* in the Ross Sea: Distribution and abundance, *J. Mar. Syst.*, 166, 97–107, doi:10.1016/j.jmarsys.2016.05.007, 2017.
- 715 Stukel, M. R., Kahru, M., Benitez-Nelson, C. R., Decima, M., Goericke, R., Landry, M. R., and Ohman, M. D.: Using lagrangian-based process studies to test satellite algorithms of vertical



- carbon flux in the eastern North Pacific Ocean, *J. Geophys. Res. Ocean.*, *120*, 1–15, doi:10.1002/2015JC011264, 2015.
- Thomalla, S. J., Ogunkoya, A. G., Vichi, M., and Swart, S.: Using Optical Sensors on Gliders to Estimate Phytoplankton Carbon Concentrations and Chlorophyll-to-Carbon Ratios in the Southern Ocean, *Front. Mar. Sci.*, *4*(February), 1–19, doi:10.3389/fmars.2017.00034, 2017.
- 720 Vaillancourt, R. D., Marra, J., Barber, R. T., and Smith, W. O.: Primary productivity and in situ quantum yields in the Ross Sea and Pacific Sector of the Antarctic Circumpolar Current, *Deep-Sea Res. II*, *50*(3–4), 559–578, doi:10.1016/S0967-0645(02)00584-2, 2003.
- Vallverdú, J.: What are Simulations? An Epistemological Approach, *Procedia Technol.*, *13*, 6–15, doi:10.1016/j.protcy.2014.02.003, 2014.
- 725 Ward, B. A., Friedrichs, M. A. M., Anderson, T. R., and Oschlies, A.: Parameter optimisation techniques and the problem of underdetermination in marine biogeochemical models, *J. Mar. Syst.*, *81*(1–2), 34–43, doi:10.1016/j.jmarsys.2009.12.005, 2010.
- Xiao, Y., and Friedrichs, M. A. M.: Using biogeochemical data assimilation to assess the relative skill of multiple ecosystem models in the Mid-Atlantic Bight: effects of increasing the complexity of the planktonic food web, *Biogeosciences*, *11*(11), 3015–3030, doi:10.5194/bg-11-3015-2014, 2014a.
- 730 Xiao, Y., and Friedrichs, M. A. M.: The assimilation of satellite-derived data into a one-dimensional lower trophic level marine ecosystem model, *J. Geophys. Res. Ocean.*, *119*, 2691–2712, doi:10.1002/2013JC009433, 2014b.
- 735 Yool, A., Popova, E. E., and Anderson, T. R.: Medusa-1.0: a new intermediate complexity plankton ecosystem model for the global domain, *Geosci. Model Dev.*, *4*(2), 381–417, doi:10.5194/gmd-4-381-2011, 2011.

740 **Table 1:** Eight parameters optimized in this analysis.

Parameter Name	Initial value (Kaufman et al., 2017a)	Bounds (lower, upper)
Diatom max growth rate at 0°C	0.375 (d ⁻¹)	0.09375, 1.5
<i>P. antarctica</i> solitary cells C:Chl ratio	30 (gC gChl ⁻¹)	7.5, 120
<i>P. antarctica</i> colonies max growth rate at 0°C	0.5 (d ⁻¹)	0.125, 2
<i>P. antarctica</i> solitary cells max growth rate at 0°C	0.5 (d ⁻¹)	0.125, 2
Diatom C:Chl ratio	150 (gC gChl ⁻¹)	37.5, 600
Fast detritus sinking fraction of diatom losses	0.75	0.5, 0.95
<i>P. antarctica</i> colonies max sinking rate	20 (m d ⁻¹)	5, 80
<i>P. antarctica</i> colonies C:Chl ratio	40 (gC gChl ⁻¹)	10, 160



745 **Table 2:** Time, depth, and time-space resolution of glider-based observations of Chl and POC assimilated for each experiment.

Experiment	Depth (m)	Temporal Resolution	Spatial Area(s)
Expt 1a: Full Assimilation	0 - 50	Hourly	Full glider track
Expt 1b: Latitudinal Assim.	0 - 50	Hourly	North, Central, South Latitudinal bands
Expt 1c: Longitudinal Assim.	0 - 50	Hourly	East, Central, West Longitudinal bands
Expt 2a: Glider Assimilation	0 - 50	~ twice per day, separated at a minimum of 12 hours.	Full glider track
Expt 2b: Cruise-based Assim.	0 - 50	3 days in a row, and then another 3 consecutive days two weeks later	Full glider track
Expt 2c: Satellite-based Assim.	0 - 5	1 day every two weeks	Full glider track



750 **Table 3:** Depth- and time-integrated primary production (PP), carbon export flux at 200 m, and costs for the No Assimilation run (cost = 0.77), Experiment #1 and #2.

Simulation name	PP (g C m⁻² y⁻¹)	Export (g C m⁻² y⁻¹)	Predictive Cost (\mathcal{J}_P)	Assim. Cost (\mathcal{J}_A)
No assimilation	111.7	18.8	-	-
Expt 1a: Full Assimilation	104.2	27.2	-	0.41
Expt 1b: Latitudinal Assim.	101.8* ±3.3	26.1 ±2.1	0.49 ±0.13	0.43 ±0.14
Expt 1c: Longitudinal Assim.	103.2 ±2.1	26.9 ±2.1	0.50 ±0.10	0.46 ±0.13
Expt 2a: Glider Assim.	103.7 ±1.8	27.0 ±1.2	0.43 ±0.01	0.43 ±0.03
Expt 2b: Cruise-based Assim.	113.1 ±22.3	24.8 ±6.6	1.24 ±0.95	0.52 ±0.19
Expt 2c: Satellite-based Assim.	114.1 ±10.7	16.7 ±2.7	1.04 ±0.36	0.26 ±0.16

* costs represent mean ± one standard deviation of assimilative runs.

755



760

Table 4: Initial parameter values (No Assimilation) and optimal parameter values after conducting the Full Assimilation, Glider, Cruise-based, and Satellite-based Assimilation experiments

Parameter Name	Initial Value	Expt 1a Full Assimilation	Expt 2a Glider*	Expt 2b Cruise-based*	Expt 2c Satellite-based*
Diatom max growth rate at 0°C (d ⁻¹)	0.375	0.40	0.43 ±0.01	0.42 ±0.15	0.41 ±0.09
<i>P. antarctica</i> solitary cells C:Chl ratio (gC gChl ⁻¹)	30	29.7	25.84 ±5.16	37.3 ±26.7	51.5 ±26.8
<i>P. antarctica</i> colonies max growth rate at 0°C (d ⁻¹)	0.5	0.29	0.22 ±0.10	0.45 ±0.58	0.29 ±0.17
<i>P. antarctica</i> solitary cells max growth rate at 0°C (d ⁻¹)	0.5	0.39	0.45 ±0.06	0.75 ±0.70	0.79 ±0.51
Diatom C:Chl ratio (gC gChl ⁻¹)	150	176.4	166.6 ±50.17	252.4 ±164.28	374.86 ±187.82
Fast detritus sinking fraction of diatom losses	0.75	0.87	0.86 ±0.05	0.86 ±0.11	0.62 ±0.14
<i>P. antarctica</i> colonies max sinking rate (m d ⁻¹)	20	10.7	10.1 ±3.66	20.1 ±20.5	12.8 ±9.27
<i>P. antarctica</i> colonies C:Chl ratio (gC gChl ⁻¹)	40	14.0	14.2 ±2.29	42.7 ±41.6	34.3 ±26.5

* mean ± one standard deviation of assimilative runs.




Figure 1: Southern Ross Sea showing transect locations where the glider was at the surface. The color of each glider dive indicates the date. Bathymetric contours are shown at 200-m intervals, as obtained from the bedmap2 bathymetric data [Fretwell *et al.*, 2013].




Figure 2: Variance explained in model outputs by parameters during sensitivity tests using Latin hypercube sampling of parameter space. Only parameters with at least one r^2 value greater than or equal to 0.01 (vertical dotted line) are shown.




Figure 3: (a) Minimum costs and (b) normalized parameter values in numerical twin experiments, illustrating that the assimilation procedure is unable to successfully recover the true parameter values when more than eight parameters are optimized. One data point in three of the experiments (#s 19, 20, and 21) exceeds the y-axis upper limit in the lower (b) panel.




Figure 4: Locations of glider observations assimilated in (a) Experiment 1b –latitudinal bands, and (b) Experiment 1c – longitudinal bands. Colors represent the three spatial bands of data assimilated.




Figure 5 (a-f): Upper 50 m mean concentrations of (a,c,e) Chl and (b,d,f) POC for various experiments assimilating the full glider and from different spatial areas (Table 2): (a,b) Experiment 1a, (c,d) Experiment 1b - latitude bands, and (e,f) Experiment 1c - longitude bands. For reference, model results for the Full Assimilation case (orange lines), and glider data (black lines) with shading (gray) representing one standard deviation are included in each panel. Colored boxes at the top of each panel indicate times of assimilated observations.




Figure 6: Upper 50 m mean concentrations of the three phytoplankton groups in terms of (a,c) Chl and (b,d) POC for the No Assimilation case (a,b) and the Full Assimilation case (c,d). The glider data are shown (black line) with shading (gray) that represents one standard deviation daily.




Figure 7: Distributions of (a) depth- and time-integrated production and (b) carbon export flux at 200 m for each assimilation experiment (Table 2). The median value for each experiment is indicated by a horizontal light-blue line. Each box extends vertically from the 1st to 3rd quartile, and the whiskers extend from the lowest to highest values. Individual values are shown as grey dots. For reference, production and export estimates from the No Assimilation (solid blue line) and Full Assimilation (dashed gray line) cases are included in each panel.




Figure 8 (a-f): Upper 50 m mean concentrations of (a,c,e) Chl and (b,d,f) POC for various experiments assimilating subsets characteristic of the original glider data, cruise-based observations and satellite-based observations (Table 2): (a,b) Experiment 2a - glider observations, (c,d) Experiment 2b - cruise-based, and (e,f) Experiment 2c - satellite-based. For reference, model results for the Full Assimilation case (orange lines), and glider data (black lines) with shading (gray) representing one standard deviation are included in each panel.