

Interactive comment on “Geomorphic influences on the contribution of vegetation to soil C accumulation and accretion in *Spartina alterniflora* marshes” by Tracy Elsey-Quirk and Viktoria Unger

Anonymous Referee #3

Received and published: 6 October 2017

This paper is a substantial and interesting addition to the literature and I think that it could be publishable for Biogeosciences Discussions pending some revisions. The study reports correlations between geomorphic variation in variables such as tidal elevation, tidal frame, suspended sediment, salinity, etc. with important biotic variables affecting carbon sequestration (above and below ground biomass, root addition, root-shoot ratio, and recalcitrant/labile carbon fraction), and observations of Carbon Accumulation Rate, measured by radiocesium date and previously published by Unger et al. and Boyd et al.. The observations that complex correlations between root production and drainage, and sediment trapping having multiple positive interactions with carbon burial, are compelling conclusions as they point to the complex and dynamic nature of

Printer-friendly version

Discussion paper



tidal wetland systems more generally. Predicting behavior outside of well-studied plots and sites is a large concern of the community and this paper speaks to many difficulties in those efforts.

I think what sets this paper apart from much of the literature is how well monitored all of the sites are. All locations have measured elevation, inundation, and soil properties. This should be commended and in many ways is close to an ideal salt marsh carbon dynamics study design.

I have three major critiques of the paper, somewhat overlapping. 1. There is not enough available methods data for the calculation of tidal datums from the water loggers. I found some of the inclusion of comparing NAVD88 elevation, MHW and MLW hard to follow, especially when these were used as proxies for multiple hydrologic properties. I was a bit taken aback at how much variation there were in datums that are located fairly close together. Could this be because of the short 1.5-year time period? 'No data' values deflating MLW datums? Etc? Is there really that much local variation in datums? I would like to see more information before making a judgment there.

2. This leads me to my second critique. A lot of the correlation analysis could be paired down. First because of what I discussed in one, maybe some of these measures are redundant or could be reduced to more directly causal variables.

For example, by converting site elevations (NAVD88) into dimensionless elevation $z = (\text{Elevation} - \text{MTL}) / (\text{MHHW} - \text{MTL})$ [Swanson et al., 2012], or focusing on flooding depth and # of floods a year (maybe converted to volume of water / unit time) since those are probably a clearer functional driver for organic and inorganic sedimentation. Maybe there's a better metric for drainage, such as an average time between inundation events.

The number of correlations discussed makes the text especially figure 7 a bit cluttered and hard to decipher on a quick read through (although there are benefits to being thorough). Statistically, the number of correlations presented is problematic. Which leads me to point 3.

[Printer-friendly version](#)[Discussion paper](#)

3. There are many correlations presented, but the threshold for significance (0.05) is not adjusted for multiple testing. The more correlations your run, the higher chance of getting false positives. Researchers often address this by using the Bonferroni correction, or some related correction (there are some arguments that the Bonferroni is too restrictive). The fact that many of the regressions presented are barely significant at the 0.05 level and many would no longer be significant after correcting for multiple testing. Maybe a combination of reducing the number of variables tested to a few functionally important variables and adjusting for multiple testing could allow the authors to focus more on the very clearly significant driving geomorphic variables.

Line comments: 63: This is an excellent point that does not get enough attention in the literature. 64: Is there any literature you could cite in the hazards literature or other ecosystem-climate change dynamics that discuss complexities in projecting system resilience? 132: What is the time frame of RSLR? Same as cores, or total length of the gauging period? 277: If dimensionless elevation may be a better fit than using MHW and MLW. Alternatively inundation time, the number of inundation events or cumulative annual mass of water seem like they would be much better variables to use as there is process-knowledge involved. 427: Is sediment trapping by biomass part of this positive feedback?

Tab 1 - What is MHW MWL and MLW relative to NAVD88? Station datum? Tab 3 - Far left column a bit hard to read. Maybe fix in formatting. Fig. 1 - Howe a basemap with better definition. Maybe one that emphasizes the differences between uplands systems and wetlands. I would delete the service layer credits and put it into the figure caption. The map looks low resolution.

Citations: Swanson, Kathleen M., et al. "Wetland accretion rate model of ecosystem resilience (WARMER) and its application to habitat sustainability for endangered species in the San Francisco Estuary." *Estuaries and Coasts* 37.2 (2014): 476-492.

Abdi, H. (2007). Bonferroni and Šidák corrections for multiple comparisons. *Encyclo-*

[Printer-friendly version](#)[Discussion paper](#)

pedia of measurement and statistics, 3, 103-107.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2017-268>, 2017.

BGD

Interactive
comment

Printer-friendly version

Discussion paper

