



1 Virus mediated transfer of nitrogen from heterotrophic bacteria to
2 phytoplankton

3 Emma J. Shelford¹ and Curtis A. Suttle^{1,2,3,4,5}

4 ¹Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia,
5 Vancouver, Canada

6 ²Department of Botany, University of British Columbia, Vancouver, Canada

7 ³Department of Microbiology and Immunology, University of British Columbia, Vancouver,
8 Canada

9 ⁴Institute of Oceans and Fisheries, University of British Columbia, Vancouver, Canada

10 ⁵Canadian Institute for Advanced Research, Toronto, Canada.

11

12 Correspondence to: C. A. Suttle (suttle@science.ubc.ca)

13

14



1 Abstract

2

3 Lytic infection of bacteria by viruses releases nutrients during cell lysis and stimulates the
4 growth of primary producers, but the path by which these nutrients flow from lysates to primary
5 producers has not been traced. This study examines the remineralisation of nitrogen (N) from
6 bacterial lysates by heterotrophic bacteria and its transfer to primary producers. In laboratory
7 trials, *Vibrio* sp. strain PWH3a was infected with a lytic virus (PWH3a-P1) and the resulting 36.0
8 μM of dissolved organic N (DON) in the lysate was added to cultures containing cyanobacteria
9 (*Synechococcus* sp. strain DC2), and a natural bacterial assemblage. Based on the increase in
10 cyanobacteria, 74% (26.5 μM N) of the DON in the lysate was remineralised and taken up by
11 *Synechococcus* sp. strain DC2 cells. Lysate from *Vibrio* sp. strain PWH3a labeled with $^{15}\text{NH}_4^+$
12 was also added to seawater containing natural microbial communities, and in four field
13 experiments, stable isotope analysis indicated that the uptake of labeled N was 0.09 to 0.70 μmol
14 N per μg of chlorophyll *a*. The results from these experiments demonstrate that DON from
15 bacterial lysate can be efficiently remineralised and transferred to phytoplankton, and provides
16 further evidence that the viral shunt is an important link in nitrogen recycling in aquatic systems.

17

18 1 Introduction

19 Viruses are significant mortality agents of bacteria and phytoplankton in the ocean, consequently
20 affecting pathways and rates of nutrient cycling (Fuhrman, 1999; Suttle, 2005, 2007; Wilhelm
21 and Suttle, 1999) when particulate and dissolved organic matter (DOM) is released into the water
22 during cell lysis (Gobler et al., 1997; Middelboe and Jørgensen, 2006). Major constituents of the



1 released material are free and combined amino acids (Middelboe and Jørgensen, 2006), which
2 are taken up and metabolised by bacteria (Middelboe et al., 1996, 2003). When the C:N of DOM
3 is low relative to the nutritional requirements of heterotrophic bacteria, ammonium may be
4 released (Goldman et al., 1987; Hollibaugh, 1978), which evidence suggests can support
5 phytoplankton growth (Haaber and Middelboe, 2009; Shelford et al., 2012; Weinbauer et al.,
6 2011).

7 High rates of bacterial mortality from viral lysis imply a continuous and substantial flux of DOM
8 from lysates into seawater. Weinbauer et al. (2011) provided evidence of the importance of this
9 flux by showing that reducing viral abundance decreased the growth of *Synechococcus*, the
10 dominant primary producer during their experiments in the Gulf of Mexico and Mediterranean
11 Sea. It was postulated that *Synechococcus* growth may have been directly stimulated by uptake
12 of dissolved organic nutrients released by lysis, or indirectly through the incorporation of these
13 organics by uninfected bacteria and subsequent remineralisation of inorganic nutrients. Evidence
14 that mineralization of DOM and release of ammonium by uninfected bacteria stimulates
15 phytoplankton growth was shown by Shelford et al. (2012).

16 The present contribution demonstrates, in the laboratory and field, that uninfected bacteria
17 metabolise dissolved organic N (DON) released as the result of viral lysis of bacteria and release
18 ammonium that supports the growth of phytoplankton.

19



1 2 **Methods**

2 2.1 **Laboratory cultures**

3 A semi-continuous culture of *Synechococcus* sp. strain DC2 (Bigelow, CCMP #1334; WH7803)
4 was grown on artificial seawater (Berges et al., 2001), modified by adding 5 M bicine (Healey
5 and Hendzel, 1979), 124 μM NH_4Cl instead of nitrate, and 13 μM K_2HPO_4 . Cultures were
6 maintained at 19°C and continuous light ($42 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$). Experiments were started
7 when cultures entered nutrient limitation near the end of exponential growth, as determined by
8 epifluorescence microscopy counts.

9 The gram negative marine bacterium *Vibrio* sp. strain PWH3a, also known as *Vibrio natriegens*
10 strain PWH3a (Suttle and Chen, 1992; Weinbauer et al., 1997) and *Vibrio alginolyticus* strain
11 PWH3a (Poorvin et al. 2011) was grown on artificial seawater with 5 mM bicine, 500 μM
12 NH_4Cl , 100 μM K_2HPO_4 , and 1 mM glucose as a carbon source for a C:N:P ratio of 60:5:1. The
13 cultures were grown at 25°C and continuously mixed at 100 rpm.

14 Bacteriophage PWH3a-P1 was added in eight-fold excess abundance (multiplicity of infection of
15 8:1) to cultures of *Vibrio* sp. strain PWH3a at the end of exponential growth, as determined by
16 absorbance at 660 nm (Ultraspex spectrophotometer, Biochrom, United Kingdom). The culture
17 was incubated with the virus until absorbance decreased to 20% of the initial (~7 h). The lysate
18 (Lys) was filtered through a 0.22 μm pore-size Durapore membrane (Millipore, Billerica, MA)
19 and kept at 4°C for approximately 20 h. The number of cells lysed prior to filtration was
20 determined by flow cytometry as described below (section 2.4.1). The amount of DON released
21 was determined by the number of cells lysed multiplied by the measured cellular N quota for



1 *Vibrio* sp. strain PWH3a, 2.54 fmol cell⁻¹ as described below (section 2.4.4). The result is the
2 amount of total N released by lysis of *Vibrio* sp. strain PWH3a.

3 Lysates for field experiments were prepared as above, except that *Vibrio* sp. strain PWH3a was
4 grown on ¹⁵NH₄Cl instead of ¹⁴NH₄Cl (90+ atom % ¹⁵N, Isotec, Miamisburg, OH), and the
5 filtered lysate was kept at 4°C for 2 to 5 d until the experiments were initiated.

6 **2.2 Growth of *Synechococcus* sp. strain DC2 on lysate from *Vibrio* sp. strain** 7 **PWH3a**

8 Water was collected from Queen Charlotte Sound (51° 02.37N, 127° 52.38W) on 2 Oct 2011.
9 Temperature and salinity were 11°C and 32, respectively. Nitrate and phosphate concentrations
10 were 21.3 and 1.9 μM, respectively. The water was ultrafiltered using a 30 kDa molecular weight
11 cut-off tangential flow cartridge (Prep/Scale, Millipore, Billerica, MA) after pre-filtration
12 through a 0.45 μm pore-size polyvinylidene fluoride (PVDF) Durapore membrane (Millipore,
13 Billerica, MA) as outlined in Suttle et al. (1991). The ultrafiltrate was stored in the dark at 4°C
14 for a year until used in experiments. Bacteria in the ultrafiltrate grew to a density of 4.18 × 10⁵
15 cells ml⁻¹, as determined by flow cytometry (described in section 2.4.1), and were used for the
16 regeneration experiment described below.

17 The experiment used *Synechococcus* sp. strain DC2, lysate from *Vibrio* sp. strain
18 PWH3a, and the bacterial assemblage (Bac) in ultrafiltrate from Queen Charlotte Sound in the
19 following six combinations (Figure 1): 1) DC2+Bac+lysate was the experimental treatment with
20 DON from lysate; 2) DC2+Bac was a control for *Synechococcus* sp. strain DC2 growth in the
21 presence of the bacterial assemblage without a DON source from lysate; 3) DC2+lysate was a
22 control for bacterial remineralisation in the *Synechococcus* sp. strain DC2 culture; 4) Bac+lysate



1 was a control to quantify ammonium remineralisation by the bacterial assemblage with the
2 addition of lysate; 5) Bac only was a control to determine the ammonium concentration of the
3 bacterial assemblage by itself; 6) DC2 only was a control to determine the ammonium
4 concentration and increase in cell number activity of *Synechococcus* sp. strain DC2 by itself. All
5 treatments were in triplicate in 1L polycarbonate Erlenmeyer flasks (Corning, New York). To
6 each appropriate treatment was added 10 ml of *Synechococcus* sp. strain DC2 culture, 100 ml of
7 bacterial assemblage, and/or 10 ml of lysate. The experimental treatment volume was 200 ml,
8 and volumes of control treatments were topped up to 200 ml by adding nitrate- and phosphate-
9 free artificial seawater to control treatments.

10 Treatments were incubated in the laboratory at 19°C under continuous light (42 μmol
11 quanta $\text{m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation) and sampled daily for cell abundance and
12 ammonium concentration.

13 **2.3 Uptake of ^{15}N from lysates in the field**

14 Water was collected from the surface at Saanich Inlet (SI), the Fraser River Plume (FRP),
15 Semiahoo Bay (SB) and Jericho Pier (JP) in southern coastal British Columbia (Table 1). SI and
16 FRP samples were collected using Go-Flo bottles mounted on a rosette, which held a SBE 25
17 CTD (Sea-Bird, Bellevue, WA) for measuring temperature and salinity. SB and JP samples were
18 collected by hand using a 20 l carboy rinsed with 10% HCl and ultrapure water, and temperature
19 and salinity were measured with a hand-held thermometer and refractometer. Samples for
20 phytoplankton identification were collected from the whole water and preserved with Lugol's
21 acidic iodine solution (Edler and Elbrächter, 2010). The water was filtered through 118- μm
22 mesh-size Nitex screening to remove large particulates.



1 For each experiment, 0.22- μm filtered lysate from *Vibrio* sp. strain PWH3a was added to
2 Nitex-filtered seawater (SW+lysate) and compared to a control containing 0.22- μm filtered
3 seawater. For SB and JP, a third treatment included 0.22- μm filtered seawater and 0.9 μM
4 $^{15}\text{NH}_4\text{Cl}$ (SW+N), to confirm that N stimulated production and that another factor was not
5 limiting phytoplankton growth, and to mirror the estimated N contributed by the lysate in the
6 SW+lysate treatment. Treatments were in triplicate. SI and FRP experiments were done in an on-
7 deck flow-through incubator with neutral density screening to reduce sunlight to approximate *in*
8 *situ* irradiance. For SB and JP, experiments were done at 19°C and 42 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$
9 continuous irradiance. Samples for cell abundance and ammonium concentration were collected
10 every 6 to 9 h for 1 to 2 d; samples for chlorophyll *a* were collected at the initial and final time
11 points, and samples for PO^{15}N were collected at the final time point.

12 **2.4 Sample Analysis**

13 **2.4.1 Cell and virus counts**

14 One ml samples were fixed with a final concentration of 0.5% glutaraldehyde for bacteria and
15 viruses, and 2% formaldehyde for phytoplankton, and flash frozen in liquid nitrogen and stored
16 at -80°C. Bacteria and viruses were stained with SYBR Green I (Sigma-Aldrich, St. Louis, MO),
17 while picophytoplankton were left unstained, immediately before counting by flow cytometry
18 (FACSCalibur, Becton Dickinson, Franklin Lakes, NJ) following the procedures of Brussaard
19 (2004), Gasol and Del Giorgio (2000), and Olson et al. (1993). Larger phytoplankton were
20 preserved with Lugol's solution, and the dominant taxa identified using a settling chamber (Edler
21 and Elbrächter, 2010) and light microscope (Axiovert 10, Zeiss, Canada).



1 2.4.2 Nutrient and chlorophyll *a* collection and analysis

2 Samples of 25 ml were filtered through acid-rinsed syringes fitted with MilliQ-soaked 0.45- μ m
3 pore size, 25-mm diameter, cellulose-nitrate filters (Whatman, United Kingdom) in a Swinnex
4 filter holder (Millipore, Billerica, MA). The first 15 ml were used to rinse the 15 ml acid-rinsed
5 polypropylene screw-cap collection tubes (Sarstedt, Germany), before collecting and freezing the
6 final 10 ml at -20°C for subsequent nutrient analysis. The filters were folded, placed in
7 aluminum foil packages, and frozen desiccated at -20°C until chlorophyll *a* was extracted using
8 90% acetone and sonication, and analysed fluorometrically (Turner Designs, 10AU fluorometer,
9 Sunnyvale, CA) following Parsons et al. (1984).

10 Nitrate+nitrite and phosphate concentrations were analyzed on a Bran & Luebbe
11 AutoAnalyzer 3 using air-segmented continuous-flow analysis (Technicon, Oakland, CA), while
12 ammonium concentrations were determined fluorometrically (Holmes et al., 1999) using a TD-
13 700 fluorometer (Turner Designs, Sunnyvale, CA).

14 2.4.3 PO¹⁵N size-fractionation, collection and determination of ¹⁵N enrichment

15 For the field experiments, the ¹⁵N incorporated was determined by collecting the particulate
16 material onto combusted (450°C for 4 h) glass-fibre (GF) filters (nominal pore size of 0.7 μ m,
17 Whatman, United Kingdom), which were subsequently dried at 50°C for 2 d. Prior to collection
18 onto the GF filters, the SI and FRP samples were poured into a Sterifil (Millipore, Germany)
19 filtration tower fitted with a 1- μ m pore-size polycarbonate filter (AMD Manufacturing, Canada).
20 While the sample was being gently filtered, the phytoplankton were kept in suspension and
21 rinsed, while the volume was maintained by adding 200 ml of ultrafiltrate. In this way, cells



1 captured by the 1- μm filter were retained for analysis of isotopic enrichment, while smaller cells
2 passed through. Samples for cell counts were taken before and after washing to determine the
3 proportion of cells lost by this process. Samples that were not rinsed with ultrafiltrate were also
4 filtered onto combusted GF filters to estimate the amount of ^{15}N uptake that was due to bacteria
5 that passed through the 1- μm pore size membrane. The SW+N treatment (SB and JP) was
6 filtered directly onto GF filters without rinsing. The $\delta^{15}\text{N}$ -PON and total PON on the filters was
7 determined at the Stable Isotope Laboratory at Boston University on a GV Instruments IsoPrime
8 isotope-ratio mass spectrometer and a Eurovector elemental analyzer, calibrated against
9 atmospheric N_2 and IAEA standards N-1, N-2, and N-3 (replicate analysis within $\pm 0.2\%$). Due
10 to instrument error, the stable-isotope data for SI and FRP are not replicated; whereas, duplicates
11 for JP and triplicates for SB were measured.

12 2.4.4 Particulate C and N analysis

13 N cell quotas of *Synechococcus* sp. strain DC2 and *Vibrio* sp. strain PWH3a were determined
14 from cultures grown using the same media and conditions as described in section 2.1. Cultures
15 entering stationary phase for *Synechococcus* sp. strain DC2 as determined by epifluorescence
16 microscopy counts, and mid-exponential phase for *Vibrio* sp. strain PWH3a as determined by
17 absorbance, were filtered onto combusted GF filters, dried at 50°C for two days, and placed in a
18 desiccator until packaged into aluminum foil and analyzed on a CHN Elemental Analyser (Carlo
19 Erba NA-1500).



1 2.4.5 Calculations and statistical analysis

2 Total uptake rates of ^{15}N in the field experiments were calculated by dividing the total particulate
3 ^{15}N on the filters by the volume filtered. Normalised uptake rates were calculated by dividing the
4 total uptake rate by the initial chlorophyll *a* concentration for each incubation. The percent
5 contribution of cells $>1\ \mu\text{m}$ to total ^{15}N uptake was calculated by dividing the total particulate
6 ^{15}N on the filters of the ‘washed’ samples (adjusted for decrease in phytoplankton abundance) by
7 the ^{15}N on the unwashed samples.

8 The differences between initial and final time points for *Synechococcus* abundances and
9 ammonium concentrations were normally distributed and with equal variances according to
10 Shapiro-Wilk and Levene’s tests, respectively; hence, the significance of the results was
11 analysed using one-way analysis of variance (ANOVA). The differences between initial and
12 final time points for bacterial abundances in the laboratory experiment, while normally
13 distributed, had unequal variances, and therefore a Kruskal-Wallis test was performed to test for
14 significant differences.

15 The differences between initial and final time points for ammonium and chlorophyll *a*
16 concentrations and bacteria and picophytoplankton abundances were normally distributed
17 (except for the picophytoplankton data for treatment SW in SI, which had only two samples),
18 and with equal variances (except the bacteria data for SI and JP). Significant differences between
19 treatments were determined using two-tailed Student’s *t* tests. Data which did not meet the
20 assumptions for the Student’s *t* test were analysed for significant differences using the non-
21 parametric Mann-Whitney *U* test.



1 Because the final time point of the bacterial data for the SW+lysate treatment in SI was
2 approximately six-fold lower than the previous time point, concurrent with a significant increase
3 in viral abundance (data not shown), significance between the two treatments was calculated for
4 the differences between the initial and the fourth time points. The data were normally distributed
5 and had equal variances (Shapiro-Wilk and Levene's tests), and so a Student's *t* test was run.

6

7 **3 RESULTS**

8

9 **3.1 Response of N-limited *Synechococcus* sp. strain DC2 to the addition of** 10 **lysate**

11 There was a greater increase in the abundance of N-limited *Synechococcus* sp. strain DC2 cells
12 exposed to lysate and a remineralising bacterial assemblage than when grown with each
13 component separately; the greatest increase occurred in the treatment DC2+Bac+lysate (Figure
14 2A), with final abundances significantly different among all treatments ($p < 0.05$). Ammonium
15 decreased from $\sim 1.0 \mu\text{M}$ to less than $0.2 \mu\text{M}$ except in the Bac+lysate treatment, in which it
16 increased to $8.24 \pm 0.04 \mu\text{M}$ (Figure 2B), significantly higher than in the other treatments
17 ($p < 0.05$). Bacterial abundance increased in all treatments, but the greatest increases occurred in
18 treatments with lysate addition (Figure 2C).

19 The proportion of N released by lysis of cultures of *Vibrio* sp. strain PWH3a that was taken up
20 by *Synechococcus* sp. strain DC2 cells was calculated using the N cell quotas for *Vibrio* sp. strain
21 PWH3a of $2.54 \text{ fmol cell}^{-1}$, and for *Synechococcus* sp. strain DC2 of $7.83 \text{ fmol cell}^{-1}$. Based on



1 the decrease in *Vibrio* sp. strain PWH3a cells from viral infection, multiplied by the N cell quota
2 of *Vibrio* sp. strain PWH3a, approximately 36.0 μM N was added to each incubation from
3 bacterial lysis (Table 2). The mean percent of N from lysate that was taken up by *Synechococcus*
4 sp. strain DC2 cells via remineralisation by bacteria was 74%.

5 **3.2 Uptake of N from bacterial lysate by primary producers in field studies**

6 Lysate from ^{15}N -labelled *Vibrio* sp. strain PWH3a cultures was added to seawater from Saanich
7 Inlet (SI), the Fraser River Plume (FRP), Semiahoo Bay (SB), and Jericho Pier (JP; Table 1).
8 Changes were followed in ammonium concentrations and in abundances of bacteria (Figure 3).
9 Bacterial abundances in SW+lysate treatments were significantly higher than in samples without
10 lysate (SW) by the final time point for FRP, SB, and JP ($p < 0.05$). In the SI sample, bacterial
11 abundances in the treatment with lysate added (SW+lysate) increased almost tenfold before
12 decreasing to below initial values by the final time point, concurrent with an increase in viral
13 abundance (data not shown). There were no significant differences between experimental and
14 control treatments for picophytoplankton abundances or for chlorophyll *a* concentrations in the
15 other stations (data not shown). Ammonium concentrations in SI, SB, and JP decreased to less
16 than 0.2 μM . In FRP, the SW+lysate treatment decreased to less than 0.04 μM before climbing
17 again to 0.25 μM , correlated with a spike in bacterial abundance. There was significantly greater
18 drawdown of ammonium in the SW+lysate treatment in every station except SB ($p < 0.05$). The
19 increase of bacteria and greater ammonium uptake at most stations indicates that bacterial
20 growth and overall uptake of ammonium was enhanced in these samples by the addition of
21 bacterial lysate.



1 There was uptake of lysate-derived N by cells $>1 \mu\text{m}$, which would be comprised primarily of
2 phytoplankton. The addition of lysate to the SW+lysate treatment for each station contributed
3 approximately $67.7 \mu\text{M N}$ for SI and FRP, and $0.44 \mu\text{M N}$ for SB and JP (Table 2). In order to
4 add a more reasonable concentration of N to the samples, the concentration of lysate N added to
5 SB and JP was greatly reduced compared to the other two previous experiments. Stable isotope
6 data collected from particulate organic material collected on $0.7\text{-}\mu\text{m}$ pore-size glass-fibre filters
7 indicated uptake of lysate by the particulate fraction (Table 3; Figure 4). From the ^{15}N data, the
8 calculated contribution to the total uptake by cells in the $>1 \mu\text{m}$ size fraction was 46.3% (SI),
9 47.6% (FRP), and 100% (SB and JP). The large ($>50\%$) contribution of the bacterial size fraction
10 to ^{15}N uptake in SI and FRP (Figure 4) corresponded to an increase in bacterial abundances
11 (Figure 3).

12 There was confirmed uptake of $^{15}\text{NH}_4$ in the treatment with ^{15}N addition (SW+N), with $0.61 \pm$
13 $0.20 \mu\text{M N}$ (SB) and $0.44 \pm 0.26 \mu\text{M N}$ (JP) taken up into the particulate fraction (corrected for
14 90 atom % ^{15}N ; data not shown). Any ammonium produced by remineralisation of lysate in the
15 SW+Lys treatment, therefore, could be used by the microbes at these two stations.

16

17 **4 DISCUSSION**

18

19 Experiments in the laboratory and with natural systems show that N in viral lysates can be
20 remineralised by bacterial communities to fuel primary production. Studies have shown that lysis
21 by viruses produces bioavailable dissolved organic matter (e.g. Middelboe et al. 2003, Poorvin et
22 al. 2004), and that phytoplankton lysate can be remineralised (Gobler et al., 1997). Other studies



1 have provided evidence that ammonium from remineralisation may stimulate primary production
2 (Haaber and Middelboe, 2009; Shelford et al., 2012; Weinbauer et al., 2011); yet, data are
3 lacking on the mechanism involved in the transfer of N to primary producers from bacterial
4 lysates produced by viral infection. This study shows that phytoplankton in culture and in
5 environmental samples take up N from bacterial lysates, and that ammonium produced through
6 bacterial remineralisation of DON enables that uptake.

7 **4.1 Remineralisation of nitrogen in bacterial lysates**

8 The increase in bacterial abundance in all experiments, along with the production of ammonium
9 in the laboratory experiment, establishes that bacterial lysates produced through viral infection
10 are available for bacterial growth and potential ammonium remineralisation. In the laboratory
11 experiment, every treatment with added lysate had a greater increase of bacteria than treatments
12 without lysate, indicating that the bacteria used the added lysate for growth. For all field studies,
13 bacterial abundances increased significantly in the treatments with lysate over the controls
14 ($p < 0.05$; Figure 3), demonstrating uptake of lysis products by the bacterial communities.

15 Ammonium concentration increased eightfold in laboratory treatments with lysate added to
16 bacteria (Bac+lysate; Figure 2B) from remineralisation of N in the lysate by the bacterial
17 community. In the other treatments, either lysate was not added as a source of DON (treatment
18 Bac), or *Synechococcus* sp. strain DC2 was present and consumed the ammonium that was
19 produced (the remainder of the treatments). There was no measured ammonium production in the
20 field studies except in the Fraser River Plume (FRP; Figure 3). The increase in ammonium at the
21 final time point in FRP may be due to the concurrent sharp increase in bacteria, likely increasing



1 ammonium remineralisation. Viral lysates can be used by the bacterial community for growth,
2 and excess N remineralised to produce ammonium.

3 **4.2 Phytoplankton uptake of remineralised nitrogen**

4 This study shows that remineralised N from virus lysis of bacteria can fuel the growth of primary
5 producers. Observations of increased ammonium production in the presence of viral lysis
6 (Haaber and Middelboe, 2009; Shelford et al., 2012), are extended by this study which
7 demonstrates that bacteria remineralise the organic N in lysates and produce ammonium, which
8 is then taken up by phytoplankton. The use of a cultivated bacterium for lysate production limits
9 generalisation of quantitative data to environmental systems; however, the observation that 74%
10 of the N in bacterial lysates was taken up by cultures of *Synechococcus* sp. strain DC2 provides
11 strong evidence that N from lysates may provide a large portion of the N taken up by
12 phytoplankton.

13 DON from bacterial lysates was remineralised by the bacterial assemblage into ammonium and
14 used to fuel primary production. In the laboratory experiment, the increase in ammonium in the
15 Bac+lysate treatment did not occur in the treatment with *Synechococcus* sp. strain DC2
16 (DC2+Bac+lysate; Figure 2B), indicating that the remineralised ammonium was taken up by
17 *Synechococcus* sp. strain DC2; this is consistent with the concurrent increase in *Synechococcus*
18 sp. strain DC2 cells (Figure 2A). The uptake of ^{15}N in seawater samples by organisms $>1\ \mu\text{m}$
19 (Figure 4) also demonstrates that remineralised N in bacterial lysate supports primary
20 production, as suggested previously (Shelford et al., 2012; Weinbauer et al., 2011).

21 All four field studies showed that ^{15}N from labeled bacterial lysate was taken up by the $>1\text{-}\mu\text{m}$
22 fraction (Figure 4). This is likely largely the result of uptake by phytoplankton which dominate



1 this size fraction. As well, these rates represent minimum estimates, as ^{15}N released through
2 bacterial remineralisation or phytoplankton exudation (Bronk and Ward, 2000) is not included.
3 Although the field experiments do not provide quantitative estimates of the amount of N in lysate
4 that can be taken up, the results corroborate those from the laboratory experiment. An anomalous
5 result for the SB experiment was that the estimated amount of N taken up ($0.67 \pm 0.02 \mu\text{M N}$;
6 corrected for 90 atom % ^{15}N ; Figure 4) was higher than the calculated amount of N in the lysate
7 addition ($0.44 \mu\text{M N}$; Table 2); however, the lysate contribution is an estimate calculated from
8 the cellular N quota and the reduction in cell number of *Vibrio* sp. strain PWH3a, and some error
9 may derive from these calculations and from a lack of replicates in the cell number data.
10 Nonetheless, it is reasonable to conclude that most of the lysate added in the SB and JP
11 experiments was incorporated by primary producers, since at both stations 100% of the estimated
12 ^{15}N uptake was into the $>1 \mu\text{m}$ fraction.

13 Nonetheless, not all of the N in lysates was taken up by primary producers. Although bacteria
14 can remineralise lysate (Bac+lysate; Figure 2B), and *Synechococcus* sp. strain DC2 can use
15 remineralised N from co-occurring bacteria (DC2+lysate; Figure 2A), in the laboratory only 74%
16 of the N in the lysate contributed to an increase in *Synechococcus* sp. strain DC2 abundance. As
17 well, in the FRP and SI samples, only about half of the ^{15}N uptake was into the $>1 \mu\text{m}$ fraction.
18 Discrepancies between N uptake and phytoplankton growth may in part be because
19 phytoplankton have been observed to take up almost 74% more N than the net amount
20 incorporated (Bronk et al., 1994); thus, in our experiments more N may have been taken up than
21 was measured, because of DON release. Bronk et al. (1994) also measured DON turnover times
22 in seawater samples of 4 to 18 d, much longer than in our incubations; however, the DON in
23 seawater would also include phytoplankton exudates and refractory DON that would have a



1 different composition than bacterial lysates. Hence, the incubation times in our study may not
2 have been long enough to allow for uptake of more recalcitrant DON, which might have been
3 incorporated with time (Middelboe and Jørgensen, 2006). For SI and FRP, the lysate was added
4 at a relatively high concentrations (Table 2), which may also have contributed to incomplete
5 incorporation. In contrast, the lysate was added at a much lower concentration in the SB and JP
6 experiments, and the ^{15}N in the lysate was completely recovered in the particulate fraction,
7 indicating that the DON in the lysate was incorporated when added at more natural
8 concentrations. Finally, the incomplete transfer of N from lysate of *Vibrio* sp. strain PWH3a to
9 *Synechococcus* sp. strain DC2 in the laboratory, relative to the complete uptake of ^{15}N at SB and
10 JP, may indicate that complete remineralisation of the DON depends on the make-up of the
11 bacterial assemblage, or that some phytoplankton in natural waters may take up DON directly
12 (Bronk et al., 2007).

13 The increase in *Synechococcus* sp. strain DC2 abundance in the laboratory experiment (Figure
14 2A) can be explained by the different sources of N present in the experimental treatment
15 (DC2+Bac+lysate). A portion of the increase can be attributed to the $21.3\ \mu\text{M}$ of nitrate present
16 in the Bac ultrafiltrate, a portion to the remineralisation of ammonium by the added bacterial
17 assemblage, and another portion to uptake of ammonium from remineralisation by contaminating
18 bacteria in the non-axenic *Synechococcus* sp. strain DC2 culture. Despite evidence for uptake of
19 DON directly by phytoplankton in field experiments (see Bronk et al. 2007 for a review),
20 preliminary studies to the current one showed no evidence for uptake of *Vibrio* sp. strain PWH3a
21 lysate uptake by axenic *Synechococcus* sp. strain DC2. If considered together, the increase of
22 *Synechococcus* sp. strain DC2 in the experimental treatment (DC2+Bac+lysate) can be
23 accounted for by the effects from each of these control treatments.



1 **4.3 Ecological implications and future directions**

2 Some studies have focused on the influence of phytoplankton lysates on fueling bacterial
3 production (Gobler et al., 1997; Haaber and Middelboe, 2009). Phytoplankton lysates can be
4 very important in certain situations, such as during a bloom termination (Brussaard et al., 2005),
5 but bacterial lysates are produced constantly throughout the ocean. Suttle (1994) estimated that
6 10 to 20% of heterotrophic bacteria are lysed every day by viruses, which can vary widely
7 dependent on location and conditions (e.g. Wilhelm et al. 1998, 2002). With an estimated 26-70
8 Pg C yr⁻¹ of bacterial production in the euphotic zone (Ducklow and Carlson, 1992), a bacterial
9 C:N of approximately 5 (Goldman et al., 1987), and loss rates of 10-20% from viral lysis,
10 approximately 0.52-2.8 Pg N yr⁻¹ is released from viral lysis of bacteria in the photic zone.
11 Primary production C demand is estimated to be 49.3 Pg C yr⁻¹ (Ducklow and Carlson, 1992),
12 corresponding to 7.4 Pg N yr⁻¹ according to the Redfield ratio (106C:16N). Therefore, an
13 estimated 7-38% of global primary production can be supported from bacterial lysates from viral
14 infection (Figure 5). This is a substantial source of recycled N available to fuel primary
15 production, especially in regions that are N-limited.

16 The ability of primary producers to use N from bacterial remineralisation of DOM from viral
17 lysis of bacteria indicates that viruses are not simply C sinks that disrupt trophic levels (Azam
18 and Worden, 2004), but are important facilitators in N recycling pathways. Traditional food
19 chain models state that C and other nutrients flow from primary producers to higher trophic
20 levels. The introduction of the microbial loop (Azam et al., 1983) included bacterial dynamics,
21 whereby DOM produced by the members of the traditional food chain is taken up by bacteria and
22 reintroduced to the food web instead of being lost to the system. The viral shunt (Wilhelm and



1 Suttle, 1999) introduced viruses as a ‘short-circuit,’ removing particulate C from primary
2 producers, consumers, and bacteria to the pool of organic matter. This model emphasises viruses
3 as a loss mechanism of foodweb C; however, implicit in these models is that nutrients, as well as
4 C, must also be released by viral lysis, and that N and P are likely recycled with greater
5 efficiency than C (Suttle, 2007). The loss of C from the particulate pool is clear, but there is
6 evidence of increased productivity in the presence of viruses. In Fe-limited regions of the eastern
7 Pacific Ocean, viruses were shown to liberate Fe into the DOM pool, where it was available for
8 uptake by primary producers (Poore et al., 2004). Weinbauer et al. (2011) provided evidence
9 for this phenomenon when they removed the viral fraction from a cyanobacteria bloom and
10 primary production ceased. Shelford et al. (2012) confirmed that observation by demonstrating a
11 reduction in both ammonium remineralisation and phytoplankton abundance with removal of
12 viruses. The current study demonstrated in four separate field experiments that N is transferred
13 from bacterial lysates to phytoplankton biomass. Weinbauer et al. (2011) hypothesised that
14 primary production decreased in their experiments due to either lysates providing a direct source
15 of nutrients for the *Synechococcus* blooms, or through bacterial remineralisation providing
16 inorganic nutrients. This study supports the interpretation that remobilization of nutrients by viral
17 lysis of bacteria and subsequent remineralisation by uninfected bacteria supported the observed
18 growth of *Synechococcus*.

19



1 **Author contributions**

2 E. Shelford and C. A. Suttle designed the experiments. E. Shelford performed the experiments
3 and analysed the samples except for the stable isotope data which was analysed by the Stable
4 Isotope Facility at UC Davis, and the nitrogen cell content which was analysed by Maureen Soon
5 at UBC. E. Shelford prepared the manuscript with contributions from C. A. Suttle.

6

7 **Competing Interests**

8 The authors declare that they have no conflict of interest.

9

10 **Acknowledgements**

11 We acknowledge the crew of the CGS Vector; Maureen Soon of UBC for nutrient analyses;
12 Chris Payne for analyses and cruise technician expertise; Amy Chan, Julia Gustavsen, Marli
13 Vlok, and Danielle Winget for help with experimental set-up, logistics, and discussions; Maite
14 Maldonado, Philippe Tortell, Cheryl Chow, Caroline Chénard, Julia Gustavsen, and Jan Finke
15 for their insights on the manuscript. This work was supported by awards from NSERC, UBC,
16 and the BRITE program to E. Shelford, and by an NSERC Discovery Grant to C. A. Suttle.

17



1 References

- 2 Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyer-Reil, L. A. and Thingstad, F.: The
- 3 ecological role of water-column microbes in the sea, *Mar. Ecol. Prog. Ser.*, 10, 257–263, 1983.
- 4 Azam, F. and Worden, A. Z.: Microbes, molecules, and marine ecosystems, *Science*, 303, 1622–
- 5 1624, 2004.
- 6 Berges, J. A., Franklin, D. J. and Harrison, P. J.: Evolution of an artificial seawater medium:
- 7 improvements in enriched seawater, artificial water over the last two decades, *J. Phycol.*, 37,
- 8 1138–1145, 2001.
- 9 Bronk, D. A., Glibert, P. M. and Ward, B. B.: Nitrogen uptake, dissolved organic nitrogen
- 10 release, and new production., *Science*, 265, 1843–1846, doi:10.1126/science.265.5180.1843,
- 11 1994.
- 12 Bronk, D. A., See, J. H., Bradley, P. and Killberg, L.: DON as a source of bioavailable nitrogen
- 13 for phytoplankton, *Biogeosciences*, 4, 283–296, 2007.
- 14 Bronk, D. A. and Ward, B. B.: Magnitude of dissolved organic nitrogen release relative to gross
- 15 nitrogen uptake in marine systems, *Limnol. Oceanogr.*, 45, 1879–1883, 2000.
- 16 Brussaard, C. P. D.: Optimization of procedures for counting viruses by flow cytometry, *Appl.*
- 17 *Environ. Microbiol.*, 70, 1506–1513, doi:10.1128/AEM.70.3.1506, 2004.
- 18 Brussaard, C. P. D., Kuipers, B. and Veldhuis, M. J. W.: A mesocosm study of *Phaeocystis*
- 19 *globosa* population dynamics, *Harmful Algae*, 4, 859–874, doi:10.1016/j.hal.2004.12.015, 2005.



- 1 Ducklow, H. W. and Carlson, C. A.: Oceanic bacterial production, in advances in microbial
2 ecology, vol. 12, edited by K. Marshall, pp. 113–181, Plenum Press, New York, 1992.
- 3 Edler, L. and Elbrächter, M.: The Utermöhl method for quantitative phytoplankton analysis, in
4 Microscopic and molecular methods for quantitative phytoplankton analysis, edited by B.
5 Karlson, C. Cusack, and E. Bresnan, pp. 13–20, UNESCO, Paris, 2010.
- 6 Fuhrman, J. A.: Marine viruses and their biogeochemical and ecological effects, *Nature*, 399,
7 541–548, 1999.
- 8 Gasol, J. M. and Del Giorgio, P. A.: Using flow cytometry for counting natural planktonic
9 bacteria and understanding the structure of planktonic bacterial communities, *Sci. Mar.*, 64, 197-
10 224, 2000.
- 11 Gobler, C. J., Hutchins, D. A., Fisher, N. S., Coper, E. M. and Sañudo-Wilhelmy, S.: Release
12 and bioavailability of C, N, P, Se, and Fe following viral lysis of a marine chrysophyte, *Limnol.*
13 *Oceanogr.*, 42, 1492–1504, 1997.
- 14 Goldman, J. C., Caron, D. A. and Dennett, M. R.: Regulation of gross growth efficiency and
15 ammonium regeneration in bacteria by substrate C:N ratio, *Limnol. Oceanogr.*, 32, 1239–1252,
16 1987.
- 17 Haaber, J. and Middelboe, M.: Viral lysis of *Phaeocystis pouchetii*: implications for algal
18 population dynamics and heterotrophic C, N and P cycling, *ISME J.*, 3, 430–441,
19 doi:10.1038/ismej.2008.125, 2009.



- 1 Healey, F. P. and Hendzel, L. L.: Indicators of phosphorus and nitrogen deficiency in five algae
2 in culture, *J. Fish. Res. Board Canada*, 36, 1364–1369, 1979.
- 3 Hollibaugh, J. T.: Nitrogen regeneration during the degradation of several amino acids by
4 plankton communities collected near Halifax, Nova Scotia, *Mar. Biol.*, 45, 191–201, 1978.
- 5 Holmes, R. M., Aminot, A., Kerouel, R., Hooker, B. A. and Peterso, B. J.: A simple and precise
6 method for measuring ammonium and marine and freshwater ecosystems, *Can. J. Fish. Aquat.
7 Sci.*, 56, 1801–1808, 1999.
- 8 Middelboe, M. and Jørgensen, N. O. G.: Viral lysis of bacteria: an important source of dissolved
9 amino acids and cell wall compounds, *J. Mar. Biol. Assoc. UK*, 86, 605–612, 2006.
- 10 Middelboe, M., Jørgensen, N. O. G. and Kroer, N.: Effects of viruses on nutrient turnover and
11 growth efficiency of noninfected marine bacterioplankton, *Appl. Environ. Microbiol.*, 62, 1991–
12 1997, 1996.
- 13 Middelboe, M., Riemann, L., Steward, G., Hansen, V. and Nybroe, O.: Virus-induced transfer of
14 organic carbon between marine bacteria in a model community, *Aquat. Microb. Ecol.*, 33, 1–10,
15 2003.
- 16 Olson, R. J., Zettler, E. R. and D, D. M.: Phytoplankton analysis using flow cytometry, in
17 handbook of methods in aquatic microbial ecology, pp. 175–186, Lewis Publishers, Boca Raton.,
18 1993.
- 19 Parsons, T. R., Maita, Y. and Lalli, C. M.: Manual of chemical and biological methods for
20 seawater analysis, Pergamon Press., 1984.



- 1 Poorvin, L., Rinta-Kanto, J. M., Hutchins, D. A. and Wilhelm, S. W.: Viral release of iron and its
2 bioavailability to marine plankton, *Limnol. Oceanogr.*, 49, 1734–1741, 2004.
- 3 Poorvin L., Sander S. G., Velasquez I., Ibisanni E., LeClerc G. R., Wilhelm S.W.: A comparison
4 of Fe bioavailability and binding of a catechol siderophore with virus-mediated lysates from
5 the marine bacterium *Vibrio alginolyticus* PWH3a. *J. Exp. Mar. Biol. Ecol.*, 399, 43-47, 2011.
- 6 Shelford, E. J., Middelboe, M., Møller, E. F. and Suttle, C. A.: Virus-driven nitrogen cycling
7 enhances phytoplankton growth, *Aquat. Microb. Ecol.*, 66, 41–46, doi:10.3354/ame01553, 2012.
- 8 Suttle, C. A.: The significance of viruses to mortality in aquatic microbial communities, *Microb.*
9 *Ecol.*, 28, 237–243, doi:10.1007/BF00166813, 1994.
- 10 Suttle, C. A.: Viruses in the sea., *Nature*, 437, 356–361, doi:10.1038/nature04160, 2005.
- 11 Suttle, C. A.: Marine viruses--major players in the global ecosystem., *Nat. Rev. Microbiol.*, 5,
12 801–812, doi:10.1038/nrmicro1750, 2007.
- 13 Suttle, C. A., Chan, A. M., and Cottrell, M. T. Use of ultrafiltration to isolate viruses from
14 seawater which are pathogens of marine phytoplankton. *Appl. Environ. Microbiol.* 57, 721-726,
15 1991.
- 16 Suttle, C. A. and Chen, F.: Mechanisms and rates of decay of marine viruses in seawater., *Appl.*
17 *Environ. Microbiol.*, 58, 3721–3729, 1992.
- 18 Weinbauer, M. G., Bonilla-Findji, O., Chan, A. M., Dolan, J. R., Short, S. M., Šimek, K.,
19 Wilhelm, S. W. and Suttle, C. A.: *Synechococcus* growth in the ocean may depend on the lysis of
20 heterotrophic bacteria, *J. Plankton Res.*, 33, 1465–1476, doi:10.1093/plankt/fbr041, 2011.



- 1 Weinbauer, M. G., Wilhelm, S. W., Suttle, C. A. and Garza, D. R.: Photoreactivation
2 compensates for UV damage and restores infectivity to natural marine virus communities., Appl.
3 Environ. Microbiol., 63, 2200–2205, 1997.
- 4 Wilhelm, S. W., Brigden, S. M. and Suttle, C. A.: A dilution technique for the direct
5 measurement of viral production: a comparison in stratified and tidally mixed coastal waters.,
6 Microb. Ecol., 43, 168–173, doi:10.1007/s00248-001-1021-9, 2002.
- 7 Wilhelm, S. W. and Suttle, C. A.: Viruses and nutrient cycles in the sea, Bioscience, 49, 781–
8 788, doi:10.2307/1313569, 1999.
- 9 Wilhelm, S. W., Weinbauer, M. G., Suttle, C. A. and Jeffrey, W. H.: The role of sunlight in the
10 removal and repair of viruses in the sea, Limnol. Oceanogr., 43, 586–592, 1998.
- 11
- 12
- 13



- 1 Table 1 Locations and environmental parameters for field sampling stations: Saanich Inlet (SI),
 2 Fraser River Plume (FRP, Semiahoo Bay (SB), and Jericho Pier (JP). Environmental and
 3 biological conditions measured include temperature (Temp), salinity (Sal), nitrate (NO_3^-),
 4 phosphate (PO_4^{2-}), ammonium (NH_4^+), and the dominant phytoplankton.

Station	Location	Date	Temp (°C)	Sal	NO_3^- (μM)	PO_4^{2-} (μM)	NH_4^+ (μM)	Dominant phytoplankton
SI	48.592°N, -123.505°W	13 Sep 2012	14	27.9	1.24	0.75	0.54	Mixed assemblage
FRP	49.072°N, -123.402°W	13 Sep 2012	11	28.0	10.9	1.12	0.07	<i>Phaeocystis</i> , <i>Skeletonema</i> , <i>Leptocylindrus</i>
SB	49.013°N, -123.037°W	26 Aug 2013	20.2	29	0.00	2.40	0.08	Cyanobacteria Unknown ciliate
JP	49.277°N, -123.202°W	27 Aug 2013	19.5	21	0.00	0.20	0.11	Diatoms (mixed assemblage)

5



- 1 Table 2 *Vibrio* sp. strain PWH3a dynamics during the creation of lysates by viral infection
 2 (Laboratory = experiments with cultured *Synechococcus* sp. strain DC2, SI = Saanich Inlet, FRP
 3 = Fraser River Plume, SB = Semiahoo Bay, JP = Jericho Pier).

	Laboratory	SI and FRP	SB and JP
<i>Vibrio</i> sp. strain PWH3a abundance pre-virus (cell l ⁻¹)	3.96×10^{11}	7.87×10^{11}	7.89×10^{10}
<i>Vibrio</i> sp. strain PWH3a abundance post-virus incubation (cell l ⁻¹)	1.12×10^{11}	9.43×10^{10}	6.38×10^{10}
Multiplicity of infection (MOI)	1	0.41	5
Total N release (μM)	721	1760	38.4
N addition to experimental incubations (μM)	36.0	67.7	0.44

4

5



- 1 Table 3 Uptake by the particulate fraction of ^{15}N from bacterial lysate from four field stations,
2 and the same uptake normalised by initial chlorophyll *a* concentrations. Error measurements are
3 standard error of triplicate incubations for SB, and range of duplicate incubations for JP. SI and
4 FRP are singleton measurements.


	Total uptake of ^{15}N (μM)	Uptake of ^{15}N normalised by chlorophyll <i>a</i> ($\mu\text{mol } \mu\text{g}^{-1}$)
Saanich Inlet (SI)	1.67	0.27
Fraser River Plume (FRP)	1.76	0.70
Semiahoo Bay (SB)	0.60 ± 0.02	0.70 ± 0.16
Jericho Pier (JP)	0.50 ± 0.1	0.09 ± 0.00

5

6



1



2

3

4

5

6


7

8


9

Figure 1 Diagram of treatments in the laboratory experiment examining increase in *Synechococcus* sp. strain DC2 abundance, in the presence of bacterial lysate from *Vibrio* sp. strain PWH3a (Lys) and an environmental bacterial assemblage (Bac). Lys, while a component of some of the treatments, was not a treatment by itself (indicated by parentheses). The * is an indication that DC2 is not axenic, and therefore contains some heterotrophic bacteria.


9



1



2



3

4 Figure 2 Time series of (A) *Synechococcus* sp. strain DC2 abundance, (B) ammonium
5 concentration, and (C) bacterial abundance in the laboratory experiment. Error bars are standard
6 error from triplicate incubations.

7




Figure 3 Time-course of ammonium concentrations and bacterial abundances. Solid and dashed lines indicate treatments with lysate addition (SW+lysate), or without (SW), respectively. Error bars are standard errors of triplicate incubations. Asterisks indicate significant differences between treatments. Note that the scales are different among plots.

1

2

3


4

5

6



1



2

3

4

5

6


7

8

9

Figure 4 Uptake of ¹⁵N normalised by initial chlorophyll *a* concentrations during the experiment by the >1 µm and <1 µm fractions in treatments with lysate addition for Saanich Inlet (SI), Fraser River Plume (FRP), Semiahoo Bay (SB), and Jericho Pier (JP). Error bars are standard error of triplicate incubations for SB, and range of duplicate incubations for JP. SI and FRP are singleton measurements.

9



2 Figure 5 Viruses release $0.52-2.8 \text{ Pg N yr}^{-1}$ globally in the euphotic ocean from lysis of
3 heterotrophic prokaryotes. That dissolved organic N (DON) is available for remineralisation by
4 other prokaryotes into ammonium, which constitutes a large proportion of the total N demand of
5 primary producers.