

1 **Benthic Archaea as potential sources of tetraether membrane**
2 **lipids in sediments across an oxygen minimum zone**

3 Marc A. Besseling^{1*}, Ellen C. Hopmans¹, R. Christine Boschman¹, Jaap S. Sinninghe Damsté^{1,2}, and
4 Laura Villanueva¹

5 ¹NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry, and
6 Utrecht University. P.O. Box 59, NL-1790 AB Den Burg, The Netherlands.

7 ²Utrecht University, Faculty of Geosciences, Department of Earth sciences, P.O. Box 80.021, 3508 TA Utrecht, The
8 Netherlands

9 *Correspondence to:* Marc A. Besseling (marc.besseling@nioz.nl)

10 **Abstract.** Benthic Archaea comprise a significant part of the total prokaryotic biomass in marine sediments. Recent
11 genomic surveys suggest they are largely involved in anaerobic processing of organic matter but the distribution and
12 abundance of these archaeal groups is still largely unknown. Archaeal membrane lipids composed of isoprenoid
13 diethers or tetraethers (glycerol dibiphytanyl glycerol tetraether, GDGT) are often used as archaeal biomarkers. Here,
14 we compare the archaeal diversity and intact polar lipid (IPL) composition in both surface (0–0.5 cm) and subsurface
15 (10–12 cm) sediments recovered within, just below, and well below the oxygen minimum zone (OMZ) of the Arabian
16 Sea. Archaeal 16S rRNA gene amplicon sequencing revealed a predominance of Thaumarchaeota (Marine Group I,
17 MG-I) in oxygenated sediments. Quantification of archaeal 16S rRNA and ammonia monooxygenase (*amoA*) of
18 Thaumarchaeota genes and their transcripts indicated the presence of an active *in situ* benthic population, which
19 coincided with a high relative abundance of hexose phosphohexose crenarchaeol, a specific biomarker for living
20 Thaumarchaeota. On the other hand, anoxic surface sediments within the OMZ and all subsurface sediments were
21 dominated by archaea belonging to the Miscellaneous Crenarchaeota Group (MCG), the Thermoplasmatales and
22 archaea of the DPANN superphylum. Members of the MCG were diverse with a dominance of subgroup MCG-12 in
23 anoxic surface sediments. This coincided with a high relative abundance of IPL GDGT-0 with an unknown polar head
24 group. Subsurface anoxic sediments were characterized by higher relative abundance of GDGT-0, 2 and 3 with
25 dihexose IPL-types, as well as GDGT-0 with a cyclopentanetetraol molecule and a hexose, as well as the presence of
26 specific MCG subgroups, suggesting that these groups could be the biological sources of these archaeal lipids.

27 INTRODUCTION

28 Archaea are ubiquitous microorganisms in the marine system (DeLong et al., 1994; Delong and Pace, 2001; Schleper et
29 al., 2005). They occur in diverse environments, e.g. hydrothermal vents (Stetter et al., 1990), the marine water column
30 (Karner et al., 2001; Massana et al., 2004), in the underlying sediments (Lloyd et al., 2013; Teske and Sørensen, 2008),
31 and well below the seafloor (Biddle et al., 2006; Lipp et al., 2008), where they are considered key players in diverse
32 biogeochemical processes (Offre et al., 2013, and references cited therein). Specifically marine sediments have been
33 shown to contain a highly diverse archaeal community (Lloyd et al., 2013; Spang et al., 2017; Teske, 2013; Teske and
34 Sørensen, 2008). The ammonia-oxidizing Thaumarchaeota of the marine group I.1a (further referred to as MG-I) is
35 probably the most widely studied archaeal group in marine sediments. However, in comparison with studies of marine
36 pelagic Thaumarchaeota, the diversity and distribution of benthic Thaumarchaeota is still not well established (e.g.
37 Durbin & Teske, 2010; Jorgenson et al., 2012; Learman et al., 2016). Genomic studies have revealed the existence of
38 uncultured archaeal groups other than Thaumarchaeota in marine, predominantly anoxic, sediments such as the
39 Miscellaneous Crenarchaeota Group (MCG; Meng et al., 2014), archaea of the DPANN superphylum (composed of
40 Micrarchaeota, Diapherotrites, Aenigmarchaeota, Nanohaloarchaeota, Parvarchaeota, Nanoarchaeota, Pacearchaeota
41 and Woearchaeota; Castelle et al., 2015; Rinke et al., 2013) and the Marine Benthic Group (MBG) B (Teske &
42 Sørensen, 2008), and D (Lloyd et al., 2013). In the case of the archaea belonging to the groups of the MCG and MBG-
43 D, metagenomic studies suggest that they are able to degrade extracellular proteins and aromatic compounds (Lloyd et
44 al., 2013; Meng et al., 2014).

45 Archaeal diversity is currently determined through nucleic acid-based methods but the characterization of other cellular
46 biomarkers such as membrane lipids has proven to be also effective in tracking the presence of archaeal groups in
47 different ecosystems (e.g. Coolen et al., 2004a; Ingalls et al., 2012; Meador et al., 2015; Pitcher et al., 2011b; Sturt et
48 al., 2004). One of the advantages of using lipid-based methods to determine the presence of archaeal groups is that
49 lipids can be preserved in the sedimentary record. Therefore, they can also be used as biomarkers of the presence and
50 metabolic potential of these microorganisms in past environments. On the contrary, other biomolecules like DNA have
51 a more rapid turnover and they cannot be used for this purpose. In recent years, intact polar lipids (IPLs) have
52 increasingly been applied for tracing 'living' bacteria and archaea in the environment (Lipp et al., 2008; Lipp and
53 Hinrichs, 2009; Rossel et al., 2008). IPLs with polar head groups are present in living cells but upon cell lysis the polar
54 head groups are lost, releasing core lipids (CLs) that may be preserved in the fossil record. Since IPLs degrade
55 relatively quickly after cell death (Harvey et al., 1986), it is possible to associate the presence of IPLs in the
56 environment with the occurrence of their living producers (Lipp and Hinrichs, 2009; Schubotz et al., 2009).

57 Archaeal membrane lipids are typically a variation of two main structures, *sn*-2,3-diphytanylglycerol diether (archaeol)
58 with phytanyl (C₂₀) chains in a bilayer structure, and *sn*-2,3-dibiphytanyl diglycerol tetraether (glycerol dibiphytanyl
59 glycerol tetraether, GDGT), in which the two glycerol moieties are connected by two C₄₀ isoprenoid chains, allowing

60 the formation of a monolayer membrane (Koga and Morii, 2007). GDGTs containing 0–4 cyclopentane moieties (Fig.
61 S1) are usually not exclusive to a specific archaeal group (Schouten et al., 2013) with the exception of the GDGT
62 crenarchaeol, containing 4 cyclopentane and one cyclohexane moiety, which is deemed to be exclusive to the
63 Thaumarchaeota phylum (De La Torre et al., 2008; Sinninghe Damsté et al., 2002, 2012). Recently, Lincoln et al.
64 (2014) proposed the Marine Group II as potential producers of crenarchaeol. However, this is still debated (Lincoln et
65 al., 2014b; Schouten et al., 2014). The newly described archaeal groups detected by genetic methods are yet uncultured,
66 therefore, their membrane lipid composition remains unknown.

67 In this study, we determined the archaeal diversity in a marine benthic system along a strong gradient in bottom water
68 oxygen concentrations and compared it with the diversity of archaeal lipids. We aimed to characterize changes in the
69 archaeal benthic community under different physicochemical conditions, as well as to provide clues on the potential
70 archaeal lipid biomarkers produced by uncultured benthic archaea. We analyzed sediments (surface 0–0.5 cm, and
71 subsurface 10–12 cm) of the Murray ridge in the Arabian Sea, which is impinged by one of the strongest present-day
72 oxygen minimum zones (OMZ). Previous studies observed changes in the diversity of archaeal lipids in the same
73 environmental setting in sediments under different oxygen and nutrient concentrations (Lengger et al., 2012; 2014). In
74 our study, we expand the repertoire of archaeal lipid diversity previously detected by Lengger et al. (2012; 2014) by
75 analyzing these sediments with High Resolution Accurate Mass/Mass spectrometry (UHPLC-HRAM MS). In addition,
76 we determined the archaeal diversity by means of 16S rRNA gene amplicon sequencing, as well as the abundance and
77 potential activity of specific archaeal groups by quantitative PCR (QPCR) of 16S rRNA and the metabolic gene coding
78 for the ammonia monooxygenase (*amoA* gene) of Thaumarchaeota.

79 **MATERIAL and METHODS**

80 **Sampling**

81 Sediments were collected in the Northern Arabian Sea during the PASOM cruise in January 2009 with *R/V Pelagia*.
82 Sediment cores obtained with a multicorer were taken on the Murray ridge at four depths, 885 m below sea level (mbsl)
83 (within the OMZ), at 1306 mbsl (just below the OMZ), at 2470 mbsl and 3003 mbsl (both well below the OMZ) as
84 previously described by Lengger et al. (2012). Upon retrieval the cores were sliced in 0.5 cm resolution for the first 2
85 cm and at 2 cm resolution beyond 10 cm below the surface, and stored at -80°C until further analysis. For an overview
86 of the surface sediments physicochemical conditions see Table 1.

87 **Lipid extraction and analysis**

88 Total lipids were extracted from surface (upper 0–0.5 cm) and subsurface (10–12 cm) sediments after freeze-drying
89 using a modified Bligh and Dyer method (Bligh and Dyer, 1959) as previously described by Lengger et al. (2014). C₁₆-
90 PAF (1-O-hexadecyl-2-acetyl-sn-glycero-3-phosphocholine) was added to the extracts as an internal standard and the

91 extracts were dried under a stream of nitrogen. The extracts with the added standard were then dissolved by adding
92 solvent (hexane:isopropanol:H₂O 718:271:10 [v/v/v/v]) and filtered through a 0.45 µm, 4 mm-diameter True
93 Regenerated Cellulose syringe filter (Grace Davison, Columbia, MD, USA).

94 IPLs were analyzed according to Sturt et al. (2004) with some modifications. An Ultimate 3000 RS UHPLC, equipped
95 with thermostated auto-injector and column oven, coupled to a Q Exactive Orbitrap MS with Ion Max source with
96 heated electrospray ionization (HESI) probe (Thermo Fisher Scientific, Waltham, MA), was used. Separation was
97 achieved on a YMC-Triart Diol-HILIC column (250 x 2.0 mm, 1.9 µm particles, pore size 12 nm; YMC Co., Ltd,
98 Kyoto, Japan) maintained at 30 °C. The following elution program was used with a flow rate of 0.2 mL min⁻¹: 100% A
99 for 5 min, followed by a linear gradient to 66% A: 34% B in 20 min, maintained for 15 min, followed by a linear
100 gradient to 40% A: 60% B in 15 min, followed by a linear gradient to 30% A: 70% B in 10 min, where A = hexane/2-
101 propanol/formic acid/14.8 M NH_{3aq} (79:20:0.12:0.04 [v/v/v/v]) and B = 2-propanol/water/formic acid/ 14.8 M NH_{3aq}
102 (88:10:0.12:0.04 [v/v/v/v]). Total run time was 70 min with a re-equilibration period of 20 min in between runs. HESI
103 settings were as follows: sheath gas (N₂) pressure 35 (arbitrary units), auxiliary gas (N₂) pressure 10 (arbitrary units),
104 auxiliary gas (N₂) T 50 °C, sweep gas (N₂) pressure 10 (arbitrary units), spray voltage 4.0 kV (positive ion ESI),
105 capillary temperature 275 °C, S-Lens 70 V. IPLs were analyzed with a mass range of *m/z* 375 to 2000 (resolving power
106 70,000 at *m/z* 200), followed by data dependent MS² (resolving power 17,500 ppm at *m/z* 200), in which the ten most
107 abundant masses in the mass spectrum (with the exclusion of isotope peaks) were fragmented (stepped normalized
108 collision energy 15, 22.5, 30; isolation window 1.0 *m/z*). A dynamic exclusion window of 6 sec was used as well as an
109 inclusion list with a mass tolerance of 3 ppm to target specific compounds (Table S1). The Q Exactive Orbitrap MS was
110 calibrated within a mass accuracy range of 1 using the Thermo Scientific Pierce LTQ Velos ESI Positive Ion
111 Calibration Solution (containing a mixture of caffeine, MRFA, Ultramark 1621, and *N*-butylamine in an acetonitrile-
112 methanol-acetic acid solution).

113 Peak areas for each individual IPL were determined by integrating the combined mass chromatogram (within 3 ppm) of
114 the monoisotopic and first isotope peak of all relevant adducts formed (protonated, ammoniated and/or sodiated adducts
115 may be formed in different proportions depending on the type of IPL). PAF was used as internal standard to
116 continuously monitor MS performance and to assess matrix effects. Reported peak areas have been corrected for these
117 effects. Absolute quantification of IPL GDGTs was not possible due to a lack of standards. Peak areas were not
118 corrected for any possible differences in response factors between the various classes of IPL-crenarchaeol. IPLs with
119 the same headgroup but with the regioisomer of crenarchaeol instead of crenarchaeol as the CL co-elute on the
120 chromatographic system used here and any peak area reported for a crenarchaeol IPL thus represents the sum of both
121 isomers.

122 To rule out any degradation of the GDGT-IPLs during storage of the sediments at -20°C, the anoxic surface sediment
123 sample at 885 mbsl was also analyzed according to the method previously used by Lengger et al. (2012). The IPL

124 fraction was separated from the CLs with the use of a silica column and elution with MeOH (Lengger et al., 2012). This
125 IPL fraction was hydrolyzed for 3 h and analyzed by HPLC-APCI/MS (according to Hopmans et al., 2016) and the IPL
126 derived CL-GDGT distribution was compared with previously published data.

127 **Nucleic acids extraction, cDNA synthesis and quantitative PCR (QPCR) analyses**

128 Sediment was centrifuged and the excess of water was removed by pipetting before proceeding with the extraction of
129 nucleic acids from the sediment. DNA/RNA of surface (0–0.5 cm) and subsurface (10–12 cm) sediments was extracted
130 with the RNA PowerSoil® Total Isolation Kit plus the DNA elution accessory (Mo Bio Laboratories, Carlsbad, CA).
131 Concentration of DNA and RNA were quantified by Nanodrop (Thermo Scientific, Waltham, MA) and Fluorometric
132 with Quant-iT™ PicoGreen® dsDNA Assay Kit (Life technologies, Netherlands). RNA extracts were treated with
133 DNase and reverse-transcribed to cDNA as described by Pitcher et al. (2011). Quantification of archaeal 16S rRNA
134 gene copies and *amoA* gene copies were estimated by QPCR by using the following primers; Parch519F and ARC915R
135 (archaeal 16S rRNA gene), CrenAmoAQ-F and CrenAmoAModR (*amoA* gene), as previously described (Pitcher et al.,
136 2011). For details on the QPCR conditions, efficiency and R² of the QPCR assays see Table S2.

137 **16S rRNA gene amplicon sequencing, analysis, and phylogeny**

138 PCR reactions were performed with the universal, Bacteria and Archaea, primers S-D-Arch-0159-a-S-15 and S-D-Bact-
139 785-a-A-21 (Klindworth et al., 2013) as previously described in Moore et al. (2015). The archaeal 16S rRNA gene
140 amplicon sequences were analyzed by QIIME v1.9 (Caporaso et al., 2010). Raw sequences were demultiplexed and
141 then quality-filtered with a minimum quality score of 25, length between 250–350, and allowing maximum two errors
142 in the barcode sequence. Taxonomy was assigned based on blast and the SILVA database version 123 (Altschul et al.,
143 1990; Quast et al., 2013). Representative operational taxonomic units (OTUs, clusters of reads with 97% similarity) of
144 archaeal groups were extracted through filter_taxa_from_otu_table.py and filter_fasta.py with QIIME (Caporaso et al.,
145 2010). The phylogenetic affiliation of the partial archaeal 16S rRNA gene sequences was compared to release 123 of
146 the Silva NR SSU Ref database (<http://www.arb-silva.de/>; Quast et al., 2013) using the ARB software package (Ludwig
147 et al., 2004). Sequences were added to the reference tree supplied by the Silva database using the ARB Parsimony tool.
148 MCG intragroup phylogeny for representative sequences of OTUs affiliated to the MCG lineage was carried out in
149 ARB (Ludwig et al., 2004). Sequences were added by parsimony to a previously-built phylogenetic tree composed of
150 reference sequences of the 17 MCG subgroups known so far (Kubo et al., 2012). Affiliation of any 16S rRNA gene
151 sequences to a given subgroup was done assuming a similarity cutoff of $\geq 85\%$.

152 **Cloning, sequencing and phylogeny of the archaeal *amoA* gene**

153 Amplification of the archaeal *amoA* gene was performed as described by Yakimov et al., (2011). PCR reaction mixture
154 was the following (final concentration): Q-solution 1× (PCR additive, Qiagen); PCR buffer 1×; BSA (200 $\mu\text{g ml}^{-1}$);
155 dNTPs (20 μM); primers (0.2 $\text{pmol } \mu\text{l}^{-1}$); MgCl_2 (1.5 mM); 1.25 U Taq polymerase (Qiagen, Valencia, CA, USA). PCR

156 conditions for these amplifications were the following: 95°C, 5 min; 35 × [95°C, 1 min; 55°C, 1 min; 72°C, 1 min];
157 final extension 72°C, 5 min. PCR products were gel purified (QIAquick gel purification kit, Qiagen) and cloned in the
158 TOPO-TA cloning® kit from Invitrogen (Carlsbad, CA, USA) and transformed in *E. coli* TOP10 cells following the
159 manufacturer's recommendations. Recombinant clones plasmid DNAs were purified by Qiagen Miniprep kit and
160 screening by sequencing (n ≥ 30) using M13R primer by Macrogen Europe Inc. (Amsterdam, The Netherlands).
161 Obtained archaeal *amoA* protein sequences were aligned with already annotated *amoA* sequences by using the Muscle
162 application (Edgar, 2004). Phylogenetic trees were constructed with the Neighbor-Joining method (Saitou and Nei,
163 1987) and evolutionary distances computed using the Poisson correction method with a bootstrap test of 1,000
164 replicates.

165 **RESULTS**

166 In this study, we analyzed both IPLs and DNA/RNA extracts from sediments previously collected along the Arabian
167 Sea Murray Ridge within the OMZ (885 mbsl), just below the lower interface (1306 mbsl), and well below the OMZ
168 (2470 and 3003 mbsl). The surface sediment (0-0.5 cm) at 885 mbsl was fully anoxic, however, the surface sediments
169 below the OMZ were partly oxygenated (1306 mbsl), and fully oxygenated at 2470 and 3003 mbsl (Table 1). The
170 subsurface sediments (10-12 cm) were fully anoxic at all stations (Table 1). For more details on the physicochemical
171 conditions in these sediments see Table 1.

172 **Archaeal IPL-GDGTs in the surface and subsurface sediments**

173 A range of IPL-GDGTs (GDGT-0 to 4 and crenarchaeol) with the IPL-types monohexose (MH), dihexose (DH) and
174 hexose-phosphohexose (HPH) was detected in surface and subsurface sediments across the Arabian Sea OMZ (Table
175 2). For the DH GDGT-0 two structural isomers (type-I with two hexose moieties at both ends of the CL, and type-II
176 with one dihexose moiety; Table 2) were detected and identified based on their mass spectral characteristics (Fig. S2).
177 These isomers were previously also reported in thaumarchaeotal cultures (Elling et al., 2014, 2017). In addition,
178 GDGT-0 with both an ether-bound cyclopentanetetraol moiety and a hexose moiety as head groups was identified (Fig.
179 S2) in some sediments (Table 2). This IPL was previously reported as a glycerol dibiphytanyl nonitol tetraether
180 (GDNT; de Rosa et al. 1983) but was later shown to contain a 2-hydroxymethyl-1-(2,3-dihydroxypropoxy)-2,3,4,5-
181 cyclopentanetetraol moiety by Sugai et al., (1995) on the basis of NMR spectroscopy characterization.

182 In the surface sediment at 885 mbsl, crenarchaeol IPLs were dominant (44.7% of all detected IPL-GDGTs), occurring
183 predominantly with DH as IPL-type (with a hexose head group on both ends; 43.1%; Table 2). IPL-GDGT-2 was the
184 second most abundant (29.6%), also mainly consisting of the IPL-type DH (29.5%; Table 2). IPL-GDGT-0, -1, -3 and -
185 4 were occurring with relative abundances of 0.3%, 1.7%, 17.8% and 6.1%, respectively (Table 2). Overall, the

186 majority (98.1%; Table 3) of IPL-GDGTs in surface sediment at 885 mbsl with IPL-type DH (all with a hexose
187 molecule on both ends of the CL).

188 The surface sediment at 1306 mbsl contained mostly IPL-GDGT-0 (37.6% of all detected IPL-GDGTs), almost entirely
189 with the IPL-type HPH (36.6% of the total; Table 2). Slightly less abundant was the IPL-crenarchaeol (35.6%), with the
190 IPL-types HPH (18.7%) and DH type-I (15.5%) in equal amounts and with a minor relative abundance with MH
191 (1.4%). Overall, the IPL-GDGTs in surface sediment at 1306 mbsl mainly contained the IPL-types HPH (55.4%; Table
192 3) and DH (42.0%; Table 3).

193 Well below the OMZ, surface sediments from 2470 and 3003 mbsl were both dominated by IPL-GDGT-0 (71.9 and
194 80.8% of all detected IPL-GDGTs, respectively), predominantly with IPL-type HPH (Table 2; Fig. 1a). The IPL-
195 crenarchaeol had a lower relative abundance (26.6 and 17.6%, respectively) and again was dominated by the member
196 with IPL-type HPH (Table 2). The other IPL-GDGTs occurred in minor quantities (<1%). Overall, IPL-type HPH was,
197 thus, by far the most abundant head group detected in surface sediments at 2470 and 3003 mbsl (97.7% and 97.4%,
198 respectively), in contrast to the other two surface sediments studied (Table 3).

199 In all subsurface (10-12 cm) sediments (i.e. at 885, 1306, 2470 and 3003 mbsl) the most abundant IPL-GDGTs were
200 DH-crenarchaeol ($28.9\pm 3.8\%$; Table 2) and DH-GDGT-2 ($25.5\pm 3.5\%$; Table 2). DH was also the most commonly
201 observed IPL-type attached to GDGT-3 and GDGT-4 (Table 2). Overall the distributions of the IPL-GDGTs in all
202 subsurface sediments were relatively similar (Fig. 1a) in comparison to the substantial changes observed at the surface
203 (cf. Fig. 1a). Overall, the IPL-type DH was the predominant one detected in subsurface sediment with a relative
204 abundance ranging from 68.8% at 3003 mbsl to 92.9% at 885 mbsl (Table 3). In contrast to all other sediments, in the
205 subsurface sediments at 885 mbsl and 1306 mbsl, two different isomers (Fig. S2) of the DH-GDGT-0 were detected
206 (Table 2). DH type-I (0.9% at 1306 mbsl) is also found in the other surface and subsurface sediments and in
207 combination with other core GDGT structures, whereas the other isomer (DH type-II) only occurs (7.8% at 885 mbsl;
208 1.8% at 1306 mbsl; Table 2; Fig. S2b). In addition, these subsurface sediments also contain small amounts of GDGT-0
209 with cyclopentanetetraol and MH head groups (IPL-type HCP; 1.6% at 885 mbsl; 0.4% at 1306 mbsl; Table 2; Fig.
210 S2c).

211 We also determined the IPL-derived CL-GDGTs in the 885 mbsl surface sediment following the method of Lengger et
212 al. (2012), in order to exclude IPL degradation within the stored samples. The CL-GDGTs composition derived from
213 freshly obtained IPL showed the same distribution ($r=0.99$, $p < 0.001$) as reported previously (Lengger et al., 2012).

214

215 **Archaeal diversity in the surface and subsurface sediment**

216 Different archaeal groups were detected in surface and subsurface sediment across the Arabian sea OMZ. The surface
217 sediment at 885 mbsl, contained archaeal 16S rRNA gene sequences that were assigned to several archaeal groups (Fig.
218 1b). The most dominant group was MCG (Total 30.5%, 12.2% attributed to C3; also known as MCG-15, Kubo et al.,

219 2012). Another major group found was the DPANN Woesearchaeota Deep sea Hydrothermal Vent Group 6 (DHVEG-
220 6, 20.3%; Fig. 1b; Castelle et al., 2015). Marine Benthic Group (MBG) -B, -D and -E were also present with 12.2%,
221 7.7% and 6.9% of the archaeal 16S rRNA gene reads, respectively (Fig. 1b). Sequences affiliated to the Marine
222 Hydrothermal Vent Group (MHVG, 8.1%) of the phylum Euryarchaeota were also detected (Fig. 1b). Other groups,
223 with lower relative abundances, were Thermoplasmatales groups ANT06-05 (5.7%) and F2apm1A36 (3.3%) and the
224 DPANN Aenigmarchaeota (previously named Deep Sea Euryarchaeotic Group, DSEG; 1.6%; Fig. 1b).

225 Below the OMZ, in partly and fully oxygenated surface sediments at 1306, 2470 and 3003 mbsl (Table 1), the most
226 dominant archaeal group was Thaumarchaeota MG-I with relative abundances of 81.5%, 89.7% and 100%, respectively
227 (Fig. 1b). At 1306 mbsl other archaeal groups, such as MHVG (5.6%), Thermoplasmatales ASC21 (3.2%), DHVEG-6
228 (2.9%), MBG-B (2.4%) and MCG (1.3%) made up the rest of the archaeal community (Fig. 1b). At 2470 mbsl
229 DHVEG-6 (1.1%) was still detectable besides the MG-I (Fig. 1b).

230 In the subsurface sediments (10–12 cm), only the DNA extracted from the sediments at 885 and 1306 mbsl gave a
231 positive amplification signal. The archaeal composition of the subsurface (10–12 cm) sediments at 885 mbsl and 1306
232 mbsl was similar (Fig. 1b; Pearson correlation coefficient of 0.95), with most of the 16S rRNA gene reads classified
233 within the MCG (47.5% and 48.4%, respectively). Other archaeal groups, such as MBG-D (14.4% and 5.7%,
234 respectively), MBG-B (10.1% and 4.4%), the Woesearchaeota (7.8% and 10.4%), were also detected with comparable
235 relative abundances (Fig. 1b). Other archaeal groups such as Thaumarchaeota Terrestrial hot spring, the Euryarchaeota
236 MHVG, MBG-E and the Aenigmarchaeota were detected but at low (< 10%) relative abundance (Fig. 1b). Only minor
237 amount of reads were classified as Thaumarchaeota MG-I (0.5% at 1306 mbsl) (Fig. 1b).

238 Considering the high relative abundance of the MCG detected in the surface sediment at 885 mbsl, as well as in the
239 subsurface (10–12 cm) sediments at 885 mbsl and 1306 mbsl (between 30.5–48.4% of total archaeal 16S rRNA gene
240 reads detected in those samples), we performed phylogenetic analyses to determine the diversity of subgroups of the
241 MCG within these sediments. A total of 57 representative 16S rRNA gene reads assigned to MCG were extracted from
242 the dataset and incorporated in a MCG phylogenetic tree of Fillol et al. (2015) (Fig. 2). The majority of MCG 16S
243 rRNA gene reads from the 885 mbsl surface sediment (77.3%; Table 4) clustered in subgroup 15. In the 885 mbsl
244 subsurface sediment, the majority of MCG reads clustered within subgroups 8 and 15 (33.6% and 19.6%, respectively;
245 Table 4). In the 1306 mbsl surface sediment there was only a low relative abundance of MCG (Fig. 1b); all MCG
246 archaea detected clustered in subgroup 15 (Table 4). On the other hand, in the 1306 mbsl subsurface sediment the reads
247 clustered in subgroups 15, 2 and 14 (34.3%, 10.9% and 10.9%, respectively; Fig. 2).

248 As the Thaumarchaeota MGI was dominant in oxygenated sediments at 1306, 2470 and 3003 mbsl (Fig. 1b), we further
249 analyzed the diversity of this group by performing a more detailed phylogeny of the recovered 16S rRNA gene reads
250 attributed to this group. Five OTUs dominated the Thaumarchaeota MG1 (Table 5); we will refer to them as OTU-1 to -

251 5. OTU-1, 2, 3 and 5 were phylogenetically closely related to other known benthic Thaumarchaeota MGI species, such
252 as ‘*Ca. Nitrosoarchaeum koreensis* MY1’ or environmental 16S rRNA gene sequences from marine sediments (Fig.3).
253 On the other hand, OTU-4 clustered with 16S rRNA gene sequences from pelagic Thaumarchaeota MGI species, like
254 *Ca. Nitrosopelagicus brevis*, and also clustered with 16S rRNA sequences recovered from seawater SPM (Fig. 3).
255 OTU-3 was the most abundant OTU in the surface sediment at 1306, 2470, and 3003 mbsl with a relative abundance of
256 44-68% (Table 5). At 1306 mbsl OTU-4 was the second most abundant (35.1%). This OTU had a much lower relative
257 abundance (1.6% and 0.0%) at 2470 and 3003 mbsl, respectively (Table 5). The relative abundance of OTU-2 increased
258 with increasing sampling station depth (Table 5), OTU-1 and 5 had an abundance <5% in the surface sediments (Table
259 5).

260 The diversity of Thaumarchaeota MG1 was further assessed by amplification, cloning and sequencing of the archaeal
261 *amoA* gene. Most of the *amoA* gene sequences from surface (27 out of 29 clones) and subsurface sediment at 885 mbsl
262 (9 out of 10 clones) and just one from the surface sediment from 1306 mbsl (1 out of 58 clones) were closely related
263 with *amoA* gene sequences previously recovered from SPM at 1050 mbsl from this area of the Arabian Sea (Villanueva
264 et al., 2014). Phylogenetically they fall within the ‘Water column B, subsurface water’ *amoA* clade as defined by
265 Francis et al. (2005) (Fig. 4). At 1306 and 3003 mbsl (surface and subsurface) the majority of recovered *amoA* gene
266 sequences clustered within the ‘shallow water/sediment’ clade (100 and 98.3%, respectively) and are closely related
267 with *amoA* gene sequences from water column SPM at 170 mbsl (Villanueva et al., 2014) as well as *amoA* gene coding
268 sequences previously detected in sediments (Villanueva et al., 2014; Fig. 4). Of all recovered *amoA* gene sequences
269 from 885 mbsl only a small fraction (8.3%) clustered within the ‘shallow water/sediment’ clade (Fig. 4).

270 **Abundance and potential activity of archaea in surface and subsurface sediments**

271 The abundance of archaeal 16S rRNA gene copies in the surface sediments of different stations varied slightly: it was
272 lowest at 1306 mbsl (9.8×10^9 copies g^{-1} sediment) and highest at 2470 mbsl (1.5×10^{11} ; Fig. 5a). The potential
273 activity, based on the 16S rRNA gene transcripts of the archaeal 16S rRNA gene, was the lowest at 2470 mbsl (5×10^4
274 transcripts g^{-1} sediment), while a higher potential activity was detected at 885, 1306 and 3003 mbsl ($0.9-42 \times 10^7$; Fig.
275 5a). The abundance of archaeal 16S rRNA gene copies in the subsurface sediment varied also within one and a half
276 order of magnitude ($1.1-54 \times 10^9$; Fig. 5c), with a decrease with increasing water depth. The potential activity showed
277 less variation within the subsurface sediments ($1.2-22 \times 10^7$ 16S rRNA gene transcripts g^{-1} of sediment; Fig. 5c) than in
278 the surface sediments.

279 The abundance of Thaumarchaeota was estimated by quantifying the archaeal *amoA* gene copies. The highest
280 abundance of *amoA* gene copies in surface sediment was detected at 2470 mbsl (1.0×10^9 copies g^{-1} sediment), and the
281 lowest at 885 mbsl (5×10^4 ; Fig. 5b). *AmoA* gene transcripts in surface sediment were under the detection limit at 885
282 mbsl but were detected below the OMZ with 4×10^2 , 2.3×10^6 and 8×10^3 gene transcripts g^{-1} of sediment at 1306,
283 2470 and 3003 mbsl, respectively (Fig. 5b). In subsurface sediments, the abundance of *amoA* gene copies was low at

284 885 and 1306 mbsl ($5.4\text{-}19 \times 10^2$ gene transcripts g^{-1} sediment) and higher at 2470 and 3003 mbsl (4.1×10^5 , 5.4×10^6 ,
285 respectively; Fig. 5d). *AmoA* gene transcripts were not detected in the subsurface sediments (Fig. 5d).

286 **DISCUSSION**

287 In this study, we assessed the changes in benthic archaeal diversity and abundance in sediments of the Arabian Sea
288 oxygen minimum zone along a gradient in bottom water oxygen concentrations. The steep Murray Ridge protrudes the
289 OMZ, allowing the study of sediments deposited under varying bottom water oxygen concentrations. All these
290 sediments receive organic matter (OM), the most important fuel for benthic prokaryotic activity in sediments. This OM
291 is produced in a relatively small area of the ocean (i.e. the station within the OMZ, at 885 mbsl, and well below the
292 OMZ, at 3003 mbsl, are only 110 km apart) and, therefore, is likely composed of the same primary photosynthate.
293 However, due to differences in the degree of mineralization resulting from different exposure to oxic conditions in the
294 water column, there were differences in OM quality. OM in the sediments within the OMZ has a higher biochemical
295 “quality” based on amino acid composition and intact phytopigments compared to OM in the sediments below the OMZ
296 (Koho et al., 2013). Therefore, changes in the quality and flux of OM received by the different sediment niches could
297 also affect the archaeal community composition as several of the archaeal groups (i.e. MCG and MBG-D) reported here
298 have been suggested to use OM as carbon source in anoxic conditions (Lloyd et al., 2013).

299 **Effect of oxygen availability on archaeal diversity and abundance in the surface sediments**

300 We detected large differences in archaeal diversity between the surface sediment deposited within the OMZ and those
301 deposited below the OMZ. In contrast to the diverse anaerobic archaeal community in the surface of 885 mbsl, in
302 surface sediments at 1306, 2470 and 3003 mbsl, Thaumarchaeota MGI were dominant, representing 80-100% of the
303 archaeal population (Fig. 1). This clear difference in the benthic archaeal population in the surface sediments can be
304 attributed to the oxygen availability as Thaumarchaeota are known to require oxygen for their metabolism (i.e.
305 nitrification; Könneke et al., 2005). In fact, the oxygen penetration depth (OPD) was observed to be 3, 10, and 19 mm
306 in sediments at 1306, 2470, and 3003 mbsl, respectively, while in sediments at 885 mbsl, the OPD was barely 0.1 mm
307 (Table S1; Kraal et al., 2012). The surface (0-5 mm) sediment at 1306 mbsl was not fully oxygenated (OPD of 3 mm),
308 which probably explains the detection in relatively low abundance (ca. 20%) of the anaerobic archaea that thrive in the
309 anoxic sediment from 885 mbsl. The low OPD at 1306 mbsl also explains the low *amoA* gene expression in comparison
310 with the deeper surface sediments (Figs. 5b,d). Overall this indicates the presence of Thaumarchaeota with lower
311 activity in the surface sediments at 1306 mbsl (Fig. 5). Within the Thaumarchaeota MGI group, we also detected
312 changes in the relative abundance of specific OTUs in the surface sediments at 1306, 2470 and 3003 mbsl (Table 5).
313 For example, OTU-2 becomes progressively more abundant with increasing water depth, suggesting that this OTU is
314 favored at the higher oxygen concentrations found in the surface sediment at 3003 mbsl. OTU-4 was closely affiliated

315 with '*Ca. Nitrosopelagicus brevis*', a pelagic MG-I member, which indicates that this DNA is most likely derived from
316 the overlying water column (Table 5), and thus should be considered to represent fossil DNA.

317 High *amoA* gene abundances were detected in the surface sediment at 2470 and 3003 mbsl, while values in the surface
318 of 885 mbsl were approximately three orders of magnitude less. The lack of oxygen in the surface sediments at 885
319 mbsl and in the subsurface sediments, as well as undetectable *amoA* gene transcripts at those depths, suggest that in
320 these cases the *amoA* gene DNA signal is fossil. It is well known that under anoxic conditions DNA of marine pelagic
321 microbes may become preserved in sediments even for periods of thousands of years (Boere et al., 2011; Coolen et al.,
322 2004b). The fossil origin of the Thaumarchaeotal *amoA* gene is supported by the phylogenetic affiliation of the *amoA*
323 gene fragments amplified from the 885 mbsl surface sediment, as those sequences were closely related to *amoA* gene
324 sequences detected in the suspended particulate matter in the subsurface water column (Villanueva et al., 2015), thus
325 suggesting that the detected DNA originated from pelagic Thaumarchaeota present in the subsurface water column, as
326 proposed for the presence of OTU-4 16S rRNA gene sequences in the surface sediment (see earlier).

327 There is a discrepancy between the 16S rRNA gene copy numbers and the *amoA* gene copy numbers within the
328 sediments (Fig. 5). *AmoA* gene copies were consistently lower than the 16S rRNA gene copies, even within sediments
329 that were completely dominated by Thaumarchaeota MG-I. This may be caused by the *amoA* gene primer mismatches
330 and/or the disparity of gene copy numbers within the archaeal genomes (Park et al., 2008).

331 In the anoxic surface sediment at 885 mbsl (within the OMZ), we detected a highly diverse archaeal population
332 composed of MCG, Thermoplasmatales, MBG-B, -D and -E, Woesearchaeota, and MHVG. Archaeal groups such as
333 MCG and MBG-B and E have been previously described in anoxic marine sediments, where they have been suggested
334 to be involved in anaerobic OM degradation (e.g. Biddle et al., 2006; Inagaki et al., 2003; Castelle et al., 2015).
335 Members of the DPANN Woesearchaeota were only present in the surface sediment at 885 mbsl but not in the
336 subsurface anoxic sediments at 885 and 1306 mbsl, suggesting that their presence here is not solely dependent on the
337 absence of oxygen but possibly also on the OM composition and availability in surface and subsurface sediments.
338 Alternatively, the DPANN Woesearchaeota 16S rRNA gene signal could also originate from the water column and
339 deposited in the surface sediment at 885 mbsl as fossil DNA as observed for the case of Thaumarchaeota as mentioned
340 above.

341 **Archaeal community composition in the anoxic subsurface sediments**

342 The archaeal diversity in the subsurface sediment (10–12 cm) from both 885 and 1306 mbsl (i.e. dominated by MCG,
343 MBG-B, -D and -E) is similar to that observed in the surface sediment at 885 mbsl. This supports that oxygen
344 availability is an important factor for determining the diversification of archaeal groups (Fig. 1b). MCG, one of the
345 dominant archaeal groups in these sediments, showed substantial differences in the distribution of its subgroups (Table
346 4). All subsurface sediments had a high intra-group diversity of MCG. This is in contrast with the surface sediment at
347 885 and 1306 mbsl where a high relative abundance of the subgroup MCG-15 is detected (Table 4). A recent study with

348 enrichments of estuarine sediments showed evidence that MCG-8 are capable of growth with lignin as an energy source
349 (Yu et al., 2018). MCG-8 was the most or one of the most abundant MCG groups in our studied subsurface sediments
350 (Table 4). Substantial lignin concentration were previously not detected in the Arabian Sea, which indicated a relatively
351 invariant terrestrial OM contribution (Cowie et al., 1999). Therefore it remains uncertain which physicochemical
352 factor(s) influence the MCG-8 subgroup in the studied subsurface sediments. A recent survey of the ecological niches
353 and substrate preferences of the MCG in estuarine sediments based on genomic data pointed to MCG-6 archaea as
354 degraders of complex extracellular carbohydrate polymers (plant-derived), while subgroups 1, 7, 15 and 17 have mainly
355 the potential to degrade detrital proteins (Lazar et al., 2016). Lazar et al. (2016) also described the presence of
356 aminopeptidases coded in the genome bin of MCG-15, suggesting that this subgroup could be specialized in
357 degradation of extracellular peptides in comparison with the other MCG subgroups, which would be restricted to the
358 use of amino acid and oligopeptides. Considering the dominance of the MCG-15 subgroup in the surface sediments
359 analyzed in this study (Table 4), we hypothesize that the proteinaceous OM deposited in the surface sediment, which
360 mainly originates from photosynthate, is still quite undegraded. This would favor the MCG-15 in this niche, fueling its
361 metabolism by the degradation of peptides extracellularly, while in subsurface sediments, other MCG groups such as 2,
362 8 and 14 would be more favored.

363 The archaeal 16S rRNA gene abundance in the subsurface sediments progressively declined with increasing water
364 depth, while the potential activity was similar. This can be due to the expected decrease in the flux of OM being
365 delivered to these anoxic sediments layers attributed to higher degradation of OM in oxygenated bottom waters and the
366 progressively larger oxic zone in the sediments (Lengger et al., 2012; Nierop et al., 2017). This results in lower organic
367 carbon concentrations and a decreased biochemical quality of the OM (Koho et al., 2013; Nierop et al., 2017) to sustain
368 the heterotrophic archaeal population inhabiting the anoxic subsurface sediments. Also the presence or lack of
369 macrofauna in the analyzed sediments would have an effect on the OM composition, sediments within the OMZ are less
370 prone to bioturbation which most likely resulted in higher OM preservation (Koho et al., 2013). Differences in the OM
371 biochemical composition can influence the microbial community composition as was shown recently for North Sea
372 sediments (Oni et al., 2015).

373 **Benthic archaea as potential sources for archaeal IPLs**

374 Archaeal lipids in surface and deeper sediments of the Murray Ridge (Lengger et al., 2012, 2014), as well as in the
375 overlying water column (Pitcher et al., 2011; Schouten et al., 2012), have been studied earlier. The study by Lengger et
376 al. (2012) was limited to the determination of MH-, DH- and HPH-crenarchaeol with HPLC/ ESI-MS² using a specific
377 selected reaction monitoring method (SRM; Pitcher et al., 2011). A follow-up study of Lengger et al. (2014) reported
378 MH-, DH- and HPH-IPLs with multiple CL-GDGTs. In our study, we expanded the screening for IPLs carrying
379 different polar head groups in combination with multiple CLs using high resolution accurate mass/mass spectrometry
380 (see Table S1). By applying this method, we were able to target a broader range of IPLs in these sediments. This allows

381 a more direct comparison with the archaeal diversity detected by gene-based methods. Note that the study of Lengger et
382 al. (2014) used a different sampling resolution (surface sediment used was 0–2 compared to our 0–0.5 cm) and our
383 results can, therefore, not be directly compared.

384 The fully oxygenated surface sediments showed a dominance of GDGT-0 and crenarchaeol mostly with HPH as IPL-
385 type (Table 2). This is the expected IPL-GDGT signature of Thaumarchaeota as previously observed in pure cultures
386 (Elling et al., 2017; Pitcher et al., 2010; Schouten et al., 2008; Sinninghe Damsté et al., 2012). The predominance of
387 the HPH IPL-type in surface (0–2 cm) sediments was previously interpreted as an indication of the presence of an
388 active Thaumarchaeotal population synthesizing membrane lipids *in situ* (Lengger et al., 2012, 2014), taking into
389 account the labile nature of sedimentary phospholipids (Harvey et al., 1986; Schouten et al., 2010; Xie et al., 2013).
390 Elling et al. (2014) showed, in a Thaumarchaeota culture experiment, that a high HPH-crenarchaeal abundance was an
391 indication of metabolically active Thaumarchaeota. The hypothesis by Lengger et al. (2012, 2014) is strongly supported
392 by our data because (i) the archaeal community in the oxygenated surface (0–0.5 cm) sediments is dominated by
393 Thaumarchaeota (Fig. 1) and (ii) the high abundance of thaumarchaeotal *amoA* gene copies and gene transcripts
394 detected in the oxygenated surface sediments. On the other hand, in the anoxic surface sediment at 885 mbsl,
395 crenarchaeol was predominantly present with DH as the predominant IPL-type (Table 2). This is considered to be a
396 fossil signal of Thaumarchaeota deposited from the water column due to a higher preservation potential of glycolipid
397 head groups (as present in DH) as previously suggested (Lengger et al., 2012, 2014).. Xie et al. (2013) showed a high
398 preservation potential for glycosidic ether lipids in a 300-day incubation study. However the differences in degradation
399 rates between phospholipid versus glycolipid GDGTs still need to be determined, especially on longer time scales that
400 apply to sediments. Nevertheless, the presence of *amoA* gene sequences in the 885 mbsl surface sediment, which are
401 derived from the deeper water column, as well as the much lower *amoA* gene abundance and lack of *amoA* gene
402 expression (Fig. 5b, d) supports the contention that the crenarchaeol IPLs in the surface sediment at 885 mbsl are
403 predominantly fossil since evidence for active Thaumarchaeota is lacking.

404 The low relative abundance of GDGT-0 IPLs in the surface sediment at 885 mbsl (Table 2) is remarkable. Only MH-
405 GDGT-0 was detected in low relative abundance (0.3 %), whereas any other of the IPL-types with GDGT-0 as CL that
406 were screened for in our study (Table S2; Fig. 1b) was absent. In contrast, Lengger et al. (2014) reported a significant
407 amount of IPL-derived CL-GDGT-0 (i.e. 18.5% of total CL GDGTs) when the head groups of the IPLs are released by
408 acid hydrolysis and the remaining CLs were analyzed in a surface (0–0.5 cm) sediment from the same site. We re-
409 analyzed the IPL-derived CL-GDGT composition in the surface (0–0.5 cm) sediment at 88 mbsl and recovered an
410 identical CL-GDGT distribution as reported by Lengger et al. (2014). The discrepancy between CL and IPL distribution
411 may partly be explained by the underestimation of MH IPLs by our method. To assess the MH IPLs underestimation,
412 two North Atlantic suspended particulate matter samples analyzed with the normal phase method (this study) and the

413 reverse phase method (Wörmer et al., 2013). This underestimation of MH IPLs with the used normal phase method
414 could be on average 10 fold compared to the reverse phase method (Fig. S4). However, the difference in response factor
415 for the different IPL types is not sufficient to explain the discrepancy between IPL-derived CL-GDGT-0 and IPL
416 GDGT-0 relative abundances. Therefore, we speculate it is due to the presence of an IPL-type with unknown head
417 groups not included in our analytical window. This unknown IPL GDGT-0 may originate from any of the archaeal
418 groups present in the surface sediment at 885 mbsl, such as MCG, Thermoplasmatales, MBG-B, MBG-E and
419 Euryarchaeota MHVG. DPANN Woese archaeota is also relatively abundant in the surface sediments at 885 mbsl (Fig.
420 1) but recent studies suggest that their small genomes lack the genes coding for the enzymes of the archaeal lipid
421 biosynthetic pathway (Jahn et al., 2004; Podar et al., 2013; Villanueva et al., 2017; Waters et al., 2003). Therefore, they
422 are not expected to contribute to the IPL-GDGT pool. Ruling out the Woese archaeota as a possible source of IPL-
423 GDGTs, the IPL GDGT-0 with unknown polar head group(s) in the surface sediment at 885 mbsl may be attributed to
424 the MCG, which make up 30.5% of the archaeal 16S rRNA gene reads in this sediment. Most of these MCG archaea
425 fall into subgroup MCG-15 (Table 4). Previous studies proposed butanetriol dibiphytanyl glycerol tetraethers (BDGTs)
426 as putative biomarker of the MCG based on the correlation between the presence of these components and MCG in
427 estuarine sediments (Meador et al., 2014). However, we did not detect any IPL BDGTs in the sediments analyzed in our
428 study. Buckles et al. (2013) suggested that members of the MCG and Crenarchaeota group 1.2 could be the biological
429 source of IPL GDGT-0 found in the anoxic hypolimnion of a tropical lake. Considering these evidences, it is possible
430 that the unknown IPL GDGT-0 present in the surface sediment at 885 mbsl could be a biomarker for MCG.

431 In subsurface sediments, the IPL GDGT distribution was remarkably different from that detected in the surface
432 oxygenated sediment as higher relative abundances of GDGT-1, 2, 3 and 4 were detected in detriment of GDGT-0,
433 similar to the distribution detected in the surface sediment at 885 mbsl. This may represent new archaeal production in
434 the anoxic sediments, selective preservation of archaeal lipids produced in the water column and surface sediments, or
435 both. The HCP GDGT-0 and two isomers of the DH GDGT-0 (Fig. S2) were detected in the subsurface sediments at
436 885 and 1306 mbsl but not in those from deeper waters (Table 2). Since these IPLs were not detected in the surface
437 sediments, it is likely that they are produced *in situ*. Unfortunately, we only obtained information on the archaeal
438 community composition of the subsurface sediments at shallow water depth, so we cannot compare these with the
439 subsurface sediments from deeper waters that lack these DH moieties, which could have provided a clue towards the
440 archaeal source of these IPLs. An IPL composed of GDGT-0 with a cyclopentanetetraol head group has been
441 previously detected in cultures of the hyperthermophilic crenarchaeal *Sulfolobales* (Langworthy et al., 1974; Sturt et al.,
442 2004). However, members of the *Sulfolobales* were not detected in our 16S rRNA gene amplicon sequencing data. We
443 also detected a high relative abundance of MCG (up to 48.4% relative abundance) in the subsurface sediment at 885 and
444 1306 mbsl (Fig. 1). The diversity of the MCG population in the subsurface sediments was much higher in comparison
445 with the diversity in surface sediments at 885 mbsl as sequences closely related to the MCG subgroups, 2, 8, 10, 14, 5b,

446 15, and 17 were detected both in the 885 mbsl and 1306 mbsl subsurface sediments (Fig. 2). This presence of these
447 different MCG members, likely caused by niche differentiation (see before), may be the origin of the unusual DH-
448 GDGT-0 isomer and the HCP-GDGT-0 IPL that we detected within the subsurface sediments at 885 and 1306 mbsl.

449 CONCLUSIONS

450 By using a combined 16S rRNA gene amplicon sequencing and IPL analysis with high-resolution accurate mass/mass
451 spectrometry we have unraveled the high diversity of benthic archaea harbored in oxygenated and anoxic sediments of
452 the Arabian Sea, as well as widening our detection window of archaeal intact polar lipids. DNA-based analyses
453 revealed a dominance of active benthic *in situ* Thaumarchaeota in those sediment where oxygen was present, which
454 coincided with high relative abundance of the HPH-crenarchaeol previously suggested to be a marker of living
455 Thaumarchaeota. In the anoxic marine sediments analyzed here, members of the MCG, DPANN and Euryarchaeota
456 Thermoplasmatales dominated. We also observed a high diversity within the MCG with a more diverse population in
457 subsurface sediments. Subsurface anoxic sediments had a high relative abundance of IPL GDGT-1, 2, and 3 with DH
458 headgroups, which could either be attributed to fossil signal due to the more recalcitrant nature of the glycosidic bonds
459 or being IPLs synthesized by the archaeal groups detected in those sediments. In addition, IPL GDGT-0 was also
460 detected with a hexose head group on both ends of the core lipid, two hexoses on one end, and a cyclopentanetetraol
461 molecule bound to the core lipid and a hexose attached to it. Members of the DPANN could possibly be ruled out of
462 making those lipids due to the lack of lipid biosynthetic pathway. Dominant archaeal members in those sediments such
463 as the MCG and Thermoplasmatales, could be potential biological sources of these IPLs. To conclude, this combined
464 approach has shed light on the possible biological sources of specific archaeal IPLs and also detected a highly diverse
465 benthic archaeal community.

466 Acknowledgments

467 Elda Panoto is thanked for assistance with molecular analyses. We would like to thank the captain and crew of the RV
468 *Pelagia* as well as the cruise leader, technicians and scientists participating in cruise 64PE301. This PASOM cruise was
469 funded by the Earth and Life Science and Research Council (ALW) with financial aid from the Netherlands
470 Organization for Scientific Research (NWO) (grant 817.01.015) to Prof. G.J. Reichart (PI). NIOZ is acknowledged for
471 the studentship of MAB. This research was further supported by the NESSC and SIAM Gravitation Grants
472 (024.002.001 and 024.002.002) from the Dutch Ministry of Education, Culture and Science (OCW) and the European
473 Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement
474 n° 694569—MICROLIPIDS) to JSSD.

475 **References**

- 476 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J.: Basic local alignment search tool., *J. Mol. Biol.*,
477 215(3), 403–10, doi:10.1016/S0022-2836(05)80360-2, 1990.
- 478 Biddle, J. F., Lipp, J. S., Lever, M. a, Lloyd, K. G., Sørensen, K. B., Anderson, R., Fredricks, H. F., Elvert, M., Kelly,
479 T. J., Schrag, D. P., Sogin, M. L., Brenchley, J. E., Teske, A., House, C. H. and Hinrichs, K.-U.: Heterotrophic Archaea
480 dominate sedimentary subsurface ecosystems off Peru., *Proc. Natl. Acad. Sci. U. S. A.*, 103(10), 3846–3851,
481 doi:10.1073/pnas.0600035103, 2006.
- 482 Bligh, E. G., Dyer, W. J.: A rapid method of total lipid extraction and Purification, *Can. J. Biochem. Physiol.*, 37, 911–
483 917, doi:10.1139/o59-099., 1959.
- 484 Boere, A. C., Rijpstra, W. I. C., de Lange, G. J., Sinninghe Damsté, J. S. and Coolen, M. J. L.: Preservation potential of
485 ancient plankton DNA in Pleistocene marine sediments, *Geobiology*, 9, 377–393, doi:10.1111/j.1472-
486 4669.2011.00290.x, 2011.
- 487 Buckles, L. K., Villanueva, L., Weijers, J. W. H., Verschuren, D. and Sinninghe Damsté, J. S.: Linking isoprenoidal
488 GDGT membrane lipid distributions with gene abundances of ammonia-oxidizing Thaumarchaeota and uncultured
489 crenarchaeotal groups in the water column of a tropical lake (Lake Challa, East Africa), *Environ. Microbiol.*, 2, 1–18,
490 doi:10.1111/1462-2920.12118, 2013.
- 491 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G.,
492 Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A.,
493 McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J.,
494 Yatsunencko, T., Zaneveld, J. and Knight, R.: correspondEnce QIIME allows analysis of high- throughput community
495 sequencing data Intensity normalization improves color calling in SOLiD sequencing, *Nat. Publ. Gr.*, 7(5), 335–336,
496 doi:10.1038/nmeth0510-335, 2010.
- 497 Castelle, C. J., Wrighton, K. C., Thomas, B. C., Hug, L. A., Brown, C. T., Wilkins, M. J., Frischkorn, K. R., Tringe, S.
498 G., Singh, A., Markillie, L. M., Taylor, R. C., Williams, K. H. and Banfield, J. F.: Genomic Expansion of Domain
499 Archaea Highlights Roles for Organisms from New Phyla in Anaerobic Carbon Cycling, *Curr. Biol.*, 25, 1–12,
500 doi:10.1016/j.cub.2015.01.014, 2015.
- 501 Coolen, M. J. L., Muyzer, G., Rijpstra, W. I. C., Schouten, S., Volkman, J. K. and Sinninghe Damsté, J. S.: Combined
502 DNA and lipid analyses of sediments reveal changes in Holocene haptophyte and diatom populations in an Antarctic
503 lake, *Earth Planet. Sci. Lett.*, 223(1–2), 225–239, doi:10.1016/j.epsl.2004.04.014, 2004a.
- 504 Coolen, M. J. L., Hopmans, E. C., Rijpstra, W. I. C., Muyzer, G., Schouten, S., Volkman, J. K. and Sinninghe Damsté,
505 J. S.: Evolution of the methane cycle in Ace Lake (Antarctica) during the Holocene: response of methanogens and
506 methanotrophs to environmental change, *Org. Geochem.*, 35(10), 1151–1167, doi:10.1016/j.orggeochem.2004.06.009,
507 2004b.
- 508 Cowie, G. L., Calvert, S. E., Pedersen, T. F., Schulz, H. and Von Rad, U.: Organic content and preservational controls
509 in surficial shelf and slope sediments from the Arabian Sea (Pakistan margin), *Mar. Geol.*, 161(1), 23–38,
510 doi:10.1016/S0025-3227(99)00053-5, 1999.
- 511 Delong, E. F. and Pace, N. R.: Environmental Diversity of Bacteria and Archaea, *Soc. Syst. Biol.*, 50(4), 470–478,
512 doi:10.1080/10635150118513, 2001.
- 513 DeLong, E. F., Ying Wu, K., Prézelin, B. B. and Jovine, R. V. M.: High abundance of Archaea in Antarctic marine
514 picoplankton, *Nature*, 371, 695–697, doi:10.1038/371695a0, 1994.
- 515 Durbin, A. M. and Teske, A.: Sediment-associated microdiversity within the Marine Group I Crenarchaeota, *Environ.*
516 *Microbiol. Rep.*, 2(5), 693–703, doi:10.1111/j.1758-2229.2010.00163.x, 2010.
- 517 Edgar, R. C.: MUSCLE: Multiple sequence alignment with high accuracy and high throughput, *Nucleic Acids Res.*,
518 32(5), 1792–1797, doi:10.1093/nar/gkh340, 2004.
- 519 Elling, F. J., Konneke, M., Lipp, J. S., Becker, K. W., Gagen, E. J. and Hinrichs, K. U.: Effects of growth phase on the
520 membrane lipid composition of the thaumarchaeon *Nitrosopumilus maritimus* and their implications for archaeal lipid
521 distributions in the marine environment, *Geochim. Cosmochim. Acta*, 141, 579–597, doi:10.1016/j.gca.2014.07.005,
522 2014.
- 523 Elling, F. J., Könneke, M., Mußmann, M., Greve, A. and Hinrichs, K.-U.: Influence of temperature, pH, and salinity on
524 membrane lipid composition and TEX86 of marine planktonic thaumarchaeal isolates, *Geochim. Cosmochim. Acta*,
525 171, 238–255, doi:10.1016/j.gca.2015.09.004, 2015.
- 526 Elling, F. J., Könneke, M., Nicol, G. W., Stieglmeier, M., Bayer, B., Spieck, E., de la Torre, J. R., Becker, K. W.,
527 Thomm, M., Prosser, J. I., Herndl, G. J., Schleper, C. and Hinrichs, K. U.: Chemotaxonomic characterisation of the
528 thaumarchaeal lipidome, *Environ. Microbiol.*, 19(7), 2681–2700, doi:10.1111/1462-2920.13759, 2017.

- 529 Fillol, M., Auguet, J.-C., Casamayor, E. O. and Borrego, C. M.: Insights in the ecology and evolutionary history of the
530 Miscellaneous Crenarchaeotic Group lineage., *ISME J.*, 1–13, doi:10.1038/ismej.2015.143, 2015.
- 531 Harvey, H. R., Fallon, R. D. and Patton, J. S.: The effect of organic matter and oxygen on the degradation of bacterial
532 membrane lipids in marine sediments, *Geochim. Cosmochim. Acta*, 50, 795–804, doi:10.1016/0016-7037(86)90355-8,
533 1986.
- 534 Hopmans, E. C., Schouten, S. and Sinninghe, J. S.: The effect of improved chromatography on GDGT-based
535 palaeoproxies, *Org. Geochem.*, 93, 1–6, doi:10.1016/j.orggeochem.2015.12.006, 2016.
- 536 Inagaki, F., Suzuki, M., Takai, K., Oida, H., Sakamoto, T., Aoki, K., Nealson, K. H. and Horikoshi, K.: Microbial
537 Communities Associated with Geological Horizons in Coastal Subseafloor Sediments from the Sea of Okhotsk
538 Microbial Communities Associated with Geological Horizons in Coastal Subseafloor Sediments from the Sea of
539 Okhotsk, *Appl. Environ. Microbiol.*, 69(12), 7224–7235, doi:10.1128/AEM.69.12.7224, 2003.
- 540 Ingalls, A. E., Huguet, C. and Truxal, L. T.: Distribution of intact and core membrane lipids of archaeal glycerol dialkyl
541 glycerol tetraethers among size-fractionated particulate organic matter in Hood Canal, Puget Sound, *Appl. Environ.*
542 *Microbiol.*, 78(5), 1480–1490, doi:10.1128/AEM.07016-11, 2012.
- 543 Jahn, U., Summons, R., Sturt, H., Grosjean, E. and Huber, H.: Composition of the lipids of Nanoarchaeum equitans and
544 their origin from its host Ignicoccus sp. strain KIN4/I, *Arch. Microbiol.*, 182(5), 404–413, doi:10.1007/s00203-004-
545 0725-x, 2004.
- 546 Jorgenson, S. L., Hannisdal, B., Lanzén, A., Baumberger, T., Flesland, K., Fonseca, R., Øvreås, L., Steen, I. H.,
547 Thorseth, I. H., Pedersen, R. B. and Schleper, C.: Correlating microbial community profiles with geochemical data in
548 highly stratified sediments from the Arctic Mid-Ocean Ridge, *Proc. Natl. Acad. Sci.*, 109(42), E2846–E2855,
549 doi:10.1594/PANGAEA.786687, 2012.
- 550 Karner, M. B., DeLong, E. F. and Karl, D. M.: Archaeal dominance in the mesopelagic zone of the Pacific Ocean,
551 *Nature*, 409(6819), 507–510, doi:10.1038/35054051, 2001.
- 552 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glockner, F. O.: Evaluation of general
553 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies, *Nucleic*
554 *Acids Res.*, 41(1), 1–11, doi:10.1093/nar/gks808, 2013.
- 555 Koga, Y. and Morii, H.: Biosynthesis of ether-type polar lipids in archaea and evolutionary considerations., *Microbiol.*
556 *Mol. Biol. Rev.*, 71(1), 97–120, doi:10.1128/MMBR.00033-06, 2007.
- 557 Koho, K. A., Nierop, K. G. J., Moodley, L., Middelburg, J. J., Pozzato, L., Soetaert, K., Van Der Plicht, J. and Reichart,
558 G. J.: Microbial bioavailability regulates organic matter preservation in marine sediments, *Biogeosciences*, 10(2),
559 1131–1141, doi:10.5194/bg-10-1131-2013, 2013.
- 560 Könneke, M., Bernhard, A. E., de la Torre, J. R., Walker, C. B., Waterbury, J. B. and Stahl, D. A.: Isolation of an
561 autotrophic ammonia-oxidizing marine archaeon, *Nature*, 437(22), 543–546, doi:10.1038/nature03911, 2005.
- 562 Kraal, P., Slomp, C. P., Reed, D. C., Reichart, G.-J. and Poulton, S. W.: Sedimentary phosphorus and iron cycling in
563 and below the oxygen minimum zone of the northern Arabian Sea, *Biogeosciences*, 9(7), 2603–2624, doi:10.5194/bg-9-
564 2603-2012, 2012.
- 565 Kubo, K., Lloyd, K. G., F Biddle, J., Amann, R., Teske, A. and Knittel, K.: Archaea of the Miscellaneous
566 Crenarchaeotal Group are abundant, diverse and widespread in marine sediments, *ISME J.*, 6(10), 1949–1965,
567 doi:10.1038/ismej.2012.37, 2012.
- 568 De La Torre, J. R., Walker, C. B., Ingalls, A. E., Könneke, M. and Stahl, D. A.: Cultivation of a thermophilic ammonia
569 oxidizing archaeon synthesizing crenarchaeol, *Environ. Microbiol.*, 10(3), 810–818, doi:10.1111/j.1462-
570 2920.2007.01506.x, 2008.
- 571 Langworthy, T. A., Mayberry, W. R. and Smith, P. F.: Long-chain glycerol diether and polyol dialkyl glycerol triether
572 lipids of *Sulfolobus acidocaldarius*, *J. Bacteriol.*, 119(1), 106–116, 1974.
- 573 Lazar, C. S., Baker, B. J., Seitz, K., Hyde, A. S., Dick, G. J., Hinrichs, K. U. and Teske, A. P.: Genomic evidence for
574 distinct carbon substrate preferences and ecological niches of Bathyarchaeota in estuarine sediments, *Environ.*
575 *Microbiol.*, 18(4), 1200–1211, doi:10.1111/1462-2920.13142, 2016.
- 576 Learman, D. R., Henson, M. W., Thrash, J. C., Temperton, B., Brannock, P. M., Santos, S. R., Mahon, A. R. and
577 Halanaych, K. M.: Biogeochemical and microbial variation across 5500 km of Antarctic surface sediment implicates
578 organic matter as a driver of benthic community structure, *Front. Microbiol.*, 7, 1–11, doi:10.3389/fmicb.2016.00284,
579 2016.
- 580 Lengger, S. K., Hopmans, E. C., Reichart, G.-J., Nierop, K. G. J., Sinninghe Damsté, J. S. and Schouten, S.: Intact polar
581 and core glycerol dibiphytanyl glycerol tetraether lipids in the Arabian Sea oxygen minimum zone. Part II: Selective
582 preservation and degradation in sediments and consequences for the TEX86, *Geochim. Cosmochim. Acta*, 98, 244–258,

583 doi:10.1016/j.gca.2012.05.003, 2012.

584 Lengger, S. K., Hopmans, E. C., Sinninghe Damsté, J. S. and Schouten, S.: Impact of sedimentary degradation and deep
585 water column production on GDGT abundance and distribution in surface sediments in the Arabian Sea: Implications
586 for the TEX86 paleothermometer, *Geochim. Cosmochim. Acta*, 142, 386–399, doi:10.1016/j.gca.2014.07.013, 2014.

587 Lincoln, S. a, Wai, B., Eppley, J. M., Church, M. J., Summons, R. E. and DeLong, E. F.: Planktonic Euryarchaeota are
588 a significant source of archaeal tetraether lipids in the ocean., *Proc. Natl. Acad. Sci. U. S. A.*, 111(27), 9858–63,
589 doi:10.1073/pnas.1409439111, 2014a.

590 Lincoln, S. a, Wai, B., Eppley, J. M., Curch, M. J., Summons, R. E. and Delong, E. F.: Reply to Schouten et al. : Marine
591 Group II planktonic Euryarchaeota are significant contributors to tetraether lipids in the ocean, *Proc. Natl. Acad. Sci.*,
592 111(41), 4286, doi:10.1073/pnas.1416736111, 2014b.

593 Lipp, J. S. and Hinrichs, K.-U.: Structural diversity and fate of intact polar lipids in marine sediments, *Geochim.*
594 *Cosmochim. Acta*, 73(22), 6816–6833, doi:10.1016/j.gca.2009.08.003, 2009.

595 Lipp, J. S., Morono, Y., Inagaki, F. and Hinrichs, K.-U.: Significant contribution of Archaea to extant biomass in
596 marine subsurface sediments., *Nature*, 454(August), 991–994, doi:10.1038/nature07174, 2008.

597 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. a, Steen, A. D., Stepanauskas, R., Richter, M.,
598 Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.: Predominant archaea in marine sediments degrade detrital
599 proteins., *Nature*, 496(7444), 215–8, doi:10.1038/nature12033, 2013.

600 Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadukumar, Buchner, A., Lai, T., Steppi, S., Jobb, G.,
601 Förster, W., Brettske, I., Gerber, S., Ginhart, A. W., Gross, O., Grumann, S., Hermann, S., Jost, R., König, A., Liss, T.,
602 Lüßmann, R., May, M., Nonhoff, B., Reichel, B., Strehlow, R., Stamatakis, A., Stuckmann, N., Vilbig, A., Lenke, M.,
603 Ludwig, T., Bode, A. and Schleifer, K.-H.: ARB: a software environment for sequence data., *Nucleic Acids Res.*, 32(4),
604 1363–71, doi:10.1093/nar/gkh293, 2004.

605 Massana, R., Castresana, J., Balague, V., Guillou, L., Romari, K., Groisillier, A., Valentin, K. and Pedros-Alio, C.:
606 Phylogenetic and Ecological Analysis of Novel Marine Stramenopiles, *Appl. Environ. Microbiol.*, 70(6), 3528–3534,
607 doi:10.1128/AEM.70.6.3528, 2004.

608 Meador, T. B., Bowles, M., Lazar, C. S., Zhu, C., Teske, A. and Hinrichs, K.-U.: The archaeal lipidome in estuarine
609 sediment dominated by members of the Miscellaneous Crenarchaeotal Group, *Environ. Microbiol.*, 17(7), 2441–2458,
610 doi:10.1111/1462-2920.12716, 2015.

611 Meng, J., Xu, J., Qin, D., He, Y., Xiao, X. and Wang, F.: Genetic and functional properties of uncultivated MCG
612 archaea assessed by metagenome and gene expression analyses, *ISME J.*, 8(3), 650–659, doi:10.1038/ismej.2013.174,
613 2014.

614 Moore, E. K., Villanueva, L., Hopmans, E. C., Rijpstra, W. I. C., Mets, A., Dedysh, S. N. and Sinninghe Damsté, J. S.:
615 Abundant trimethylornithine lipids and specific gene sequences are indicative of planctomycete importance at the
616 oxic/anoxic interface in sphagnum-dominated northern wetlands, *Appl. Environ. Microbiol.*, 81(18), 6333–6344,
617 doi:10.1128/AEM.00324-15, 2015.

618 Nierop, K. G. J., Reichart, G., Veld, H. and Sinninghe Damsté, J. S.: The influence of oxygen exposure time on the
619 composition of macromolecular organic matter as revealed by surface sediments on the Murray Ridge (Arabian Sea),
620 *Geochim. Cosmochim. Acta*, 206, 40–56, 2017.

621 Offre, P., Spang, A. and Schleper, C.: Archaea in Biogeochemical Cycles, *Annu. Rev. Microbiol.*, 437–457,
622 doi:10.1146/annurev-micro-092412-155614, 2013.

623 Oni, O. E., Schmidt, F., Miyatake, T., Kasten, S., Witt, M., Hinrichs, K.-U. and Friedrich, M. W.: Microbial
624 Communities and Organic Matter Composition in Surface and Subsurface Sediments of the Helgoland Mud Area,
625 North Sea, *Front. Microbiol.*, 6(November), 1–16, doi:10.3389/fmicb.2015.01290, 2015.

626 Park, S.-J., Park, B.-J. and Rhee, S.-K.: Comparative analysis of archaeal 16S rRNA and amoA genes to estimate the
627 abundance and diversity of ammonia-oxidizing archaea in marine sediments, *Extremophiles*, 12(4), 605–615,
628 doi:10.1007/s00792-008-0165-7, 2008.

629 Pitcher, A., Rychlik, N., Hopmans, E. C., Spieck, E., Rijpstra, W. I. C., Ossebaar, J., Schouten, S., Wagner, M. and
630 Sinninghe Damsté, J. S.: Crenarchaeol dominates the membrane lipids of *Candidatus Nitrososphaera gargensis*, a
631 thermophilic group I.1b Archaeon, *ISME J.*, 4(4), 542–52, doi:10.1038/ismej.2009.138, 2010.

632 Pitcher, A., Villanueva, L., Hopmans, E. C., Schouten, S., Reichart, G.-J. and Sinninghe Damsté, J. S.: Niche
633 segregation of ammonia-oxidizing archaea and anammox bacteria in the Arabian Sea oxygen minimum zone., *ISME J.*,
634 5(12), 1896–904, doi:10.1038/ismej.2011.60, 2011.

635 Podar, M., Makarova, K. S., Graham, D. E., Wolf, Y. I., Koonin, E. V and Reysenbach, A.-L.: Insights into archaeal
636 evolution and symbiosis from the genomes of a nanoarchaeon and its inferred crenarchaeal host from Obsidian Pool,

- 637 Yellowstone National Park., *Biol. Direct*, 8(1), 9, doi:10.1186/1745-6150-8-9, 2013.
- 638 Qin, W., Carlson, L. T., Armbrust, E. V., Devol, A. H., Moffett, J. W., Stahl, D. A. and Ingalls, A. E.: Confounding
639 effects of oxygen and temperature on the TEX₈₆ signature of marine Thaumarchaeota, *Proc. Natl. Acad. Sci.*, 112(35),
640 10979–10984, doi:10.1073/pnas.1501568112, 2015.
- 641 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glockner, F. O.: The SILVA
642 ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic Acids Res.*,
643 41(November 2012), D590–D596, doi:10.1093/nar/gks1219, 2013.
- 644 Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N. N., Anderson, I. J., Cheng, J.-F., Darling, A. E., Malfatti, S., Swan,
645 B. K., Gies, E. a, Dodsworth, J. a, Hedlund, B. P., Tsiamis, G., Sievert, S. M., Liu, W.-T., Eisen, J. a, Hallam, S. J.,
646 Kyrpides, N. C., Stepanauskas, R., Rubin, E. M., Hugenholtz, P. and Woyke, T.: Insights into the phylogeny and coding
647 potential of microbial dark matter., *Nature*, 499(7459), 431–437, doi:10.1038/nature12352, 2013.
- 648 De Rosa, M. and Gambacorta, A.: The lipids of archaeobacteria, *The lipids of archaeobacteria*, 27, 153–175, 1988.
- 649 Rossel, P. E., Lipp, J. S., Fredricks, H. F., Arnds, J., Boetius, A., Elvert, M. and Hinrichs, K. U.: Intact polar lipids of
650 anaerobic methanotrophic archaea and associated bacteria, *Org. Geochem.*, 39(8), 992–999,
651 doi:10.1016/j.orggeochem.2008.02.021, 2008.
- 652 Saitou, N. and Nei, M.: The Neighbor-joining Method: A New Method for Reconstructing Phylogenetic Trees', *Mol.*
653 *Biol. Evol.*, 4(4), 406–425, doi:citeulike-article-id:93683, 1987.
- 654 Schleper, C., Jurgens, G. and Jonuscheit, M.: Genomic studies of uncultivated archaea, *Nat. Rev. Microbiol.*, 3(6), 479–
655 488, doi:10.1038/nrmicro1159, 2005.
- 656 Schouten, S., Hopmans, E. C., Baas, M., Boumann, H., Standfest, S., Könneke, M., Stahl, D. a. and Sinninghe Damsté,
657 J. S.: Intact membrane lipids of “*Candidatus Nitrosopumilus maritimus*,” a cultivated representative of the cosmopolitan
658 mesophilic group I crenarchaeota, *Appl. Environ. Microbiol.*, 74(8), 2433–2440, doi:10.1128/AEM.01709-07, 2008.
- 659 Schouten, S., Middelburg, J. J., Hopmans, E. C. and Sinninghe Damsté, J. S.: Fossilization and degradation of intact
660 polar lipids in deep subsurface sediments: A theoretical approach, *Geochim. Cosmochim. Acta*, 74(13), 3806–3814,
661 doi:10.1016/j.gca.2010.03.029, 2010.
- 662 Schouten, S., Pitcher, A., Hopmans, E. C., Villanueva, L., van Bleijswijk, J. and Sinninghe Damsté, J. S.: Intact polar
663 and core glycerol dibiphytanyl glycerol tetraether lipids in the Arabian Sea oxygen minimum zone: I. Selective
664 preservation and degradation in the water column and consequences for the TEX₈₆, *Geochim. Cosmochim. Acta*, 98,
665 228–243, doi:10.1016/j.gca.2012.05.002, 2012.
- 666 Schouten, S., Hopmans, E. C. and Sinninghe Damsté, J. S.: The organic geochemistry of glycerol dialkyl glycerol
667 tetraether lipids: A review, *Org. Geochem.*, 54, 19–61, doi:10.1016/j.orggeochem.2012.09.006, 2013.
- 668 Schouten, S., Villanueva, L., Hopmans, E. C., van der Meer, M. T. J. and Sinninghe Damsté, J. S.: Are Marine Group II
669 Euryarchaeota significant contributors to tetraether lipids in the ocean?, *Proc. Natl. Acad. Sci. U. S. A.*, 111(41), E4285,
670 doi:10.1073/pnas.1416176111, 2014.
- 671 Schubotz, F., Wakeham, S. G., Lipp, J. S., Fredricks, H. F. and Hinrichs, K.-U.: Detection of microbial biomass by
672 intact polar membrane lipid analysis in the water column and surface sediments of the Black Sea, *Environ. Microbiol.*,
673 11(10), 2720–2734, doi:10.1111/j.1462-2920.2009.01999.x, 2009.
- 674 Sinninghe Damsté, J. S., Schouten, S., Hopmans, E. C., van Duin, A. C. T. and Geenevasen, J. A. J.: Crenarchaeol: the
675 characteristic core glycerol dibiphytanyl glycerol tetraether membrane lipid of cosmopolitan pelagic crenarchaeota, *J.*
676 *Lipid Res.*, 43, 1641–1651, doi:10.1194/jlr.M200148-JLR200, 2002.
- 677 Sinninghe Damsté, J. S., Rijpstra, W. I. C., Hopmans, E. C., Jung, M.-Y., Kim, J.-G., Rhee, S.-K., Stieglmeier, M. and
678 Schleper, C.: Intact Polar and Core Glycerol Dibiphytanyl Glycerol Tetraether Lipids of Group I.1a and I.1b
679 Thaumarchaeota in Soil, *Appl. Environ. Microbiol.*, 78(19), 6866–6874, doi:10.1128/AEM.01681-12, 2012.
- 680 Spang, A., Caceres, E. F. and Ettema, T. J. G.: Genomic exploration of the diversity, ecology, and evolution of the
681 archaeal domain of life, *Science (80-.)*, 357(6351), doi:10.1126/science.aaf3883, 2017.
- 682 Stetter, K. O., Fiala, G., Huber, G., Huber, R. and Segerer, a: Hyperthermophilic microorganisms, *FEMS Microbiol.*
683 *Rev.*, 75, 117–124, doi:10.1111/j.1574-6968.1990.tb04089.x, 1990.
- 684 Sturt, H. F., Summons, R. E., Smith, K., Elvert, M. and Hinrichs, K.-U.: Intact polar membrane lipids in prokaryotes
685 and sediments deciphered by high-performance liquid chromatography/electrospray ionization multistage mass
686 spectrometry—new biomarkers for biogeochemistry and microbial ecology, *Rapid Commun. Mass Spectrom.*, 18(6),
687 617–628, doi:10.1002/rcm.1378, 2004.
- 688 Sugai, A., Sakuma, R., Fukuda, I., Kurosawa, N. and Itoh, Y. H.: The Structure of the Core Polyol of the Ether Lipids
689 from *Sulfolobus acidocaldarius*, *Lipids*, 30(4), 339–344, doi:10.1007/BF02536042, 1995.

690 Teske, A.: Marine deep sediment microbial communities., 2013.

691 Teske, A. and Sørensen, K. B.: Uncultured archaea in deep marine subsurface sediments: have we caught them all?,
692 *ISME J.*, 2(1), 3–18, doi:10.1038/ismej.2007.90, 2008.

693 Villanueva, L., Schouten, S. and Sinninghe Damsté, J. S.: Depth-related distribution of a key gene of the tetraether lipid
694 biosynthetic pathway in marine Thaumarchaeota., *Environ. Microbiol.*, 17(10), 3527–3539, doi:10.1111/1462-
695 2920.12508, 2015.

696 Villanueva, L., Schouten, S. and Sinninghe Damsté, J. S.: Phylogenomic analysis of lipid biosynthetic genes of Archaea
697 shed light on the “lipid divide,” *Environ. Microbiol.*, 19(1), 54–69, doi:10.1111/1462-2920.13361, 2017.

698 Waters, E., Hohn, M. J., Ahel, I., Graham, D. E., Adams, M. D., Barnstead, M., Beeson, K. Y., Bibbs, L., Bolanos, R.,
699 Keller, M., Kretz, K., Lin, X., Mathur, E., Ni, J., Podar, M., Richardson, T., Sutton, G. G., Simon, M., Soll, D., Stetter,
700 K. O., Short, J. M. and Noordewier, M.: The genome of *Nanoarchaeum equitans*: insights into early archaeal evolution
701 and derived parasitism., *Proc. Natl. Acad. Sci. U. S. A.*, 100(22), 12984–8, doi:10.1073/pnas.1735403100, 2003.

702 Wörmer, L., Lipp, J. S., Schröder, J. M. and Hinrichs, K. U.: Application of two new LC-ESI-MS methods for
703 improved detection of intact polar lipids (IPLs) in environmental samples, *Org. Geochem.*, 59, 10–21,
704 doi:10.1016/j.orggeochem.2013.03.004, 2013.

705 Xie, S., Lipp, J. S., Wegener, G., Ferdelman, T. G. and Hinrichs, K.-U.: Turnover of microbial lipids in the deep
706 biosphere and growth of benthic archaeal populations, *Proc. Natl. Acad. Sci.*, 110(15), 6010–6014,
707 doi:10.1073/pnas.1218569110, 2013.

708 Yakimov, M. M., Cono, V. La, Smedile, F., DeLuca, T. H., Juárez, S., Ciordia, S., Fernández, M., Albar, J. P., Ferrer,
709 M., Golyshin, P. N. and Giuliano, L.: Contribution of crenarchaeal autotrophic ammonia oxidizers to the dark primary
710 production in Tyrrhenian deep waters (Central Mediterranean Sea)., *ISME J.*, 5(6), 945–61,
711 doi:10.1038/ismej.2010.197, 2011.

712 Yu, T., Wu, W., Liang, W., Lever, M. A., Hinrichs, K. and Wang, F.: Growth of sedimentary Bathyarchaeota on lignin
713 as an energy source, *Proc. Natl. Acad. Sci.*, 1–6, doi:10.1073/pnas.1718854115, 2018.

714

715

716 **Figure legends**

717 **Fig. 1.** (A) Relative abundances of the IPL-GDGTs (sum of the IPL-types MH, DH and HPH) for the different core
718 GDGTs in the surface (0-0.5 cm) and subsurface sediments (10-12 cm) and (B) the archaeal community composition as
719 revealed by 16S rRNA gene reads (with average abundance above of > 1%) in the surface sediments at 885, 1306,
720 2470, and 3003 mbsl and in the subsurface sediments at 885 and 1306 mbsl.

721 **Fig. 2.** Maximum likelihood phylogenetic tree of the archaeal groups MCG+C3 (modified from Fillol et al., 2015).
722 Extracted OTUs from the Arabian Sea sediments assigned as MCG were inserted in the tree. The number of detected
723 reads per OTU per samples are indicated. Per MCG subgroup the relative abundance is given as detected at the different
724 stations and sediments depths, this is also noted in Table 4. Scale bar represents a 2% sequence dissimilarity.

725 **Fig. 3.** Maximum likelihood phylogenetic tree of MG-I OTUs recovered within the sediment based on the 16S rRNA
726 gene (colored in blue). Sequences from cultured representatives of Thaumarchaeota MG-I are indicated in red.
727 Environmental sequences of MG-I members are indicated in black with their origin specified. The relative abundances
728 of the various OTUs are listed in Table 4. Scale bar represents a 2% sequence dissimilarity.

729 **Fig. 4.** Maximum likelihood phylogenetic tree of *amoA* gene coding sequences recovered from surface (S; 0-0.5 cm)
730 and subsurface (SS; 10-12 cm) sediments (colored in blue) at 885 mbsl, 1306 mbsl and 3003 mbsl (155 clones). *AmoA*
731 gene coding sequences recovered from SPM (colored in orange) at 170 mbsl (28 clones), SPM at 1050 (25 clones)
732 reported by Villanueva et al. (2014). ** indicates *amoA* gene sequences recovered from surface sediments at 3003 mbsl
733 previously reported in Villanueva et al., (2015). Scale bar represents a 2% sequence dissimilarity.

734 **Fig. 5.** Abundance of Thaumarchaeotal 16S rRNA (A,C) and *amoA* (B,D) gene fragment copies per gram of dry weight
735 in the surface sediment (0-0.5 cm) (A,B) and the subsurface sediment (10-12 cm) (C,D). Black bars indicate the amount
736 of DNA 16S rRNA or *amoA* gene fragment copies and the gray bars indicate the RNA (gene transcripts) of 16S rRNA
737 or *amoA* gene fragment copies. Error bars indicate standard deviation based on $n = 3$ experimental replicates.

738

739
740

Table 1. Bottom water temperature and bottom water oxygen (BWO) concentration, oxygen penetration depth (OPD) in the sediment, and TOC content and pore water composition of the surface (0-0.5 cm) sediment^a

Station (mbsl)	T (°C)	BWO ($\mu\text{mol}\cdot\text{L}^{-1}$)	OPD (mm)	TOC (wt %)	NH_4^+ (μM)	NO_2^- (μM)	NO_3^- (μM)	HPO_4^{2-} (μM)
885	10	2.0	0.1	5.6 (\pm 0.2)	2	1.2	1.3	9.2
1306	6.7	14.3	2.9	2.9 (\pm 0.1)	2.6 [*]	0.1 [*]	36.2 [*]	5.6
2470	2.1	63.8	9.8	0.8 (\pm 0.1)	- ^b	-	-	-
3003	1.4	82.9	19	0.7 (\pm 0.1)	55.6	8.3	46.2	3.8

^aData from Kraal et al. (2012) and Lengger et al. (2014)

^b no data available

741

Table 2. Total IPL abundance and heatmap^a of the relative abundance (%) of the detected IPLs and sum (not color coded) per IPL-GDGT in the sediments studied.

Sediment	Depth (mbsl)	GDGT-0						GDGT-1				GDGT-2			
		MH	DH		HCP ^c	HPH	Sum	MH	DH	HPH	Sum	MH	DH	HPH	Sum
		I ^b		II ^b				I ^b				I ^b			
Surface (0-0.5 cm)	885	0.3	ND ^d	ND	ND	ND	0.3	0.1	1.6	ND	1.7	0.1	29.5	ND	29.6
	1306	1.1	ND	ND	ND	36.6	37.6	0.1	1.5	0.2	1.7	ND	15.4	ND	15.4
	2470	0.2	0.1	ND	ND	71.5	71.9	0.0	0.1	0.4	0.5	ND	0.8	ND	0.8
	3003	0.5	0.1	ND	ND	80.3	80.8	ND	0.2	ND	0.2	ND	0.9	ND	0.9
Subsurface (10-12 cm)	885	0.3	ND	7.8	1.6	2.1	11.9	0.1	1.7	0.1	1.9	0.2	27.0	ND	27.1
	1306	2.2	0.9	1.8	0.4	2.1	7.4	0.2	6.7	ND	6.9	0.1	29.7	ND	29.7
	2470	4.3	2.7	ND	ND	18.6	25.6	0.1	5.8	ND	5.9	ND	23.2	ND	23.2
	3003	9.1	3.4	ND	ND	13.0	25.5	0.2	4.3	ND	4.6	ND	21.9	ND	21.9

744

Sediment	Depth (mbsl)	GDGT-3				GDGT-4				Crenarchaeol				IPL abundance [au . g sed dw ⁻¹] ^e	
		MH	DH	HPH	Sum	MH	DH	HPH	Sum	MH	DH	HPH	Sum		
		I ^b				I ^b				I ^b					
Surface (0-0.5 cm)	885	ND	17.8	ND	17.8	ND	6.1	ND	6.1	1.3	43.1	0.3	44.6	2.7E+09	
	1306	0.0	6.9	ND	6.9	ND	2.7	ND	2.7	1.4	15.5	18.7	35.6	1.2E+10	
	2470	ND	0.2	ND	0.2	ND	0.0	ND	0.0	0.2	0.6	25.8	26.6	2.2E+09	
	3003	ND	0.4	ND	0.4	ND	0.0	ND	0.0	0.4	0.2	17.1	17.6	1.3E+10	
Subsurface (10-12 cm)	885	0.1	15.9	ND	15.9	ND	9.4	ND	9.4	1.1	31.1	1.5	33.8	2.0E+09	
	1306	0.0	14.5	ND	14.5	ND	6.1	ND	6.1	2.7	32.4	0.4	35.5	2.2E+09	
	2470	ND	9.6	ND	9.6	ND	2.9	ND	2.9	3.5	28.3	1.0	32.8	7.8E+08	
	3003	ND	9.7	ND	9.7	ND	5.6	ND	5.6	8.2	23.9	0.6	32.7	1.6E+09	

745

^a Green colors indicate a low relative abundance, red colors indicate a high relative abundance

746

^b DH isomers were detected as a GDGT with a glycosidically-bound hexose moiety on both ends of the core (I) and with one glycosidically-bound dihexose moiety on one end (II).

747

^c HCP is an IPL-type with an ether-bound cyclopentanetetraol moiety on one end and an hexose moiety on the other (previously reported as GDNT; e.g. De Rosa and Gambacorta, 1988; Sturt et al., 2004).

748

749

^d ND = not detected

750

^e Response area of summed IPLs given in au (arbitrary units) per gram of dry weight (dw) sediment.

751

752 **Table 3. Relative abundance of IPL-GDGTs grouped by polar head group^a.**

Sample	Depth (mbsl)	MH	DH	HCP	HPH
Surface (0-0.5 cm)	885	1.7%	98.1%	0.0%	0.3%
	1306	2.6%	42.0%	0.0%	55.4%
	2470	0.5%	1.8%	0.0%	97.7%
	3003	0.8%	1.8%	0.0%	97.4%
Subsurface (10-12 cm)	885	1.8%	92.9%	1.6%	3.7%
	1306	5.2%	91.9%	0.4%	2.5%
	2470	7.9%	72.6%	0.0%	19.6%
	3003	17.6%	68.8%	0.0%	13.6%

753 ^aPolar head group types detected: MH = monohexose, DH = dihexose, both isomers combined, HCP = monohexose and

754 cyclopentanetetraol, HPH = monohexose and phosphohexose.

755 ^bND = not detected

756

757 **Table 4. Relative abundance (in %) of MCG- and C3-assigned 16S rRNA gene reads relative to total archaeal**
 758 **reads and distribution (in %) of various subgroups for a station within and a station just below the OMZ**

759

Subgroup	885 mbsl		1306 mbsl	
	Surface (0—0.5 cm)	Subsurface (10-12 cm)	Surface (0-0.5 cm)	Subsurface (10-12 cm)
Total	30.5	47.5	1.3	48.8
1		4.6		8.6
2		9.7		10.9
3		<1		2.3
4		<1		
5b		<1		
8	2.3	33.6		10.3
10		13.4		4.0
12	13.6	7.7		8.0
13		1.2		2.3
14	2.3	3.1		10.9
15	77.3	19.6	100	34.3
17	4.5	5.7		8.6

760

761 **Table 5.** Total Thaumarchaeota MG-I 16S rRNA gene reads and distribution per OTU (%) in surface sediments.

	Depth (mbsl)			
	885	1306	2470	3003
Total reads	0	915	1341	1305
OTU ID #1	n.a. ^a	4.3	2.5	3.0
OTU ID #2	n.a.	3.9	8.1	13.6
OTU ID #3	n.a.	43.6	67.6	61.8
OTU ID #4	n.a.	35.1	1.6	0
OTU ID #5	n.a.	3.3	4.7	2.1

762 ^an.a. = not applicable

763

764 **Fig. 1.**

765

766

767

768

769

770

771

772

773

774

775

776


777

778

779


780


IPL-GDGTs





B

Archaeal community composition









Thaumarchaeota MG-I

0.2

