

**Soil properties determine the elevational patterns of base cations and
micronutrients in plant-soil system up to the upper limits of trees and shrubs**

Ruzhen Wang ^{1,a}, Xue Wang ^{1,a}, Yong Jiang ^{1,*}, Artemi Cerdà ², Jinfei Yin ¹, Heyong

5 Liu ¹, Xue Feng ¹, Zhan Shi ¹, Feike A. Dijkstra ³, Mai-He Li ^{1,4}

¹ State Engineering Laboratory of Soil Nutrient Management, Institute of Applied
Ecology, Chinese Academy of Sciences, Shenyang 110016, China

² Soil Erosion and Degradation Research Group, Department of Geography,
University of Valencia, Valencia, Spain

10 ³ Sydney Institute of Agriculture, School of Life and Environmental Sciences, The
University of Sydney, Sydney, NSW 2006, Australia

⁴ Forest dynamics, Swiss Federal Research Institute WSL, Zuercherstrasse 111,
CH-8903 Birmensdorf, Switzerland

^a These authors contributed equally to this work.

15 * Correspondence: jiangyong@iae.ac.cn; Tel.: +86 24 83970902 (Yong Jiang)

Abstract

To understand whether base cations and micronutrients in the plant-soil system change with elevation, we investigated the patterns of base cations and micronutrients 20 in both soils and plant tissues along three elevational gradients in three climate zones in China. Base cations (Ca, Mg and K) and micronutrients (Fe, Mn and Zn) were determined in soils, trees and shrubs growing at lower and middle elevations as well as at their upper limits on Balang (subtropical, SW China), Qilian (dry-temperate, NW China) and Changbai (wet-temperate, NE China) mountains. No consistent elevational 25 patterns were found for base cation and micronutrient concentrations in both soils and plant tissues (leaves, roots, shoots and stem sapwood). Soil pH, soil organic carbon (SOC), total soil nitrogen (TN), the SOC to TN ratio (C:N), and soil extractable nitrogen (NO_3^- and NH_4^+) determined the elevational patterns of soil exchangeable Ca and Mg and available Fe, Mn and Zn. However, the controlling role of soil pH and 30 SOC was not universal as revealed by their weak correlations with soil base cations under tree canopies at the wet-temperate mountain and with micronutrients under both tree and shrub canopies at the dry-temperate mountain. In most cases, soil base cation and micronutrient availabilities played fundamental roles in determining the base cation and micronutrient concentrations in plant tissues. An exceptional case existed 35 for the decoupling of leaf K and Fe with their availabilities in the soil. Our results highlight the importance of soil physicochemical properties (mainly SOC, C:N and pH) rather than elevation (i.e., canopy cover and environmental factors, especially temperature), in determining base cation and micronutrient availabilities in soils and

subsequently their concentrations in plant tissues.

40 **Key words** base cation, micronutrient, plant tissue, soil physicochemical property,
shrubline, treeline

1 Introduction

Base cations and micronutrients are essential for soil fertility and plant

45 physioecological processes of photosynthesis, metabolism, growth and productivity

(Salisbury and Ross, 1992). For instance, exchangeable calcium (Ca) and magnesium

(Mg) are predominant base cations responsible in buffering soil acidity, and

deficiency of these nutrients can occur in terrestrial ecosystems (Naples and Fisk,

2010; Baribault et al., 2012; Sardans and Peñuelas, 2015), particularly when they are

50 exposed to acid rain. Micronutrient deficiency, on the other hand, occurs more

frequently, for instance, when replenishment of micronutrients via litter

decomposition do not keep pace with output processes of plant uptake and leaching

(White and Zasoski, 1999; Hernandez-Apaolaza, 2014). High soil pH can limit the

availability of micronutrients of iron (Fe), manganese (Mn) and zinc (Zn) (Reisenauer,

55 1988; Lucena, 2000; Rengel, 2007), while low soil pH can induce toxicities of trace

metals constraining terrestrial net primary productivity (He et al., 2005; Reisenauer,

1988; Tian et al., 2016). The concentrations of soil base cations and available

micronutrients were suggested to be positively and negatively correlated with soil pH,

respectively, but both positively correlated with soil organic matter (SOM)

60 concentration (Reisenauer, 1988; Wang et al., 2017). Quantifying base cation and

micronutrient concentrations in soils and plant tissues (leaves, roots, shoots, and stems),

can help understand the nutritional status and potential deficiencies of micronutrients

during plant growth (Richardson, 2004). However, little attention has been paid to

base cation and micronutrient availabilities as well as their inputs and outputs. Under

65 changing environmental conditions (Rengel, 2007).

The plant distribution and growth along elevational gradients reflect changes in environmental conditions (Li et al., 2003, 2004, 2006, 2008a,b; Zhu et al., 2012a,b).

Plants growing at high elevation, especially close to their upper limits, are expected to be highly sensitive to climate change, in particular to global warming (Noble, 1993).

70 Physiological studies of treeline trees have mainly focused on macronutrients such as nitrogen (N), phosphorus (P), and Ca (Richardson, 2004; Liptzin et al., 2013; Mayor et al., 2017), while there are few data available for micronutrients in plant-soil systems along elevational gradients (Wang et al., 2009).

Two hypotheses have been proposed to explain nutrient accumulation and/or nutrient deficiency in plant tissues at high elevations (Oleksyn et al., 2002; Richardson, 2004). First, the decrease in temperature with increasing elevation declines soil microbial activity and plant metabolism, and thus constrains soil nutrient cycling and plant uptake processes (Körner & Paulsen, 2004; Thébaud et al., 2014).

80 At the alpine treeline, low temperature slows down microbial-mediated litter decomposition and thus reduces nutrient supply to plants (van den Driessche, 1974; Richardson, 2004). Second, another paradigm exists that plants retain higher nutrient concentrations in their tissues to maintain metabolic capacity and to avoid cold injury at higher elevations with cold growth conditions (Oleksyn et al., 2002). Whether plants growing at high elevations have higher or lower base cation and micronutrient concentrations is still unclear.

85 The change in environmental conditions along elevational gradients, including

temperature and growing season length (Barry, 1981), provides a unique opportunity to examine the spatial distribution of base cations and micronutrients in plant-soil systems. Uncertainties still exists whether soil properties, plant cover, or

90 environmental factors determine the base cations and micronutrients in plant-soil systems. We, therefore, studied the base cation (Ca, Mg, and K) and micronutrient (Fe, Mn, and Zn) concentrations in plant-soil systems along elevational gradients up to the alpine treeline and shrubline in sub-tropic, dry temperate and wet temperate climate zones in China. Soil base cation and micronutrient concentrations can increase

95 through soil weathering and decomposition of organic matter, but can decrease with plant uptake and loss through leaching. At low elevations, forests have closed

canopies and become increasingly more open with elevation. We therefore

hypothesized that soil base cation and micronutrient concentrations increase with increasing elevation because plant uptake decreases more than the supply through

100 weathering and decomposition with elevation. We also expected that plants at higher elevations would have greater base cation and micronutrient concentrations in their

tissues (leaves, roots, shoots, and stem sapwood) to maintain physio-ecological

processes in a colder environment. To test these hypotheses, we collected soil and

plant samples along three elevational gradients from lower elevations up to the alpine

105 treeline or shrubline in three climate zones in China, and studied the Ca, Mg, K, Fe,

Mn, and Zn concentrations in plant-soil systems.

2 Materials and methods

2.1 Site description and sample collection

110 Study sites were located in three climate zones (summarized in Table S1): Balang mountain with a subtropical climate located in Wolong Nature Reserve (“subtropical Mt. ”, 102 °52'-103 °24'E, 30 °45'-31 °25'N) in southwestern China, Qilian mountain located in the dry-temperate climate zone (“dry-temperate Mt. ”, 102 °58'-103 °01'E, 37 °14'-37 °20'N) in northwestern China, and Changbai mountain with a wet-temperate 115 climate (“wet-temperate Mt.”, 126 °55'-129 °00'E, 41 °23'-42'36 °N) located in northeastern China (see Fig. S1). Three distinct sites were chosen to find the general patterns for base cations and micronutrients along elevational gradients across climate scales rather than to investigate the comparability among study sites. The subtropical Mt. is influenced by warm-wet monsoon masses in summer and continental air masses 120 in winter (Li et al., 2012). The mean annual precipitation (MAP) of the subtropical Mt. is about 846 mm monitored by Dengsheng Meteorological Station at 2730 m (Li et al., 2012). For the dry-temperate Mt., the MAP is 435 mm, which is monitored by the Qilian weather station at 2787 m altitude (Qiang et al., 2003). The wet-temperate Mt. 125 is located in a typical continental temperate monsoon climate zone with MAP increasing from 632 to 1154 mm along the elevational gradient from 530 to 2200 m (Shen et al., 2013).

In this study, the alpine treeline and shrubline are defined as the upper limit of 130 obvious trees and shrubs, respectively. The trees that were investigated are *Abies faxoniana* (elevation range 2860-3670 m) for the subtropical Mt., *Picea crassifolia* (elevation range 2540-3250 m) for the dry-temperate Mt., and *Betula ermanii*

(elevation range 1700-2030 m) for the wet-temperate Mt.. The shrubs are *Quercus aquifolioides* (elevation range 2840-3590 m) for the subtropical, *Salix gilashanica* (elevation range 3020-3540 m) for the dry-temperate, and *Vaccinium uliginosum* (elevation range 1430-2380 m) for the wet-temperate Mts. The targeted treeline trees
135 and shrubline shrubs are dominant and common species for each study site. The soils from the three sampling sites of subtropical, dry-temperate, and wet-temperate Mts. were classified as Umbric Cryic Cambisols, Calcaric Ustic Cambisols and Andic Gelic Cambisols, respectively (IUSS Working Group WRB, 2014).

Plant tissue samples of current-year mature leaves, roots (< 2 mm), stem sapwood, 140 and shoots (twigs) from trees and shrubs were collected at lower and middle elevations, as well as at the upper limits. At each elevation, 6 plots (10 m × 10 m) were selected to serve as 6 replicates on southern slopes. Within each plot, 6-10 tree or shrubs of similar height were randomly selected for tissue sampling. Soils (0-10 cm) were directly collected under the canopy of trees or shrubs sampled for each plot 145 using a 3-cm diameter corer. Both plant and soil samples were homogenized and composited within each plot. Samples were collected at the middle of July for subtropical Mt., at the beginning of August for dry-temperate Mt., and at end of August for wet-temperate Mt. in 2014. The main characteristics of the three study sites are summarized in Table S1.

150

2.2 Chemical analysis

The soil samples were separated into two subsamples with one subsample being

air-dried to constant weight and the other one stored at 4 °C for further analyses. For subtropical and wet-temperate Mts., soil organic carbon (SOC) and total nitrogen (TN)

155 was determined on ground soils using an elemental analyzer (Vario MACRO Cube,

Elementar, Germany). For dry-temperate Mt., the ground soil samples were treated

with 12 M HCl according to Wang et al. (2015) to remove inorganic C before organic

C determination on the elemental analyzer. Soil NO_3^- -N and NH_4^+ -N were extracted

from fresh soils with a 2 M KCl solution and measured using an AutoAnalyser III

160 continuous Flow Analyzer (Bran & Luebbe, Norderstedt, Germany). Soil total

inorganic nitrogen (TIN) was the sum up of extractable NO_3^- -N and NH_4^+ -N. Soil

Olsen phosphorus (P) was quantified by colorimetric analysis after extraction with a

0.5 M NaHCO_3 solution (Olsen et al., 1954).

A subsample of 5 g soil was used to determine soil pH in a 1:5 (w/v) soil-to-water

165 extract. Soil exchangeable base cations were extracted with a 1 M ammonium acetate

solution (Wang et al., 2017). Soil available micronutrients were extracted by

diethylenetriamine pentaacetic acid (DTPA) according to Lü et al. (2016). Briefly, 10

g of soil was extracted by 20 ml 0.005 M DTPA + 0.01 M CaCl_2 + 0.1 M TEA

(triethanolamine) (pH 7.0). The soil-solution suspension was shaken for 2 h at 180

170 rpm and then filtered through ash-free filter paper. The concentrations of base cations

and micronutrients were determined using an atomic absorption spectrometer (AAS,

Shimadzu, Japan).

Plant samples of leave¹, roots, shoots and stem sapwood were oven-dried² and ground for base cation and micronutrient analyses. Root samples were washed prior to

175 being oven-dried. To determine total base cation and trace element concentrations, 0.2 g plant samples were digested with a mixture of acids of HNO_3 and HClO_4 (5:1, v/v) on a hot plate. After the mixture turned into clear solution, the digests were decanted into a 50 ml volumetric flasks and the volume was adjusted into 50 ml. The concentrations of Ca, Mg, K, Fe, Mn, and Zn were determined by the AAS (Shimadzu, 180 Japan).

2.3 Statistical analyses

Normality of data was determined using the Kolmogorov-Smirnov test, and homogeneity of variances using Levene's test. Two-way ANOVAs were executed to 185 determine the effects of plant life form (tree or shrub), elevation position and their interactions on soil pH, SOC, soil exchangeable base cations and available micronutrients, and total base cations and micronutrients in plant tissues. We assigned sampling site as a random factor in the statistics.  this study aimed to test the general elevational patterns instead of site-specific heterogeneity of base cations and 190 micronutrients in plant-soil system across three sites. Within each site, the effect of elevation on measured parameters was determined by multiple comparison  with a Tukey design for soils and each life form. Pearson correlation analysis was used to determine the relationships between measured parameters. All statistical analyses were performed in SPSS 16.0 (SPSS, Inc., Chicago, IL, USA) and statistical 195 significance was accepted at $P < 0.05$.

3 Results

3.1 Soil pH and SOC

Soil pH was significantly different among elevational positions (Table 1). For both

200 subtropical and dry-temperate Mt.  soil pH decreased with increasing elevation under tree canopy, while it was the opposite trend under shrub canopy (Fig. 1a). For wet-temperate Mt., the upper limit of shrubs had significantly higher soil pH (Fig. 1a).

For all three sites, SOC concentration showed a hump-shaped trend with the highest value at the middle elevation under tree canopy (Fig. 1b). Under shrubs, SOC concentration significantly increased with increasing elevation for subtropical and wet-temperate Mts., while it was the lowest at the upper limit of shrubs for dry-temperate Mt. (Fig. 1b).

3.2 Changes in soil base cations and available micronutrients

210 Soil exchangeable Ca and Mg decreased with increasing elevation under tree canopy of subtropical and wet-temperate Mts. and under shrubs of dry-temperate Mt. (not for Mg) (Fig. 2a,b). However, they showed the opposite trend under shrubs of subtropical (not for Mg) and wet-temperate Mts. and under trees of dry-temperate Mt. (Fig. 2a,b). Soil exchangeable K decreased with increasing elevation under tree and shrub canopies at subtropical Mt. and under trees at dry-temperate Mt. (Fig. 2c).

215 Soil available Fe was significantly affected by elevation position (Table 1, Fig. 2d). The upper limit had the lowest concentration under both tree and shrub canopies for subtropical Mt. and under shrub canopy for wet-temperate Mt. (Fig. 2d). For

dry-temperate Mt., soil available Fe significantly increased with increasing elevation
220 under both tree and shrub canopies (Fig. 2a). For subtropical Mt., soil available Mn
was significantly higher at the middle elevation under tree canopies. Soil available Mn
decreased with increasing elevation under shrub and tree canopies for subtropical and
wet-temperate Mts., respectively, while it showed the opposite trend under both tree
and shrub canopies of dry-temperate Mt. and under shrubs of wet-temperate Mt. (Fig.
225 2e). Soil available Zn was significantly affected by plant life form and the interactive
effect between life form and elevation position (Table 1). Specifically, soils at the
upper limit had the highest available Zn under shrubs at wet-temperate Mt..

3.3 Base cations in plants

230 For subtropical Mt., a significant decrease of Ca concentration was detected in the
leaves of trees and shrubs (Fig. 3a), roots and shoots of trees (Fig. 3b,c), and stem
sapwood of shrubs (Fig. 3d). For dry-temperate Mt., Ca concentration decreased with
increasing elevation in roots, shoots and stem sapwood of trees (Fig. 3b,c,d), and in
shoots and stem sapwood of shrubs (Fig. 3c,d). For wet-temperate Mt., shoot Ca
235 concentration decreased with increasing elevation for trees (Fig. 3c). Along with
increasing elevation, a significant decrease of Mg was found in shrub leaves, tree
roots, shrub shoots and stem sapwood at sub-tropical Mt. (Fig. 3e,f,g,h), and in roots,
shoots and stem sapwood of trees at dry-temperate Mt. (Fig. 3f,g,h), and in leaves and
shoots of both trees and shrubs at wet-temperate Mt. (Fig. 3e,g). With the increase in
240 elevation, K concentration significantly decreased in leaves of trees, roots and stem

sapwood of both trees and shrubs at subtropical Mt. (Fig. 3i,j,l), in tree shoots of dry-temperate Mt. (Fig. 3k), and in leaves of both trees and shrubs at wet-temperate Mt. (Fig. 3i).

245 **3.4 Micronutrients in plants**

For subtropical Mt., Fe concentrations in leaves (Fig. 4a) and roots (Fig. 4b) showed a similar trend with soil available Fe, with the highest values at the middle elevation for both trees and shrubs. For dry-temperate Mt., the highest Fe concentrations were found at the lower elevation in leaves (Fig. 4a), roots (Fig. 4b), shoots (Fig. 4c) and 250 stem sapwood (Fig. 4d) for trees and in shoots (Fig. 4c) and stem sapwood (Fig. 4d) for shrubs. For wet-temperate Mt., Fe concentration was the highest in tree shoots at the middle elevation (Fig. 4c), in shrub leaves at lower elevation (Fig. 4a), and in roots, shoots and stem sapwood of shrubs at the upper limit of trees (Fig. 4b,c,d).

The Mn concentration decreased with increasing elevation in leaves and shoots of 255 both trees and shrubs at subtropical Mt. (Fig. 4e,g), in stem sapwood of shrubs at both subtropical and dry-temperate Mts. (Fig. 4h). The Mn concentration increased with increasing elevation in leaves of trees at dry-temperate Mt. (Fig. 4e), in roots of both trees and shrubs at wet-temperate Mt. (Fig. 4f), in shoots of shrubs and stem sapwood of trees and shrubs at wet-temperate Mt. (Fig. 4g,h).

260 The Zn concentration was the highest at middle elevation for trees in leaves at wet-temperate Mt. (Fig. 4i), in roots at dry-temperate Mt. (Fig. 4j), in shoots at wet-temperate Mt. (Fig. 4k) and in stem sapwood at subtropical Mt. (Fig. 4l). With the

increase in elevation, a decrease of Zn concentration was found in roots of trees at subtropical Mt. (Fig. 4j) and in stem sapwood of shrubs at dry-temperate Mt. (Fig. 4l); 265 however, an increase of Zn was found in shrub roots at wet-temperate Mt., in shoots of trees at dry-temperate Mt. and shrubs at wet-temperate Mt. (Fig. 4k), and in stem sapwood of trees at dry-temperate Mt. and shrubs at wet-temperate Mt. (Fig. 4l).

3.5 Correlations

270 Across all sampling sites and plant life forms, both soil exchangeable Ca and Mg were positively correlated with soil pH (Fig. S2a,b) and TN (Table 2), while they were negatively correlated with soil C:N, NO_3^- , and NH_4^+ (Table 2). For wet-temperate Mt., both soil pH and SOC showed no relationship with soil exchangeable Ca and Mg under tree canopies, although SOC was positively related to exchangeable K (Table 275 S2). Negative correlations were found for both Mg and K concentrations between stems and leaves (both $p < 0.01$; Table S3). However, Mg and K concentrations in roots showed no correlation with that in leaves (Table S3).

When analyzing data across sampling sites and plant life forms, soil available Fe, Mn and Zn were negatively correlated with soil pH ($p < 0.01$; Table 2, Fig. S2c,d,e), 280 and soil available Fe and Zn were positively correlated with SOC ($p < 0.01$; Table 2). However, available micronutrients had no relationships with both soil pH and SOC at dry-temperate Mt., except for a positive correlation between soil pH and available Mn under shrub canopies (Table S2). For both Mn and Zn concentrations, significant and positive correlations were found between soil and plant tissues (Table S3). Soil

285 available Fe was negatively correlated with Fe concentrations in shoots and stems
(both $p < 0.01$; Table S3).

4 Discussion

4.1 Elevational patterns of base cations and available micronutrients in soils and 290 relationships with pH and SOC

Contrary to our first hypothesis, no consistent elevational patterns were detected for
soil exchangeable base cations and available micronutrients under either trees or
shrubs. Inconsistent elevational patterns of soil base cations and available
micronutrients indicated that plant uptake of these nutrients did not necessarily
295 decrease more than nutrient supply at higher elevation due to more open canopies.

Our results suggest that soil physiochemical parameters were the dominant
contributors and more important than environmental gradients affecting elevational
patterns of soil exchangeable base cations and available micronutrients (Fig. S2, Table
2). For instance, soil available Fe, Mn and Zn followed patterns of SOC under trees
300 along the elevational gradient at subtropical Mt. (Table S2), while for shrubs at
subtropical Mt., soil pH, instead of SOC, regulated elevational patterns of soil
available Fe, Mn and Zn (Table S2). Our findings are consistent with a vast amount of
previous studies confirming the pivotal role of soil pH and SOC concentration in
determining soil base cation and micronutrient availabilities (Sharma et al., 2004; Lü
305 et al., 2016; Wang et al., 2017). However, the fundamental roles of SOC and soil pH
in controlling soil base cation and micronutrient availabilities, was not universal as

suggested by the relatively weak relationships of soil pH and SOC with soil base cations under tree canopies at wet-temperate Mt. and with micronutrients under both tree and shrub canopies at dry-temperate Mt. (Table S2). This could indicate species-
310 and life form-specific effects on soil base cation and micronutrient availabilities.

Other soil parameters, such as C:N and extractable NO_3^- and NH_4^+ also influenced the availability of base cations (Table 2). Soil C:N ratio serves as an indicator of SOM decomposition status where more decomposed SOM possesses a lower C:N ratio (Sollins et al., 2009) and a higher content of negatively charged functional groups (i.e. phenolic, carboxyl, and hydroxyl groups) (Haberhauer et al.,
315 1998). In this study, negative correlations between soil C:N and base cations (Table 2) suggest that more decomposed SOM is beneficial for the retention of soil base cations. Furthermore, soil with a higher level of extractable NO_3^- predisposes cations to leach accompanied by loss of NO_3^- (Cremer and Prietzel, 2017). Therefore, significant
320 negative correlations were detected between soil NO_3^- and base cations in this study (Table 2). Soil extractable NH_4^+ was also negatively correlated to exchange with Ca and Mg, possibly because NH_4^+ can exchange with base cations on surface soil colloids into soil solution thereby enhancing their loss (Wang et al., 2015; Cusack et al., 2016).

325 A negative correlation between soil pH and soil available micronutrients (Table 2) might be due to precipitation of micronutrient cations at higher soil pH (Rengel, 2007). Indeed, solubility of micronutrients was suggested to decrease from 100-fold (for Mn and Zn) to 1000-fold (for Fe) with one-unit increase of soil pH (Rengel, 2001). Soil

organic matter plays an important role in micronutrient retention due to its negative charge (He et al., 2005; Wang et al., 2015, 2017). This may be a reason for the positive relationships between SOC and micronutrients (although not always significant, Table 2). While no general patterns were found for distribution of micronutrients under both tree and shrub canopies with elevation, our results suggest that the determinants of soil micronutrient availabilities were soil pH and SOC concentration, which are reflections of long-term climatic conditions, plant-soil interactions, and biogeochemical processes (Sinsabaugh et al., 2008).

4.2 Elevational patterns of base cations and micronutrients in plants and plant-soil system

In contrast to our second hypothesis, both trees and shrubs at higher elevation did not necessarily contain higher base cation and micronutrient concentrations in their tissues. No general patterns were found for base cations and micronutrients in both trees and shrubs along elevational gradients across the three sites (Fig. 3,4). Even normalizing the data to per unit concentration of soil available nutrients, there were still no consistent elevational patterns for both base cations and micronutrients in plant tissues (Fig. S3). This suggests that base cation and micronutrient concentrations in plants are influenced by other factors besides elevation-induced changes in temperature, precipitation, specific nutrient absorption characteristics of different tissues, soil base cation and micronutrient availabilities and other edaphic properties (Campo-Alves, 2003; Richardson, 2004). Another explanation could be that initial

differences in soil properties (*e.g.*, parent material) among climate zones were larger than the effects of elevation. Soil base cation and micronutrient availabilities were an important factor influencing their concentrations in plant tissues across all plant species and sampling sites (Table S3). Similar results were found for macronutrients (i.e. nitrogen and phosphorus) suggesting that “plants are what they root in” (Elser et al., 2010; Han et al., 2014). However, plant nutrients did not covary with soil nutrients along a 2200 km-long climatic gradient in grasslands of northern China (Luo et al., 2015, 2016). The discrepancy of our study with Luo et al. (2015, 2016) might be driven by different ecosystem types (forest vs. grassland), dominant climatic factor gradients (temperature vs. precipitation), and different soil properties. The studies of Luo et al. (2015, 2016) were conducted in grassland ecosystems where precipitation played an essential role in nutrient concentrations in plant-soil systems. Moreover, base cation and micronutrient cycling processes are likely to be different between high-organic and fine-grained forest soils in our study versus low-organic and sandy grassland soils in Luo et al. (2016). Inconsistently elevational patterns in plant nutrient concentrations could also be derived from the fact that individual plant species reinforced patterns of soil nutrient availabilities in their vicinity causing a positive feedback between plant and soil (Hobbie, 1992).

The topic of base cation and micronutrient translocation in intact plant is important as it deals with the movement of micronutrients from root to the leaves for physiological activities, such as photosynthesis (Welch and Shuman, 1995). Also, it is an important process in determining plant chemical composition and subsequently

litter quality, litter decomposition and nutrient release (Sun et al., 2016). Given earlier findings that transport of base cations from roots to the leaves in woody plants is slow (van der Heijden et al., 2015), we found no significant correlation for both Mg and K between roots and leaves (Table S3). However, negative relationships of stem Mg vs. leaf Mg and stem K vs. leaf K suggest that the plant internal pools of base cations could act as sources of base cation supply for leaves (Weatherall et al., 2006).

Translocation of base cations within plant tissues is one of the main physiological

mechanisms buffering low nutrient availabilities in soils (van der Heijden et al., 2015).

For instance, supplementation of Mg is a critical process to maintain photosynthesis in

forests growing on acid and cation poor soils (Verbruggen and Hermans, 2013). In

support of this, we found significantly positive correlations between Mg and soluble

sugar concentrations (one of the main photosynthate) in leaves across the three sites

(Fig. S4a), while relationships were more pronounced at wet-temperate Mt. (Fig. S4b)

where soil pH and exchangeable Mg was the lowest (Fig. 1a, 2b).

Unlike Ca, Mg, Mn and Zn, the concentrations of K and Fe in plant leaves

decoupled with their availabilities in the soil (Table S3), which may suggest that not

only availability of these nutrients in soils affect their leaf concentrations, but that also

other environmental factors (*e.g.*, temperature) played more important roles in

affecting plant nutrition (van den Driessche, 1974). We do not know why this

decoupling only occurred for K and Fe, but possibly factors such as temperature

constrained soil microbial activity and plant metabolism (Körner and Paulsen, 2004)

and subsequently uptake of these nutrients by plants. On the other hand, plants often

395 increase nutrient uptake to compensate for decreased metabolism at low temperature (Reich and Oleksyn, 2004). Thus, these opposite effects of temperature on K and Fe concentrations in plant tissues may have obscured their relationships with K and Fe availability in the soil along the elevational gradients. While plant nutrient concentrations were mainly influenced by nutrient availabilities in the soil and by 400 plant-internal translocation processes, we found no consistent evidence that plants accumulate more base cations and micronutrients in their tissues to better adapt to cold environments at higher elevation.

5 Conclusions

405 We did not find consistent elevational patterns of base cations and micronutrients in plant-soil systems along three different elevation transects up to the alpine treeline and shrubline in different regions of China. Rather, our results highlight the essential roles of specific edaphic properties of soil pH, SOC and extractable nitrate and ammonium in regulating soil base cation and micronutrient availabilities across 410 climate zones. Soil available base cations and micronutrients were mostly positively correlated with concentrations of base cations and micronutrients in plant tissues, except for K and Fe. Our results suggest that base cation and micronutrient concentrations in plants (trees and shrubs) growing at their upper limits are largely controlled by their availabilities in the soil rather than by plant adaptations to cold 415 environment at higher elevations.

Acknowledgments

This was financially supported by the National Natural Science Foundation of China (41371076, 31570501) and the Strategic Priority Research Program of the Chinese

420 Academy of Sciences (XDB15010302).

Data Availability

Data sets for this paper can be obtained via personal communication.

425 **Author contributions**

Mai-He Li designed the study; Xue Wang, Heyong Liu, Jinfei Yin and Zhan Shi did

the plant and soil measurements; Ruzhen Wang and Xue Wang analyzed the data;

Ruzhen Wang wrote the manuscript; Mai-He Li, Feike A. Dijkstra and Artemi Cerdà

revised the manuscript; Mai-He Li and Yong Jiang provided funding and laboratory

430 facilities for this study.

References

Barry, R. G.: Mountain weather and climate. Methuen, London and New York, 1981.

Baribault, T. W., Kobe, R. K., Finley, A. O.: Tropical tree growth is correlated with

435 soil phosphorus, potassium, and calcium, though not for legumes. *Ecol. Monogr.*,

82, 189-203, doi:10.1890/11-1013.1, 2012.

Campo-Alves J.: Nutrient availability and fluxes along a toposequence with tropical

dry forest in Mexico. *Agrociencia*, 37, 211-219, 2003.

Cremer, M., and Prietzel, J.: Soil acidity and exchangeable base cation stocks under
440 pure and mixed stands of European beech, Douglas fir and Norway spruce. *Plant
Soil*, 415, 393-405, doi:10.1007/s11104-017-3177-1, 2017.

Cusack, D. F., Macy, J., and McDowell, W. H.: Nitrogen additions mobilize soil base
cations in two tropical forests. *Biogeochemistry*, 128, 67-88,
doi:10.1007/s10533-016-0195-7, 2016.

445 Elser, J. J., Fagan, W. F., Kerkhoff, A. J., Swenson, N. G., and Enquist, B. J.:
Biological stoichiometry of plant production: metabolism, scaling and ecological
response to global change. *New Phytol.*, 186, 593-608,
doi:10.1111/j.1469-8137.2010.03214.x, 2010.

Han, X., Sistla, S.A., Zhang, Y., Lü, X., and Han, X.: Hierarchical responses of plant
450 stoichiometry to nitrogen deposition and mowing in a temperate steppe. *Plant Soil*,
382, 175-187, doi:10.1007/s11104-014-2154-1, 2014.

He, Z., Yang, X., and Stoffella, P. J.: Trace elements in agroecosystems and impacts
on the environment. *J. Trace Elem. Med. Biol.*, 19, 125-140,
doi:10.1016/j.jtemb.2005.02.010, 2005.

455 Hernandez-Apaolaza, L.: Can silicon partially alleviate micronutrient deficiency in
plants? A review. *Planta*, 240, 447-458, doi:10.1007/s00425-014-2119-x, 2014.

IUSS Working Group WRB: World Reference Base for Soil Resource 2014.
International soil classification system for naming soils and creating legends for
soil maps. *World Soil Resources Reports No. 106*, FAO, Rome, 2014.

460 Kochian, L. V.: Mechanisms of micronutrient uptake, translocation, and interactions

in plants. In: Mortvedt JJ, Cox FR, Shuman LM, Welch RM (eds) *Micronutrients in Agriculture*. Soil Science Society of America, Madison pp 229-296, 1991.

Körner, C., and Paulsen, J.: A world-wide study of high altitude treeline temperatures. *J. Biogeogr.*, 31, 713-732, doi:10.1111/j.1365-2699.2003.01043.x, 2004.

465 Laudicina, V. A., Novara, A., Barbera, V., Egli, M., and Badalucco, L.: Long-term tillage and cropping system effects on chemical and biochemical characteristics of soil organic matter in a Mediterranean semiarid environment. *Land Degrad. Dev.*, 26, 45-53, doi:10.1002/ldr.2293, 2015.

Li, M.-H., Yang, J., and Kräuchi, N.: Growth responses of *Picea abies* and *Larix decidua* to elevation in subalpine areas of Tyrol, Austria. *Can. J. For. Res.*, 33, 653-662, doi:<https://doi.org/10.1139/x02-202>, 2003.

470 Li, M.-H., and Yang, J.: Effects of elevation and microsite on growth of *Pinus cembra* in the subalpine zone of the Austrian Alps. *Ann. For. Sci.*, 61, 319-325, doi:10.1051/forest:2004025, 2004.

475 Li MH, Kräuchi N, and Dobbertin M.: Biomass distribution of different-aged needles in young and old *Pinus cembra* trees at highland and lowland sites. *Trees*, 20, 611-618, <https://doi.org/10.1007/s00468-006-0076-0>, 2006.

Li, M.-H., Xiao, W., Shi, P., Wang, S., Zhong, Y., Liu, X., Wang, X., Cai, X., and Shi, Z.: Nitrogen and carbon source-sink relationships in trees at the Himalayan treelines compared with lower elevations. *Plant Cell Environ.*, 31, 1377-1387, doi:10.1111/j.1365-3040.2008.01848.x, 2008.

480 Li, M.-H., Xiao, W., Wang, S., Cheng, G., Cherubini, P., Cai, X., Liu, X., Wang, X.,

and Zhu, W.: Mobile carbohydrates in Himalayan treeline trees I. Evidence for carbon gain limitation but not for growth limitation. *Tree Physiol.*, 28, 1287-1296, 485 doi:<https://doi.org/10.1093/treephys/28.8.1287>, 2008.

Li, Z., Liu, G., Fu, B., Hu, C., Luo, S., Liu, X., and He, F.: Anomalous temperature-growth response of *Abies faxoniana* to sustained freezing stress along elevational gradients in China's Western Sichuan Province. *Trees*, 26, 1373-1388, doi:10.1007/s00468-012-0712-9, 2012.

490 Liptzin, D., Sanford, Jr. R. L., and Seastedt, T. R.: Spatial patterns of total and available N and P at alpine treeline. *Plant Soil*, 365, 127-140, doi:10.1007/s11104-012-1379-0, 2013.

Lucena, J.J.: Effects of bicarbonate, nitrate and other environmental factors on iron deficiency chlorosis: A review. *J. Plant Nutri.*, 23, 1591-1606, 495 doi:10.1080/01904160009382126, 2000.

Luo, W., Elser, J. J., Lü, X., Wang, Z., Bai, E., Yan, C., Wang, C., Li, M.-H., Zimmermann, N. E., Han, X., Xu, Z., Li, H., Wu, Y., and Jiang, Y.: Plant nutrients do not covary with soil nutrients under changing climatic conditions. *Global Biogeochem. Cy.*, 29, 1298-1308, doi:10.1002/2015GB005089, 2015.

500 Luo, W., Sardans, J., Dijkstra, F. A., Peñuelas, J., Lü, X., Wu, H., Li, M.-H., Bai, E., Wang, Z., Han, X., and Jiang, Y.: Thresholds in decoupled soil-plant elements under changing climatic conditions. *Plant Soil*, 409, 159-173, doi:10.1007/s11104-016-2955-5, 2016.

Lü, L., Wang, R., Liu, H., Yin, J., Xiao, J., Wang, Z., Zhao, Y., Yu, G., Han, X., and

505 Jiang, Y.: Effect of soil coarseness on soil base cations and available micronutrients
in a semi-arid sandy grassland. *Solid Earth*, 7, 549-556, doi:10.5194/se-7-549-2016,
2016.

Martens, D. C., and Westermann, D. T.: Fertilizer applications for correcting
510 micronutrient deficiency. In: Mortvedt JJ, Cox FR, Shuman LM, Welch RM (eds)
Micronutrients in agriculture. Soil Science Society of America, Madison, pp
549-592, 1991.

Mayor, J. R., Sanders, N. J., Classen, A. T., Bardgett, R. D., Clément, J. C., Fajardo,
A., Lavorel, S., Sundqvist, M. K., Bahn, M., Chisholm, C., Cieraad, E., Gedalof, Z.,
Grigulis, K., Kudo, G., Oberski, D. L., and Wardle, D. A.: Elevation alters
515 ecosystem properties across temperate treelines globally. *Nature*, 542, 91-95,
doi:10.1038/nature21027, 2017.

McLaughlin, M. J., Parker, D. R., and Clarke, J. M.: Metals and micronutrients-food
safety issues. *Field Crop Res.*, 60, 143-163,
doi:[https://doi.org/10.1016/S0378-4290\(98\)00137-3](https://doi.org/10.1016/S0378-4290(98)00137-3), 1999.

Naples, B. K., Fisk, M. C.: Belowground insights into nutrient limitation in northern
520 hardwood forests. *Biogeochemistry*, 97, 109-121, doi:10.1007/s10533-009-9354-4,
2010.

Noble, I. R.: A model of the responses of ecotones to climate change. *Ecol. Appl.*, 3,
396-403, doi:10.2307/1941908, 1993.

Oleksyn, J., Reich, P. B., Zytkowiak, R., Karolewski, P., and Tjoelker, M. G.: Needle
525 nutrients in geographically diverse *Pinus sylvestris* L. populations. *Ann. For.*

Science, 59, 1-18, doi: <https://doi.org/10.1051/forest:2001001>, 2002.

Olsen, S. R., Watanabe, F. S., Cosper, H. R., Larson, W. E., and Nelson, L. B.:
Residual phosphorus availability in long-term rotations on calcareous soils. Soil
530 Sci., 78, 141-152, 1954.

Qiang, W., Wang, X., Chen, T., Feng, H., An, L., He, Y., and Wang, G.: Variations of
stomatal density and carbon isotope values of *Picea crassifolia* at different altitudes
in the Qilian Mountains. Trees, 17, 258-262, doi:10.1007/s00468-002-0235-x,
2003.

535 Reisenauer, H. M.: Determination of plant-available soil manganese. In: Graham RD,
Hannam RJ, Uren NC (eds) Manganese in soils and plants. Kluwer Academic
Publisher, Dordrecht, pp 87-98, 1988.

Reich, P. B., and Oleksyn, J.: Global patterns of plant leaf N and P in relation to
temperature and latitude. P. Natl. Acad. Sci. USA, 101, 11001-11006,
540 doi:10.1073/pnas.0403588101, 2004.

Rengel, Z.: Genotypic differences in micronutrient use efficiency in crops. Commun.
Soil Sci. Plant Anal., 32, 1163-1186, doi:<http://dx.doi.org/10.1081/CSS-100104107>,
2001.

Rengel, Z.: Cycling of micronutrients in terrestrial ecosystem. In: Marschner P,
545 Rengel Z (eds) Nutrient cycling in terrestrial ecosystems. Springer-Verlag, Berlin,
pp 93-113, 2007.

Richardson, A. D.: Foliar chemistry of balsam fir and red spruce in relation to
elevation and the canopy light gradient in the mountains of the northeastern United

States. *Plant Soil*, 260, 291-299, doi:10.1023/B:PLSO.0000030179.02819.85, 2004.

550 Salisbury, F. B., and Ross, C. W.: Mineral nutrition. In: *Plant Physiology*, Wadsworth Publishing Company, Belmont, pp 116-135, 1992.

Sardans, J., Peñuelas, J.: Potassium: a neglected nutrient in global change. *Global Ecol. Biogeogr.*, 24, 261-275, doi:10.1111/geb.12259, 2015.

Sharma, B. D., Arora, H., Kumar, R., and Nayyar, V. K.: Relationships between soil characteristics and total and DTPA-extractable micronutrients in Inceptisols of

555 Punjab. *Commun. Soil Sci. Plant Anal.*, 35, 799-818, doi:<http://dx.doi.org/10.1081/CSS-120030359>, 2004.

Shen, C., Xiong, J., Zhang, H., Feng, Y., Lin, X., Li, X., Liang, W., and Chu, H.: Soil pH drives the spatial distribution of bacterial communities along elevation on

560 Changbai Mountain. *Soil Biol. Biochem.*, 57, 204-211, doi:<http://dx.doi.org/10.1016/j.soilbio.2012.07.013>, 2013.

Sinsabaugh, R. L., Lauber, C. L., Weintraub, M. N., Ahmed, B., Allison, S. D., Crenshaw, C., Contosta, A. R., Cusack, D., Frey, S., Gallo, M. E., Gartner, T. B., Hobbie, S. E., Holland, K., Keeler, B. L., Powers, J. S., Stursova, M.,

565 Takacs-Vesbach, C., Waldrop, M. P., Wallenstein, M. D., Zak, D. R., and Zeglin, L. H.: Stoichiometry of soil enzyme activity at global scale. *Ecol. Lett.*, 11, 1252-1264, doi:10.1111/j.1461-0248.2008.01245.x, 2008.

Sollins, P., Kramer, M. G., Swanston, C., Lajtha, K., Filley, T., Aufdenkampe, A. K., Wagai, R., and Bowden, R. D.: Sequential density fractionation across soils of

570 contrasting mineralogy: evidence for both microbial-and mineral-controlled soil

organic matter stabilization. *Biogeochemistry*, 96, 209-231,

doi:10.1007/s10533-009-9359-z, 2009.

Sun, T., Dong, L., Zhang, L., Wu, Z., Wang, Q., Li, Y., Zhang, H., and Wang, Z.: Early stage fine-root decomposition and its relationship with root order and soil depth in a

575 *Larix gmelinii* plantation. *Forests*, 7, 234, doi:10.3390/f7100234, 2016.

Thébault, A., Clément, J. C., Ibanez, S., Roy, J., Geremia, R. A., Pérez, C. A., Buttler, A., Estienne, Y., and Lavorel, S.: Nitrogen limitation and microbial diversity at the treeline. *Oikos*, 123, 729-740, doi:10.1111/j.1600-0706.2013.00860.x, 2014.

Tian, Q., Liu, N., Bai, W., Li, L., Chen, J., Reich, P. B., Yu, Q., Guo, D., Smith, M. D.,

580 Knapp, A. K., Cheng, W., Lu, P., Gao, Y., Yang, A., Wang, T., Li, X., Wang, Z., Ma, Y., Han, X., and Zhang, W.: A novel soil manganese mechanism drives plant species loss with increased nitrogen deposition in a temperate steppe. *Ecology*, 97, 65-74, doi:10.1890/15-0917.1, 2016.

van den Driessche, R.: Prediction of mineral nutrient status of trees by foliar analysis.

585 *Bot. Rev.*, 40, 347-394, doi:10.1007/BF02860066, 1974.

van der Heijden, G., Dambrine, E., Pollier, B., Zeller, B., Ranger, J., and Legout, A.:

Mg and Ca uptake by roots in relation to depth and allocation to aboveground tissues: results from an isotopic labeling study in a beech forest on base-poor soil. *Biogeochemistry*, 122, 375-393, doi:10.1007/s10533-014-0047-2, 2015.

590 Verbruggen, N., and Hermans, C.: Physiological and molecular responses to magnesium nutritional imbalance in plants. *Plant Soil*, 368, 87-99, doi:10.1007/s11104-013-1589-0, 2013.

Wang, R., Dungait, J. A. J., Creamer, C. A., Cai, J., Li, B., Xu, Z., Zhang, Y., Ma, Y., and Jiang, Y.: Carbon and nitrogen dynamics in soil aggregates under long-term
595 nitrogen and water addition in a temperate steppe. *Soil Sci. Soc. Am. J.*, 79, 527-535, doi:10.2136/sssaj2014.09.0351, 2015.

Wang, R., Dungait, J. A. J., Buss, H. L., Yang, S., Zhang, Y., Xu, Z., and Jiang, Y.: Base cations and micronutrients in soil aggregates as affected by enhanced nitrogen
600 and water inputs in a semi-arid steppe grassland. *Sci. Total Environ.*, 575, 564-572, doi:<http://dx.doi.org/10.1016/j.scitotenv.2016.09.018>, 2017.

Wang, X., Cheng, G., Zhong, X., and Li, M. H.: Trace elements in subalpine forest soils on the eastern edge of the Tibetan Plateau, China. *Environmental Geology* 58, 635-643, <https://doi.org/10.1007/s00254-008-1538-z>, 2009.

Weatherall, A., Proe, M. F., Craig, J., Cameron, A. D., and Midwood, A. J.: Internal
605 cycling of nitrogen, potassium and magnesium in young Sitka spruce. *Tree Physiol.*, 26, 673-680, doi:<https://doi.org/10.1093/treephys/26.5.673>, 2006.

White, J. G., and Zasoski, R. J.: Mapping soil micronutrients. *Field Crop. Res.*, 60, 11-26, doi:10.1093/aob/mcg164, 1999.

Welch, R. M., and Shuman, L.: Micronutrient nutrition of plants. *Crit. Rev. Plant Sci.*,
610 14, 49-82, doi:<http://dx.doi.org/10.1080/07352689509701922>, 1995.

Zhu, W., Xiang, J., Wang, S., and Li, M.-H.: Resprouting ability and mobile carbohydrate reserves in an oak shrubland decline with increasing elevation on the eastern edge of the Qinghai-Tibet Plateau. *Forest Ecol. Manag.*, 278, 118-126, doi:<https://doi.org/10.1016/j.foreco.2012.04.032>, 2012.

615 Zhu, W., Cao, M., Wang, S., Xiao, W., and Li, M.-H.: Seasonal dynamics of mobile
carbon supply in *Quercus aquifolioides* at the upper elevational limit. PLoS ONE, 7,
e34213, doi:<https://doi.org/10.1371/journal.pone.0034213>, 2012.

Tables 

Table 1 Two-way ANOVAs (*F* values) of the effect of plant life form (L, tree or shrub), elevation position (E), and their interactions on soil pH,

620 soil organic carbon (SOC), base cations and micronutrients in soils (exchangeable/available form) and plants (total) across sampling sites.

	soil pH	SOC	soil Ca	Soil Mg	Soil K	leaf Ca	root Ca	shoot Ca	stem Ca	leaf Mg	root Mg	Shoot t Mg	stem Mg	leaf K	root K	shoot K	stem K	soil Fe	soil Mn	soil Zn	leaf Fe	root Fe	shoot Fe	stem Fe	leaf Mn	root Mn	shoot Mn	stem Mn	leaf Zn	root Zn	shoot Zn
L	0.18	5.75*	1.0 2	8.15 ** 4**	24.6	9.74	1.69	10.8 5**	39.3 6** 1**	93.4	8.35	15.8 2**	118.5	12.5 7	6.4 1	63.5 6	87.0 0**	8.72 0	2.6 6.89 7	64.6 2	0.7 2	0.1 2	8.54 27.0 5	40.6 7** 7	25.1 3** 7	11.8 3** 1	2.0 0	4.4 0	80.6 0*		
E	3.16*	10.1	1.2 2**	1.71	15.2	7.82	3.85	8.36*	1.17	0.83	5.85	2.42	10.47	9.30*	0.2 2	0.48	0.85	5.75 4	1.5 0.01	0.02	1.8 0	0.5 0	2.40	7.13 2.63	1.54	0.53	0.67 1	0.7 7	1.6 3.61*		
L×E	15.5 7**	3.34*	2.4 0	0.55	8.61*	2.24	8.55	6.61*	1.11	1.82	7.20	5.23*	0.37	0.96	1.0 1	2.20	2.84	0.10	1.6 9	4.42 0.36	0.36	3.4 0*	3.7 3*	0.61	0.15	2.81	1.22	2.37	0.90 3	1.9 4	1.3 2.12

* and ** indicate significant level at $P < 0.05$ and 0.01, respectively.

Table 2 Regression statistics relating soil base cations and micronutrients to other soil physicochemical parameters under all trees and shrubs across three sampling sites.

625 The TN, C:N, TIN, and Olsen P represent soil total nitrogen, SOC to TN ratio, total inorganic nitrogen, and Olsen phosphorus.

	Ca	Mg	K	Fe	Mn	Zn
pH	0.94 **	0.87 **	-0.06	-0.64 **	-0.36 **	-0.73 **
SOC	0.12	0.24*	0.18	0.30**	0.07	0.37 **
TN	0.39 **	0.51 **	0.18	-0.01	0.03	0.11
C:N	-0.45**	-0.44**	-0.11	0.60 **	0.02	0.44**
NO ₃ ⁻	-0.24*	-0.29 **	0.28 **	-0.05	0.02	0.08
NH ₄ ⁺	-0.38**	-0.33**	-0.10	0.46**	0.15	0.63 **
TIN	-0.38**	-0.39**	0.19*	0.17	0.08	0.36**
Olsen P	0.10	0.18	0.49 **	-0.05	0.04	0.03

* and ** indicate significant level at $P < 0.05$ and 0.01, respectively.

Figures

630




Fig. 1 Soil pH values (a) and soil organic carbon (SOC) concentration (b) at lower and middle elevations as well as at the upper limit of treelines or shrublines for each of the three sites. Different letters indicate significant differences among three elevations within treeline or shrubline for each site.

635





Fig. 2 Concentrations of soil exchangeable base cations of Ca (a), Mg (b) and K (c)


and available micronutrients of Fe (d), Mn (e) and Zn (f) at lower and middle

640 elevations as well as at the upper limit of trees or shrubs for each of the three sites.

Different letters indicate significant differences among three elevations within treeline or shrubline for each site.



645 **Fig. 3** Base cation concentrations of Ca (a, b, c, d), Mg (e, f, g, h) and K (i, j, k, l) in
 plant tissues of leaf, root, shoot and stem sapwood at lower and middle elevations as
 well as at the upper limit of trees or shrubs for each of the three sites. Different letters
 indicate significant differences among three elevations within treeline or shrubline for
 each site.



650

Fig. 4 Micronutrient concentrations of Fe (a, b, c, d), Mn (e, f, g, h) and Zn (i, j, k, l)

in plant tissues of leaf, root, shoot and stem sapwood at lower and middle elevations

as well as at the upper limit of trees or shrubs for each of the three sites. Different

letters indicate significant differences among three elevations within treeline or

655 shrubline for each site.