

# ***Interactive comment on “C<sub>5</sub> glycolipids of heterocystous cyanobacteria track symbiont abundance in the diatom *Hemiaulus hauckii* across the tropical north Atlantic” by Nicole J. Bale et al.***

**Nicole J. Bale et al.**

nicole.bale@nioz.nl

Received and published: 3 October 2017

We thank the reviewer for their thoughtful and constructive review. We are pleased he/she found our data set “valuable” and that “the topic of the manuscript is definitely of great interest”. We found the reviewer’s edits useful and have addressed their points below.

1. In the result section I would recommend to group the different stations, as already done to a certain degree in paragraph 3.4. (“high-salinity open ocean sites”, “coastal-

Printer-friendly version

Discussion paper



shelf stations”, etc.). The naming of individual stations with only their number, paired with a very detailed description of concentrations at each of them, makes it hard to filter out the most relevant trends.

We appreciate this idea of grouping the stations and will edit as per suggested.

2. Short chain (C26) C5 HGs were initially described by Wörmer et al. (2012) in freshwater systems and a culture, before the description of longer chain C5 HGs by Schouten et al. (2013) in symbionts. As Wörmer et al (2012) only described C26 HGS, I would generally recommend to clearly differentiate between long- and short-chain C5 HGs throughout the text, e.g. in the conclusions “long-chain C5 HGs provide a robust, reliable method for detecting DDAs”. Such a differentiation would make the authors’ statements much more robust, as it eliminates potential interference from the short chain C5 HGs.

Indeed this is an important differentiation, we will edit to make this clear throughout the manuscript that the C5 HGs associated with DDAs have C30 and C32 chains as opposed to the C26 chain seen in the study of Wörmer et al. (2012).

3. More importantly, I think that the discussion of the correlation between long-chain C5 HGs and different DDAs and free-living cyanobacteria needs to be improved to solidify the claim of a diagnostic relationship. For example, cross-plots and regression curves should be shown, instead of only stating  $r$  and  $p$  values. Based on these values alone, actually a strong correlation of C5 HGs is also observed with *Trichodesmium* colonies, and this harms the proposed biomarker potential. Even though I may share the authors’ opinion that this regression might be coincidental and due to the co-occurrence of *Trichodesmium* and DDAs at certain stations, a better effort to demonstrate the specific correlation between DDAs and C5 HGs is mandatory. In this sense I would for example recommend to pay special attention to the values which deviate from the regression. Assuming that several source organisms for C5 HGs exist, the fact that one potential producer (e.g. *Rhizoselenia* symbionts) is not abundant when C5 HGs are

[Printer-friendly version](#)[Discussion paper](#)

highly concentrated does not imply that it is not a potential source organism, as other producers may be present (e.g. *Hemiaulus* symbionts). This concept is hinted at in l. 281-290, but should be expanded. On the other hand, abundance of an organism without corresponding HG abundance (e.g. maxima of *Trichodesmium* colonies) is a much more robust factor to rule out a potential source organism. In this sense it might also be interesting to plot a combined regression line for all DDAs vs C5 HGs.

We agree that presenting the data as regression curves would strengthen our assertion that long-chain C5 HGs track the abundance of certain marine DDAs. We realize that only stating  $r$  and  $p$  values from Pearson Correlations doesn't give a full picture of the data. We have plotted the regressions as suggested (see attached Figure). We will include them in the supplement of the revised manuscript and describe them in the text. Indeed, looking at them brings an extra dimension to the results/discussion. While the number of *Hemiaulus* symbionts has the best correlation with the C5 HG concentration ( $r^2 = 0.62$ ), that of the *Trichodesmium* colonies isn't far off ( $r^2 = 0.48$ ). However, there is a clear outlier in the *Hemiaulus* symbiont vs. C5 HG concentration plot, which is station 8 at 10 m water depth. As the DDAs and *Trichodesmium* are all surface dwellers (upper 5 m) we assume that this point related to dead/detrital HGs, whereas viable DDAs were not visible under the microscope. Hence we also plotted the four regressions for only surface data ( $n=19$ ). Here the correlation between the number of *Hemiaulus* and the C5 HG concentration is stronger still ( $r^2 = 0.97$ ), and again so is that of the *Trichodesmium* colonies ( $r^2 = 0.94$ ). On closer examination of these regressions it is clear that one station, again station 8, with unusually high levels of both *Hemiaulus* symbionts and *Trichodesmium* colonies (station 8) drives both these correlations. Removal of station 8 from the regressions showed that the number of *Hemiaulus* symbionts has yet again the best correlation with the C5 HG concentration ( $r^2 = 0.67$ ) whereas the correlation with *Trichodesmium* colonies has disappeared ( $n=0.03$ ), as would be expected. Interestingly in this sample subset there is a stronger correlation between *Rhizosolenia* symbionts and the C5 HG concentration ( $r^2 = 0.56$ ). We will include this information in the revised discussion.

[Printer-friendly version](#)[Discussion paper](#)

4. Finally, the authors claim that the analyzed compounds are ribose-containing, but I couldn't find any description of how the sugar moiety has been characterized. If they haven't been described I would rather use the term pentose (as hexose is used for the C6 compounds).

We did not include this information in this manuscript as it was published previously but the sugar was identified as part of the study of Schouten et al. 2013. "Comparison of the retention time and mass spectrum of an authentic methyl ribose standard established that the C5 sugar was ribose, though the stereo-configuration of the sugar could not be determined. However, since L-ribose does not occur in nature and is only produced synthetically (e.g. Hu et al., 2011) we assume that the sugar moiety is D-ribose. Thus, glycolipid VII was identified as 1-(O-ribose)-3,27,29-triacontanetriol." We will include clearer reference to this structural information arising from the previous study in our manuscript.

Another issue is that the manuscript preparation sometimes seems a little careless, and I would appreciate a thorough revision. Some minor comments and edits include:

Text is sometimes indented, sometimes not.

This was our understanding of the BGS style. Don't indent first paragraphs but then indent thereafter. We will double check the indenting requirements and further check our indentations.

I.12-13: "have a thickened cell walls" please correct use of singular/plural

We will make this edit.

I.14: use singular form "cyanobacterium" or plural verb form "make"

We will make this edit.

I.43: please specify that you are referring to heterocystous cyanobacteria "all heterocystous, non-symbiotic cyanobacteria"

[Printer-friendly version](#)[Discussion paper](#)

We will make this edit.

I.45: It might be better to already mention here that short chain C5 HGs have been described in a non-symbiotic cyanobacterial culture, not only C6 HGs.

We will mention this here.

I.52: It is confusing to state that the “first study of the C5 HGs in the natural environment” was Bale et al. 2015 while providing the fact that “HG with a C5 sugar moiety” were identified in freshwater environments three years earlier.

We will edit this to make it clear that C5 HGs have been detected twice previously in a marine and fresh water environment, but that is the first study in the marine environment that correlates their presence with DDA cell counts.

I.111: Station number is missing, maybe 10?

Yes it should have been Station 10. We edit to correct this mistake.

I.114: add “each”, “For each sediment”

We will make this edit.

I.120: “freeze dried filtered seawater”. I guess the lipids are extracted from the filters, not from the filtered seawater, right?

This is correct. We will edit this phrase.

I.145: Just out of interest, have the authors tried to increase flow to shorten analysis

When developing the LC-MS method we examined the flow rate and found it to be suitable for separating a wide range of IPLs and maximizing the ionization stability. We do not discuss the LC-MS method development in this manuscript as it was not within the scope of the paper. We appreciate the reviewer’s suggestion and will keep it in mind when reviewing the method for further application.

I.149: at which m/z is resolution measured?

The resolution stated in our method (70,000 rpm) is for m/z 200. We will include this information in our revised manuscript.

I.178: is 36.3 a value for salinity?

We will insert 'salinity of' into this sentence

I.180: "(Fig. 3c,d)" close parentheses

We will make this edit.

I.180: "NO<sub>3</sub>+NO<sub>2</sub>" (no subscript for "+")

We will make this edit.

I.212: may be rephrased: "Free Trichodesmium trichomes were broadly distributed (Fig. 4d) and often occurred

We will make this edit.

i.266: delete space: "Hemiaulus hauckii-Richelina"

We will make this edit.

I.290-293: The separation of DDAs depending on salinity with the current data is unclear, as the authors state. Therefore I would delete this topic and also the corresponding figure.

As the reviewer finds this topic and figure unclear we will remove the figure (6) and simplify this text to clarify our point.

I.304-308: I think the sampling-volume explanation is a little confusing. Couldn't the authors just state that sensitivity of the chemical biomarker method is much more sensitive than the microscopic approach?

We will edit this section to make it briefer and clearer to the reader

I.347: please rephrase to avoid the term "vegetal"

[Printer-friendly version](#)

[Discussion paper](#)



We will replace this word with “planktonic organisms”.

I.352: add “regarding” or similar: “difference regarding the limit of detection”

We will make this edit.

I.600: “Trichodesmium” should be in italics

We will make this edit.

I.603: use “dashed” instead of “broken”?

We will make this edit.

I.611-612, Table 1: “\*” is not defined. Please define BMWL and DCM, even though they are already defined in the text, table should be informative on its own. Same for figure S1, where actually BML is used.

We will make these edits to the Table and Fig S1.

I.629: are Trichodesmium colonies expressed as colonies or trichomes/ml?

We will make it clear in this Supplement legend that Trichodesium was enumerated both in terms of Trichodesium colonies L-1 and as free Trichodesium trichomes L-1.

Figures: Please use larger fonts.

We will increase font size in figures

Figure 2-4: I think it would be better to place the axis legend (e.g. Salinity in figure 2) to the right, instead of on top of the figure. Especially in fig 3 and 4 this makes it easier to identify what is shown.

We will make these edits to the figures

Figure 4: (d) is used twice, for panel (d) and what should be panel (e). Why is C32 C5 HG shown as %? Wouldn't it be more informative to show concentration?

[Printer-friendly version](#)

[Discussion paper](#)



We will edit the panel numbering and change the figure to show concentration of C32 C5 HG rather than percent of total HG.

---

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2017-300>, 2017.

**BGD**

---

Interactive  
comment


Printer-friendly version

Discussion paper






A) All data (n=54)



B) Only surface data (n=19)



C) Only surface data without station 8 (n=18)




Fig. 1. Regression curves for reply to Reviewer 2

Printer-friendly version

Discussion paper

